There and back again: the dynamic evolution of panarthropod germ cell specification mechanisms

Jonchee A. Kao[1†], Emily L. Rivard[1†], Rishabh R. Kapoor[2], Cassandra G. Extavour[1,3,4] *

1. Department of Molecular and Cellular Biology, Harvard University, Cambridge MA, USA
2. Program for Systems, Synthetic and Quantitative Biology, Harvard University, Boston MA, USA
3. Department of Organismic and Evolutionary Biology, Harvard University, Cambridge MA, USA
4. Howard Hughes Medical Institute, Chevy Chase MD, USA

† equal contribution

* correspondence: extavour@oeb.harvard.edu

Abstract

Germ cells enable the reproduction of an organism and the continuity of its lineage.

These crucial cells are segregated from the soma at different times and places and via

distinct mechanisms across the animal tree of life. Making sense of the evolution of

germ cell specification across animals is complicated by the difficulty of making

meaningful comparisons across the diversity of animal embryonic development. Here,

we characterize germ cell specification within a clade as old as all animals, which

encompasses massive animal biodiversity but within which it is still possible to conduct

comparative embryogenesis: the Panarthropoda. We amass data from centuries of

studies describing the timing and mechanisms of germ cell formation, and use these

data to propose novel hypotheses about the trajectory of evolution in this process.

Furthermore, we speculate on the mechanisms underlying these evolutionary dynamics

by considering the relationships among germ cell specification, concurrent

developmental processes, and the germ line gene network. Collectively, this study

derives new insights from a rich historical database of embryological observations, and

has broad implications for understanding the evolution of metazoan germ cells.

Keywords: germ plasm, pole cells, inductive signaling, BMP, cytoplasmic determinant

I. Introduction

During development, animals specify the germ line, a collection of cells that will later give rise to the next generation. The process of germ cell specification across animals has intrigued researchers for more than 150 years. Whereas historical researchers relied primarily on observational data alone, we can now investigate these processes using functional genetic tools. These new techniques allow us not only to identify germ cells with more certainty, but also to gain more insight into the molecular mechanisms of germ cell specification. Several hypotheses have been proposed to explain evolutionary transitions between these mechanisms across animals. However, such hypotheses are difficult to test because embryonic development of animals is too different across vast evolutionary time scales to be meaningfully compared. Moreover, there is disagreement in the field about whether existing well-studied mechanisms of specification are truly the only modes, and whether they are discrete, mutually exclusive mechanisms. Further, there are conflicting reports about the impact of germ cell specification mode on the evolution of genomes, body plans, and fitness (Evans et al., 2014; Whittle and Extavour, 2016; Whittle and Extavour, 2017) . Here, we examine variation in germ cell specification across the most speciose phylum of animals, within which the general features of embryogenesis can nevertheless be meaningfully compared: the Panarthropoda. This clade displays striking diversity in the temporal and spatial origin of germ cells, as well as multiple transitions between specification mechanisms, offering an ideal system to study evolution and regulation of germ cell specification.

In this study, we first review the existing observational data regarding embryonic

germ line development across panarthropods. We are primarily interested in the first

embryonic appearance of germ cells before they enter the gonad, during which time

they are referred to as *primordial germ cells*. We identify three categories of temporally

distinct developmental context in which primordial germ cells are first detectable during

embryogenesis and analyze the distribution of these three categories in their

phylogenetic context. We then review perturbational experimental studies aimed at

uncovering the mechanisms of germ cell specification. Finally, we discuss the

implications of these data on our understanding of the evolution of germ cell

specification in panarthropods, and more broadly across Metazoa.

I.i. How do we identify germ cells?

Historically, embryologists studied early development using histological

preparations of sectioned embryos. With light microscopy, primordial germ cells can be

readily distinguished from surrounding somatic cells by their rounder and larger nuclei

(Nieuwkoop and Sutasurya, 1979). The chromatin of germ cells often appears more

diffuse than that of somatic cells, and the cytoplasm, especially cortical or perinuclear

cytoplasm, may contain electron-dense, non-membrane bound granular material (Eddy,

1975). However, tracking these cells throughout development using fixed samples and

morphological features alone is not always technically possible.

Since the late 20[th] century, studies assessing gene expression patterns unique to

or diagnostic of germ cells (henceforth referred to as "molecular markers") have been

instrumental in identifying germ cells at earlier stages than previously believed. The

most commonly used markers are the mRNA or protein products of the genes *vasa*,

nanos, *tudor*, and *piwi* (detectable by *in situ* hybridization or antibody staining,

respectively) (Ewen-Campen et al., 2010). The use of these molecular germ cell

markers can enable their earlier identification, since some germ cells express their

encoding genes before their morphological differentiation becomes detectable. In

addition, these markers facilitate tracking of germ cells over the course of development.

However, using molecular markers is a biased approach, as it relies on previous

knowledge of germ line gene expression from other species. Furthermore, interpretation

of molecular marker data can be complicated by the fact that their encoding genes are

sometimes also expressed by somatic cells (e.g. Yajima and Wessel, 2011).

II. Germ cell formation in panarthropods

In our review of the embryological literature from 1860 to the present, we

identified three categories of primordial germ cell specification in panarthropods, based

on the timing of specification during embryogenesis (Figure 1). We describe these

categories below, along with relevant details of embryogenesis for context.

II.i. Category 1: PGC specification at early cleavage and blastoderm stages

After fertilization, embryogenesis begins with *cleavage*, a series of mitotic

divisions of the zygote (Gilbert, 1997). In most arthropods, cleavage leads to the

formation of the *blastoderm*, a relatively uniform layer of cells at the egg cortex,

surrounding a central yolk mass populated by a small number of yolk cells (Anderson,

1973; Scholtz and Wolff, 2013). The first category of primordial germ cell specification in

panarthropods occurs during or before the formation of the blastoderm.

Insects

In most insects, the blastoderm is initially *syncytial*, with no cell membranes

separating the nuclei (Anderson, 1973; Johannsen and Butt, 1941). In some insects, the

primordial germ cells are the first cells to form, appearing at the posterior pole before

the rest of the blastoderm is cellularized (Figure 1A1). Primordial germ cells that appear

in this manner are called *pole cells*. Pole cells are most commonly seen in species of

Diptera, Coleoptera, and Hymenoptera, although they have also been reported in

Thysanoptera and Dermaptera (Table S1).

Pole cells were first reported in species of the midge *Chironomus* by Weismann

(Weismann, 1863a), although Metchnikoff (1866) was the first to follow their migration

into the gonad. In many insects, the cytoplasm at the posterior pole of the embryo is

morphologically distinct from that of the rest of the embryo, often with visible granules

(Hegner, 1914; Zissler, 1992). The close association of this granular cytoplasm with

pole cells led researchers to speculate that the granular material contained germ line

determinants (e.g. Hegner, 1909a; Zissler, 1992); we will describe experimental data

testing this hypothesis below. It was long believed that pole cells gave rise to somatic

cell types in addition to the germ cells (Anderson, 1962a; Anderson, 1962b; Poulson,

1947; Poulson, 1950). However, careful lineage tracing in the fruit fly *Drosophila*

melanogaster showed that all pole cells are determined as germ cells, and any that

become lost during their migration die (Underwood et al., 1980).

While the function of pole cells as primordial germ cells appears conserved, the

specific details of their formation can vary considerably across species. For example,

pole cells differ in which cleavage cycle they cellularize. Pole cells form during the ninth

mitotic cycle of *D. melanogaster* (Huettner, 1923) and during the sixth mitotic cycle in

the parasitic wasp *Nasonia vitripennis* (Buchta et al., 2013). To our knowledge, the

earliest observed formation of pole cells is during the third cleavage cycle in the midges

Miastor metraloas and *Chironomus sp*. (Hegner, 1914; Weismann, 1863b). The number

of pole cells formed is also variable: a single pole cell forms in embryos of the latter

midges (Hegner, 1914; Weismann, 1863b), whereas four appear in the fruit fly

Bactrocera tryoni (Anderson, 1962a) and sixteen appear in the beetle *Calligrapha*

multipunctata (Hegner, 1909b).

After the pole cells form, they enter the egg and migrate to the location of the

future embryonic gonad, which may be quite far from the posterior pole. Several

different migration pathways have been described. In *D. melanogaster* and many other

Diptera, the pole cells are carried into the egg during posterior midgut invagination, then

migrate through the epithelium of the posterior midgut to enter the abdominal mesoderm

and associate with the developing somatic gonad (Metschnikow, 1866; Sonnenblick,

1941). In Hymenoptera, pole cells are internalized with the rest of the mesoderm during

gastrulation (Gatenby, 1917). Yet another variation is observed in chrysomelid beetles

(Hegner, 1909a), where germ cells are internalized with the rest of the germ band as

the extraembryonic membranes fold over the germ band.

Other arthropods

Primordial germ cells are also specified early in Collembola, an early-diverging

branch of non-insect hexapods (Du et al., 2024) REF. Embryogenesis in these animals

begins with a period of holoblastic cleavage, and certain blastomeres on the basal (yolk)

side of the blastoderm can be recognized as likely primordial germ cells very early

(Jura, 1967; Klag, 1982). The cytoplasm of these germ cells contains granular material

visible by light and electron microscopy (Klag, 1982) that resembles the cytoplasm

containing germ line determinants observed in insects (Hegner, 1914; Zissler, 1992).

Rather than maintaining contact with the germ band as in insects, collembolan germ

cells migrate into the center of the yolk after their specification until eventually migrating

to associate with the abdominal mesoderm, from which the embryonic gonad develops

(Jura, 1967).

Among non-hexapod Pancrustacea, embryos of Copepoda, Amphipoda,

Decapoda, and some Cladocera undergo stereotypical holoblastic cleavage, allowing

precise tracking of cell lineages and the identification of a specific cleavage blastomere

as the progenitor of primordial germ cells (Gerberding et al., 2002; Hertzler, 2005; Kühn,

1913; McClendon, 1907). In amphipods and decapods, a putative germ plasm has been

identified in the freshly laid egg, termed the RNA-containing body (Gupta and Extavour,

2013) or intracellular body (Biffis et al., 2009; Chen et al., 2021). During cleavage, this

region of cytoplasm is inherited asymmetrically by the blastomeres that will give rise to

the primordial germ cells (Gupta and Extavour, 2013) (Figure 1A2).

In chelicerates, candidate primordial germ cells are first detectable with

molecular markers in the yolk of blastoderm-stage embryos in the mite *Tetranychus*

urtichae (Dearden et al., 2003). As the germ band forms, the primordial germ cells

associate with the posterior germ band near the primordia of the fourth walking legs. We

are not aware of any other chelicerate or myriapod species in which primordial germ cell

specification has been proposed to occur before gastrulation.

II.ii. Category 2: PGC specification at gastrulation

After formation of the blastoderm, a portion of the blastoderm is specified as the

presumptive embryonic region, variably referred to as the *germ band*, *germ rudiment*,

embryonic rudiment, or *germ disc* depending on its initial form (Anderson, 1973;

Johannsen and Butt, 1941). The rest of the blastoderm differentiates into one or more

extraembryonic membranes (Johannsen and Butt, 1941). Soon after this specification,

gastrulation occurs, whereby some of the blastoderm cells are internalized and become

the mesoderm. In insects, mesoderm internalization primarily occurs along the ventral

midline of the germ band (Leptin, 2004; Roth, 2004). In other arthropods, mesoderm

may be internalized through a *blastopore* near the germ rudiment (Chaw et al., 2007;

Gerberding and Patel, 2004), or by epiboly (Edgar et al., 2015). The second category of

primordial germ cell differentiation occurs at or very close to the time of gastrulation.

Insects

Among insects, this timing of primordial germ cell specification is most common

among members of Hemiptera, Psocodea, and Lepidoptera, although it is also reported

in species of Dermaptera, Blattodea, Hymenoptera, Megaloptera, Strepsiptera,

Coleoptera, Mecoptera, and Diptera. We observed three different spatial patterns of

germ cell specification around the time of gastrulation: before germ band formation, at

the posterior of the germ band after its formation, or at the ventral midline of the

blastoderm just before gastrulation. In Hemiptera, Psocodea, and tenebrionid beetles

(Coleoptera), germ cells are first detectable at the ventral posterior region of the cellular

blastoderm in an invaginating structure called the *posterior pit* (Figure 1B) (Butt, 1949a;

Goss, 1952a; Mellanby, 1935a; Schröder, 2006; Shinji, 1919). These cells appear to

form on the side of the blastoderm epithelium facing the yolk, rather than on the outside

of the blastoderm like the pole cells described in Category 1. In Hemiptera and

Psocodea, the invagination continues into the egg towards the anterior, forming a

definitive germ band with germ cells at its posterior tip (Figure 1B1). Germ cells in

tenebrionid beetles move further inside the embryo once the amnion (ventral

extraembryonic membrane) has enclosed the entire ventral surface of the germ band,

similar to the internalization seen in Coleoptera that specify their germ cells as pole

cells. Following their formation, the germ cells migrate along the dorsal side of the germ

band toward the anterior, eventually settling into the somatic gonad in the abdominal

mesoderm.

In several species with this timing of germ cell specification, primordial germ cells

were first identified at the posterior tip of the germ band after it has detached from the

blastoderm [*e.g.*, *Periplaneta orientalis* (Blattodea) (Heymons, 1895), *Sialis mitsuhashii*

(Megaloptera) (Suzuki et al., 1981), *Nematus ribesii* (Hymenoptera) (Singh, 1967), and

Mecoptera (Suzuki, 1990)]. Early studies of tenebrionid beetles described this pattern of

germ cell differentiation as well (Rempel and Church, 1969; Ullmann, 1964). However,

more recent work using genetic markers suggests that primordial germ cells in at least

one such beetle, *Tribolium castaneum*, appear earlier, during the cellular blastoderm

stage in a posterior pit much like the Hemiptera (Schröder, 2006). This raises the

possibility that in the other tenebrionid species listed above, primordial germ cells may

likewise differentiate from the blastoderm, with the reported later appearance in the

germ band simply reflecting limitations in older detection methods.

In contrast to the posterior pit, germ cells differentiate along the ventral midline of

the blastoderm in Lepidoptera, separating from the blastoderm before gastrulation to lie

on the dorsal side of the presumptive germ band in the presumptive abdominal region

(Ando and Tanaka, 1980; Miya, 1958a; Presser and Rutschky, 1957; Tanaka, 1987;

Woodworth, 1889). Unlike when germ cells form at the posterior pit or at the posterior

tip of the germ band, in this case there does not appear to be long-range germ cell

migration, since the germ cells apparently form where the gonad eventually forms in the

abdomen.

Other arthropods

Among non-insect Pancrustacea, primordial germ cells are the first cells to be

internalized from the blastopore lip in Mysida (Manton, 1928), Anomopoda (Cannon,

1921), and the copepod orders Siphonostomatoida (McClendon, 1907), Harpacticoida

(Witschi, 1934), and Cyclopoida (Schimkewitsch, 1896). In Isopoda, primordial germ

cells differentiate from the mesendoderm soon after internalization (Figure 1B2)

(Goodrich, 1939a; Needham, 1942; Strömberg, 1965; Wolff, 2009). Thus, primordial

germ cell differentiation seems to be tied to gastrulation in these crustacean orders.

In chelicerates, primordial germ cells have been identified based on

morphological criteria at the blastopore in Opiliones (Figure 1B3) (Faussek, 1888;

Faussek, 1892; Holm, 1947) and Scorpiones (Brauer, 1894). A recent study using

molecular markers supported this interpretation in Opiliones (Gainett et al., 2022).

Putative primordial germ cells have also been identified at the blastopore in the

centipedes *Scolopendra cingulata* and *Strigamia maritima* (Myriapoda) (Green and

Akam, 2014; Heymons, 1901).

Tardigrade, germ cells are the first cells to enter the blastopore during

gastrulation, based on data from both morphological characteristics (Hejnol and

Schnabel, 2005) and molecular markers (Heikes et al., 2023) (Figure 1B4). Lineage

tracing in these embryos suggests that the germ cells are derived from both

blastomeres of the two-cell embryo (Gabriel et al., 2007; Hejnol and Schnabel, 2005).

Finally, onychophoran germ cells in different species emerge either at the

blastopore lip during gastrulation (Manton, 1949) or at the posterior end of the germ

band during elongation (Manton, 1949; Mayer and Tait, 2009; Sedgwick, 1887). We are

not aware of any studies using molecular markers to track germ cell formation in

Onychophora (Figure 1B5).

II.iii. Category 3: PGC specification from mesoderm

After gastrulation, the germ band elongates and becomes segmented, and the

mesoderm becomes organized into pairs of somites in each segment. In most living

arthropods, each somite forms a small *coelom* enclosing a *coelomic cavity* (Anderson,

1973; Johannsen and Butt, 1941). The coelomic cavities are eventually lost as the

mesoderm differentiates into different cell types and reorganizes into the primordia of

different organ systems, including the somatic gonad (Anderson, 1973; Johannsen and

Butt, 1941). The final category of primordial germ cell differentiation timing

encompasses species in which primordial germ cells differentiate from the mesoderm at

this comparatively late stage of embryogenesis.

Insects

In insects, this category includes members of Archaeognatha, Zygentoma,

Orthoptera (Figure 1C1), Mantodea, Blattodea, Trichoptera, Coleoptera, and

Lepidoptera. Primordial germ cells may differentiate before formation of the coelomic

cavities [e.g., Orthoptera (Ewen-Campen et al., 2013a), Mantodea (Görg, 1959)], after

formation of the coelomic cavities [e.g., Archaeognatha (Larink, 1969), Zygentoma

(Woodland, 1957)], or after the coelomic cavities are no longer recognizable [e.g.,

Trichoptera (Miyakawa, 1974), *Odontotaenius disjunctus* (Coleoptera) (Krause, 1947),

Apis mellifera (Hymenoptera) (Dearden, 2006; Nelson, 1915)]. In this developmental

timing category, primordial germ cells are usually specified in the same abdominal

segments where the somatic gonad forms, so there is no long-range movement of

primordial germ cells after their specification.

Other arthropods

Differentiation of primordial germ cells from the mesoderm has also been

described in Leptostraca (Figure 1C2) (Manton, 1934), Anaspidacea (Hickman, 1936),

and Branchiopoda (Anderson, 1967; Mitsumoto and Makioka, 2002; Sudler, 1899).

However, data for stages of crustacean development late enough to include

differentiation of the trunk mesoderm from which primordial germ cells might arise are

sparse. This relative scarcity of early embryonic descriptions may be in part because

many crustaceans hatch as a nauplius, comprising only the head segments (Scholtz,

2000), and germ cells might not be specified until later larval stages, after formation of

the trunk. Many studies of naupliar and larval development in crustaceans are focused

on organogenesis of the gonad as a whole, without specific reference to the origin of

primordial germ cells (Cannon, 1924; Ikuta and Makioka, 1994; Mitsumoto and Makioka,

2002).

Classical studies on myriapods suggest a late differentiation of primordial germ

cells from the mesoderm in three of the four myriapod classes [Pauropoda (Tiegs,

1947), Symphyla (Tiegs, 1940), and Diplopoda (Dohle, 1964)], although only one

species was reported from each class in these studies. Germ cells in these classes

have not been investigated with molecular markers to our knowledge.

Among chelicerates, primordial germ cells differentiate from the mesoderm of the

opisthosomal segments in spiders (Figure 1C3) (Gough, 1902; Kautzsch, 1910;

Schimkewitsch, 1903). Data from molecular markers are consistent with this

observation (Schwager et al., 2015). Primordial germ cells are first detectable in the

mesoderm of the prosoma in pycnogonids (Alexeeva and Tamberg, 2021; Alexeeva et

al., 2018; Miyazaki and Makioka, 2012).

II.iv. Summary :evolutionary trends from late to early germ cell specification

In his monograph on embryogenesis in Dermaptera and Orthoptera, Richard

Heymons (Heymons, 1895) proposed that "the formation of sex cells at the rear end of

the embryonic primordium, which has now been clearly demonstrated for

representatives of very different groups, is probably generalizable for all insects." Rather

than making broad statements such as this, we summarize the data regarding the

timing of germ cell specification in the context of currently accepted panarthropod

phylogeny (Ballesteros et al., 2022; Fernández et al., 2018; Giribet and Edgecombe,

2019; Misof et al., 2014; Schwentner et al., 2017) in Figures 2–3. This shows that in

early diverging euarthropod orders, primordial germ cells tend to be first observed later

in embryogenesis (Category 3). The timing of first germ cell appearance evolved to be

earlier in development in multiple lineages independently (Dermaptera, Condylognatha,

and Holometabola within insects; Copepoda, Malacostraca, and Cladocera within non-

hexapod Pancrustacea; Chilopoda within myriapods; and Opiliones, Scorpiones, and

mites within chelicerates). These data are consistent with the hypothesis that

differentiation from the mesoderm is the ancestral mode of primordial germ cell

specification in Euarthropoda, a hypothesis which we formally test below.

Conflicting interpretations in the literature about the timing of germ cell

specification have sometimes resulted from technical advances enabling identification of

germ cells at stages earlier than previously detected. For example, early work on the

silk moth *Bombyx mori* (Lepidoptera) reported primordial germ cells emerging from the

mesoderm of coelomic sacs in late embryogenesis (Category 3) (Toyama, 1902), while

later studies identified primordial germ cells on the posterior side of the ventral midline

of the early germ band (Category 2) (Miya, 1953). The onset of primordial germ cell

specification before gastrulation was later further supported by data from molecular

markers (Nakao, 1999; Nakao et al., 2006). Therefore, it is possible that species that

are currently classified in Category 3 may be reclassified in Categories 2 or 1 by future

work deploying molecular markers. This caveat notwithstanding, we note that molecular

markers and functional genetic studies often corroborate late germ cell specification

[see for example *Gryllus bimaculatus* (Orthoptera, Insecta) (Ewen-Campen et al.,

2013a), *A. mellifera* (Hymenoptera, Insecta) (Dearden, 2006), *Parasteatoda*

tepidariorum (Araneae, Chelicerata) (Schwager et al., 2015)]. We therefore do not

assume that the hypothesis of ancestral late germ cell specification is an artifact of

exclusive use of morphological identification methods.

Going forward, data from key early diverging lineages should be gathered to

further test our hypothesis about the ancestral timing of primordial germ cell

specification. Given the phylogenetic position of the Palaeoptera as basal branches of

the hexapod tree, data about primordial germ cell specification in these orders would be

especially useful to test our hypothesis with respect to hexapods, as would more

information from the ametabolous Archaeognatha and Zygentoma or the non-insect

hexapod Diplura. Similarly, data about primordial germ cell specification are missing

from Oligostraca, the most basally branching clade of Pancrustacea, which includes

Ostracoda and Branchiura.

Outside of Pancrustacea, the chelicerate phylogeny remains unstable (Sharma

and Gavish-Regev, 2025), and embryological data are too scattered for us to suggest

trends in primordial germ cell specification within this subphylum. Data regarding

primordial germ cell differentiation are even sparser for myriapods: we are aware of

relevant data from only five species across the four myriapod classes (Dohle, 1964;

Green and Akam, 2014; Heymons, 1901; Tiegs, 1940; Tiegs, 1947). Studies of

primordial germ cells in a broader range of both chelicerates and myriapods, as well as

detailed embryological studies in unexplored taxa, will therefore be needed to fully

understand the developmental context and evolutionary trends of germ cell specification

in these subphyla.

III. Molecular and genetic mechanisms of germ cell specification in

panarthropods

Beyond studying where and when germ cells first appear during arthropod

embryogenesis, researchers have sought to understand how they are specified. Germ

cell specification in animals has historically been classified into two mechanisms:

maternal inheritance and zygotic induction (Extavour and Akam, 2003; Nieuwkoop and

Sutasurya, 1981). In maternal inheritance, germ cell determinants accumulate in a

specific region of the oocyte cytoplasm, forming a germ plasm. Cells that inherit

sufficient germ plasm components after the embryo divides become primordial germ

cells. In embryos that undergo zygotic induction, all cells are in principle competent to

become germ cells at the beginning of embryogenesis. Certain cells later receive

signals from neighboring cells that induce them to take on germ cell identity. In the

following section, we will describe how historical observational data and perturbational

experimental approaches have offered increasingly robust evidence to distinguish

between these mechanisms in several panarthropod species.

III.i. Maternal inheritance

In the early twentieth century, researchers speculated that the electron-dense,

membraneless granules frequently found in the cytoplasm of primordial germ cells were

responsible for their specification. Experiments in chrysomelid beetles provided early

evidence that the posterior cytoplasm of the oocyte or early embryo contains germ cell

determinants. Burning or puncturing the posterior poles of beetle eggs with needles to

damage or remove posterior cytoplasm reduced or eliminated primordial germ cells

(Hegner, 1908; Hegner, 1911), demonstrating that this cytoplasm played a causal, and

nor simply correlative, role in primordial germ cell specification. Studies in *D.*

melanogaster offered further evidence for the necessity of germ plasm. Removing or

destroying the posterior cytoplasm of *D. melanogaster* eggs via UV irradiation

demonstrated this region's requirement for germ cell formation (Geigy, 1931).

Subsequent centrifugation experiments in *D. melanogaster* embryos revealed that

shifting the granular structures in posterior cytoplasm away from the posterior pole

prevented germ cell formation or led to germ cells forming in a different part of the

embryo (Jazdowska-Zagrodzińska, 1966).

Experiments transplanting specific regions of cytoplasm between *D.*

melanogaster embryos showed that the contents of the posterior cytoplasm were

sufficient to specify germ cells. Transplanting posterior cytoplasm from undamaged *D.*

melanogaster embryos into sterile ones in which the posterior pole had been damaged

with UV-irradiation rescued fertility, while transplanting anterior cytoplasm did not

(Okada et al., 1974). Further transplantation experiments demonstrated that posterior

pole plasm introduced into the anterior of another embryo caused formation of ectopic

pole cells that gave rise to functional germ cells when transplanted again to the

posterior of a third, sterile embryo (Illmensee and Mahowald, 1974). Similar

experiments transferring the posterior cytoplasm from late-stage oocytes into embryos

revealed that the oocyte pole plasm could specify functional primordial germ cells

(Illmensee et al., 1976). This result revealed that the necessary germ cell determinants

in *D. melanogaster* are deposited during oogenesis. These foundational studies

demonstrated that maternally derived cytoplasmic granules are both necessary and

sufficient for germ cell formation in some species.

Contemporary studies testing for maternal inheritance often ask whether known

germ line gene products accumulate during late oogenesis and early embryogenesis in

the region where primordial germ cells first arise. For example, antibody staining or *in*

situ hybridization experiments to detect germ line molecules like Vasa protein or *nanos*

mRNA have suggested the presence of germ plasm in mosquitoes (Juhn and James,

2006; Juhn et al., 2008), scuttle flies (Wotton et al., 2014), ants (Khila and Abouheif,

2008; Lynch et al., 2011; Rafiqi et al., 2020), wasps (Kharel et al., 2024; Lynch et al.,

2011), weevils (Lynch et al., 2011), aphids (Chang et al., 2006; Chang et al., 2007;

Chen et al., 2009; Lin et al., 2014), amphipods (Gupta and Extavour, 2013), and prawns

(Chen et al., 2021). Transcriptomic analyses of different portions of embryos and

oocytes (Carter et al., 2015; Quan and Lynch, 2016; Yoon et al., 2019) have also

revealed asymmetric localization of molecules associated with primordial germ cell

specification.

However, expression data showing asymmetric localization of putative germ

plasm components are not sufficient to demonstrate that these molecules function in

primordial germ cell specification. For this, functional genetic techniques are needed to

manipulate gene activity and test the effect on germ plasm assembly and germ line

development. The function of germ plasm components in panarthropods has been most

thoroughly studied in *D. melanogaster* (Lehmann, 2016; Mahowald, 2001; Trcek and

Lehmann, 2019). The membraneless ribonucleoprotein granules at the posterior of *D.*

melanogaster oocytes and early embryos are nucleated by the protein Oskar, which is

both necessary and sufficient for germ plasm assembly (Ephrussi and Lehmann, 1992;

Kistler et al., 2018). Oskar-mediated localization of core germ line molecules such as

Vasa, Tudor, Aubergine, and *nanos* to germ granules is essential for primordial germ

cell specification and maintenance of germ cell identity during cell migration and gonad

formation in these fruit flies (Lehmann, 2016). The roles of these germ plasm molecules

in establishing the *D. melanogaster* germ line have been extensively reviewed

elsewhere (Lehmann, 2016; Mahowald, 2001).

Analysis of germ plasm composition and localization in species beyond *D.*

melanogaster has revealed both conserved and rapidly evolving features. Within the

Drosophila genus, germ granule size, morphology (Counce, 1963; Mahowald, 1968;

Rivard et al., 2025), and the relative proportions of different granule components vary

widely (Doyle et al., 2023). The wasp *N. vitripennis* also has a well-characterized germ

plasm in the form of a single large structure called the oosome (Bull, 1982). Like

Drosophila, N. vitripennis requires *oskar* to assemble a germ plasm containing many of

the same molecules (e.g., Vasa, Tudor, and *nanos*) (Lynch and Desplan, 2010; Lynch

et al., 2011). However, while *D. melanogaster* has many small germ granules

(Mahowald, 1962), the *N. vitripennis* oosome has a distinct protein organization and

transcript composition compared to that observed in *D. melanogaster* (Kemph et al.,

2025; Kharel et al., 2024; Lynch et al., 2011). In several crustacean species, the germ

plasm appears as a large, round granule in the cortex of embryos and contains

conserved germ plasm markers (Chen et al., 2021; Grattan et al., 2013; Gupta and

Extavour, 2013). Studying germ plasm in additional species will be needed to clarify

whether and how variation in germ plasm makeup and organization influences the post-

transcriptional regulation and function of germ line determinants.

III.ii. Zygotic induction

Experimentally demonstrating inductive germ cell specification is more

technically challenging than detecting germ plasm accumulation. Consequently, most

studies have inferred or hypothesized zygotic induction based on absence of evidence

of germ plasm. For example, *in situ* hybridization experiments fail to reveal asymmetric

localization of putative germ line gene transcripts in oocytes and early embryos of

various species, including milkweed bugs (Ewen-Campen et al., 2013b), crickets

(Ewen-Campen et al., 2013a), honeybees (Dearden, 2006), mites (Dearden et al.,

2003), spiders (Schwager et al., 2015), centipedes (Green and Akam, 2014), and

tardigrades (Heikes et al., 2023). Instead, key germ line genes are expressed in a

subset of cells later in embryogenesis, coincident with reports of the first primordial

germ cells based on morphological criteria. However, we note that even in the well-

accepted case of mouse primordial germ cell induction (Saitou and Yamaji, 2010), it

remains formally possible that some unknown, and therefore untested, germ line

determinant, or factor that makes cells competent to respond to later germ line induction

signals, is asymmetrically localized and preferentially inherited by germ cells.

This caveat notwithstanding, there are multiple examples of experimental support

for induction rather than inheritance across panarthropods. Maternal knockdown of

putative germ plasm components in milkweed bugs and crickets reveals the

dispensability of maternal products of these genes for primordial germ cell formation

(Ewen-Campen et al., 2013b; Ewen-Campen et al., 2013a). Laser ablation of single

cells in two and four cell stage embryos of the tardigrade *Thuliana stephaniae* does not

prevent primordial germ cell specification, also suggesting that this species lacks a

cytoplasmic region with inherited germ line determinants (Hejnol and Schnabel, 2005).

Observations of germ cell formation in embryos with tissue duplications can be

interpreted as evidence against germ plasm and consistent with induction. For example,

in the drain fly *Clogmia albipunctata* and the flour beetle *T. castaneum*, genetic

manipulations producing double abdomen embryos lead to germ cell formation at both

anterior and posterior ends (Ansari et al., 2018; Yoon et al., 2019). Similarly, naturally

occurring "twin" embryos of the stick insect *Carausius morosus* results in duplicated

embryos fused along their anterior-posterior axis, that possess germ cells(Cavallin,

1971). In contrast, double abdomen embryos of *D. melanogaster* only produce germ

cells at the true posterior, since germ plasm is absent at the ectopic posterior end

(Lasko and Ashburner, 1990).

Among panarthropods, experiments in the cricket *G. bimaculatus* have provided

the most direct evidence for zygotic induction of primordial germ cells. Knockdown of

the mesoderm specification factor *twist* led to loss of primordial germ cells, suggesting

that mesoderm contributes to germ cell specification and/or maintenance (Ewen-

Campen et al., 2013a) and consistent with morphological evidence that orthopteran

primordial germ cells first arise from the embryonic mesoderm (Wheeler, 1893). The

BMP signaling pathway, known to induce germ cells in mice (Hopf et al., 2011; Lawson

et al., 1999; Ying and Zhao, 2001; Ying et al., 2000), is also required for cricket germ

cell specification. Specifically, BMP signaling and the downstream transcription factor

Blimp-1 induce a subset of mesodermal cells to become primordial germ cells in

crickets (Donoughe et al., 2014; Nakamura and Extavour, 2016).

The shared use of both BMP and *Blimp-1* in mice and crickets may have

convergently evolved, or could suggest that this signaling mechanism is ancestral in

bilaterian inductive primordial germ cell specification (Lochab and Extavour, 2017).

Determining whether BMP and its downstream target genes are required for germ line

specification in more species will be necessary to distinguish between these

possibilities. Surprisingly few signaling pathways contribute to developmental fate

decisions in animals (Pires-daSilva and Sommer, 2003), increasing the likelihood of

evolutionary convergence, and limiting the number of candidate signaling pathways to

test for roles in primordial germ cell specification in different species.

III.iii. Intermediate and flexible strategies of germ cell specification

Evidence from some animal species suggests that there may not be a strict

dichotomy between inheritance and induction (discussed by Seervai and Wessel, 2013).

These mechanisms may instead represent two ends of a spectrum, and they may not

be mutually exclusive. Below we discuss cases of panarthropod species for which data

suggests that their mechanisms of germ cell specification may fall somewhere along

this continuum.

In the silk moth *B. mori*, historical studies have generated conflicting evidence

about how germ cells are specified. Germ granules were not detected in early light

microscopy experiments, but germ cells failed to form after cauterization of the ventral

side of cleavage stage embryos, suggesting this region could contain germ cell

determinants (Miya, 1958a). Subsequent studies found that Vasa protein is not

asymmetrically localized before cellularization, again suggesting a lack of germ plasm

(Nakao et al., 2006). However, the mRNAs of one of four *B. mori nanos* homologs,

nanosO, localizes along the ventral midline of syncytial embryos immediately following

egg-laying, and is inherited by the primordial germ cells (Nakao et al., 2008).

Intriguingly, researchers found that maternal and zygotic *nanosO* act redundantly in

primordial germ cell specification; loss of one or the other reduces but does not

completely eliminate germ cells, whereas loss of both prevents germ cell formation

(Nakao and Takasu, 2019). The possibility that both inheritance and induction

mechanisms contribute to *B. mori* primordial germ cell specification suggests that this

species could offer insight into transitions in germ cell specification mechanisms within

Lepidoptera or in insects more broadly.

Although *D. melanogaster* is traditionally characterized as a species using a

germ plasm, zygotic signaling was recently suggested to contribute to germ cell

specification in this species (Colonnetta et al., 2022). Loss-of-function mutations in the

BMP family morphogen Dpp lead to partial loss of transcriptional quiescence and

mislocalization of germ plasm in the newly formed pole cells (Colonnetta et al., 2022).

Furthermore, loss of BMP signaling leads to activation of the normally suppressed

terminal signaling pathway in pole cells, which in turn disrupts their migration

(Colonnetta et al., 2022). These observations illustrate a universal conceptual challenge

in the field of developmental biology: distinguishing between, or even consistently

defining, cell type specification and maintenance. Although pole cells can form upon

disruption of BMP signaling, they do not contain the proper molecular components to

act as primordial germ cells. We suggest it may be most useful to define primordial

germ cell specification mechanisms as those that ensure both the physical formation of

germ cell precursors (including adoption of specific morphologies and activation of

specific gene regulatory states), and the competence of those cells to take the next

developmental steps toward joining the gonad. Under this definition, we would indeed

have to concede that both germ plasm and zygotic inductive mechanisms play roles in

D. melanogaster germ cell specification.

The ability to post-embryonically regenerate a germ line initially specified via

maternal inheritance has also been reported in multiple invertebrate species, including

the sea squirt *Ciona intestinalis* (Takamura et al., 2002), the earthworm *Enchytraeus*

japonensis (Tadokoro et al., 2006), and the marine worm *Capitella teleta* (Dannenberg

and Seaver, 2018) (see also Özpolat, 2024 for a review of this phenomenon in

Annelida)). The wasp *Pimpla turionellae* (Hymenoptera) has an oosome required for

germ cell specification: damaging this material results in no pole cells, which are the

primordial germ cells of this species (Achtelig and Krause, 1971). However, oosome-

deficient embryos lacking pole cells develop into larvae with germ cells in their gonad,

indicating that an additional mechanism must regenerate germ cells later in

development. Evidence of germ line regeneration has also been reported in the

amphipod *Parhyale hawaiensis*, which has a germ plasm that is asymmetrically

segregated to the *g* micromere and its lineage (Extavour, 2005; Gerberding et al., 2002;

Gupta and Extavour, 2013). Ablation of the *g* micromere eliminates primordial germ

cells in the embryo (Extavour, 2005), but the germ line is replaced post-embryonically

with germ cells that appear to derive from the mesoderm (Kaczmarczyk, 2014). The

authors of this study speculated that these replacement germ cells specifically derive

from cells of the somatic gonad(Kaczmarczyk, 2014) , reminiscent of a theory popular

for the first half of the 20th century, that primordial germ cells in most animals derived

from the inner epithelium of the somatic gonad (Everett, 1944). Species capable of

regenerating primordial germ cells present the opportunity to investigate the relative

contributions of and potential interactions between inheritance- and induction-based

mechanisms in a single species. Further, these species may represent living examples

of the intermediates that are hypothesized to exist in evolutionary transitions between

modes of germ line specification (Extavour, 2007).

III.iv. Summary: convergent evolution of panarthopod germ plasm

The phylogenetic distribution of germ cell specification mechanisms across

panarthropods reveals a complex evolutionary history. Data for tardigrades,

chelicerates, and myriapods are consistent with induction as the ancestral mechanism

of panarthropod primordial germ cell specification. Pancrustacea are the most heavily

represented group in our dataset. The late origin of primordial germ cells in several

species suggests induction may be widely used, but there appear to be independent

origins of germ plasm in Malacostraca and Branchiopoda. The use of induction and

germ plasm in different decapod species suggests that further sampling could reveal

patterns in the evolution of germ cell formation within this order (Biffis et al., 2009; Chen

et al., 2021; Terao, 1929). However, functional experiments and broader taxonomic

sampling will be necessary to confirm mechanisms and reconstruct their evolutionary

trajectories across Panarthropoda as a whole.

Reminiscent of the broader pattern we previously reported across animals

(Extavour, 2007; Extavour and Akam, 2003; Whittle and Extavour, 2017), in some insect

orders we noted that early diverging lineages deployed induction, while later lineages

appeared to use germ plasm. This pattern is well exemplified in Coleoptera, (Kobayashi

et al., 2006; Komatsu and Kobayashi, 2012; Krause, 1947; Niikura et al., 2017;

Schröder, 2006; Ullmann, 1964), with pole cells forming only in Chrysomeloidea and

Curculionidae (Brauer, 1925; Hegner, 1908; Lynch et al., 2011). In addition, in

Carabidae, a basally branching beetle family that lacks pole cells, *oskar* orthologs have

been identified, decoupling the presence of *oskar* from germ plasm in beetles (Blondel

et al., 2021; Kobayashi et al., 2013). These observations raise the possibility that germ

plasm may not be ancestral to Coleoptera. When considered alongside the absence of

germ plasm in most Lepidoptera and the observation of late-arising, Category 3 germ

cells in Trichoptera (Miyakawa, 1974) and Category 2 germ cells in Mecoptera (Suzuki,

1990), these patterns are consistent with a novel hypothesis in which germ plasm

evolved independently multiple times across insects, rather than only once in the

Holometabola as we and others have previously proposed (Ewen-Campen et al., 2012;

Ewen-Campen et al., 2013a; Lynch et al., 2011). Our previous work had suggested the

independent evolution of germ plasm at the phylum level (Extavour and Akam, 2003),

but this new hypothesis is the first, to our knowledge, to suggest such widespread

convergent evolution of maternal inheritance-based germ cell specification within a

phylum.

To test this hypothesis, we performed ancestral character reconstruction using

both maximum parsimony and maximum likelihood approaches. For this analysis, we

designated species as utilizing inheritance or induction based on experimental evidence

when available or inferred from observational data describing the timing and location of

germ cell specification (Table S1). The results of the parsimony analysis inferred that

the ancestral panarthropod specification mode was induction, with a minimum of 24

changes over the history of the panarthropods (Figure S1; Table S3). When considering

all equally parsimonious character histories, changes from induction to maternal

inheritance (min = 18, max = 23, average = 20.5) were the most common, but reverse

transitions were also inferred (min = 2, max = 7, average = 4.5). At a minimum,

reversions from induction to maternal inheritance were inferred in the honeybee *A.*

mellifera (Hymenoptera) and drain fly *C. albipunctata* (Diptera). We note that all

maximum parsimony reconstructions assign induction as the ancestral state of

Holometabola (node 11), which requires many fewer total transitions within

Holometabola (N = 12) than would be required if germ plasm evolved a single time in

the last common ancestor of Holometabola (N = 20).

The marginal ancestral reconstructions of our maximum likelihood approach are

consistent with the parsimony analysis described above (see Table S3 for a

comparison), again providing strong statistical support for induction as the ancestral

specification mechanism in Panarthropoda, Arthropoda, and Insecta, with multiple

independent instances of germ plasm evolution and reversion to induction (Figure S2;

see Table S7 for marginal probabilities). We further find strong evidence that induction

was the ancestral specification mechanism in Holometabola (probability 80.6%),

followed by independent transitions to germ plasm in the common ancestors of Diptera

and Mecoptera (or independently within Mecoptera, Siphonaptera, and Diptera), derived

lineages of Lepidoptera, in the common ancestor of the families Chrysomelidae and

Curculionidae (within Coleoptera), and in the common ancestor of Hymenoptera.

Outside of Holometabola, we infer independent transitions to germ plasm in thrips,

aphids, earwigs, Collembola (which may have undergone multiple independent

transitions within the class), Decapoda, the last common ancestor of Hyalidae and

Talitridae (order Amphipoda), and in the chelicerate *Tetranychus urticae*.

IV. Evolution of panarthropod germ cell specification

In the following section, we discuss the developmental and genetic factors that might

contribute to evolutionary transitions in germ cell specification across Panarthropoda.

IV.i. The relationship between germ cells and somatic germ layers

In arthropods with late formation of primordial germ cells (Category 3), primordial

germ cells appear to differentiate directly from the mesoderm (Evans, 1901; Hickman,

1936; Nelson, 1915). Consistent with this hypothesis, abolishing mesoderm prevents

germ cell specification in the cricket *G. bimaculatus* (Ewen-Campen et al., 2013a).

Germ cell differentiation occurs concurrently with the differentiation of many other

mesodermal cell types, including the somatic gonad, musculature, heart, and fat body.

We suggest that the types of mechanisms that specify germ cells are not distinct in

principle from those specifying any other mesodermal cell type in these animals.

The germ layer origin of germ cells is less clear in arthropods whose germ cells

appear before or very close to the time of gastrulation (Category 2). For example,

primordial germ cells in many crustaceans are described as being the first to enter the

blastopore during gastrulation (e.g. Cannon, 1921; Goodrich, 1939b; Manton, 1928),

and usually considered by the authors to be derived from the mesendoderm. In

contrast, germ cells in Category 2 insects arise before gastrulation (Mellanby, 1935b;

Ullmann, 1964), consistent with the hypothesis that germ cells are distinct from the

mesoderm and may not derive from any germ layer.

In crustaceans in which a putative germ plasm has been identified, the cleavage

pattern is stereotypical, and cell lineages are available, the germ cell lineage is sister to

the mesoderm or the mesendoderm [Amphipoda (Extavour, 2005; Gerberding et al.,

2002; Scholtz and Wolff, 2002; Wolff and Scholtz, 2002), Euphausiacea (Alwes and

Scholtz, 2014), Decapoda (Hertzler, 2005)], but never to the ectoderm. In addition,

removing the putative germ plasm from embryos of the amphipod *P. hawaiensis* makes

the *g* micromere morphologically indistinguishable from the other micromeres, which

give rise to mesoderm or endoderm in unperturbed embryos (Gupta and Extavour,

2013). This result suggests that in the absence of germ cell determinants, the *g*

micromere, normally the germ line progenitor, could instead adopt mesendodermal fate.

This hypothesis has not been directly tested in this amphipod because gastrulation fails

after ablation of the putative germ plasm (Gupta and Extavour, 2013). Together, these

data suggest a close association between the germ cell and mesendodermal lineages in

these crustaceans. This relationship is apparent beyond Pancrustacea, since germ cells

induced late in embryogenesis arise from the mesoderm (Category 3). We thus

speculate that germ cells in panarthropods may be ancestrally derived from the

mesodermal lineage. We further speculate that inductive germ cell specification is

ancestral to panarthropods, and present a formal test of this hypothesis below.

IV.ii. Evolutionary shifts in the spatial and temporal origin of primordial germ cells

We have seen significant variation in the spatial and temporal origin of primordial

germ cells across panarthropods (Figures 2-3; Table S1). However, regardless of when

primordial germ cells first appear, we find a common stage across many panarthropods

in which primordial germ cells are situated in bilateral segmental clusters across

multiple trunk segments. Soon after this stage, the segmental clusters on each side fuse

together into bilaterally symmetrical genital ridges, corresponding to the primordial

embryonic gonad (Miya, 1958b; Nelsen, 1934; Seidel, 1924; Ullmann, 1964).

Most arthropods undergo "short germ" development, wherein the blastoderm

tissue generates only anterior segments, with posterior segments being added

sequentially to the posterior end of the germ band later in development. In contrast, the

"long germ" mode of development seen in *D. melanogaster*, wherein the blastoderm

tissue immediately generates all body segments, is an evolutionary novelty found only

in some insects (Clark et al., 2019; Krause, 1939). In Category 1 or 2 arthropods with

short germ development, germ cells are often specified before the formation of the

segments that will include the future embryonic gonad. In Category 2 insects, germ cells

often associate with the posterior end of the germ band and then migrate toward the

anterior as the germ band elongates (Goss, 1952b; Hegner, 1909b; Heming and

Huebner, 1994; Paz, 1958; Suzuki, 1990; Ullmann, 1964). This migratory path is so

common among insects that Johanssen and Butt included it in their description of

embryogenesis in a stereotypical insect (Johannsen and Butt, 1941).

The association of germ cells with the posterior end of the embryo may have had

other evolutionary consequences in insects. Many insects, both long and short germ,

specify the body plan early in development via maternal deposition of axial patterning

determinants, independent of the mechanism of germ cell specification (reviewed in

Lynch, 2019; Rosenberg et al., 2009). For example, the short germ grasshopper

Schistocerca americana exhibits asymmetric localization of the posterior patterning

determinant *nanos* in the absence of a germ plasm (Lall et al., 2003). However, insects

with germ plasm that includes axial patterning determinants, like *D. melanogaster*, often

exhibit long germ development. Based on these observations, we employed

phylogenetic logistic regression and Pagel's test of evolutionary rate covariation (Pagel,

1994) to test the hypothesis that the acquisition of long germ band development

influences the evolutionary rate of germ cell mechanism transitions or vice versa, and

found statistically significant evidence of this association using both methods (Figure

S3; Figure S4). All Pagel covariation models assuming coupled evolutionary rates

among these two characters are significantly better fits to the data than the null model of

independent evolution (likelihood ratio test p-value < 1e−3; Figure S4; Table S8). We

note that the bidirectional and GB-dependent models perform similarly to the GC-

dependent model, suggesting both that long germ band development may drive the

evolution of germ plasm, and that germ plasm may drive transitions to long germ band

development.

We further hypothesize that a spatial shift in germ cell specification to the

posterior end of the embryo in Category 2 insects created a condition in which germ line

determinants are localized to the same embryonic region as posterior patterning

determinants (Figure 4B). Subsequently, germ line determinants may have hitchhiked

onto the asymmetric localization machinery, coupling germ cell specification to posterior

patterning and thus promoting the evolution of germ plasm (Figure 4C). Under this

hypothesis, the germ plasm of *D. melanogaster* would represent the final stage of this

evolutionary scenario, as its germ granules include both germ line and abdominal

patterning determinants (Ephrussi and Lehmann, 1992; Trcek et al., 2015; Wang and

Lehmann, 1991).

Nasonia vitripennis is an intriguing case to study the relationship between

anteroposterior patterning and germ cell specification. The posterior cytoplasm of this

wasp has two populations of *nanos*: one that is incorporated into the oosome and

eventually into pole cells, and another that functions in axial patterning (Lynch and

Desplan, 2010). *nanos* localization to the oocyte posterior in this wasp depends on

actin-dependent anchoring (Olesnicky and Desplan, 2007), just as in *D. melanogaster*

(Forrest and Gavis, 2003), suggesting a conserved localization mechanism between

these two species. We speculate that *N. vitripennis* represents an evolutionary

intermediate between species where anteroposterior patterning is entirely independent

of germ cell formation (Figure 4A) and those where these two processes are fully linked

(Figure 4C). Our evolutionary hypothesis can be tested further by examining the

mechanisms of localization of posterior patterning determinants and germ line

determinants in more insect species to understand how existing cytoskeletal machinery

for one developmental process may have been coopted for use in another.

IV.iii. Potential gene regulatory basis of evolutionary shifts in the molecular

mechanism of primordial germ cell specification

Beyond the timing and position of germ cell origin, evolutionary changes in the

localization of gene products and timing of gene expression may also underlie

transitions in the mechanism of specification (Extavour, 2007; Srouji and Extavour,

2010). Many of the genes important for specifying germ cells are conserved across

metazoans (Ewen-Campen et al., 2010), but they exhibit distinct temporal and spatial

expression patterns across species. In species that use inheritance, the germ line gene

products are expressed during oogenesis and persist in the early embryonic cytoplasm

to be inherited by the primordial germ cells. In species that induce germ cells, these

genes are either not expressed in oocytes or the gene products are not asymmetrically

localized so they are not preferentially inherited by particular cells. As we have

suggested previously (Extavour, 2007; Srouji and Extavour, 2010), relatively simple

regulatory modifications affecting post-transcriptional regulation, the cytoskeleton, or

RNA or protein localization could cause germ line gene product retention in oocytes,

thereby enabling the transition from induction to inheritance.

The insect-specific gene *oskar* offers a potential example of gene network

evolution driving shifts in germ cell specification mechanism. Since *oskar* is likely a

nucleator of germ plasm in multiple insects (Kemph and Lynch, 2022; Lynch et al.,

2011) and has been lost multiple times since its origin prior to the divergence of

Pterogyta from Ametabola (Blondel et al., 2021), we hypothesized that *oskar* gene

presence might be correlated with germ line specification mode in a given taxon (Figure

S3). A related hypothesis, that the evolutionary advent of *oskar* gave rise to

holometabolous germ plasm, was previously proposed based on the co-occurrence of

oskar transcript and germ plasm in *N. vitripennis* and *D. melanogaster*, as well as the

absence of both *oskar* homologs and f germ plasm in *T. castaneum*, *A. mellifera*, *C.*

albipunctata, and most Lepidoptera (Lynch et al., 2011). The results of applying

phylogenetic logistic regression to this hypothesis (Figure S3; Table S9) suggest a

strong and statistically significant association between *oskar* presence and germ-plasm

mode (coefficient = 2.5827, adjusted odds-ratio: = 13.28, p-value = 0.00059) and no

association of germ-plasm mode with BUSCO completeness (coefficient = 0.228, p-

value = 0.368). The association between *oskar* presence and germ-plasm mode

remains strong and statistically significant even after removing 24 within-family

substitutions made when neither genomic nor transcriptomic data were available for a

species (coefficient = 2.56, adjusted odds ratio = 12.929, p-value = 0.00026). We

emphasize that this observed association does not imply that *oskar* is necessary or

sufficient for germ plasm formation; indeed, there are insect species without *oskar* that

nevertheless have germ plasm and species with *oskar* that lack germ plasm (Figure

S3), as we discuss further below.

In contrast to prior hypotheses from our group and others of a single origin of

germ plasm at the base of the Holometabola (Ewen-Campen et al., 2012; Ewen-

Campen et al., 2013a; Kemph and Lynch, 2022; Lynch et al., 2011), our current

likelihood and parsimony reconstruction of induction as ancestral to Holometabola

(Figure 5; Figures S1-S2) implies that *oskar* has been independently recruited as a

germ plasm nucleator at least twice in Hymenoptera (Lynch et al., 2011) and Diptera

(Ephrussi and Lehmann, 1992). The strong association between *oskar* presence and

germ plasm (Figure S3) implies that it may have been co-opted for germ plasm

nucleation even more times across insects. We note that such "gene reuse" is a

common feature of convergent evolution across multiple biological processes (Chan et

al., 2010; Martin and Orgogozo, 2013; Morris, 2009; Reed et al., 2011; Satterlee et al.,

2024). Since *oskar* is co-expressed with germ plasm components in nervous systems of

hemimetabolous and holometabolous insects (Ewen-Campen et al., 2012; Kulkarni et

al., 2023; Xu et al., 2013), we posit that independent acquisitions of germ plasm in

distinct lineages may have proceeded via parallel cooption of a pre-existing gene

interaction network. Future studies should aim to determine whether each independent

origin of germ plasm in the Holometabola arose by co-option of *oskar* to the germ line,

or by *oskar*-independent mechanisms of germ plasm assembly.

The results of our analysis (Figure S3; Table S9) suggest that *oskar*'s functional

evolution may have contributed to a transition from induction to inheritance, but

precisely when and how *oskar* acquired a germ line role remains unclear. Morphological

data suggest that thrips use germ plasm (Haga, 1985; Heming, 1979), and *oskar*

homologs have been found in this order (Blondel et al., 2021). If *oskar* plays a role in

thrips germ cell specification, it could support the entirely novel hypothesis that *oskar*

evolved its germ line determinant properties before the evolution of insect

metamorphosis.

IV.v. The case for novel panarthropod germ plasm nucleators

Recruitment of *oskar* to the germ line is not the only mechanism explaining the

evolution of germ plasm in arthropods. The hemimetabolous pea aphid *Acyrthosiphon*

pisum lacks *oskar* but has a germ plasm containing Vasa and Nanos (Chang et al.,

2006; Chen et al., 2009; Lin et al., 2014), which are asymmetrically localized by

unknown mechanisms. Vasa is also asymmetrically localized to a putative germ plasm

in amphipod and decapod species (Chen et al., 2021; Gupta and Extavour, 2013),

which must also have evolved via distinct mechanisms since *oskar* is an insect-specific

gene (Blondel et al., 2021). Germ plasm nucleators have convergently evolved in

diverse animal phyla (Aoki et al., 2016; Bontems et al., 2009; Kulkarni and Extavour,

2017; Lee et al., 2020; Marlow and Mullins, 2008; Scholl et al., 2024). Similarly, we posit

that germ plasm nucleators other than *oskar* must also exist across the panarthropods,

awaiting discovery.

V. Concluding thoughts

By reinterpreting the meticulous work of generations of classical embryologists

through the lens of modern developmental genetics and an improved panarthropod

phylogeny, we have generated novel, testable evolutionary hypotheses about the

mechanisms that specify the germ line, with implications for the genetic basis of

convergence in evolution that extend beyond this superphylum. The advent of new

genetic tools in an increasing number of panarthropod model species place us in an

exciting era to tackle age-old questions about the evolution of panarthropod germ cell

specification, as well as the impact of these mechanisms of genome evolution and body

patterning in the most successful group of animals on earth.

Acknowledgements

We thank members of the Extavour Lab for helpful discussions, Kira Heikes for the image of tardigrade germ cells (Figure 1B4), and Prashant Sharma for the image of harvestmen germ cells (Figure 1B3).

Data Availability

All data sources are listed in the accompanying Supplementary Material and/or are available at

https://github.com/rishabhrajkapoor/panarthropoda_gc_specification_evolution, commit ID 25384e7.

Author Contributions

All authors gathered data and performed data analysis and interpretation. JAK, ELR and RRK wrote the first draft of the manuscript. CGE conceived of the study, obtained funding for the study, supervised its execution, and reviewed and edited the manuscript.

Funding

This study was supported by NSF award IOS-2220747, funds from Harvard University, and from the Howard Hughes Medical Institute awarded to CGE. Additional support was provided by the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number #1764269), the Harvard Quantitative Biology Initiative, and a Herchel Smith Graduate Fellowship awarded to ELR, and by an NSF Graduate Research Training Fellowship to RRK. CGE is an Investigator of the Howard Hughes Medical Institute.

Conflicts of Interest

The authors declare no conflicts of interests.

A. Category 1

A1 *Drosophila melanogaster* (Diptera)

A2 *Parhyale hawaiensis* (Amphipoda)

B. Category 2

B1 *Oncopeltus fasciatus* (Hemiptera)

B2 *Limnoria lignorum* (Isopoda)

B3 *Phalangium opilio* (Opiliones)

B4 *Hypsibius exemplaris* (Tardigrada)

B5 *Peripatopsis moseleyi* (Onychophora)

C. Category 3

C1 *Gryllus bimaculatus* (Orthoptera)

C2 *Nebalia bipes* (Leptostraca)

C3 *Parasteatoda tepidariorum* (Araneae)

Figure 1 (previous page). Primordial germ cell origin in the context of panarthropod
embryogenesis. In all schematics, magenta represents the approximate location of germ plasm or
primordial germ cells; light gray represents yolk or extraembryonic tissue.
(**A1**) Schematic: In the fruit fly *Drosophila melanogaster*, germ plasm accumulates at the posterior of
syncytial embryos and is inherited by pole cells that migrate to the ultimate site of the embryonic gonad
(Illmensee and Mahowald, 1974). Data: TEM micrograph displays pole cells budding in *D. melanogaster*
embryos. The pole cell nuclei (N) have characteristically diffuse chromatin, and polar granules (P) are
distributed throughout the cytoplasm. Image reproduced with permission from (Mahowald, 1968).
(**A2**) Schematic: In the amphipod *Parhyale hawaiensis*, germ plasm accumulates in a region of the
embryo called the RNA-containing body, which is inherited by the *g* micromere in the 8-cell stage
(Extavour, 2005; Gupta and Extavour, 2013). All descendants of the *g* micromere are primordial germ
cells. Data: Primordial germ cells can be visualized by immunohistochemistry with a cross-reactive
antibody raised against recombinant Vasa protein from both *D. melanogaster* and the grasshopper
Schistocerca americana (Chang et al., 2002).
(**B1**) Schematic: In the milkweed bug *Oncopeltus fasciatus*, primordial germ cells first appear in the
ventral posterior blastoderm in an invaginating structure called the posterior pit (Ewen-Campen et al.,
2013b). Primordial germ cells remain for some time at the posterior tip of the germ band as it invaginates
into the yolk. The germ cells migrate along the germ band toward the anterior and settle in segments A4–
A6. Data: Light micrograph of sectioned *O. fasciatus* embryos shows germ cells in the posterior pit in the
late blastoderm stage. Image reproduced with permission from (Butt, 1949b).
(**B2**) Schematic: In the isopod *Limnoria lignorum*, primordial germ cells are first identifiable among the
mesendodermal mass just after the beginning of gastrulation (Strömberg, 1968). Primordial germ cells
remain associated with the mesoderm throughout development. Data: Light micrograph of a sectioned *L.*
lignorum embryo shows primordial germ cells among the mesoderm in the extended germ band stage,
distinguished by their lighter-staining cytoplasm and larger nuclei. Image reproduced with permission from
(Strömberg, 1968).
(**B3**) Schematic: In the harvestman *Phalangium opilio*, primordial germ cells are first identifiable at the
posterior end of the germ band soon after the germ band forms (Faussek, 1888; Gainett et al., 2022).
Primordial germ cells remain at the posterior end of the germ band at least until the fourth opisthosomal
segment is formed, after which the germ cells can be found in that segment. Data: Primordial germ cells
can be identified by staining with the nucleic acid dye Hoechst 33342; germ cell chromatin is more diffuse
compared with that of surrounding cells. Image kindly provided by Prashant Sharma (University of
Wisconsin, Madison).
(**B4**) Schematic: In the tardigrade *Hypsibius exemplaris*, primordial germ cells are first identifiable in the
32-cell stage and are the first cells to be internalized during gastrulation. Data: Primordial germ cells can
be visualized by *in situ* hybridization against *He-vasa*. Image kindly provided by Kira Heikes (Duke
University).
(**B5**) Schematic: In the velvet worm *Peripatopsis moseleyi*, primordial germ cells are first identifiable near
the blastopore at the beginning of gastrulation. As the embryo elongates and more segments form, the
primordial germ cells migrate to more anterior segments. Data: Light micrograph of a *P. moseleyi* embryo
shows primordial germ cells soon after their internalization during gastrulation, distinguished by their
lighter-staining nuclei and cytoplasm. Image reproduced with permission from (Manton, 1949).
(**C1**) Schematic: In the cricket *Gryllus bimaculatus*, primordial germ cells are first detectable among the
mesoderm in segments A2–A4 in embryonic stage 6, after most abdominal segments have been
specified (Ewen-Campen et al., 2013a). Data: Primordial germ cells can be visualized by
immunohistochemistry with an antibody against *Gb*-Vasa.
(**C2**) Schematic: In the leptostracan *Nebalia bipes*, primordial germ cells are first identifiable in the
mesoderm in the thorax and abdomen during the extended germ band stage (Manton, 1934). Data: Light
micrograph of a sectioned *N. bipes* embryo shows primordial germ cells among the mesoderm in the
extended germ band, distinguished by their lighter-staining cytoplasm and larger nuclei. Image
reproduced with permission from (Manton, 1934).
(**C3**) Schematic: In the house spider *Parasteatoda tepidariorum*, primordial germ cells are first identifiable
in segmental clusters in the opisthosomal mesoderm after all segments have formed (Schwager et al.,
2015). Data: Primordial germ cells can be visualized by immunohistochemistry using an antibody raised
against *Pt*-Vasa.

Clade

Collembola
Ametabola *Insecta*
Hemimetabola
Holometabola

Timing category

- Category 1
- Category 2
- Category 3

Evidence of mechanism

- Germ plasm (exp.)
- Germ plasm (obs.)
- Induction (exp.)

Figure 2 (previous page). Germ cell specification across Hexapoda. Phylogeny of studied hexapod
lineages marking the origin category and the mechanism of primordial germ cell specification.
Phylogenetic relationships in the cladogram are derived from published relationships described in the
Supplementary Material (Misof et al., 2014)and branch lengths are not to scale. Data on the timing of
germ cell specification in the first column. Dark purple = Category 1 (cleavage/blastoderm), medium
purple = Category 2 (gastrulation), light purple = Category 3 (mesoderm differentiation). Data about the
mechanism of germ cell specification are found in the second column. Dark pink = experimental evidence
supports the use of germ plasm, light pink = morphological evidence of germ plasm, blue = experimental
evidence of induction. Data are described in more detail in Table S1.

Figure 3 (previous page). Panarthropod germ cell specification outside of Hexapoda. Phylogeny of
studied non-hexapod panarthropod lineages marking the origin category and mechanism of primordial
germ cell specification. Phylogenetic relationships are derived from published relationships described in
the Supplementary Material (Misof et al., 2014)and branch lengths are not to scale. Colors are the
same as in Figure 2. Data are described in more detail in Table S1.

A AP patterning and germ cell determination are spatially and temporally separate

B AP patterning and germ cell determinants overlap spatially and temporally

C germ plasm controls both AP patterning and germ cell determination

938
939
Figure 4. Hypothesized links between spatial and temporal dynamics of germ cell origination and the mechanism of germ cell
specification. A) In species that induce germ cells later in embryogenesis (Category 3), the asymmetrically localized anterior–posterior (AP)
patterning determinants (blue) and germ cell determination factors (pink) are spatially and temporally separate. B) The shift to earlier germ cell
specification at the embryonic posterior in Category 2 insects enabled overlap between the patterning and germ cell determinants. C) Subsequent
hitchhiking of the asymmetric localization machinery by germ line determinants could couple germ cell specification and posterior patterning and
thus promote the evolution of germ plasm.

946
Figure 5. Evolutionary dynamics of panarthropod germ cell specification. Phylogenetic relationships
among all study species are derived from published relationships described in the Supplementary
Material, and branch lengths are not to scale. Boxes to the right of the cladogram designate the inferred
mechanism of germ cell specification (pink = germ plasm, blue = induction). Pink asterisks denote
instances of germ plasm evolution and pink pluses mark ambiguous cases of germ plasm gain. Blue
asterisks denote instances of germ plasm loss and blue pluses mark ambiguous cases of germ plasm
loss.
954

References

Achtelig, M. and Krause, G. (1971). Experimente am ungefurchten Ei von Pimpla turionellae L. (Hymenoptera) zur Funktionsanalyse des Oosombereichs. *Wilhelm Roux Arch. Dev. Biol.* **167**, 164–182.

Alexeeva, N. and Tamberg, Y. (2021). Anatomical changes in postembryonic development of Pycnogonum litorale. *J. Morphol.* **282**, 329–354.

Alexeeva, N., Tamberg, Y. and Shunatova, N. (2018). Postembryonic development of pycnogonids: A deeper look inside. *Arthropod Struct. Dev.* **47**, 299–317.

Alwes, F. and Scholtz, G. (2014). The early development of the onychopod cladoceran Bythotrephes longimanus (Crustacea, Branchiopoda). *Front. Zool.* **11**, 10.

Anderson, D. T. (1962a). The embryology of Dacus tryoni (Frogg.) [Diptera, Trypetidae (= Tephritidae)], the Queensland fruit-fly. *Development* **10**, 248–292.

Anderson, D. T. (1962b). The epigenetics of the larva in Diptera. *Acta Zoöl.* **43**, 221–228.

Anderson, D. T. (1967). Larval development and segment formation in the branchiopod crustaceans Limnadia stanleyana King (Conchostraca) and Artemia salina (L.) (Anostraca). *Aust. J. Zool.* **15**, 47–91.

Anderson, D. T. (1973). *Embryology and phylogeny in annelids and arthropods*. Oxford, UK: Pergamon Press.

Ando, H. and Tanaka, M. (1980). Early embryonic development of the primitive moths, Endoclyta signifier Walker and E. excrescens Butler (Lepidoptera: Hepialidae). *International Journal of Insect Morphology and Embryology* **9**, 67–77.

Ansari, S., Troelenberg, N., Dao, V. A., Richter, T., Bucher, G. and Klingler, M. (2018). Double abdomen in a short-germ insect: Zygotic control of axis formation revealed in the beetle Tribolium castaneum. *Proc. Natl. Acad. Sci.* **115**, 1819–1824.

Aoki, S. T., Kershner, A. M., Bingman, C. A., Wickens, M. and Kimble, J. (2016). PGL germ granule assembly protein is a base-specific, single-stranded RNase. *Proc Natl Acad Sci U S A* **113**, 201524400.

Ballesteros, J. A., Santibáñez-López, C. E., Baker, C. M., Benavides, L. R., Cunha, T. J., Gainett, G., Ontano, A. Z., Setton, E. V. W., Arango, C. P., Gavish-Regev, E., et al. (2022). Comprehensive Species Sampling and Sophisticated Algorithmic Approaches Refute the Monophyly of Arachnida. *Mol Biol Evol* **39**, msac021.

Biffis, C., Alwes, F. and Scholtz, G. (2009). Cleavage and gastrulation of the dendrobranchiate shrimp Penaeus monodon (Crustacea, Malacostraca, Decapoda). *Arthropod Struct. Dev.* **38**, 527–540.

Blondel, L., Besse, S., Rivard, E. L., Ylla, G. and Extavour, C. G. (2021). Evolution of a cytoplasmic determinant: Evidence for the biochemical basis of functional evolution of the novel germ line regulator Oskar. *Mol. Biol. Evol.* **38**, 5491–5513.

Bontems, F., Stein, A., Marlow, F., Lyautey, J., Gupta, T., Mullins, M. C. and Dosch, R. (2009). Bucky Ball Organizes Germ Plasm Assembly in Zebrafish. *Curr Biol* **19**, 414–422.

Brauer, A. (1894). Beiträge zur Kenntnis der Entwicklungsgeschichte des Skorpions. *Z wiss. Zool.* **57**, 402–432.

Brauer, A. (1925). Studies on the embryology of Bruchus quadrimaculatus, Fabr. *Ann. Entomol. Soc. Am.* **18**, 283–312.

Buchta, T., Özüak, O., Stappert, D., Roth, S. and Lynch, J. A. (2013). Patterning the dorsal–ventral axis of the wasp Nasonia vitripennis. *Dev Biol* **381**, 189–202.

Bull, A. L. (1982). Stages of Living Embryos in the Jewel Wasp Mormoniella (Nasonia) vitripennis (Walker) (Hymenoptera: Pteromalidae). *Int. J. Insect Morphol. Embryol.* **11**, 1–23.

Butt, F. H. (1949a). Embryology of the Milkweed Bug, Oncopeltus fasciatus (Hemiptera). *Cornell Experiment Station Memoir* 3–43.

Butt, F. (1949b). *Embryology of the milkweed bug, Oncopeltus fasciatus (Hemiptera)*. Ithaca, NY: Cornell University Agricultural Experiment Station.

Cannon, H. G. (1921). The Early Development of the summer egg of a Cladoceran (Simocephalus vetulus). *J. Cell Sci.* **s2-65**, 627–642.

Cannon, H. G. (1924). On the development of an estherid crustacean. *Philos. Trans. R. Soc. Lond. Ser B, Contain. Pap. a Biol. Character* **212**, 395–430.

Carter, J.-M., Gibbs, M. and Breuker, C. J. (2015). Divergent RNA Localisation Patterns of Maternal Genes Regulating Embryonic Patterning in the Butterfly Pararge aegeria. *PLoS ONE* **10**, e0144471.

Cavallin, M. (1971). ["Substitutive polyembryony" and the problem of germinal line origin in the phasma Carausius morosus Br]. *C. R. Hebd. des seances l'Acad. des Sci. Ser. D: Sci. Nat.* **272**, 462–5.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D., Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., et al. (2010). Adaptive Evolution of Pelvic Reduction in Sticklebacks by Recurrent Deletion of a *Pitx1* Enhancer. *Science* **327**, 302–305.

Chang, C., Dearden, P. and Akam, M. (2002). Germ Line Development in the Grasshopper Schistocerca gregaria: vasa As a Marker. *Dev. Biol.* **252**, 100–118.

Chang, C., Lee, W.-C., Cook, C. E., Lin, G.-W. and Chang, T. (2006). Germ-plasm specification and germline development in the parthenogenetic pea aphid Acyrthosiphon pisum: Vasa and Nanos as markers. *Int J Dev Biol* **50**, 413–421.

Chang, C., Lin, G., Cook, C. E., Horng, S., Lee, H. and Huang, T. (2007). Apvasa marks germ-cell migration in the parthenogenetic pea aphid Acyrthosiphon pisum (Hemiptera: Aphidoidea). *Dev. Genes Evol.* **217**, 275–287.

Chaw, R. C., Vance, E. and Black, S. D. (2007). Gastrulation in the spider Zygiella x-notata involves three distinct phases of cell internalization. *Dev Dynam* **236**, 3484–3495.

Chen, R. P.-Y., Shih, C.-L., Lin, G.-W., Cook, C. E., Huang, T.-Y. and Chang, C.-C. (2009). Developmental expression of Apnanos during oogenesis and embryogenesis in the parthenogenetic pea aphid Acyrthosiphon pisum. *Int J Dev Biol* **53**, 169–176.

Chen, Y., Fang, X., Tian, X.-Q., Cui, Z., Feng, H.-Y. and Qiu, G.-F. (2021). Germ plasm and the origin of the primordial germ cells in the oriental river prawn, Macrobrachium nipponense. *Cell Tissue Res.* **386**, 559–569.

Clark, E., Peel, A. D. and Akam, M. (2019). Arthropod segmentation. *Development* **146**, dev170480.

Colonnetta, M. M., Goyal, Y., Johnson, H. E., Syal, S., Schedl, P. and Deshpande, G. (2022). Preformation and epigenesis converge to specify primordial germ cell fate in the early Drosophila embryo. *PLoS Genet.* **18**, e1010002.

Counce, S. J. (1963). Developmental morphology of polar granules in Drosophila. Including observations on pole cell behavior and distribution during embryogenesis. *J. Morphol.* **112**, 129–145.

Dannenberg, L. C. and Seaver, E. C. (2018). Regeneration of the germline in the annelid Capitella teleta. *Dev. Biol.* **440**, 74–87.

Dearden, P. K. (2006). Germ cell development in the Honeybee (Apis mellifera); Vasa and Nanos expression. *BMC Dev. Biol.* **6**, 6.

Dearden, P., Grbic, M. and Donly, C. (2003). Vasa expression and germ-cell
specification in the spider mite Tetranychus urticae. *Dev. Genes Evol.* **212**, 599–603.

Dohle, W. (1964). Die Embryonalentwicklung von Glomeris marginata (Villers) im
Vergleich zur Entwicklung anderer Diplopoden. *Zool. Jahrb. Anat.* **81**, 241–310.

Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D. A., Henderson, L. and
Extavour, C. G. (2014). BMP signaling is required for the generation of primordial
germ cells in an insect. *Proc. Natl. Acad. Sci.* **111**, 4133–4138.

Doyle, D. A., Burian, F. N., Aharoni, B., Klinder, A. J., Menzel, M. M., Nifras, G. C.
C., Shabazz-Henry, A. L., Palma, B. U., Hidalgo, G. A., Sottolano, C. J., et al.
(2023). Germ Granule Evolution Provides Mechanistic Insight into Drosophila
Germline Development. *Mol Biol Evol* **40**, msad174.

Du, S., Tihelka, E., Yu, D., Chen, W.-J., Bu, Y., Cai, C., Engel, M. S., Luan, Y.-X. and
Zhang, F. (2024). Revisiting the four Hexapoda classes: Protura as the sister group
to all other hexapods. *Proc. Natl. Acad. Sci.* **121**, e2408775121.

Eddy, E. M. (1975). Germ Plasm and the Differentiation of the Germ Cell Line.
International Review of Cytology **43**, 229–280.

Edgar, A., Bates, C., Larkin, K. and Black, S. (2015). Gastrulation occurs in multiple
phases at two distinct sites in Latrodectus and Cheiracanthium spiders. *EvoDevo* **6**,
33.

Ephrussi, A. and Lehmann, R. (1992). Induction of germ cell formation by oskar.
Nature **358**, 387–392.

Evans, R. (1901). On the Malayan Species of Onychophora. Part II - The Development
of Eoperipatus weldoni. *Q. J. Microscop. Sci.* **45**, 41–88.

Evans, T., Wade, C. M., Chapman, F. A., Johnson, A. D. and Loose, M. (2014).
Acquisition of Germ Plasm Accelerates Vertebrate Evolution. *Science* **344**, 200–203.

Everett, N. B. (1944). The present status of the germ-cell problem in vertebrates. *Biol*
Rev **20**, 45–55.

Ewen-Campen, B., Schwager, E. E. and Extavour, C. G. M. (2010). The molecular
machinery of germ line specification. *Mol. Reprod. Dev.* **77**, 3–18.

Ewen-Campen, B., Srouji, J. R., Schwager, E. E. and Extavour, C. G. (2012). oskar
Predates the Evolution of Germ Plasm in Insects. *Curr Biol* **22**, 2278–2283.

Ewen-Campen, B., Donoughe, S., Clarke, D. N. and Extavour, C. G. (2013a). Germ
cell specification requires zygotic mechanisms rather than germ plasm in a basally
branching insect. *Curr. Biol.* **23**, 835–842.

Ewen-Campen, B., Jones, T. E. M. and Extavour, C. G. (2013b). Evidence against a
germ plasm in the milkweed bug Oncopeltus fasciatus, a hemimetabolous insect.
Biol. Open **2**, 556–568.

Extavour, C. G. (2005). The fate of isolated blastomeres with respect to germ cell
formation in the amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* **277**, 387–402.

Extavour, C. G. M. (2007). Evolution of the bilaterian germ line: lineage origin and
modulation of specification mechanisms. *Integr Comp Biol* **47**, 770–785.

Extavour, C. G. and Akam, M. (2003). Mechanisms of germ cell specification across
the metazoans: epigenesis and preformation. *Development* **130**, 5869–5884.

Faussek, V. (1888). Ueber die embryonale Entwicklung der Geschlechtsorgane bei der
Afterspinne (Phalangium). *Biol. Zentralbl.* **8**, 359–363.

Faussek, V. (1892). Zur Anatomie und Embryologie der Phalangiden. *Biol. Zentralbl.*
12, 1–8.

Fernández, R., Edgecombe, G. D. and Giribet, G. (2018). Phylogenomics illuminates
the backbone of the Myriapoda Tree of Life and reconciles morphological and
molecular phylogenies. *Sci. Rep.* **8**, 83.

Forrest, K. M. and Gavis, E. R. (2003). Live Imaging of Endogenous RNA Reveals a
Diffusion and Entrapment Mechanism for nanos mRNA Localization in Drosophila.
Curr Biol **13**, 1159–1168.

Gabriel, W. N., McNuff, R., Patel, S. K., Gregory, T. R., Jeck, W. R., Jones, C. D.
and Goldstein, B. (2007). The tardigrade Hypsibius dujardini, a new model for
studying the evolution of development. *Dev. Biol.* **312**, 545–559.

Gainett, G., Crawford, A. R., Klementz, B. C., So, C., Baker, C. M., Setton, E. V. W.
and Sharma, P. P. (2022). Eggs to long-legs: embryonic staging of the harvestman
Phalangium opilio (Opiliones), an emerging model arachnid. *Front. Zool.* **19**, 11.

Gatenby, J. B. (1917). The embryonic development of Trichogramma evanescens,
Westw., monembryonic egg parasite of Donacia simplex, Fab. *J. Cell Sci.* **s2-62**,
149–187.

Geigy, R. (1931). Action de l'ultra-violet sur le pôle germinal dans l'oeuf de Drosophila
melanogaster (Castration et mutabilité). *Rev. suisse Zoöl.* **38**, 187–285.

Gerberding, M. and Patel, N. H. (2004). Chapter 6. Gastrulation in Crustaceans: germ layers and cell lineages. In *Gastrulation: From Cells to Embryo* (ed. Stern, C.), p. Cold Spring Harbor Laboratory Press.

Gerberding, M., Browne, W. E. and Patel, N. H. (2002). Cell lineage analysis of the amphipod crustacean Parhyale hawaiensis reveals an early restriction of cell fates. *Development* **129**, 5789–5801.

Gilbert, S. (1997). *Embryology: Constructing the Organism.* Sunderland, MA, USA: Sinauer Associates Inc.

Giribet, G. and Edgecombe, G. D. (2019). The Phylogeny and Evolutionary History of Arthropods. *Curr Biol* **29**, R592–R602.

Goodrich, A. L. (1939a). The origin and fate of the entoderm elements in the embryogeny of Porcellio laevis latr. and Armadillidium nasatum B. L. (Isopoda). *J. Morphol.* **64**, 401–429.

Goodrich, A. L. (1939b). The origin and fate of the entoderm elements in the embryogeny of Porcellio laevis Latr. and Armadillidium nasutum B.L. (Isopoda). *Journal of Morphology* **64**, 401–426.

Görg, I. (1959). Untersuchungen am Keim von Hierodula (Rhombodera) crassa Giglio Tos, ein Beitrag zur Embryologie der Mantiden. *Dtsch. Entomol. Z.* **6**, 389–450.

Goss, R. J. (1952a). The early embryology of the book louse, Liposcelis divergens Badonnel (Psocoptera; liposcelidae). *J. Morphol.* **91**, 135–167.

Goss, R. J. (1952b). The early embryology of the book louse, Liposcelis divergens Badonnel (Psocoptera; Liposcelidae). *Journal of Morphology* **91**, 135–167.

Gough, L. H. (1902). The development of Admetus pumilio, Koch: a contribution to the embryology of the Pedipalps. *J. Cell Sci.* **s2-45**, 595–630.

Grattan, R. M., McCulloch, R. J., Sellars, M. J. and Hertzler, P. L. (2013). Ultrastructure of putative germ granules in the penaeid shrimp Marsupenaeus japonicus. *Arthropod Struct. Dev.* **42**, 153–164.

Green, J. E. and Akam, M. (2014). Germ cells of the centipede Strigamia maritima are specified early in embryonic development. *Dev. Biol.* **392**, 419–430.

Gupta, T. and Extavour, C. G. (2013). Identification of a putative germ plasm in the amphipod Parhyale hawaiensis. *EvoDevo* **4**, 34.

Haga, K. (1985). Oogenesis and embryogenesis of the idolothripine thrips, Bactrothrips brevitubus (Thysanoptera, Phlaeothripidae). In *Recent Advances in Insect Embryology in Japan*, pp. 45–106. Tsukuba: ISEBU.

Hegner, R. W. (1908). Effects of removing the germ-cell determinants from the eggs of some chrysomelid beetles. Preliminary report. *Biol. Bull.* **16**, 19–26.

Hegner, R. W. (1909a). The effects of centrifugal force upon the eggs of some chrysomelid beetles. *J. Exp. Zoöl.* **6**, 507–552.

Hegner, R. W. (1909b). The origin and early history of the germ-cells in some chrysomelid beetles. *J. Morphol.* **20**, 231–296.

Hegner, R. W. (1911). Experiments with chrysomelid beetles. III. The effects of killing parts of the eggs of Leptinotarsa decemlineata. *Biol Bull* **20**, 237–251.

Hegner, R. W. (1914). Studies on germ cells. I. The history of the germ cells in insects with special reference to the Keimbahn-determinants. II. The origin and significance of the Keimbahn-determinants in animals. *J. Morphol.* **25**, 375–509.

Heikes, K. L., Game, M., Smith, F. W. and Goldstein, B. (2023). The embryonic origin of primordial germ cells in the tardigrade Hypsibius exemplaris. *Dev. Biol.* **497**, 42–58.

Hejnol, A. and Schnabel, R. (2005). The eutardigrade Thulinia stephaniae has an indeterminate development and the potential to regulate early blastomere ablations. *Development* **132**, 1349–1361.

Heming, B. S. (1979). Origin and fate of germ cells in male and female embryos of Haplothrips verbasci (Osborn) (Insecta, Thysanoptera, Phlaeothripidae). *J. Morphol.* **160**, 323–343.

Heming, B. S. and Huebner, E. (1994). Development of the germ cells and reproductive primordia in male and female embryos of Rhodnius prolixus Stål (Hemiptera: Reduviidae). *Can. J. Zool.* **72**, 1100–1119.

Hertzler, P. L. (2005). Cleavage and gastrulation in the shrimp Penaeus (Litopenaeus) vannamei (Malacostraca, Decapoda, Dendrobranchiata). *Arthropod Struct. Dev.* **34**, 455–469.

Heymons, R. (1895). *Die Embryonalentwickelung von Dermapteren und Orthopteren unter Besonderer Berücksichtigung der Keimblätterbildung*. Jena: Jena Verlag von Gustav Fischer.

Heymons, R. (1901). Die Entwicklungsgeschichte der Scolopender. *Zoologica* **33**, 1–244.

Hickman, V. V. (1936). The embryology of the syncarid crustacean, Anaspides
tasmaniae. *Pap. Proc. Roy. Soc. Tasman.* 1–36.

Holm, Å. (1947). On the development of Opilio parietinus Deg. *Zool. bidr. Uppsala* 409–
422.

Hopf, C., Viebahn, C. and Püschel, B. (2011). BMP signals and the transcriptional
repressor BLIMP1 during germline segregation in the mammalian embryo. *Dev*
Genes Evol **221**, 209.

Huettner, A. F. (1923). The origin of the germ cells in Drosophila melanogaster. *Journal*
of Morphology **2**, 385–423.

Ikuta, K. and Makioka, T. (1994). Notes on the postembryonic development of the
ovary in Argulus japonicus (Crustacea: Branchiura). *Proc Arthropod Embryol Soc Jpn*
29, 15–17.

Illmensee, K. and Mahowald, A. P. (1974). Transplantation of posterior polar plasm in
Drosophila. Induction of germ cells at the anterior pole of the egg. *Proc. Natl. Acad.*
Sci. **71**, 1016–1020.

Illmensee, K., Mahowald, A. P. and Loomis, M. R. (1976). The ontogeny of germ
plasm during oogenesis in Drosophila. *Dev Biol* **49**, 40–65.

Jazdowska-Zagrodzińska, B. (1966). Experimental studies on the role of "polar
granules" in the segregation of pole cells in Drosophila melanogaster. *J Embryol Exp*
Morph **16**, 391–9.

Johannsen, O. and Butt, F. (1941). *Embryology of insects and myriapods: the*
developmental history of insects, centipedes, and millepedes from egg deposition to
hatching. McGraw-Hill, Inc.

Juhn, J. and James, A. A. (2006). oskar gene expression in the vector mosquitoes,
Anopheles gambiae and Aedes aegypti. *Insect Mol. Biol.* **15**, 363–372.

Juhn, J., Marinotti, O., Calvo, E. and James, A. A. (2008). Gene structure and
expression of nanos (nos) and oskar (osk) orthologues of the vector mosquito, Culex
quinquefasciatus. *Insect Mol. Biol.* **17**, 545–552.

Jura, C. (1967). Origin of germ cells and gonads formation in embryogenesis of
Tetrodontophora bielanensis (Collembola). *Acta Biol Crac Zool* **10**, 97–103.

Kaczmarczyk, A. N. (2014). Germline maintenance and regeneration in the amphipod
crustacean, Parhyale hawaiensis.

Kautzsch, G. (1910). Über die Entwicklung von Agelena labyrinthica Clerck. II. Teil. *Zool. Jahrb. Anat.* **30**, 535–602.

Kemph, A. and Lynch, J. A. (2022). Evolution of germ plasm assembly and function among the insects. *Curr Opin Insect Sci* **50**, 100883.

Kemph, A., Kharel, K., Tindell, S. J., Arkov, A. L. and Lynch, J. A. (2025). Novel structure and composition of the unusually large germline determinant of the wasp Nasonia vitripennis. *Mol. Biol. Cell* **36**, ar55.

Kharel, K., Tindell, S. J., Kemph, A., Schmidtke, R., Alexander, E., Lynch, J. A. and Arkov, A. L. (2024). Dynamic protein assembly and architecture of the large solitary membraneless organelle during germline development in the wasp Nasonia vitripennis. *Development* **151**, dev202877.

Khila, A. and Abouheif, E. (2008). Reproductive constraint is a developmental mechanism that maintains social harmony in advanced ant societies. *Proc National Acad Sci* **105**, 17884–17889.

Kistler, K. E., Trcek, T., Hurd, T. R., Chen, R., Liang, F.-X., Sall, J., Kato, M. and Lehmann, R. (2018). Phase transitioned nuclear Oskar promotes cell division of Drosophila primordial germ cells. *eLife* **7**, e37949.

Klag, J. (1982). Germ line of Tetrodontophora bielanensis (Insecta, Collembola). Ultrastructural study on the origin of primordial germ cells. *Development* **72**, 183–195.

Kobayashi, Y., Watanabe, T. and Suzuki, H. (2006). Embryonic development of the firefly Pyrocoelia rufa Olivier (Insecta: Coleoptera, Lampyridae), with special reference to its hibernal diapause. *Proc Arthropod Embryol Soc Jpn* **41**, 47–53.

Kobayashi, Y., Niikura, K., Oosawa, Y. and Takami, Y. (2013). Embryonic development of Carabus insulicola (Insecta, Coleoptera, Carabidae) with special reference to external morphology and tangible evidence for the subcoxal theory. *J. Morphol.* **274**, 1323–1352.

Komatsu, S. and Kobayashi, Y. (2012). Embryonic development of a whirligig beetle, Dineutus mellyi, with special reference to external morphology (Insecta: Coleoptera, Gyrinidae). *J. Morphol.* **273**, 541–560.

Krause, G. (1939). Die Eitypen der Insekten. *Biol. Zentralbl.* **59**, 495–536.

Krause, J. B. (1947). The development of the gonads of the wood-eating beetle, Passalus cornutus Fabricius. *Ann. Entomol. Soc. Am.* **40**, 172–202.

Kühn, A. (1913). Die Sonderung der Keimesbezirke in der Entwicklung der Sommereier
von Polyphemus pediculus de Geer. *Zool. Jahrb. Anat.* **35**, 243–340.

Kulkarni, A. and Extavour, C. G. (2017). Convergent evolution of germ granule
nucleators: A hypothesis. *Stem Cell Res.* **24**, 188–194.

Kulkarni, A., Ewen-Campen, B., Terao, K., Matsumoto, Y., Li, Y., Watanabe, T.,
Kao, J. A., Parhad, S. S., Ylla, G., Mizunami, M., et al. (2023). oskar acts with the
transcription factor Creb to regulate long-term memory in crickets. *Proc. Natl. Acad.*
Sci. **120**, e2218506120.

Lall, S., Ludwig, M. Z. and Patel, N. H. (2003). Nanos Plays a Conserved Role in Axial
Patterning outside of the Diptera. *Curr Biol* **13**, 224–229.

Larink, O. (1969). Zur Entwicklungsgeschichte von Petrobius brevistylis (Thysanura,
Insecta). *Helgoländer Wiss. Meeresunters.* **19**, 111–155.

Lasko, P. F. and Ashburner, M. (1990). Posterior localization of vasa protein
correlates with, but is not sufficient for, pole cell development. *Gene Dev* **4**, 905–921.

Lawson, K. A., Dunn, N. R., Roelen, B. A. J., Zeinstra, L. M., Davis, A. M., Wright,
C. V. E., Korving, J. P. W. F. M. and Hogan, B. L. M. (1999). Bmp4 is required for
the generation of primordial germ cells in the mouse embryo. *Gene Dev* **13**, 424–
436.

Lee, C.-Y. S., Putnam, A., Lu, T., He, S., Ouyang, J. P. T. and Seydoux, G. (2020).
Recruitment of mRNAs to P granules by condensation with intrinsically-disordered
proteins. *eLife* **9**, e52896.

Lehmann, R. (2016). Germ Plasm Biogenesis - An Oskar-Centric Perspective. *Curr*
Top Dev Biol **116**, 679–707.

Leptin, M. (2004). Chapter 7: Gastrulation in Drosophila. In *Gastrulation: From Cells to*
Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor Laboratory
Press.

Lin, G., Cook, C. E., Miura, T. and Chang, C. (2014). Posterior localization of ApVas1
positions the preformed germ plasm in the sexual oviparous pea aphid Acyrthosiphon
pisum. *EvoDevo* **5**, 18.

Lochab, A. K. and Extavour, C. G. (2017). Bone Morphogenetic Protein (BMP)
signaling in animal reproductive system development and function. *Dev Biol* **427**,
258–269.

Lynch, J. A. (2019). Evolution of maternal control of axial patterning in insects. *Curr*
Opin Insect Sci **31**, 37–42.

Lynch, J. A. and Desplan, C. (2010). Novel modes of localization and function of
nanos in the wasp Nasonia. *Development* **137**, 3813–3821.

Lynch, J. A., Özüak, O., Khila, A., Abouheif, E., Desplan, C. and Roth, S. (2011).
The Phylogenetic Origin of oskar Coincided with the Origin of Maternally Provisioned
Germ Plasm and Pole Cells at the Base of the Holometabola. *PLoS Genet.* **7**,
e1002029.

Mahowald, A. P. (1962). Fine structure of pole cells and polar granules in Drosophila
melanogaster. *J Exp Zool* **151**, 201–215.

Mahowald, A. P. (1968). Polar granules of Drosophila. II. Ultrastructural changes during
early embryogenesis. *J. Exp. Zool.* **167**, 237–261.

Mahowald, A. P. (2001). Assembly of the Drosophila germ plasm. *Int Rev Cytol* **203**,
187–213.

Manton, S. M. (1928). On the Embryology of a Mysid Crustacean, Hemimysis
lamornae. *Philos. Trans. R. Soc. Lond. Ser. B, Contain. Pap. a Biol. Character* **216**,
363–463.

Manton, S. M. (1934). On the embryology of the crustacean Nebalia bipes.
Philosophical Transactions Royal Soc Lond Ser B **223**, 163–238.

Manton, S. M. (1949). Studies on the Onychophora VII. The early embryonic stages of
Peripatopsis, and some general considerations concerning the morphology and
phylogeny of the Arthropoda. *Philos. Trans. R. Soc. Lond. Ser. B, Biol. Sci.* **233**,
483–580.

Marlow, F. L. and Mullins, M. C. (2008). Bucky ball functions in Balbiani body
assembly and animal–vegetal polarity in the oocyte and follicle cell layer in zebrafish.
Dev Biol **321**, 40–50.

Martin, A. and Orgogozo, V. (2013). The loci of repeated evolution: A catalog of
genetic hotspots of phenotypic variation. *Evolution* **67**, 1235–1250.

Mayer, G. and Tait, N. N. (2009). Position and development of oocytes in velvet worms
shed light on the evolution of the ovary in Onychophora and Arthropoda. *Zool. J.
Linn. Soc. Lond.* **157**, 17–33.

McClendon, J. F. (1907). On the development of parasitic copepods. Part II. *Biol. Bull.*
12, 53–88.

Mellanby, H. (1935a). The early embryonic development of Rhodnius prolixus
(Hemiptera, Heteroptera). *J. Cell Sci.* **s2-78**, 71–90.

Mellanby, H. (1935b). The Early Embryonic Development of Rhodnius prolixus
(Hemiptera, Heteroptera). *Q. J. Microscop. Sci.* **309**, 71–90.

Metschnikow, E. (1866). *Embryologische studien an insecten*. Leipzig, Germany:
Verlag von Wilhelm Engelmann.

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C., Frandsen,
P. B., Ware, J., Flouri, T., Beutel, R. G., et al. (2014). Phylogenomics resolves the
timing and pattern of insect evolution. *Science* **346**, 763–767.

Mitsumoto, H. and Makioka, T. (2002). Preliminary notes on the structure and
postembryonic development of gonads in two tadpole shrimps Triops longicaudatus
and T. numidicus (Crustacea: Branchiopoda, Notostraca). *Proc. Arthropod. Embryol.*
Soc. Jpn. **37**, 13–17.

Miya, K. (1953). The presumptive genital region at the blastoderm stage of the silkworm
egg. *J Fac Agr Iwate Univ* **1**, 223–227.

Miya, K. (1958a). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part I. Differentiation of Germ Cells. *J. Fac. Agr. Iwate*
Univ. 436–467.

Miya, K. (1958b). Studies on the embryonic development of the gonad in the silkworm,
Bombyx mori L. Part II. Formation of the gonad. *J Fac Agr Iwate Univ* **4**, 126–151.

Miyakawa, K. (1974). The embryology of the caddisfly Stenopsyche griseipennis
MacLachlan (Trichoptera, Stenopsychidae): IV. Organogenesis: mesodermal
derivatives. *Kontyû* **42**, 451–466.

Miyazaki, K. and Makioka, T. (2012). Postembryonic development of the female
reproductive system in the pycnogonid, Propallene longiceps (Pycnogonida,
Callipallenidae). *Invertebr. Reprod. Dev.* **56**, 287–292.

Morris, S. C. (2009). The predictability of evolution: glimpses into a post-Darwinian
world. *Die Naturwissenschaften* **96**, 1313–37.

Nakamura, T. and Extavour, C. G. (2016). The transcriptional repressor Blimp-1 acts
downstream of BMP signaling to generate primordial germ cells in the cricket Gryllus
bimaculatus. *Development* **143**, 255–263.

Nakao, H. (1999). Isolation and characterization of a Bombyx vasa-like gene. *Dev.*
Genes Evol. **209**, 312–316.

Nakao, H. and Takasu, Y. (2019). Complexities in Bombyx germ cell formation process
revealed by Bm-nosO (a Bombyx homolog of nanos) knockout. *Dev. Biol.* **445**, 29–
36.

Nakao, H., Hatakeyama, M., Lee, J. M., Shimoda, M. and Kanda, T. (2006).
Expression pattern of Bombyx vasa-like (BmVLG) protein and its implications in germ
cell development. *Dev. Genes Evol.* **216**, 94–99.

Nakao, H., Matsumoto, T., Oba, Y., Niimi, T. and Yaginuma, T. (2008). Germ cell
specification and early embryonic patterning in Bombyx mori as revealed by nanos
orthologues. *Evol Dev* **10**, 546–554.

Needham, A. E. (1942). The structure and development of the segmental excretory
organs of Asellus aquaticus (Linné). *J. Cell Sci.* **s2-83**, 205–243.

Nelsen, O. E. (1934). The segregation of the germ cells in the grasshopper, Melanoplus
differentialis (Acrididae; Orthoptera). *J. Morphol.* **55**, 545–575.

Nelson, J. A. (1915). *The embryology of the honey bee*. Princeton: Princeton University
Press.

Nieuwkoop, P. D. and Sutasurya, L. A. (1979). *Primordial germ cells in the chordates:
embryogenesis and phylogenesis*. Cambridge University Press.

Nieuwkoop, P. D. and Sutasurya, L. A. (1981). *Primordial germ cells in the
invertebrates: from epigenesis to preformation*. Cambridge: Cambridge University
Press.

Niikura, K., Hirasawa, K., Inoda, T. and Kobayashi, Y. (2017). Embryonic
development of a diving beetle, Hydaticus pacificus Aubé (Insecta: Coleoptera;
Dytiscidae): external morphology and phylogenetic implications. *Proc Arthropod
Embryol Soc Jpn* **48**, 19–32.

Okada, M., Kleinman, I. A. and Schneiderman, H. A. (1974). Restoration of fertility in
sterilized Drosophila eggs by transplantation of polar cytoplasm. *Dev. Biol.* **37**, 43–
54.

Olesnicky, E. C. and Desplan, C. (2007). Distinct mechanisms for mRNA localization
during embryonic axis specification in the wasp Nasonia. *Dev Biol* **306**, 134–142.

Özpolat, B. D. (2024). Annelids as models of germ cell and gonad regeneration. *J. Exp.
Zoöl. Part B: Mol. Dev. Evol.* **342**, 126–143.

Pagel, M. (1994). Detecting correlated evolution on phylogenies: a general method for
the comparative analysis of discrete characters. *Proc. R. Soc. Lond. Ser. B: Biol. Sci.*
255, 37–45.

Paz, J. de la (1958). Studies on the embryogeny of Oncopeltus fasciatus (Dallas)
(Hemiptera: Lygaeidae).

Pires-daSilva, A. and Sommer, R. J. (2003). The evolution of signalling pathways in
animal development. *Nat Rev Genet* **4**, 39–49.

Poulson, D. F. (1947). The pole cells of Diptera, their fate and significance. *Proc Natl*
Acad Sci U S A **33**, 182–4.

Poulson, D. F. (1950). Histogenesis, organogenesis, and differentiation in the embryo
of Drosophila melanogaster Meigen.pp. 168–274. New York: Wiley.

Presser, B. D. and Rutschky, C. W. (1957). The embryonic development of the corn
earworm, Heliothis zea (Boddie) (Lepidoptera, Phalaenidae). *Ann. Entomol. Soc. Am.*
50, 133–164.

Quan, H. and Lynch, J. A. (2016). The evolution of insect germline specification
strategies. *Curr. Opin. Insect Sci.* **13**, 99–105.

Rafiqi, Ab. M., Rajakumar, A. and Abouheif, E. (2020). Origin and elaboration of a
major evolutionary transition in individuality. *Nature* **585**, 239–244.

Reed, R. D., Papa, R., Martin, A., Hines, H. M., Counterman, B. A., Pardo-Diaz, C.,
Jiggins, C. D., Chamberlain, N. L., Kronforst, M. R., Chen, R., et al. (2011). optix
drives the repeated convergent evolution of butterfly wing pattern mimicry. *Science*
333, 1137–41.

Rempel, J. G. and Church, N. S. (1969). The embryology of Lytta viridana Le Conte
(Coleoptera: Meloidae). V. The blastoderm, germ layers, and body segments. *Can. J.*
Zool. **47**, 1157–1171.

Rivard, E. L., Srouji, J., Repouliou, A. and Extavour, C. G. (2025). Protein sequence
evolution underlies interspecies incompatibility of a cell fate determinant. *bioRxiv*.

Rosenberg, M. I., Lynch, J. A. and Desplan, C. (2009). Heads and tails: Evolution of
antero-posterior patterning in insects. *Biochimica Et Biophysica Acta Bba - Gene*
Regul Mech **1789**, 333–342.

Roth, S. (2004). Chapter 8: Gastrulation in Other Insects. In *Gastrulation: From Cells to*
Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor Laboratory
Press.

Saitou, M. and Yamaji, M. (2010). Germ cell specification in mice: signaling,
transcription regulation, and epigenetic consequences. *Reproduction* **139**, 931–942.

Satterlee, J. W., Alonso, D., Gramazio, P., Jenike, K. M., He, J., Arrones, A.,
Villanueva, G., Plazas, M., Ramakrishnan, S., Benoit, M., et al. (2024).
Convergent evolution of plant prickles by repeated gene co-option over deep time.
Science **385**, eado1663–eado1663.

Schimkewitsch, W. (1896). Studien über parasitische Copepoden. *Z. wiss. Zool.* **61**,
339–362.

Schimkewitsch, V. M. (1903). Über die Entwicklung von Thelyphonus caudatus (L.).
Zool. Anz. **26**, 665–685.

Scholl, A., Liu, Y. and Seydoux, G. (2024). Caenorhabditis elegans germ granules
accumulate hundreds of low translation mRNAs with no systematic preference for
germ cell fate regulators. *Dev. (Camb., Engl.)* **151**, dev202575.

Scholtz, G. (2000). Evolution of the nauplius stage in malacostracan crustaceans*. *J
Zool Syst Evol Res* **38**, 175–187.

Scholtz, G. and Wolff, C. (2002). Cleavage, gastrulation, and germ disc formation of
the amphipod Orchestia cavimana (Crustacea, Malacostraca, Peracarida). *Contrib.
Zool.* **71**, 9–28.

Scholtz, G. and Wolff, C. (2013). Chapter 4: Arthropod Embryology: Cleavage and
Germ Band Development. In *Arthropod Biology and Evolution* (ed. Minelli, A.), pp.
63–89.

Schröder, R. (2006). vasa mRNA accumulates at the posterior pole during blastoderm
formation in the flour beetle Tribolium castaneum. *Dev. Genes Evol.* **216**, 277.

Schwager, E. E., Meng, Y. and Extavour, C. G. (2015). vasa and piwi are required for
mitotic integrity in early embryogenesis in the spider Parasteatoda tepidariorum. *Dev.
Biol.* **402**, 276–290.

Schwentner, M., Combosch, D. J., Nelson, J. P. and Giribet, G. (2017). A
Phylogenomic Solution to the Origin of Insects by Resolving Crustacean-Hexapod
Relationships. *Curr Biol* **27**, 1818-1824.e5.

Sedgwick, A. (1887). The development of the Cape species of Peripatus: Part III. On
the changes from stage A to stage F. *J. Cell Sci.* **s2-27**, 467–550.

Seervai, R. N. H. and Wessel, G. M. (2013). Lessons for inductive germline
determination. *Mol Reprod Dev* **80**, 590–609.

Seidel, F. (1924). Die geschlechtsorgane in der embryonalen entwicklung von
Pyrrhocoris apterus L. *Z. Morphol. Ökol. Tiere* **1**, 429–506.

Sharma, P. P. and Gavish-Regev, E. (2025). The Evolutionary Biology of Chelicerata.
Annu. Rev. Entomol. **70**, 143–163.

Shinji, G. O. (1919). Embryology of coccids, with especial reference to the formation of
the ovary, origin and differentiation of the germ cells, germ layers, rudiments of the
midgut, and the intracellular symbiotic organisms. *J. Morphol.* **33**, 72–167.

Singh, J. P. (1967). Early embryonic development of gonads in Labidura riparia (Pallas)
(Labiduridae: Dermaptera). *Agra Univ. J. Res.* **1621**, 67–76.

Sonnenblick, B. P. (1941). Germ cell movements and sex differentiation of the gonads
in the Drosophila embryo. *Proc Natl Acad Sci U S A* **27**, 484–9.

Srouji, J. and Extavour, C. (2010). Redefining stem cells and assembling germ plasm:
key transitions in the evolution of the germ line.pp. 360–397. Enfield, NH: Science
Publishers.

Strömberg, J.-O. (1965). On the embryology of the isopod Idotea. *Ark. Zool.* **2**, 421–
473.

Strömberg, J.-O. (1968). Segmentation and organogenesis in Limnoria lignorum
(Rathke) (Isopoda). *Ark. Zool.* **20**, 91.

Sudler, M. T. (1899). The development of Penilia schmackeri Richard. *Proc. Boston
Soc. Nat. Hist.* **29**, 109–131.

Suzuki, N. (1990). Embryology of the Mecoptera (Panorpidae, Panorpodidae,
Bittacidae and Boreidae). *Bull. Sugadaira Montane Res. Cent.* **11**, 1–87.

Suzuki, N., Shimizu, S. and Ando, H. (1981). Early embryology of the alderfly, Sialis
mitsuhashii Okamoto (Megaloptera: Sialidae). *Int. J. Insect Morphol. Embryol.* **10**,
409–418.

Tadokoro, R., Sugio, M., Kutsuna, J., Tochinai, S. and Takahashi, Y. (2006). Early
Segregation of Germ and Somatic Lineages during Gonadal Regeneration in the
Annelid Enchytraeus japonensis. *Curr Biol* **16**, 1012–1017.

Takamura, K., Fujimura, M. and Yamaguchi, Y. (2002). Primordial germ cells
originate from the endodermal strand cells in the ascidian Ciona intestinalis. *Dev
Genes Evol* **212**, 11–18.

Tanaka, M. (1987). Differentiation and Behaviour of Primordial Germ Cells during the
Early Embryonic Development of Parnassius glacialis Butler, Luehdorfia japonica
Leech and Byasa (Atrophaneura) alcinous alcinous Klug (Lepidoptera: Papilionidae).
Recent Advances in Insect Embryology in Japan and Poland 255–266.

Terao, A. (1929). On the embryonic development of the spiny lobster, Panulirus
japonicus (v. Siebold). *Jpn J Zool* **2**, 387–449.

Tiegs, O. W. (1940). The embryology and affinities of the Symphyla, based on a study
of Hanseniella agilis. *J. Cell Sci.* **s2-82**, 1–208.

Tiegs, O. W. (1947). The development and affinities of the Pauropoda, based on a
study of Pauropus silvaticus, Part I. *J. Cell Sci.* **s3-88**, 165–267.

Toyama, K. (1902). Contributions to the Study of Silk-Worms. I. On the Embryology of
the Silk-Worms. *Bull. Coll. Agric. Tokyo Imp. Univ.* **5**, 73–118.

Trcek, T. and Lehmann, R. (2019). Germ granules in Drosophila. *Traffic* **20**, 650–660.

Trcek, T., Grosch, M., York, A., Shroff, H., Lionnet, T. and Lehmann, R. (2015).
Drosophila germ granules are structured and contain homotypic mRNA clusters. *Nat.*
Commun. **6**, 7962.

Ullmann, S. L. (1964). The origin and structure of the mesoderm and the formation of
the coelomic sacs in Tenebrio molitor L. [Insecta, Coleoptera]. *Philosophical*
Transactions Royal Soc Lond Ser B Biological Sci **248**, 245–277.

Underwood, E. M., Caulton, J. H., Allis, C. D. and Mahowald, A. P. (1980).
Developmental fate of pole cells in Drosophila melanogaster. *Dev. Biol.* **77**, 303–314.

Wang, C. and Lehmann, R. (1991). Nanos is the localized posterior determinant in
Drosophila. *Cell* **66**, 637–647.

Weismann, A. (1863a). Die Entwickelung der Dipteren im Ei, nach Beobachtungen an
Chironomous spec., Musca vomitoria und Pulex canis. *Zeitschrift für*
wissenschaftliche Zoologie **13**, 107–220.

Weismann, A. (1863b). Die Entwicklung der Dipteren im Ei nach Beobachtungen an
Chironomus,spec., Musca vomitoria und Pulex canis. *Z wiss Zool* **13**,.

Wheeler, W. M. (1893). A Contribution to Insect Embryology. *J. Morphol.* **8**, 1–160.

Whittle, C. A. and Extavour, C. G. (2016). Refuting the hypothesis that the acquisition
of germ plasm accelerates animal evolution. *Nat Commun* **7**, 12637.

Whittle, C. A. and Extavour, C. G. (2017). Causes and evolutionary consequences of
primordial germ-cell specification mode in metazoans. *Proc National Acad Sci* **114**,
5784–5791.

Witschi, E. (1934). On determinative cleavage and yolk formation in the harpactid
copepod Tisbe furcata (Baird). *Biol. Bull.* **67**, 335–340.

Wolff, C. (2009). The embryonic development of the malacostracan crustacean
Porcellio scaber (Isopoda, Oniscidea). *Dev. Genes Evol.* **219**, 545–564.

Wolff, C. and Scholtz, G. (2002). Cell Lineage, Axis Formation, and the Origin of Germ
Layers in the Amphipod Crustacean Orchestia cavimana. *Dev. Biol.* **250**, 44–58.

Woodland, J. T. (1957). A contribution to our knowledge of lepismatid development. *J.*
Morphol. **101**, 523–577.

Woodworth, C. W. (1889). Studies on the embryological development of Euvanessa
antiopa. In *The Butterflies of the Eastern United States and Canada, with special*
reference to New England, pp. 95–104. Cambridge, MA: Published by the Author.

Wotton, K. R., Jiménez-Guri, E., Matheu, B. G. and Jaeger, J. (2014). A staging
scheme for the development of the scuttle fly Megaselia abdita. *PLoS ONE* **9**,
e84421.

Xu, X., Brechbiel, J. L. and Gavis, E. R. (2013). Dynein-Dependent Transport of
nanos RNA in Drosophila Sensory Neurons Requires Rumpelstiltskin and the Germ
Plasm Organizer Oskar. *J Neurosci* **33**, 14791–14800.

Yajima, M. and Wessel, G. M. (2011). The DEAD-box RNA helicase Vasa functions in
embryonic mitotic progression in the sea urchin. *Development* **138**, 2217–2222.

Ying, Y. and Zhao, G.-Q. (2001). Cooperation of Endoderm-Derived BMP2 and
Extraembryonic Ectoderm-Derived BMP4 in Primordial Germ Cell Generation in the
Mouse. *Dev Biol* **232**, 484–492.

Ying, Y., Liu, X. M., Marble, A., Lawson, K. A. and Zhao, G. Q. (2000). Requirement
of Bmp8b for the generation of primordial germ cells in the mouse. *Mol. Endocrinol.*
14, 1053–63.

Yoon, Y., Klomp, J., Martin-Martin, I., Criscione, F., Calvo, E., Ribeiro, J. and
Schmidt-Ott, U. (2019). Embryo polarity in moth flies and mosquitoes relies on
distinct old genes with localized transcript isoforms. *eLife* **8**, e46711.

Zissler, D. (1992). From egg to pole cells: Ultrastructural aspects of early cleavage and
germ cell determination in insects. *Microsc. Res. Tech.* **22**, 49–74.

1538