There and back again: the dynamic evolution of panarthropod germ cell specification mechanisms

Jonchee A. Kao[1][†], Emily L. Rivard[1][†], Rishabh R. Kapoor[2], Cassandra G. Extavour[1,3,4] *

1. Department of Molecular and Cellular Biology, Harvard University, Cambridge MA, USA
2. Program for Systems, Synthetic and Quantitative Biology, Harvard University, Boston MA, USA
3. Department of Organismic and Evolutionary Biology, Harvard University, Cambridge MA, USA
4. Howard Hughes Medical Institute, Chevy Chase MD, USA

† equal contribution

* correspondence: extavour@oeb.harvard.edu

Abstract

Germ cells enable the reproduction of an organism and the continuity of its lineage.

These crucial cells are segregated from the soma at different times and places and via

distinct mechanisms across the animal tree of life. Making sense of the evolution of

germ cell specification across animals is complicated by the difficulty of making

meaningful comparisons across the diversity of animal embryonic development. Here,

we characterize germ cell specification within an ancient clade, that encompasses

massive animal biodiversity but within which we can conduct meaningful comparative

embryogenesis: the Panarthropoda. We amass data from centuries of studies

describing the timing and mechanisms of germ cell formation, and apply ancestral state

reconstruction to these data to propose novel hypotheses about the trajectory of

evolution in this process. Furthermore, we speculate on the mechanisms underlying

these evolutionary dynamics by considering the relationships among germ cell

specification, concurrent developmental processes, and the germ line gene network.

Collectively, this study derives new insights from a rich historical database of

embryological observations, and has broad implications for understanding the evolution

of metazoan germ cells.

Keywords: germ plasm, pole cells, inductive signaling, BMP, cytoplasmic determinant

I. Introduction

During development, animals specify the germ line, a collection of cells that will later give rise to the next generation. The specification and function of germ cells across animals has intrigued researchers for at least two millennia (Aristotle) . Whereas historical researchers relied primarily on observational data alone (Bounoure, 1939; Wolff, 1962) , we can now investigate these processes using functional genetic tools. These new techniques allow us not only to identify germ cells with more certainty, but also to gain more insight into the molecular mechanisms of germ cell specification. Several hypotheses have been proposed to explain evolutionary transitions between these mechanisms across animals (Buss, 1982; Buss, 1983; Crother et al., 2007; Crother et al., 2016; Evans et al., 2014; Extavour, 2007; Johnson et al., 2011; Whittle and Extavour, 2016; Whittle and Extavour, 2017) . However, such hypotheses are difficult to test because embryonic development of animals is too different across vast evolutionary time scales to be meaningfully compared. Moreover, there is disagreement in the field about whether existing well-studied mechanisms of specification are truly the only modes, and whether they are discrete, mutually exclusive mechanisms. Further, there are conflicting reports about the impact of germ cell specification mode on the evolution of genomes, body plans, and fitness (Evans et al., 2014; Otto and Hastings, 1998; Queller, 2000; Whittle and Extavour, 2016; Whittle and Extavour, 2017). Here, we examine variation in germ cell specification across the most speciose phylum of animals, wherein the general features of embryogenesis can nevertheless be meaningfully compared: the Panarthropoda. This clade displays striking diversity in the temporal and spatial origin of germ cells, as well as multiple transitions between

specification mechanisms, offering an ideal system to study evolution and regulation of

germ cell specification.

In this study, we first review the existing observational data regarding embryonic

germ line development across panarthropods. We are primarily interested in the first

embryonic appearance of germ cells before they enter the gonad, during which time we

refer to them as *primordial germ cells*. We identify three categories of temporally distinct

developmental context in which primordial germ cells are first detectable during

embryogenesis, and analyze the distribution of these three categories in their

phylogenetic context. We then review perturbational experimental studies aimed at

uncovering the mechanisms of germ cell specification. Finally, we discuss the

implications of these data on our understanding of the evolution of germ cell

specification in panarthropods, and more broadly across Metazoa.

I.i. How do we identify germ cells?

Historically, embryologists studied early development using histological

preparations of sectioned embryos (His, 1868) . With light microscopy, primordial germ

cells can be readily distinguished from surrounding somatic cells by their rounder and

larger nuclei (Nieuwkoop and Sutasurya, 1979). The chromatin of germ cells often

appears more diffuse than that of somatic cells, and the cytoplasm, especially cortical or

perinuclear cytoplasm, may contain electron-dense, non-membrane bound granular

material (Eddy, 1975). However, tracking these cells throughout development using

fixed samples and morphological features alone is not always technically possible.

Since the late 20th century, studies assessing gene expression patterns unique to

or diagnostic of germ cells (henceforth referred to as "molecular markers") have been

instrumental in identifying germ cells at earlier stages than previously recognized. The

most commonly used markers are the mRNA or protein products of the genes *vasa*,

nanos, *tudor*, and *piwi* (detectable by *in situ* hybridization or antibody staining,

respectively) (Ewen-Campen et al., 2010). The use of molecular markers can enable

germ cell identification at earlier developmental stages, since some germ cells express

their encoding genes before their morphological differentiation becomes unambiguously

detectable (see for example Ewen-Campen et al., 2013a) . In addition, these markers

facilitate tracking of germ cells over the course of development. However, using

molecular markers is a biased approach, as it relies on previous knowledge of germ line

gene expression from other species. Furthermore, interpretation of molecular marker

data can be complicated by pleiotropy, as their encoding genes may also be expressed

by somatic cells (e.g. Yajima and Wessel, 2011).

II. Germ cell formation in panarthropods

In our review of the embryological literature from 1860 to the present, we

identified three categories of primordial germ cell specification in panarthropods, based

on the timing of specification during embryogenesis (Figure 1). We describe these

categories below, along with relevant details of embryogenesis for context.

II.i. Category 1: PGC specification at early cleavage and blastoderm stages

After fertilization, embryogenesis begins with *cleavage*, a series of mitotic

divisions of the zygote (Gilbert, 1997). In most arthropods, cleavage leads to the

formation of the *blastoderm*, a relatively uniform layer of cells at the egg cortex,

surrounding a central yolk mass populated by a small number of yolk cells (Anderson,

1973; Scholtz and Wolff, 2013). The first category of primordial germ cell specification in

panarthropods occurs during or before the formation of the blastoderm (Figure 1A).

Insects

In most insects, the blastoderm is initially *syncytial*, with no cell membranes

separating the nuclei from each other (Anderson, 1973; Johannsen and Butt, 1941). In

some insects, the primordial germ cells are the first cells to form, appearing at the

posterior pole before the rest of the blastoderm is cellularized (Figure 1A1). Primordial

germ cells that appear in this manner are called *pole cells*. Pole cells are most

commonly seen in species of Diptera (flies, midges and mosquitoes), Coleoptera

(beetles), and Hymenoptera (bees, ants and wasps), although they have also been

reported in Thysanoptera (thrips) and Dermaptera (earwigs) (Table S1).

Pole cells were first reported in species of the midge *Chironomus* by Weismann

(Weismann, 1863), although Metschnikow (1866) was the first to follow their migration

into the gonad. In many insects, the cytoplasm at the posterior pole of the embryo is

morphologically distinct from that of the rest of the embryo, often with visible granules

(Hegner, 1914; Zissler, 1992). The close association of this granular cytoplasm with

pole cells led researchers to speculate that the granular material contained germ line

determinants (e.g. Hegner, 1909a; Zissler, 1992); we will describe experimental data

testing this hypothesis in Part III below. It was long believed that pole cells gave rise to

somatic cell types in addition to the germ cells (Anderson, 1962a; Anderson, 1962b;

Poulson, 1947; Poulson, 1950). However, careful lineage tracing in the fruit fly

Drosophila melanogaster showed that all pole cells are determined as germ cells, and

any that become lost during their migration die (Underwood et al., 1980).

While the function of pole cells as primordial germ cells appears conserved, the

specific details of their formation can vary considerably across species. For example,

pole cells differ in which cleavage cycle they cellularize. Pole cells form during the ninth

mitotic cycle of *D. melanogaster* (Huettner, 1923) and during the sixth mitotic cycle in

the parasitic wasp *Nasonia vitripennis* (Buchta et al., 2013). To our knowledge, the

earliest observed formation of pole cells is during the third cleavage cycle in the midges

Miastor metraloas and *Chironomus sp*. (Hegner, 1914; Weismann, 1863). The number

of pole cells formed is also variable: a single pole cell forms in embryos of the latter

midges (Hegner, 1914; Weismann, 1863), whereas four appear in the fruit fly

Bactrocera tryoni (Anderson, 1962a) and sixteen appear in the beetle *Calligrapha*

multipunctata (Hegner, 1909b).

After the pole cells form, they enter the egg and migrate to the location of the

future embryonic gonad, which is quite far from the posterior pole (Santos and

Lehmann, 2004) . Several different migration pathways have been described. In *D.*

melanogaster and many other Diptera, the pole cells are carried into the egg during

posterior midgut invagination, then migrate through the epithelium of the posterior

midgut to enter the abdominal mesoderm and associate with the developing somatic

gonad (Metschnikow, 1866; Sonnenblick, 1941). In Hymenoptera, pole cells are

internalized with the rest of the mesoderm during gastrulation (Gatenby, 1917). Yet

another variation is observed in Chrysomelidae (leaf beetles) (Hegner, 1909a), where

germ cells are internalized with the rest of the germ band as the extraembryonic

membranes fold over the germ band.

Other arthropods

Primordial germ cells are also specified early in Collembola (springtails), an

early-diverging branch of non-insect hexapods (Du et al., 2024). Embryogenesis in

these animals begins with a period of holoblastic cleavage, and certain blastomeres on

the basal (yolk) side of the blastoderm can be recognized as likely primordial germ cells

very early in development (Jura, 1967; Klag, 1982). The cytoplasm of these cells

contains granular material visible by light and electron microscopy (Klag, 1982),

resembling the cytoplasm containing germ line determinants observed in insects

(Hegner, 1914; Zissler, 1992). Rather than maintaining contact with the germ band as in

insects, collembolan germ cells migrate into the center of the yolk after their

specification until eventually migrating to associate with the abdominal mesoderm, from

which the embryonic gonad develops (Jura, 1967).

Among non-hexapod Pancrustacea, embryos of Copepoda (copepods),

Amphipoda (beach hoppers), Decapoda (crabs, lobsters and shrimp), and some

Cladocera (water fleas) undergo stereotypical *holoblastic cleavage*, allowing precise

tracking of cell lineages and the identification of a specific cleavage blastomere as the

progenitor of primordial germ cells (Gerberding et al., 2002; Hertzler, 2005; Kühn, 1913;

McClendon, 1907). In amphipods and decapods, a putative germ plasm has been

identified in the freshly laid egg, termed the RNA-containing body (Gupta and Extavour,

2013) or intracellular body (Biffis et al., 2009; Chen et al., 2021). During cleavage, this

region of cytoplasm is inherited asymmetrically by the blastomeres that will give rise to

the primordial germ cells (Gupta and Extavour, 2013) (Figure 1A2).

In chelicerates (spiders, scorpions, ticks and mites), candidate primordial germ

cells are first detectable with molecular markers in the yolk of blastoderm-stage

embryos in the mite *Tetranychus urticae* (Dearden et al., 2003). As the germ band

forms, the primordial germ cells associate with the posterior germ band near the

primordia of the fourth walking legs (Dearden et al., 2003). We are not aware of any

other chelicerate or myriapod (centipedes and millipedes) species in which primordial

germ cell specification has been proposed to occur before gastrulation.

II.ii. Category 2: PGC specification at gastrulation

After formation of the blastoderm, a portion of the blastoderm is specified as the

presumptive embryonic region, variably referred to as the *germ band*, *germ rudiment*,

embryonic rudiment, or *germ disc* depending on its initial form (Anderson, 1973;

Johannsen and Butt, 1941). The rest of the blastoderm differentiates into one or more

extraembryonic membranes (Johannsen and Butt, 1941). Soon after this specification,

gastrulation occurs, whereby some of the blastoderm cells are internalized and become

the mesoderm (Stern, 2004). In insects, mesoderm internalization primarily occurs

along the ventral midline of the germ band (Leptin, 2004; Roth, 2004). In other

arthropods, mesoderm may be internalized through a *blastopore* near the germ

rudiment (Chaw et al., 2007; Gerberding and Patel, 2004), or by epiboly (Edgar et al.,

2015). The second category of primordial germ cell differentiation occurs at or very

close to the time of gastrulation (Figure 1B).

Insects

Among insects, this timing of primordial germ cell specification is most common

among members of Hemiptera (true bugs), Psocodea (booklice), and Lepidoptera

(moths and butterflies), although it is also reported in species of Dermaptera, Blattodea

(cockroaches), Hymenoptera, Megaloptera (alderflies), Strepsiptera (twisted-wing flies),

Coleoptera, Mecoptera (scorpion flies), and Diptera. We observed three different spatial

patterns of germ cell specification around the time of gastrulation: before germ band

formation, at the posterior of the germ band after its formation, or at the ventral midline

of the blastoderm just before gastrulation. In Hemiptera, Psocodea, and Tenebrionidae

(darkling beetles; Coleoptera), germ cells are first detectable at the ventral posterior

region of the cellular blastoderm in an invaginating structure called the *posterior pit*

(Figure 1B1) (Butt, 1949; Goss, 1952; Mellanby, 1935a; Schröder, 2006; Shinji, 1919).

These cells appear to form on the side of the blastoderm epithelium facing the yolk,

rather than on the outside of the blastoderm like the pole cells described in Category 1.

In Hemiptera and Psocodea, the invagination continues into the egg towards the

anterior, forming a definitive germ band with germ cells at its posterior tip (Figure 1B1).

Germ cells in tenebrionid beetles move further inside the embryo once the amnion

(ventral extraembryonic membrane) has enclosed the entire ventral surface of the germ

band, similar to the internalization seen in coleopteran species that specify their germ

cells as pole cells (e.g. the weevils *Callosobruchus maculatus* (Brauer, 1925; Brauer,

1949; Quan, 2018) and *Acanthoscelides obtectus* (Jung, 1966; Lynch et al., 2011)).

Following their formation, the germ cells migrate along the dorsal side of the germ band

toward the anterior, eventually settling into the somatic gonad in the abdominal

mesoderm.

In several species with this timing of germ cell specification, primordial germ cells

were first identified at the posterior tip of the germ band after its detachment from the

blastoderm [*e.g.*, *Periplaneta orientalis* (Blattodea) (Heymons, 1895), *Sialis mitsuhashii*

(Megaloptera) (Suzuki et al., 1981), *Nematus ribesii* (Hymenoptera) (Singh, 1967), and

Mecoptera (Suzuki, 1990)]. Early studies of tenebrionid beetles described this pattern of

germ cell differentiation as well (Rempel and Church, 1969; Ullmann, 1964). However,

later work using molecular markers suggests that primordial germ cells in at least one

such beetle, *Tribolium castaneum*, appear earlier, during the cellular blastoderm stage

in a posterior pit much like the Hemiptera (Schröder, 2006). This raises the possibility

that in the other tenebrionid species listed above, primordial germ cells may likewise

differentiate from the blastoderm, with the reported later appearance in the germ band

simply reflecting limitations in detection methods.

In contrast to the posterior pit, germ cells differentiate along the ventral midline of

the blastoderm in Lepidoptera, separating from the blastoderm before gastrulation to lie

on the dorsal side of the presumptive germ band in the presumptive abdominal region

(Ando and Tanaka, 1980; Miya, 1958a; Presser and Rutschky, 1957; Tanaka, 1987;

Woodworth, 1889). Unlike when germ cells form at the posterior pit or at the posterior

tip of the germ band, in this case there does not appear to be long-range germ cell

migration, since the germ cells apparently differentiate in the same abdominal location

as the gonad.

Other arthropods

Among non-insect Pancrustacea, primordial germ cells are the first cells to be

internalized from the blastopore lip in Mysida (opossum shrimp) (Manton, 1928),

Anomopoda (water fleas) (Cannon, 1921), and the copepod orders Siphonostomatoida

(sea lice) (McClendon, 1907), Harpacticoida (Witschi, 1934), and Cyclopoida

(Schimkewitsch, 1896). In Isopoda (woodlice and pill bugs), primordial germ cells

differentiate from the mesendoderm soon after internalization (Figure 1B2) (Goodrich,

1939; Needham, 1942; Strömberg, 1965; Wolff, 2009). Thus, primordial germ cell

differentiation seems to be tied to gastrulation in these crustacean orders.

In chelicerates, primordial germ cells have been identified based on

morphological criteria at the blastopore in Opiliones (harvestmen) (Figure 1B3)

(Faussek, 1888; Faussek, 1892; Holm, 1947) and Scorpiones (Brauer, 1894), and a

recent study using molecular markers supported this interpretation (Gainett et al., 2022).

Putative primordial germ cells have also been identified at the blastopore in the

centipedes *Scolopendra cingulata* and *Strigamia maritima* (Myriapoda) (Green and

Akam, 2014; Heymons, 1901).

In Tardigrada (water bears), germ cells are the first cells to enter the blastopore

during gastrulation, based on data from both morphological characteristics (Hejnol and

Schnabel, 2005) and molecular markers (Heikes et al., 2023) (Figure 1B4). Lineage

tracing in these embryos suggests that the germ cells are derived from both

blastomeres of the two-cell embryo (Gabriel et al., 2007; Hejnol and Schnabel, 2005).

Finally, onychophoran (velvet worm) germ cells in different species emerge either

at the blastopore lip during gastrulation (Manton, 1949) or at the posterior end of the

germ band during elongation (Manton, 1949; Mayer and Tait, 2009; Sedgwick, 1887).

We are not aware of any studies using molecular markers to track germ cell formation in

Onychophora (Figure 1B5).

II.iii. Category 3: PGC specification from mesoderm

After gastrulation, the germ band elongates and becomes segmented, and the

mesoderm becomes organized into pairs of somites in each segment. In most living

arthropods, each somite forms a small *coelom* enclosing a *coelomic cavity* (Anderson,

1973; Johannsen and Butt, 1941). The coelomic cavities are eventually lost as the

mesoderm differentiates into different cell types and reorganizes into the primordia of

the internal organ systems, including the somatic gonad (Anderson, 1973; Johannsen

and Butt, 1941). The final category of primordial germ cell differentiation timing

encompasses species in which primordial germ cells differentiate from the mesoderm at

this comparatively late stage of embryogenesis (Figure 1C).

Insects

In insects, this category includes members of Archaeognatha (jumping

bristletails), Zygentoma (silverfish and firebrats), Orthoptera (crickets and grasshoppers)

(Figure 1C1), Mantodea (praying mantises), Blattodea, Trichoptera (caddisflies),

Coleoptera, and Lepidoptera. Primordial germ cells may differentiate before formation of

the coelomic cavities [e.g., Orthoptera (Ewen-Campen et al., 2013b), Mantodea (Görg,

1959)], after formation of the coelomic cavities [e.g., Archaeognatha (Larink, 1969),

Zygentoma (Woodland, 1957)], or after the coelomic cavities are no longer recognizable

[e.g., Trichoptera (Miyakawa, 1974), *Odontotaenius disjunctus* (Coleoptera) (Krause,

1947), *Apis mellifera* (Hymenoptera) (Dearden, 2006; Nelson, 1915)]. In this

developmental timing category, primordial germ cells are usually specified in the same

abdominal segments where the somatic gonad forms, so there is no long-range

movement of primordial germ cells after their specification.

Other arthropods

Differentiation of primordial germ cells from the mesoderm has also been

described in Leptostraca (mud shrimp) (Figure 1C2) (Manton, 1934), Anaspidacea

(mountain shrimp) (Hickman, 1936), and Branchiopoda (fairy, clam and tadpole shrimp,

and water fleas) (Anderson, 1967; Mitsumoto and Makioka, 2002; Sudler, 1899).

However, data for stages of crustacean development late enough to include

differentiation of the trunk mesoderm from which primordial germ cells might arise are

sparse. This relative scarcity of early embryonic descriptions may be in part because

many crustaceans hatch as a nauplius, comprising only the head segments (Scholtz,

2000), and germ cells might not be specified until later larval stages, after formation of

the trunk. Many studies of naupliar and larval development in crustaceans are focused

on organogenesis of the gonad as a whole, without specific reference to the origin of

primordial germ cells (Cannon, 1924; Ikuta and Makioka, 1994; Mitsumoto and Makioka,

2002).

Classical studies on myriapods suggest a late differentiation of primordial germ

cells from the mesoderm in three of the four myriapod classes [Pauropoda (Tiegs,

1947), Symphyla (pseudocentipedes) (Tiegs, 1940), and Diplopoda (millipedes) (Dohle,

1964)], although only one species was reported from each class in these studies. Germ

cells in these classes have not been investigated with molecular markers to our

knowledge.

Among chelicerates, primordial germ cells differentiate from the mesoderm of the

opisthosomal segments in spiders (Figure 1C3) (Gough, 1902; Kautzsch, 1910;

Schimkewitsch, 1903). Data from molecular markers are consistent with this

observation (Schwager et al., 2015). Primordial germ cells are first detectable in the

mesoderm of the prosoma in pycnogonids (sea spiders) (Alexeeva and Tamberg, 2021;

Alexeeva et al., 2018; Miyazaki and Makioka, 2012).

II.iv. Summary: evolutionary trends from late to early germ cell specification

In his monograph on embryogenesis in Dermaptera and Orthoptera, Richard

Heymons (Heymons, 1895) proposed that "the formation of sex cells at the rear end of

the embryonic primordium, which has now been clearly demonstrated for

representatives of very different groups, is probably generalizable for all insects." Rather

than making broad statements such as this, we summarize the data regarding the

timing of germ cell specification (Table S1) in the context of currently accepted

panarthropod phylogeny (Ballesteros et al., 2022; Fernández et al., 2018; Giribet and

Edgecombe, 2019; Misof et al., 2014; Schwentner et al., 2017) in Figures 2 and 3. This

shows that in early diverging euarthropod orders, primordial germ cells tend to be first

observed later in embryogenesis (Category 3). The timing of first germ cell appearance

evolved to be earlier in development in multiple lineages independently (Dermaptera,

Condylognatha (true bugs and thrips), and Holometabola (undergoing complete

metamorphosis) within insects; Copepoda, Malacostraca (the largest class of

crustaceans), and Cladocera within non-hexapod Pancrustacea; Chilopoda (centipedes)

within myriapods; and Opiliones, scorpions, and mites within chelicerates). These data

are consistent with the hypothesis that differentiation from the mesoderm is the

ancestral mode of primordial germ cell specification in Euarthropoda.

Conflicting interpretations in the literature about the timing of germ cell

specification have sometimes resulted from technical advances enabling identification of

germ cells at stages earlier than previously detected. For example, early work on the

silk moth *Bombyx mori* (Lepidoptera) reported primordial germ cells emerging from the

mesoderm of coelomic cavities in late embryogenesis (Category 3) (Toyama, 1902),

while later studies identified primordial germ cells on the posterior side of the ventral

midline of the early germ band (Category 2) (Miya, 1953). The onset of primordial germ

cell specification before gastrulation was later further supported by data from molecular

markers (Nakao, 1999; Nakao et al., 2006). Therefore, it is possible that species that

are currently classified in Category 3 may be reclassified in Categories 2 or 1 by future

work deploying molecular markers. This caveat notwithstanding, we note that molecular

markers and functional genetic studies often corroborate late germ cell specification

[see for example *Gryllus bimaculatus* (Orthoptera, Insecta) (Ewen-Campen et al.,

2013b), *A. mellifera* (Hymenoptera, Insecta) (Dearden, 2006), *Parasteatoda*

tepidariorum (Araneae, Chelicerata) (Schwager et al., 2015)]. We therefore do not

assume that the hypothesis of ancestral late germ cell specification is an artifact of

exclusive use of morphological identification methods.

Going forward, data from key early diverging lineages should be gathered to

further test our hypothesis about the ancestral timing of primordial germ cell

specification. Given the phylogenetic position of the Palaeoptera (mayflies, dragonflies

and damselflies) as early diverging branches of the hexapod tree (Misof et al., 2014) ,

data about primordial germ cell specification in these orders would be especially useful

to test our hypothesis with respect to hexapods, as would more information from the

ametabolous Archaeognatha and Zygentoma or the non-insect hexapod Diplura (two-

pronged bristletails). Similarly, data about primordial germ cell specification are missing

from Oligostraca, the earliest diverging clade of Pancrustacea, which includes

Ostracoda (seed shrimp) and Branchiura (fish lice).

Outside of Pancrustacea, the chelicerate phylogeny remains unstable (Sharma

and Gavish-Regev, 2025), and embryological data are too scattered for us to suggest

trends in primordial germ cell specification within this subphylum. Data regarding

primordial germ cell differentiation are even sparser for myriapods: we are aware of

relevant data from only five species across the four myriapod classes (Dohle, 1964;

Green and Akam, 2014; Heymons, 1901; Tiegs, 1940; Tiegs, 1947). Studies of

primordial germ cells in a broader range of both chelicerates and myriapods, as well as

detailed embryological studies in unexplored taxa, will therefore be needed to fully

understand the developmental context and evolutionary trends of germ cell specification

in these subphyla.

III. Molecular and genetic mechanisms of germ cell specification in

panarthropods

Beyond studying where and when germ cells first appear during arthropod

embryogenesis, researchers have sought to understand how they are specified. Germ

cell specification in animals has historically been classified into two mechanisms:

maternal inheritance and zygotic induction (Extavour and Akam, 2003; Nieuwkoop and

Sutasurya, 1981). In maternal inheritance, germ cell determinants accumulate in a

specific region of the oocyte cytoplasm, forming a germ plasm. Cells that inherit

sufficient germ plasm components after the embryo divides become primordial germ

cells. In embryos that undergo zygotic induction, all cells are in principle competent to

become germ cells at the beginning of embryogenesis. Certain cells later receive

signals from neighboring cells that induce them to take on germ cell identity. In the

following section, we will describe how historical observational data and perturbational

experimental approaches have offered increasingly robust evidence to distinguish

between these mechanisms in several panarthropod species.

III.i. Maternal inheritance

In the early twentieth century, researchers speculated that the electron-dense,

membraneless granules frequently found in the cytoplasm of primordial germ cells were

responsible for their specification. Experiments in chrysomelid beetles provided early

evidence that the posterior cytoplasm of the oocyte or early embryo contains germ cell

determinants. Burning or puncturing the posterior poles of beetle eggs with needles to

damage or remove posterior cytoplasm reduced or eliminated primordial germ cells

(Hegner, 1908; Hegner, 1911), demonstrating that this cytoplasm played a causal, and

not simply correlative, role in primordial germ cell specification. Studies in *D.*

melanogaster offered further evidence for the necessity of germ plasm. Removing or

destroying the posterior cytoplasm of *D. melanogaster* eggs via UV irradiation

demonstrated this region's requirement for germ cell formation (Geigy, 1931).

Subsequent centrifugation experiments in *D. melanogaster* embryos revealed that

shifting the granular structures in posterior cytoplasm away from the posterior pole

prevented germ cell formation or led to germ cells forming in a different part of the

embryo (Jazdowska-Zagrodzińska, 1966).

Experiments transplanting specific regions of cytoplasm between *D.*

melanogaster embryos showed that the contents of the posterior cytoplasm were

sufficient to specify germ cells. Transplanting posterior cytoplasm from undamaged *D.*

melanogaster embryos into sterile ones in which the posterior pole had been damaged

with UV irradiation rescued fertility, while transplanting anterior cytoplasm did not

(Okada et al., 1974). Further transplantation experiments demonstrated that posterior

pole plasm introduced into the anterior of another embryo caused formation of ectopic

pole cells that gave rise to functional germ cells when transplanted again to the

posterior of a third, sterile embryo (Illmensee and Mahowald, 1974). Similar

experiments transferring the posterior cytoplasm from late-stage oocytes into embryos

revealed that the oocyte pole plasm could specify functional primordial germ cells

(Illmensee et al., 1976). This result revealed that the necessary germ cell determinants

in *D. melanogaster* are deposited during oogenesis. These foundational studies

demonstrated that maternally derived cytoplasmic granules are both necessary and

sufficient for germ cell formation in some species.

Contemporary studies testing for maternal inheritance often ask whether known

germ line gene products accumulate during late oogenesis or early embryogenesis in

the region where primordial germ cells first arise. For example, antibody staining or *in*

situ hybridization experiments to detect germ line molecules like Vasa protein or *nanos*

mRNA have suggested the presence of germ plasm in mosquitoes (Juhn and James,

2006; Juhn et al., 2008), scuttle flies (Wotton et al., 2014), ants (Khila and Abouheif,

2008; Lynch et al., 2011; Rafiqi et al., 2020), wasps (Kharel et al., 2024; Lynch et al.,

2011), weevils (Lynch et al., 2011), aphids (Chang et al., 2006; Chang et al., 2007;

Chen et al., 2009; Lin et al., 2014), amphipods (Gupta and Extavour, 2013), and prawns

(Chen et al., 2021). Transcriptomic analyses of different portions of embryos and

oocytes (Carter et al., 2015; Quan and Lynch, 2016; Yoon et al., 2019) have also

revealed asymmetric localization of molecules associated with primordial germ cell

specification.

However, expression data showing asymmetric localization of putative germ

plasm components are not sufficient to demonstrate that these molecules function in

primordial germ cell specification. For this, functional genetic techniques are needed to

manipulate gene activity and test the effect on germ plasm assembly and germ line

development. The function of germ plasm components in panarthropods has been most

thoroughly studied in *D. melanogaster* (Lehmann, 2016; Mahowald, 2001; Trcek and

Lehmann, 2019). The membraneless ribonucleoprotein granules at the posterior of *D.*

melanogaster oocytes and early embryos are nucleated by the protein Oskar, which is

both necessary and sufficient for germ plasm assembly (Ephrussi and Lehmann, 1992;

Kistler et al., 2018). Oskar-mediated localization of core germ line molecules such as

Vasa, Tudor, Aubergine, and *nanos* to germ granules is essential for primordial germ

cell specification and maintenance of germ cell identity during cell migration and gonad

formation in these fruit flies (Lehmann, 2016). The roles of these germ plasm molecules

in establishing the *D. melanogaster* germ line have been extensively reviewed

elsewhere (Lehmann, 2016; Mahowald, 2001).

Analysis of germ plasm composition and localization in species beyond *D.*

melanogaster has revealed both conserved and rapidly evolving features. Within the

Drosophila genus, germ granule size, morphology (Counce, 1963; Mahowald, 1968;

Rivard et al., 2025), and the relative proportions of different granule components vary

widely (Doyle et al., 2023). The wasp *N. vitripennis* also has a well-characterized germ

plasm in the form of a single large structure called the oosome (Bull, 1982). Like *D.*

melanogaster, N. vitripennis requires *oskar* to assemble a germ plasm containing many

of the same molecules (e.g., Vasa, Tudor, and *nanos*) (Lynch and Desplan, 2010;

Lynch et al., 2011). However, while *D. melanogaster* has many small germ granules

(Mahowald, 1962), the *N. vitripennis* oosome has a distinct protein organization and

transcript composition compared to that observed in *D. melanogaster* (Kemph et al.,

2025; Kharel et al., 2024; Lynch et al., 2011). In several crustacean species, the germ

plasm appears as a large, round granule in the cortex of embryos and contains

conserved germ plasm markers (Chen et al., 2021; Grattan et al., 2013; Gupta and

Extavour, 2013). Studying germ plasm in additional species will be needed to clarify

whether and how variation in germ plasm makeup and organization influences the post-

transcriptional regulation and function of germ line determinants.

III.ii. Zygotic induction

Experimentally demonstrating inductive germ cell specification is more

technically challenging than detecting germ plasm accumulation. Consequently, most

studies have inferred or hypothesized zygotic induction based on absence of evidence

of germ plasm. For example, *in situ* hybridization experiments fail to reveal asymmetric

localization of putative germ line gene transcripts in oocytes and early embryos of

various species, including milkweed bugs (Ewen-Campen et al., 2013a), crickets

(Ewen-Campen et al., 2013b), honeybees (Dearden, 2006), mites (Dearden et al.,

2003), spiders (Schwager et al., 2015), centipedes (Green and Akam, 2014), and

tardigrades (Heikes et al., 2023). Instead, key germ line genes are expressed in a

subset of cells later in embryogenesis, coincident with reports of the first primordial

germ cells based on morphological criteria (see Part II, Table S1). However, we note

that even in the well-accepted case of mouse primordial germ cell induction (Saitou and

Yamaji, 2010), it remains formally possible that some unknown, and therefore untested,

germ line determinant, or factor that makes cells competent to respond to later germ

line induction signals, is asymmetrically localized and preferentially inherited by germ

cells.

This caveat notwithstanding, there are multiple examples of experimental support

for induction rather than inheritance across panarthropods. Maternal knockdown of

putative germ plasm components in milkweed bugs and crickets reveals the

dispensability of maternal products of these genes for primordial germ cell formation

(Ewen-Campen et al., 2013a; Ewen-Campen et al., 2013b). Laser ablation of single

cells in two and four cell stage embryos of the tardigrade *Thuliana stephaniae* does not

prevent primordial germ cell specification, also suggesting that this species lacks a

cytoplasmic region with inherited germ line determinants (Hejnol and Schnabel, 2005).

Observations of germ cell formation in embryos with tissue duplications can be

interpreted as evidence against germ plasm and consistent with induction. For example,

in the drain fly *Clogmia albipunctata* and the flour beetle *T. castaneum*, genetic

manipulations producing double abdomen embryos lead to germ cell formation at both

anterior and posterior ends (Ansari et al., 2018; Yoon et al., 2019). Similarly, naturally

occurring "twins" of the stick insect *Carausius morosus* are duplicated embryos fused

along their anterior–posterior axis, that possess germ cells (Cavallin, 1971). In contrast,

double abdomen embryos of *D. melanogaster* only produce germ cells at the true

posterior, since germ plasm is absent at the ectopic posterior end (Lasko and

Ashburner, 1990).

Among panarthropods, experiments in the cricket *G. bimaculatus* have provided

the most direct evidence for zygotic induction of primordial germ cells. Knockdown of

the mesoderm specification factor *twist* leads to loss of primordial germ cells,

suggesting that mesoderm contributes to germ cell specification and/or maintenance

(Ewen-Campen et al., 2013b) and consistent with morphological evidence that

orthopteran primordial germ cells first arise from the embryonic mesoderm (Wheeler,

1893). The bone morphogenetic protein (BMP) signaling pathway, known to induce

germ cells in mice (Hopf et al., 2011; Lawson et al., 1999; Ying and Zhao, 2001; Ying et

al., 2000), is also required for cricket germ cell specification. Specifically, BMP signaling

and the downstream transcription factor Blimp-1 induce a subset of mesodermal cells to

become primordial germ cells in crickets (Donoughe et al., 2014; Nakamura and

Extavour, 2016).

The shared use of both BMP and Blimp-1 in mice and crickets may have

convergently evolved, or could suggest that this signaling mechanism is ancestral in

bilaterian inductive primordial germ cell specification (Lochab and Extavour, 2017).

Determining whether BMP and its downstream target genes are required for germ line

specification in more species will be necessary to distinguish between these

possibilities. Surprisingly few signaling pathways contribute to developmental fate

decisions in animals (Pires-daSilva and Sommer, 2003), increasing the likelihood of

evolutionary convergence, and limiting the number of candidates to evaluate for roles in

primordial germ cell induction in different species.

III.iii. Intermediate and flexible strategies of germ cell specification

Evidence from some animal species suggests that there may not be a strict

dichotomy between inheritance and induction (discussed by Seervai and Wessel, 2013).

These mechanisms could instead represent two ends of a spectrum, and they may not

be mutually exclusive. Below we discuss cases of panarthropod species for which data

suggest that their mechanisms of germ cell specification may fall somewhere along this

continuum.

In the silk moth *B. mori*, historical studies have generated conflicting evidence

about how germ cells are specified. Germ granules were not detected in early light

microscopy experiments, but germ cells failed to form after cauterization of the ventral

side of cleavage stage embryos, suggesting this region could contain germ cell

determinants (Miya, 1958a). Subsequent studies found that Vasa protein is not

asymmetrically localized before cellularization, again suggesting a lack of germ plasm

(Nakao et al., 2006). However, the mRNA of one of four *B. mori nanos* homologs,

nanosO, localizes along the ventral midline of syncytial embryos immediately following

egg-laying, and is inherited by the primordial germ cells (Nakao et al., 2008).

Intriguingly, researchers found that maternal and zygotic *nanosO* act redundantly in

primordial germ cell specification; loss of one or the other reduces but does not

completely eliminate germ cells, whereas loss of both prevents germ cell formation

(Nakao and Takasu, 2019). The possibility that both inheritance and induction

mechanisms contribute to *B. mori* primordial germ cell specification suggests that this

species could offer insight into transitions in germ cell specification mechanisms within

Lepidoptera or in insects more broadly.

Although *D. melanogaster* is traditionally characterized as a species using a

germ plasm, zygotic signaling was recently suggested to contribute to germ cell

specification in this species (Colonnetta et al., 2022). Loss-of-function mutations in the

BMP family morphogen Dpp lead to partial loss of transcriptional quiescence and

mislocalization of germ plasm in the newly formed pole cells (Colonnetta et al., 2022).

Furthermore, loss of BMP signaling leads to activation of the normally suppressed

terminal signaling pathway in pole cells, which in turn disrupts their migration

(Colonnetta et al., 2022). These observations illustrate a universal conceptual challenge

in the field of developmental biology: distinguishing between, or even consistently

defining, cell type specification and maintenance. Although pole cells can form upon

disruption of BMP signaling, they do not contain the proper molecular components to

act as primordial germ cells (Colonnetta et al., 2022). We suggest it may be most useful

to define primordial germ cell specification mechanisms as those that ensure both the

physical formation of germ cell precursors (including adoption of specific morphologies

and activation of characteristic gene regulatory states), and the competence of those

cells to take the next developmental steps toward joining the gonad. Under this

definition, we would indeed have to concede that both germ plasm and zygotic inductive

mechanisms play roles in *D. melanogaster* germ cell specification.

The ability to post-embryonically regenerate a germ line initially specified via

maternal inheritance has also been reported in multiple invertebrate species, including

the sea squirt *Ciona intestinalis* (Takamura et al., 2002), the earthworm *Enchytraeus*

japonensis (Tadokoro et al., 2006), and the marine worm *Capitella teleta* (Dannenberg

and Seaver, 2018) (see also Özpolat, 2024 for a review of this phenomenon in

Annelida). The wasp *Pimpla turionellae* (Hymenoptera) has an oosome (Meng, 1968)

required for germ cell specification: damaging this material results in no pole cells,

which are the primordial germ cells of this species (Achtelig and Krause, 1971).

However, oosome-deficient embryos lacking pole cells develop into larvae with germ

cells in their gonad, indicating that an additional mechanism must regenerate germ cells

later in development (Achtelig and Krause, 1971). Evidence of germ line regeneration

has also been reported in the amphipod *Parhyale hawaiensis* (Kaczmarczyk, 2014) ,

which has a germ plasm that is asymmetrically segregated to the *g* micromere and its

lineage (Extavour, 2005; Gerberding et al., 2002; Gupta and Extavour, 2013). Ablation

of the *g* micromere eliminates primordial germ cells in the embryo (Extavour, 2005), but

the germ line is replaced post-embryonically with germ cells that appear to derive from

the mesoderm (Kaczmarczyk, 2014). The authors of this study speculated that these

replacement germ cells specifically derive from cells of the somatic gonad

(Kaczmarczyk, 2014), reminiscent of a theory popular for the first half of the 20th

century, that primordial germ cells in most animals derived from the inner epithelium of

the somatic gonad (Everett, 1944; Hargitt, 1919; Heys, 1931). Species capable of

regenerating primordial germ cells present the opportunity to investigate the relative

contributions of and potential interactions between inheritance- and induction-based

mechanisms in a single species. Further, these species may represent living examples

of the intermediates that are hypothesized to exist in evolutionary transitions between

modes of germ line specification (Extavour, 2007).

IV. Evolutionary drivers of panarthropod germ cell specification

In the following section, we discuss the evolutionary dynamics of germ cell specification

mechanisms across Panarthropoda, and propose potential developmental and genetic

factors that might contribute to these dynamics.

IV.i. Convergent evolution of panarthropod germ plasm

The phylogenetic distribution of germ cell specification mechanisms across

panarthropods reveals a complex evolutionary history. Data for tardigrades,

chelicerates, and myriapods are consistent with induction as the ancestral mechanism

of panarthropod primordial germ cell specification (Figure 3). Pancrustacea are the most

heavily represented group in our dataset (Table S1). The late origin of primordial germ

cells in several species suggests induction may be widely used, but there appear to be

independent origins of germ plasm in Malacostraca and Branchiopoda. The use of

induction and germ plasm in different decapod species suggests that further sampling

could reveal patterns in the evolution of germ cell formation within this order (Biffis et al.,

2009; Chen et al., 2021; Terao, 1929). However, functional experiments and broader

taxonomic sampling will be necessary to confirm mechanisms and reconstruct their

evolutionary trajectories across Panarthropoda as a whole.

Reminiscent of a broader pattern we and others previously reported across

animals (Extavour, 2007; Extavour and Akam, 2003; Johnson et al., 2003; Whittle and

Extavour, 2017), in some insect orders we noted that early diverging lineages deployed

induction, while later branching lineages appeared to use germ plasm (Figure 2, 3). This

pattern is well exemplified in Coleoptera, (Kobayashi et al., 2006; Komatsu and

Kobayashi, 2012; Krause, 1947; Niikura et al., 2017; Schröder, 2006; Ullmann, 1964),

with pole cells forming only in Chrysomeloidea (including long-horned and leaf beetles)

and Curculionidae (Brauer, 1925; Hegner, 1908; Lynch et al., 2011). These

observations raise the possibility that germ plasm may not be ancestral to Coleoptera.

When considered alongside the absence of germ plasm in most Lepidoptera, the

observation of late-arising, Category 3 germ cells in Trichoptera (Miyakawa, 1974) and

Category 2 germ cells in Mecoptera (Suzuki, 1990), and the distinct morphologies of *N.*

vitripennis and *D. melanogaster* germ plasm (see Part III.i), these patterns are

consistent with a novel hypothesis in which germ plasm evolved independently multiple

times across insects, rather than only once in the Holometabola as we and others have

previously proposed (Ewen-Campen et al., 2012; Ewen-Campen et al., 2013b; Lynch et

al., 2011). Our previous work had suggested the independent evolution of germ plasm

at the phylum level (Extavour and Akam, 2003), but this new hypothesis is the first, to

our knowledge, to suggest such widespread convergent evolution of maternal

inheritance-based germ cell specification within a phylum.

To test this hypothesis, we performed ancestral character reconstruction using

both maximum parsimony and maximum likelihood approaches. For this analysis, we

designated species as using inheritance or induction based on experimental evidence

when available or inferred from observational data describing the timing and location of

germ cell specification (Supplementary Materials, Table S1). The results of a parsimony

analysis inferred that the ancestral panarthropod specification mode was induction, with

a minimum of 24 changes over the history of the panarthropods (Figure S1; Table S3).

When considering all equally parsimonious character histories, changes from induction

to maternal inheritance (min = 18, max = 23, average = 20.5) were the most common,

but reverse transitions were also inferred (min = 2, max = 7, average = 4.5). At a

minimum, reversions from maternal inheritance to induction were inferred in the

honeybee *A. mellifera* (Hymenoptera) and the drain fly *C. albipunctata* (Diptera). We

note that all maximum parsimony reconstructions assign induction as the ancestral state

of Holometabola (Figure S1, node 11), which requires many fewer total transitions

within Holometabola (N = 12) than would be required if germ plasm evolved a single

time in the last common ancestor of Holometabola (N = 20) as previously proposed

(Ewen-Campen et al., 2013b; Lynch et al., 2011) .

Marginal ancestral reconstructions of a maximum likelihood approach also

proved consistent with the parsimony analysis described above (see Table S3 for a

comparison), again providing strong statistical support for induction as the ancestral

specification mechanism in Panarthropoda, Arthropoda, and Insecta, with multiple

independent instances of germ plasm evolution and reversion to induction (Figure S2;

see Table S7 for marginal probabilities). We further find strong evidence that induction

was the ancestral specification mechanism in Holometabola (probability 80.6%),

followed by independent transitions to germ plasm in the common ancestors of Diptera

and Mecoptera (or independently within Mecoptera, Siphonaptera (fleas), and Diptera),

derived lineages of Lepidoptera, in the common ancestor of Chrysomelidae and

Curculionidae (Coleoptera), and in the common ancestor of Hymenoptera. Outside of

Holometabola, we infer independent transitions to germ plasm in Thysanoptera,

Hemiptera, Dermaptera, Collembola (which may have undergone multiple independent

transitions within the class), Decapoda, the last common ancestor of Hyalidae and

Talitridae (Amphipoda), and in the chelicerate *Tetranychus urticae*. Collectively, these

qualitative and quantitative analyses suggest a previously unappreciated dynamism in

the evolutionary history of germ cell specification within Panarthropoda.

IV.ii. The relationship between germ cells and somatic germ layers

In arthropods with late formation of primordial germ cells (Category 3), primordial

germ cells appear to differentiate directly from the mesoderm (Evans, 1901; Hickman,

1936; Nelson, 1915). Consistent with this hypothesis, abolishing mesoderm prevents

germ cell specification in the cricket *G. bimaculatus* (Ewen-Campen et al., 2013b).

Germ cell differentiation occurs concurrently with the differentiation of many other

mesodermal cell types, including the somatic gonad, musculature, heart, and fat body

(Johannsen and Butt, 1941). We suggest that the types of mechanisms that specify

germ cells are not distinct in principle from those specifying any other mesodermal cell

type in these animals.

The germ layer origin of germ cells is less clear in arthropods whose germ cells

appear before or very close to the time of gastrulation (Category 2). For example,

primordial germ cells in many crustaceans are described as being the first to enter the

blastopore during gastrulation (e.g. Cannon, 1921; Goodrich, 1939; Manton, 1928), and

are usually considered by researchers to be derived from the mesendoderm. In

contrast, germ cells in Category 2 insects arise before gastrulation (Mellanby, 1935b;

Ullmann, 1964), consistent with the hypothesis that germ cells are distinct from the

mesoderm and may not derive from any germ layer.

In crustaceans in which a putative germ plasm has been identified, cleavage

patterns are stereotypical, and cell lineages are available, the germ cell lineage is sister

to the mesoderm or the mesendoderm [Amphipoda (Extavour, 2005; Gerberding et al.,

2002; Scholtz and Wolff, 2002; Wolff and Scholtz, 2002), Euphausiacea (krill) (Alwes

and Scholtz, 2014), Decapoda (Hertzler, 2005)], but never to the ectoderm. In addition,

removing the putative germ plasm from embryos of the amphipod *P. hawaiensis* makes

the *g* micromere morphologically indistinguishable from the other micromeres, which

give rise to mesoderm or endoderm in unperturbed embryos (Gupta and Extavour,

2013). This result suggests that in the absence of germ cell determinants, the *g*

micromere, normally the germ line progenitor, could instead adopt mesendodermal fate.

This hypothesis has not been directly tested in this amphipod because gastrulation fails

after ablation of the putative germ plasm (Gupta and Extavour, 2013). Together, these

data suggest a close association between the germ cell and mesendodermal lineages in

these crustaceans. This relationship is apparent beyond Pancrustacea, since germ cells

induced late in embryogenesis arise from the mesoderm (Category 3). We thus

speculate that germ cells in panarthropods may be ancestrally derived from the

mesodermal lineage, consistent with our inference that inductive germ cell specification

is ancestral to panarthropods (see Part IV.i above).

IV.iii. Evolutionary shifts in the spatial and temporal origin of primordial germ

cells

We have seen significant variation in the spatial and temporal origin of primordial

germ cells across panarthropods (Figures 2–3; Table S1). However, regardless of when

primordial germ cells first appear, we find a common stage across many panarthropods

in which primordial germ cells are situated in bilateral segmental clusters across

multiple trunk segments. Soon after this stage, the segmental clusters on each side fuse

together into bilaterally symmetrical genital ridges, corresponding to the primordial

embryonic gonad (Miya, 1958b; Nelsen, 1934; Seidel, 1924; Ullmann, 1964).

Most arthropods undergo "short germ" development, wherein the blastoderm

tissue generates only anterior segments, with posterior segments being added

sequentially to the posterior end of the germ band later in development (Krause, 1939) .

In contrast, the "long germ" mode of development seen in *D. melanogaster*, wherein the

blastoderm tissue immediately generates all body segments, is an evolutionary novelty

found only in some insects (Clark et al., 2019; Krause, 1939). In Category 1 or 2

arthropods with short germ development, germ cells are often specified before the

formation of the segments that will include the future embryonic gonad. In Category 2

insects, germ cells often associate with the posterior end of the germ band and then

migrate toward the anterior as the germ band elongates (Goss, 1953; Hegner, 1909b;

Heming and Huebner, 1994; Paz, 1958; Suzuki, 1990; Ullmann, 1964). This migratory

path is so common among insects that Johanssen and Butt included it in their

description of embryogenesis in a stereotypical insect (Johannsen and Butt, 1941).

The association of germ cells with the posterior end of the embryo may have had

other evolutionary consequences in insects. Many insects, both long and short germ,

specify the body plan early in development via maternal deposition of axial patterning

determinants, independent of the mechanism of germ cell specification (reviewed in

Lynch, 2019; Rosenberg et al., 2009). For example, the short germ grasshopper

Schistocerca americana exhibits asymmetric localization of the posterior patterning

determinant *nanos* in the absence of a germ plasm (Lall et al., 2003). However, insects

with germ plasm that includes axial patterning determinants, like *D. melanogaster*, often

exhibit long germ development. Based on these observations, we employed

phylogenetic logistic regression and Pagel's test of evolutionary rate covariation (Pagel,

1994) to test the hypothesis that the acquisition of long germ band development

influences germ cell mechanism transitions or vice versa, and found statistically

significant evidence of this association using both methods (Figure S3; Figure S4). All

Pagel covariation models assuming coupled evolutionary rates among these two

characters are significantly better fits to the data than the null model of independent

evolution (likelihood ratio test p-value < 1e−3; Figure S4; Table S8). We note that the

bidirectional and GB-dependent models perform similarly to the GC-dependent model,

suggesting both that long germ band development may drive the evolution of germ

plasm, and that germ plasm may drive transitions to long germ band development.

We further hypothesize that a spatial shift in germ cell specification to the

posterior end of the embryo in Category 2 insects created a condition in which germ line

determinants are localized to the same embryonic region as posterior patterning

determinants (Figure 4B). Subsequently, germ line determinants may have hitchhiked

onto the asymmetric localization machinery, coupling germ cell specification to posterior

patterning and thus promoting the evolution of germ plasm (Figure 4C). Under this

hypothesis, the germ plasm of *D. melanogaster* would represent the final stage of this

evolutionary scenario, as its germ granules include both germ line and abdominal

patterning determinants (Ephrussi and Lehmann, 1992; Trcek et al., 2015; Wang and

Lehmann, 1991).

N. vitripennis is an intriguing case to study the relationship between

anteroposterior patterning and germ cell specification. The posterior cytoplasm of this

wasp has two populations of *nanos* mRNA: one that is incorporated into the oosome

and eventually into pole cells, and another that functions in axial patterning (Lynch and

Desplan, 2010). *nanos* localization to the oocyte posterior in this wasp depends on

actin-dependent anchoring (Olesnicky and Desplan, 2007), just as in *D. melanogaster*

(Forrest and Gavis, 2003), suggesting a conserved localization mechanism between

these two species. We speculate that *N. vitripennis* represents an evolutionary

intermediate between species where anteroposterior patterning is entirely independent

of germ cell formation (Figure 4A) and those where these two processes are fully linked

(Figure 4C). Our evolutionary hypothesis could be tested further by examining the

mechanisms of localization of posterior patterning determinants and germ line

determinants in more insect species to understand how existing cytoskeletal machinery

for one developmental process may have been coopted for use in another.

IV.iv. Potential genetic basis of evolutionary shifts in the molecular mechanism of

primordial germ cell specification

777   Beyond the timing and position of germ cell origin, evolutionary changes in the

localization of gene products and timing of gene expression may also underlie

transitions in the mechanism of specification (Extavour, 2007; Srouji and Extavour,

2010). Many of the genes important for specifying germ cells are conserved across

metazoans (Ewen-Campen et al., 2010), but they exhibit distinct temporal and spatial

expression patterns across species. In species that use inheritance, the germ line gene

products are expressed during oogenesis and persist in the early embryonic cytoplasm

to be inherited by the primordial germ cells. In species that induce germ cells, these

genes are either not expressed in oocytes or the gene products are neither

asymmetrically nor preferentially inherited by particular cells. As we have suggested

previously (Extavour, 2007; Srouji and Extavour, 2010), relatively simple regulatory

modifications affecting post-transcriptional regulation, the cytoskeleton, or RNA or

protein localization could cause germ line gene product retention in oocytes, thereby

enabling the transition from induction to inheritance.

The insect-specific gene *oskar* offers a potential example of gene network

evolution driving shifts in germ cell specification mechanism. Since *oskar* is likely a

nucleator of germ plasm in multiple insects (Ephrussi et al., 1991; Jones and

Macdonald, 2007; Kemph and Lynch, 2022; Lynch et al., 2011; Rivard et al., 2025) and

has been lost multiple times since its origin prior to the divergence of Pterogyta from

Ametabola (Blondel et al., 2021), we hypothesized that *oskar* gene presence in the

genome might be correlated with germ line specification mode in a given taxon (Figure

S3). A related hypothesis, that the evolutionary advent of the *oskar* gene facilitated the

evolution of holometabolous germ plasm, was previously proposed based on the co-

occurrence of *oskar* transcript and germ plasm in *N. vitripennis* and *D. melanogaster*, as

well as the absence of both *oskar* homologs and germ plasm in *T. castaneum*, *A.*

mellifera, *C. albipunctata*, most Lepidoptera, and the hemimetabolous genomic

resources available at the time (Lynch et al., 2011). The results of applying phylogenetic

logistic regression to this hypothesis (Figure S3; Table S9) suggest a strong and

statistically significant association between *oskar* presence and maternal inheritance

(coefficient = 2.5827, adjusted odds-ratio: = 13.28, p-value = 0.00059) We emphasize

that this observed association does not imply that *oskar* is necessary or sufficient for

germ plasm formation; indeed, there are insect species without *oskar* that nevertheless

have germ plasm and species with *oskar* that lack germ plasm (Figure S3), as we

discuss in Part IV.v below.

In contrast to prior hypotheses from our group and others of a single origin of

germ plasm at the base of the Holometabola (Ewen-Campen et al., 2012; Ewen-

Campen et al., 2013b; Kemph and Lynch, 2022; Lynch et al., 2011), our current

likelihood and parsimony reconstruction of induction as ancestral to Holometabola

(Figure 5; Figures S1–S2) implies that *oskar* has been independently recruited as a

germ plasm nucleator at least twice, in Hymenoptera (Lynch et al., 2011) and in Diptera

(Ephrussi and Lehmann, 1992). The strong association between *oskar* presence and

germ plasm (Figure S3) implies that it may have been co-opted for germ plasm

nucleation even more times across insects. We note that such "gene reuse" is a

common feature of convergent evolution across multiple biological processes (Chan et

al., 2010; Martin and Orgogozo, 2013; Morris, 2009; Reed et al., 2011; Satterlee et al.,

2024). Since *oskar* is co-expressed with germ plasm components in nervous systems of

hemimetabolous and holometabolous insects (Ewen-Campen et al., 2012; Kulkarni et

al., 2023; Xu et al., 2013), we posit that independent acquisitions of germ plasm in

distinct lineages may have proceeded via parallel cooption of a pre-existing gene

interaction network. Future studies should aim to determine whether each independent

origin of germ plasm in the Holometabola arose by co-option of *oskar* to the germ line,

or by *oskar*-independent mechanisms of germ plasm assembly.

The results of our analysis (Figure S3; Table S9) suggest that *oskar*'s functional

evolution may have contributed to a transition from induction to inheritance, but

precisely when and how *oskar* acquired a germ line role remains unclear. Morphological

data suggest that thrips use germ plasm (Haga, 1985; Heming, 1979), and *oskar*

homologs have been found in this order (Blondel et al., 2021). If *oskar* plays a role in

thrips germ cell specification, it could support the entirely novel hypothesis that *oskar*

evolved its germ line determinant properties before the evolution of complete insect

metamorphosis.

IV.v. The case for novel panarthropod germ plasm nucleators

Recruitment of *oskar* to the germ line is not the only mechanism explaining the

evolution of germ plasm in arthropods. In the genomes of species of Carabidae (ground

beetles), an early diverging beetle family that lacks pole cells, *oskar* orthologs have

been identified, decoupling the presence of *oskar* from germ plasm in Holometabola

(Blondel et al., 2021; Kobayashi et al., 2013). Similarly, the genome of the

hemimetabolous pea aphid *Acyrthosiphon pisum* lacks an *oskar* ortholog but their

embryos have a germ plasm containing Vasa and Nanos proteins (Chang et al., 2006;

Chen et al., 2009; Lin et al., 2014), which are asymmetrically localized by unknown

mechanisms. Vasa is also asymmetrically localized to a putative germ plasm in

amphipod and decapod species (Chen et al., 2021; Gupta and Extavour, 2013), which

is also likely to have evolved via distinct mechanisms since *oskar* is an insect-specific

gene (Blondel et al., 2021). Outside of arthropods, germ plasm nucleators have

convergently evolved in diverse animal phyla (Aoki et al., 2016; Bontems et al., 2009;

Kulkarni and Extavour, 2017; Lee et al., 2020; Marlow and Mullins, 2008; Scholl et al.,

2024). Similarly, we posit that germ plasm nucleators other than *oskar* must also exist

across the panarthropods, awaiting discovery.

V. Concluding thoughts

By reinterpreting the meticulous work of generations of classical embryologists

through the lens of modern developmental genetics and an improved panarthropod

phylogeny, we have generated novel, testable evolutionary hypotheses about the

mechanisms that specify the germ line, with implications for the genetic basis of

convergence in evolution that extend beyond this superphylum. The advent of new

genetic tools in an increasing number of panarthropod model species places us in an

exciting era to tackle age-old questions about the evolution of panarthropod germ cell

specification, as well as the impact of these mechanisms of genome evolution and body

patterning in the most successful group of animals on earth.

Acknowledgements

We thank members of the Extavour Lab for helpful discussions, Kira Heikes for the image of tardigrade germ cells (Figure 1B4), and Prashant Sharma for the image of harvestmen germ cells (Figure 1B3).

Data Availability

All data sources are listed in the accompanying Supplementary Material and/or are available at

https://github.com/rishabhrajkapoor/panarthropoda_gc_specification_evolution, commit ID ff9945c.

Author Contributions

All authors gathered data and performed data analysis and interpretation. JAK, ELR and RRK wrote the first draft of the manuscript. CGE conceived of the study, obtained funding for the study, supervised its execution, and reviewed and edited the manuscript.

Funding

This study was supported by NSF award IOS-2220747, funds from Harvard University, and from the Howard Hughes Medical Institute awarded to CGE. Additional support was provided by the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number #1764269), the Harvard Quantitative Biology Initiative, a Herchel Smith Graduate Fellowship awarded to ELR, and an NSF Graduate Research Training Fellowship to RRK. CGE is an Investigator of the Howard Hughes Medical Institute.

Conflicts of Interest

The authors declare no conflicts of interests.

Figure 1 (previous page). Primordial germ cell origin in the context of panarthropod
embryogenesis. In all schematics, magenta represents the approximate location of germ plasm or
primordial germ cells; gray represents yolk or extraembryonic tissue, blue box in schematic shows the
stage/location where germ cells are detectable in the micrographs and drawings on the right. (**A1**)
Schematic: In the fruit fly *Drosophila melanogaster*, germ plasm accumulates at the posterior of syncytial
embryos and is inherited by pole cells that migrate to the ultimate site of the embryonic gonad (Illmensee
and Mahowald, 1974). Data: TEM micrograph displaying pole cells budding in *D. melanogaster* embryos.
The pole cell nuclei (N) have characteristically diffuse chromatin, and polar granules (P) are distributed
throughout the cytoplasm. Image reproduced with permission from (Mahowald, 1968). M: mitochondria;
N: nuclei; P: polar granules. (**A2**) Schematic: In the amphipod *Parhyale hawaiensis*, germ plasm
accumulates in a region of the embryo called the RNA-containing body, which is inherited by the *g*
micromere at the 8-cell stage (Extavour, 2005; Gupta and Extavour, 2013). All descendants of *g* are
primordial germ cells. Data: Primordial germ cells visualized by immunohistochemistry with a cross-
reactive antibody raised against recombinant Vasa protein from both *D. melanogaster* and the
grasshopper *Schistocerca americana* (Chang et al., 2002). (**B1**) Schematic: In the milkweed bug
Oncopeltus fasciatus, primordial germ cells first appear in the ventral posterior blastoderm in an
invaginating structure called the posterior pit (Ewen-Campen et al., 2013a). Primordial germ cells remain
at the posterior tip of the germ band as it invaginates into the yolk. The germ cells migrate along the germ
band toward the anterior to segments A4–A6. Data: Hand drawing of a light micrograph of a sectioned
late blastoderm stage *O. fasciatus* embryo showing germ cells in the posterior pit . Image reproduced with
permission from (Butt, 1949). gb: germ band; gc: germ cell. (**B2**) Schematic: In the isopod *Limnoria
lignorum*, primordial germ cells are first identifiable among the mesendodermal mass just after the
beginning of gastrulation (Strömberg, 1968). Primordial germ cells remain associated with the mesoderm
throughout development. Data: Light micrograph of a sectioned *L. lignorum* embryo showing primordial
germ cells distinguished by their lighter-staining cytoplasm and larger nuclei among the mesoderm at the
extended germ band stage. Image reproduced with permission from (Strömberg, 1968). vi: vitellophage;
gen ru: genital rudiment. (**B3**) Schematic: In the harvestman *Phalangium opilio*, primordial germ cells are
first identifiable at the posterior end of the germ band soon after the germ band forms (Faussek, 1888;
Gainett et al., 2022). Primordial germ cells remain at the posterior end of the germ band at least until the
fourth opisthosomal segment is formed, after which the germ cells can be found in that segment. Data: ;
Chromatin (visualized with the nucleic acid dye Hoechst 33342) is more diffuse in primordial germ cells
than in surrounding cells. Image kindly provided by Prashant Sharma (University of Wisconsin, Madison).
Green: alpha-tubulin. (**B4**) Schematic: In the tardigrade *Hypsibius exemplaris*, primordial germ cells are
first identifiable in the 32-cell stage and are the first cells to be internalized during gastrulation. Data:
Primordial germ cells visualized by *in situ* hybridization against *He-vasa*. Image kindly provided by Kira
Heikes (Duke University). (**B5**) Schematic: In the velvet worm *Peripatopsis moseleyi*, primordial germ
cells are first identifiable near the blastopore at the beginning of gastrulation. As the embryo elongates
and more segments form, the primordial germ cells migrate to more anterior segments. Data: Hand
drawing of a light micrograph of a *P. moseleyi* embryo showing primordial germ cells soon after their
internalization during gastrulation, distinguished by their lighter-staining nuclei and cytoplasm. Image
reproduced with permission from (Manton, 1949). g: germ cells; en: endoderm; b.a.: blastoporal area.
(**C1**) Schematic: In the cricket *Gryllus bimaculatus*, primordial germ cells are first detectable among the
mesoderm in segments A2–A4 at embryonic stage 6, after most abdominal segments have been
specified (Ewen-Campen et al., 2013b). Data: Primordial germ cells visualized by immunohistochemistry
with an antibody against *Gb*-Piwi. (**C2**) Schematic: In the leptostracan *Nebalia bipes*, primordial germ
cells are first identifiable in the mesoderm in the thorax and abdomen during the extended germ band
stage (Manton, 1934). Data: hand drawing of a light micrograph of a sectioned *N. bipes* embryo showing
primordial germ cells among the mesoderm in the extended germ band, distinguished by their lighter-
staining cytoplasm and larger nuclei. Image reproduced with permission from (Manton, 1934). G: genital
rudiment; l.c.m.: circular muscle round liver lobe; th.8: eighth thoracic segment. (**C3**) Schematic: In the
house spider *Parasteatoda tepidariorum*, primordial germ cells are first identifiable in segmental clusters
in the opisthosomal mesoderm after all segments have formed (Schwager et al., 2015). Data: Primordial
germ cells visualized by immunohistochemistry using an antibody raised against *Pt*-Vasa.

Clade

Collembola
Ametabola
Hemimetabola
Holometabola

Insecta

Timing category

■ Category 1
■ Category 2
■ Category 3

Evidence of mechanism

■ Germ plasm (exp.)
■ Germ plasm (obs.)
■ Induction (exp.)

Figure 2 (previous page). Germ cell specification across Hexapoda. Phylogeny of studied hexapod lineages indicating the primordial germ cell origin timing category and the inferred mechanism of primordial germ cell specification. Phylogenetic relationships in the cladogram are derived from published relationships listed in Table S2; branch lengths are not to scale. Data on the timing of germ cell specification are shown in the first column as follows: dark purple = Category 1 (cleavage/blastoderm), medium purple = Category 2 (gastrulation), light purple = Category 3 (mesoderm differentiation). Data on the mechanism of germ cell specification, inferred as described in Supplementary Materials, are shown in the second column: dark pink = experimental evidence supports the use of germ plasm, light pink = morphological evidence of germ plasm, blue = experimental evidence of induction. Data and data sources are described in more detail in Table S1.

Clade

Tardigrada
Onychophora
Chelicerata
Myriapoda
Pancrustacea

Arthropoda

Timing category

■ Category 1
■ Category 2
■ Category 3

Evidence of mechanism

■ Germ plasm (exp.)
■ Germ plasm (obs.)
■ Induction (exp.)

Figure 3 (previous page). Panarthropod germ cell specification outside of Hexapoda. Phylogeny of
studied non-hexapod panarthropod lineages indicating the primordial germ cell origin timing category and
the inferred mechanism of primordial germ cell specification. Phylogenetic relationships in the cladogram
are derived from published relationships listed in Table S2; branch lengths are not to scale. Data on the
timing of germ cell specification are shown in the first column as follows: dark purple = Category 1
(cleavage/blastoderm), medium purple = Category 2 (gastrulation), light purple = Category 3 (mesoderm
differentiation). Data on the mechanism of germ cell specification, inferred as described in Supplementary
Materials, are shown in the second column: dark pink = experimental evidence supports the use of germ
plasm, light pink = morphological evidence of germ plasm, blue = experimental evidence of induction.
Data and data sources are described in more detail in Table S1.

A — AP patterning and germ cell determination are spatially and temporally separate

B — AP patterning and germ cell determinants overlap spatially and temporally

C — germ plasm controls both AP patterning and germ cell determination

Figure 4. Hypothesized links between spatial and temporal dynamics of germ cell origination and the mechanism of germ cell specification. (**A**) In species that induce germ cells later in embryogenesis (Category 3), the asymmetrically localized anterior–posterior (AP) patterning determinants (blue) and germ cell determination factors (pink) are spatially and temporally separate. (**B**) Proposed shift to earlier germ cell specification at the embryonic posterior in Category 2 insects enabled overlap between the patterning and germ cell determinants. (**C**) Subsequent hitchhiking of the asymmetric localization machinery by germ line determinants could have coupled germ cell specification and posterior patterning and thus promoted the evolution of germ plasm.

977
978
Figure 5. Evolutionary dynamics of panarthropod germ cell specification. Phylogenetic relationships
among all study species (Table S1) are derived from published relationships listed in Table S2; branch
lengths are not to scale. Colored lines to the right of each branch in the cladogram designate the
mechanism of germ cell specification (inferred as described in Supplementary Materials) as follows: pink
= germ plasm, blue = induction. Pink asterisks denote proposed instances of germ plasm evolution and
pink pluses mark ambiguous cases of germ plasm gain. Blue asterisks denote proposed instances of
germ plasm loss and blue pluses mark ambiguous cases of germ plasm loss.
986

References

Achtelig, M. and Krause, G. (1971). Experimente am ungefurchten Ei von Pimpla
turionellae L. (Hymenoptera) zur Funktionsanalyse des Oosombereichs. *Wilhelm*
Roux Arch. Dev. Biol. **167**, 164–182.

Alexeeva, N. and Tamberg, Y. (2021). Anatomical changes in postembryonic
development of Pycnogonum litorale. *J. Morphol.* **282**, 329–354.

Alexeeva, N., Tamberg, Y. and Shunatova, N. (2018). Postembryonic development of
pycnogonids: A deeper look inside. *Arthropod Struct. Dev.* **47**, 299–317.

Alwes, F. and Scholtz, G. (2014). The early development of the onychopod cladoceran
Bythotrephes longimanus (Crustacea, Branchiopoda). *Front. Zool.* **11**, 10.

Anderson, D. T. (1962a). The embryology of Dacus tryoni (Frogg.) [Diptera, Trypetidae
(= Tephritidae)], the Queensland fruit-fly. *Development* **10**, 248–292.

Anderson, D. T. (1962b). The Epigenetics of the Larva in Diptera. *Acta Zool.* **43**, 221–
228.

Anderson, D. T. (1967). Larval development and segment formation in the branchiopod
crustaceans Limnadia stanleyana King (Conchostraca) and Artemia salina (L.)
(Anostraca). *Aust. J. Zool.* **15**, 47–91.

Anderson, D. T. (1973). *Embryology and Phylogeny in Annelids and Arthropods.*
Oxford, UK: Pergamon Press.

Ando, H. and Tanaka, M. (1980). Early embryonic development of the primitive moths,
Endoclyta signifier Walker and E. excrescens Butler (Lepidoptera: Hepialidae). *Int. J.*
Ins. Morph. Embryol. **9**, 67–77.

Ansari, S., Troelenberg, N., Dao, V. A., Richter, T., Bucher, G. and Klingler, M.
(2018). Double abdomen in a short-germ insect: Zygotic control of axis formation
revealed in the beetle Tribolium castaneum. *Proc. Natl. Acad. Sci.* **115**, 1819–1824.

Aoki, S. T., Kershner, A. M., Bingman, C. A., Wickens, M. and Kimble, J. (2016).
PGL germ granule assembly protein is a base-specific, single-stranded RNase. *Proc*
Natl Acad Sci U S A **113**, 201524400.

Aristotle *De generatione animalia*.

Ballesteros, J. A., Santibáñez-López, C. E., Baker, C. M., Benavides, L. R., Cunha,
T. J., Gainett, G., Ontano, A. Z., Setton, E. V. W., Arango, C. P., Gavish-Regev,

E., et al. (2022). Comprehensive Species Sampling and Sophisticated Algorithmic
Approaches Refute the Monophyly of Arachnida. *Mol Biol Evol* **39**, msac021.

Biffis, C., Alwes, F. and Scholtz, G. (2009). Cleavage and gastrulation of the
dendrobranchiate shrimp Penaeus monodon (Crustacea, Malacostraca, Decapoda).
Arthropod Struct. Dev. **38**, 527–540.

Blondel, L., Besse, S., Rivard, E. L., Ylla, G. and Extavour, C. G. (2021). Evolution of
a cytoplasmic determinant: Evidence for the biochemical basis of functional evolution
of the novel germ line regulator Oskar. *Mol. Biol. Evol.* **38**, 5491–5513.

Bontems, F., Stein, A., Marlow, F., Lyautey, J., Gupta, T., Mullins, M. C. and Dosch,
R. (2009). Bucky Ball Organizes Germ Plasm Assembly in Zebrafish. *Curr. Biol.* **19**,
414–422.

Bounoure, L. (1939). *L'origine des cellules reproductrices et le problème de la lignée*
germinale. Paris, France: Gauthier-Villars.

Brauer, A. (1894). Beiträge zur Kenntnis der Entwicklungsgeschichte des Skorpions. *Z*
wiss. Zool. **57**, 402–432.

Brauer, A. (1925). Studies on the embryology of Bruchus quadrimaculatus, Fabr. *Ann.*
Entomol. Soc. Am. **18**, 283–312.

Brauer, A. (1949). Localization of presumptive areas in the blastoderm of the pea
beetle Callosobruchus maculatus Fabr., as determined by ultraviolet (2537 Å)
irradiation injury. *J. Exp. Zool.* **112**, 165–193.

Buchta, T., Özüak, O., Stappert, D., Roth, S. and Lynch, J. A. (2013). Patterning the
dorsal–ventral axis of the wasp Nasonia vitripennis. *Dev Biol* **381**, 189–202.

Bull, A. L. (1982). Stages of Living Embryos in the Jewel Wasp Mormoniella (Nasonia)
vitripennis (Walker) (Hymenoptera: Pteromalidae). *Int. J. Insect Morphol. Embryol.*
11, 1–23.

Buss, L. W. (1982). Somatic cell parasitism and the evolution of somatic tissue
compatibility. *Proc. Natl. Acad. Sci.* **79**, 5337–5341.

Buss, L. W. (1983). Evolution, development, and the units of selection. *Proc. Natl.*
Acad. Sci. **80**, 1387–1391.

Butt, F. H. (1949). Embryology of the Milkweed Bug, Oncopeltus fasciatus (Hemiptera).
Cornell Experiment Station Memoir 3–43.

Cannon, H. G. (1921). The Early Development of the summer egg of a Cladoceran
(Simocephalus vetulus). *J. Cell Sci.* **s2-65**, 627–642.

Cannon, H. G. (1924). On the development of an estherid crustacean. *Philos. Trans. R.*
Soc. Lond. Ser. B **212**, 395–430.

Carter, J.-M., Gibbs, M. and Breuker, C. J. (2015). Divergent RNA Localisation
Patterns of Maternal Genes Regulating Embryonic Patterning in the Butterfly Pararge
aegeria. *PLoS ONE* **10**, e0144471.

Cavallin, M. (1971). ["Substitutive polyembryony" and the problem of germinal line
origin in the phasma Carausius morosus Br]. *C. R. Hebd. des seances l'Acad. des*
Sci. Ser. D: Sci. Nat. **272**, 462–5.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Shapiro, M. D., Brady, S. D.,
Southwick, A. M., Absher, D. M., Grimwood, J., Schmutz, J., et al. (2010).
Adaptive Evolution of Pelvic Reduction in Sticklebacks by Recurrent Deletion of a
Pitx1 Enhancer. *Science* **327**, 302–305.

Chang, C., Dearden, P. and Akam, M. (2002). Germ Line Development in the
Grasshopper Schistocerca gregaria: vasa As a Marker. *Dev. Biol.* **252**, 100–118.

Chang, C., Lee, W.-C., Cook, C. E., Lin, G.-W. and Chang, T. (2006). Germ-plasm
specification and germline development in the parthenogenetic pea aphid
Acyrthosiphon pisum: Vasa and Nanos as markers. *Int. J. Dev. Biol.* **50**, 413–421.

Chang, C., Lin, G., Cook, C. E., Horng, S., Lee, H. and Huang, T. (2007). Apvasa
marks germ-cell migration in the parthenogenetic pea aphid Acyrthosiphon pisum
(Hemiptera: Aphidoidea). *Dev. Genes Evol.* **217**, 275–287.

Chaw, R. C., Vance, E. and Black, S. D. (2007). Gastrulation in the spider Zygiella x-
notata involves three distinct phases of cell internalization. *Dev. Dynam.* **236**, 3484–
3495.

Chen, R. P.-Y., Shih, C.-L., Lin, G.-W., Cook, C. E., Huang, T.-Y. and Chang, C.-C.
(2009). Developmental expression of Apnanos during oogenesis and embryogenesis
in the parthenogenetic pea aphid Acyrthosiphon pisum. *Int. J. Dev. Biol.* **53**, 169–
176.

Chen, Y., Fang, X., Tian, X.-Q., Cui, Z., Feng, H.-Y. and Qiu, G.-F. (2021).
Germ plasm and the origin of the primordial germ cells in the oriental river prawn,
Macrobrachium nipponense. *Cell Tissue Res.* **386**, 559–569.

Clark, E., Peel, A. D. and Akam, M. (2019). Arthropod segmentation. *Development*
146, dev170480.

Colonnetta, M. M., Goyal, Y., Johnson, H. E., Syal, S., Schedl, P. and Deshpande,
G. (2022). Preformation and epigenesis converge to specify primordial germ cell fate
in the early Drosophila embryo. *PLoS Genet.* **18**, e1010002.

Counce, S. J. (1963). Developmental morphology of polar granules in Drosophila.
Including observations on pole cell behavior and distribution during embryogenesis.
J. Morphol. **112**, 129–145.

Crother, B. I., White, M. E. and Johnson, A. D. (2007). Inferring developmental
constraint and constraint release: Primordial germ cell determination mechanisms as
examples. *J Theor Biol* **248**, 322–330.

Crother, B. I., White, M. E. and Johnson, A. D. (2016). Diversification and Germ-Line
Determination Revisited: Linking Developmental Mechanism with Species Richness.
Frontiers Ecol Evol **4**, 26.

Dannenberg, L. C. and Seaver, E. C. (2018). Regeneration of the germline in the
annelid Capitella teleta. *Dev. Biol.* **440**, 74–87.

Dearden, P. K. (2006). Germ cell development in the Honeybee (Apis mellifera); Vasa
and Nanos expression. *BMC Dev. Biol.* **6**, 6.

Dearden, P., Grbic, M. and Donly, C. (2003). Vasa expression and germ-cell
specification in the spider mite Tetranychus urticae. *Dev. Genes Evol.* **212**, 599–603.

Dohle, W. (1964). Die Embryonalentwicklung von Glomeris marginata (Villers) im
Vergleich zur Entwicklung anderer Diplopoden. *Zool. Jahrb. Anat.* **81**, 241–310.

Donoughe, S., Nakamura, T., Ewen-Campen, B., Green, D. A., Henderson, L. and
Extavour, C. G. (2014). BMP signaling is required for the generation of primordial
germ cells in an insect. *Proc. Natl. Acad. Sci.* **111**, 4133–4138.

Doyle, D. A., Burian, F. N., Aharoni, B., Klinder, A. J., Menzel, M. M., Nifras, G. C.
C., Shabazz-Henry, A. L., Palma, B. U., Hidalgo, G. A., Sottolano, C. J., et al.
(2023). Germ Granule Evolution Provides Mechanistic Insight into Drosophila
Germline Development. *Mol. Biol. Evol.* **40**, msad174.

Du, S., Tihelka, E., Yu, D., Chen, W.-J., Bu, Y., Cai, C., Engel, M. S., Luan, Y.-X. and
Zhang, F. (2024). Revisiting the four Hexapoda classes: Protura as the sister group
to all other hexapods. *Proc. Natl. Acad. Sci.* **121**, e2408775121.

Eddy, E. M. (1975). Germ Plasm and the Differentiation of the Germ Cell Line.
International Review of Cytology **43**, 229–280.

Edgar, A., Bates, C., Larkin, K. and Black, S. (2015). Gastrulation occurs in multiple
phases at two distinct sites in Latrodectus and Cheiracanthium spiders. *EvoDevo* **6**,
33.

Ephrussi, A. and Lehmann, R. (1992). Induction of germ cell formation by oskar.
Nature **358**, 387–392.

Ephrussi, A., Dickinson, L. K. and Lehmann, R. (1991). oskar organizes the germ
plasm and directs localization of the posterior determinant nanos. *Cell* **66**, 37–50.

Evans, R. (1901). On the Malayan Species of Onychophora. Part II - The Development
of Eoperipatus weldoni. *Q. J. Microscop. Sci.* **45**, 41–88.

Evans, T., Wade, C. M., Chapman, F. A., Johnson, A. D. and Loose, M. (2014).
Acquisition of Germ Plasm Accelerates Vertebrate Evolution. *Science* **344**, 200–203.

Everett, N. B. (1944). The present status of the germ-cell problem in vertebrates. *Biol*
Rev **20**, 45–55.

Ewen-Campen, B., Schwager, E. E. and Extavour, C. G. M. (2010). The molecular
machinery of germ line specification. *Mol. Reprod. Dev.* **77**, 3–18.

Ewen-Campen, B., Srouji, J. R., Schwager, E. E. and Extavour, C. G. (2012). oskar
Predates the Evolution of Germ Plasm in Insects. *Curr Biol* **22**, 2278–2283.

Ewen-Campen, B., Jones, T. E. M. and Extavour, C. G. (2013a). Evidence against a
germ plasm in the milkweed bug Oncopeltus fasciatus, a hemimetabolous insect.
Biol. Open **2**, 556–568.

Ewen-Campen, B., Donoughe, S., Clarke, D. N. and Extavour, C. G. (2013b). Germ
cell specification requires zygotic mechanisms rather than germ plasm in a basally
branching insect. *Curr. Biol.* **23**, 835–842.

Extavour, C. G. (2005). The fate of isolated blastomeres with respect to germ cell
formation in the amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* **277**, 387–402.

Extavour, C. G. M. (2007). Evolution of the bilaterian germ line: lineage origin and
modulation of specification mechanisms. *Integr. Comp. Biol.* **47**, 770–785.

Extavour, C. G. and Akam, M. (2003). Mechanisms of germ cell specification across
the metazoans: epigenesis and preformation. *Development* **130**, 5869–5884.

Faussek, V. (1888). Ueber die embryonale Entwicklung der Geschlechtsorgane bei der
Afterspinne (Phalangium). *Biol. Zentralbl.* **8**, 359–363.

Faussek, V. (1892). Zur Anatomie und Embryologie der Phalangiden. *Biol. Zentralbl.*
12, 1–8.

Fernández, R., Edgecombe, G. D. and Giribet, G. (2018). Phylogenomics illuminates
the backbone of the Myriapoda Tree of Life and reconciles morphological and
molecular phylogenies. *Sci. Rep.* **8**, 83.

Forrest, K. M. and Gavis, E. R. (2003). Live Imaging of Endogenous RNA Reveals a
Diffusion and Entrapment Mechanism for nanos mRNA Localization in Drosophila.
Curr. Biol. **13**, 1159–1168.

Gabriel, W. N., McNuff, R., Patel, S. K., Gregory, T. R., Jeck, W. R., Jones, C. D.
and Goldstein, B. (2007). The tardigrade Hypsibius dujardini, a new model for
studying the evolution of development. *Dev. Biol.* **312**, 545–559.

Gainett, G., Crawford, A. R., Klementz, B. C., So, C., Baker, C. M., Setton, E. V. W.
and Sharma, P. P. (2022). Eggs to long-legs: embryonic staging of the harvestman
Phalangium opilio (Opiliones), an emerging model arachnid. *Front. Zool.* **19**, 11.

Gatenby, J. B. (1917). The embryonic development of Trichogramma evanescens,
Westw., monembryonic egg parasite of Donacia simplex, Fab. *J. Cell Sci.* **s2-62**,
149–187.

Geigy, R. (1931). Action de l'ultra-violet sur le pôle germinal dans l'oeuf de Drosophila
melanogaster (Castration et mutabilité). *Rev. Suisse Zool.* **38**, 187–285.

Gerberding, M. and Patel, N. H. (2004). Chapter 6. Gastrulation in Crustaceans: germ
layers and cell lineages. In *Gastrulation: From Cells to Embryo* (ed. Stern, C.), p.
Cold Spring Harbor Laboratory Press.

Gerberding, M., Browne, W. E. and Patel, N. H. (2002). Cell lineage analysis of the
amphipod crustacean Parhyale hawaiensis reveals an early restriction of cell fates.
Development **129**, 5789–5801.

Gilbert, S. (1997). *Embryology: Constructing the Organism*. Sunderland, MA, USA:
Sinauer Associates Inc.

Giribet, G. and Edgecombe, G. D. (2019). The Phylogeny and Evolutionary History of
Arthropods. *Curr. Biol.* **29**, R592–R602.

Goodrich, A. L. (1939). The origin and fate of the entoderm elements in the
embryogeny of Porcellio laevis Latr. and Armadillidium nasutum B.L. (Isopoda). *J.*
Morphol. **64**, 401–426.

Görg, I. (1959). Untersuchungen am Keim von Hierodula (Rhombodera) crassa Giglio
Tos, ein Beitrag zur Embryologie der Mantiden. *Dtsch. Entomol. Z.* **6**, 389–450.

Goss, R. J. (1952). The early embryology of the book louse, Liposcelis divergens
Badonnel (Psocoptera; Liposcelidae). *J. Morphol.* **91**, 135–167.

Goss, R. J. (1953). The advanced embryology of the book louse, Liposcelis divergens
Badonnel (Psocoptera; Liposcelidae). *J. Morphol.* **92**, 157–205.

Gough, L. H. (1902). The development of Admetus pumilio, Koch: a contribution to the
embryology of the Pedipalps. *J. Cell Sci.* **s2-45**, 595–630.

Grattan, R. M., McCulloch, R. J., Sellars, M. J. and Hertzler, P. L. (2013).
Ultrastructure of putative germ granules in the penaeid shrimp Marsupenaeus
japonicus. *Arthropod Struct. Dev.* **42**, 153–164.

Green, J. E. and Akam, M. (2014). Germ cells of the centipede Strigamia maritima are
specified early in embryonic development. *Dev. Biol.* **392**, 419–430.

Gupta, T. and Extavour, C. G. (2013). Identification of a putative germ plasm in the
amphipod Parhyale hawaiensis. *EvoDevo* **4**, 34.

Haga, K. (1985). Oogenesis and embryogenesis of the idolothripine thrips, Bactrothrips
brevitubus (Thysanoptera, Phlaeothripidae). In *Recent Advances in Insect*
Embryology in Japan, pp. 45–106. Tsukuba: ISEBU.

Hargitt, G. T. (1919). Germ Cells of Coelenterates. VI. General considerations,
discussion, conclusions. *Journal of Morphology* **33**, 1–59.

Hegner, R. W. (1908). Effects of removing the germ-cell determinants from the eggs of
some chrysomelid beetles. Preliminary report. *Biol. Bull.* **16**, 19–26.

Hegner, R. W. (1909a). The effects of centrifugal force upon the eggs of some
chrysomelid beetles. *J. Exp. Zool.* **6**, 507–552.

Hegner, R. W. (1909b). The origin and early history of the germ-cells in some
chrysomelid beetles. *J. Morphol.* **20**, 231–296.

Hegner, R. W. (1911). Experiments with chrysomelid beetles. III. The effects of killing
parts of the eggs of Leptinotarsa decemlineata. *Biol. Bull.* **20**, 237–251.

Hegner, R. W. (1914). Studies on germ cells. I. The history of the germ cells in insects
with special reference to the Keimbahn-determinants. II. The origin and significance
of the Keimbahn-determinants in animals. *J. Morphol.* **25**, 375–509.

Heikes, K. L., Game, M., Smith, F. W. and Goldstein, B. (2023). The embryonic origin
of primordial germ cells in the tardigrade Hypsibius exemplaris. *Dev. Biol.* **497**, 42–
58.

Hejnol, A. and Schnabel, R. (2005). The eutardigrade Thulinia stephaniae has an
indeterminate development and the potential to regulate early blastomere ablations.
Development **132**, 1349–1361.

Heming, B. S. (1979). Origin and fate of germ cells in male and female embryos of
Haplothrips verbasci (Osborn) (Insecta, Thysanoptera, Phlaeothripidae). *J. Morphol.*
160, 323–343.

Heming, B. S. and Huebner, E. (1994). Development of the germ cells and
reproductive primordia in male and female embryos of Rhodnius prolixus Stål
(Hemiptera: Reduviidae). *Can. J. Zool.* **72**, 1100–1119.

Hertzler, P. L. (2005). Cleavage and gastrulation in the shrimp Penaeus (Litopenaeus)
vannamei (Malacostraca, Decapoda, Dendrobranchiata). *Arthropod Struct. Dev.* **34**,
455–469.

Heymons, R. (1895). *Die Embryonalentwickelung von Dermapteren und Orthopteren*
unter Besonderer Berücksichtigung der Keimblätterbildung. Jena: Jena Verlag von
Gustav Fischer.

Heymons, R. (1901). Die Entwicklungsgeschichte der Scolopender. *Zoologica* **33**, 1–
244.

Heys, F. (1931). The Problem of the Origin of Germ Cells. *Q Rev Biology* **6**, 1–45.

Hickman, V. (1936). The embryology of the Syncarid Crustacean, Anaspides
tasmaniae. *Pap. Proc. R. Soc. Tasman.* **1936**, 1–36.

His, W. (1868). *Untersuchungen über die erste Anlage des Wirbelthierleibes: die erste*
Entwickelung des Hünchens im Ei. Leipzig: F. C. W. Vogel.

Holm, Å. (1947). On the development of Opilio parietinus Deg. *Zool. bidr. Uppsala* 409–
422.

Hopf, C., Viebahn, C. and Püschel, B. (2011). BMP signals and the transcriptional
repressor BLIMP1 during germline segregation in the mammalian embryo. *Dev.*
Genes Evol. **221**, 209.

Huettner, A. F. (1923). The origin of the germ cells in Drosophila melanogaster. *J.*
Morphol. **2**, 385–423.

Ikuta, K. and Makioka, T. (1994). Notes on the postembryonic development of the
ovary in Argulus japonicus (Crustacea: Branchiura). *Proc. Arthropod. Embryol. Soc.*
Jpn. **29**, 15–17.

Illmensee, K. and Mahowald, A. P. (1974). Transplantation of posterior polar plasm in
Drosophila. Induction of germ cells at the anterior pole of the egg. *Proc. Natl. Acad.*
Sci. **71**, 1016–1020.

Illmensee, K., Mahowald, A. P. and Loomis, M. R. (1976). The ontogeny of germ
plasm during oogenesis in Drosophila. *Dev. Biol.* **49**, 40–65.

Jazdowska-Zagrodzińska, B. (1966). Experimental studies on the role of "polar
granules" in the segregation of pole cells in Drosophila melanogaster. *J. Embryol.*
Exp. Morph. **16**, 391–9.

Johannsen, O. A. and Butt, F. H. (1941). *Embryology of Insects and Myriapods: The*
developmental history of insects, centipedes, and millepedes from egg deposition to
hatching. New York: McGraw-Hill, Inc.

Johnson, A. D., Drum, M., Bachvarova, R. F., Masi, T., White, M. E. and Crother, B.
I. (2003). Evolution of predetermined germ cells in vertebrate embryos: implications
for macroevolution. *Evol Dev* **5**, 414–431.

Johnson, A. D., Richardson, E., Bachvarova, R. F. and Crother, B. I. (2011).
Evolution of the germ line–soma relationship in vertebrate embryos. *Reproduction*
141, 291–300.

Jones, J. R. and Macdonald, P. M. (2007). Oskar controls morphology of polar
granules and nuclear bodies in Drosophila. *Development* **134**, 233–236.

Juhn, J. and James, A. A. (2006). oskar gene expression in the vector mosquitoes,
Anopheles gambiae and Aedes aegypti. *Insect Mol. Biol.* **15**, 363–372.

Juhn, J., Marinotti, O., Calvo, E. and James, A. A. (2008). Gene structure and
expression of nanos (nos) and oskar (osk) orthologues of the vector mosquito, Culex
quinquefasciatus. *Insect Mol. Biol.* **17**, 545–552.

Jung, E. (1966). Untersuchungen am Ei des Speisebohnenkäfers Bruchidius obtectus
Say (Coleoptera). *Z. Morphol. Ökol. Tiere* **56**, 444–480.

Jura, C. (1967). Origin of germ cells and gonads formation in embryogenesis of
Tetrodontophora bielanensis (Collembola). *Acta Biol. Crac. Zool.* **10**, 97–103.

Kaczmarczyk, A. N. (2014). Germline maintenance and regeneration in the amphipod
crustacean, Parhyale hawaiensis.

Kautzsch, G. (1910). Über die Entwicklung von Agelena labyrinthica Clerck. II. Teil.
Zool. Jahrb. Anat. **30**, 535–602.

Kemph, A. and Lynch, J. A. (2022). Evolution of germ plasm assembly and function
among the insects. *Curr. Opin. Insect Sci.* **50**, 100883.

Kemph, A., Kharel, K., Tindell, S. J., Arkov, A. L. and Lynch, J. A. (2025). Novel structure and composition of the unusually large germline determinant of the wasp Nasonia vitripennis. *Mol. Biol. Cell* **36**, ar55.

Kharel, K., Tindell, S. J., Kemph, A., Schmidtke, R., Alexander, E., Lynch, J. A. and Arkov, A. L. (2024). Dynamic protein assembly and architecture of the large solitary membraneless organelle during germline development in the wasp Nasonia vitripennis. *Development* **151**, dev202877.

Khila, A. and Abouheif, E. (2008). Reproductive constraint is a developmental mechanism that maintains social harmony in advanced ant societies. *Proc. Natl. Acad. Sci.* **105**, 17884–17889.

Kistler, K. E., Trcek, T., Hurd, T. R., Chen, R., Liang, F.-X., Sall, J., Kato, M. and Lehmann, R. (2018). Phase transitioned nuclear Oskar promotes cell division of Drosophila primordial germ cells. *eLife* **7**, e37949.

Klag, J. (1982). Germ line of Tetrodontophora bielanensis (Insecta, Collembola). Ultrastructural study on the origin of primordial germ cells. *Development* **72**, 183–195.

Kobayashi, Y., Watanabe, T. and Suzuki, H. (2006). Embryonic development of the firefly Pyrocoelia rufa Olivier (Insecta: Coleoptera, Lampyridae), with special reference to its hibernal diapause. *Proc. Arthropod. Embryol.Soc. Jpn.* **41**, 47–53.

Kobayashi, Y., Niikura, K., Oosawa, Y. and Takami, Y. (2013). Embryonic development of Carabus insulicola (Insecta, Coleoptera, Carabidae) with special reference to external morphology and tangible evidence for the subcoxal theory. *J. Morphol.* **274**, 1323–1352.

Komatsu, S. and Kobayashi, Y. (2012). Embryonic development of a whirligig beetle, Dineutus mellyi, with special reference to external morphology (Insecta: Coleoptera, Gyrinidae). *J. Morphol.* **273**, 541–560.

Krause, G. (1939). Die Eitypen der Insekten. *Biol. Zentralbl.* **59**, 495–536.

Krause, J. B. (1947). The development of the gonads of the wood-eating beetle, Passalus cornutus Fabricius. *Ann. Entomol. Soc. Am.* **40**, 172–202.

Kühn, A. (1913). Die Sonderung der Keimesbezirke in der Entwicklung der Sommereier von Polyphemus pediculus de Geer. *Zool. Jahrb. Anat.* **35**, 243–340.

Kulkarni, A. and Extavour, C. G. (2017). Convergent evolution of germ granule nucleators: A hypothesis. *Stem Cell Res.* **24**, 188–194.

Kulkarni, A., Ewen-Campen, B., Terao, K., Matsumoto, Y., Li, Y., Watanabe, T.,
Kao, J. A., Parhad, S. S., Ylla, G., Mizunami, M., et al. (2023). oskar acts with the
transcription factor Creb to regulate long-term memory in crickets. *Proc. Natl. Acad.*
Sci. **120**, e2218506120.

Lall, S., Ludwig, M. Z. and Patel, N. H. (2003). Nanos Plays a Conserved Role in Axial
Patterning outside of the Diptera. *Curr. Biol.* **13**, 224–229.

Larink, O. (1969). Zur Entwicklungsgeschichte von Petrobius brevistylis (Thysanura,
Insecta). *Helgoländer Wiss. Meeresunters.* **19**, 111–155.

Lasko, P. F. and Ashburner, M. (1990). Posterior localization of vasa protein
correlates with, but is not sufficient for, pole cell development. *Genes Dev.* **4**, 905–
921.

Lawson, K. A., Dunn, N. R., Roelen, B. A. J., Zeinstra, L. M., Davis, A. M., Wright,
C. V. E., Korving, J. P. W. F. M. and Hogan, B. L. M. (1999). Bmp4 is required for
the generation of primordial germ cells in the mouse embryo. *Genes Dev.* **13**, 424–
436.

Lee, C.-Y. S., Putnam, A., Lu, T., He, S., Ouyang, J. P. T. and Seydoux, G. (2020).
Recruitment of mRNAs to P granules by condensation with intrinsically-disordered
proteins. *eLife* **9**, e52896.

Lehmann, R. (2016). Germ Plasm Biogenesis - An Oskar-Centric Perspective. *Curr.*
Top. Dev. Biol. **116**, 679–707.

Leptin, M. (2004). Chapter 7: Gastrulation in Drosophila. In *Gastrulation: From Cells to*
Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor Laboratory
Press.

Lin, G., Cook, C. E., Miura, T. and Chang, C. (2014). Posterior localization of ApVas1
positions the preformed germ plasm in the sexual oviparous pea aphid Acyrthosiphon
pisum. *EvoDevo* **5**, 18.

Lochab, A. K. and Extavour, C. G. (2017). Bone Morphogenetic Protein (BMP)
signaling in animal reproductive system development and function. *Dev. Biol.* **427**,
258–269.

Lynch, J. A. (2019). Evolution of maternal control of axial patterning in insects. *Curr.*
Opin. Insect Sci. **31**, 37–42.

Lynch, J. A. and Desplan, C. (2010). Novel modes of localization and function of
nanos in the wasp Nasonia. *Development* **137**, 3813–3821.

Lynch, J. A., Özüak, O., Khila, A., Abouheif, E., Desplan, C. and Roth, S. (2011).
The Phylogenetic Origin of oskar Coincided with the Origin of Maternally Provisioned
Germ Plasm and Pole Cells at the Base of the Holometabola. *PLoS Genet.* **7**,
e1002029.

Mahowald, A. P. (1962). Fine structure of pole cells and polar granules in Drosophila
melanogaster. *J. Exp. Zool.* **151**, 201–215.

Mahowald, A. P. (1968). Polar granules of Drosophila. II. Ultrastructural changes during
early embryogenesis. *J. Exp. Zool.* **167**, 237–261.

Mahowald, A. P. (2001). Assembly of the Drosophila germ plasm. *Int. Rev. Cytol.* **203**,
187–213.

Manton, S. M. (1928). On the Embryology of a Mysid Crustacean, Hemimysis
lamornae. *Philos. Trans. R. Soc. B.* **216**, 363–463.

Manton, S. M. (1934). On the embryology of the crustacean Nebalia bipes. *Philos.*
Trans. R. Soc. B. **223**, 163–238.

Manton, S. M. (1949). Studies on the Onychophora VII. The early embryonic stages of
Peripatopsis, and some general considerations concerning the morphology and
phylogeny of the Arthropoda. *Philos. Trans. R. Soc. B.* **233**, 483–580.

Marlow, F. L. and Mullins, M. C. (2008). Bucky ball functions in Balbiani body
assembly and animal–vegetal polarity in the oocyte and follicle cell layer in zebrafish.
Dev. Biol. **321**, 40–50.

Martin, A. and Orgogozo, V. (2013). The loci of repeated evolution: A catalog of
genetic hotspots of phenotypic variation. *Evolution* **67**, 1235–1250.

Mayer, G. and Tait, N. N. (2009). Position and development of oocytes in velvet worms
shed light on the evolution of the ovary in Onychophora and Arthropoda. *Zool. J.*
Linn. Soc. Lond. **157**, 17–33.

McClendon, J. F. (1907). On the development of parasitic copepods. Part II. *Biol. Bull.*
12, 53–88.

Mellanby, H. (1935a). The early embryonic development of Rhodnius prolixus
(Hemiptera, Heteroptera). *J. Cell Sci.* **s2-78**, 71–90.

Mellanby, H. (1935b). The Early Embryonic Development of Rhodnius prolixus
(Hemiptera, Heteroptera). *Q. J. Microscop. Sci.* **309**, 71–90.

Meng, C. (1968). Strukturwandel und histochemische Befunde insbesondere am
Oosom während der Oogenese und nach der Ablage des Eies von Pimpla turionellae

L. (Hymenoptera, Ichneumonidae). *Wilhelm Roux' Arch. für Entwicklungsmechanik*
Org. **161**, 162–208.

Metschnikow, E. (1866). *Embryologische studien an insecten*. Leipzig, Germany:
Verlag von Wilhelm Engelmann.

Misof, B., Liu, S., Meusemann, K., Peters, R. S., Donath, A., Mayer, C., Frandsen,
P. B., Ware, J., Flouri, T., Beutel, R. G., et al. (2014). Phylogenomics resolves the
timing and pattern of insect evolution. *Science* **346**, 763–767.

Mitsumoto, H. and Makioka, T. (2002). Preliminary notes on the structure and
postembryonic development of gonads in two tadpole shrimps Triops longicaudatus
and T. numidicus (Crustacea: Branchiopoda, Notostraca). *Proc. Arthropod. Embryol.*
Soc. Jpn. **37**, 13–17.

Miya, K. (1953). The presumptive genital region at the blastoderm stage of the silkworm
egg. *J. Fac. Agr. Iwate Univ.* **1**, 223–227.

Miya, K. (1958a). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part I. Differentiation of Germ Cells. *J. Fac. Agr. Iwate*
Univ. 436–467.

Miya, K. (1958b). Studies on the Embryonic Development of the Gonad in the
Silkworm, Bombyx mori L. Part II. Formation of the Gonad. *J. Fac. Agr. Iwate Univ.*
126–151.

Miyakawa, K. (1974). The embryology of the caddisfly Stenopsyche griseipennis
MacLachlan (Trichoptera, Stenopsychidae): IV. Organogenesis: mesodermal
derivatives. *Kontyû* **42**, 451–466.

Miyazaki, K. and Makioka, T. (2012). Postembryonic development of the female
reproductive system in the pycnogonid, Propallene longiceps (Pycnogonida,
Callipallenidae). *Invertebr. Reprod. Dev.* **56**, 287–292.

Morris, S. C. (2009). The predictability of evolution: glimpses into a post-Darwinian
world. *Die Naturwissenschaften* **96**, 1313–37.

Nakamura, T. and Extavour, C. G. (2016). The transcriptional repressor Blimp-1 acts
downstream of BMP signaling to generate primordial germ cells in the cricket Gryllus
bimaculatus. *Development* **143**, 255–263.

Nakao, H. (1999). Isolation and characterization of a Bombyx vasa-like gene. *Dev.*
Genes Evol. **209**, 312–316.

Nakao, H. and Takasu, Y. (2019). Complexities in Bombyx germ cell formation process
revealed by Bm-nosO (a Bombyx homolog of nanos) knockout. *Dev. Biol.* **445**, 29–
36.

Nakao, H., Hatakeyama, M., Lee, J. M., Shimoda, M. and Kanda, T. (2006).
Expression pattern of Bombyx vasa-like (BmVLG) protein and its implications in germ
cell development. *Dev. Genes Evol.* **216**, 94–99.

Nakao, H., Matsumoto, T., Oba, Y., Niimi, T. and Yaginuma, T. (2008). Germ cell
specification and early embryonic patterning in Bombyx mori as revealed by nanos
orthologues. *Evol. Dev.* **10**, 546–554.

Needham, A. E. (1942). The structure and development of the segmental excretory
organs of Asellus aquaticus (Linné). *J. Cell Sci.* **s2-83**, 205–243.

Nelsen, O. E. (1934). The segregation of the germ cells in the grasshopper, Melanoplus
differentialis (Acrididae; Orthoptera). *J. Morphol.* **55**, 545–575.

Nelson, J. A. (1915). *The Embryology of the Honey Bee*. Princeton: Princeton
University Press.

Nieuwkoop, P. D. and Sutasurya, L. A. (1979). *Primordial Germ Cells in the*
Chordates: embryogenesis and phylogenesis. Cambridge University Press.

Nieuwkoop, P. D. and Sutasurya, L. A. (1981). *Primordial Germ Cells in the*
Invertebrates: from epigenesis to preformation. Cambridge: Cambridge University
Press.

Niikura, K., Hirasawa, K., Inoda, T. and Kobayashi, Y. (2017). Embryonic
Development of a Diving Beetle, Hydaticus pacificus Aubé(Insecta: Coleoptera;
Dytiscidae): External Morphology and Phylogenetic Implications. *Proc. Arthropod.*
Embryol. Soc. Jpn. **48**, 19–32.

Okada, M., Kleinman, I. A. and Schneiderman, H. A. (1974). Restoration of fertility in
sterilized Drosophila eggs by transplantation of polar cytoplasm. *Dev. Biol.* **37**, 43–
54.

Olesnicky, E. C. and Desplan, C. (2007). Distinct mechanisms for mRNA localization
during embryonic axis specification in the wasp Nasonia. *Dev Biol* **306**, 134–142.

Otto, S. P. and Hastings, I. M. (1998). Mutation and selection within the individual.
Genetica **102–103**, 507.

Özpolat, B. D. (2024). Annelids as models of germ cell and gonad regeneration. *J. Exp.*
Zool. Part B **342**, 126–143.

Pagel, M. (1994). Detecting correlated evolution on phylogenies: a general method for
the comparative analysis of discrete characters. *Proc. Roy. Soc. London, Ser. B,*
Biol. Sci **255**, 37–45.

Paz, J. de la (1958). Studies on the embryogeny of Oncopeltus fasciatus (Dallas)
(Hemiptera: Lygaeidae).

Pires-daSilva, A. and Sommer, R. J. (2003). The evolution of signalling pathways in
animal development. *Nat. Rev. Genet.* **4**, 39–49.

Poulson, D. F. (1947). The pole cells of Diptera, their fate and significance. *Proc. Natl.*
Acad. Sci. **33**, 182–4.

Poulson, D. F. (1950). Histogenesis, organogenesis, and differentiation in the embryo
of Drosophila melanogaster Meigen. In *Biology of Drosophila* (ed. Demerec, M.), pp.
168–274. New York: Hafner Pub. Co.

Presser, B. D. and Rutschky, C. W. (1957). The embryonic development of the corn
earworm, Heliothis zea (Boddie) (Lepidoptera, Phalaenidae). *Ann. Entomol. Soc. Am.*
50, 133–164.

Quan, H. (2018). Evolution of insect germline specification.

Quan, H. and Lynch, J. A. (2016). The evolution of insect germline specification
strategies. *Curr. Opin. Insect Sci.* **13**, 99–105.

Queller, D. C. (2000). Relatedness and the fraternal major transitions. *Philos. Trans. R.*
Soc. Lond. Ser. B: Biol. Sci. **355**, 1647–1655.

Rafiqi, Ab. M., Rajakumar, A. and Abouheif, E. (2020). Origin and elaboration of a
major evolutionary transition in individuality. *Nature* **585**, 239–244.

Reed, R. D., Papa, R., Martin, A., Hines, H. M., Counterman, B. A., Pardo-Diaz, C.,
Jiggins, C. D., Chamberlain, N. L., Kronforst, M. R., Chen, R., et al. (2011). optix
drives the repeated convergent evolution of butterfly wing pattern mimicry. *Science*
333, 1137–41.

Rempel, J. G. and Church, N. S. (1969). The embryology of Lytta viridana Le Conte
(Coleoptera: Meloidae). V. The blastoderm, germ layers, and body segments. *Can. J.*
Zool. **47**, 1157–1171.

Rivard, E. L., Srouji, J., Repouliou, A. and Extavour, C. G. (2025). Protein sequence
evolution underlies interspecies incompatibility of a cell fate determinant. *bioRxiv*
10.1101/2025.08.02.668269, 1–53.

Rosenberg, M. I., Lynch, J. A. and Desplan, C. (2009). Heads and tails: Evolution of
antero-posterior patterning in insects. *BBA* **1789**, 333–342.

Roth, S. (2004). Chapter 8: Gastrulation in Other Insects. In *Gastrulation: From Cells to*
Embryo (ed. Stern, C.), p. New York, NY, USA: Cold Spring Harbor Laboratory
Press.

Saitou, M. and Yamaji, M. (2010). Germ cell specification in mice: signaling,
transcription regulation, and epigenetic consequences. *Reproduction* **139**, 931–942.

Santos, A. C. and Lehmann, R. (2004). Germ Cell Specification and Migration in
Drosophila and beyond. *Curr Biol* **14**, R578–R589.

Satterlee, J. W., Alonso, D., Gramazio, P., Jenike, K. M., He, J., Arrones, A.,
Villanueva, G., Plazas, M., Ramakrishnan, S., Benoit, M., et al. (2024).
Convergent evolution of plant prickles by repeated gene co-option over deep time.
Science **385**, eado1663–eado1663.

Schimkewitsch, W. (1896). Studien über parasitische Copepoden. *Z. wiss. Zool.* **61**,
339–362.

Schimkewitsch, V. M. (1903). Über die Entwicklung von Thelyphonus caudatus (L.).
Zool. Anz. **26**, 665–685.

Scholl, A., Liu, Y. and Seydoux, G. (2024). Caenorhabditis elegans germ granules
accumulate hundreds of low translation mRNAs with no systematic preference for
germ cell fate regulators. *Development* **151**, dev202575.

Scholtz, G. (2000). Evolution of the nauplius stage in malacostracan crustaceans. *J.*
Zool. Syst. Evol. Res. **38**, 175–187.

Scholtz, G. and Wolff, C. (2002). Cleavage, gastrulation, and germ disc formation of
the amphipod Orchestia cavimana (Crustacea, Malacostraca, Peracarida). *Contrib.*
Zool. **71**, 9–28.

Scholtz, G. and Wolff, C. (2013). Chapter 4: Arthropod Embryology: Cleavage and
Germ Band Development. In *Arthropod Biology and Evolution* (ed. Minelli, A.), pp.
63–89.

Schröder, R. (2006). vasa mRNA accumulates at the posterior pole during blastoderm
formation in the flour beetle Tribolium castaneum. *Dev. Genes Evol.* **216**, 277.

Schwager, E. E., Meng, Y. and Extavour, C. G. (2015). vasa and piwi are required for
mitotic integrity in early embryogenesis in the spider Parasteatoda tepidariorum. *Dev.*
Biol. **402**, 276–290.

Schwentner, M., Combosch, D. J., Nelson, J. P. and Giribet, G. (2017). A
Phylogenomic Solution to the Origin of Insects by Resolving Crustacean-Hexapod
Relationships. *Curr. Biol.* **27**, 1818-1824.e5.

Sedgwick, A. (1887). The development of the Cape species of Peripatus: Part III. On
the changes from stage A to stage F. *J. Cell Sci.* **s2-27**, 467–550.

Seervai, R. N. H. and Wessel, G. M. (2013). Lessons for inductive germline
determination. *Mol. Reprod. Dev.* **80**, 590–609.

Seidel, F. (1924). Die geschlechtsorgane in der embryonalen entwicklung von
Pyrrhocoris apterus L. *Z. Morphol. Ökol. Tiere* **1**, 429–506.

Sharma, P. P. and Gavish-Regev, E. (2025). The Evolutionary Biology of Chelicerata.
Annu. Rev. Entomol. **70**, 143–163.

Shinji, G. O. (1919). Embryology of coccids, with especial reference to the formation of
the ovary, origin and differentiation of the germ cells, germ layers, rudiments of the
midgut, and the intracellular symbiotic organisms. *J. Morphol.* **33**, 72–167.

Singh, J. P. (1967). Early embryonic development of gonads in Labidura riparia (Pallas)
(Labiduridae: Dermaptera). *Agra Univ. J. Res.* **1621**, 67–76.

Sonnenblick, B. P. (1941). Germ cell movements and sex differentiation of the gonads
in the Drosophila embryo. *Proc. Natl. Acad. Sci.* **27**, 484–9.

Srouji, J. and Extavour, C. G. (2010). Redefining Stem Cells and Assembling Germ
Plasm: Key Transitions in the Evolution of the Germ Line. In *Key Innovations in*
Animal Evolution (ed. DeSalle, R.) and Schierwater, B.), pp. 360–397. Enfield, NH:
CRC Press.

Stern, C. D. (2004). *Gastrulation: From Cells to Embryo*. (ed. Stern, C. D.) New York,
USA: Cold Spring Harbor Laboratory Press.

Strömberg, J.-O. (1965). On the embryology of the isopod Idotea. *Ark. Zool.* **2**, 421–
473.

Strömberg, J.-O. (1968). Segmentation and organogenesis in Limnoria lignorum
(Rathke) (Isopoda). *Ark. Zool.* **20**, 91.

Sudler, M. T. (1899). The development of Penilia schmackeri Richard. *Proc. Boston*
Soc. Nat. Hist. **29**, 109–131.

Suzuki, N. (1990). Embryology of the Mecoptera (Panorpidae, Panorpodidae,
Bittacidae and Boreidae). *Bull. Sugadaira Montane Res. Cent.* **11**, 1–87.

Suzuki, N., Shimizu, S. and Ando, H. (1981). Early embryology of the alderfly, Sialis
mitsuhashii Okamoto (Megaloptera: Sialidae). *Int. J. Insect Morphol. Embryol.* **10**,
409–418.

Tadokoro, R., Sugio, M., Kutsuna, J., Tochinai, S. and Takahashi, Y. (2006). Early
Segregation of Germ and Somatic Lineages during Gonadal Regeneration in the
Annelid Enchytraeus japonensis. *Curr. Biol.* **16**, 1012–1017.

Takamura, K., Fujimura, M. and Yamaguchi, Y. (2002). Primordial germ cells
originate from the endodermal strand cells in the ascidian Ciona intestinalis. *Dev.*
Genes Evol. **212**, 11–18.

Tanaka, M. (1987). Differentiation and Behaviour of Primordial Germ Cells during the
Early Embryonic Development of Parnassius glacialis Butler, Luehdorfia japonica
Leech and Byasa (Atrophaneura) alcinous alcinous Klug (Lepidoptera: Papilionidae).
Recent Advances in Insect Embryology in Japan and Poland 255–266.

Terao, A. (1929). On the embryonic development of the spiny lobster, Panulirus
japonicus (v. Siebold). *Jpn. J. Zool.* **2**, 387–449.

Tiegs, O. W. (1940). The embryology and affinities of the Symphyla, based on a study
of Hanseniella agilis. *J. Cell Sci.* **s2-82**, 1–208.

Tiegs, O. W. (1947). The development and affinities of the Pauropoda, based on a
study of Pauropus silvaticus, Part I. *J. Cell Sci.* **s3-88**, 165–267.

Toyama, K. (1902). Contributions to the Study of Silk-Worms. I. On the Embryology of
the Silk-Worms. *Bull. Coll. Agric. Tokyo Imp. Univ.* **5**, 73–118.

Trcek, T. and Lehmann, R. (2019). Germ granules in Drosophila. *Traffic* **20**, 650–660.

Trcek, T., Grosch, M., York, A., Shroff, H., Lionnet, T. and Lehmann, R. (2015).
Drosophila germ granules are structured and contain homotypic mRNA clusters. *Nat.*
Commun. **6**, 7962.

Ullmann, S. L. (1964). The origin and structure of the mesoderm and the formation of
the coelomic sacs in Tenebrio molitor L. [Insecta, Coleoptera]. *Philos. Trans. R. Soc.*
Lond., B **248**, 245–277.

Underwood, E. M., Caulton, J. H., Allis, C. D. and Mahowald, A. P. (1980).
Developmental fate of pole cells in Drosophila melanogaster. *Dev. Biol.* **77**, 303–314.

Wang, C. and Lehmann, R. (1991). Nanos is the localized posterior determinant in
Drosophila. *Cell* **66**, 637–647.

Weismann, A. (1863). Die Entwickelung der Dipteren im Ei, nach Beobachtungen an
Chironomous spec., Musca vomitoria und Pulex canis. *Zeit. wiss. Zool.* **13**, 107–220.

Wheeler, W. M. (1893). A Contribution to Insect Embryology. *J. Morphol.* **8**, 1–160.

Whittle, C. A. and Extavour, C. G. (2016). Refuting the hypothesis that the acquisition
of germ plasm accelerates animal evolution. *Nat. Commun.* **7**, 12637.

Whittle, C. A. and Extavour, C. G. (2017). Causes and evolutionary consequences of
primordial germ-cell specification mode in metazoans. *Proc. Natl. Acad. Sci.* **114**,
5784–5791.

Witschi, E. (1934). On determinative cleavage and yolk formation in the harpactid
copepod Tisbe furcata (Baird). *Biol. Bull.* **67**, 335–340.

Wolff, E. (1962). *L' origine de la lignée germinale chez les vertébrés et chez quelques
groupes d'invertébrés*. Paris, France: Éditions Hermann.

Wolff, C. (2009). The embryonic development of the malacostracan crustacean
Porcellio scaber (Isopoda, Oniscidea). *Dev. Genes Evol.* **219**, 545–564.

Wolff, C. and Scholtz, G. (2002). Cell Lineage, Axis Formation, and the Origin of Germ
Layers in the Amphipod Crustacean Orchestia cavimana. *Dev. Biol.* **250**, 44–58.

Woodland, J. T. (1957). A contribution to our knowledge of lepismatid development. *J.
Morphol.* **101**, 523–577.

Woodworth, C. W. (1889). Studies on the embryological development of Euvanessa
antiopa. In *The Butterflies of the Eastern United States and Canada, with special
reference to New England*, pp. 95–104. Cambridge, MA: Published by the Author.

Wotton, K. R., Jiménez-Guri, E., Matheu, B. G. and Jaeger, J. (2014). A staging
scheme for the development of the scuttle fly Megaselia abdita. *PLoS ONE* **9**,
e84421.

Xu, X., Brechbiel, J. L. and Gavis, E. R. (2013). Dynein-Dependent Transport of
nanos RNA in Drosophila Sensory Neurons Requires Rumpelstiltskin and the Germ
Plasm Organizer Oskar. *J. Neurosci.* **33**, 14791–14800.

Yajima, M. and Wessel, G. M. (2011). The DEAD-box RNA helicase Vasa functions in
embryonic mitotic progression in the sea urchin. *Development* **138**, 2217–2222.

Ying, Y. and Zhao, G.-Q. (2001). Cooperation of Endoderm-Derived BMP2 and
Extraembryonic Ectoderm-Derived BMP4 in Primordial Germ Cell Generation in the
Mouse. *Dev. Biol.* **232**, 484–492.

Ying, Y., Liu, X. M., Marble, A., Lawson, K. A. and Zhao, G. Q. (2000). Requirement
of Bmp8b for the generation of primordial germ cells in the mouse. *Mol. Endocrinol.*
14, 1053–63.

Yoon, Y., Klomp, J., Martin-Martin, I., Criscione, F., Calvo, E., Ribeiro, J. and
Schmidt-Ott, U. (2019). Embryo polarity in moth flies and mosquitoes relies on
distinct old genes with localized transcript isoforms. *eLife* **8**, e46711.

Zissler, D. (1992). From egg to pole cells: Ultrastructural aspects of early cleavage and
germ cell determination in insects. *Microsc. Res. Tech.* **22**, 49–74.

1613