Discovery and genomics of H$_2$-oxidizing/O$_2$-reducing *Deferribacterota*

ectosymbiotic with protists in the guts of termites and a *Cryptocercus*

cockroach

Naoya Maruoka[1], Rinpei Kudo[1], Katsura Igai[1], Michiru Shimizu[2], Masahiro Yuki[2],

Moriya Ohkuma[2], Yuichi Hongoh[1,2*]

[1]School of Life Science and Technology, Institute of Science Tokyo, Tokyo, Japan

[2]Japan Collection of Microorganisms/Microbe Division, RIKEN BioResource Research

Center, Ibaraki, Japan

*Corresponding author: Yuichi Hongoh (yhongo@life.isct.ac.jp), Department of Life Science

and Technology, Institute of Science Tokyo, Ookayama 2-12-1 W3-48, Meguro-ku, Tokyo

152-8550, Japan

Running headline: *Deferribacterota* attached to protists

E-mail: maruoka.n.aa@m.titech.ac.jp; bananariyasu@gmail.com; kigai@life.isct.ac.jp;

michiru.tahara@riken.jp; masahiro.yuki@a.riken.jp; mohkuma@riken.jp

Abstract

Members of the phylum *Deferribacterota* inhabit diverse environments, but their symbiosis

with protists has never been reported. We discovered an ectosymbiotic clade of

Deferribacterota specifically associated with spirotrichonymphid protists in the guts of the

termites *Reticulitermes speratus* and *Hodotermopsis sjostedti* and trichonymphid protists in

the gut of the wood-feeding cockroach *Cryptocercus punctulatus*. The ectosymbiotic

Deferribacterota were spiral shaped and attached to 16–91% of the host protist cells. These

formed a monophyletic cluster within an uncultured insect gut-associated family-level clade,

which is sister to the vertebrate gut-associated family *Mucispirillaceae*. The complete genome

of an ectosymbiotic *Deferribacterota* was obtained from a *Trichonympha acuta* cell in a *C.*

punctulatus gut and analyzed together with a single-cell amplified genome of another

ectosymbiotic *Deferribacterota* associated with *Holomastigotes* sp. in the gut of *R. speratus*.

Genome analyses suggest that these *Deferribacterota* ferment monosaccharides and conduct

fumarate and oxidative respiration with H_2 as an electron donor. They thus possibly

contribute to the removal of hydrogen and oxygen to protect the fermentative activity of the

protist hosts. The ectosymbionts possess reduced signal transduction gene repertoires,

implying that the association has provided a relatively stable environment for these bacteria.

The ectosymbionts likely possess flagella with an unusually expanded number of flagellin

variants up to 40, which may reflect an adaptation to their ectosymbiotic lifestyle. We propose

a novel genus, *Termitispirillum*, for these ectosymbionts and a novel family,

Termitispirillaceae, for the insect-gut clade, under SeqCode. Our findings provide new

insights into the ecology and evolution of *Deferribacterota*.

Keywords: *Deferribacterota*, protists, termites, *Cryptocercus*, symbiosis, genomics; gut

bacteria

Introduction

Deferribacterota is a phylum of Gram-negative curved rods or spiral-shaped bacteria [1, 2], currently comprising one class (*Deferribacteres*), one order (*Deferribacterales*), and seven families (*Calditerrivibrionaceae*, *Deferribacteraceae*, *Deferrivibrionaceae*, *Flexistipitaceae*, *Geovibrionaceae*, *Mucispirillaceae*, and "*Candidatus* Microvillispirillaceae") [3–5]. The first five families have been isolated from a wide variety of environments, including deep-sea hydrothermal vents [6], activated sludges [7], terrestrial hot springs [8], and subsurface oil reservoirs [9], whereas the latter two families have exclusively been discovered in animal intestines.

Mucispirillaceae inhabit the guts of vertebrates, including rodents [10], chickens [11], and turkeys [12]. *Mucispirillum schaedleri*, the sole cultured species in the family *Mucispirillaceae*, inhabits the guts of rodents and is reportedly involved in host inflammation [10, 13]. "*Candidatus* Microvillispirillaceae" is an uncultured family represented by a single species, "*Candidatus* Rimicarispirillum atlantis", which occupies the space among microvilli of the ectoperitrophic space in the midgut of the deep-sea hydrothermal vent shrimp genus *Rimicaris* [5, 14, 15]. "*Candidatus* Rimicarispirillum atlantis" can degrade chitin and presumably contributes to the turnover of chitin released during shrimp molt [5, 16]. Members of *Deferribacterota* have also been detected by 16S rRNA amplicon sequencing and metagenomic analyses as minor residents in the guts of insects, including the Auckland tree wētā [17], cockroaches [18, 19], and termites [20–22].

Termites thrive solely on dead plant matter with the aid of their hindgut microbiota [23]. In "lower" termites, i.e., all families except Termitidae ("higher" termites), and their closest relative, the cockroach genus *Cryptocercus*, the digestion of lignocellulose is largely attributable to the gut protists belonging to the phylum *Parabasalia* or *Preaxostyla* [24, 25]. Most of these protists harbor symbiotic prokaryotes in their cytoplasm, nucleoplasm, and/or

on the cell surface [26]. It has been suggested that these prokaryotic symbionts contribute to

the provision of nitrogenous compounds [27–34] and/or hydrogen removal to promote

fermentative activities in the gut [29, 35–38].

During our previous single-cell genomic survey of the bacterial microbiota in the gut of the

termite *Reticulitermes speratus* [39], a single-cell amplified genome (SAG), designated

MS499-41, was obtained and phylogenetically assigned to *Deferribacterota*. Our preliminary

fluorescence in situ hybridization (FISH) analysis implied that the bacterium is an

ectosymbiont of a gut protist; however, symbiosis between *Deferribacterota* and protists has

never been reported thus far. Thus, in the present study, we aimed to clarify the localization

and phylogenetic diversity of *Deferribacterota* in the guts of termites and cockroaches as well

as their metabolic capacities and symbiotic roles based on their genome sequences. This study

provides evolutionary and functional perspectives on the novel, uncultured *Deferribacterota*

clade associated with cellulolytic gut protists.

Materials and Methods

Termite and cockroach collection

The Japanese subterranean termite *R. speratus* (family Heterotermitidae) was collected in

Tsukuba, Ibaraki Prefecture, and Ogose, Saitama Prefecture in Japan. The damp-wood termite

Hodotermopsis sjostedti (family Archotermopsidae) was collected in Yakushima, Kagoshima

Prefecture in Japan. The wood-feeding cockroach *Cryptocercus punctulatus* was collected in

the Black Rock Mountain State Park, Georgia, U.S.A. under permit numbers 192019, 202022,

and 222017. The termites and cockroaches were kept with fragments of their nest logs in the

laboratory.

16S rRNA gene amplicon sequencing and phylogenetic analysis

DNA was extracted from the entire guts of two *C. punctulatus* adults individually using the

Power Soil DNA Isolation Kit (MoBio Laboratories). The V3–V4 regions of 16S rRNA (ca.

450 bp) genes were amplified by PCR using the prokaryote-universal primer set Pro341F and

Pro805R (Table S1). The library preparation, sequencing on a MiSeq System (Illumina), and

amplicon data processing were conducted as previously described [40]. The amplicon

sequence variants (ASVs) obtained in this study and in our previous study using 60 termite

and eight cockroach species [40] were phylogenetically classified using SINA v1.7.2 [41]

with the database SILVA SSURef NR99 release 138.2. The ASVs assigned to

Deferribacterota were aligned with reference sequences using MAFFT v7 [42]. Ambiguously

aligned regions were trimmed using trimAl v1.2 [43]. Maximum likelihood trees were

constructed with 1,000 ultrafast bootstrap resampling using IQ-TREE v1.6.12 [44]. The best-

fit nucleotide substitution models were selected using ModelFinder [45].

FISH

The entire guts were removed from worker termites and adult cockroaches, and the gut

contents were suspended in Trager's solution U [46]. The samples were fixed in 4%

paraformaldehyde for over 8 hrs at 4°C, rinsed twice with double-distilled water, and placed

on MAS-GP glass slides (Matsunami, Japan). FISH targeting 16S rRNA was performed as

previously described [47, 48] using oligonucleotide probes labeled with Texas Red or 6-

carboxyfluorescein (6FAM). Specific oligonucleotide probes (Table S2) were designed using

ARB [49]. The specimens were observed under an Olympus BX51 epifluorescence

microscope.

Collection of protist cells, Sanger sequencing of small subunit (SSU) rRNA genes, and

phylogenetic analysis

Termites and cockroaches were dissected, and the gut contents were suspended in sterile

Trager's solution U containing 0.01% bovine serum albumin (BSA). Single protist cells were

collected in PCR tubes using a TransferMan NK2 micromanipulator (Eppendorf) equipped

with a hand-pulled glass capillary. The collected protist cells were subjected to whole genome

amplification (WGA) using a GenomiPhi V2 DNA Amplification Kit (GE Healthcare Life

Science). Near full-length 16S and 18S rRNA genes were amplified from the WGA products

by PCR using Phusion High-Fidelity DNA polymerase (New England Biolabs). PCR was

performed with initial denaturation at 98°C for 3 min, followed by 25 cycles of denaturation

at 98□ for 30 sec, annealing at 50–70°C for 30 sec, and extension at 72□ for 1 min with a

final extension at 72□ for 10 min. For *Holomastigotes* samples, nested PCR was performed

for the amplification of the 18S rRNA genes using primer sets listed in Table S1 and the

above program. The PCR products were cloned using a Zero Blunt TOPO PCR Cloning Kit

(Thermo Fisher Scientific) after purification using a MonoFas DNA Purification Kit I

(ANIMOS, Japan). Sanger sequencing of randomly chosen clones was conducted using a

BigDye Terminator v3.1 Cycle Sequencing Kit (Applied Biosystems) on an ABI 3730

Genetic Analyzer. The SSU rRNA gene trees of *Deferribacterota* and their host protists were

constructed as described above.

Genome sequencing and assembly

A SAG of *Deferribacterota* (MS499-41: DRR253682) was obtained in our previous study

from the gut content of *R. speratus* using the MDA-in-AGM method and a MiSeq System

with a MiSeq Reagent Kit V3 (600 cycles) [39]. In the present study, the MiSeq raw reads

were subjected to quality trimming using Cutadapt 3.7 [50] and Prinseq-lite 0.20.4 [51] and

were assembled using SPAdes v3.14.0 [52] with the --sc option. Contaminating contigs were

identified and removed using the Contig Annotation Tool v5.2.3 [53] and BLASTn searches

of the NCBI non-redundant nucleotide database [54]. Only contigs >500 bp were retained.

To acquire the genome sequence of *Deferribacterota* associated with *Trichonympha acuta*,

the hindgut content of *C. punctulatus* was suspended in sterile Trager's solution U containing

0.01% BSA, and the membrane portion of a single *T. acuta* cell was separated using a Micro

Feather Blade K-715 attached to the micromanipulator and collected in a PCR tube. The

sample was subjected to WGA using EquiPhi29 DNA Polymerase (Thermo Fisher Scientific)

as previously described [55]. The WGA product was purified using a Zymo Genomic DNA

Clean & Concentrator kit (ZYMO Research). Sequencing libraries were prepared using a

QIAseq FX DNA Library Kit (QIAGEN) for the MiSeq System and a Native Barcoding Kit

24 V14 for a MinION System (Oxford Nanopore Technologies). Prior to the latter library

preparation, debranching, single-strand DNA digestion, purification, and size selection that

retained fragments of >2 kbp were conducted as previously described [55]. Sequencing was

performed on the MiSeq System as above and on the MinION System with a FLO-MIN114

(R10.4.1) flow cell. After base calling using dorado and quality-filtering using nanoq [56], the

MinION reads were assembled using Raven [57]. The resulting circular contig was polished

one time using Pilon v1.24 [58] with the quality-filtered MiSeq reads.

The genome quality was assessed using CheckM2 [59]. Circular genomes were visualized

using Proksee [60]. The average nucleotide identity (ANI) and average amino acid identity

(AAI) were calculated using FastANI [61] and EzAAI [62], respectively. The Skew Index

(SkewI), which indicates the degree of GC skew, was calculated using SkewIT [63].

Phylogenomic analysis

The phylogenomic analysis was conducted using GToTree [64] as the following process.

Seventy-four single-copy marker genes conserved among *Bacteria* [64] were extracted from

Deferribacterota genomes using Prodigal [65] and HMMER [66]. Only genomes that

contained ≥50% of the marker genes were retained as references. The deduced amino acid

sequences were aligned using MUSCLE [67], trimmed using trimAl v1.2, and concatenated to

construct a supermatrix. A maximum likelihood tree was constructed using IQ-TREE as

above. The phylogenetic classification was confirmed using the Genome Taxonomy Database

(GTDB) [68].

Functional annotation

The reconstructed genomes were annotated using DFAST [69], KAAS [70], and DIAMOND

[71]. Operon structures were predicted using Operon-mapper [72]. CRISPR-Cas systems

were identified using CRISPRCasFinder (https://crisprcas.i2bc.paris-

saclay.fr/CrisprCasFinder/Index). Protein-coding sequences (CDSs) were assigned to clusters

of orthologous groups (COGs) using RPS-BLAST v2.13.0+ [73]. Principal component

analysis (PCA) was performed using the sklearn.decomposition module of the Scikit-learn

library [74]. Orthologous genes among the *Deferribacterota* genomes were identified using

OrthoFinder [75]. Principal coordinate analysis (PCoA) based on orthogroup composition was

performed using Phyloseq [76]. The functional annotation of each orthogroup was performed

using GhostKOALA [77]. The orthologous genes characteristically associated with a

phylogenetic group were identified using Scoary2 [78]. A maximum likelihood tree of the

flagellin genes was constructed based on the deduced amino acid sequences with >350 aa as

described above.

Results

Phylogenetic diversity and relative abundance of *Deferribacterota* in termite and

cockroach guts

The phylogenetic diversity and relative abundance of *Deferribacterota* in the guts of termites

and cockroaches were examined based on the 16S rRNA gene amplicon data derived from the

guts of diverse termite and cockroach species. We identified 72 ASVs assigned to

Deferribacterota from 47 out of the 68 termite and cockroach species (Fig. S1). Their relative

abundance was consistently low (0.00–1.29%). These ASVs formed a monophyletic cluster

with sequences derived from other insect guts (Fig. S1). The corresponding 16S rRNA region

of the SAG MS499-41 was identical to the sequence of ASV13241 detected in the guts of *R.*

speratus (relative abundance: 0.05%) and *Hodotermopsis* sp. (0.06%) (Fig. S1). A near full-

length 16S rRNA sequence, RsTz2-092 (TACW01000013), previously obtained from the

same *R. speratus* gut sample, showed 99.9% similarity to that of MS499-41. The bacterial

species represented by the SAG MS499-41, ASV13241, and RsTz2-092 was designated here

as "*Candidatus* Termitispirillum reticulitermitis" (abbreviated as *Te. reticulitermiti*s) and

described below.

Localization, morphology, and infection rate

We designed probe RsTz2-092-190 (Table S2), which is specific to the RsTz2-092 subclade

within the insect-gut clade (Fig. S1). FISH using this probe detected long, curved rods or

spiral-shaped bacteria, which were specifically attached to the cell surface of the parabasalid

protist *Holomastigotes* sp. (family *Spirotrichonymphidae*) in the gut of *R. speratus* (Fig. 1A,

B). Morphologically similar bacterial cells were also detected with this probe specifically on

the cell surface of *Holomastigotes* sp. (Fig. 1C, D) and *Brugerollina cincta*

(*Spirotrichonymphidae*) (Fig. 1E, F) in the gut of *Hodotermopsis sjostedti*. These

ectosymbionts appeared to be randomly distributed across the host cell surface (Fig. 1B, D, F)

together with spirochaetes, and their cell dimensions were ca. 3–10 µm in length and 0.4 µm

in width. Among the cells of the protist host species, 15.8–91.4% harbored ectosymbionts

detected with this probe (Table 1).

We then designed probe Deferri-term-661 (Table S2), which is specific to most members of

the insect-gut clade, except for the RsTz2-092 subclade (Fig. S1). When FISH with this probe

was performed for the gut content of *C. punctulatus*, spiral cells with dimensions 3–13 µm in

length and 0.4 µm in width were detected specifically on the anterior surface area of the

parabasalids *Trichonympha lata* (Fig. 1G, H) and *Trichonympha acuta* (Fig. S2A, B) (family

Trichonymphidae). Of the *T. lata* and *T. acuta* cells in a *C. punctulatus* gut, 67.6% harbored

the ectosymbiotic *Deferribacterota* (Table 1). Ectosymbiotic spirochaetes were also detected

on the same host protist cells by FISH using a spirochaete-specific probe (Figs. 1H and S2B;

Table S2). No specific signals were detected in the gut contents of *R. speratus* and *H. sjostedti*

with probe Deferri-term-661.

Phylogenetic analysis based on near full-length 16S rRNA genes

We collected single cells of these host protist species, i.e., *Holomastigotes* spp., *B. cincta*, *T.*

acuta, and *T. lata*, based on their morphological characteristics and performed WGA. Near

full-length 18S rRNA and 16S rRNA genes were amplified by PCR and the products were

cloned and subjected to Sanger sequencing. We confirmed the taxonomic assignment of the

host protist species based on the 18S rRNA genes (Fig. S3A, B). The 16S rRNA genes of

Deferribacterota formed a monophyletic cluster of sequences within the insect-gut clade (Fig.

2A). The clone RsHol6-1, obtained from *Holomastigotes* sp. in an *R. speratus* gut, showed

99.9% sequence similarity to the clone RsTz2-092. The clones HsHol1-2 from

Holomastigotes sp. and HsBr1-3 from *B. cincta* in the gut of *Hodotermopsis sjostedti* were

identical and showed 98.7% similarity to RsHol6-1 and RsTz2-092 (Fig. 2A). The clones

CpT15-2 and CpT06-9 were obtained from *T. acuta* and *T. lata*, respectively, in the gut of *C.*

punctulatus. These two sequences shared 97.9% identity with each other (Fig. 2A) and

showed ≥99.3% similarities to ASV149, which was detected in one *C. punctulatus* gut

(relative abundance: 0.21%) (Fig. S1).

General features of *Termitispirillum* genomes and phylogenomics

In addition to the SAG MS499-41 (1,578,178 bp, 79% estimated completeness, 44% GC

content) of *Te. reticulitermitis* from the gut of *R. speratus*, a circular chromosome (CpT32-4:

2,151,577 bp, 40% GC content) was reconstructed from a single *T. acuta* cell sample (Table

2). The 16S rRNA gene sequence of CpT32-4 shared 98.0% and 98.4% identity with CpT15-

2 and CpT06-9, respectively (Fig. 2A). The bacterium represented by the genome CpT32-4

was designated here as *Termitispirillum cryptocerci* and described below under SeqCode. The

total sequence length and GC content of the *Te. reticulitermitis* and *Te. cryptocerci* genomes

were similar to those of other insect-gut clade species (Fig. S4). The GC skew pattern of the

Te. cryptocerci genome is distinct from other genomes of *Deferribacterota*; the SkewI of *Te.*

cryptocerci is only 0.176, whereas the others range 0.733–0.914 (Fig. S5). The *Te.*

cryptocerci genome encodes CRISPR and associated genes (Table S3), whereas they were not

found in the *Te. reticulitermitis* genome.

The genomes of *Te. reticulitermitis* and *Te. cryptocerci* were most closely related to

metagenome-assembled genomes (MAGs) from the guts of higher termites [79, 80], which

generally do not harbor cellulolytic gut protists. These and two additional MAGs from lower

termites (Pcl387_bin.162 and Ly174_bin.14) [33] formed a monophyletic cluster (Fig. 2B).

This clade corresponds to the uncultured family "JAISNW01" in the GTDB and is

phylogenetically sister to the vertebrate-gut inhabitants *Mucispirillaceae* (Fig. 2B). The ANI

and AAI values between the *Te. reticulitermitis* and *Te. cryptocerci* genomes were <70% and

62.0%, respectively (Fig. S6A, B). Thus, these two bacteria are distinct species but in the

same genus according to the proposed criterion of genus-level similarity (ANI ≤ 95%; AAI ≥

60–85%) [61, 81].

Metabolic capacities of *Te. cryptocerci* and *Te. reticulitermitis*

Energy conservation and carbon metabolism

The *Te. cryptocerci* genome contains a complete set of genes for glycolysis and

gluconeogenesis (Fig. 3A, B). *Termitispirillum cryptocerci* likely imports monosaccharides

and sugar acids, such as fructose, xylose, mannose, N-acetylglucosamine, and uronate, via

phosphotransferase systems (PTS), ABC transporters, or transporters of major facilitator

superfamilies (MFS). It also likely imports and utilizes glycerol, lactate, citrate, and

unidentified carboxylates. Citrate can be metabolized to acetate and oxaloacetate by the action

of citrate lyase (CitDEF) (Fig. 3A). It is predicted that pyruvate is fermented to acetate,

ethanol, lactate, and H_2 using monomeric iron hydrogenase B1/B3 and electron-bifurcating

iron hydrogenase HydABC (Fig. 3A, B). The *Te. reticulitermitis* genome lacks two genes in

the glycolysis/gluconeogenesis pathway and all genes encoding the transporters for sugars

and other carbon sources, except for those transporting uronates and carboxylates (Figs 3A, B,

and S7), possibly due to its low completeness (Table 2). Thus, the main carbon source of *Te.*

reticulitermitis remains unclear. No genes encoding hydrolytic enzymes for extracellular

polysaccharides were identified in either genome.

In addition to substrate-level phosphorylation through glycolysis and oxidation to acetate,

Te. cryptocerci has the potential to generate ATP via the agmatine deiminase pathway with

the agmatine/putrescine antiporter AguD [82] (Fig. 3A). The *Te. cryptocerci* genome is

equipped with several systems to generate H^+/Na^+ membrane potential to produce ATP via

FoF_1-type ATPase. The genome encodes the membrane-bound [NiFe]-hydrogenase

HybOABC, the Rnf complex, Na^+-transporting oxaloacetate/methylmalonyl-CoA

decarboxylase (OadABC), and the H^+/Na^+ antiporter NhaB. It also encodes fumarate

reductase (FrdABCD) with the fumarate/succinate antiporter DcuB and the succinate exporter

DcuC, and in addition, cbb_3-type cytochrome c oxidase (Fig. 3A). All these components

except for NhaB and FrdABCD were identified in the *Te. reticulitermitis* genome. Thus, the

ectosymbionts likely use H_2 as an electron donor and fumarate and oxygen as electron

acceptors for respiration. Considering the very low concentration of oxygen in the hindguts of

termites and cockroaches, except for their peripheral regions [83], the cbb_3-type cytochrome c

oxidase may function not only in ATP synthesis via microaerobic respiration but also in

oxygen removal to protect anaerobic reactions. The *Te. reticulitermitis* genome encodes a

gene for periplasmic octaheme nitrite reductase (Onr) (Figs. 3A and S7), which possibly

contributes to the detoxification of nitrite by dissimilatory reduction to ammonium, as

suggested in *Desulfovibrio* species [84]. The genomes of these ectosymbionts contain several

other genes for oxygen tolerance (Fig. S7).

Nitrogen metabolism

No genes encoding a nitrogenase were identified in the *Te. cryptocerci* and *Te. reticulitermitis*

genomes. Both genomes encode several transporters for amino acids and di/oligo-peptides,

which are likely the main nitrogen sources (Fig. 3A). The *Te. cryptocerci* genome also

encodes transporters for nucleosides and uracil although both genomes apparently can de

novo synthesize nucleotides. Both genomes lack the biosynthetic pathways for branched-

chain amino acids, and only the *aroE, aroK, aroA, and aroC* genes were identified in the

pathway for aromatic amino acids. Although an operon for the tryptophan synthase genes

trpA and *trpB* was found in the *Te. cryptocerci* genome, both are pseudogenized.

Termitispirillum cryptocerci cannot synthesize histidine, whereas *Te. reticulitermitis*

possesses its complete biosynthetic pathway. These biosynthetic pathways, absent in *Te.*

cryptocerci and/or *Te. reticulitermitis*, are typically retained by most other *Deferribacterota*

(Fig. 3B). In the *Te. cryptocerci* genome, the biosynthetic pathways for riboflavin,

tetrahydrofolate, cobalamin, and heme are complete, whereas the biosynthesis of

pantothenate/coenzyme A (CoA), nicotinamide adenine dinucleotide (NAD), biotin,

pyridoxin 5'-phosphate, and menaquinone requires precursors (Fig. S7).

Sulfur metabolism

Termitispirillum cryptocerci potentially takes up sulfate, which is subsequently reduced to

sulfite via adenosine 5'-phosphosulfate (APS) and 3'-phosphoadenosine 5'-phosphosulfate

(PAPS). It is predicted that sulfite is reduced to sulfide by the action of assimilatory sulfite

reductase (Sir) in cooperation with the heterodisulfide reductase complex HdrABC-FlxABCD

(Fig. S8A, B) [85]. Sulfides can be assimilated into methionine and cysteine. No genes

involved in sulfur assimilation were detected in the *Te. reticulitermitis* genome (Fig. 3A, B).

Cell wall, secretion systems, and motility

Both genomes encode the biosynthetic pathways for peptidoglycan and lipopolysaccharides

and also contain the gene sets encoding proteins for flagellar assembly and chemotaxis,

similar to the cases in other *Deferribacterota* [3, 5, 6, 13] (Fig. 3A, B). Genes for the

secretory (Sec) system and the twin arginine translocation (Tat) pathway, which facilitate

protein transport, were present, whereas no other secretion systems were identified (Fig. 3A).

Comparison of genomic compositions with other members of Deferribacterota

PCA based on the relative abundance of COG categories exhibited the clustering of the gut-

inhabiting clades, i.e., the insect-gut clade, *Mucispirillaceae*, and "*Ca*. Microvillispirillaceae"

(Fig. 4). Among the COG categories, a decrease in the ratio and number of genes in category

[T] (signal transduction mechanisms) was prominent in *Te. cryptocerci* and *Te.*

reticulitermitis (Figs 4 and S9A, B). In PCoA based on the composition of all orthologs (Fig.

S10A, B), the genomes of the insect-gut clade formed a cluster and were separated from those

of other animal-gut inhabitants.

Most genomes of the insect-gut clade and *Mucispirillaceae* lack components for nitrogen

fixation and NADH dehydrogenase complex I (Fig. 3B). Among the *Deferribacterota*, *Te.*

cryptocerci exceptionally has capacity to use diverse carbon and energy sources (Fig. S7) and

to generate H_2 as a fermentation product (Fig. 3B). In addition, the agmatine deiminase

pathway is almost unique to *Te. cryptocerci* in *Deferribacterota*, and its ability to assimilate

sulfate is rare among the gut inhabitants. The biosynthetic capacity for amino acids and

cofactors is reduced in the insect-gut clade and especially in the ectosymbionts *Te.*

cryptocerci and *Te. reticulitermitis*. This tendency is also conspicuous in the shrimp-gut

symbiont "*Ca*. Rimicarispirillum atlantis" in "*Ca*. Microvillispirillaceae" (Figs 3B and S7).

Termitispirillum cryptocerci and *Te. reticulitermitis* possess an unusually high copy number

of flagellin genes compared with other *Deferribacterota* species (*Te. cryptocerci*: 40 copies,

Te. reticulitermitis: 15 copies, others: 0–7 copies) (Figs 3B and S11).

Discussion

In this study, we discovered the first examples of ectosymbiosis between *Deferribacterota*

and protists. The ectosymbiotic *Deferribacterota* are associated with specific parabasalid

protist species in the guts of termites and *Cryptocercus* cockroaches, although not all cells of

the host protist species are colonized. The monophyly of these bacteria suggests that their

ability to establish ectosymbiosis with protist cells may have evolved in the gut microbiota of

the common ancestor of termites and *Cryptocercus*, i.e., before ~150 million years [86, 87],

after the insect-gut clade diverged from *Mucispirillaceae*, a vertebrate gut-associated family

that includes *Mucispirillum schaedleri*, a member of the Schaedler flora [88].

The *Deferribacterota* detected in the guts of higher termites lacking cellulolytic gut protists

are most likely free-swimming bacteria with flagella (Fig. 3B). The ectosymbionts *Te.*

cryptocerci and *Te. reticulitermitis* possess genes encoding proteins for flagellar assembly

and chemotaxis; therefore, these bacteria also swim, possibly targeting host protists, and

attach to the host cell surfaces. The expansion of the number of flagellin gene variants in the

genomes of *Te. cryptocerci* and *Te. reticulitermitis*, up to 40, is noteworthy. In many bacteria,

flagella act not only for swimming but also for other roles such as adhesion and colonization

[89]. It has been suggested that such multiple functions of flagella are associated with the

structural variations of flagellin [90]. In addition, in pathogenic bacteria, shifts in the

expression of flagellin variants to avoid or confuse recognition of their flagella by the host

cell were previously reported [89]. We thus hypothesize that the large repertoire of flagellin

variants may be used for multiple functions, such as swimming and adhesion, and possibly

also to avoid recognition and elimination by the protist host.

Most protist cells in the guts of termites and *Cryptocercus* cockroaches harbor

ectosymbiotic bacteria of diverse lineages, which likely play various important roles in the

multilayered symbiotic system [26]. Our genome analyses indicated that *Te. cryptocerci* and

Te. reticulitermitis are not able to fix dinitrogen. In addition, they have limited capacity to

synthesize amino acids (Fig. 3B), whereas their ability to synthesize several cofactors, such as

riboflavin, cobalamin, and tetrahydrofolate, may contribute to the nutrition of the protist and

termite hosts (Fig. S7). Although *Te. cryptocerci* has the potential to assimilate diverse carbon

sources, including plant-derived monosaccharides and uronate, it does not possess secretory

enzymes for the hydrolysis of polysaccharides (Fig. 3A).

As *Te. cryptocerci* and *Te. reticulitermitis* possess the ability to consume H_2 as an energy

source and use O_2 as an electron acceptor, these ectosymbionts possibly play a role in the

removal of these molecules to promote the fermentative activities of their anaerobic protist

hosts. This is in concordance with the localization of *Te. cryptocerci* confined to the anterior

part of *Trichonympha* cells, where hydrogenosomes are densely localized [91], although this

restriction may alternatively reflect higher phagocytotic activity in the posterior region. This

localization pattern was previously observed in H_2-oxidizing ectosymbiotic *Desulfovibrio*

associated with *Trichonympha* in termite guts [36, 37]. Thus, the ectosymbiotic

Deferribacterota are potentially beneficial to their protist hosts, yet they are not consistently

maintained across all host cells. This suggests that they are not essential for the host survival.

The reduction in the number of signal transduction-related genes, with no prominent

genome size reduction, was previously found in several obligate ectosymbionts of protists in

the guts of termites and *Cryptocercus* cockroaches [37]. This trait is shared by *Te. cryptocerci*

and *Te. reticulitermitis*; it is implied that these bacteria almost entirely live as ectosymbionts,

with free-living or infectious stages before colonizing a host occurring only transiently. The

unusual GC skew pattern of *Te. cryptocerci* suggests a non-canonical chromosome replication

system and/or massive genome rearrangements, as previously found in endosymbiotic

bacteria of termite-gut protists [27, 55, 92] (Fig. S5). However, as no complete genomes

except for *Te. cryptocerci* were obtained, it remains unclear whether this GC skew pattern is a

common trait in the insect-gut clade or attributable to the ectosymbiosis. Additional genome

sequences of ectosymbiotic *Deferribacterota* and ecological characterization of other

members of the insect-gut clade will further provide insights into the evolution of

ectosymbiosis from free-swimming gut bacteria.

We here describe the novel genus *Termitispirillum* (represented by the type species *Te.*

cryptocerci) and a novel family, *Termitispirillaceae*, for the insect-gut clade, under the Code

of Nomenclature of Prokaryotes Described from Sequence Data (SeqCode) [93]. We also

propose the novel species "*Candidatus* Termitispirillum reticulitermitis" and "*Candidatus*

Termitispirillum hodotermopsidis", which lack high-quality genomic information, as

recommended for uncultured bacteria under the International Code of Nomenclature of

Prokaryotes [94].

Description of *Termitispirillaceae* fam. nov.

Termitispirillaceae (Ter.mi.ti.spi.ril.la'ce.ae. N.L. neut. n. *Termitispirillum*, the type genus of

the family; *-aceae*, suffix denoting a family; N.L. fem. pl. n. *Termitispirillaceae*, the family of

Termitispirillum). Members of this family are uncultured and inhabit the guts of insects,

including termites and *Cryptocercus* cockroaches. This family is a sister clade of the family

Mucispirillaceae in the phylum *Deferribacterota*. The type genus of this family is

Termitispirillum.

Description of *Termitispirillum* gen. nov.

Termitispirillum (Ter.mi.ti.spi.ril'lum. L. masc. n. *termes*, termite; L. neut. dim. n. *spirillum*, a

small spiral; N.L. neut. n. *Termitispirillum*, a spiral bacterium from termites). Members of this

genus are uncultured and spiral shaped or curved long rods that specifically attach to

parabasalid protists in the guts of termites and *Cryptocercus* cockroaches. Cells measure 3–13

μm in length and 0.4 μm in width. Genomic analyses indicated that these bacteria possess a

Gram-negative-type cell wall and chemoheterotrophic metabolism with fermentation and

respiratory pathways. Motility was inferred from the presence of genes involved in flagellar

assembly and chemotaxis. The type species is *Termitispirillum cryptocerci* with its genome

sequence as the type material.

Description of *Termitispirillum cryptocerci* sp. nov.

Termitispirillum cryptocerci (cryp.to.cer'ci. N.L. gen. n. *cryptocerci*, of *Cryptocercus*,

referring to the host cockroach genus). The bacterium specifically attaches to the cell surfaces

of *Trichonympha acuta* and *Trichonympha lata* in the gut of *Cryptocercus punctulatus*. The

cell dimensions are 3–13 µm by 0.4 µm. This assignment is based on the complete genome

sequence (AP043677) and detection by FISH using the oligonucleotide probe Deferri-term-

661 (Table S2). This species corresponds to CpT32-4.

Description of "*Candidatus* Termitispirillum reticulitermitis" sp. nov.

Termitispirillum reticulitermitis (re.ti.cu.li.ter'mi.tis. N.L. gen. masc. n. *reticulitermitis*, of

Reticulitermes, referring to the host termite genus). The bacterium specifically attaches to the

cell surface of *Holomastigotes* sp. in the gut of *Reticulitermes speratus*. The cell dimensions

are 3–10 µm by 0.4 µm. This assignment is based on the 16S rRNA gene sequence

(LC866559) and specific detection by FISH using the oligonucleotide probe RsTz2-092-190

(Table S2). The single-cell genome MS499-41 (BAAIHG010000001–BAAIHG010000254)

was obtained. This species includes the 16S rRNA phylotypes RsTz2-092 and RsHol6-1.

Description of "*Candidatus* Termitispirillum hodotermopsidis" sp. nov.

Termitispirillum hodotermopsidis (ho.do.ter.mop'si.dis. N.L. fem. n. *hodotermopsidis*, of

Hodotermopsis, referring to the host termite genus). The bacterium specifically attaches to the

cell surfaces of *Holomastigotes* sp. and *Brugerollina cincta* in the gut of *Hodotermopsis*

sjostedti. The cell dimensions are 3–10 µm by 0.4 µm. This assignment is based on the 16S

rRNA gene sequences (LC866557, LC866558) and specific detection by FISH using the

oligonucleotide probe RsTz2-092-190 (Table S2). This species includes the 16S rRNA

phylotypes HsHol1-2 and HsBr1-3.

Acknowledgements

We are grateful to Gaku Tokuda, Aram Mikaelyan, and Melbert Schwartz for cooperation in cockroach collection. We also thank Shu Onouchi, Hirokazu Kuwahara, and Kazuki Takahashi for assisting with experiments and Takumi Murakami for handling sequence data. Sanger sequencing was performed in the RIKEN CBS and the Biomaterial Analysis Center in the Tokyo Institute of Technology.

Author contributions

Naoya Maruoka: conceptualization, methodology, investigation, formal analysis, data curation, visualization, resource, writing – original draft, funding acquisition. Rinpei Kudo: investigation, data curation. Katsura Igai: investigation, data curation. Masahiro Yuki: methodology, investigation, data curation, resource. Michiru Shimizu: methodology, investigation, data curation, resource. Moriya Ohkuma: resource, writing – review and editing, funding acquisition. Yuichi Hongoh: conceptualization, methodology, resource, formal analysis, writing – original draft, writing – review and editing, funding acquisition, project administration, supervision.

Conflicts of interest

The authors declare no conflict of interest.

Funding

This study was financially supported by JST (Japan Science and Technology Agency) SPRING for PhD course students to N.M. (JPMJSP2106), by NEXT and KAKENHI grants from JSPS (Japan Society for the Promotion of Science) to Y.H. (GS009, 22241046,

Data availability

498   The genome sequences obtained in this study are available under the BioProject PRJDB20420

499   in DDBJ. The accession numbers of the SSU rRNA genes appears under DDBJ accession

500   numbers LC866555–64 and LC875758.

References

1.   Jumas-Bilak E, Roudière L, Marchandin, H. Despcription of "*Synergistetes*" phyl. nov.

and emended description of the phylum "*Deferribacteres*" and of the family

Syntrophomonadaceae, phylum "*Firmicutes*". *Int J Syst Evol Microbiol* 2009;**59**:1028–

35. https://doi.org/10.1099/ijs.0.006718-0

2.   Oren A, Garrity GM. Valid publication of the names of forty-two phyla of prokaryotes.

Int J Syst Evol Microbiol 2021;**71**:e005056. https://doi.org/10.1099/ijsem.0.005056

3.   Spring S, Rohde M, Bunk B *et al*. New insights into the energy metabolism and

taxonomy of *Deferribacteres* revealed by the characterization of a new isolate from a

hypersaline microbial mat. *Environ Microbiol* 2022;**24**:2543–75.

https://doi.org/10.1111/1462-2920.15999

4.   Zavarzina DG, Prokofeva MI, Pikhtereva VA *et al*. *Deferrivibrio essentukiensis* sp.

nov., gen. nov., a representative of *Deferrivibrionaceae* fam. nov., isolated from the

subsurface aquifer of caucasian mineral drinking waters. *Microbiology (Russian Fed)*

2022;**91**:143–59. https://doi.org/10.1134/S0026261722020114

5.   Aubé J, Cambon-Bonavita MA, Velo-Suárez L *et al*. A novel and dual digestive

symbiosis scales up the nutrition and immune system of the holobiont *Rimicaris*

exoculata. *Microbiome* 2022;**10**:e189. https://doi.org/10.1186/s40168-022-01380-2

6.   Takaki Y, Shimamura S, Nakagawa S *et al*. Bacterial lifestyle in a deep-sea

hydrothermal vent chimney revealed by the genome sequence of the thermophilic

bacterium *Deferribacter desulfuricans* SSM1. *DNA Res* 2010;**17**:123–37.

https://doi.org/10.1093/dnares/dsq005

7.   Rauschenbach I, Posternak V, Cantarella P *et al*. *Seleniivibrio woodruffii* gen. nov., sp.

nov., a selenate- and arsenate-respiring bacterium in the *Deferribacteraceae*. *Int J Syst*

Evol Microbiol 2013;**63**:3659–65. https://doi.org/10.1099/ijs.0.043547-0

8.   Iino T, Nakagawa T, Mori K *et al*. *Calditerrivibrio nitroreducens* gen. nov., sp. nov., a

thermophilic, nitrate-reducing bacterium isolated from a terrestrial hot spring in Japan.

Int J Syst Evol Microbiol 2008;**58**:1675–9. https://doi.org/10.1099/ijs.0.65714-0

9.   Tamazawa S, Mayumi D, Mochimaru H *et al*. *Petrothermobacter organivorans* gen.

nov., sp. nov., a thermophilic, strictly anaerobic bacterium of the phylum

Deferribacteres isolated from a deep subsurface oil reservoir. *Int J Syst Evol Microbiol*

2017;**67**:3982–6. https://doi.org/10.1099/ijsem.0.002234

10.   Robertson BR, O'Rourke JL, Neilan BA *et al*. *Mucispirillum schaedleri* gen. nov., sp.

nov., a spiral-shaped bacterium colonizing the mucus layer of the gastrointestinal tract

of laboratory rodents. *Int J Syst Evol Microbiol* 2005;**55**:1199–204.

https://doi.org/10.1099/ijs.0.63472-0

11.   Gilroy R, Ravi A, Getino M *et al*. Extensive microbial diversity within the chicken gut

microbiome revealed by metagenomics and culture. *PeerJ* 2021;**9**:e10941.

https://doi.org/10.7717/peerj.10941

12.   Scupham AJ, Patton TG, Bent E *et al*. Comparison of the cecal microbiota of domestic

542   and wild turkeys. *Microb Ecol* 2008;**56**:322–31. https://doi.org/10.1111/1462-

543   2920.70010

544   13.   Loy A, Pfann C, Steinberger M *et al*. Lifestyle and horizontal gene transfer-mediated

545   evolution of *Mucispirillum schaedleri*, a core member of the murine gut microbiota.

546   *mSystems* 2017;**2**:e00171-16. https://doi.org/10.1128/msystems.00171-16

547   14.   Durand L, Zbinden M, Cueff-Gauchard V *et al*. Microbial diversity associated with the

548   hydrothermal shrimp *Rimicaris exoculata* gut and occurrence of a resident microbial

549   community. *FEMS Microbiol. Ecol* 2010;**71**:291–303. https://doi.org/10.1111/j.1574-

550   6941.2009.00806.x

551   15.   Guéganton M, Rouxel O, Durand L *et al*. Anatomy and symbiosis of the digestive

552   system of the vent shrimp *Rimicaris exoculata* and *Rimicaris chacei* revealed through

553   imaging approaches. *Front Mar Sci* 2022;**9**:e903748.

554   https://doi.org/10.3389/fmars.2022.903748

555   16.   Qi L, Shi M, Zhu FC *et al*. Genomic evidence for the first symbiotic Deferribacterota, a

556   novel gut symbiont from the deep-sea hydrothermal vent shrimp *Rimicaris kairei*.

557   *Front Microbiol* 2023;**14**:e1179935. https://doi.org/10.3389/fmicb.2023.1179935

558   17.   Waite DW, Dsouza M, Biswas K *et al*. Microbial community structure in the gut of the

559   New Zealand insect Auckland tree weta (*Hemideina thoracica*). *Arch Microbiol*

560   2015;**197**:603–12. https://doi.org/10.1007/s00203-015-1094-3

561   18.   Schauer C, Thompson CL, Brune A. The bacterial community in the gut of the

562   cockroach *Shelfordella lateralis* reflects the close evolutionary relatedness of

563   cockroaches and termites. *Appl Environ Microbiol* 2012;**78**:2758–67.

564   https://doi.org/10.1128/AEM.07788-11

19.   Berlanga M, Llorens C, Comas J *et al.* Gut bacterial community of the xylophagous

cockroaches *Cryptocercus punctulatus* and *Parasphaeria boleiriana*. *PLOS ONE*

2016;**11**:e0152400. https://doi.org/10.1371/journal.pone.0152400

20.   Warnecke F, Luginbühl P, Ivanova N *et al.* Metagenomic and functional analysis of

hindgut microbiota of a wood-feeding higher termite. *Nature* 2007;**450**:560–5.

https://doi.org/10.1038/nature06269

21.   Liu N, Zhang L, Zhou H *et al.* Metagenomic insights into metabolic capacities of the

gut microbiota in a fungus-cultivating termite (*Odontotermes yunnanensis*). *PLOS*

ONE 2013;**8**:e69184. https://doi.org/10.1371/journal.pone.0069184

22.   King JHP, Mahadi NM, Bong CFJ *et al*. Bacterial microbiome of *Coptotermes*

curvignathus (Isoptera: Rhinotermitidae) reflects the coevolution of species and dietary

pattern. *Insect Sci.* 2014;**21**:584–96. https://doi.org/10.1111/1744-7917.12061

23.   Brune A. Symbiotic digestion of lignocellulose in termite guts. *Nat. Rev. Microbiol.*

2014;**12**:168–80. https://doi.org/10.1038/nrmicro3182

24.   Cleveland LR, Hall SR, Sanders EP *et al.* The wood-feeding roach *Cryptocercus*, its

protozoa, and the symbiosis between protozoa and roach. *Mem Am Acad Arts Sci*

1934;**17**:185–342. https://doi.org/10.2307/25058198

25.   Ohkuma M, Brune A. Diversity, structure, and evolution of the termite gut microbial

community. In: Bignell D, Roisin Y, Lo N (ed.), *Biology of Termites: a Modern*

Synthesis. Switzerland: Springer Nature, 2011, 413–38. https://doi.org/10.1007/978-90-

481-3977-4

26.   Hongoh Y, Ohkuma M. Termite gut flagellates and their methanogenic and eubacterial

symbionts. In: Hackstein JHP (ed.), *(Endo)symbiotic Methanogenic Archaea*.

Switzerland: Springer Nature, 2018, 55–80. https://doi.org/10.1007/978-3-319-98836-

8_5

27. Hongoh Y, Sharma VK, Prakash T *et al.* Complete genome of the uncultured Termite

Group 1 bacteria in a single host protist cell. *Proc Natl Acad Sci USA* 2008;**105**:5555–

60. https://doi.org/10.1073/pnas.0801389105

28. Hongoh Y, Sharma VK, Prakash T *et al.* Genome of an endosymbiont coupling N_2

fixation to cellulolysis within protist cells in termite gut. *Science* 2008;**322**:1108–9.

https://doi.org/10.1126/science.1165578

29. Ohkuma M, Noda S, Hattori S *et al.* Acetogenesis from H_2 plus CO_2 and nitrogen

fixation by an endosymbiotic spirochete of a termite-gut cellulolytic protist. *Proc Natl*

Acad Sci USA 2015;**112:**10224–30. https://doi.org/10.1073/pnas.1423979112

30. Tai V, Carpenter KJ, Weber PK *et al.* Genome evolution and nitrogen fixation in

bacterial ectosymbionts of a protist inhabiting wood-feeding cockroaches. *Appl*

Environ Microbiol 2016;**82**:4682–95. https://doi.org/10.1128/AEM.00611-16

31. Strassert JFH, Mikaelyan A, Woyke T *et al.* Genome analysis of '*Candidatus* Ancillula

trichonymphae', first representative of a deep-branching clade of Bifidobacteriales,

strengthens evidence for convergent evolution in flagellate endosymbionts. *Environ*

Microbiol Rep 2016;**8**:865–73. https://doi.org/10.1111/1758-2229.12451

32. Treitli SC, Kolisko M, Husník F *et al.* Revealing the metabolic capacity of

Streblomastix strix and its bacterial symbionts using single-cell metagenomics. *Proc*

Natl Acad Sci USA 2019;**116**:19675–84. https://doi.org/10.1073/pnas.1910793116

33. Mies US, Hervé V, Kropp T *et al.* Genome reduction and horizontal gene transfer in

the evolution of Endomicrobia—rise and fall of an intracellular symbiosis with termite

gut flagellates. *mBio* 2024;**15**:e00826-24. https://doi.org/10.1128/mbio.00826-24

34. Silva JK, Hervé V, Mies US *et al.* A novel lineage of endosymbiotic *Actinomycetales*:

genome reduction and acquisition of new functions in *Bifidobacteriaceae* associated

with termite gut flagellates. *Environ Microbiol* 2025;**27**:e70010.

https://doi.org/10.1111/1462-2920.70010

35.   Ikeda-Ohtsubo W, Strassert JFH, Köhler T *et al.* '*Candidatus* Adiutrix intracellularis',

an endosymbiont of termite gut flagellates, is the first representative of a deep-

branching clade of *Deltaproteobacteria* and a putative homoacetogen. *Environ*

Microbiol 2016;**18**:2548–64. https://doi.org/10.1111/1462-2920.13234

36.   Kuwahara H, Yuki M, Izawa K *et al.* Genome of '*Ca.* Desulfovibrio trichonymphae',

an H_2-oxidizing bacterium in a tripartite symbiotic system within a protist cell in the

termite gut. *ISME J* 2017;**11**:766–76. https://doi.org/10.1038/ismej.2016.143

37.   Takeuchi M, Kuwahara H, Murakami T *et al.* Parallel reductive genome evolution in

Desulfovibrio ectosymbionts independently acquired by *Trichonympha* protists in the

termite gut. *ISME J* 2020;**14**:2288–301. https://doi.org/10.1038/s41396-020-0688-1

38.   Kaneko M, Omori T, Igai K *et al.* Facultative endosymbiosis between cellulolytic

protists and methanogenic archaea in the gut of the Formosan termite *Coptotermes*

formosanus. *ISME Commun* 2024;**4**:ycae097. https://doi.org/10.1093/ismeco/ycae097

39.   Aoki H, Yuki M, Shimizu M *et al.* Agarose gel microcapsules enable easy-to-prepare,

picolitre-scale, single-cell genomics, yielding high-coverage genome sequences. *Sci*

Rep 2022;**12**:e17014. https://doi.org/10.1038/s41598-022-20923-z

40.   Utami YD, Kuwahara H, Murakami T *et al.* Phylogenetic diversity and single-cell

genome analysis of "*Melainabacteria*", a non-photosynthetic cyanobacterial group, in

the termite gut. *Microbes Environ* 2018;**33**:50–7.

https://doi.org/10.1264/jsme2.ME17137

41.   Pruesse E, Peplies J, Glöckner FO. SINA: accurate high-throughput multiple sequence

alignment of ribosomal RNA genes. *Bioinformatics* 2012;**28**:1823–9.

https://doi.org/10.1093/bioinformatics/bts252

42.   Katoh K, Misawa K, Kuma KI *et al.* MAFFT: a novel method for rapid multiple

sequence alignment based on fast Fourier transform. *Nucleic Acids Res* 2002;**30**:3059–

66. https://doi.org/10.1093/nar/gkf436

43.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated

alignment trimming in large-scale phylogenetic analyses. *Bioinformatics*

2009;**25**:1972–3. https://doi.org/10.1093/bioinformatics/btp348

44.   Nguyen LT, Schmidt HA, von Haeseler A *et al.* IQ-TREE: a fast and effective

stochastic algorithm for estimating maximum-likelihood phylogenies. *Mol Biol Evol*

2015;**32**:268–74. https://doi.org/10.1093/molbev/msu300

45.   Kalyaanamoorthy S, Minh BQ, Wong TKF *et al*. ModelFinder: fast model selection for

accurate phylogenetic estimates. *Nat Methods* 2017;**14**:587–9.

https://doi.org/10.1038/nmeth.4285

46.   Trager W. The cultivation of a cellulose-digesting flagellate, *Trichomonas termopsidis*,

and of certain other termite protozoa. *Biol Bull* 1934;**66**:182–90.

47.   Noda S, Ohkuma M, Yamada A *et al.* Phylogenetic position and in situ identification

of ectosymbiotic spirochetes on protists in the termite gut. *Appl Environ Microbiol*

2003;**69**:625–33. https://doi.org/10.1128/AEM.69.1.625-633.2003

48.   Murakami T, Onouchi S, Igai K *et al.* Ectosymbiotic bacterial microbiota densely

colonize the surface of thelastomatid nematodes in the gut of the wood-feeding

cockroach *Panesthia angustipennis*. *FEMS Microbiol Ecol* 2019;**95**:fiy238.

https://doi.org/10.1093/femsec/fiy238

49.   Ludwig W, Strunk O, Westram R *et al.* ARB: a software environment for sequence

data. *Nucleic Acids Res* 2004;**32**:1363–71. https://doi.org/10.1093/nar/gkh293

50.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing

reads. *EMBnet J* 2011;**17**:10–12. https://doi.org/10.14806/ej.17.1.200

51.   Schmieder R, Edwards R. Quality control and preprocessing of metagenomic datasets.

Bioinformatics 2011;**27**:863–4. https://doi.org/10.1093/bioinformatics/btr026

52.   Bankevich A, Nurk S, Antipov D *et al.* SPAdes: a new genome assembly algorithm

and its applications to single-cell sequencing. *J Comput Biol* 2012;**19**:455–77.

https://doi.org/10.1089/cmb.2012.0021

53.   von Meijenfeldt FAB, Arkhipova K, Cambuy DD *et al.* Robust taxonomic

classification of uncharted microbial sequences and bins with CAT and BAT. *Genome*

Biol 2019;**20**:e217. https://doi.org/10.1186/s13059-019-1817-x

54.   Camacho C, Coulouris G, Avagyan V *et al.* BLAST+: architecture and applications.

BMC Bioinformatics 2009;**10**:e421. https://doi.org/10.1186/1471-2105-10-421

55.   Takahashi K, Kuwahara H, Horikawa Y *et al.* Emergence of putative energy parasites

within *Clostridia* revealed by genome analysis of a novel endosymbiotic clade. *ISME J*

2023;**17**:1895–1906. https://doi.org/10.1038/s41396-023-01502-0

56.   Steinig E, Coin L. Nanoq: ultra-fast quality control for nanopore reads. *J Open Source*

Softw 2022;**7**:e2991. https://doi.org/10.21105/joss.02991

57.   Vaser R, Šikić M. Time- and memory-efficient genome assembly with Raven. *Nat*

Comput Sci 2021;**1**:332–6. https://doi.org/10.1038/s43588-021-00073-4

58.   Walker BJ, Abeel T, Shea T *et al.* Pilon: an integrated tool for comprehensive

microbial variant detection and genome assembly improvement. *PLOS ONE*

2014;**9**:e112963. https://doi.org/10.1371/journal.pone.0112963

59.   Chklovski A, Parks DH, Woodcroft BJ *et al.* CheckM2: a rapid, scalable and accurate

tool for assessing microbial genome quality using machine learning. *Nat Methods*

2023;**20**:1203–12. https://doi.org/10.1038/s41592-023-01940-w

60.   Grant JR, Enns E, Marinier E *et al.* Proksee: in-depth characterization and visualization

of bacterial genomes. *Nucleic Acids Res* 2023;**51**:W484–92.

https://doi.org/10.1093/nar/gkad326

61.   Jain C, Rodriguez-R LM, Phillippy AM *et al.* High throughput ANI analysis of 90K

prokaryotic genomes reveals clear species boundaries. *Nat Commun* 2018;**9**:e5114.

https://doi.org/10.1038/s41467-018-07641-9

62.   Kim D, Park S, Chun J. Introducing EzAAI: a pipeline for high throughput calculations

of prokaryotic average amino acid identity. *J Microbiol* 2021;**59**:476–80.

https://doi.org/10.1007/s12275-021-1154-0

63.   Lu J, Salzberg SL. SkewIT: the Skew Index Test for large-scale GC Skew analysis of

bacterial genomes. *PLOS Comput Biol* 2020;**16**:e1008439.

https://doi.org/10.1371/journal.pcbi.1008439

64.   Lee MD. GToTree: a user-friendly workflow for phylogenomics. *Bioinformatics*

2019;**35**:4162–4. https://doi.org/10.1093/bioinformatics/btz188

65.   Hyatt D, Chen G, LoCascio PF *et al.* Prodigal: prokaryotic gene recognition and

translation initiation site identification. *BMC Bioinformatics* 2010;**11**:e119.

https://doi.org/10.1186/1471-2105-11-119

66.   Eddy SR. Accelerated profile HMM searches. *PLOS Comput. Biol.* 2011;**7**:e1002195.

https://doi.org/10.1371/journal.pcbi.1002195

67.   Edgar RC. Muscle5: high-accuracy alignment ensembles enable unbiased assessments

of sequence homology and phylogeny. *Nat Commun* 2022;**13**:e6989.

https://doi.org/10.1038/s41467-022-34630-w

68.   Parks DH, Chuvochina M, Rinke C *et al.* GTDB: an ongoing census of bacterial and

archaeal diversity through a phylogenetically consistent, rank normalized and complete

genome-based taxonomy. *Nucleic Acids Res* 2022;**50**:D785–94.

https://doi.org/10.1093/nar/gkab776

69.   Tanizawa Y, Fujisawa T, Nakamura Y. DFAST: a flexible prokaryotic genome

annotation pipeline for faster genome publication. *Bioinformatics* 2018;**34**:1037–9.

https://doi.org/10.1093/bioinformatics/btx713

70.   Moriya Y, Itoh M, Okuda S *et al.* KAAS: an automatic genome annotation and

pathway reconstruction server. *Nucleic Acids Res* 2007;**35**:182–5.

https://doi.org/10.1093/nar/gkm321

71.   Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.

Nat Methods 2015;**12**:59–60. https://doi.org/10.1038/nmeth.3176

72.   Taboada B, Estrada K, Ciria R *et al.* Operon-mapper: a web server for precise operon

identification in bacterial and archaeal genomes. *Bioinformatics* 2018;**34**:4118–20.

https://doi.org/10.1093/bioinformatics/bty496

73.   Marchler-Bauer A, Panchenko AR, Shoemarker BA *et al.* CDD: a database of

conserved domain alignments with links to domain three-dimensional structure.

Nucleic Acids Res 2002;**30**:281–3. https://doi.org/10.1093/nar/30.1.281

74.   Pedregosa F, Varoquaux G, Gramfort A *et al.* Scikit-learn: machine learning in Python.

J Mach Learn Res 2011;**12**:2825–30.

75.   Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative

genomics. *Genome Biol* 2019;**20**:e238. https://doi.org/10.1186/s13059-019-1832-y

76.   McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis

and graphics of microbiome census data. *PLOS ONE* 2013;**8**:e61217.

https://doi.org/10.1371/journal.pone.0061217

77.   Kanehisa M, Sato Y, Morishima K. BlastKOALA and GhostKOALA: KEGG tools for

735   functional characterization of genome and metagenome sequences. *J Mol Biol*

736   2016;4**28**:726–31. https://doi.org/10.1016/j.jmb.2015.11.006

737   78.   Roder T, Pimentel G, Fuchsmann P *et al.* Scoary2: rapid association of phenotypic

738   multi-omics data with microbial pan-genomes. *Genome Biol* 2024;**25**:e93.

739   https://doi.org/10.1186/s13059-024-03233-7

740   79.   Hervé V, Liu P, Dietrich C *et al.* Phylogenomic analysis of 589 metagenome-

741   assembled genomes encompassing all major prokaryotic lineages from the gut of

742   higher termites. *PeerJ* 2020;**8**:e8614. https://doi.org/10.7717/peerj.8614

743   80.   Arora J, Kinjo Y, Šobotník J *et al.* The functional evolution of termite gut microbiota.

744   *Microbiome* 2022;**10**:e78. https://doi.org/10.1186/s40168-022-01258-3

745   81.   Riesco R, Trujillo ME. Update on the proposed minimal standards for the use of

746   genome data for the taxonomy of prokaryotes. *Int J Syst Evol Microbiol*

747   2024;**74**:e006300. https://doi.org/10.1099/ijsem.0.006300

748   82.   del Rio B, Linares DM, Ladero V *et al.* Putrescine production via the agmatine

749   deiminase pathway increases the growth of *Lactococcus lactis* and causes the

750   alkalinization of the culture medium. *Appl Microbiol Biotechnol* 2015;**99**:897–905.

751   https://doi.org/10.1007/s00253-014-6130-8

752   83.   Brune A. Termite guts: the world's smallest bioreactors. *Trends Biotechnol* 1998;**16**:

753   16–21. https://doi.org/10.1016/S0167-7799(97)01151-7

754   84.   Besson S, Almeida MG, Silveira CM. Nitrite reduction in bacteria: a comprehensive

755   view of nitrite reductases. *Coord Chem Rev* 2022;**464**:e214560.

756   https://doi.org/10.1016/j.ccr.2022.214560

757   85.   Ramos AR, Grein F, Oliveira GP *et al.* The FlxABCD-HdrABC proteins correspond to

758   a novel NADH dehydrogenase/heterodisulfide reductase widespread in anaerobic

bacteria and involved in ethanol metabolism in *Desulfovibrio vulgaris* Hildenborough.

Environ Microbiol 2015;**17**:2288–305. https://doi.org/10.1111/1462-2920.12689

86. Bourguignon T, Lo N, Cameron SL *et al.* The evolutionary history of termites as

inferred from 66 mitochondrial genomes. *Mol. Biol. Evol.* 2015;**32**:406–21.

https://doi.org/10.1093/molbev/msu308

87. Evangelista DA, Wipfler B, Béthoux O *et al.* An integrative phylogenomic approach

illuminates the evolutionary history of cockroaches and termites (Blattodea). *Proc R*

Soc B Biol Sci 2019;**286**:e20182076. https://doi.org/10.1098/rspb.2018.2076

88. Wymore Brand M, Wannemuehler MJ, Phillips GJ, Proctor A, Overstreet AM, Jergens

AE, et al. The altered Schaedler flora: continued applications of a defined murine

microbial community. *ILAR J* 2015;**56**:169–78. https://doi.org/10.1093/ilar/ilv012

89. Chaban B, Hughes HV, Beeby M. The flagellum in bacterial pathogens: for motility

and a whole lot more. *Semin Cell Dev Biol* 2015;**46**:91–103.

https://doi.org/10.1016/j.semcdb.2015.10.032

90. Fields JL, Zhang H, Bellis NF *et al.* Structural diversity and clustering of bacterial

flagellar outer domains. *Nat Commun* 2024;**15**:e9500. https://doi.org/10.1038/s41467-

024-53923-w

91. Carpenter KJ, Chow L, Keeling PJ. Morphology, phylogeny, and diversity of

Trichonympha (Parabasalia: Hypermastigida) of the wood-feeding cockroach

Cryptocercus punctulatus. *J Eukaryot Microbiol* 2009;**56**:305–13.

https://doi.org/10.1111/j.1550-7408.2009.00406.x

92. Izawa K, Kuwahara H, Kihara K *et al.* Comparison of intracellular "*Ca.*

Endomicrobium trichonymphae" genomovars illuminates the requirement and decay of

defense systems against foreign DNA. *Genome Biol Evol* 2016;**8**:3099–107.

https://doi.org/10.1093/gbe/evw227

93. Hedlund BP, Chuvochina M, Hugenholts P. *et al*. SecCode: a nomenclatural code for

prokaryotes described from sequence data. *Nat Microbiol* 2022;**7**:1702–8.

https://doi.org/10.1038/s41564-022-01214-9

94. Murray RG, Stackebrandt E. Taxonomic note: implementation of the provisional status

Candidatus for incompletely described procaryotes. *Int J Syst Bacteriol* 1995;**45**:186–

7. https://doi.org/10.1099/00207713-45-1-18

Legends to Figures

Figure 1. Detection of *Deferribacterota* associated with protists in the guts of termites and a

Cryptocercus cockroach. Phase-contrast images of (A) *Holomastigotes* sp. from

Reticulitermes speratus, (C) *Holomastigotes* sp. from *Hodotermopsis sjostedti*, (E)

Brugerollina cincta from *H. sjostedti*, and (G) *Trichonympha lata* in *Cryptocercus*

punctulatus are shown. *Deferribacterota* were detected by FISH using probe RsTz2-092-190

labeled with 6FAM (green) (B, D, F) and probe Deferri-term-661 labeled with 6FAM (green)

(H), respectively. Spirochaetes were detected using probe Spiro-36 labeled with Texas red

(red), and the images were overlayed (B, D, F, H). Bars indicate 10 μm and 5 μm (inset for

magnification).

Figure 2. Phylogenetic positions of the ectosymbiotic *Deferribacterota*. (A) 16S rRNA gene

tree constructed using 993 nucleotide sites with the GTR+F+I+G4 substitution model and an

outgroup (AB729138). (B) Phylogenomic tree constructed using 12,478 amino acid sites of

concatenated, partitioned 74 marker genes, with substitution models selected for each part and

an outgroup (GCF_000469585 and GCF_000177635). Family-level taxonomic codes

according to GTDB are indicated in parentheses. Filled and open circles indicate nodes with

ultrafast bootstrap value ≥95% and ≥80% in panel A, and 100% and ≥90% in panel B,

respectively. Asterisks indicate sequences obtained by Arora et al. (2022).

Figure 3. Predicted metabolic capacities of *Termitispirillum cryptocerci* (CpT32-4) and

"*Candidatus* Termitispirillum reticulitermitis" (MS499-41). (A) Outline of predicted

metabolic pathways based on genome sequences. Pathways detected exclusively in *Te.*

cryptocerci or *Te. reticulitermitis* are indicated in blue and red, respectively. (B) Comparisons

of repertoires of genes and pathways among members of *Deferribacterota*. The completeness

of pathways is indicated in a heatmap. Estimated genome completeness, total sequence length,

and the number of flagellin genes are also shown.

Figure 4. Comparation of genomic contents among members of *Deferribacterota*, including

Termitispirillum cryptocerci (CpT32-4) and "*Candidatus* Termitispirillum reticulitermitis"

(MS499-41). Principal component analysis based on the relative abundance of the clusters of

orthologous groups (COGs) was performed. The COG categories: [C] Energy production and

conversion, [D] Cell cycle control, cell division, chromosome partitioning, [E] Amino acid

transport and metabolism, [F] Nucleotide transport and metabolism, [G] Carbohydrate

transport and metabolism, [H] Coenzyme transport and metabolism, [I] Lipid transport and

metabolism, [J] Translation, ribosomal structure and biogenesis, [K] Transcription, [L]

Replication, recombination and repair, [M] Cell wall/membrane/envelope biogenesis, [N] Cell

motility, [O] Posttranslational modification, protein turnover, chaperones, [P] Inorganic ion

transport and metabolism, [Q] Secondary metabolites biosynthesis, transport and catabolism,

[R] General function prediction only, [S] Function unknown, [T] Signal transduction

mechanisms, [U] Intracellular trafficking, secretion, and vesicular transport, [V] Defense

mechanisms, [W] Extracellular structures, [X] Mobilome: prophages, transposons.

Table 1. Host protist species and frequency of association with *Deferribacterota*.

Host termite and cockroach	Host protist	Ratio of associated protist cells*
Reticulitermes speratus (from Ogose)	*Holomastigotes* sp.	91.4% (n = 35)
R. speratus (from Tsukuba)	*Holomastigotes* sp.	15.8% (n = 19)
Hodotermopsis sjostedti	*Holomastigotes* sp.	44.4% (n = 27)
	Brugerollina cincta	62.9% (n = 35)
Cryptocercus punctulatus	*Trichonympha lata* and *Trichonympha acuta***	67.6% (n = 34)

* "n" means the number of surveyed cells of the host protist species.

** Counted together because of difficulty in clearly discriminating these two species in a FISH specimen.

Table 2. General genome features of MS499-41, CpT32-4, and related *Deferribacterota*.

	MS499-41	CpT32-4	*Mucispirillum schaedleri*	"Rimicarispirillum atlantis"
Family	Insect-gut clade	Insect-gut clade	*Mucispirillaceae*	"Microvillispirillaceae"
Localization	Ectosymbiont of *Holomastigotes* sp.	Ectosymbiont of *T. acuta*	Mucus layer in the mouse gut	Space among microvilli in the shrimp gut
Total length (Mb)	1.58	2.15	2.35	1.34
Contigs	254	1 (circular)	1 (circular)	49
Completeness (%)*	78.6	Complete	Complete	90.7
Contamination (%)*	0.5	-	-	0.0
GC content (%)	44.1	39.7	31.1	47.4
CDS	1,363	1,931	2,145	1,072
Coding density (%)	78.8	89.3	87.9	84.5
rRNA	4	6	9	3
tRNA	36	40	42	33

* Estimated using CheckM2.

A

- *Flexistipes sinusarabici* isolated from deep-sea brine water NR_074881 — **Flexistipitaceae**
- M_Rainbow_MAG_00001 from vent shrimp gut CAMZNS010000014 — **"Ca. Microvillispirillaceae"**
- *Deferribacter autotrophicus* isolated from deep-sea hydrothermal vent EU407777 — **Deferribacteraceae**
- *Deferribacter desulfuricans* isolated from deep-sea hydrothermal vent AP011529
- *Calditerrivibrio nitroreducens* isolated from hot spring water AB364234 — **Calditerrivibrionaceae**
- *Petrothermobacter organivorans* isolated from deep subsurface oil reservoir LC155975
- *Deferrivibrio essentukiensis* isolated from meniral water deposit JAJAFU000000000 — **Deferrivibrionaceae**
- *Seleniivibrio woodruffii* isolated from activated sludge JN107635 — **Geovibrionaceae**
- *Denitrovibrio acetiphilus* isolated from oil reservoir model column AF146526
- *Geovibrio thiophilus* isolated from methanogenic mixed culture AJ299402
- *Limisalsivibrio acetivorans* isolated from hypersaline microbial mat KC631812
- *Mucispirillum schaedleri* isolated from mouse gut AY387670 — **Mucispirillaceae**

Insect-gut clade
- OTU76 from primary infected root canals of humans OR045488
- Pn-107 from *Pericapritermes nitobei* TACT01000007
- Mm-134 from *Macrotermes malaccensis* TACC01000043
- Gs-38 from *Glyptotermes satsumensis* TABQ01000056
- MgKI3d002E04 from *Coptotermes curvignathus* KP690862
- HETt-144 from *Heterotermes tenuior* TABS01000001
- HETt-113 from *H. tenuior* TABS01000018
- ATW054 from *Hemideina thoracia* KF318238
- PBg1-126 from *Protaetia brevitarsis* AY791243
- SL69 from *Shelfordella lateralis* JN680688
- Of-092 from *Odontotermes formosanus* TACL01000070
- Of-050 from *O. formosanus* TACL01000069
- 08102902-Y3GUT-16S-1 15 from *Odontotermes yunnanensis* JN619912
- MgMjR-062 from *Macrotermes gilvus* AB234550
- Of-099 from *O. formosanus* TACL01000071
- ORi-134 from *Oriensubulitermes inanis* TACK01000020
- 290cost002-P3L-421 from *Nasutitermes* sp. EF454802
- LONI-063 from *Longipeditermes longipes* TABZ01000024
- **CpT06-9 from *Trichonympha lata* in *Cryptocercus punctulatus* LC866555**
- **CpT15-2 from *Trichonympha acuta* in *C. punctulatus* LC866556**
- **CpT32-4 from *Trichonympha acuta* in *C. punctulatus* AP043677**
- **HsHol1-2 from *Holomastigotes* sp. in *Hodotermopsis sjostedti* LC866557**
- **HsBr1-3 from *Brugerollina cincta* in *H. sjostedti* LC866558**
- RsTz2-092 from *Reticulitermes speratus* TACW01000013
- **RsHol6-1 from *Holomastigotes* sp. in *R. speratus* LC866559**
- **MS499-41 from *R. speratus* BAAIHG010000001**

0.1

B

- "Ca. Rimicarispirillum atlantis" from vent shrimp gut GCA_947313855 — **"Ca. Microvillispirillaceae" (f__CAMZNJ01)**
- M2_7_5_bin.3 from production fluids GCA_020054865
- Deferribacteraceae bacterium V6Fe1 isolated from shallow hydrothermal vent GCA_022813675 — **Deferrivibrionaceae**
- *Deferrivibrio essentukiensis* isolated from mineral water deposit GCA_020480685
- W28 from hot spring water GCA_020217115 — **Calditerrivibrionaceae**
- ZAV-05 from hot spring water GCA_002878065
- *Calditerrivibrio nitroreducens* isolated from hot spring water GCA_000183405
- UBA1873 from crude oil GCA_002338145 — **Flexistipitaceae**
- *Flexistipes sinusarabici* isolated from deep-sea brine water GCA_000218625
- UBA4806 from tailing pond GCA_002402635
- *Deferribacter autotrophicus* isolated from deep-sea hydrothermal fields GCA_008362905 — **Deferribacteraceae**
- *Deferribacter desulfuricans* isolated from deep-sea hydrothermal vent GCA_000010985
- *Limisalsivibrio acetivorans* from hypersaline cyanobacterial mat GCA_000421105
- Deferri_01 from cellulose digester GCA_001603805
- OH_SFB_213 from wild boar feces GCA_023395255
- *Geovibrio thiophilus* isolated from drainage ditch GCA_004087915
- *Denitrovibrio acetiphilus* from oil reservoir model column GCA_000025725 — **Geovibrionaceae**
- UBA6799 from waste water GCA_002452335
- *Seleniivibrio woodruffii* isolated from activated sludge GCA_004339245
- UBA6793 from waste water GCA_002452405
- S2C1195 from waste water GCA_021209385
- BRZ_RS__bin62 from chicken gut GCA_944325215
- BRZ_AT__bin62 from chicken gut GCA_944324495 — **Mucispirillaceae**
- "Ca. Mucispirillum faecigallinarum" from chicken feces GCA_019116905
- BRZ_NI__bin29 from chicken gut GCA_944326705
- *Mucispirillum schaedleri* isolated from mouse gut GCA_000487995
- Pcl387_bin.162 from *Procryptotermes leewardensis* GCA_031278965
- Iy174_bin.14 from *Incisitermes snyderi* GCA_031276055
- Coa324_bin.33 from *Coatitermes kartaboensis* GCA_031256375 — **Insect-gut clade (f__JAISNW01)**
- 272-90.bin.133 from *Pericapritermes parvus**
- TD116_bin.76 from *Odontotermes* sp. GCA_031285415
- 301-30.bin.171 from *Odontotermes* sp.*
- **CpT32-4 from *Trichonympha acuta* in *Cryptocercus punctulatus***
- **MS499-41 from *Reticulitermes speratus***

0.1