Unique Unbiased Median Solution for Even Sample Sizes

Abstract

Data in experimental biology are frequently marred by outliers and asymmetric distributions. The median, being a robust estimator of central tendency, is less sensitive to outliers than the mean. However, for ranked datasets with an even number of observations, the conventional median—calculated as the average of the two middle values—can introduce bias by implicitly assuming symmetry in the data distribution. This study aims to identify a median estimator that is unbiased.

To derive the unbiased median estimator, we minimized the sum of residuals raised to a rational power approaching one. We compared the properties of the unbiased and conventional medians using Poisson-distributed datasets. Random samples were generated with the Mersenne Twister algorithm implemented in IgorPro software (WaveMetrics Inc., Oregon).

For odd sample sizes, the unbiased median coincides with the conventional median (the middle value). For even sample sizes, the unbiased median is defined as the value that equalizes the product of distances to data points above and below it—a definition that differs from the conventional median in asymmetric distributions. Although both median estimators tend to underestimate the mean of Poisson-distributed data, the unbiased median is consistently closer to the expected value. Additionally, the unbiased median exhibits lower variance compared to the conventional median.

Thus, for even sample sizes, the proposed unbiased median provides a central tendency measure that is unbiased, more accurate, and has reduced variance relative to the conventional median.

Keywords: *unbiased median, Poisson, well-posed problem.*

Introduction

The median was first introduced in 1755 by the distinguished physicist, astronomer, and mathematician Roger Joseph Boscovich as a method to fit a line to astronomical observations (Stigler, 1990). Boscovich sought the value that minimizes the sum of absolute deviations, discovering that this value separates the lower half of a dataset from the upper half. For ordinal data—where elements are ranked relative to one another but not necessarily measured on an absolute scale—only the median and mode serve as appropriate measures of central tendency (Stevens, 1946). Consequently, the median is an especially valuable statistical measure for biological data, which are often ordinal, asymmetrically distributed, and contain outliers.

The median proves to be a powerful tool in biological data analysis whenever a robust measure of central tendency is required to represent a "typical" value, such as cell size, metabolite concentration, or dispersal distance (McDonald, 2014). It is also commonly used to determine typical survival times in populations (Lethaby *et al*., 2013). In pharmacology, median effective dose (ED50) and median toxic dose (TD50)—which indicate doses causing specific effects or toxicity in 50% of a population—are essential for assessing drug safety and efficacy (Matsumura *et al*., 2013).

For datasets with an odd number of samples, the median—defined as the value minimizing the sum of absolute deviations—is simply the middle value. However, with an even number of samples, this approach leads to an ill-posed problem: the solution is not unique, as any point between the two middle values minimizes the sum of absolute deviations. Notably, there is no historical record of when or who first defined the median for even-sized samples as the average of the two middle values. Moreover, this conventional definition is poorly justified and

only appropriate under the strong assumption of a symmetric data distribution—an assumption

rarely valid for most biological datasets.

In this study, we demonstrate that the problem of defining the median for even sample

sizes can be well-posed; specifically, the location of the median can be uniquely determined

without bias toward symmetrical distributions. To assess the performance of this unbiased

median, we compared it with the conventional median by measuring the central tendency of

Poisson-distributed datasets. The Poisson process was chosen for its fundamental role in

modeling random independent events in biology, such as genetic mutations (Balin and Cascalho,

2010), cell divisions (Summers *et al*., 2015), and cancer development (Belikov *et al*., 2021). Our

results show that the unbiased median provides a more accurate estimate of central tendency and

exhibits lower variance than the conventional median.

58   **Methods and Results**

Jacques Hadamard (1902) defined a problem as *well-posed* if its solution exists, is

unique, and depends continuously on the input data. These three criteria have since become

fundamental concepts in mathematics. In contrast, finding the median for an even number of

samples by minimizing the sum of absolute deviations is an *ill-posed* problem because the

solution is not unique and does not depend continuously on the data. However, by framing this

ill-posed problem as a limiting case of a well-posed problem—specifically, by minimizing the

sum of residuals raised to the power of a rational number that approaches one—we can obtain a

solution that is both unique and depends continuously on the data.

Formally, consider a ranked dataset $X = \{x_1 < x_2 < ... x_M\}$ consisting of M measurements.

We seek an estimate α that minimizes the objective function:

$$69 \qquad SR = \sum_{i=1}^{M} (x_i - \alpha)^{\frac{2N}{2N-1}} \qquad (1)$$

70   where N is a natural number.

71   The even numerator $2N$ in the exponent ensures that each term in the sum (1) is non-negative. To

72   find the minimum of the sum of residuals SR defined by equation (1), we take its first derivative

73   with respect to α:

$$74 \qquad \frac{\partial SR}{\partial \alpha} = -\frac{2N}{2N-1} \sum_{i=1}^{M} (x_i - \alpha)^{\frac{1}{2N-1}} \qquad (2)$$

75   and set it equal to zero:

$$76 \qquad \sum_{i=1}^{M} (x_i - \alpha)^{\frac{1}{2N-1}} = 0 \qquad (3)$$

77   In a special case where $N = 1$, the equation (3) holds true when the estimate α equals the

78   arithmetic mean of the dataset:

$$79 \qquad \alpha|_{N=1} = \frac{1}{M} \sum_{i=1}^{M} x_i \qquad (4)$$

80   The mean provides the best estimate when the distribution of the dataset X is

81   symmetrical. For asymmetrical distributions, we transform each term in the derivative (equation

82   3) as follows:

$$83 \qquad (x_i - \alpha)^{\frac{1}{2N-1}} = \mathrm{sgn}(x_i - \alpha) \cdot \exp \frac{\ln|x_i - \alpha|}{2N-1} \qquad (5)$$

84   where the sign function "sgn($*$)" is:

$$85 \qquad \mathrm{sgn}(x_i - \alpha) = \begin{cases} -1 & \text{at } x_i < \alpha \\ 0 & \text{at } x_i = \alpha \\ 1 & \text{at } x_i > \alpha \end{cases} \qquad (6)$$

86   For large values of N ($N \gg 1$), the exponent in equation (5) can be expanded into a Taylor series:

$$\exp \frac{\ln|x_i - \alpha|}{2N - 1} = 1 + \frac{\ln|x_i - \alpha|}{2N - 1} + O(N^{-2}) \tag{7}$$

88   Therefore, each term in the sum from equation (3) can be approximated as:

$$(x_i - \alpha)^{\frac{1}{2N-1}} = \text{sgn}(x_i - \alpha) \cdot \left(1 + \frac{\ln|x_i - \alpha|}{2N - 1}\right) + O(N^{-2}) \tag{8}$$

90   Neglecting terms of order $O(N^{-2})$ for $N \gg 1$, and substituting these approximations back

91   into equation (3), the condition reduces to:

$$\sum_{i=1}^{M} \text{sgn}(x_i - \alpha) \cdot \left(1 + \frac{\ln|x_i - \alpha|}{2N - 1}\right) = 0 \tag{9}$$

94   For an even number of samples $M = 2K$, the equation (9) holds true at $x_K < \alpha < x_{K+1}$:

$$\sum_{i=1}^{K} \left(1 + \frac{\ln(\alpha - x_i)}{2N - 1}\right) = \sum_{i=K+1}^{2K} \left(1 + \frac{\ln(x_i - \alpha)}{2N - 1}\right) \tag{10}$$

96   As the numbers of ones in the sums of left and right sides of the equation (10) are equal, they

97   cancel each other, and the equation simplifies to:

$$\prod_{i=1}^{K} (\alpha - x_i) = \prod_{i=K+1}^{2K} (x_i - \alpha) \tag{11}$$

99   from which the estimate α at $K = 1, 2$ can be found as follows:

$$\alpha = \begin{cases} \dfrac{x_1 + x_2}{2} & \text{at } K = 1 \\ \dfrac{x_1 x_2 - x_3 x_4}{x_1 + x_2 - x_3 - x_4} & \text{at } K = 2 \end{cases} \tag{12}$$

101   At $K = 3$ to 4, the estimate α is a root of the third-degree polynomial, whereas at $K \geq 5$ α can be

102   calculated iteratively.

103   For an odd number of samples $M = 2K+1$, the equation (9) can be rearranged into:

104
$$\sum_{i=1}^{K}\left(1 + \frac{\ln(\alpha - x_i)}{2N - 1}\right) + (x_{K+1} - \alpha)^{\frac{1}{2N-1}} = \sum_{i=K+2}^{2K+1}\left(1 + \frac{\ln(x_i - \alpha)}{2N - 1}\right) \quad (13)$$

105   from which the residual $x_{K+1} - \alpha$ is found as:

106
$$x_{K+1} - \alpha = \left[\frac{1}{2N - 1}\ln\left(\prod_{i=1}^{K}\frac{(x_{K+i+1} - \alpha)}{(\alpha - x_i)}\right)\right]^{2N-1} \quad (14)$$

107   At $N \gg 1$, the right part of the equation (14) approaches zero, thus leaving:

108
$$\alpha = x_{K+1} \quad (15)$$

109   that matches the conventional median of the dataset with an odd number of samples.

110   Note that at $N \longrightarrow \infty$ the equation (1) can be simplified:

111
$$\lim_{N\to\infty}\sum_{i=1}^{M}(x_i - \alpha)^{\frac{2N}{2N-1}} = \sum_{i=1}^{M}|x_i - \alpha| \quad (16)$$

Thus, as $N \longrightarrow \infty$, the estimate α minimizes the sum of distances to the values in dataset

X. Since this estimate α from equation (11) is not biased toward the arithmetic mean of the

middle values and minimizes the sum of distances (equation 16), we refer to it as the "unbiased

median".

To evaluate the performance of the proposed unbiased median on datasets with

asymmetric distributions, we performed computer simulations of Poisson processes with

probability mass function $f(k;\lambda)$:

119
$$f(k;\lambda) = \frac{\lambda^k e^{-\lambda}}{k!} \quad (17)$$

where k denotes the number of events and λ is the expected value. Random Poisson-distributed

samples were generated using IgorPro software (WaveMetrics Inc., Oregon), which employs the

Mersenne Twister pseudo-random number generator. The unbiased median α was calculated in

IgorPro by numerically solving equation (11) with an accuracy of 10^{-6}.

To illustrate the differences among the unbiased median, conventional median, and mean,

we plotted their corresponding central tendency estimates for datasets with varying sample sizes

(Figure 1). Despite the considerable noise inherent to the Poisson distribution, it is evident that

the unbiased median (◆) often either coincides with the conventional median (◆)—notably at

sample sizes n = 10, 12, 14 in panel **A** and n = 8, 10 in panel **B**—or tends to shift closer to the

mean (◆). Importantly, the unbiased median frequently aligns more closely with the mean.

Figure 1. *Unbiased Median vs. Conventional Median and Mean*. Central tendency

estimates for Poisson-distributed datasets—evaluated as mean (◆), unbiased median (◆), and

conventional median (◆)—are plotted against sample size for expected values λ of 2, 4, and 8

(dashed lines) in panels **A**, **B**, and **C**, respectively. Notably, the unbiased median either coincides

with the conventional median (e.g., at sample sizes *n* = 10, 12, 14 in panel **A** and *n* = 8 and 10 in

panel **B**) or is shifted toward the mean, highlighting its tendency to align more closely with the

mean value as sample characteristics vary.

The tendency of the unbiased median to lean toward the mean suggests that its values

may be closer to the mean. To evaluate this hypothesis, we conducted a large number of

simulations ($n = 3 \cdot 10^7$) and computed the arithmetic means of the central tendency estimates to

reduce noise. Here and in subsequent analyses, we consider the arithmetic means of three

estimators: the arithmetic mean ($\blacklozenge$), the conventional median ($\color{red}\blacklozenge$), and the unbiased median ($\color{green}\blacklozenge$)

(Fig. 2). As expected for the Poisson process, the arithmetic mean ($\blacklozenge$) closely matches the

expected value λ (Fig. 2), with standard deviations (SD) from λ of $1.9 \cdot 10^{-5}$, $2.7 \cdot 10^{-5}$, $4.1 \cdot 10^{-5}$,

$3.6 \cdot 10^{-5}$, and $5.0 \cdot 10^{-5}$ for $\lambda = 1, 2, 3, 4$, and 5, respectively. Since all these SDs are greater than

10^{-6}, which is the numerical precision for calculating the unbiased median, they set the lower

bound for inaccuracy in subsequent analyses. Importantly, compared to the conventional median

($\color{red}\blacklozenge$), the unbiased median ($\color{green}\blacklozenge$) consistently provides estimates that are nearer to the expected

value λ.

Figure 2. *Underestimation of the Expected Value by Medians*. The arithmetic mean

(calculated from $3 \cdot 10^7$ simulations) of central tendency estimates for Poisson-distributed datasets

with expected values λ ranging from 1 to 5 (bottom to top) is plotted against the number of

samples. Central tendencies are evaluated as the mean (◆), unbiased median (◆), and

conventional median (◆). Note that both median estimators tend to underestimate the expected

value λ, but the unbiased median consistently provides estimates closer to the mean than the

conventional median.

Notably, the central tendency measured by the medians is significantly underestimated

for small sample sizes (Fig. 2) because medians are insensitive to the extreme positive values

that are intrinsic to the Poisson process. These extreme values are not outliers; rather, they are a

natural part of the distribution. As a result, the medians' disregard of these values leads to an

underestimation of the true central tendency. Therefore, using the mean—which correctly

estimates the expected value of a Poisson-distributed dataset—is more appropriate here than

using the median, which tends to underestimate it.

However, if these extreme values were true outliers, the median would provide a more

accurate measure of central tendency than the mean. Thus, the underestimation of the mean by

the medians (see Fig. 2) reflects their ability to mitigate the influence of extreme values. This

effect is most pronounced when the median reaches a local minimum in central tendency. To

quantify this effect, we plotted the largest differences between the medians and the mean

(referred to as undershoots) against the expected value λ (Fig. 3A). The undershoots for the

unbiased median (**+**) were fitted with an exponential function: $a \cdot \exp(-\lambda/\tau) - b$ (shown in green),

where $a = 0.0667 \pm 0.0019$, $\tau = 1.41 \pm 0.11$, $b = 0.1507 \pm 0.0009$ (here and further, the data are given

as a mean ± SD). Similarly, the undershoots for the conventional median (**×**) were fitted with the

same exponential form (shown in red), with parameters $a = 0.0313 \pm 0.0012$, $\tau = 1.50 \pm 0.16$, $b =$

0.1608 ± 0.0006. The average largest undershoot was smaller for the unbiased median compared

to the conventional median significantly (-0.1381 ± 0.0126 vs. -0.1544 ± 0.0061, n = 5, $P = 0.005$,

Student's paired t-test). Furthermore, the difference between the baselines ($-b$) of the exponential

fits was even more significant (-0.1507 ± 0.0009 vs. -0.1608 ± 0.0006, n = 5, $P = 1.3 \cdot 10^{-8}$,

Student's paired t-test). The key takeaway is that although both conventional and unbiased

medians reduce the influence of extreme values, the unbiased median yields a more accurate

estimate of the central tendency.

The sample sizes at which the largest undershoots occur in Poisson-distributed datasets

are presented in Fig. 3B. These sample sizes were fitted with linear functions of the form $a \cdot \lambda + b$,

yielding parameters $a = 6\pm0$, $b = 4\pm0$ for the unbiased median (shown in green), and $a = 5.4\pm0.2$,

$b = 3.00\pm0.66$ for the conventional median (shown in red). A key finding is that the largest

undershoot for the unbiased median consistently occurs at a smaller sample size compared to the

conventional median (Student's paired t-test, $P = 0.002$, n = 5).

Figure 3. *Reduced underestimation of the expected value by the unbiased median*. (*A*)

The largest difference (undershoot) between the estimated central tendency (using either the

unbiased or conventional median) and the expected value (λ) is plotted against λ. Undershoots

for the unbiased median (**+**) and conventional median (**×**) medians were fitted with the

exponential $a \cdot \exp(-\lambda/\tau) - b$. For the unbiased median (green), $a = 0.0667\pm0.0019$, $\tau =$

1.41 ± 0.11, $b = 0.1507\pm0.0009$. For the conventional median (red), $a = 0.0313\pm0.0012$, $\tau =$

1.50 ± 0.16, $b = 0.1608\pm0.0006$. On average, the largest undershoot was significantly smaller for

the unbiased median than for the conventional median (-0.1381 ± 0.0126 vs. -0.1544 ± 0.0061, n =

5, $P = 0.005$, Student's paired t-test). (**B**), The sample size at which the maximum undershoot

occurs for the unbiased (**+**) and conventional (**×**) medians is plotted against λ. These values were

fitted with a line $a \cdot \lambda + b$. For the unbiased median (green): $a = 6 \pm 0$, $b = 4 \pm 0$. For the

conventional median (red), $a = 5.4 \pm 0.2$, $b = 3.00 \pm 0.66$. Notably, the largest undershoot for the

unbiased median consistently occurs at a smaller sample size than for the conventional median

($P = 0.002$, n = 5, Student's paired t-test).

242   To determine when the unbiased median offers the greatest advantage over the

conventional median, we plotted the difference between these two estimators shown in Fig. 2

against sample size (Fig. 4A). The results indicate that this difference reaches its maximum at

sample sizes of 4 or 6 (Fig. 4A). Thus, the benefit of using the unbiased median—demonstrated

previously in Fig. 3—is particularly pronounced for small datasets containing 4 or 6 samples.

This advantage becomes even more significant at lower expected values λ, as shown in Fig. 4B:

here, the largest combined difference between the medians at sample sizes 4 and 6 (♦) is plotted

versus λ and fitted with the exponential function $a \cdot \exp(-\lambda/\tau) + b$, where $a = 0.0553 \pm 0.0007$, $\tau =$

0.409 ± 0.024, and $b = 0.02678 \pm 0.00003$ (n = 2).

Figure 4. *Peak Difference between Unbiased and Conventional Medians.* (***A***) The

differences between central tendency estimates using the unbiased and conventional medians are

plotted for expected values λ ranging from 1 to 5. Note that these differences reach their

maximum at sample sizes of 4 and 6. (***B***) The largest combined difference at sample sizes 4 and

6 (♦) is plotted against the expected value λ and fitted with an exponential function (grey):

$a \cdot \exp(-\lambda/\tau) + b$, where $a = 0.0553 \pm 0.0007$, $\tau = 0.409 \pm 0.024$, and $b = 0.02678 \pm 0.00003$ (n = 2).

Note that lower expected values correspond to greater differences between the two medians.

      In data analysis, variance—which quantifies the spread or dispersion of values—is as

essential a measure as central tendency, as it reveals how much data deviate from the average

and enables assessment of reliability, consistency, and interpretability (Weisberg, 1992). For

Poisson-distributed datasets with sample size *n*, the theoretical variance is λ/*n*; thus, multiplying

variance by *n* provides a convenient normalization for comparing results across different sample

sizes. To compare the variances of the central tendency estimators—the unbiased median,

conventional median, and mean—we normalized each variance by multiplying it by *n* and

plotted these values against sample size (Fig. 5). Normalized variances for central tendency

measures in Poisson datasets, evaluated as the mean (black), conventional median (red), and

unbiased median (green), are shown for expected values λ ranging from 1 to 5 (Fig. 5). As

expected for the Poisson process, the arithmetic mean (♦) closely matches the expected value λ

(see Fig. 2), with standard deviations from the expected value of $2.3 \cdot 10^{-4}$, $5.3 \cdot 10^{-4}$, $8.2 \cdot 10^{-4}$,

$1.1 \cdot 10^{-3}$, and $1.3 \cdot 10^{-3}$ for λ = 1, 2, 3, 4, and 5, respectively. These standard deviation values

represent the lower bound of inaccuracy in subsequent analyses.

Note that at small sample sizes, the normalized variances of both medians overestimate

the expected value λ, with the unbiased median consistently exhibiting lower variance than the

conventional median. Notably, as sample size increases, the variances of both medians

eventually fall below that of the mean beyond certain thresholds (as indicated with circles),

highlighting the greater efficiency of median-based estimators for larger sample sizes. The

earlier crossover point for the unbiased median (on average by 5.7 ± 1.1 samples, P = 0.0003, n =

5, Student's paired t-test) demonstrates its greater efficiency compared to the conventional

median.

Remarkably, at small sample sizes, the normalized variances of both median estimators

exceed the expected value λ, with the unbiased median consistently displaying lower variance

than the conventional median. As sample size increases, the variances of both medians

eventually drop below that of the mean at specific thresholds (indicated by circles in Fig. 5),

illustrating the greater efficiency of median-based estimators for larger datasets. Importantly, the

unbiased median achieves this crossover point significantly earlier—by an average of 5.7 ± 1.1

samples (P = 0.0003, n = 5, Student's paired t-test)—confirming its superior efficiency

compared to the conventional median.

Figure 5. *Variance overestimation by medians at small sample sizes*. Normalized

variances of central tendency estimators for Poisson-distributed datasets—computed as the mean

(black), conventional median (red), and unbiased median (green)—are shown for expected

values λ from 1 to 5. At small sample sizes, the variances of both median estimators exceed λ,

with the unbiased median consistently exhibiting lower variance than the conventional median.

As the sample size increases, the variances of both medians eventually drop below that of the

mean, beyond sample-size thresholds indicated by blue circles—demonstrating the greater

efficiency of median-based estimators at larger sample sizes. Notably, for the unbiased median,

this crossover point occurs an average of 5.7 ± 1.1 samples earlier than for the conventional

median ($P = 0.0003$, n = 5, Student's paired t-test).

It may seem counterintuitive that the variances of the medians overestimated the variance

of the mean, since one might expect that the medians' tendency to disregard extreme values in

Poisson datasets would reduce, rather than increase, their variance. However, when recognizing

that, at small sample sizes, the medians systematically underestimate central tendency (Fig. 2),

the corresponding overestimation of the deviation from the central tendency (Fig. 5) is a logical

consequence rather than an unexpected result. The advantage of the medians in disregarding

extreme values becomes evident at large sample sizes: the central tendency estimates converge to

the expected value, and the variance of the medians falls below that of the means beyond the

encircled regions in Fig. 5.

Normalizing the variances in Figure 5 facilitated comparison across large sample sizes.

However, for practical data analysis, it is essential to also consider the absolute (non-normalized)

variances, as their magnitudes provide direct information about variability in the original

measurement scale. To quantify how much the variance estimates from the medians exceed that

of the mean, we plotted the difference between the variances of the medians and the variance of

the mean as a function of sample size (Fig. 6A). The relative variance estimate from the

conventional median (shown in red) peaked consistently at a sample size of 6 across all expected

values λ. In contrast, the unbiased median (shown in green) exhibited peak relative variance at

sample size 6 for $\lambda = 1$ and 5; at sample size 8 for $\lambda = 3$ and 4; and at sample size 10 for $\lambda = 2$.

The peaks in the relative variances represent the points of greatest inefficiency of the medians

compared to the mean. To illustrate this, we plotted the peak values (referred to as overshoots)

against the expected value λ (Fig. 6B). These overshoots were fitted with a quadratic polynomial,

$a \cdot \lambda^2 + b \cdot \lambda + c$, yielding parameters: for the conventional median (red), $a = 0.0024 \pm 0.0009$, $b =$

0.0297 ± 0.0055, $c = 0.0636 \pm 0.0073$, and for the unbiased median (green), $a = 0.0008 \pm 0.0002$, $b =$

0.0177±0.0013, c = 0.0561±0.0017. On average, the largest overshoot was significantly smaller

for the unbiased median compared to the conventional median (0.1186±0.0361 vs.

0.1788±0.0696, n = 5, P = 0.016, Student's paired t-test), demonstrating the superior

performance of the unbiased median.

Notably, the relative variances were highest for sample sizes between 8 and 34,

coinciding with the greatest underestimation of central tendency by the medians (see Figs. 2 and

3C). As the medians approached the mean at larger sample sizes (Fig. 2), the corresponding

relative variances also dropped below zero at the sample sizes indicated in Fig. 6C. The sample

sizes were fitted with a line $a \cdot \lambda - b$: for the conventional median (red), a = 63.63±0.26, b =

6.78±0.85; for the unbiased median (green), a = 62.95±0.27, b = 10.43±0.90.

Figure 6. *Peaks and zeros of the relative variances*. (*A*) The relative variance estimate

from the conventional median (shown in red) peaked at the sample size of 6 at all expected

values λ, whereas that of the unbiased median (shown in green) peaked at samples sizes of 6 (at

$\lambda = 1, 5$), 8 (at $\lambda = 3, 4$), and 10 (at $\lambda = 2$). (*B*) The largest overshoots for the conventional ($\times$) and

unbiased (+) medians are plotted against the expected value λ. The overshoots were fitted with a

quadratic polynomial $a \cdot \lambda^2 + b \cdot \lambda + c$, yielding parameters: for the conventional median (red),

$a = 0.0024 \pm 0.0009$, $b = 0.0297 \pm 0.0055$, $c = 0.0636 \pm 0.0073$, and for the unbiased median (green),

$a = 0.0008 \pm 0.0002$, $b = 0.0177 \pm 0.0013$, $c = 0.0561 \pm 0.0017$. On average, the largest overshoot

was significantly smaller for the unbiased median than for the conventional median

(0.1186 ± 0.0361 vs. 0.1788 ± 0.0696, n = 5, $P = 0.016$, Student's paired t-test). *C*, The number of

samples at zero relative variances of conventional ($\times$) or unbiased (+) medians. The sample sizes

were fitted with a line $a \cdot \lambda - b$, where $a = 63.63 \pm 0.26$, $b = 6.78 \pm 0.85$ for conventional median

(red), and $a = 62.95 \pm 0.27$, $b = 10.43 \pm 0.90$ for unbiased median (green).

One can observe that the relative variance estimate of the unbiased median was

consistently lower than that of the conventional median for the same expected value λ (Figs.

6A,B), indicating a clear advantage in using the unbiased median. To identify the conditions

under which the unbiased median provides the greatest benefit, we analyzed the difference

between the variances of the unbiased and conventional medians (Fig. 7A). This difference was

the largest at sample sizes of 4 and 6. To further explore how this difference varies with sample

size, we plotted the combined variance difference at sample sizes 4 and 6 against the expected

value λ (Fig. 7B). The largest difference for the number of samples 4 and 6 (combined, n = 2)

was fitted with a line $a \cdot \lambda - b$, where $a = 0.0219 \pm 0.0001$, $b = 0.0004 \pm 0.0003$.

400   **Figure 7**. *Difference in variances of unbiased and conventional median*. (***A***) Difference

between the normalized variances of central tendency of Poisson's datasets (shown in Fig. 5)

evaluated as unbiased and conventional medians are shown for expected values λ from 1 to 5.

Note that the difference is the largest at the number of samples 4 and 6. (***B***) The largest

difference (♦) shown in panel (***A***) for the number of samples 4 and 6 (combined, n = 2) for

expected values λ from 1 to 5 is fitted with a line $a \cdot \lambda - b$, where $a = 0.0219 \pm 0.0001$, $b =$

0.0004 ± 0.0003.

Discussion

In this study, we proposed a novel median estimator for even samples sizes that, unlike the conventional median, is unbiased for asymmetrical distributions. This unbiased estimator was developed by re-framing the original *ill-posed* problem as a special case of a *well-posed* one: "finding the median by minimizing the sum of deviations raised to a power that approaches unity". Compared to the conventional median, the proposed median is a more accurate, lower-variance estimator of central tendency for asymmetrically distributed datasets.

We refer to the proposed median estimator as the *unbiased* median to distinguish it from the conventional median. The term "unbiased" was chosen for two main reasons. First, the conventional median—defined as the arithmetic mean of the two middle values—implicitly assumes that the data distribution is symmetrical, thereby introducing bias when this assumption does not hold. In contrast, the unbiased median, as calculated from equation (11), makes no such assumption and is therefore free from bias toward symmetrical distributions. Second, the name is an analogy to the concept of bias in statistics. For instance, the sample variance formula uses a correction (dividing by $n - 1$) because using the sample mean would otherwise create a biased estimate of the true population variance. In a similar way, the conventional median can be seen as a biased estimator of the population mean if the data's distribution is asymmetrical.

In the Poisson distribution (equation 17), k represents the number of events per measurement interval and must therefore be an integer. However, the expected number of events λ—defined as the arithmetic mean—can take any positive rational value. We assessed the properties of medians for Poisson-distributed datasets at discrete expectation values ($\lambda = 1, 2, 3, 4,$ and 5). In practice, however, λ can range from values near zero to very large numbers,

depending on the underlying process. To enable readers to interpolate between these discrete

values and, where appropriate, extrapolate beyond them, we fitted the data with suitable

functions (see Figs. 3, 4, 6, and 8). For example, extrapolating the exponential fits from Fig. 3A

predicts that the maximal difference between the central tendencies measured by the medians

occurs at $\lambda \ll 1$ and a sample size of four (see Fig. 4A).

While the difference between the central tendencies measured by the unbiased and

conventional medians is statistically significant, it remains to be determined whether this

difference is substantial enough to be perceptually meaningful. In other words, does a

quantitative difference result in a qualitative change? To address this, we simulated the capture

of an image of a circle using a 2D photodetector array. Since the maximal difference between the

medians occurs at $\lambda \ll 1$ and a sample size of four (see Figs. 3A and 4A), we set the expected

values for both photon flux and photodetector dark current to $\lambda = 1/5$. The resulting images were

then filtered using the conventional median, the unbiased median, and the mean of 2×2

neighboring pixels. For enhanced visualization, the grayscale images were mapped onto a

rainbow color scale: red indicates the lowest gray values, yellow to green corresponds to

intermediate values, and blue represents the highest values. Figure 8A displays the dark current

of the photodetector array; Figure 8B shows the incident photon flux; and Figure 8C presents the

combined image (sum of dark current and photon flux). Panels *D*, *E*, *F* represent the image from

panel *C* being filtered using conventional median, unbiased median, and mean, respectively. The

images in panels *A*, *B*, and *C* are all scaled to the same maximum value (the highest pixel in

panel *C*); similarly, images in panels *D*, *E*, and *F* are scaled to the maximum value in panel *F*.

Figure 8. *Qualitative Advantage of the Unbiased Median*. For clarity, grayscale images were mapped to a rainbow color scale: red denotes the lowest gray values, yellow to green correspond to intermediate values, and blue indicates the highest values. (*A*) Dark current image of the photodetector array (expected value $\lambda = 1/5$). (*B*) Incident photon flux shaped as a circle ($\lambda = 1/5$). (*C*) Combined output of the photodetector array (sum of dark current and photon flux). Panels *A*, *B*, and *C* are all scaled to the same maximum value (the highest pixel value in panel *C*). Panels *D*, *E*, *F* show the results of filtering the image from panel *C* using a 2×2 neighborhood with the conventional median (*D*), unbiased median (*E*), and mean (*F*), respectively. Images in *D*, *E*, and *F* are scaled to the maximum value in panel *F*. Note that brightness and contrast increase from the conventional median (panel *D*) to the unbiased median (panel *E*), and to the mean (panel *F*).

Remarkably, all filtered images in panels D, E, and F reveal the circle with much greater

clarity than the unfiltered image in panel C. Brightness and contrast progressively increase from

the conventional median (panel D), to the unbiased median (panel E), and finally to the mean

(panel F). As expected, the mean produces the brightest image, since both median estimators

tend to underestimate the mean (Figs. 2 and 3). However, the unbiased median produces a

distinctly brighter image than the conventional median, underscoring its reduced underestimation

of the mean. These results demonstrate that the unbiased median provides a more accurate

estimate of central tendency than the conventional median. Thus, the quantitative improvement is

sufficient to result in a qualitative difference that is perceptible to the naked eye.

Strictly speaking, using medians to analyze Poisson-distributed data is not ideal, since

such data inherently lack outliers (Haight 1967). By disregarding extreme positive values—

which are a natural part of the Poisson distribution—medians tend to underestimate the expected

value (see Figs. 2 & 3) and, as a consequence, overestimate the variance (see Figs. 5 & 6; also:

Huang 1991; Hozo *et al*., 2005). In real-world datasets, it is often difficult to determine whether

outliers are present. For instance, in scenarios where outliers are unlikely—such as measuring

photon counts from a photodetector with zero dark current—the median systematically

underestimates the central tendency, as illustrated in Figs. 2 and 3. Conversely, when genuine

outliers are present and serve as the primary source of asymmetry in the data, using the median

to estimate central tendency becomes particularly advantageous—especially under the similar

conditions in which it exhibits the greatest underestimation for Poisson-distributed data.

Overall, the results demonstrate the clear advantage of using the unbiased median over

the conventional one: (i) the unbiased median (equation 11) provides a unique solution to a well-

posed problem, free from any bias toward symmetrical distributions; (ii) central tendency of

asymmetrically distributed data measured as unbiased median is more accurate (Figs. 2, 3A, 4A);

(iii) the variance of central tendency measured as unbiased median is lower (Figs. 5, 6A, 7A);

(iv) the superior results provided by the unbiased median can be evident upon visual inspection

(Fig. 8). The only limitation of the unbiased median is that, for sample sizes of ten or more, its

calculation requires iterative methods based on equation (11), making it considerably more

computationally demanding than the conventional median, which is obtained simply by

averaging the two middle values.

Acknowledgements

This study was funded by grant R01AG066905 to S.A. Thomas from the National Institutes of

Health.

References

Balin SJ, Cascalho M. The rate of mutation of a single gene. *Nucleic Acids Res*. 2010

38(5):1575-82. doi: 10.1093/nar/gkp1119. PMID: 20007603

Belikov AV, Vyatkin A, Leonov SV. The Erlang distribution approximates the age distribution

of incidence of childhood and young adulthood cancers. 2021 *PeerJ* 9:e11976. doi:

10.7717/peerj.11976. eCollection. PMID: 34434669

Hadamard J. Sur les Problèmes aux Dérivées Partielles et Leur Signification Physique. 1902

Princeton University Bulletin, 13, 49-52.

Haight FA. Handbook of the Poisson Distribution. 1967 New York, NY, US: John Wiley &

Sons. ISBN: 9780471339328, 0471339326. 168 pages.

Hozo SP, Djulbegovic B, Hozo I. Estimating the mean and variance from the median, range, and

the size of a sample. 2005 *BMC Med Res Methodol*. **5**:13. doi: 10.1186/1471-2288-5-13. PMID:

15840177

Huang JS. Estimating the variance of the sample median, discrete case. 1991 *Statistics &*

Probability Letters, **11**:4, p. 291-298. doi: 10.1016/0167-7152(91)90038-S

Lethaby CD, Picton S, Kinsey SE, Phillips R, van Laar M, Feltbower RG. A systematic review

of time to diagnosis in children and young adults with cancer. 2013 *Arch Dis Child.* 98(5):349-

55. doi: 10.1136/archdischild-2012-303034. PMID: 23476000

Matsumura N, Kikuchi-Utsumi K, Sakamaki K, Watabe M, Aoyama K, Nakaki T.

Anticonvulsant action of indazole. 2013 *Epilepsy Res*. **104**(3):203-16. doi:

10.1016/j.eplepsyres.2012.11.001. PMID: 23219048

McDonald JH. Handbook of Biological Statistics, 3rd ed. 2014 Sparky House Publishing,

Baltimore, Maryland. http://www.biostathandbook.com/HandbookBioStatThird.pdf

Stevens SS. On the Theory of Scales of Measurement. 1946 *Science*, **103**(2684): 677–680.

PMID: 17750512 DOI: 10.1126/science.103.2684.677.

Stigler SM. The History of Statistics: The Measurement of Uncertainty before 1900. 1990 ISBN-

13 978-0674403413; Reprint by Belknap Press. 428 pages.

Summers HD, Wills JW, Brown MR, Rees P. Poisson-event-based analysis of cell proliferation.

2015 *Cytometry A*, **87**(5):385–392. doi: 10.1002/cyto.a.22620. PMID: 25572722

Weisberg HF. Central Tendency and Variability. 1992 SAGE Publications, Inc. (Quantitative

Applications in the Social Sciences, No. 83). ISBN 13: 9780803940079

Figure_1

Figure_2

Figure_3

Figure_4

Figure_5

Figure_6

Figure_7

Figure_8