Cross-fostering affects microglia and cell death in the hippocampus of female and

male degu pups

Gurprince Attlas[1], Mallory Duprey[2], Cassandra Carlson[1], Kaja S. Arusha[3], Krystle D. Boadi[3], Sabrina S.

Ellah[3], Daniela Kim[3], Charlis Raineki[1,2], Carolyn M. Bauer[3], Paula Duarte-Guterman[1,2]

1. Department of Psychology, Brock University, St. Catharines, ON, Canada

2. Centre for Neuroscience, Brock University, St. Catharines, ON, Canada

3. Department of Biology, Swarthmore College, PA USA

Corresponding authors:

Dr. Paula Duarte Guterman, pduarteg@brocku.ca

Dr. Carolyn Bauer, cbauer1@swarthmore.edu

Keywords: neurogenesis; dentate gyrus; apoptosis; early life stress; sex differences

Abstract

Parental care is essential for social, behavioural, and neural development of offspring. Disruption of parental care, for example through parental separation, can negatively affect offspring development in rodents. While previous work has focused on maternal and paternal deprivation, the effects of cross-fostering, another form of parental-offspring instability, on brain development remain poorly understood. Such disruptions are thought to induce stress, which in turn can suppress neurogenesis and increases inflammation and apoptosis in the hippocampus, with effects that often differ between sexes. Given that degus (*Octodon degus*) are born precocial and form strong attachments with their parents early in life, this study aimed to investigate the effect of cross-fostering on hippocampal development in female and male degu pups. At postnatal day 8, degus were assigned to either control (pups remained with parents and littermates), partial cross-foster (one pup per litter was cross-fostered), or full cross-foster (the entire litter was cross-fostered) conditions. At weaning (5-weeks-old), offspring brains were collected for immunohistochemistry to examine dentate gyrus volume, density of pyknotic cells, density of immature neurons, and number and morphology of microglia. Both types of cross-fostering reduced hippocampal cell death in both sexes relative to controls, while no significant effects were observed in the dentate gyrus volume or density of immature neurons. Cross-fostering did not affect total microglia density in either sex however full cross-fostered females had fewer amoeboid microglia compared to female controls. Together, these findings indicate that cross-fostering affects hippocampal microglia and cell death in pups, potentially disrupting circuit refinement and plasticity during development, with effects that vary by cross-fostering type, sex, and hippocampal region.

Introduction

Early life stress affects offspring development in a range of mammalian species, including humans (Gimsa et al., 2022; Kentner et al., 2010; Priebe et al., 2005; Raineki et al., 2019). In rodents, there are different paradigms which induce early life stress that include, but are not limited to, changing the parental environment via maternal separation, paternal deprivation, or cross-fostering (Aguggia et al., 2013; Boccia and Pedersen, 2001; George et al., 2010; Madison et al., 2022; Oomen et al., 2011). Any disruption to typical parent-offspring interactions can lead to alterations in offspring neurodevelopment (Champagne and Meaney, 2001). During postnatal development, there is high growth and refinement of brain circuits. At a mechanistic level, there are high rates of neurogenesis (i.e. production of new neurons) and regulated cell death (i.e., apoptosis; Hayashi et al., 2008; Loi et al., 2017; Sierra et al., 2014). Moreover, microglia, the neuroimmune cells of the brain, play a very important role in refining circuits during early development, either by removing cells through phagocytosis or by supporting neuronal development and survival (Diaz-Aparicio et al., 2020; Sierra et al., 2014).

One of the brain regions that is most sensitive to early life stressors is the hippocampus. It is also one of the few regions where neurogenesis continues during the postnatal period (Kempermann, 2016). Past studies have shown that parental separation affects hippocampal neurogenesis, cell death (apoptosis), and microglia in the offspring, but the effects may depend on age, sex of the offspring, and type of separation (maternal/paternal, acute/chronic). For instance, acute maternal separation (24 h on postnatal day 3) reduces the numbers of immature neurons in the hippocampus of female rats at weaning, while increasing immature neurons in males (Oomen et al., 2009). In contrast, chronic maternal separation has been shown to reduce neurogenesis in male rats (Hulshof et al., 2011). Early life stressors can also affect programmed

cell death. For example, maternal separation increases apoptosis in the dentate gyrus of rat pups

(Yang et al., 2017). Thus, research suggests a potential link between dysregulated neurogenesis

and apoptosis in the dentate gyrus due to early life stress, potentially mediated by microglial

activity. Indeed, brief daily maternal separation increases both the number of and phagocytic

activity of microglia in the hippocampus of juvenile male mice (Delpech et al., 2016; Johnson

and Kaffman, 2018; Walker et al., 2013). In addition to removing cells through phagocytosis,

microglia can also promote the production and survival of new neurons (neurogenesis) through

the secretion of trophic factors (Diaz-Aparicio et al., 2020). Here we aimed to understand how

these three key components—neurogenesis, cell death, and microglia—are regulated following a

stressor during development.

Most studies to date have focused on the effects of maternal and paternal deprivation on

offspring development, whereas the impact of other disruptions of parental-offspring interactions

has received much less attention. Cross-fostering disrupts parent-offspring attachment, but unlike

maternal separation or paternal deprivation, it does not rely on separation and removing pup's

access to a caring adult. Past research has shown that cross-fostering affects offspring body

weight, emotional behaviour, cardiovascular and metabolic functions, and the endocrine stress

response in different rodents species, including mice, degus, and rats (Arusha et al., 2024;

Bartolomucci et al., 2004; Gomez-Serrano et al., 2001; Guan et al., 2023; Lu et al., 2009;

Matthews et al., 2011). Degus (*Octodon degus)* are a particularly suitable model for studying the

effects of cross-fostering on offspring development because they practice communal care and

pups are born precocial and form strong attachments early in life with both parents and

littermates (Colonnello et al., 2011; Guan et al., 2023). Importantly, the presence of siblings

during cross-fostering may buffer the effects of separation from biological parents (Arusha et al.,

2024). Accordingly, in the current study, we investigated the effects of cross-fostering either the

entire litter (full cross-fostering) or a single pup (partial cross-fostering) on hippocampal

development in female and male degus. We measured hippocampal development by assessing

dentate gyrus volume, postnatal neurogenesis, apoptosis, and microglia prevalence and

morphology. We hypothesized that (1) cross-fostering would impact hippocampal development

by reducing neurogenesis, increasing cell death and microglial numbers, and altering the

microglia morphological profile in both sexes; and (2) partial cross-fostering would be more

detrimental to hippocampal development compared to cross-fostering the entire litter.

Methods

Study animals and breeding

Breeding pairs of captive-born *Octodon degus* and their pups were housed in the animal

facility at Swarthmore College, in 51 x 25 x 33 cm acrylic terrariums containing corn cob or

paper bedding allowing for visual observations. Animals were maintained under standard

laboratory conditions (12:12 light/dark cycle, 23 ± 1 ˚C) with *ad libitum* access to water and

food. Alfalfa hay was provided three times per week, and cages were cleaned weekly.

Degu females (n = 21) were paired with unrelated males (n = 21) for mating and remained

together until pup weaning. All experiments were conducted in strict compliance with

Institutional Animal Care and Use Committee at Swarthmore College and were carried out

according to the guidelines of the Association for Assessment and Accreditation of Laboratory

Animal Care.

Experimental treatment

Pups were cross-fostered at postnatal day (PND) 8, thus allowing for sufficient time for

the development of parental and social attachments within the nest (Colonnello et al., 2011;

Guan et al., 2023). The date of parturition was taken into consideration while assigning the

treatment groups as previously described (Arusha et al., 2024). Litters were assigned to one

treatment condition: control (N=5 litters), full cross-fostering (N=4 litters), and partial cross-

fostering (N=12 litters). In control litters, pups remained with biological parents and biological

siblings for the duration of the experiment. In full cross-foster litters, all pups from one litter

were exchanged with all pups from another such that pups were raised with unfamiliar parents

but with biological siblings. In partial cross-foster litters, a single pup was exchanged with one

pup of the same sex from another litter such that the pup was raised with both unfamiliar parents

and unfamiliar siblings. Pups were fostered between litters of similar sizes and ages (<2 days in

age difference). Between 1-2 pups per sex per group were used for subsequent analyses, with the

exception of the male full cross-foster condition that included 3 male pups from the same litter

(out of 4 litters). Sample sizes were decided based on previous work (Arusha et al., 2024; Guan

et al., 2023) and are presented in Table 1 and each of the figure captions.

Tissue collection

Euthanasia was performed at weaning (~5 weeks of age). Pups received 0.2 mL of

undiluted Euthasol (pentobarbital sodium and phenytoin sodium solution), and brains were

immediately dissected and split into the right and left hemispheres. The right hemispheres were

placed on dry ice and frozen at –80˚C for future analysis, whereas the left hemispheres (used in

the present study) were immersion-fixed in 4% paraformaldehyde for 24h at 4˚C. Following

fixation, brains were then transferred to a 30% sucrose solution containing 0.1% sodium azide

and stored at 4˚C until sectioning. Coronal sections of 30 µm thickness were cut using a cryostat

at –20˚C. Hippocampal sections were collected in a seven-tube series containing an antifreeze

solution (ethylene glycol, glycerol and 0.1M phosphate buffer saline with a pH of 7.4) and stored

at –20˚C. All three neural measurements (volume, neurogenesis, and microglia) were performed

in parallel series from the same left hemisphere for all animals.

Cresyl violet staining

Brain sections were stained with Cresyl violet to determine the effects of cross-fostering

on dentate gyrus volume and to count pyknotic cells. Hippocampal tissue from one of the series

was mounted onto Superfrost microscope slides. Cresyl violet staining was performed after the

tissue had completely dried on the slides. The staining process started by dipping the slides into

distilled water three times to remove residual salts, then submerged in 0.2% Cresyl violet acetate

solution for 4 minutes. Following staining, the slides were placed in a differentiator solution,

comprising 70% ethanol and seven drops of glacial acetic acid, for 30 seconds. The slides were

then sequentially dehydrated in 95% ethanol (twice, for 1 minute each), followed by 100%

ethanol for 1 minute. Finally, slides were cleared in xylene (twice, for 5 minutes each) and

cover-slipped using Permount.

Doublecortin immunohistochemistry

Brain sections were stained with doublecortin (DCX) to assess the effects of cross-

fostering on postnatal neurogenesis. All staining procedures were conducted in multi-well plates

under gentle shaking. To begin, sections were washed in 0.1M phosphate buffer saline (PBS, pH

= 7.4) to remove antifreeze and then incubated in 0.6% hydrogen peroxide in PBS for 30

minutes. Sections were then rinsed with PBS and then incubated for 22h at 4˚C in rabbit anti-

DCX primary antibody (1:200 dilution; Cell Signalling, catalogue number: 10124618) diluted in

0.1M PBS containing 3% of normal goat serum (NGS) and 0.4% Triton-X. After incubation,

tissue was rinsed in 0.1M PBS, then incubated for 18h at 4°C in biotinylated goat anti-rabbit

secondary antibody (1:500 dilution; Vector laboratories) diluted in 0.1M PBS. This was followed

by washes in 0.1M PBS and incubation for 4 hours in an avidin-biotin horseradish peroxidase

complex solution (1:1000 dilution; ABC kit, Vector Laboratories). After rinsing in 0.1M PBS,

tissue was incubated in diaminobenzidine in the presence of nickel (DAB Peroxidase Substrate

Kit, Vector Laboratories) for 3 minutes. Stained sections were mounted on Superfrost

microscope slides and allowed to dry for 2–3 days. Slides were then dipped in distilled water

three times to remove salts, dehydrated sequentially in 70% ethanol (1 minute), 95% ethanol (1

minute), and 100% ethanol (1 minute), then cleared in xylene (twice, 5 minutes each). Finally,

coverslips were applied using Permount.

Iba-1 immunohistochemistry

176   Brain sections were stained for ionized calcium-binding adaptor molecule (Iba-1) to

access the effects of cross-fostering on microglia. All staining procedures were conducted in

multi-well plates under gentle shaking. Sections were first washed in 0.1M PBS (pH = 7.4) to

remove antifreeze, followed by an incubation in 0.6% hydrogen peroxide in PBS for 30 minutes.

Tissue was then rinsed with PBS and incubated for 22h at 4°C in rabbit anti-Iba-1 primary

antibody (1:1000 dilution, Wako, catalogue number: 019-19741), diluted in 0.1M PBS

containing 3% NGS and 0.4% Triton-X. The following day, sections were rinsed in 0.1M PBS,

then incubated for ~18h in biotinylated goat anti-rabbit secondary antibody (1:500 dilution,

Vector laboratories) diluted in 0.1M PBS. This was followed by washes in 0.1M PBS and

incubation for 4h in an avidin-biotin horseradish peroxidase complex solution (1:1000 dilution;

ABC kit, Vector Laboratories). Tissue was rinsed in 0.1M PBS followed by incubation in

diaminobenzidine in the presence of nickel (DAB Peroxidase Substrate Kit, Vector Laboratories)

for three minutes. Stained sections were mounted on Superfrost microscope slides and allowed to

dry for 2–3 days. Slides were dipped in distilled water three times to remove salts, dehydrated

sequentially in 70% ethanol (1 minute), 95% ethanol (1 minute), and 100% ethanol (1 minute),

then cleared in xylene (twice, 5 minutes each). Coverslips were applied using Permount.

Microscopy, Imaging, and Quantification

194   All analyses were performed by a researcher blinded to experimental conditions.

Brightfield images of the entire left hippocampus stained with Cresyl violet were taken using a

standard light microscope (Olympus BX53) equipped with a camera (ORCA-spark, Hamamatsu)

using the 4X objective. The granule cell layer (GCL) and hilus were traced separately using

ImageJ to obtain areas in μm^2. Volumes of GCL and hilus were calculated using the Cavalieri's

principle: the measured areas were multiplied by the section thickness (30 µm), by 7 (reflecting

the 7 sections collect for the hippocampus), and finally by 2 (to present the volume for both

hemispheres). The dentate gyrus volume was calculated by adding the volume of the GCL and

the hilus. For the pyknotic cell count, two dorsal and two ventral sections of the dentate gyrus

were selected for each animal. Ventral sections were defined as those located between interaural-

incisor coordinates 2.32 and –0.32 of the degu brain atlas (Kumazawa-Manita et al., 2018). We

counted the number of pyknotic cells in the GCL using a 100X oil immersion objective. During

counting, the fine focus knob was used to scan through the entire thickness of each section to

ensure no cells were missed.

208   For the analysis of immature neurons, pups exhibited a very high expression of DCX,

typical for this developmental period, but making it challenging to reliably delineate and count

individual immature neurons (Figure 2A and B). Therefore, we used optical density as an index

of DCX expression. In addition, although morphological classification of DCX+ cells can

provide insight into neuronal maturation, the high expression of DCX also prevented us from

reliably scoring individual cell processes. Brightfield images were taken from two dorsal and two

ventral sections using an Olympus BX53 microscope with a 4X objective. Optical density (OD)

of DCX labeling was quantified using ImageJ. The GCL was traced in each image, and grey

values were obtained. The Rodbard equation was employed to calculate the OD for both the

GCL and background region (corpus callosum). For each section, background OD was

subtracted from the GCL OD. The average corrected OD for both image densities was used in

statistical analysis.

For microglia, brightfield images were taken using an Olympus BX53 microscope with a

20X objective. Two images from the dorsal and two from the ventral dentate gyrus per animal

were taken. We used a standardized region of interest (490.4 x 368.5 μm; see placement in

Figure 3A), which included the GCL and hilus regions. We counted the total number of

microglia present and categorized them into three morphological types—ameboid, intermediate,

and ramified—using ImageJ, as we have done in the past (Duarte-Guterman et al., 2023).

Ramified microglia were defined by a small cell body with at least three processes, each at least

twice the length of the cell body diameter. Intermediate microglia were defined by cell bodies

with 1–2 processes, each no longer than twice the diameter of the cell body. Ameboid microglia

were defined as a dark cell body with very small or no visible processes. All microglia were fit

into the best or closest category by an experimenter that was blinded to the treatment groups. The

number of microglia was added to get the total number of microglia per individual and

normalised to region of interest to obtain a density measure. For one partial cross-fostered male,

due to tissue loss during immunostaining, we did not have dorsal sections for Iba-1 analysis. One

control female brain did not have two ventral section images due to missing tissue sections,

therefore, data of only one section were utilized in the analysis.

Statistical Analysis

All statistics were performed in IBM SPSS, Version 30.0.0.0 and data visualization was

performed in R (R Core Team, 2021) using the ggplot2 package (Wickham, 2016). ANOVA

assumptions of normality and homogeneity of the variance were checked with visual inspections

of QQ plots and residual plots, and Kolmogorov-Smirnov and Levene's tests, respectively. The

volumes of the dentate gyrus, hilus, and GCL were analysed using analysis of covariance

(ANCOVAs) with treatment (control, partial, and full cross-foster) and sex (female, male) as

between-subject factors, and litter size and number of females as covariates. For DCX OD,

microglia density, microglia morphology, and pyknotic cell density, data were analysed using

repeated measures ANOVAs, with treatment and sex as between subject factors, region (dorsal

and ventral) as a within-subject factor, and litter size and number of females as covariates.

Including litter size and sex ratio as covariates ensured that treatment-related effects were not

confounded by natural variation in litter composition. Including these covariates in the analysis

did not affect the main patterns of the results. For all tests, we present the estimated marginal

means and the raw data. Post hoc comparisons for the repeated-measures ANOVA were conducted

using the Sidak correction. For the repeated measures ANOVAs, if the assumption of sphericity

was violated, the degrees of freedom were corrected using the Greenhouse-Geisser estimate of

sphericity. Pearson r correlations were performed between cell types, separately for females and

males and the different treatment groups. P-values < 0.05 were considered statistically significant.

Results

Cross-fostering did not alter dentate gyrus volume or DCX optical density

Cross-fostering did not affect the volume of the dentate gyrus in female and male pups, (all p's > 0.526) (Figure 1). Similarly, cross-fostering did not affect the volume of the hilus (all p's > 0.292) or the GCL (all p's > 0.595; Table 1). We did not find any sex differences for any of the volumes (dentate gyrus, hilus, or GCL) (all p's > 0.292; Table 1).

To examine the effects of cross-fostering on postnatal neurogenesis, we measured the OD of DCX in the dorsal and ventral regions of the hippocampus. We found a significant interaction between treatment, sex, and region (F (2,32) = 4.212, p = 0.024; η^2 = 0.208). Post hoc analysis indicated that partial cross-fostered females had a trend for significantly higher DCX OD than full cross-fostered females in the ventral hippocampus (p=0.059; Figure 2). However, this trend was largely driven by a single outlier in the partial cross-fostered group; after winsorising this value to the nearest non-extreme value, the contrast was no longer significant (p = 0.458). There were no significant differences between partial cross-fostered females and control females (p = 0.168), or between full cross-fostered females and control females (p = 0.157) in the ventral hippocampus. There were no differences between males across treatment conditions in DCX OD in the ventral hippocampus (all p's > 0.072). In the dorsal hippocampus, cross-fostering did not affect DCX OD in either females or males (all p's > 0.101).

Furthermore, we detected sex differences in DCX OD in the partial cross-fostered and full cross-fostered pups. In the ventral hippocampus, partial cross-fostered females had higher DCX OD than males (p = 0.032) although this sex difference disappears when the partial cross fostered female outlier is replaced with the nearest non-extreme value. In contrast, full cross-

fostered females had lower OD compared to males (p = 0.023). In controls, no sex differences in

DCX OD were found in either the dorsal or ventral hippocampus (all p's > 0.429).

Cross-fostering altered microglia counts and morphology in the dentate gyrus of degu pups

Cross-fostering did not affect the density of Iba-1-positive cells in the dentate gyrus of

degu pups when taking into account litter size and sex ratio in the analysis (all p's > 0.149,

including the effect of covariates; Figure 3). Without these covariates, there was a significant

treatment by region interaction: (F (2,33) = 3.824, p = 0.033), however post-hoc analyses did not

reveal any significant differences between treatments.

While total Iba-1 density was not affected by cross-fostering, we analysed the density of

the three morphological states of Iba-1 expressing cells. Cross-fostering also affected the density

of ameboid Iba-1-positive cells with the effect being sex-specific but across both dorsal and

ventral regions (treatment by sex interaction: F (2,31) = 4.531, p = 0.019; η^2 = 0.226, Figure 4).

Full cross-fostered female pups had fewer ameboid microglia compared to female controls (p =

0.026). Partial cross-fostering did not affect the density of ameboid Iba-1 positive cells in female

pups compared to control or full cross-fostering conditions (all p's > 0.449). No significant

differences were found in the density of ameboid microglia in males (all p's > 0.733). No

significant differences were observed in the density of ramified or intermediate microglia across

cross-fostering conditions (all p's > 0.089).

Cross-fostering differentially altered pyknotic cell density in dorsal and ventral hippocampus

regions

Cross-fostering affected pyknotic cell density differently in the dorsal and ventral

hippocampus (treatment by region interaction: $F_{(2,32)} = 5.058$, $p = 0.012$, $\eta^2 = 0.240$, Figure 5).

In the dorsal and ventral hippocampus, full cross-fostered pups had fewer pyknotic cells

compared to controls ($p < 0.001$ and $p = 0.025$, respectively). Partial cross-fostered pups also had

fewer pyknotic cells compared to controls but only in the dorsal hippocampus ($p = 0.002$) No sex

differences were found in pyknotic cell density in either dorsal or ventral regions (all p's >

0.127).

Correlations between cell types

We examined correlations between microglia density and morphology, DCX OD, and

pyknotic cell density within each cross-fostering condition, separately for males and females and

hippocampal region. In the dorsal region, we found density of ramified microglia was

significantly correlated with DCX OD in partial cross-fostered males only (Pearson $r = 0.914$, p

$= 0.03$). No other significant associations between these endpoints in either sex were found in the

dorsal region (all p's > 0.10). In the ventral region, intermediate microglia density was correlated

with DCX OD in partial cross-fostered females only (Pearson $r = 0.948$, $p = 0.014$). There were

no other significant correlations in the ventral region (p's > 0.08).

Discussion

The purpose of this study was to examine the effects of cross-fostering on neural

development in female and male degu pups. Degus are born precocial and form strong

attachments early in life, making them a great model for studying the effects of early-life

adversity and social bonding during development (Arusha et al., 2024; Guan et al., 2023). We

found that cross-fostering affected microglia morphology and pyknotic cell density, but not

dentate gyrus volume or neurogenesis. Full cross-fostering was associated with a reduction in

ameboid microglia in female pups in both dorsal and ventral regions. Full cross-fostered pups,

regardless of sex, had a lower density of pyknotic cells compared to controls in the dorsal and

ventral regions. However, partial cross-fostering reduced pyknotic cell density in both sexes but

only in the dorsal region. Overall, this study demonstrates that cross-fostering affects

hippocampal microglia and cell death in a type of cross-fostering-, sex-, and region-specific

manner in degu pups.

Cross-fostering did not affect dentate gyrus volume in pups

Cross-fostering did not impact the volume of the dentate gyrus in degu pups. Previous

work in other rodents such as rats, mice, and California mice has been mixed. Some studies find

reductions in the dentate gyrus volume of pups exposed to early life stressors such as paternal

deprivation (Madison et al., 2022), chronic unpredictable stress (Herpfer et al., 2012), and

scarcity-adversity model using low bedding (Raineki et al., 2019). However, other studies find

no effect after exposure to chronic unpredictable stress (Huot et al., 2002) or maternal separation

(Loi et al., 2017). Overall, our findings highlight the importance of considering both the type of

stressor and the species when evaluating the effects of early life stressors on hippocampus

development. Notably, in the current experiment, maternal and paternal licking and grooming

were not affected by cross-fostering in degus (partial or full cross-fostering), although cross-

fostered mothers spent more time off the nest compared to control and partial cross-foster groups

(Bauer et al., in prep). It is possible that having a relatively more stable parental care

environment (e.g., compared to separation paradigms) may help protect against the effects of

cross-fostering on dentate gyrus volume in pups.

Interestingly, we did not find sex differences in dentate gyrus volume in degu pups. This

is in contrast with findings in mice, where males have larger hippocampal volume compared to

females between PND 10 and PND 17, and again at PND 65 (Qiu et al., 2018). To our

knowledge, our study is the first one to examine hippocampal volume in degu pups. Further

research is required in both juvenile and adult degus to understand sex differences in

hippocampal development.

Cross-fostering does not affect neurogenesis in pups

We examined the effects of cross-fostering on postnatal neurogenesis using DCX OD as

a marker of immature neurons. We found that cross-fostering did not affect DCX OD compared

to controls. Previous work has found that maternal separation leads to a reduction in the number

of immature neurons at weaning in female, but not male rats (Oomen et al., 2009). In biparental

species, paternal deprivation either has no effect on neurogenesis (cell survival) in adult

California mice (Madison et al., 2022), or decreases cell survival in adult females only (Glasper

et al., 2018). In mandarin voles, paternal deprivation is associated with a reduction in immature

neurons in adulthood compared to biparentally raised pups (He et al., 2018). Overall, these

findings suggest that changes in neurogenesis following parental separation are not always

observed in all studies and results can vary across sexes and potentially with age. Our

observations were limited to a single developmental time point and one neurogenesis marker

using optical density. It remains possible that cross-fostering may affect the number of immature

neurons, without affecting the density of DCX, or that maturation and survival of new neurons

are affected, or that changes are manifested later in adolescence or adulthood.

Cross-fostering alters pyknotic cell density in pups

The current study revealed noteworthy alterations in pyknotic cell density in the dentate

gyrus depending on the type of cross-fostering and region. We found that full cross-fostered

pups, regardless of sex, exhibited fewer pyknotic cells compared to controls in both regions of

the hippocampus, while partial cross-fostered pups had reduced pyknotic cell density in the

dorsal region only. In contrast, previous studies in rats have shown that maternal separation can

increase cell death, although the effects depend on timing of exposure. For example, maternal

separation increases hippocampal cell death after an acute 24-hour separation at PND12, but not

at PND20 (Zhang et al., 2002). Increases in hippocampal cell death have also been reported

following chronic maternal separation in rats (Baek et al., 2012). These discrepancies between

previous work and the current study may reflect species differences or differences in the type and

severity of the early life stressor, as discussed earlier. Apoptosis is a crucial developmental

process that contributes to the formation and refinement of neural circuits, and alterations in this

process can influence neuron number and brain function (Forger, 2009; Majcher-Maślanka et al.,

2019). Thus, the observed reduction in pyknotic cell density in cross-fostered pups could indicate

a decreased level of apoptosis and insufficient cell pruning, potentially resulting in overcrowding

of neurons and altered neural circuitry.

Partial cross-fostering affected the dorsal and ventral regions differently in pups. The

dorsal hippocampus is primarily involved in spatial learning and other cognitive aspects of

memory, whereas the ventral hippocampus is associated with stress and anxiety regulation

(Bannerman et al., 2004; Fanselow and Dong, 2010; Kheirbek and Hen, 2011). Further research

is needed to determine whether cross-fostering-related changes in cell death during

development translate into changes in cognition, mood or other hippocampal-related behaviours

in degus.

Cross-fostering altered microglia numbers and morphology in pups

We found that cross-fostering altered the morphology of microglia in the hippocampus

but not overall microglia density. Microglia play a crucial role in the central nervous system,

contributing to synaptic remodeling through pruning, as well as supporting neuronal survival

via the release of neurotrophic factors (Cornell et al., 2022; Sierra et al., 2014). Although we did

not observe changes in overall microglia numbers, previous work has shown that an adverse

early life environment due to a combination of minimal bedding, frequent cage changes, and a

Western diet increases microglial numbers in the mouse hippocampus at weaning (Cohen et al.,

2016). Similarly, maternal separation in rats (Banqueri et al., 2019) and paternal deprivation in

California mice (Madison et al., 2022) have also been associated with increased microglia in the

offspring hippocampus.

Instead, in the current study, we found that cross-fostering affected the morphology of

microglia in a sex-dependent manner. Full cross-fostered female pups had fewer ameboid

microglia compared to control females regardless of hippocampal region. Ameboid microglia are

typically associated with phagocytosis, synaptic pruning, and the secretion of pro-inflammatory

cytokines and are characterized by a rounded morphology lacking the long processes often

observed in ramified microglia (Burke et al., 2016; Colonna and Butovsky, 2017; Franco and

Fernández-Suárez, 2015; Paolicelli et al., 2022). The current literature agrees that microglia vary

from ameboid to ramified states, but each morphology does not have exclusive functions. The

ability to transition between different stages allows microglia to maintain a balanced immune

response (Paolicelli et al., 2022). One possible interpretation of our results is that the reduction in

ameboid microglia in full cross-fostered female pups may reflect lower levels of phagocytosis,

potentially contributing to the reduced pyknotic cell density observed. However, this may not be

the only contributing factor because cross-fostering also decreased pyknotic cell density in males

with no changes in microglia morphology. In addition, there were no significant correlations

between microglia (numbers and morphology) and pyknotic cell density in our samples, although

the results may be limited by the small sample size. Overall, while cross-fostering clearly

affected microglial morphology in the female hippocampus, whether these changes directly lead

to apoptosis remain to be elucidated.

Brain measures depend on type of cross-fostering

Our results indicate that full cross-fostered pups showed greater effects on cell death and

microglia morphology, and to some extent, neurogenesis compared to partial cross-fostered pups.

These findings were contrary to our hypothesis and previous work demonstrating that presence

of a familiar sibling (when the entire litter is transferred) reduces the effect of cross-fostering on

circulating cortisol levels and weight gain, compared to pups that were cross-fostered on their

own (Arusha et al., 2024). In adult rats, the effects of corticosterone on hippocampal

neurogenesis vary depending on the dose, with studies showing increases, decreases, or no effect

(reviewed in Duarte-Guterman et al., 2017). It is possible that a lower-stress condition (such as

full cross-fostering based on cortisol levels and weight gain; Arusha et al., 2024) may

paradoxically produce greater effects on the hippocampal development compared to a moderate

stressor (such as partial cross-fostering). Further work is needed to investigate the impact of

stress hormones on hippocampal development in degus.

Conclusion

Because degus are precocial and fathers, in addition to mothers, may also contribute to offspring care, studies examining cross-fostering in this species may provide a more translationally relevant model for understanding how early-life caregiving environments, such as those experienced in adoption and foster care, shape developmental outcomes. In this context, our results demonstrated that cross-fostering did not affect volume or postnatal neurogenesis in the hippocampus. However, cross-fostering affected pyknotic cell density and microglia morphology, in a type of cross-fostering-, region- and sex-specific manner. Our findings complement earlier work with the same cross-foster degu model to examine its effects on HPA axis regulation (Guan et al., 2023; Arusha et al., 2024) and they further advance our understanding of how this early life stressor influences brain development. Finally, the results also highlight the importance of including both sexes and diverse forms of early-life stressors, such as cross-fostering paradigms. Further work is needed to determine the long-term effects of cross-fostering on brain function and potential behavioural consequences.

Acknowledgments

This research was supported by the Canada Research Chair Program to PDG and Swarthmore College to CMB. GA was funded by Ontario Graduate Scholarship and Faculty of Social Sciences at Brock University. SE was funded by Women's Science Research Fellowship and KB, DK, and KA were funded by Swarthmore College Summer Opportunities and Research Fellowships. We thank the animal care staff at Swarthmore College for technical assistance.

460 **Table 1**. Sample size for each group, and volume (mm^3) of the dentate gyrus, hilus, and granule cell

461 layer (GCL) in female and male degu pups in control, partial, and full cross-fostering conditions.

462 Estimated marginal means are presented ± SEM.

	Sample Size		Dentate Gyrus		Hilus		GCL	
	Females	Males	Females	Males	Females	Males	Females	Males
Control	7	7	6.32±0.42	6.26±0.40	3.41±0.30	3.51±0.28	2.91±0.19	2.75±0.17
Partial	5	6	6.14±0.52	6.38±0.44	3.55±0.37	3.61±0.31	2.59±0.23	2.77±0.19
Full	6	9	6.05±0.45	6.81±0.39	3.26±0.32	3.89±0.27	2.80±0.20	2.91±0.17

FIGURE CAPTIONS

Figure 1. Representative image of the dorsal (A) and ventral (B) hippocampus in degu pups. The

granule cell layer (GCL) and hilus are marked. Dentate gyrus volume (C) was not affected by partial

or full cross-fostering in male and female pups. Individual data points are shown with the estimated

marginal means (±SEM). Sample sizes were the following: control n=7 females, n=7 males; partial

cross-foster n=5 females, n=6 males; full cross-foster n=6 females, n=9 males.

Figure 2. A) Representative image of DCX expression in the dorsal (A) and ventral (B)

hippocampus of degu pups. Scale bars equal to 200 μm or 50 μm. C) DCX optical density (OD) in

the dorsal (D) and ventral (C) regions in female and male degus. Cross-fostering did not affect DCX

OD in either region in both sexes. Individual data points are shown with the estimated marginal

means (±SEM). Sample sizes were the following: control n=7 females, n=7 males; partial cross-

foster n=5 females, n=6 males; full cross-foster n=6 females, n=9 males.

Figure 3. (A) A depiction of Iba-1 expression within the region of interest (black rectangle box) in

the dentate gyrus of the hippocampus. Total number of microglia in the dorsal (B) and ventral (C)

hippocampus. There were no significant differences across cross-fostering groups in either region.

Individual data points are shown with the estimated marginal means (±SEM). Sample sizes were the

following: control n=7 females, n=7 males; partial cross-foster n=5 females, n=6 males (one male

sample was damaged and did not have dorsal sections); full cross-foster n=6 females, n=9 males.

Figure 4. Representative images of microglia morphology showing (A) ameboid, (B) intermediate,

and (C) ramified microglia (scale bar = 20 µm). (D) Full cross-fostered female pups had a lower

ameboid microglia density than control (* $p < 0.05$). Cross-fostering did not affect the density of

intermediate (E) or ramified microglia (F). Individual data points are shown with the estimated

marginal means (±SEM). Sample sizes were the following: control n=7 females, n=7 males; partial

cross-foster n=5 females, n=6 males (one male sample was damaged and did not have dorsal

sections); full cross-foster n=6 females, n=9 males.

Figure 5. (A) Representative image of pyknotic cells (arrows) in the granule cell layer in degu pups

(scale bar = 20 µm). Pyknotic cell density was affected by cross-fostering in the dorsal (B) and

ventral hippocampus (C). In the dorsal region, full cross-fostered pups had the lowest density of

pyknotic cells, followed by partial cross-fostered pups when compared to controls (** $p < 0.01$, ***

$p < 0.001$). In the ventral region, full cross-fostered pups had a lower density of pyknotic cells

compared to controls (* $p<0.05$). Individual data points are shown with the estimated marginal

means (±SEM). Sample sizes were the following: control n=7 females, n=7 males; partial cross-

foster n=5 females, n=6 males; full cross-foster n=6 females, n=9 males.

References

Aguggia, J.P., Suárez, M.M., Rivarola, M.A., 2013. Early maternal separation: neurobehavioral consequences in mother rats. Behav. Brain Res. 248, 25–31. https://doi.org/10.1016/j.bbr.2013.03.040

Arusha, K.S., Boadi, K.D., Ellah, S.S., Kim, D., Bauer, C.M., 2024. Sibling presence during fostering ameliorates endocrine stress profile changes in a social rodent species (Octodon degus) in a sex-specific manner. Horm. Behav. 166, 105660. https://doi.org/10.1016/j.yhbeh.2024.105660

Baek, S.-S., Jun, T.-W., Kim, K.-J., Shin, M.-S., Kang, S.-Y., Kim, C.-J., 2012. Effects of postnatal treadmill exercise on apoptotic neuronal cell death and cell proliferation of maternal-separated rat pups. Brain Dev. 34, 45–56. https://doi.org/10.1016/j.braindev.2011.01.011

Bannerman, D.M., Rawlins, J.N.P., McHugh, S.B., Deacon, R.M.J., Yee, B.K., Bast, T., Zhang, W.-N., Pothuizen, H.H.J., Feldon, J., 2004. Regional dissociations within the hippocampus—memory and anxiety. Neurosci. Biobehav. Rev., Festschrift in Honour of Jeffrey Gray - Issue 1: Anxiety and Neuroticism 28, 273–283. https://doi.org/10.1016/j.neubiorev.2004.03.004

Banqueri, M., Méndez, M., Gómez-Lázaro, E., Arias, J., 2019. Early life stress by repeated maternal separation induces long-term neuroinflammatory response in glial cells of male rats. Stress Amst. Neth. 22. https://doi.org/10.1080/10253890.2019.1604666

Bartolomucci, A., Gioiosa, L., Chirieleison, A., Ceresini, G., Parmigiani, S., Palanza, P., 2004. Cross fostering in mice: behavioral and physiological carry-over effects in adulthood. Genes Brain Behav. 3, 115–122. https://doi.org/10.1111/j.1601-183X.2003.00059.x

Boccia, M.L., Pedersen, C.A., 2001. Brief vs. long maternal separations in infancy: contrasting relationships with adult maternal behavior and lactation levels of aggression and anxiety. Psychoneuroendocrinology 26, 657–672. https://doi.org/10.1016/S0306-4530(01)00019-1

Burke, N.N., Fan, C.Y., Trang, T., 2016. Microglia in health and pain: impact of noxious early life events. Exp. Physiol. 101, 1003–1021. https://doi.org/10.1113/EP085714

Champagne, F., Meaney, M.J., 2001. Like mother, like daughter: evidence for non-genomic transmission of parental behavior and stress responsivity. Prog. Brain Res. 133, 287–302. https://doi.org/10.1016/S0079-6123(01)33022-4

Cohen, S., Ke, X., Liu, Q., Fu, Q., Majnik, A., Lane, R., 2016. Adverse early life environment increases hippocampal microglia abundance in conjunction with decreased neural stem cells in juvenile mice. Int. J. Dev. Neurosci. 55, 56–65. https://doi.org/10.1016/j.ijdevneu.2016.09.010

Colonna, M., Butovsky, O., 2017. Microglia function in the central nervous system during health and neurodegeneration. Annu. Rev. Immunol. 35. https://doi.org/10.1146/annurev-immunol-051116-052358

Colonnello, V., Iacobucci, P., Fuchs, T., Newberry, R.C., Panksepp, J., 2011. Octodon degus. A useful animal model for social-affective neuroscience research: Basic description of separation distress, social attachments and play. Neurosci. Biobehav. Rev. 35, 1854–1863. https://doi.org/10.1016/j.neubiorev.2011.03.014

Cornell, J., Salinas, S., Huang, H.-Y., Zhou, M., 2022. Microglia regulation of synaptic plasticity and learning and memory. Neural Regen. Res. 17, 705. https://doi.org/10.4103/1673-5374.322423

Delpech, J.-C., Wei, L., Hao, J., Yu, X., Madore, C., Butovsky, O., Kaffman, A., 2016. Early life stress perturbs the maturation of microglia in the developing hippocampus. Brain. Behav. Immun. 57, 79–93. https://doi.org/10.1016/j.bbi.2016.06.006

Diaz-Aparicio, I., Paris, I., Sierra-Torre, V., Plaza-Zabala, A., Rodríguez-Iglesias, N., Márquez-Ropero, M., Beccari, S., Huguet, P., Abiega, O., Alberdi, E., Matute, C., Bernales, I., Schulz, A., Otrokocsi, L., Sperlagh, B., Happonen, K.E., Lemke, G., Maletic-Savatic, M., Valero, J., Sierra, A., 2020. Microglia actively remodel adult hippocampal neurogenesis through the phagocytosis secretome. J. Neurosci. 40, 1453–1482. https://doi.org/10.1523/JNEUROSCI.0993-19.2019

Duarte-Guterman, P., Yagi, S., Epp, J.R., Brummelte, S., Wainwright, S.R., Barha, C.K., Chow, C., Barker, J.M., Galea, L.A.M., 2017. Hormone Regulation of Neurogenesis Across the Lifespan, in: Hormones, Brain and Behavior. Elsevier, pp. 373–410. https://doi.org/10.1016/B978-0-12-803592-4.00108-5

Fanselow, M.S., Dong, H.-W., 2010. Are the Dorsal and Ventral Hippocampus Functionally Distinct Structures? Neuron 65, 7–19. https://doi.org/10.1016/j.neuron.2009.11.031

Forger, N., 2009. Control of cell number in the sexually dimorphic brain and spinal cord. J. Neuroendocrinol. 21. https://doi.org/10.1111/j.1365-2826.2009.01825.x

Franco, R., Fernández-Suárez, D., 2015. Alternatively activated microglia and macrophages in the central nervous system. Prog. Neurobiol. 131, 65–86. https://doi.org/10.1016/j.pneurobio.2015.05.003

George, E.D., Bordner, K.A., Elwafi, H.M., Simen, A.A., 2010. Maternal separation with early weaning: a novel mouse model of early life neglect. BMC Neurosci. 11, 123. https://doi.org/10.1186/1471-2202-11-123

Gimsa, U., Brückmann, R., Tuchscherer, A., Tuchscherer, M., Kanitz, E., 2022. Early-life maternal deprivation affects the mother-offspring relationship in domestic pigs, as well as the neuroendocrine development and coping behavior of piglets. Front. Behav. Neurosci. 16. https://doi.org/10.3389/fnbeh.2022.980350

Glasper, E.R., Hyer, M.M., Hunter, T.J., 2018. Enduring Effects of Paternal Deprivation in California Mice (Peromyscus californicus): Behavioral Dysfunction and Sex-Dependent Alterations in Hippocampal New Cell Survival. Front. Behav. Neurosci. 12, 20. https://doi.org/10.3389/fnbeh.2018.00020

Gomez-Serrano, M., Tonelli, L., Listwak, S., Sternberg, E., Riley, A., 2001. Effects of cross fostering on open-field behavior, acoustic startle, lipopolysaccharide-induced corticosterone release, and body weight in Lewis and Fischer rats. Behav. Genet. 31. https://doi.org/10.1023/a:1012742405141

Guan, Q.S., Arusha, K.S., Caramihai, N., Stukal, I., Bauer, C.M., 2023. Fostered offspring develop hyper-reactive endocrine stress responses in a plural-breeding rodent, Octodon degus. Gen. Comp. Endocrinol. 336, 114259. https://doi.org/10.1016/j.ygcen.2023.114259

Hayashi, F., Takashima, N., Murayama, A., Inokuchi, K., 2008. Decreased postnatal neurogenesis in the hippocampus combined with stress experience during adolescence is accompanied by an enhanced incidence of behavioral pathologies in adult mice. Mol. Brain 1. https://doi.org/10.1186/1756-6606-1-22

He, Z., Guo, Q., Yang, Y., Wang, L., Zhang, S., Yuan, W., Li, L., Zhang, J., Hou, W., Yang, J.,
Jia, R., Tai, F., 2018. Pre-weaning paternal deprivation impairs social recognition and
alters hippocampal neurogenesis and spine density in adult mandarin voles. Neurobiol.
Learn. Mem. 155, 452–462. https://doi.org/10.1016/j.nlm.2018.09.006
Herpfer, I., Hezel, H., Reichardt, W., Clark, K., Geiger, J., Gross, C.M., Heyer, A., Neagu, V.,
Bhatia, H., Atas, H.C., Fiebich, B.L., Bischofberger, J., Haas, C.A., Lieb, K., Normann,
C., 2012. Early life stress differentially modulates distinct forms of brain plasticity in
young and adult mice. PLOS ONE 7, e46004.
https://doi.org/10.1371/journal.pone.0046004
Hulshof, H.J., Novati, A., Sgoifo, A., Luiten, P.G.M., den Boer, J.A., Meerlo, P., 2011. Maternal
separation decreases adult hippocampal cell proliferation and impairs cognitive
performance but has little effect on stress sensitivity and anxiety in adult Wistar rats.
Behav. Brain Res. 216, 552–560. https://doi.org/10.1016/j.bbr.2010.08.038
Huot, R.L., Plotsky, P.M., Lenox, R.H., McNamara, R.K., 2002. Neonatal maternal separation
reduces hippocampal mossy fiber density in adult Long Evans rats. Brain Res. 950, 52–
63. https://doi.org/10.1016/S0006-8993(02)02985-2
Johnson, F.K., Kaffman, A., 2018. Early life stress perturbs the function of microglia in the
developing rodent brain: new insights and future challenges. Brain. Behav. Immun. 69,
18–27. https://doi.org/10.1016/j.bbi.2017.06.008
Kempermann, G., 2016. Adult Neurogenesis: An Evolutionary Perspective. Cold Spring Harb.
Perspect. Biol. 8, a018986. https://doi.org/10.1101/cshperspect.a018986
Kentner, A.C., Abizaid, A., Bielajew, C., 2010. Modeling Dad: Animal models of paternal
behavior. Neurosci. Biobehav. Rev. 34, 438–451.
https://doi.org/10.1016/j.neubiorev.2009.08.010
Kheirbek, M.A., Hen, R., 2011. Dorsal vs ventral hippocampal neurogenesis: Implications for
cognition and mood. Neuropsychopharmacology 36, 373–374.
https://doi.org/10.1038/npp.2010.148
Kumazawa-Manita, N., Hashikawa, T., Iriki, A., 2018. The 3D stereotaxic brain Atlas of the
degu., Brain Science. Springer Japan, Tokyo. https://doi.org/10.1007/978-4-431-56615-
1_1
Loi, M., Mossink, J.C.L., Meerhoff, G.F., Blaauwen, J.L.D., Lucassen, P.J., Joëls, M., 2017.
Effects of early-life stress on cognitive function and hippocampal structure in female
rodents. Neuroscience 342, 101–119. https://doi.org/10.1016/j.neuroscience.2015.08.024
Lu, L., Mamiya, T., Lu, P., Niwa, M., Mouri, A., Zou, L.-B., Nagai, T., Hiramatsu, M.,
Nabeshima, T., 2009. The long-lasting effects of cross-fostering on the emotional
behavior in ICR mice. Behav. Brain Res. 198, 172–178.
https://doi.org/10.1016/j.bbr.2008.10.031
Madison, F.N., Palin, N., Whitaker, A., Glasper, E.R., 2022. Sex-specific effects of neonatal
paternal deprivation on microglial cell density in adult California mouse (Peromyscus
californicus) dentate gyrus. Brain. Behav. Immun. 106, 1–10.
https://doi.org/10.1016/j.bbi.2022.07.161
Majcher-Maślanka, I., Solarz, A., Chocyk, A., 2019. Maternal separation disturbs postnatal
development of the medial prefrontal cortex and affects the number of neurons and glial
cells in adolescent rats. Neuroscience 423, 131–147.
https://doi.org/10.1016/j.neuroscience.2019.10.033

Matthews, P.A., Samuelsson, A.-M., Seed, P., Pombo, J., Oben, J.A., Poston, L., Taylor, P.D., 2011. Fostering in mice induces cardiovascular and metabolic dysfunction in adulthood. J. Physiol. 589, 3969–3981. https://doi.org/10.1113/jphysiol.2011.212324

Oomen, C.A., Girardi, C.E.N., Cahyadi, R., Verbeek, E.C., Krugers, H., Joëls, M., Lucassen, P.J., 2009. Opposite effects of early maternal deprivation on neurogenesis in male versus female rats. PLOS ONE 4, e3675. https://doi.org/10.1371/journal.pone.0003675

Oomen, C.A., Soeters, H., Audureau, N., Vermunt, L., van Hasselt, F.N., Manders, E.M.M., Joëls, M., Krugers, H., Lucassen, P.J., 2011. Early maternal deprivation affects dentate gyrus structure and emotional learning in adult female rats. Psychopharmacology (Berl.) 214, 249–260. https://doi.org/10.1007/s00213-010-1922-8

Paolicelli, R.C., Sierra, A., Stevens, B., Tremblay, M.-E., Aguzzi, A., Ajami, B., Amit, I., Audinat, E., Bechmann, I., Bennett, M., Bennett, F., Bessis, A., Biber, K., Bilbo, S., Blurton-Jones, M., Boddeke, E., Brites, D., Brône, B., Brown, G.C., Butovsky, O., Carson, M.J., Castellano, B., Colonna, M., Cowley, S.A., Cunningham, C., Davalos, D., De Jager, P.L., de Strooper, B., Denes, A., Eggen, B.J.L., Eyo, U., Galea, E., Garel, S., Ginhoux, F., Glass, C.K., Gokce, O., Gomez-Nicola, D., González, B., Gordon, S., Graeber, M.B., Greenhalgh, A.D., Gressens, P., Greter, M., Gutmann, D.H., Haass, C., Heneka, M.T., Heppner, F.L., Hong, S., Hume, D.A., Jung, S., Kettenmann, H., Kipnis, J., Koyama, R., Lemke, G., Lynch, M., Majewska, A., Malcangio, M., Malm, T., Mancuso, R., Masuda, T., Matteoli, M., McColl, B.W., Miron, V.E., Molofsky, A.V., Monje, M., Mracsko, E., Nadjar, A., Neher, J.J., Neniskyte, U., Neumann, H., Noda, M., Peng, B., Peri, F., Perry, V.H., Popovich, P.G., Pridans, C., Priller, J., Prinz, M., Ragozzino, D., Ransohoff, R.M., Salter, M.W., Schaefer, A., Schafer, D.P., Schwartz, M., Simons, M., Smith, C.J., Streit, W.J., Tay, T.L., Tsai, L.-H., Verkhratsky, A., von Bernhardi, R., Wake, H., Wittamer, V., Wolf, S.A., Wu, L.-J., Wyss-Coray, T., 2022. Microglia states and nomenclature: A field at its crossroads. Neuron 110, 3458–3483. https://doi.org/10.1016/j.neuron.2022.10.020

Priebe, K., Brake, W.G., Romeo, R.D., Sisti, H.M., Mueller, A., McEwen, B.S., Francis, D.D., 2005. Maternal influences on adult stress and anxiety-like behavior in C57BL/6J and BALB/cJ mice: A cross-fostering study. Dev. Psychobiol. 47, 398–407. https://doi.org/10.1002/dev.20098

Qiu, L.R., Fernandes, D.J., Szulc-Lerch, K.U., Dazai, J., Nieman, B.J., Turnbull, D.H., Foster, J.A., Palmert, M.R., Lerch, J.P., 2018. Mouse MRI shows brain areas relatively larger in males emerge before those larger in females. Nat. Commun. 9, 2615. https://doi.org/10.1038/s41467-018-04921-2

R Core Team, 2021. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Raineki, C., Opendak, M., Sarro, E., Showler, A., Bui, K., McEwen, B.S., Wilson, D.A., Sullivan, R.M., 2019. During infant maltreatment, stress targets hippocampus, but stress with mother present targets amygdala and social behavior. Proc. Natl. Acad. Sci. 116, 22821–22832. https://doi.org/10.1073/pnas.1907170116

Sierra, A., Beccari, S., Diaz-Aparicio, I., Encinas, J.M., Comeau, S., Tremblay, M.-È., 2014. Surveillance, phagocytosis, and inflammation: How never-resting microglia influence adult hippocampal neurogenesis. Neural Plast. 2014, 610343. https://doi.org/10.1155/2014/610343

681   Walker, F.R., Nilsson, M., Jones, K., 2013. Acute and chronic stress-induced disturbances of
682   microglial plasticity, phenotype and function. Curr. Drug Targets 14, 1262–1276.

683   Wickham, H., 2016. ggplot2: elegant graphics for data analysis. Springer-Verlag, New York.

684   Yang, S., Li, J., Han, L., Zhu, G., 2017. Early maternal separation promotes apoptosis in dentate
685   gyrus and alters neurological behaviors in adolescent rats. Int. J. Clin. Exp. Pathol. 10.

686   Zhang, L.-X., Levine, S., Dent, G., Zhan, Y., Xing, G., Okimoto, D., Kathleen Gordon, M., Post,
687   R.M., Smith, M.A., 2002. Maternal deprivation increases cell death in the infant rat brain.
688   Dev. Brain Res. 133, 1–11. https://doi.org/10.1016/S0926-6410(01)00118-5

Figure 1

Figure 3

Figure 4

Figure 5