Title: Distant Site Mutations in Clinical TEM Beta-Lactamase Variants Enhance Non-Covalent Binding to Ceftazidime: Insights from Spectroscopic and Biophysical Investigations.

Sandip Kumar Mukherjee [a][1][*], Padmaja Prasad Mishra [b], Mandira Mukherjee [a][*]

[a] Department of Biochemistry and Medical Biotechnology, School of Tropical Medicine,

Kolkata, West Bengal, 700073, India

[b] Chemical Sciences Division, Saha Institute of Nuclear Physics, 1/AF Bidhannagar, Kolkata,

West Bengal, 700064, India

Present address:

[1] Department of Microbiology, Immunology and Molecular Genetics, University of California,

1602 Molecular Sciences Building 609 Charles E. Young Drive, East Los Angeles, CA 90095,

USA

*Correspondence:

Dr. Mandira Mukherjee,

E-mail: mandira_71@yahoo.com

Mobile no: 91-9433948556

https://orcid.org/0000-0003-4123-1377

Dr. Sandip Kumar Mukherjee

E-mail: sansbubun@gmail.com

https://orcid.org/0000-0001-9958-5237

Abstract

β-lactamases retain the central armamentarium against the β-lactams, resulting in surge of antibiotic resistance, primarily due to the hydrolysis of the amide bond of the four-membered ring. This study aimed to investigate the binding interactions of ceftazidime (CAZ) to TEM β-lactamase variants with distant site mutations isolated from clinical setting to explore the cause of their selection and dissemination due to indiscriminate use of β-lactams. Absorbance and fluorescence spectroscopy along with biophysical experimentation indicated facilitated binding of CAZ to the active site of the mutants than the wild type. The CAZ-TEM β-lactamase mutant interactions were predominantly hydrophobic compared to H-bonding and van der Waals forces in the CAZ-wild type complex. Additionally structural alteration to justify more rigid binding of CAZ to the mutants in contrast to the wild type enzyme was established *in silico*. Therefore, acquisition of distant site mutations with respect to the active site of the β-lactamase variants that rendered conformational flexibility to accommodate CAZ was evidenced. This study provided an insight to the bioactive interaction of CAZ with TEM β-lactamase variants that probably facilitated their selection from clinical settings in response to rampant usage of β-lactams.

Key words:

TEM β-lactamase mutants, ceftazidime, absorbance spectroscopy, fluorescence quenching,

fluorescence anisotropy, Van't Hoff plot, Molecular dynamics simulation

1. Introduction:

Persistence of bacterial resistance against antimicrobial agents is a global health threat (Avci et al. 2016). Production of active site serine β-lactamase enzymes is the major mechanism of antibiotic resistant in uropathogenic *Escherichia coli* (UPEC) that catalyze the hydrolysis of β-lactam antibiotics. (Chinemerem et al. 2022,Natalia Dı´az et al.2003,Stojanoski et al.2015). Currently, based on Ambler classification scheme, β-lactamases can be grouped into four classes A, B, C and D. Class A, C, and D are serine β-lactamases (non-metallo) that utilize the serine residue to catalyze the hydrolysis of antibiotics, whereas class B enzymes are metallo-β-lactamases and need zinc ions in the active site to attack the β-lactam ring(Stojanoski et al.2015).In the case of Class A β-lactamases which is prime concerned for the clinicians due to its wide range of hydrolyzing activity against antibiotics mostly against β-lactam antibiotics where activation of catalytic Ser70 conserved residue initiates nucleophilic attack on the β-lactam carbonyl carbon and opening of the β-lactam ring to form a covalently linked acyl-enzyme intermediate and ultimately the attack leads to deacylation of the acyl-enzyme species resulting inactivation of antibiotic and regeneration of the free enzyme (Stojanoski et al.2015).TEM1 β-lactamase was identified as one of the most evolved Class A β-lactamases in hospital infection, mediated by plasmid found in *E. coli* resistant to antibiotics and also considered as the precursor of the TEM enzyme family with three major types of activity; the broad-spectrum β-lactamases, the extended spectrum β-lactamases (ESBLs) and the inhibitor resistant β-lactamases. The emergence of resistance mediated by this TEM enzyme family were most explored β-lactamases till date and it was found that their extensive structural adjustment by their sustainable mutational modifications led to the generation of distinct multiple variants with in-depth structural diversity (Singhet al.2012). To overcome β-lactam resistance extensive studies were carried out against the acylation mechanism and two types of β-lactams were introduced such as extended spectrum cephalosporins; and β-lactamase inhibitors, such as clavulanic acid, sulbactam and tazobactam [(Kumar et al.2014). Furthermore, the structural alterations in Class A β-lactamases impair the antibacterial activities of the inhibitors by covalent interaction between the active site residues in the former with the inhibitory molecules resulting in resistance against them (Drawz and Bonomo 2010). Till date ~200 TEM-variants have been reported that confers resistance to novel β-lactams and their inhibitor combinations which were introduced in the past three decades (Barlow et al, 2010. Pimenta et al,2014). Studies from the

different part of the world showed that the indiscriminate use of β-lactam-β-lactamase inhibitor
combination has increased the number of TEM β-lactamase variants; the extended-spectrum (ES)
and inhibitor-resistant phenotypes (IR) (Tooke et al.2019,Cantón et al. 2008). Previous reports
showed that mutation or substitution in the active site Ω-loop region of TEM β-lactamase not
only increased the resistance phenotype against cephalosporins but also increased the structural
flexibility imparting enlargement of the active site pocket loop that can accommodate newer β-
lactams irrespective of their -R group thus decreasing the activity of oxyimino-cephalosporin,
ceftazidime (CAZ), and all other β-lactam-β-lactamase inhibitors (Natalia Dı´az et
al.2003,Stojanoski et al.2015). Fluorescence techniques were widely used as one of the important
biophysical application that paved sensitivity, reliability and rapidity in the field of drug-protein
interactions study (Moller et al.2002,Kumar et al. 2024).

There are not many Indian studies that described existence and characterization of TEM β-
lactamase mutants that harbor mutations at sites other than the active site of the enzyme isolated
from clinical origin as a result of random drug administration. An earlier study from our
laboratory, demonstrated identification and characterization of three novel TEM β-lactamase
mutants; pUE184TEM (M184), pUE203TEM (M203), pUE210TEM (M210) from clinical
uropathogenic *E. coli* isolates with mutation at residues other than those that constituted the
active site of the enzymes (Mukherjee S K et al, 2018). In the present study bioactive interactions
of CAZ with the mutant β-lactamases under physiological conditions was investigated. The role
of distant site mutations on the conformational flexibility in the mutants to accommodate CAZ
was explored by using a collection of biophysical and *in silico* methods to obtain an insight into
selection and dissemination of the resistant β-lactamases in response to uncontrolled β-lactam
administration.

2. Materials and Methods:

Absorbance Spectroscopy:

Absorbance spectral measurement wasrecorded in UV–vis Spectrometer at a scan rate of 480
nm/min with 1 cm-path-lengthquartz cuvette in the visible wavelength range of 200-800nmat
different temperatures (298K, 303K, 308K, 311K).The titrations of the experiment were
conducted on 5 μM wild type TEM β-lactamase and its variants (M184, M203, M210)
(Mukherjee S K et al, 2018) in presence ofascendingconcentration of CAZ (5 μM-20μM) at
respective temperatures.The inner filter effect was eliminated using the relationship; I_{cor} =

$I_{obs}e^{(Aex+Aem)/2}$,where the corrected fluorescence intensity is denoted as I_{corr}, and observed

fluorescence intensity is denoted as I_{obs}. The A_{ex} and A_{em} denote absorbance values at excitation

and emission wavelengths, respectively (Swain et al. 2020)

Fluorescence Spectroscopy:

Fluorescence emission spectrum was recorded in the range of280–600 nm using Hitachi F-7000

Spectrofluorimeter with excitation at 295 nm. Both the excitation and emission bandwidth were

fixed at5 nm. Quenching experiment was carried out in presence of increasing concentration of

CAZ (5 μM-20μM) solution at different temperatures (298K, 303K, 308K, 311K) to 5 μM

concentration of the wild type and the mutant β-lactamases.Static fluorescence quenching was

determined by using the Stern–Volmer (SV) equation from steady state measurements $F_0/F = 1 +$

$K_{sv}(Q)$, where F_0 and F denote the steady-state fluorescence emission intensities in the absence

and in presence of CAZ. K_{SV} is the Stern–Volmer constant and (Q) is the concentration of the

quencher. Collisional or dynamic fluorescence quenching was determined using the modified

Stern-Volmer equation; $\log ((F_0 -F)/F)) = \log K_a + n \log (Q)$, where K_a denoted the binding

constant and n the number of binding sites.

Fluorescence lifetime measurements

The fluorescence lifetime(τ) measurements determine energy transfer at the molecular level

between interacting species. Time correlated single photon counting (TCSPC) was used to

measure the fluorescence life times of wild type and mutant β-lactamases in gradually increasing

concentration of CAZ (0 μM-20μM) in sodium phosphate buffer (pH 7.4) at 298K (Maldal S K

et al, 2012).

Quantum yield and fluorescence anisotropy

The quantum yield (Φ) or quantum efficiency of CAZ at 5μM concentration in sodium

phosphate buffer (pH 7.4) was recorded by monitoring the fluorescence emission intensityin

presence of increasing concentrations of the wild type and mutant β lactamases (0-20 μM)at

298K (Mishra et al, 2006). The fluorescence anisotropy (r) measurement of CAZ at 5μM

concentration in 5mM sodium phosphate buffer (pH 7.4) was recorded against addition of

ascending concentration of wild type and mutant proteins (5μM-20μM) in Fluromax

Spectrofluorometer. All measurements were conducted at 298K at 3 minutes of interval.

Thermodynamic properties

Van't Hoff isotherm equation was used to determine the thermodynamic parameter (kinetic and

equilibrium properties) that govern the interaction between the ligands (CAZ) and the proteins

(TEM and its mutants). The values of ΔH^0 and ΔS^0 were obtained from the *lnK* vs. 1/T plot,

where ΔG^0, ΔH^0, and ΔS^0 are depicted as standard enthalpy change, standard free energy change,

and standard entropy change respectively.

Circular Dichroism (CD) Spectroscopy:

Far and Near UV-CD spectra was recorded at room temperature for β-lactamases (TEM and its

mutants) in presence of ascending concentration of ceftazidime ranges from 0 μM-20μM

solutions using JASCO-J815 spectrometer with a temperature controller at 298K in a quartz

cuvette of 1 mm path length. Each final spectrum was recorded as an average over three scans.

The far and near-UV-CD region was recorded as 250–360 nm and 360 nm to 500 nm.CD results

were expressed as mean residue elasticity (MRE) in degree cm^2 mol^{-1}. The secondary structures

(α-helix, β-sheet, random coils) were calculated from respective MRE values (Bertucci et

al.2010) **Raman spectroscopy**

Raman scattering was recorded for β-lactamases (TEM and its mutant variants) in presence of

ascending concentration of ceftazidime ranges from 0μM-20μM. The spectral range was taken

for500–4000 cm^{-1} with an integration time of 20 s using a 670 nm red laser excitation, at

10 mW power (Miani et al.2007).

Förster resonance energy transfer (FRET) computation

FRET involves a distance dependent physical method of non-radiative transfer of energy

between an excited fluorophore (donor) and another fluorophore (acceptor) by intermolecular

dipole-dipole interactions. The energy transfer efficiency (E) signified the fraction of photons

absorbed by the donor moiety which was transferred to the acceptor. $E= 1-(F/F_0) = R_0^6/(R_0^6 + r^6)$,

where, r represented the distance between the donor and acceptor and R_0 represented the critical

energy transfer distance, at which 50% of the excitation energy was transmitted from the former

to the latter. R_0 was evaluated from the following equation; $R_0^6 = 8.79 \times 10^{-25}K^2N^{-4}$, where K^2

was the spatial orientation factor of the altered dipoles of the acceptor and the donor, N

represented the refractive index of the medium, denoted the fluorescence quantum yield of the

donor in absence of the acceptor. (Chakraborty et al.2015).

MD simulation

Molecular dynamics and simulation was performed on the TEM β-lactamase (1ZG4) and its variants (Mukherjee S K et al, 2018) in presence of CAZ to determine the binding stability of the protein–drug docked complex using Desmond v3.6 Package. Topology of the protein molecules were generated by optimized potentials for liquid simulation (OPLS) force field. Simple point charge (SPC) water model was used to perform the protein–drug complexes Simulation with a distance of 10 Å between the complex and the periodic box. System was neutralized by adding Na+/Cl– ions and placed randomly in the solvated system. To minimize and relax the protein/protein–drug complex under NPT ensemble, the default protocol of Desmond was used followed by total of nine stages among which two minimization and four short simulation (equilibration phase) steps was involved before starting the actual production time. System temperature and pressure was 300 K and 1 atmosphere, respectively, using Nose–Hoover temperature coupling and isotropic scaling, and the simulation was performed for 100 ns NPT production simulation and saving the configurations at 10 ns intervals (Mukherjee S K et al, 2021).

Simulation trajectory analysis

Simulation trajectory files were analyzed using Desmond module programs i.e. simulation quality analysis (SQA), simulation event analysis (SEA), and simulation interaction diagram (SID) to calculate energies, root-mean-square deviation (RMSD), root-mean-square fluctuation (RMSF), the radius of gyration (Rg) and the secondary structure elements (SSEs) which pave the protein structure stability. SQA was used to qualitatively validate the system stability throughout the simulated length of chemical time for the given temperature, pressure, and volume of the total simulation box, SEA was used to analyze each frame of the simulated trajectory output, whereas total SSE change in the protein structure in presence of the drug during the simulation time was estimated using SID (Abdul et al, 2016, (Matyus et al, 2006)).

Results:

Spectroscopic measurements:

UV-Vis absorption spectra of wild type and mutant β-lactamases (**Fig. 1A-D**) were recorded in presence of increasing concentrations of ceftazidime, at 298K. All protein showed a broad band at 280 nm while ceftazidime had a strong absorbance at 267 nm and a moderate absorbance at 310 nm. With gradual increase of CAZ concentration, the absorbance spectra of wild β-lactamase showed enhancement of absorption intensity.

The possibility of ground state complex formation couldnot be ruled out from the above
absorption spectra. Additionally, from **Fig. S1** it was observed that the spectra of the mixture of
CAZ and wild type β-lactamase (**curve 5**) was much intense than intensity of spectra
corresponding to the summationof the two absorption bands of the individual reacting
components (**curve 3**). The **curve 4** obtained by subtracting the **curve 3** from the **curve 5**
indicated formation of ground state complex between the two species.
The UV-Vis absorption spectra of the wild type and mutant β lactamases in presence of CAZ
were recorded at 303 K, 308 K and 311 K respectively which indicated that the proteins-CAZ
complexes were thermally stable (**Fig. S2**).
The fluorescence spectra of wild type and mutant β-lactamases at pH 7.5in the presence of 5, 10,
15, 20μM CAZ at 298K showed strong fluorescence intensity at 340 nm upon excitation at 280
nm which gradually decreased upon addition of CAZ with a slight red shift at 298 K (**Fig. 1E-H,**
Fig. S3). Similar comparable fluorescence spectra of the β-lactamases at the different
temperatures (303K.308K, 311K) were also recorded (data not shown).
The quenching phenomenon exhibited in the fluorescence emission spectra of the proteins and
protein –CAZ complexes were analyzed by Stern–Volmer (SV) plot. The non-linearity observed
in the SV plot at 298K, 303K, 308K, 311K (**Fig. 2A-D**) indicated a combination of static (ground
state) and dynamic (excited state) quenching which was validated by modified SV plots (**Fig.**
2E-H). Additionally, SV plots of quenching of the wild type in presence of increased
concentration of CAZ at different temperatures indicated a gradual shift towards dynamic mode
although mutants exhibited a decrease in their dynamic mode of quenching.
Furthermore, fluorescence lifetime changes of the wild type and the mutant proteins in presence
of different concentrations of CAZ also exhibited a steady decrease (**Fig. 3A, Table 1**).
Fluorescence quantum yield and fluorescence anisotropyof CAZ
Absolute fluorescence quantum yield was also analyzed for small molecule i.e. CAZ to measure
changes in radiative rates from changes in nonradiative rates occurring in terms of interaction.
The quantum yields of the CAZ molecule increased with the increase in concentration of proteins
(wild and all three mutant β-lactamases). The quantum yields of CAZ increased gradually from
0.08 to 0.27, 0.36, 0.36 and 0.37 for the wild type, M184, M203 and M210 β-lactamase
respectively (**Table 2**).Fluorescence anisotropy was also performed to measure the
microenvironment around CAZ in terms of rotational diffusion and interactions. The anisotropy

of the CAZ molecule was very low (r = 0.07). With increased concentration of the proteins (wild

type and all three mutant β-lactamases), fluorescence anisotropy of CAZ increased gradually. At

20μM concentration of CAZ a steady increase in r value from 0.18, 0.20, 0.21 to 0.24 in

presence of the respective β-lactamases (wild type, M184, M203, M210) respectively (**Fig. 3B**)

Binding constants, number of binding sites and mode of binding

The steady state fluorescence data were further evaluated by the binding affinity (K_b) of CAZ

molecule to the equivalent sites of wild type and mutant β-lactamases independently. For mutant

β-lactamases a progressive decrease of binding affinity (K_b) was observed with the increase of

temperature whereas in case of wild type a vice versa affinity was observed in pH 7.5.(**Fig. 4**).

The equilibrium interaction (such as, hydrogen bonds, van der Waals forces, electrostatic forces,

and/or hydrophobic associations), between ligand (CAZ) and protein molecule were determined

from the value of the thermodynamic parameter (ΔG^0, ΔH^0, ΔS^0)using van't Hoff isotherm

equation from the lnK vs. 1/T plot. Significant negative ΔG^0value, $\Delta H^0 < 0$ and $\Delta S^0 > 0$ indicated

spontaneous interaction between the reacting components. Increase in negative ΔG^0 value with

increase in positive ΔH^0 and ΔS^0 values in the mutantβ-lactamases indicated discrete

predominance of hydrophobic interaction in contrary to the wild type β-lactamase that exhibited

van der Waals and H-bond interaction with high negative $\Delta H^0$and ΔS^0 values compared to less

negative ΔG^0 value (**Fig. 4, Table 3**).

Circular Dichroism (CD) Spectroscopy:

Conformational changes of wild and mutant β-lactamases were further measured by CD

spectroscopy in the absence and presence of CAZ at pH 7.5. Results showed that the mean

residue ellipticity (MRE) of pure α-helix percentage of wild and mutant β-lactamases were

decreased upon binding of CAZ demonstrated secondary structure alternations of the proteins in

far-UV-CD region (185–250 nm) with two characteristic dips at 209 nm and 222 nm for all the

proteins respectively. Moreover, in the near-UV-CD region spectra (240–320 nm) of the wild

type and mutant proteins showed comparable minor alterations of their tertiary structure (**Fig.

S4, Table 4**)

Raman spectrum of β-lactamaseproteins

Raman intensity of wild type and mutant β-lactamases (5μM) were measured in presence of

ascending concentration of CAZ (5 μM-15 μM). Results showed a gradual increase of Raman

intensity of β-lactamases with the increase of the concentration of CAZ in comparison with the

wild type protein. There are three regions (880 cm^{-1}, the region of Tyr doublet due to H-

Bonding),(1360 cm^{-1}, CH2- CH2 deformation) and (1640-1680 cm^{-1}, characteristics of amide

peak) in the spectrum which indicated the secondary structure alternation due to molecular

interaction. A small blue shift was also observed for the mutant β lactamases in comparison with

the wild type due to secondary structure adjustment during interaction with CAZ. (**Fig. S5**)

Energy transfer between CAZ and β lactamases

The absorbance spectrum of CAZ (excited at 267 nm) and fluorescence spectrum of the TEM 1

β- lactamase at 340 nm overlapped with each other which satisfied the requisite condition of

FRET between the CAZ and the proteins **(Fig. S6)**. Moreover from the Förster theory of energy

transfer, the molecular interactions between the drug-protein complexes were evidenced from

discrete values of the energy transfer efficiency (E); 0.47, 0.94, 0.79 and 0.82 for CAZ-wild type,

CAZ-M184, CAZ-M203 and CAZ-M210 respectively which facilitated binding of CAZ to the

mutants than the wild type.

In silico **study**

Earlier *in silico* study from our laboratory indicated effective binding of CAZ to the wild type

and mutant β lactamases (Mukherjee S K et al, 2018). The stability and conformational changes

of the β lactamase in presence of CAZ were analyzed by MD simulation study.The movement of

atoms and molecules for 100 ns of simulation were monitored. In presence of CAZ, the wild

type β lactamases (1ZG4) and M184 exhibited C-α backbone deviations at about ∼1.5 to 1.75 Å

and ∼1.75 to 2.5 Å respectively throughout the simulation trajectory and an equilibrium was

maintained after 40 ns for both. However, for M203 and M210 C-α backbone deviations from

∼1.5 to 2.5 and 1.75 to 2.5 Å with an elevation at 60 ns for the latter which reached equilibrium

from 70 ns was observed. The C-α all residue deviations for M184 and M210 were ∼2.5 to 3.25

Å and ∼2.5 to 3.40 Å respectively with an elevation at 60 ns and equilibrium from 70ns in case

of the latter. In contrast, M203 showed reduced Cα backbone deviation at 30 ns akin to wild type

throughout the 100 ns simulation trajectory. However, M203 showed C-α all residue deviations

from ∼2.25 to 3.0 in comparison to wild type in presence of CAZ (**Fig. 5 A-F**)

The residue level fluctuations (RMSF) of Cα backbone for M184 and M210 in presence of CAZ

were moderately high (∼3.25 Å) in comparison to wild type at residues located between 50-80,

120-170, 200, and 225-250 respectively. However, a comparable pattern of fluctuation was

observed for M203 and wild type except at residue 150. Similarly, RMSF for Cα side chain of

the mutant proteins (M184, M203, M210) in presence of CAZ compared to wild type in presence

of the drug showed similar pattern as the RMSF for Cα backbone (**Fig. 5 G-L**).

The compactness of protein-drug interaction was ascertained by Cα Radius of gyration (Rg) plot.

The residue level Rg value fluctuation for wild type protein was within the range of ∼17.8 to

18.3 Å, whereas for mutant protein (M184) was within the range of ∼17.8 to 18.2 Å and reached

equilibrium from 40 ns onward in presence of CAZ. Furthermore, in presence of the drug the Rg

value fluctuations for M203 was ∼17.8 to 18.1 across the 100 ns simulation trajectory but for

M210, the Rg value fluctuations was ∼17.9 to 18.6 Å with a decrease in fluctuation within the

range of ∼18.2 to 18.4 Å from 60 ns onward to attain the equilibrium. The overall Rg analysis

for compactness in the mutants in presence of CAZ showed lowest radius of gyration that

demonstrated tight binding of CAZ to the mutant proteins in comparison to wild type (**Fig. 5 M-**

O).

The protein-drug interactions were further monitored by estimation of secondary structure

elements (SSE) composition (helices, strands, turns, loops) in the interacting proteins across the

simulation trajectory throughout the 100 ns time frame. SSE for each amino acid residue of the

mutants exhibited significant structural adjustment in comparison with wild type in presence of

CAZ. In M184 increase of 310-helix in residue positions 100, 160, 220, turns in residue position

50, 90, 160, coils in position 140, 200 and π-helix in positions 210 to 225 was evidenced. In

M203, the segment of the residues in positions 60-90 showed increase of 310-helix, coil structure

at residues 150-170 and 225-240. M210 showed an increase of α-helix and coil with the absence

of π-helix at positions 60-90 and 160-170 in comparison with the segment of residues in the

wild-type with increase of 310-helix and increase of π-helix at residue position 210-225

respectively (**Fig. 5 P-T**).

324   **5. Discussion**

Emergence of varied β-lactamases was reportedwhich was the primary cause of β-lactam

resistance in uropathogenic *E. coli*, the primary etiologic agent of urinary tract infection. These

β-lactamases were found to harbor mutations which paved the possibility of effective binding of

the different β-lactams and their inhibitor combinations to develop resistance. Therefore, studies

on the protein-drug interactions have become increasingly important from a pharmaceutical

perspective.Absorption spectroscopy study provide information on the binding modes of ligands
with proteins pave the possibility of ground state complex formation between protein-ligand
interfaces (Swain et al, 2020). The individual UV-Vis absorption spectrum of the CAZ and
TEM-1 β-lactamase were markedly different to that of CAZ-β-lactamase complex, indicating the
formation of new complexes between CAZ and β-lactamases. Appearance of isosbestic points at
339.4 nm, 360.2 nm, 341.4 nm and 340 nm for the TEM-1-CAZ, M184-CAZ, M203-CAZ and
M210-CAZ systems respectively suggested that an equilibrium was established between the
respective two species and reflected the formation of complexes between them. Therefore, these
observations signify that the ground state complexes are formed between respective wild type
and mutant β-lactamase proteins and CAZ. The UV-Vis absorption of the four β-lactamases
increased and a slight red-shift that occurred with an increase in concentration of CAZ, which
indicated that, the drug-protein interaction resulted in subtle conformational changes in the
protein in presence of the drug. A hyper- chromic effect and slight red-shift of the maximum
absorption peak were also separately observed with increasing CAZ concentrations in solution.
The shift of maximum wavelength signifies hydrophobic effect on Trp residue due to complex
formation between drug and respective proteins (Topala et al. 2014). However, the observations
obtained from absorption studies were not sufficient to study the interactions in detail.
Therefore, other excited state spectroscopic techniques e.g., fluorescence spectroscopy, TCSPC
etc. were preferred to study the binding mode of protein-drug interactions. Variation in
temperature produced a local change in the protein microenvironment as heat disrupted the
hydrogen bonds, non-polar hydrophobic interactions, which might produce a local change in the
Trp microenvironment and affected the spectroscopic properties of the protein. In this study, the
increase of quantum yieldof wild type and mutant β-lactamases with the ascending temperatures
represented the change in the local hydrophobicity in presence of CAZ. Moreover, a decrease in
fluorescence intensity (quenching) of the β-lactamases (wild type and mutants) with increase in
temperature in presence of CAZ alsoindicated formation of a new complex.which corroborated
with the change in the protein microenvironment (Topala et al. 2014,Ghisaidoobe et
al.2014,Butler et al.2015.Bardhan et al. 2011)
Additionally, in CAZ-β lactamases system the quenching mechanism might also be either due to
simultaneous occurrence of static (ground state) and dynamic (excited state) interaction modes.
The static quenching had already been established by ground state complex formation. However,

the decrease in fluorescence lifetime of wild type and mutant β lactamases with simultaneous
addition of CAZ accounted only for dynamic quenching supported by the linear Stern-Volmer
plot. Moreover, the positive deviation of Stern-Volmer plot for CAZ-β-lactamase systems
demonstrated the simultaneous occurrence of static and dynamic quenching. Furthermore, the
Stern-Volmer plots of the wild type β-lactamase in presence of increasing concentration of CAZ
at different temperatures indicated a gradual shift towards dynamic mode but for mutant-CAZ
complexes a sharp shift towards dynamicity indicated a more rigid interaction of the drug to the
mutant enzymes (Mukherjee S K et al, 2021Bardhan et al. 2011)
Determination of binding constant and binding stoichiometry of β-lactamases in presence of
CAZprovided evidence for both static and dynamic quenching.The thermodynamic parameters
of binding, i.e., changes in standard enthalpy (ΔH^0), entropy (ΔS^0) and Gibb's free energy (ΔG^0)
using van't Hoff isotherm exhibited the nature of interacting forces between drug and protein.
Usually, four types of forces play an important role in drug–protein interaction, like electrostatic
forces, hydrophobic forces, van der Waals interactions and hydrogen bonding (Ross et al.1981).
In this study it was found that the interaction of CAZ with the mutant proteins was
predominantly hydrophobic, however van der Waal's interactions and hydrogen bonding was the
primary interacting forces in the binding of CAZ to the wild type enzyme. Furthermore, the
anisotropy changes demonstrated flexible interactions of the drug with the mutants than the wild
type protein which also supported predominance of hydrophobic interaction between the drug
and mutant enzymes (Zhao et al. 2019).
The circular dichroism (CD) is a sensitive technique to monitor the conformational changes in
proteins upon interaction with ligand molecules (Bertucci et al.2010). CD spectra of the wild
type and mutant proteins in presence of CAZ further corroborated conformational changes in the
secondary as well as tertiary structure alterations in the protein at pH 7.0. Additionally,
assessment of energy transfer efficiency values (E) using FRET technique and Raman
spectroscopy provided evidence that the specific mutations in the M184, M203 and M210
assisted an effective binding of CAZ to the mutant β lactamases than to the wild type protein
(Chakraborty et al 2015,Lee et al 2016,,Pezzotti et al 2021)
.
In silico MD simulation analysis of the wild type and the mutant β lactamases revealed increase
in the RMSD for the C-α backbone for the mutant β lactamases which might be attributed to the

overall conformational flexibility of the protein to accommodate the drug molecule (Mukherjee et al.,2021,Gill et al. 2019]. Therefore, the overall comparison of RMSD plots for wild type and the variants showed that the C-α backbone residues in the latter maintained stabilizing interaction with the drug molecule.

Similarly, RMSF for Cα side chain of the mutant proteins (M184, M203, M210) compared to wild type in presence of the drug showed similar pattern as the RMSF for Cα backbone which further documented the enhanced conformational flexibility in the mutant β-lactamases to bind to CAZ than the wild type (Mukherjee et al.,2018,Fantini et al 2020). The Cα Radius of gyration (Rg) for compactness in the mutantβ lactamases in presence of CAZ also demonstrated a tight binding of the drug to the mutant proteins in comparison to wild type (Mukherjee et al.,2021,Galdadas et al.2021). Furthermore, the residue-based conformational changes in the mutants in presence of CAZ also indicated that the altered secondary structure elements (SSE) in the mutants probably imparted additional overall stability and rendered conformational flexibility compared to the wild type protein (Fantini et al 2020). This *in-silico* study established that the mutant-CAZ interactions were more stable and rigid with minimum conformational alterations between the interacting residues compared to the wild type-CAZ interaction (Yang et al 2020).

Conclusion

This is the first of its kind research that illustrated the comprehensive bioactive interaction of CAZ with three novel mutants of TEM β-lactamases derived from clinical isolates of uropathogenic Escherichia coli at the molecular level via spectroscopic and biophysical approach. β-lactamases remain at the forefront of driving antibiotic resistance via hydrolysis of β-lactam antibiotics, necessitating the understanding of their evolving mechanisms. We explained the role of far-site mutations in clinical TEM β-lactamase variants that enhance ceftazidime (CAZ) binding and resistance, even though they are distant from the active site. Through absorbance and fluorescence spectroscopy, coupled with biophysical measurements, we demonstrated that the mutations favor tighter, mostly hydrophobic interactions with CAZ, as opposed to hydrogen-bond-dominated binding in the wild-type enzyme. In silico

structural modeling also revealed that the mutations confer conformational flexibility,
enabling more rigid and stable accommodation of CAZ within the enzyme's active site. These
findings underscore the way that non-active site mutations can
indirectly contribute to antibiotic resistance by optimizing substrate binding dynamics. The
widespread therapeutic application of β-lactams likely selects and propagates such
variants since they provide a survival advantage under therapeutic pressures. This study
highlights the difficulty of resistance evolution and the need for new strategies
to inhibit mutational adaptations outside of traditional active-site-directed drug design.

Significance

This study represents a pivotal advancement in understanding the molecular evolution of
antibiotic resistance by describing how distant mutations in TEM β-lactamase variants enhance
ceftazidime (CAZ) resistance, a critical third-generation cephalosporin. Through spectroscopic
and biophysical approaches, this study describes how these mutations facilitate tighter,
hydrophobic-restricted binding to CAZ compared to hydrogen-bond-dependent binding of the
wild-type enzyme which opposes the traditional paradigm that resistance results from active-site
modifications alone.The findings emphasize an important evolutionary strategy that, mutations
far from the catalytic site can modulate the enzyme architecture to enhance substrate
specificity.This study highlights the urgent necessity for new approaches to fight against
resistance mechanisms beyond active-site targeting. By integrating structural, functional, and
evolutionary data, the study provides a road map for designing next-generation antimicrobials to
combat the relentless rise of resistance in Gram-negative bacteria.

References

Abdul Samad, F., Suliman, B.A., Basha, S.H., Manivasagam, T., Essa, M.M. (2016). A
Comprehensive In Silico Analysis on the Structural and Functional Impact of SNPs in the
Congenital Heart Defects Associated with NKX2-5 Gene—A Molecular Dynamic Simulation
Approach. *PLoS ONE*, 11, e0153999. https://doi.org/10.1371/journal.pone.0153999.

Avci, F.G., Altinisik, F.E., Vardar Ulu, D., Ozkirimli Olmez, E., Sariyar Akbulut, B. (2016). An
evolutionarily conserved allosteric site modulates beta-lactamase activity. *Journal of Enzyme*
Inhibition and Medicinal Chemistry, 31, 33–40. https://doi.org/10.1080/14756366.2016.1201813.

Bardhan, M., Chowdhury, J., Ganguly, T. (2011). Investigations on the interactions of
aurintricarboxylic acid with bovine serum albumin: Steady state/time resolved spectroscopic and
docking studies. *Journal of Photochemistry and Photobiology B: Biology*, 102, 11–19.
https://doi.org/10.1016/j.jphotobiol.2010.08.011.

Bertucci, C., Pistolozzi, M., De Simone, A. (2010). Circular dichroism in drug discovery and
development: an abridged review. *Anal Bioanal Chem*, 398, 155–166.
https://doi.org/10.1007/s00216-010-3959-2.

Cantón, R., Novais, A., Valverde, A., Machado, E., Peixe, L., Baquero, F., Coque, T.M. (2008).
Prevalence and spread of extended-spectrum β-lactamase-producing Enterobacteriaceae in
Europe. *Clinical Microbiology and Infection*, 14, 144–153. https://doi.org/10.1111/j.1469-
0691.2007.01850.x.

Chakraborty, B., Mitra, P., Basu, S. (2015). Spectroscopic exploration of drug–protein
interaction: a study highlighting the dependence of the magnetic field effect on inter-radical
separation distance formed during photoinduced electron transfer. *RSC Advances*, 5(99), 81533–
81545. https://doi.org/10.1039/C5RA13575C.

Chib, R., Butler, S., Raut, S., Shah, S., Borejdo, J., Gryczynski, Z., Gryczynski, I. (2015). Effect
of quencher, denaturants, temperature and pH on the fluorescent properties of BSA protected
gold nanoclusters. *Journal of Luminescence*, 168, 62–68.
https://doi.org/10.1016/j.jlumin.2015.07.030.

Chinemerem Nwobodo, D., Ugwu, M.C., Oliseloke Anie, C., Al☐Ouqaili, M.T.S., Chinedu
Ikem, J., Victor Chigozie, U., Saki, M. (2022). Antibiotic resistance: The challenges and some
emerging strategies for tackling a global menace. *Clinical Laboratory Analysis*, 36.
https://doi.org/10.1002/jcla.24655.

Díaz, N., Sordo, T.L., Merz, K.M., Suárez, D. (2003). Insights into the Acylation Mechanism of
Class A β-Lactamases from Molecular Dynamics Simulations of the TEM-1 Enzyme Complexed
with Benzylpenicillin. *J. Am. Chem. Soc.*, 125, 672–684. https://doi.org/10.1021/ja027704o.

Drawz, S.M., Bonomo, R.A. (2010). Three Decades of β-Lactamase Inhibitors. *Clin Microbiol*
Rev, 23, 160–201. https://doi.org/10.1128/CMR.00037-09.

Fantini, M., Lisi, S., De Los Rios, P., Cattaneo, A., Pastore, A. (2020). Protein Structural
Information and Evolutionary Landscape by In Vitro Evolution. *Molecular Biology and*
Evolution, 37, 1179–1192. https://doi.org/10.1093/molbev/msz256.

Galdadas, I., Qu, S., Oliveira, A.S.F., Olehnovics, E., Mack, A.R., Mojica, M.F., Agarwal, P.K.,
Tooke, C.L., Gervasio, F.L., Spencer, J., Bonomo, R.A., Mulholland, A.J., Haider, S. (2021).
Allosteric communication in class A β-lactamases occurs via cooperative coupling of loop
dynamics. *ELife*, 10, e66567. https://doi.org/10.7554/eLife.66567.

Ghisaidoobe, A., Chung, S. (2014). Intrinsic Tryptophan Fluorescence in the Detection and
Analysis of Proteins: A Focus on Förster Resonance Energy Transfer Techniques. *IJMS*, 15,
22518–22538. https://doi.org/10.3390/ijms151222518.

Gill, M., McCully, M.E. (2019). Molecular dynamics simulations suggest stabilizing mutations
in a de novo designed α/β protein. *Protein Engineering, Design and Selection*, 32, 317–329.
https://doi.org/10.1093/protein/gzaa005.

Kumar, D., Singh, A.K., Ali, M.R., Chander, Y. (2014). Antimicrobial Susceptibility Profile of
Extended Spectrum β-Lactamase (ESBL) Producing *Escherichia coli* from Various Clinical
Samples. *Infect Dis (Auckl)*, 7, IDRT.S13820. https://doi.org/10.4137/IDRT.S13820.

Kumar, V., Lakshman, P.K., Prasad, T.K., Manjunath, K., Bairy, S., Vasu, A.S., Ganavi, B., Jasti,
S., Kamariah, N. (2024). Target-based drug discovery: Applications of fluorescence techniques
in high throughput and fragment-based screening. *Heliyon*, 10(1).
https://doi.org/10.1016/j.heliyon.2023.e23864.

Lee, M.M., Peterson, B.R. (2016). Quantification of small molecule–protein interactions using
FRET between tryptophan and the pacific blue fluorophore. *ACS Omega*, 1(6), 1266–1276.
https://doi.org/10.1021/acsomega.6b00356.

Mátyus, L., Szöllősi, J., Jenei, A. (2006). Steady-state fluorescence quenching applications for
studying protein structure and dynamics. *Journal of Photochemistry and Photobiology B:*
Biology, 83, 223–236. https://doi.org/10.1016/j.jphotobiol.2005.12.017.

Mandal, P., Bardhan, M., Ganguly, T. (2012). Spectroscopic investigations to reveal the nature
of interactions between the haem protein myoglobin and the dye rhodamine 6G. *Luminescence*,
27(4), 285–291. https://doi.org/10.1002/bio.1348.

Miani, A., Raugei, S., Carloni, P., Helfand, M.S. (2007). Structure and Raman Spectrum of
Clavulanic Acid in Aqueous Solution. *J. Phys. Chem. B*, 111, 2621–2630.
https://doi.org/10.1021/jp066135u

Mishra, P.P., Patel, S., Datta, A. (2006). Effect of Increased Hydrophobicity on the Binding of
Two Model Amphiphilic Chlorin Drugs for Photodynamic Therapy with Blood Plasma and Its
Components. *J. Phys. Chem. B*, 110, 21238–21244. https://doi.org/10.1021/jp0615858.

Möller, M., Denicola, A. (2002). Protein tryptophan accessibility studied by fluorescence quenching. *Biochem. Mol. Biol. Educ.*, 30, 175–178. https://doi.org/10.1002/bmb.2002.494030030035.

Mukherjee, S.K., Mandal, R.S., Das, S., Mukherjee, M. (2018). Effect of non-β-lactams on stable variants of inhibitor-resistant TEM β-lactamase in uropathogenic *Escherichia coli*: implication for alternative therapy. *J Appl Microbiol*, 124, 667–681. https://doi.org/10.1111/jam.13671.

Mukherjee, S.K., Mukherjee, M., Mishra, P.P. (2021). Impact of Mutation on the Structural Stability and the Conformational Landscape of Inhibitor-Resistant TEM β-Lactamase: A High-Performance Molecular Dynamics Simulation Study. *J. Phys. Chem. B*, 125, 11188–11196. https://doi.org/10.1021/acs.jpcb.1c05988.

Pezzotti, G. (2021). Raman spectroscopy in cell biology and microbiology. *J Raman Spectroscopy*, 52, 2348–2443. https://doi.org/10.1002/jrs.6204.

Pimenta, A.C., Fernandes, R., Moreira, I.S. (2014). Evolution of drug resistance: insight on TEM β-lactamases structure and activity and β-lactam antibiotics. *Mini Reviews in Medicinal Chemistry*, 14(2), 111–122. https://doi.org/10.2174/1389557514666140123145809.

Ross, P.D., Subramanian, S. (1981). Thermodynamics of protein association reactions: forces contributing to stability. *Biochemistry*, 20, 3096–3102. https://doi.org/10.1021/bi00514a017.

Salverda, M.L.M., De Visser, J.A.G.M., Barlow, M. (2010). Natural evolution of TEM-1 β-lactamase: experimental reconstruction and clinical relevance. *FEMS Microbiol Rev*, 34, 1015–1036. https://doi.org/10.1111/j.1574-6976.2010.00222.x.

Singh, M.K., Dominy, B.N. (2012). The Evolution of Cefotaximase Activity in the TEM β-Lactamase. *Journal of Molecular Biology*, 415, 205–220. https://doi.org/10.1016/j.jmb.2011.10.041.

Stojanoski, V., Chow, D.-C., Hu, L., Sankaran, B., Gilbert, H.F., Prasad, B.V.V., Palzkill, T. (2015). A Triple Mutant in the Ω-loop of TEM-1 β-Lactamase Changes the Substrate Profile via a Large Conformational Change and an Altered General Base for Catalysis. *Journal of Biological Chemistry*, 290, 10382–10394. https://doi.org/10.1074/jbc.M114.633438.

Swain, B.C., Mukherjee, S.K., Rout, J., Sakshi, Mishra, P.P., Mukherjee, M., Tripathy, U. (2020). A spectroscopic and computational intervention of interaction of lysozyme with 6-mercaptopurine. *Analytical and Bioanalytical Chemistry*, 412, 2565–2577. https://doi.org/10.1007/s00216-020-02483-1.

Topală, T., Bodoki, A., Oprean, L., Oprean, R. (2014). Bovine serum albumin interactions with metal complexes. *Medicine and Pharmacy Reports*, 87, 215–219. https://doi.org/10.15386/cjmed-357.

Tooke, C.L., Hinchliffe, P., Bragginton, E.C., Colenso, C.K., Hirvonen, V.H.A., Takebayashi, Y.,
Spencer, J. (2019). β-Lactamases and β-Lactamase Inhibitors in the 21st Century. *Journal of*
Molecular Biology, 431, 3472–3500. https://doi.org/10.1016/j.jmb.2019.04.002.

Yang, J., Naik, N., Patel, J.S., Wylie, C.S., Gu, W., Huang, J., Ytreberg, F.M., Naik, M.T.,
Weinreich, D.M., Rubenstein, B.M. (2020). Predicting the viability of beta-lactamase: How
folding and binding free energies correlate with beta-lactamase fitness. *PLOS ONE*, 15,
e0233509. https://doi.org/10.1371/journal.pone.0233509.

Zhao, Q., Tao, J., Uppal, J.S., Peng, H., Wang, H., Le, X.C. (2019). Nucleic acid aptamers
improving fluorescence anisotropy and fluorescence polarization assays for small molecules.
TrAC Trends in Analytical Chemistry, 110, 401–409. https://doi.org/10.1016/j.trac.2018.11.018.

Fig. 1 Steady-state UV–vis absorption spectra and fluorescence emission spectra of TEM-β-lactamase (A,E) and

mutant β-lactamases 184M (B,F), M203 (C,G), M210 (D,H) at 5 μM concentration of the respective protein in

solution in presence of increasing concentration of CAZ (0-20 μM). Inset in (A) represented steady-state UV–vis

absorption spectra of CAZ at 20 μM at 298 K.

Fig. 1.

571   Fig.2 Stern-Volmer plot and Modified Stern-Volmer plot of Fluorescence quenching of TEM-β lactamase (wild type

572   protein) and mutant β lactamases (M184, M203, M210) at 5 µM concentration in the presence of CAZ (0-20 µM) at

573   298 K (A,E), 303K (B,F), 308K (C,G) and 313K (D,H). Nonlinear fitting analysis was performed by plotting F0/F

574   against [Q] and Log F0-F/F1 against Log{Q} respectively.

Fig.2.

Fig.3 Fluorescence lifetime (τ) of TEM-β lactamase (wild type protein) in comparison with mutant β lactamases (M184, M203 and M210) at 5 μM concentration in the presence of CAZ (0-20 μM) at temperature 298K (A) and Fluorescence anisotropy (r) plot of CAZ at 5 μM concentration in presence of increasing concentrations (5-20 μM) of the β lactamases (TEM 1, M184, M203, M210) (B).

Fig. 3.

598   Fig. 4 Van't Hoff plot for the binding of TEM-β lactamases (wild type protein) (A) in comparison with mutant β

599   lactamases M184(B), M203(C), M210(D) at 5 µM concentration in the presence of CAZ (0-20µM) concentration at

600   different temperatures 298K, 303K, 308K and 311K.

Fig. 4.

Fig. 5 Cα backbone RMSD, Cα all residue RMSD, RMSF for Cα backbone, RMSF for Cα side chain and Rg for
backbone Cα for 100 ns simulation time frame shown for the wild type TEM-1 (1ZG4) interacting with CAZ;
displayed in black, in comparison with mutant proteins M184 (A,D,G,J,M), M203 (B,E,H,K,N), and M210
(C,F,I,L,O); displayed in red respectively. Projection of SSEs throughout the MD simulation trajectory in presence
of CAZ for wild type TEM-1 (IZG4) (P), M184 (Q), M203 (R), M210 (S). SSE composition is represented by
different color codes: helices; pink, turns; green, 3-10 helix; blue, β-strands; yellow, coils; white.

Fig. 5.

Table 1:

TCSPC life time fluorescence decay profile of (A) TEM-1(wild type protein) and mutant β-lactamases (B) M184, (C) M203 and (D) M210 at 0.5 mM concentration in the absence and presence of Ceftazidime (CAZ).

A

Sample	λ_{em}	λ_{ex}	α_1	T_1	α_2	T_2	Avg	X^2
TEM-1	295	340	32.17	1.90	67.83	5.94	**5.40**	1.08
5μM CAZ			30.82	1.69	69.18	5.87	**5.39**	1.08
10μM CAZ			30.39	1.62	69.61	5.83	**5.37**	1.07
15μM CAZ			30.93	1.70	67.07	5.95	**5.33**	1.03
20μM CAZ			33.96	1.43	66.04	5.69	**5.20**	1.004

B

Sample	λ_{em}	λ_{ex}	α_1	T_1	α_2	T_2	Avg	X^2
M184	295	340	44.79	1.32	55.21	4.94	**4.29**	1.08
5μM CAZ			36.29	1.14	63.71	4.45	**4.02**	1.2008
10μM CAZ			38.58	0.851	66.52	3.94	**3.73**	1.16
15μM CAZ			34.56	0.881	65.44	3.97	**3.64**	1.207
20μM CAZ			52.23	0.718	47.77	3.99	**3.45**	1.26

C

Sample	λ_{em}	λ_{ex}	α_1	T_1	α_2	T_2	Avg	X^2
M203	295	340	34.57	1.60	65.43	5.24	**4.73**	1.14
5μM CAZ			32.89	1.46	67.11	5.06	**4.61**	1.09
10μM CAZ			30.71	1.37	69.29	4.86	**4.47**	1.004
15μM CAZ			25.99	0.809	74.01	4.16	**3.94**	1.29
20μM CAZ			33.36	0.456	66.64	3.63	**3.44**	1.23

D

Sample	λ_{em}	λ_{ex}	α_1	T_1	α_2	T_2	Avg	X^2
M210	295	340	31.38	1.57	68.62	5.26	**4.81**	1.005
5μM CAZ			29.44	1.50	70.56	5.11	**4.7**	1.12
10μM CAZ			30.19	1.35	71.81	4.88	**4.63**	1.03
15μM CAZ			29.00	1.31	71.00	4.93	**4.57**	1.01
20μM CAZ			27.61	0.870	72.39	4.40	**4.15**	1.16

Table 2:

Quantum yield profile of TEM-1(wild type protein) and mutant β-lactamases (M184, M203 andM210)at different concentration in the absence and presence of Ceftazidime (CAZ).

[CAZ]M	[β-lactamases]M		OD	Area	$\phi_{rel} = \dfrac{Area}{1-10^{-OD}}$	ϕ_{Abl}	$K_r = \dfrac{\phi}{\tau}$	$K_{nr} = \dfrac{1-\phi}{\tau}$
~5 $\Box 10^{-6}$		0	0.10925	0.226	0.009826	0.08	0.014	0.170
	TEM-1	~0.5 $\Box 10^{-5}$ M		0.483	0.02100	0.17	0.031	0.153
		~1.0 $\Box 10^{-5}$ M		0.548	0.02382	0.20	0.037	0.148
		~1.5 $\Box 10^{-5}$ M		0.655	0.02847	0.24	0.045	0.142
		~2.0 $\Box 10^{-5}$ M		0.740	0.03217	0.27	0.051	0.140
		0	0.10566	0.226	0.009826	0.08	0.018	0.214
	M184	~0.5 $\Box 10^{-5}$ M		0.651	0.02830	0.23	0.057	0.191
		~1.0 $\Box 10^{-5}$ M		0.752	0.03269	0.28	0.075	0.193
		~1.5 $\Box 10^{-5}$ M		0.856	0.03721	0.31	0.085	0.189
		~2.0 $\Box 10^{-5}$ M		0.979	0.04256	0.36	0.104	0.185
		0	0.1093	0.226	0.009826	0.08	0.017	0.194
	M203	~0.5 $\Box 10^{-5}$ M		0.648	0.02817	0.23	0.049	0.167
		~1.0 $\Box 10^{-5}$ M		0.755	0.03282	0.27	0.060	0.163
		~1.5 $\Box 10^{-5}$ M		0.840	0.03652	0.31	0.078	0.175
		~2.0 $\Box 10^{-5}$ M		0.988	0.04295	0.36	0.104	0.186
		0	0.1093	0.226	0.009826	0.08	0.016	0.191
	M210	~0.5 $\Box 10^{-5}$		0.701	0.03047	0.25	0.053	0.159

M					
~1.0 $\times 10^{-5}$ M	0.827	0.03595	0.30	0.064	0.151
~1.5 $\times 10^{-5}$ M	0.946	0.04113	0.34	0.074	0.144
~2.0 $\times 10^{-5}$ M	1.023	0.04447	0.37	0.089	0.151

Table 3:

Binding Parameters and Thermodynamic Parameters for the Binding of TEM-1(wild type protein) and mutant β-lactamases (M184, M203 and M210) with Ceftazidime (CAZ) at Different temperature.

PROTEIN	Temparature	ΔG^0(kJ mol^{-1}K^{-1})	ΔH^0 (kJ mol^{-1})	ΔS^0 (J mol^{-1} K^{-1})	Pattern of Interaction
TEM-1	298K	-265.59378	-591.644238	-10.9412905	V .W. Interaction and H-bond
	303K	-260.12313			
	308K	-254.652491			
	311K	-251.370103			
M184	298K	-284.81196	103.344092	44.2366739	Hydrophobic
	303K	-306.93029			
	308K	-329.04863			
	311K	-342.31963			
M203	298K	-323.56295	164.577702	66.0852341	Hydrophobic
	303K	-356.60557			
	308K	-389.64819			
	311K	-409.47376			
M210	298K	-276.32427	183.413275	70.8207056	Hydrophobic
	303K	-311.73462			
	308K	-347.14498			
	311K	-368.39119			

Table 4:

Variation in a-helix% of TEM-1(wild type protein) and mutant β-lactamases (M184, M203 and M210) with increase concentration of Ceftazidime (CAZ).

[β-lactamases]M	[CAZ]M	α – helix (%)
TEM-1(~5 $\times 10^{-6}$ M)	0	59.24
	~0.5 $\times 10^{-5}$ M	41
	~1.0 $\times 10^{-5}$ M	27.17
	~1.5 $\times 10^{-5}$ M	26.72
	~2.0 $\times 10^{-5}$ M	20.72
M184(~5 $\times 10^{-6}$ M)	0	44.10
	~0.5 $\times 10^{-5}$ M	42.55
	~1.0 $\times 10^{-5}$ M	34.17
	~1.5 $\times 10^{-5}$ M	33.27
	~2.0 $\times 10^{-5}$ M	14.06
M203(~5 $\times 10^{-6}$ M)	0	69.44
	~0.5 $\times 10^{-5}$ M	55.75
	~1.0 $\times 10^{-5}$ M	49.51
	~1.5 $\times 10^{-5}$ M	30.06
	~2.0 $\times 10^{-5}$ M	10.34
M210(~5 $\times 10^{-6}$ M)	0	79.10
	~0.5 $\times 10^{-5}$ M	76.75
	~1.0 $\times 10^{-5}$ M	53.31
	~1.5 $\times 10^{-5}$ M	40.72
	~2.0 $\times 10^{-5}$ M	33.68

657