Cigarette smoke sets up a pro-inflammatory circuit in the lung that induces the

hyper-activation of autoreactive T helper cells

Nuria Alvarez-Sanchez[a,b,c,1,*], Annette Haughian[a,*], Brendan Cordeiro[a,b,2,*], Deeva Uthyakumar[b,c], Elvira

Paneda[d], Thomas A. Heaney[b,c], Molly Pitkethly[a], Jhenifer Gabriela Vasquez Rojas[d], Emily Pullen[c], Elyse

Latreille[a], Chao Wang[b,c], Warren L. Lee[a], Martin Stampfli[e], Clinton Robbins[b,d], Shannon E. Dunn[b,c,¶]

[a]Keenan Research Center of St. Michael's Hospital, 209 Victoria Street, Toronto, ON, Canada, M5B
1X3

[b]Department of Immunology, University of Toronto, Medical Sciences Building, Rm 7207, 1 King's
College Circle, Toronto, ON, Canada, M5S 1A8.

[c]Sunnybrook Research Institute, 2075 Bayview Avenue, North York, ON, Canada, M4N 3M5.

[d]Toronto General Hospital Research Institute, 101 College St., Division of Advanced Diagnostics,
Room 3-906, Toronto, Ontario Canada M5G 1L7.

[e]Department of Pathology and Molecular Medicine, McMaster University, Hamilton, ON, Canada, L8S
4K1

Present address:

[1]Cell and Systems Biology, The Hospital for Sick Children (SickKids) Research Institute, Toronto,
ON, Canada

[2]Princess Margaret Cancer Research Tower, MaRS Discovery District, 101 College Street, East Tower,
3rd floor, Toronto, ON, M5G 1L7.
21

22 *co-first authors

23 ¶Corresponding author: Shannon E. Dunn, shannon.dunn@sri.utoronto.ca.
24

Conflict of interest: The authors have declared that no conflict of interest exists.

Abstract

Cigarette smoke (CS) exposure increases the risk of multiple sclerosis; however, the mechanisms are

unclear. To investigate this, we exposed C57BL6/J mice to CS or ambient air (AA) for 8 weeks using a

protocol modelling human chronic CS exposure and studied the impact of these exposures on the

development of experimental autoimmune encephalomyelitis (EAE). We found that CS increased the

incidence of neurological signs in 2D2 myelin oligodendrocyte glycoprotein (MOG) T cell receptor

transgenic mice, whereas in active EAE induced by MOG peptide and Complete Freund's adjuvant, CS

delayed the onset of disease. In both cases, the effects of CS tended to be greater in the males who also

developed greater leukocyte infiltration and expression of IL-12p40 and other cytokines. To gain insights

into the paradoxical effects of CS in these EAE models, we transferred congenically-marked pMOG-

reactive T helper cells into AA- or CS-exposed mice and examined the phenotype of these cells in the

lungs, spleen, and spinal cord. We found that CS-exposed lungs acted as a sink for the pMOG-reactive

T cells, in that more of these T cells homed to the lungs instead of the spinal cord. However, these CS-

educated pMOG Th cells exhibited a hyperactivated Th phenotype with greater expression of activation

markers, IL-17, and GM-CSF, a cytokine that is essential for EAE development. Intranasal treatment

with anti-IL-12p40 negated these effects of CS on the expression of pro-inflammatory cytokines by

pMOG Th cells. These studies suggest that CS sets up a pro-inflammatory circuit in the lung that

amplifies the encephalitogenic properties of myelin-specific T effector cells.

Key words: multiple sclerosis; experimental autoimmune encephalomyelitis; cigarette smoking; lungs;

T cells; Th1 cells; interleukin (IL)-12-p40; autoimmunity

Highlights:

• CS exposure enhances the development of spontaneous neuroimmunity.

• Myelin-reactive T helper 1 responses are enhanced by CS exposure.

• CS-exposed lung attracts myelin-specific T helper cells to this site.

• CS enhances myelin-specific Th1/Th17 cytokine production and GM-CSF through lung IL-12p40

1. Introduction

More than a billion people worldwide smoke cigarettes [1]. This habit is a major cause of preventable death [1]. Though tobacco consumption has declined in recent decades [2], consumption rose during the 20th century [3], coinciding with the rising prevalence of autoimmune diseases [4, 5]. In this regard, CS exposure has been linked to an increased incidence of MS, rheumatoid arthritis (RA), systemic lupus erythematosus (SLE), Crohn's disease, and psoriasis [6]. It has also been estimated that a third of RA cases [7], and 20% of MS cases [8] are preventable if people stopped smoking. For MS, smoking increases disease risk in a dose-dependent manner (odds ratio between 1.4-2.0), with a tendency for a greater effect in males than in females [9]. This effect of CS on the risk of MS is sustained even after 5 years after quitting smoking [9]. In addition, CS hastens disability accrual in people with established MS (pwMS), which is associated with greater disease activity and brain atrophy [10-12]. Each year of smoking post-MS diagnosis is associated with a five percent accelerated time to conversion to secondary progressive MS (SPMS), a stage of disease characterized by a steady accumulation of disability [13]. In contrast, quitting smoking delays onset of SPMS [13, 14]. Despite this strong connection between CS and MS, the biological mechanisms of how CS promotes myelin-specific autoimmunity remain unknown.

Clues to these potential mechanisms have been provided by epidemiology studies that studied the effects of different modes of tobacco consumption on MS risk in humans. While CS exposure promotes MS, consumption of Swedish snuff, a tobacco product that is absorbed through the oral mucosa, protects against this disease [9]. Switching from smoking to snuff also slows the onset of the progressive phase of MS [13]. These data suggest that it isn't the tobacco components *per se*, but rather the inhalation of combusted smoke particles into the lung that is the risk factor for the disease. Studies have also found an interaction of CS exposure with carriage of the disease-associated MHC-II risk alleles in the development of MS and RA [15-17], suggesting that the smoke particles are somehow promoting the activity of

autoreactive T helper cells in these diseases. Consistent with this idea, a study that examined the

phenotype of polyclonal T cells in the broncheolar lavage (BAL) of smokers and non-smokers with and

without MS noted that T cells showed a hyperactive phenotype in the BAL of smokers [18]. In addition,

SLE and RA, smoking promotes autoantibody generation [19-21]. In experimental models of RA, it is

also reported that CS promotes protein citrullination in the lung, a post-translational modification where

arginine is converted to citrulline [22]. Citrullinated proteins generate cryptic epitopes that serve as new

autoantigens that stimulate T cells in people with RA [23]. While protein citrullination is also increased

in the brains of people with MS (pwMS), this is not associated with increased T cell reactivity to

citrullinated antigens [24], suggesting that CS promotes MS-related autoimmunity by another

mechanism. Altogether, mounting evidence suggests that lung pathology caused by the inhalation of

tobacco smoke particles promotes T cell autoimmunity; however, these mechanisms have not yet been

elucidated.

To gain insights into these mechanisms in MS, we passively exposed male and female mice to

CS for 8 weeks; a regimen previously reported to model chronic CS exposure in humans [25]. We then

studied the impact of CS on myelin-specific autoimmunity in active (MOG p35-55/CFA-induced) and

spontaneous (2D2 MOG T cell receptor transgenic mice) EAE models of MS. We found that CS exposure

enhanced leukocyte infiltration and inflammation in the lung and that this was associated with greater

IFN-γ production by myelin-specific T helper cells in the peripheral lymphoid organs in the active model

of EAE; however, EAE onset was puzzlingly delayed in the male mice and was not affected in the female

mice. In contrast, in 2D2 MOG-specific T cell receptor transgenic mice, CS promoted the development

spontaneous neuroimmunity characterized by leukocyte infiltration in the dorsal root ganglion and spinal

nerve roots. To gain insights into these paradoxical findings, we tracked congenically-marked pMOG-

reactive Th17 cells after transfer into CS- or AA-exposed mice. We found in both males and females that

CS exposure caused a greater number of the pMOG-reactive Th17 cells to migrate to the lung, which

associated with a delayed onset of EAE. In the CS-exposed lungs, the pMOG-reactive Th cells acquired

a hyper-activated phenotype and greater expression of IL-17 and GM-CSF. These pMOG-reactive Th

cells in CS-exposed retained this greater expression of IL-17 and GM-CSF after reaching the CNS. Lung

DCs from CS-exposed mice also had a greater capacity to activate and promote cytokine production by

2D2 cells *in vitro*, an effect that was fully- or partially-prevented by neutralizing IL-12p40. Together,

these studies suggest a mechanism whereby CS promotes autoimmunity by recruiting antigen-

experienced autoreactive Th cells into the lung, where these cells are activated by a mechanism of

bystander activation.

2. Results

2.1 Chronic CS exposure induces lung inflammation

To investigate the mechanisms linking CS exposure to the promotion of myelin-specific autoimmunity,

we employed a whole-body CS exposure system and 8-week CS exposure protocol that has been

previously described to model the lung pathology and inflammatory signatures seen in human smokers

[25, 26]. Histological studies conducted on lungs from males exposed to CS- confirmed the presence of

greater leukocyte infiltration in CS- versus AA-exposed mice (Fig. S1A-C). $CD45^+$ leukocytes were

detected throughout the parenchyma and within immune microclusters surrounding the bronchioles and

blood vessels (Fig. S1A-C). To gain insights into the composition of lung immune cell infiltrates with

CS, we also conducted flow cytometry analysis on lung mononuclear cells (Fig. S1D-G). This analysis

confirmed greater $CD45^+$ leukocyte infiltration in the lungs of males with CS-exposure (Fig. S1D-E).

The numbers of eosinophils and conventional DCs were increased in both sexes with CS, whereas T and

B cells, alveolar and interstitial macrophages were significantly increased only in the males (Fig. 1G).

We also immunolocalized lymphocytes in lung sections in male CS- and AA-exposed mice. We found

that B cells were rare, and T cells were sparsely scattered throughout the lungs of AA-exposed mice. In

contrast, in CS-exposed mice, T cells were detected in para-bronchiole immune micro-clusters,

sometimes in close-proximity to B cells (Fig. S2). Taken together, these data suggested that this CS-

protocol enhanced lung inflammation, especially in the male mice.

2.2 Chronic CS exposure enhances myelin-specific Th1 responses and the incidence of spontaneous

autoimmunity in 2D2 mice, but delays the onset of active EAE in males.

We next examined the effect of this CS exposure regimen on myelin-specific Th cell responses during

EAE induced by immunization with MOG_{35-55}/Complete Freund's adjuvant (CFA). Mononuclear cells

were prepared from the draining lymph nodes and spleens of mice at 9 days post-immunization, and the

MOG_{35-55}-elicited cytokine and proliferation response of these cells was examined *ex vivo*. We observed

that CS induced a prominent increase in the pMOG-elicited IFN-γ secretion by spleen and lymph node

cells of male mice as compared to AA-exposed controls (Fig. 1A-B). pMOG-elicited TNF production

was also greater in the spleen with CS, and IL-13 was elevated in the lymph nodes (Fig. 1A-B). In

contrast, the levels of IL-17 and other Th2 cytokines were not significantly altered (Fig. 1A-B). The

increased pro-inflammatory MOG_{35-55}-Th1 response in the lymphoid organs did not relate to there being

a greater abundance of MOG_{35-55}-responding $CD4^+$ T cells (Fig. 1C-E). Greater MOG_{35-55}-specific Th1

cytokine production was also seen in the spleen, but not in lymph nodes, when similar studies were done

in female CC- and AA-exposed mice (Supplementary Fig. 3A-B). We next repeated these studies but

also injected mice with pertussis toxin and followed mice for the development of clinical signs of EAE

(Fig. 2A). Since Th1 cells can be pathogenic in EAE [27], we had anticipated that EAE would be more

severe with CS exposure. Instead, this treatment delayed and lessened the severity of EAE in the males

and had no significant effect on clinical scores in the females (Fig. 2B-E). Histological studies conducted

at end-point confirmed lessened CNS inflammation and demyelination in CS-exposed male mice (Fig.

2F-G). This effect of CS in the males was seen regardless of whether CS exposures were continued (Fig.

2A) or stopped prior to EAE induction (Supplementary Fig. 4).

To investigate whether CS could have an impact on the incidence of EAE, we also tested the

effect of CS- and AA-exposures in 2D2 MOG-specific T cell receptor (TCR) transgenic mice (Fig. 2A),

which can develop spontaneous EAE when housed under specific-pathogen-free conditions [28, 29]. We

found that classic EAE signs rarely developed in either AA- or CS-exposed mice; however, with CS,

male mice exhibited an greater incidence of hindlimb clasping upon tail elevation (Fig. 2H). Previously,

we reported that this neurological reflex develops in 2D2 mice in response to mild T cell infiltration in

the spinal nerve roots and dorsal spinal cord [30]. To confirm this, we harvested the spinal columns and

spinal cord from representative mice that developed clinical signs and from mice that remained

asymptomatic, and embedded these in paraffin, and stained spinal cord/column sections for CD45. We

observed only a small number of CD45$^+$ cells in the dorsal white matter and spinal nerve roots of the CS-

exposed 2D2 mice (Fig. 2I). The more prominent pathological finding was that 50% of dorsal root

ganglions (DRGs) of hindlimb clasping CS-exposed 2D2 mice were infiltrated with CD45$^+$ cells (Fig.

2I). In AA-exposed mice, no infiltrating cells were observed in the spinal cord or nerve roots and a

significantly lower fraction of DRGs were infiltrated (12%, Chi-square=38.1, p<0.05). Since the DRG

does not express MOG and is part of the peripheral nervous system, we suspect that this clinical

phenotype is driven by autoimmunity against neurofilament-medium, a protein that is strongly cross-

recognized by the 2D2 TCR CD4$^+$ T cells [30, 31]: Therefore, the signs we detected likely represents

neuroautoimmunity. Altogether, CS enhanced Th1 cytokine production by pMOG-reactive T cells upon

pMOG/CFA immunization and increased the incidence of spontaneous neuroimmunity, but

paradoxically delayed the onset of active EAE. In both cases, the effects were more prominent in the

male mice.

2.3. CS attracts myelin-specific Th Cells to the lung where they acquire increased pro-

inflammatory characteristics

The results in the active EAE model suggested that the MOG_{35-55}-reactive T cells were being primed

appropriately, but that CS was somehow preventing their trafficking to the CNS. Past studies

demonstrated that myelin-reactive Th cells travel through the bloodstream to the lung and bronchiole-

associated lymphoid (BALT), where they reside for several days before continuing to the CNS [32]. To

see whether the myelin-reactive Th cells were homing to a greater extent to the lungs with CS-exposure,

we employed a passive transfer model of EAE where we transferred CD45.1 congenic female MOG p35-

55-reactive cells into male CS- or AA-exposed hosts. The T cells were polarized with IL-23, which is

necessary to make these cells encephalitogenic by promoting a pathogenic Th17 phenotype. Similar to

findings in active EAE, CS-exposed mice initially showed a delay in disease onset; however, eventually,

EAE scores became equivalent between the two groups (Fig. 3A). Flow cytometry on CNS mononuclear

cells performed at endpoint revealed a similar accumulation of total leukocytes in CS- and AA-exposed

mice (Fig. 3B shows representative gating, results in Fig. 3C); however, less than half the amount of

$CD45.1^+CD4^+$ cells made it to the CNS in the CS- vs. AA-exposed mice (Fig. 3D). When we examined

cytokine expression within the transferred $CD4^+$ T cells in the CNS, we found that a greater fraction of

the cells in the CS group expressed IL-17 or co-produced IL-17 and GM-CSF (Fig. 3E-F). This result

suggests that the T cells became more highly encephalitogenic in the CS host environment.

To investigate the cause of the delayed passive transfer EAE in the males, we repeated

experiments, except harvested the lungs and spleens from mice at a time when clinical signs were delayed

in the CS group (e.g. *black arrow*, in Fig. 3A). These experiments revealed a 5-10-fold higher number

of MOG_{35-55}-specific $IL-17^+$ and $IL-17^+IFN-\gamma^+$ $CD4^+$ T cells in lungs of CS relative to the AA-exposed

mice (Fig. 3G-H). In contrast, we detected no differences between groups in the spleen (Fig. S5A). In

addition, we detected an overall increase in $IFN-\gamma^+$ by polyclonal $CD4^+$ T cells in the lung, in the absence

of pMOG, consistent with the idea that CS-exposed lung environment has Th1 promoting activities (Fig.

3I-J). In light of past reports that CS exposure can enhance the activation of T cells in the BAL in humans

and mice [18, 33], we also measured the T cell expression of activation markers as well as GM-CSF, a

cytokine that was upregulated by CS exposure in the CNS-infiltrating Th cells and that is crucial for

initiating demyelination in EAE [34]. This analysis revealed greater expression of IL-17 and a massive

increase in GM-CSF (Fig. 3I). In addition, we observed the upregulated expression of activation markers

CD25, PD-1, CD69 on T cells in the CS-exposed lung (Fig. 3J). In similar experiments performed in

females, we also detected a greater recruitment of CD45.1$^+$ CD4$^+$ T cells to the lung, and these cells also

expressed greater CD25 and CD69 expression with CS (Fig. S5B-C); however, the major difference was

that the onset of EAE was delayed only by 1-2 d (Fig. S5D). In contrast to findings for the pMOG-

reactive Th cells, the expression of activation markers was not altered on the host CD45.2 CD4$^+$ T cells

in the lung (Figure S5E). Thus, while the effects of CS in increasing T cell cytokine production were

seen for both the host and myelin-reactive Th cells, only the myelin-reactive Th cells, which had recently

been primed, showed greater expression of activation markers. To investigate whether the myelin-

reactive CD45.1 cells homed to the lymphocyte clusters in the lung, we also reserved a few lungs from

the experiment in males for CD45.1 immunostaining. We detected CD45.1$^+$ cells throughout the lung

parenchyma, but also within the para-bronchiole immune clusters that are enriched in the CS-exposed

lungs (Fig. 3K). Thus, CS-exposure attracted the MOG p35-55-reactive Th effector cells into the lungs,

where these cells became increasingly activated and expressed greater GM-CSF. At the same time, this

chemotactic pull of the CS-exposed lung on the T cells seemed to slow their entry into the CNS.

2.4. CS exposure amplifies cytokine production and activation marker expression of antigen-

activated, but not naive T cells

The increased expression of activation markers on the myelin-reactive T cells in the CS-exposed lung,

but not in the spleen, suggested that these T cells were either seeing an antigen that they recognized in

the lung or were being activated in an antigen-independent manner via a process of bystander activation.

To gain insights into this, we also phenotyped the transgenic CD4$^+$ T cells in the 2D2 mice that were

exposed to CS or AA for 8 weeks. As was the case in passive EAE, CS increased the accumulation of

total leukocytes and tended to increase the numbers of 2D2 CD4$^+$ T cells in the lungs (Fig. S6A-B);

however, in contrast to the situation in passive EAE, the majority of these 2D2 Th cells had a naive status

and the expression of activation markers was not increased on the 2D2 cells in the CS lungs (Fig. 6CD-

E). This argues against the smoke particles serving as an antigen for the Th cells. However, we did detect

that the expressions of IFN-γ and IL-17A as assessed by MFI tended to be increased in the CD44^{hi}

(memory) but not the CD44^{lo} (naive) fraction of lung 2D2 CD4$^+$ T cells. These lung-isolated 2D2 cells

also produced greater levels of cytokines upon *ex vivo* stimulation with MOG p35-55 (Supplementary

Fig. 6F-G). These effects of CS were not observed in the spleen (Fig. S6G-L), confirming that these

effects of CS were mediated through the lung. Collectively, these findings suggest that the CS-induced

inflammatory environment is attracting myelin-reactive Th cells to the lung where the pro-inflammatory

milieu is inducing bystander activation of antigen-experienced Th cells. In the absence of antigen, the

CS lung promotes pro-inflammatory cytokine production by the Th cells, but does not increase their

activation state.

2.5. Alterations in human and murine lung inflammatory signatures with CS exposure

To gain further insights into what inflammatory pathways may be upregulated in the lung with smoking,

we conducted an exploratory Gene ontology (GO) enrichment analysis on an existing data set of

differentially-expressed genes (DEGs) that were found to be upregulated with CS in the lungs in humans

[35]. GO analysis revealed the upregulation of a number of immune pathways with smoking including

"regulation of the innate immune response", "toll-like receptor signaling", "antigen processing and

presentation", "chemotaxis", "leukocyte activation in immune response", and the "initiation of cytokine

programs priming the adaptive immune response" (Fig. 4A and Supplementary Table 1). In addition to

these top GO biological processes, specific cytokine pathways were enriched with CS, including "TNF",

"IL-6", "type II IFN", "IL-12", and "IL-23" production (Figure 4A, Supplementary Table 1). To validate

these findings, we also performed Molecular Signatures Database (MSigDB) Hallmark and Reactome

enrichment on the same set of DEGs, with top immune pathways displayed (Figure 4B). The top 8

immune response pathways were similar to those identified in the GO analysis and encompassed innate

immune activation, antigen presentation, and molecular signalling programs that are relevant to T cell

programming. Leading-edge genes from these enriched pathways, such as *TLR4, TLR8, IRF5, CD86*,

and *CXCL1*, overlapped with those in curated cytokine pathways (e.g., IL-12/IL-23) (see Supplementary

Table 1), suggesting upregulation of cytokine pathways by TLRs despite weak enrichment of cytokines

themselves within the DEGs.

To see how inflammatory signatures with our murine CS exposure regimen compared to these

human lung gene signatures, we profiled cytokine and chemokine protein expression in lung lysates

prepared from AA- and CS-exposed male and female mice (Fig. 4C). Consistent with findings in this

human data set and past studies using this CS-exposure regimen in BALB/C mice [33, 36-40], we

detected greater levels of several pro-inflammatory cytokines in the lung with CS, including IL-1α, IL-

12p40, IL-33, leukemia inhibitory factor (LIF), IL-7, and GM-CSF (Fig. 4C). In contrast, IL-4, IL-5, IL-

12p70, IL-23, and IL-10 were not altered (Fig. 4C) . Similar to findings in the human smokers, we

detected a trend for increased TNF and IL-6 with CS-exposure (Fig. 4C). Levels of chemoattractants for

B cells (CXCL13), eosinophils (eotaxin), monocytes/macrophages (CCL3), and T cells (CXCL10) were

also increased with CS (Fig. 4C). When comparing the effects of CS in males and females, we found that

certain inflammatory factors were upregulated more so in the males; these factors included IL-12p40,

IL-33, eotaxin, and the B cell survival factor BAFF (Fig. 4C). The expressions of IL-3 were also more

extensively downregulated in the males with CS (Fig. 4C). The greater levels of eotaxin and BAFF could

provide an explanation for the greater eosinophil and B cell with CS in males, whereas the greater

expression of IL-12p40 in the CS-exposed mice could explain the greater propensity of the T cells to secrete IFN-γ, IL-17 and GM-CSF [41]. Though IL-33 is known to activate type 2 innate lymphoid cells (ILC2s) in the lung under conditions of allergy, in the presence of IL-12, IL-33 can promote a type 1 inflammatory axis in the lungs [42]. Therefore, we suspect that this could potentially be a further contributor to the greater IFN-γ production by the T cells in CS-exposed lungs.

2.6. CS enhances the maturation status of antigen presenting cells in the lung and these are more capable of priming myelin-specific T helper cells

Given the upregulation of the "antigen processing and presentation" pathway, and the prominent increase in the expression of CD86 in the CS-enriched DEGs [35], which also featured in "lymphocyte activation" and "lymphocyte proliferation" pathways, we examined the maturation status of conventional DC (cDCs) and macrophages in the male and female CS- and AA-exposed lungs in the context of passive transfer EAE. We detected that CS increased CD86 expression on lung murine cDCs and macrophages (*black panels*, Fig. 5A). CS also increased MHC Class II expression on lung macrophages and monocytes (Fig. 5A). A similar pattern of greater CD86 expression was seen in the female cDCs and macrophages with CS (Fig. S7). To investigate whether DCs from the CS-exposed had a different functional capacity to prime MOG p35-55-reactive T cells, we magnetically enriched CD11c+ cells from lung mononuclear cells from male CS- and AA-exposed mice and co-cultured these cells with male 2D2 CD4+ T cells *in vitro* in the presence and absence of MOG p35-55 (Fig. 5B-F). We detected that a greater percentage of 2D2 CD4+ T cells were activated, divided, and became effector Th1 and Th17 CD4+ T cells when co-cultured with CS lung CD11c+ cells as compared to AA-exposed CD11c+ cells (Fig. 5B-F). Notably, MOG p35-55 had to be added in the cultures to see this effect, consistent with the idea that the CS-exposed CD11c+ cells were not providing a source of antigen to the 2D2 cells, but rather were enhancing

the activation of pre-primed Th effector cells. This suggests that increased co-stimulatory marker

expression on lung APCs could be one factor contributing to the greater T cell activation in the CS-lungs.

2.7. Lung IL-12p40 production mediates the effect of CS exposure in enhancing Th cell cytokine

production by the myelin-reactive Th cells

In light of the known effect of IL-12p40 in enhancing Th1/Th17 cytokine production and GM-

CSF expression by pMOG Th cells (via IL-12p40 homodimer, IL-23, and IL-12p70) [41, 43], we

investigated the role of this cytokine in the effect of CS-exposed DCs on T helper cells. To this end, we

repeated the DC: T cell culture studies, except also added anti-IL-12p40 or isotype control antibody to

the cultures (Fig. 6A-d). We found that anti-IL-12p40 antibody completely reduced the effect of the CS-

DCs in enhancing T cell proliferation (Fig. 6A) and expression of IL-17 (Fig. 6D), and partly inhibited

the effect of CS-DC in raising expression of the activation marker CD25 and IFN-γ (Fig. 6B-C), but had

little impact in the co-cultures containing AA-exposed DCs (Fig. 6A-D). We also conducted *in vivo*

neutralization studies in the context of passive EAE by providing CS- or AA-exposed mice two doses of

anti-IL12p40 or isotype control antibody, intranasally on two consecutive days before the transfer of the

IL-23-polarized pMOG-reactive Th cells. The phenotype and number of the transferred pMOG-specific

Th cells were examined in the lungs at 4 days post-transfer by flow cytometry (Fig. 6E shows

representative staining). Pilot studies established that intranasal instillation of IL-12p40 at the 125 μg

specifically reduced lung IL-12p40 levels (Supplementary Fig. 8). We found that intranasal

administration of this dose of anti-IL-12p40 did not reduce the total leukocyte numbers (Fig. 6F), but did

reduce the number of donor pMOG-reactive CD4$^+$ T cells (Fig. 6G) and the donor Th17 cells in the CS-

exposed lung (Fig. 6G-H). Anti-IL-12p40 treatment also lowered the percentage of CD45.1$^+$CD4$^+$ T cell

cells secreting IL-17 and GM-CSF (Fig. 6I-J), consistent with the idea that IL-12p40 is the major factor

accounting for the greater Th1/Th17 effector cytokine production in lung with CS. This treatment,

however, did not counter the effect of CS in raising CD25 expression on the T cells (Fig. 6K), suggesting

that the T cell activation is mediated by another mechanism, likely increased co-stimulation (Fig. 5).

Again, these effects of anti-IL-12p40 were only seen in the CS-exposed lungs (Fig. 6E-K). These findings

suggest that IL-12p40 in the lung contributes to both the T cell chemoattractant and the cytokine-

promoting properties of CS on the autoreactive Th cells in the lung.

3. Discussion

The mechanisms that drive MS initiation are unknown. Learning the biology of how MS risk factors

enhance autoimmunity may shed light on the pathways involved in the initiation of this disease. Cigarette

smoking is known to promote the development of MS and other autoimmune conditions, but no unifying

mechanism has been proposed. Here, we demonstrate that CS exposure promotes leukocyte infiltration

and the expression of cytokines and chemotactic factors in the lung and induces the maturation of local

lung cDCs and macrophages. This has the effect of drawing myelin-specific Th memory cells into the

lungs, where these cells express higher levels of cytokines, including GM-CSF, and become

hyperactivated. pMOG Th cells that were educated in the CS mouse also showed greater expression of

pro-inflammatory cytokines when they reached the CNS compartment. Our studies also revealed that

lung IL-12p40 is one mediator of these effects of CS in the lung. Our findings thereby suggest that CS

is not the trigger for autoimmunity, but rather amplifies the encephalitogenic potential of myelin-specific

T memory cells via a mechanism of bystander activation. In the 2D2 model, we showed that these effects

of CS tipped the balance towards autoimmunity.

We explored the effect of CS on EAE development in three EAE models: active EAE, IL-23-

induced passive T cell transfer EAE, and in 2D2 mice that can develop spontaneous EAE. In the case of

active EAE, where the presence of heat-killed tuberculosis in CFA promotes a strong Th1 response, CS

enhanced IFN-γ and TNF production by MOG p35-55 Th cells in the peripheral lymphoid organs. In

light of past studies that myelin-reactive Th cells also traffic through the lung during active EAE [32],

we speculate that MOG p35-55-reactive Th cells acquired the greater capacity to produce IFN-γ because

they had already travelled through the pulmonary circuit at the time of sampling. Greater IFN-γ

expression was also observed with CS exposure in the host CD44^{hi}CD4$^+$ cells that had been recruited to

the lung during passive transfer EAE and in the CD44^{hi}, but not naive 2D2 TCR transgenic cells in the

CS-exposed mice. In the Th17-passive EAE model, CS instead promoted greater T cell expression of IL-

17 and GM-CSF by the Th17 effectors, essentially enhancing the expression of cytokines that these cells

were programmed to make. Altogether, our findings align with past reports that CS enhances polyclonal

T cell activation in the lungs of mice [40, 44] and promotes type I inflammation in the BAL, and IFN-

γ$^+$IL-17$^+$ and CXCR3$^+$ (Th1-like) memory T cells in the blood of human smokers [45, 46]. Our finding

that CS increased the maturation of murine lung DCs also coincides with past reports that lung DCs play

a role in enhancing Th1 and Th17 responses in human smokers with emphysema [47] and in in

experimental emphysema induced by long-term CS exposure [48]. In a longer-term, 6 month CS-

exposure model, lung DCs also promote the formation of tertiary lymphoid structures in the lung, a

pathological feature of the human chronic obstructive pulmonary disease (COPD) [49]. Altogether, these

findings suggest that cDCs in the CS-exposed lung amplify cytokine production and promote the

bystander activation of the myelin-specific T cells that are attracted to this site.

We also found that neutralizing IL-12p40 in the airways inhibited the recruitment of myelin-

specific T cells and reduced the potential of these cells to make Th1 and Th17 cytokines *in vivo*. In the

more simplified CD11c$^+$: 2D2 T cell co-culture system, anti-IL-12p40 also inhibited the effect of DCs

on the MOG-elicited proliferation and activation of 2D2 CD4$^+$ T cells. Our proteomics study detected

greater IL-12p40, but not IL-12p70 and IL-23 with CS in the murine lung, suggesting that the activities

may be mediated through the activity of the IL-12p40 homodimer. Although T cell chemoattractant

activities of IL-12p40 homodimer have not yet been reported, this cytokine is chemotactic for both

dendritic cells and macrophages (reviewed in [50]) and promotes lung inflammation during viral

infection through immune cell recruitment [51]. Previously, we also showed in the CD11c$^+$: 2D2 T cell

co-culture system that IL-12p40 levels correlated with the expression of both IFN-γ and IL-17 production

by 2D2 CD4$^+$ T cells and that neutralization of the activity of the IL-12p40 homodimer reduced T cell

production of IFN-γ, whereas the IL-17-promoting activities were mediated through IL-23 [43]. Of these

IL-12 family cytokines, only IL-12p70 and IL-23 have been reported to promote GM-CSF expression by

T cells [52, 53]. Thus, it is likely that a combination of activities of the IL-12p40 homodimer, IL-23

(p19/p40), and possibly IL-12p70 are mediating these effects of CS on cytokine production by T cells.

Consistent with this idea, past studies have described roles for both IL-12p70 and IL-23 in the

development of other CS-associated lung pathologies, including fibrosis [54] and emphysema [55].

Resolving the individual contributions of IL-12p70, IL-23, and IL-12p40$_2$ will require employing mice

that are singly or doubly deficient in the protein subunits as has been done in the context of experimental

lung infection (e.g, [56]).

Finding that anti-IL-12p40 can dampen myelin-specific Th cell cytokine production aligns with

past evidence in EAE that this cytokine plays a key role in the initiation of this disease [57, 58]. Due to

promising effects in EAE models, anti-IL12/23 p40 antagonistic antibodies were tested in two phase 2

clinical trials in MS. One trial testing subcutaneous ustekinumab failed to show efficacy of the antibody

in reducing the appearance of new brain gadolinium-enhancing lesions in relapsing-remitting MS

patients [59], while another trial that tested briakinumab noted an effect in reducing gadolinium-

enhancing lesions and disease relapses[60]. This antibody was not developed further since the efficacy

was modest compared with other MS treatments and since trials in psoriasis revealed a signal for

increased risk of serious infections, nonmelanoma skin cancers, and cardiovascular events [61]. Notably,

these trials did not segregate data by sex or smoking status, making it unclear whether anti-IL-12p40

preferentially benefited smokers or males with MS. No one has yet tested the effect of intranasal delivery

of anti-IL-12p40 in MS, although other biologics including anti-CD3 are now being tested through this

route with promising results (e.g., [62]).

The protein signatures in murine lungs revealed a prominent increase in the alarmins IL-1α and

IL-33, IL-12p40, GM-CSF with CS exposure. This cytokine signature is similar to that previously

described using the same CS-exposure protocol in mice [36, 37]. These past studies showed that the

alarmin IL-1α acts as an upstream effector in the pro-inflammatory cascade in the lung and that

neutralizing this cytokine inhibits GM-CSF induction [36], placing IL-1α upstream of GM-CSF in the

inflammatory pathway. Additionally, a study focusing on GM-CSF demonstrated that neutralizing this

cytokine prevented the accumulation of dendritic cells, the activation of $CD8^+$ T cells, and the induction

of IL-12p40 in the lungs with CS exposure [37], thus positioning GM-CSF as an intermediary in the

pathway leading to IL-12p40 in CS-induced lung inflammation. Our data further revealed that IL-33 was

elevated more significantly in male lungs with CS. Although IL-33 is primarily known for it role in

promoting expansion of type 2 innate lymphoid cells (ILC2s), especially in female lungs, during states

of health and allergy [63], in the context of cigarette smoke and influenza, increased IL-33 coupled with

decreased IL-33 receptor ST2 on ILC2s instead promotes macrophage and dendritic cell activation and

type 1 immunity [42]. In the context of CS, IL-33-deficient mice are protected from type I inflammation

in the lung [42]. Similarly, in humans, the combination of IL-12 and IL-1β is reported to induce trans-

differentiation of human ILC2s to ILC1s [64]. Consequently, it is likely that IL-33 worked together with

IL-12p40, CCL3, and eotaxin to enhance lung inflammation with cigarette smoke in males.

We noted a trend for greater leukocyte infiltration in the lungs of males with CS. This sex

difference did not impact the activation status of either the DCs or of the infiltrating pMOG-specific Th

cells, since these measures increased to a similar extent in males and females with CS. However, the

greater lung inflammation correlated with comparatively greater delay in EAE in the passive EAE model,

which is likely due to a delayed exit pMOG-reactive Th cells from the lung and delayed migration to the

CNS. By contrast, this same tendency for greater lung inflammation in males favored the development

of neuroimmunity in males in the 2D2 model. The disparate effects of CS in these models can be

reconciled when considering how pMOG Th cells are primed in each disease model. In the case of active

EAE and passive EAE, pMOG Th cells were primed in the draining lymph nodes by their antigen coupled

with a strong adjuvant, and these Th cells travel in a large wave to the CNS to initiate disease. The CS

lung, though enhancing the encephalitogenic properties of these cells, also slowed their trafficking to the

CNS. However, in the case of the 2D2 model, no external source of antigen was provided, and instead,

these cells are thought to gain antigen experience upon exposure to gut commensals and microbes at

mucosal surfaces [29, 65]. In this case, the CS lung had the effect of drawing these otherwise dormant

2D2 CD44^{hi} CD4$^+$ T cells into a pro-inflammatory environment that had the effect of tipping the balance

towards autoimmunity. We expect that this model may be closer to the scenario in human MS where

autoreactive T cells are also activated by the microbiota and EBV infection at mucosal sites.

The reasons for these sex differences in lung inflammation with CS are unclear. We did not

pursue the mechanisms for the greater expression of IL-12p40 or IL-33 in the males with CS. This is

because it has already been established that these cytokines are produced more prominently in females

due to the action of sex hormones. IL-12p40 is produced at greater levels by murine female DCs and

macrophages in response to T cell-derived signals [66-68]. This sex difference has been demonstrated to

be due to enhancing effects of estradiol and repressive effects of androgens on IL-12p40 in macrophages

and DCs [66, 67]. Similarly, females have been reported to exhibit greater activity of the IL-33/ILC2

axis in the lung at steady state and during allergic inflammation; this sex difference is due to the effects

of androgens in suppressing the lung accumulation of ILC2 cells [63, 69, 70]. We therefore speculate

that the sex difference in lung production of certain factors is regulates by effects of sex on signaling

pathways upstream. In this regard, analysis of the human CS-induced DEGs identified genes involved

in TLR signaling (TLR2, TLR4, MYD88) signaling, activation of the inflammasome (PYCARD/IL18) and activation of Dectin-1-fungal sensing pathways (CLEC7, SYK, CARD9) in the lungs with CS (Supplementary Table 1). Past studies in mice established that these same pathways are upregulated in the murine lungs with CS exposure; the activation of TLR4 is mediated by HSP60 [71], while activation of the fungal sensing pathways mediated by TLR2 and Dectin-1 are in response to greater levels of Hmgb1 [72]. Future studies will explore which of these innate pathways are more operative in the male versus female lung with CS exposure.

We observed that chronic CS in mice also promoted the formation of small T cell and T cell/B cell microclusters in the lungs. These are likely a prelude to the more organized tertiary lymphoid structures reported to develop with more prolonged CS exposures leading to COPD-like features in mice [26] and humans [65]. We showed that the CD45.1[+] MOG p35-55-reactive Th cells that we transferred homed to these clusters, suggesting that these could be sites where the MOG p35-55 T cells becoming activated. Past studies showed that treatment with a BAFF-Fc reagent could prevent the development of tertiary lymphoid structures in the lungs of mice exposed to CS for 6 months [26]. Interestingly, interstitial lung disease with prominent lymphocytic infiltration has been described as a disease feature of RA [73]. Whether this also occurs in people with MS who smoke, to our knowledge, has not yet been examined; however, the risk of MS is reported to be 6-fold higher in those who subsequently develop COPD before age 60 years [74], hinting at a potential link between lung disease and MS.

Our study also brings further attention to the lung-brain axis in the regulation of MS-related autoimmunity. A connection between the lung and CNS was first suspected when epidemiology studies detected a temporal association between respiratory tract infections and MS onset [75] and disease relapses [76-81]. However, a concrete link was not envisaged until it was demonstrated in studies in rats that myelin-reactive T cells traffic through the BALT and lung draining lymph nodes on their way to the spinal cord during passive and active EAE [32]. The transit through these organs was shown to be

important for the acquisition of expression of chemokine receptors and adhesion molecules on the T cells

that are necessary for transmigration into the CNS [32]. Other studies have demonstrated that lung

commensals and microbes also modulate EAE severity. For instance, intranasal infection of mice with

PR8 influenza exacerbates active EAE by promoting a Th1 cytokine response by myelin-reactive Th

cells and induces immune infiltration into the spinal roots in 2D2 mice, similar to the pathology that we

observed with CS [65, 82]. In addition, it has been shown that depleting the lung flora with neomycin in

mice ameliorates EAE by lowering the activation state of microglia in the CNS [83]. Finally, a murine

study identified the lung as a promising site for the delivery of antigen-coated microparticles to induce

tolerance. Interestingly, part of the benefit of antigen nanoparticle therapy in attenuating EAE related to

its ability to promote mild inflammation in the lung, which caused the T cells to be retained in the lung

instead of going to the brain [84].

Study Limitations

Despite testing CS in three different EAE models and demonstrating an effect of CS in promoting

neuroimmunity, we were unable to model an effect of CS in worsening *myelin*-specific autoimmunity.

Having such a model would be important for pre-clinical testing of therapies that aim to treat lung

inflammation in pwMS who smoke. Similar to our findings, others similarly showed that CS-exposure

delayed EAE [85], and the onset of autoimmunity in the MRL-*lpr/lp*r SLE-prone mouse strain [86].

Interestingly, in the latter model, the CS had the paradoxical effect of enhancing autoantibody generation.

Future studies will modulate the dose of CS and timing of CS exposures to minimize this chemotactic

activity while still promoting the pro-encephalitogenic activities of the T myelin-reactive cells.

In conclusion, our findings demonstrate that myelin-reactive Th cells traffic to the CS-exposed

lung where signals, including IL-12p40 and increase DC co-stimulation, and amplify T cell activation

and cytokine production, thereby providing a general mechanism for how CS may promote

autoimmunity.

4. Methods

4.1. Sex as a biological variable

Our study examined both male and female animals, and the phenotypes were compared between both males and females.

4.2. Mice

Male and female C57BL6/J mice (Stock#664), female CD45.1 congenic mice on the C57BL6/J background (PepBoy; B6.SJL-Ptprca Pepcb/BoyJ; Stock# 002014), male and female 2D2 TCR transgenic mice (2D2 mice; C57BL/6-Tg(Tcra2D2,Tcrb2D2)1Kuch/J; Stock#6912) were from the Jackson Laboratory. 2D2 mice were bred in pairings of heterozygote males with wild-type C57BL/6J females. 2D2 heterozygote offspring were genotyped by flow cytometry of PBMCs after staining for CD4 and TCR Vβ11 chain antibodies as described previously [43]. All mice were housed within a specific pathogen-free facility at the University Health Network (UHN), and work was conducted under an animal use protocol (AUP 5937) that was approved by the UHN animal care committee, following guidelines of the Canadian Council on Animal Care.

4.3. Exposures to CS and AA, and Cotinine Measurements

Adolescent mice were exposed to CS from research-grade cigarettes, with filters removed (Tobacco and Health Research Institute, University of Kentucky, Lexington, KY) using a whole-body exposure system (SIU48; Promech Lab Inc., Vintrie, Sweden) that was housed in a fume hood [33]. CS was puffed into a central chamber where freely moving mice were housed in compartments with cage mates, with food and water removed. Exposures were initiated at 7 weeks of age and consisted of two daily 50 min sessions of exposure to CS (separated by 3 h) generated from 12 serially burned cigarettes, 5 days/week for 8-10 weeks, and was preceded by an initial 3-day acclimatization period where mice were placed in the exposure box for increasing lengths of time. This CS exposure regimen was well tolerated. AA-exposed

controls were brought into the same room, and, at the time of CS exposures, placed into an empty cage

having similar dimensions as the caging system used for the CS-exposures, with water and food removed.

To estimate the dose of acute CS exposures, blood was drawn within 1 h of exposure and was centrifuged

(5000 rpm x 5 min) to collect serum, which was cryopreserved at -80°C. Cotinine was measured using

an ELISA kit (Origene) and was found to be 379.1 ± 32.2 ng/ml in CS-exposed mice and 0.4 ± 0.1 ng/ml

in the ambient mice, which is similar to levels reported in past studies using this CS-exposure protocol

[33] and for chronic human smokers [87].

4.4. Lung Pathology

Mice were anesthetized with isoflurane and perfused with 1x PBS containing 5 U/mL of heparin. The

lung and heart were dissected, and a 0.5 cm length of tubing (VWR 427411), was inserted into the trachea

and was tied in place using surgical twine. To prepare formalin-fixed paraffin sections, lungs were first

inflated with formalin by filling a 50 mL syringe with neutral buffered formalin, which was gravity-fed

through surgical tubing to a 23 G needle placed in the canula. After the lungs were fully inflated, the

trachea was tied off below the level of the cannula, and the heart and lung were stored in 10% formalin.

After 3-7 days, the lungs were grossed, processed and embedded in paraffin with the right lung embedded

longitudinally, and the left lung upper, middle, and lower lobes embedded in cross-section. H&E staining

and CD45 IHC (rat anti-CD45 Ab, clone 30F-11, BD) were conducted as previously described using our

established protocols [88]. Dual CD3 and B220 IHC was conducted via a similar method with the

following modifications. After de-waxing, antigen retrieval, and blocking [88], sections were stained

overnight at 4°C using a rabbit polyclonal anti-human CD3 Ab (1:1000, Dako). The following day,

sections were washed 3x in 0.1% PBS-Tween-20 (PBST), probed with secondary Ab (HRP-conjugated,

goat anti-rabbit Ab, clone BA-1000, Vector), washed 3 times, and then incubated with HRP-streptavidin

complexes (universal ABC-HRP kit, Vector labs) [88]. After three final washes, immune complexes

were developed using DAB (Vector SK4100) [88]. Sections were then re-blocked in blocking buffer (2%

BSA in 2% rabbit serum) that contained avidin (Avidin Biotin kit, Vector labs) and incubated overnight at 4°C with anti-B220 Ab (1:500, clone RA3-6B2, BD Biosciences) in blocking buffer that contained biotin. Slides were washed with PBST and incubated with biotinylated rabbit anti-rat secondary Ab (1:200, Vector labs) for 30 min at RT. After an additional wash, sections were incubated for 30 min at RT with Avidin-Alkaline Phosphatase (1:1000; BioShop) in 1x PBS. Slides were washed again with PBST, and antibody-AP complexes were developed using Vector Blue as a chromogen (Vector Labs). Slides were dehydrated and then cleared with Citrosolve and cover-slipped using Permount (both from Fisher).

For CD45.1 immunohistochemistry, the lungs and heart with attached trachea were harvested and canulated as described above, except that the lungs were inflated with 500 µL of a 2:3 solution of 1x PBS: OCT compound using a 23 G needle and syringe. The inflated lungs were tied off below the cannula and submerged in the PBS:OCT solution until all specimens were collected. After grossing the lungs, they were embedded in OCT in melting 2-methyl butane cooled by liquid nitrogen. OCT-embedded lungs were cut at 4 µm and stained using a CD45.1 Ab (1:1000, clone A20, Biolegend) and the mouse on mouse (M.O.M) kit (Vector Laboratories) according to kit directions. In brief, sections were air-dried for 10 min and then fixed in ice-cold methanol for 15 min. Slides were washed in 0.5% PBST (3 x 5 min) and peroxidase quenched with 3% hydrogen peroxide in methanol. Sections were blocked using the blocking solution provided in the M.O.M. kit for 30 min at RT, then washed with PBST. After secondary Ab detection, sections were incubated with streptavidin-HRP (universal ABC HRP kit, Vector labs) and then developed with DAB, counterstained with hematoxylin, dehydrated, cleared with xylene, and mounted with Permount™ according to our published protocols [88].

4.5. Isolation of Lung Mononuclear Cells

The lungs were dissected as above and placed in a conical tube containing ice-cold RPMI 1640 (Life

Technologies) and 2% FBS (Wisent). Lung lobes were transferred to a petri dish containing 5 mL of ice-

cold RPMI, scissor-minced and transferred to a 50 mL conical tube. Five mL of 2x digestion buffer

containing 300 U/mL type 1 collagenase (Worthington) and 10 U/mL DNAse I (Sigma) in RPMI were

added and the tissue was agitated for 1 h on a 37°C shaker. The cells were then dissociated through a 70

micron sieve and centrifuged at 300 x g for 10 min at 4°C. The supernatant was discarded and red blood

cells were lysed with ACK buffer (0.15 M NH_4Cl, 10 mM $KHCO_3$, 0.1 mM Na_2EDTA) for 1 min and

15 s [43], and the lysis stopped with RPMI. Cells were centrifuged again, counted and washed with FACS

buffer.

4.6. Flow Cytometry

Cells were incubated with Fc Block (5 µg/mL; BioLegend) for 15 min at 4°C and washed in FACS

buffer. Cells were incubated with the Ab cocktails detailed in Supplementary. Table 2 and either Fixable

Viability Dye eFluor 506 or LIVE/DEAD™ Fixable Aqua Dead Cell Stain (both 1:1000; Invitrogen) in

the dark for 30 min at 4°C, and then washed again. If intracellular staining was performed, cells were

fixed with 4% paraformaldehyde in 1x PBS for 10 min at 4°C, washed, and then incubated with 1x Perm

Wash buffer (BD Biosciences) in the dark for 30 min at 4°C. After washing cells three times with Perm

Wash buffer, cells were incubated with the intracellular antibodies detailed in Suppl. Table 2 diluted in

Perm Wash buffer for 30 min at 4°C in the dark, and washed with 1x Perm Wash buffer. Cells were

either analyzed and unmixed using a Sony SP6800 Spectral cell cytometer and software, or analyzed

using a Fortessa X-20 or a Cytoflex-LX cytometer, and data analyzed in FlowJo (version 10). Cell

populations were gated using fluorescent minus one (FMO) controls.

4.7. MOG p35-55/CFA Immunization and Active EAE Induction

Mice were immunized subcutaneously at two sites of the chest (50 µl/side) with an emulsion containing

100 µg MOG p35-55 (Genemed Synthesis) and CFA containing 200 µg of heat-killed *Mycobacterium*

tuberculosis (H37RA, Difco Laboratories) [43]. For active EAE induction, mice were also injected with

100-250 ng (lot-dependent) *Bordetella pertussis* toxin (PTX, #181, List Biologicals) on days 0 and 2

post-immunization. Mice were examined daily for clinical signs of EAE using a five-point scale: (0) no

clinical signs; (1) limp tail; (2) hindlimb or foot weakness; (3) complete hindlimb paralysis in one or both

hindlimbs; (4) hindlimb paralysis plus some forelimb weakness; (5) moribund or dead. Mice that died

from EAE were assigned a score of five for the remainder of the experiment. Cumulative score was

calculated for each mouse by adding up the daily scores over the period of observation [43]. Mice were

checked for the hindlimb clasping reflex after tail suspension and it was noted whether the hindlimbs

curled inwards (clasping) or remained splayed [30].

4.8. Measuring MOG p35-55 Reactive T Cell Recall Assays in Spleen and Draining Lymph Nodes

Mice were immunized with MOG p35-55/CFA, and at day 9 post-immunization, the recall proliferative

and cytokine response of the spleen and draining lymph nodes to MOG p35-55 were examined as

described previously [43]. In brief, mononuclear cells were isolated from the spleen and draining lymph

nodes and cultured in 96-well flat-bottomed plates (0.5×10^6 cells/well) with 0, 2, 5, and 10 µg/ml MOG

p35-55 in Complete RPMI Medium (CM): RPMI 1640, 2 mM L-glutamine, 1 mM sodium pyruvate, 0.1

mM nonessential amino acids, 100 U/mL of penicillin, 0.1 mg/ml of streptomycin (all from Life

Technologies), 50 µM 2-mercaptoethanol (BioShop), and 10% FBS. Cytokines were measured in culture

supernatants at the time of peak production (IFN-γ at 48 h, IL-17A at 72 h) using Ready-Set-Go ELISA

kits (ThermoFisher) or the LegendPlex Th cytokine panel (Biolegend). In addition, the number of MOG

p35-55 CD4$^+$ T cells that had expanded after immunization was estimated in the spleen by examining

the number of CD4$^+$ T cells that divided as assessed by CFSE dilution assay (Molecular Probes) after re-

stimulation *ex vivo* with MOG p35-55. For this assay, 10 million spleen cells were labelled with 0.5 µM

CFSE according to the kit instructions. Cells were then plated in duplicate in 24-well plates (2.5 x 10^6

cells/well) with MOG p35-55 (5 µg/mL) for 72 h in a 37°C incubator with 5% CO_2. Cells were then

harvested, washed with FACS buffer, and stained with LIVE/DEAD™ Fixable Aqua Dead Cell Stain

(1:1000) and the Ab cocktail detailed in Suppl. Table 2. The number of MOG p35-55 reactive Th cells

was estimated as the frequency of CD44^{hi}CD4$^+$ T cells that specifically divided in response to MOG p35-

55, multiplied by the total amount of live splenocytes counted at the hemacytometer.

4.9. Passive transfer EAE

Passive transfer studies were carried out similar to described previously [89]. Eight-week-old female

C57BL6/J mice or CD45.1 Pepboy mice were immunized with MOG p35-55/CFA, and mononuclear

cells were isolated from spleen and draining lymph nodes at day 9 post-immunization. Cells were plated

at 8 x 10^6/ml in 75 cm^2 vented tissue culture flasks with 10 µg/ml of MOG p35-55 and 10 ng/mL IL-23

(R&D Systems) for 3 days. Cells were then washed three times in 1x PBS and then 10-20 x 10^6 cells

(experiment-dependent) were injected into recipient AA- or CS-exposed male or female CD45.2

C57BL6/J mice. Exposures to CS and AA were continued throughout the experiment.

4.10. Histological Scoring of EAE

Brains and spinal cords from mice with active EAE were isolated, preserved in 10% neutral buffered

formalin (Sigma) for 7 days, processed in paraffin, and embedded in a single paraffin block (10–12 spinal

cord and 6 brain sections/block) according to our published protocol [88]. Cross-sections (5 µm) were

cut and stained with Luxol Fast Blue (LFB) to visualize myelin or CD45 IHC [88]. Percent demyelination

and CD45$^+$ leukocytes/mm^2 in white matter were measured using ImageJ [88].

4.11. CNS Mononuclear Cell Isolation

CNS mononuclear cells were isolated as described previously using collagenase digestion followed by

Percoll gradient[43]. For cytokine measurement, isolated CNS or lung mononuclear cells were incubated

with or without 20 μg/ml MOG p35-55 overnight (12 h) in a 37ºC incubator with 5% CO_2. GolgiStop

(BD Biosciences) was added in the last 6 h of culture. Cells were washed with FACS buffer and stained

for FACS as described above.

4.12. Analysis of a human lung tissue microarray dataset

To identify immune response pathways enriched in smoking patients, we leveraged a previously

published microarray-derived gene list identifying probe sets significantly altered between smoking and

non-smoking patients with lung cancer, obtained from Bossé *et al*. [35]. While raw data is publically

available in the Gene Expression Omnibus database (GSE23546), files were not annotated with patient

information, and so key parameters required to perform our analyses, such as sex and smoking status,

were not available to us, so analysis was performed on the published list of DEGs. All data were analyzed

on RStudio (version 4.4.0). GO pathway enrichment analysis was performed using the clusterProfiler R

package. Gene symbols were matched to corresponding Entrez Gene identifiers using the org.Hs.eg.db

annotation database. Statistical significance was determined using a hypergeometric test with Benjamini–

Hochberg correction for multiple hypothesis testing. GO terms were considered enriched at a false

discovery rate (FDR) < 0.05. Redundant GO terms were reduced by grouping highly overlapping

pathways and selecting representative immune-related biological processes for visualization.

Enrichment of MSigDB Hallmark and Reactome gene sets was assessed using the fgsea algorithm

applied to the ranked gene list. We obtained curated pathway collections from the Molecular Signatures

Database (MSigDB), using the msigdbr R package, which included Hallmark and Reactome pathway

gene sets. Hallmark gene sets represent curated, non-redundant biological processes, while Reactome

provides detailed signaling pathway annotations. Using the fgsea R package, genes were ranked

according to the differential expression statistic derived from smoking compared to non-smoking. The

enrichment score for each gene set was calculated using the running-sum statistic across the ranked gene

list. Significance was estimated using the adaptive multilevel splitting Monte-Carlo method implemented

in fgseaMultilevel, and FDRs were calculated using the Benjamini–Hochberg method. Gene sets were

considered significantly enriched at FDR < 0.05. Leading-edge genes contributing to each enriched

pathway were extracted from the fgsea results and used for downstream overlap analyses and

visualization. Classical GSEA could not be performed, given our inability to access the full dataset.

Results were visualized using dot plots generated with the ggplot2 and dplyr R packages. The heatmap

of gene expression Z-scores, across enriched GO immune pathways were generated using the pheatmap

R package.

4.13. Quantification of Cytokines and Chemokines in Lung Homogenates

Lungs were isolated from AA- or CS-exposed mice 9 days after immunization with MOG p35-55/CFA,

snap-frozen in liquid nitrogen and stored at -80°C. Lungs were weighed and homogenized in T-PER

buffer according to instructions (Thermo Scientific). Homogenates were incubated on ice for 1 h,

centrifuged for 10 min at 10,000 x g at 4°C, aliquoted and frozen at -80°C. Total protein was measured

in homogenates using a BCA Assay (Pierce). Cytokine and chemokine levels were measured in

supernatants by Eve Technologies using the Mouse Cytokine/Chemokine 32-Plex Discovery Assay

Array (MD32) panel, or with the individual ELISA kits: amphiregulin (RayBiotech), BAFF and CXCL13

(R&D Systems), IL-23 and IL-33 (Biolegend), and Verikine-HS Mouse IFN-α all subtype ELISA kit

(PBL Biosciences). Levels were expressed per mg of lung protein.

4.14. DC: T Cell Co-culture Assays

Mononuclear cells were isolated from pooled lungs of AA- and CS-exposed male C57BL/6J mice

(n=5/group) as described above, and DCs were enriched from lung mononuclear cells using the mouse

Pan Dendritic Cell Isolation Kit (Miltenyi). Isolated DCs (2.5×10^4) were cultured in triplicate in round-

bottom plates with 0 or 6 ng/mL LPS and 0-20 μg/mL MOG p35-55 for 6 h at 37°C and 5% CO_2 (pre-

activation phase). Splenic CD4$^+$ T cells were enriched from 2D2 male mice using the MojoSort Mouse

CD4 T Cell Isolation Kit (Biolegend). A portion of these cells were stained with Cell Proliferation Dye

eFluor450 according to the product directions (eBioscience). 2D2 CD4$^+$ T cells (5.0 x 10^4) were placed

on top of the cultured DCs. These cells were then cultured for 2-3 days at 37°C and 5% CO_2, and then

washed with FACS buffer. Cell proliferation was assessed by dye dilution in 2D2 CD44^{hi}CD4$^+$ T cells

using flow cytometry. For cytokine assessment, cells were re-stimulated with 50 ng/mL PMA and 500

ng/mL ionomycin for 4 h, with GolgiStop added in the last 2 h of culture, and then stained for FACS as

described above.

To assess the role of IL-12p40 signalling in T cell priming by lung DCs, DC:T cell co-cultures were set

up similarly, with the addition of 10 µg/mL of either anti-mouse IL-12p40 neutralizing Ab (clone C17.8,

BioLegend) or isotype control Ab (clone RTK2758, BioLegend) during the pre-activation phase when

DCs were incubated with LPS and MOG p35-55. After the co-culture of DCs and 2D2 CD4$^+$ T cells for

2-3 days at 37°C and 5% CO_2, cells were stained and analyzed as described above.

4.15. IL-12p40 Blockade in Lung

We tested the effect of anti-IL-12p40 Ab (clone 2A3; BioXCell) in neutralizing IL-12p40 in the lungs of

naive mice, which have detectable IL-12p40 protein at steady-state. This was done by administering 50,

125, or 200 µg/mouse of the InVivoMab anti-mouse IL-12p40 Ab (clone C17.8; BioxCell) to mice

(n=2/group) intranasally or intratracheally, and then euthanizing mice after 3 days for measurement of

IL-12p40 in lung, spleen, and small intestine homogenates using an IL-12p40 ELISA kit (Biolegend).

This study demonstrated that 125 µg of anti-IL-12p40 administered intranasally specifically lessened IL-

12p40 levels in the lung, while higher doses, or intratracheal administration, also led to systemic

depletion of the cytokine (Supplementary Fig. 8).

EAE was induced in AA or CS-exposed male recipients with pre-primed female MOG p35-55 reactive

CD45.1$^+$ cells. On days -2 and 1 post-transfer, the recipient C57BL/6J mice were anaesthetized with

isoflurane and received 125 µg/mouse of anti-IL-12p40 Ab or InVivoMAb rat IgG2a isotype control

(clone 2A3; BioXCell) intranasally. Mice were euthanized on day 4 post-transfer, and lungs were

collected for flow cytometry as described above.

4.16. Statistical Analysis

Data were analyzed using GraphPad Prism (v10). Data are showed as mean ± SEM. For studies that

contained 2 groups, a two-tailed Mann-Whitney U test was used to determine between-group differences.

For experiments with a 2 x 2 design (e.g., sex by smoking or smoking by treatment), data were analyzed

using a 2-way ANOVA followed by a Bonferroni post-hoc test. P values of p≤0.05 were considered

statistically significant.

4.16. Data availability

The data that support the findings of this study are available from the corresponding author, upon request.

5. Author contributions and Funding Support

N.A-S, A.H., B.C., and S.E.D. conceived and conducted experiments and analyzed data. N.A-S. drafted

the manuscript. D.X. conducted gene expression analysis and contributed to the writing of the

manuscript. E.P. conducted CS and AA exposures on mice. M.P. and E.P. conducted histological

analysis. J.V.R. conducted clinical and histological scoring for some EAE studies. E.L. and W.L.L helped

with lung histology studies. T.H. conducted ELISA assays. S.E.D., M.S., and C.R. obtained funding for

the project. All authors have reviewed the manuscript.

This work was funded by grants from the National MS society (M.S., C.R., S.E.D.) and MS Canada

(S.E.D., C.S., W.L.) and start-up funds from the St. Michael's Hospital Foundation (S.E.D.).

6. Acknowledgements

We thank Dr. Monika Lodyga, Dr. Caterina DiCiano-Oliveira and Dr. Xiaofeng Lu from research

facilities at St. Michael's hospital for help with training personnel in imaging and flow cytometry. We

thank staff at the Toronto Center for Phenogenomics for embedding and sectioning of our tissues. We

thank the staff of the University Health Network animal facility for their expert care of animals. We

thank Dr. Alan Lazarus and Dr. Veronica Miron for helpful discussions and mentorship of trainees.

7. References

1.   Taylor, A.L., Bettcher, D.W., *WHO Framework Convention on Tobacco Control: a global "good" for public health.* Bulletin of the World Health Organization, 2000, 78 (7), 2000: p. 920-929.

2.   *WHO global report on trends in prevalence of tobacco use 2000-2025, third addition.* 2019: p. 1-99.

3.   Dai, X., E. Gakidou, and A.D. Lopez, *Evolution of the global smoking epidemic over the past half century: strengthening the evidence base for policy action.* Tob Control, 2022. **31**(2): p. 129-137.

4.   Miller, F.W., *The increasing prevalence of autoimmunity and autoimmune diseases: an urgent call to action for improved understanding, diagnosis, treatment, and prevention.* Curr Opin Immunol, 2023. **80**: p. 102266.

5.   Bach, J.F., *The effect of infections on susceptibility to autoimmune and allergic diseases.* N Engl J Med, 2002. **347**(12): p. 911-20.

6.   Perricone, C., et al., *Smoke and autoimmunity: The fire behind the disease.* Autoimmun Rev, 2016. **15**(4): p. 354-74.

7.   Kallberg, H., et al., *Smoking is a major preventable risk factor for rheumatoid arthritis: estimations of risks after various exposures to cigarette smoke.* Ann Rheum Dis, 2011. **70**(3): p. 508-11.

8. Hedstrom, A.K., T. Olsson, and L. Alfredsson, *Smoking is a major preventable risk factor for*

multiple sclerosis. Mult Scler, 2016. **22**(8): p. 1021-6.

9. Hedstrom, A.K., et al., *Tobacco smoking, but not Swedish snuff use, increases the risk of multiple*

sclerosis. Neurology, 2009. **73**(9): p. 696-701.

10. Zivadinov, R., et al., *Smoking is associated with increased lesion volumes and brain atrophy in*

multiple sclerosis. Neurology, 2009. **73**(7): p. 504-10.

11. Healy, B.C., et al., *Smoking and disease progression in multiple sclerosis.* Arch Neurol, 2009.

66(7): p. 858-64.

12. Alshehri, E., et al., *The impact of cigarette smoking on cognitive processing speed and brain*

atrophy in multiple sclerosis. Mult Scler, 2023. **29**(7): p. 846-855.

13. Ramanujam, R., et al., *Effect of Smoking Cessation on Multiple Sclerosis Prognosis.* JAMA

Neurol, 2015. **72**(10): p. 1117-23.

14. Tanasescu, R., et al., *Smoking cessation and the reduction of disability progression in Multiple*

Sclerosis: a cohort study. Nicotine Tob Res, 2017.

15. Moutsianas, L., et al., *Class II HLA interactions modulate genetic risk for multiple sclerosis.* Nat

Genet, 2015. **47**(10): p. 1107-1113.

16. Karlson, E.W., et al., *Gene-environment interaction between HLA-DRB1 shared epitope and*

heavy cigarette smoking in predicting incident rheumatoid arthritis. Ann Rheum Dis, 2010. **69**(1): p. 54-

60.

17. Hedstrom, A.K., et al., *Smoking and two human leukocyte antigen genes interact to increase the*

risk for multiple sclerosis. Brain, 2011. **134**(Pt 3): p. 653-64.

18.   Ockinger, J., et al., *T-cell activation and HLA-regulated response to smoking in the deep airways*

of patients with multiple sclerosis. Clin Immunol, 2016. **169**: p. 114-120.

19.   Speyer, C.B. and K.H. Costenbader, *Cigarette smoking and the pathogenesis of systemic lupus*

erythematosus. Expert Rev Clin Immunol, 2018. **14**(6): p. 481-487.

20.   Barbhaiya, M., et al., *Cigarette smoking and the risk of systemic lupus erythematosus, overall*

and by anti-double stranded DNA antibody subtype, in the Nurses' Health Study cohorts. Ann Rheum

Dis, 2018. **77**(2): p. 196-202.

21.   Conforti, A., et al., *Beyond the joints, the extra-articular manifestations in rheumatoid arthritis.*

Autoimmun Rev, 2021. **20**(2): p. 102735.

22.   Makrygiannakis, D., et al., *Smoking increases peptidylarginine deiminase 2 enzyme expression*

in human lungs and increases citrullination in BAL cells. Ann Rheum Dis, 2008. **67**(10): p. 1488-92.

23.   Curran, A.M., et al., *Citrullination modulates antigen processing and presentation by revealing*

cryptic epitopes in rheumatoid arthritis. Nat Commun, 2023. **14**(1): p. 1061.

24.   Faigle, W., et al., *Brain Citrullination Patterns and T Cell Reactivity of Cerebrospinal Fluid-*

Derived CD4(+) T Cells in Multiple Sclerosis. Front Immunol, 2019. **10**: p. 540.

25.   Obeidat, M., et al., *The Overlap of Lung Tissue Transcriptome of Smoke Exposed Mice with*

Human Smoking and COPD. Sci Rep, 2018. **8**(1): p. 11881.

26.   Morissette, M.C., et al., *Persistence of pulmonary tertiary lymphoid tissues and anti-nuclear*

antibodies following cessation of cigarette smoke exposure. Respir Res, 2014. **15**(1): p. 49.

27.   Kroenke, M.A., et al., *IL-12- and IL-23-modulated T cells induce distinct types of EAE based on*

histology, CNS chemokine profile, and response to cytokine inhibition. J Exp Med, 2008. **205**(7): p. 1535-

41.

28.   Bettelli, E., et al., *Myelin oligodendrocyte glycoprotein-specific T cell receptor transgenic mice*

develop spontaneous autoimmune optic neuritis. J Exp Med, 2003. **197**(9): p. 1073-81.

29.   Berer, K., et al., *Commensal microbiota and myelin autoantigen cooperate to trigger autoimmune*

demyelination. Nature, 2011. **479**(7374): p. 538-41.

30.   Cahill, L.S., et al., *Aged hind-limb clasping experimental autoimmune encephalomyelitis models*

aspects of the neurodegenerative process seen in multiple sclerosis. Proc Natl Acad Sci U S A, 2019.

116(45): p. 22710-22720.

31.   Krishnamoorthy, G., et al., *Myelin-specific T cells also recognize neuronal autoantigen in a*

transgenic mouse model of multiple sclerosis. Nat Med, 2009. **15**(6): p. 626-32.

32.   Odoardi, F., et al., *T cells become licensed in the lung to enter the central nervous system.* Nature,

2012. **488**(7413): p. 675-9.

33.   Botelho, F.M., et al., *Innate immune processes are sufficient for driving cigarette smoke-induced*

inflammation in mice. Am J Respir Cell Mol Biol, 2010. **42**(4): p. 394-403.

34.   Becher, B., S. Tugues, and M. Greter, *GM-CSF: From Growth Factor to Central Mediator of*

Tissue Inflammation. Immunity, 2016. **45**(5): p. 963-973.

35.   Bosse, Y., et al., *Molecular signature of smoking in human lung tissues.* Cancer Res, 2012.

72(15): p. 3753-63.

36.   Botelho, F.M., et al., *IL-1alpha/IL-1R1 expression in chronic obstructive pulmonary disease and*

mechanistic relevance to smoke-induced neutrophilia in mice. PLoS One, 2011. **6**(12): p. e28457.

37.   Botelho, F.M., et al., *A mouse GM-CSF receptor antibody attenuates neutrophilia in mice*

exposed to cigarette smoke. Eur Respir J, 2011. **38**(2): p. 285-94.

38.   Morissette, M.C., et al., *Disruption of pulmonary lipid homeostasis drives cigarette smoke-*

induced lung inflammation in mice. Eur Respir J, 2015. **46**(5): p. 1451-60.

39.   Robbins, C.S., et al., *Cigarette smoke impacts immune inflammatory responses to influenza in*

mice. Am J Respir Crit Care Med, 2006. **174**(12): p. 1342-51.

40.   Roos, A.B., et al., *IL-17A and the Promotion of Neutrophilia in Acute Exacerbation of Chronic*

Obstructive Pulmonary Disease. Am J Respir Crit Care Med, 2015. **192**(4): p. 428-37.

41.   Rindlisbacher, L., M.N. Navarro, and B. Becher, *Inflame and restrain - the paradoxical roles of*

IL-12 and IL-23 in immunity. Nat Rev Immunol, 2026.

42.   Kearley, J., et al., *Cigarette smoke silences innate lymphoid cell function and facilitates an*

exacerbated type I interleukin-33-dependent response to infection. Immunity, 2015. **42**(3): p. 566-79.

43.   Drohomyrecky, P.C., et al., *Peroxisome Proliferator-Activated Receptor-delta Acts within*

Peripheral Myeloid Cells to Limit Th Cell Priming during Experimental Autoimmune Encephalomyelitis.

J Immunol, 2019. **203**(10): p. 2588-2601.

44.   Qiu, S.L., et al., *Cigarette Smoke Induction of Interleukin-27/WSX-1 Regulates the*

Differentiation of Th1 and Th17 Cells in a Smoking Mouse Model of Emphysema. Front Immunol, 2016.

7: p. 553.

45.   Xu, W., R. Li, and Y. Sun, *Increased IFN-gamma-producing Th17/Th1 cells and their*

association with lung function and current smoking status in patients with chronic obstructive pulmonary

disease. BMC Pulm Med, 2019. **19**(1): p. 137.

46.   Brozyna, S., et al., *Chemotactic mediators of Th1 T-cell trafficking in smokers and COPD*

patients. COPD, 2009. **6**(1): p. 4-16.

47.   Shan, M., et al., *Lung myeloid dendritic cells coordinately induce TH1 and TH17 responses in*

human emphysema. Sci Transl Med, 2009. **1**(4): p. 4ra10.

48.   Shan, M., et al., *Cigarette smoke induction of osteopontin (SPP1) mediates T(H)17 inflammation*

in human and experimental emphysema. Sci Transl Med, 2012. **4**(117): p. 117ra9.

49.   Naessens, T., et al., *Human Lung Conventional Dendritic Cells Orchestrate Lymphoid*

Neogenesis during Chronic Obstructive Pulmonary Disease. Am J Respir Crit Care Med, 2020. **202**(4):

p. 535-548.

50.   Cooper, A.M. and S.A. Khader, *IL-12p40: an inherently agonistic cytokine.* Trends Immunol,

2007. **28**(1): p. 33-8.

51.   Gounder, A.P., et al., *Interferon induced protein 35 exacerbates H5N1 influenza disease through*

the expression of IL-12p40 homodimer. PLoS Pathog, 2018. **14**(4): p. e1007001.

52.   Teng, M.W., et al., *IL-12 and IL-23 cytokines: from discovery to targeted therapies for immune-*

mediated inflammatory diseases. Nat Med, 2015. **21**(7): p. 719-29.

53.   Grifka-Walk, H.M., D.A. Giles, and B.M. Segal, *IL-12-polarized Th1 cells produce GM-CSF and*

induce EAE independent of IL-23. Eur J Immunol, 2015. **45**(10): p. 2780-6.

54.   Guo, D.H., et al., *Visceral adipose NLRP3 impairs cognition in obesity via IL-1R1 on CX3CR1+*

cells. J Clin Invest, 2020. **130**(4): p. 1961-1976.

55.   Tian, X., et al., *Blocking IL-23 Signaling Mitigates Cigarette Smoke-Induced Murine*

Emphysema. Environ Toxicol, 2024. **39**(12): p. 5334-5346.

56.   Khader, S.A., et al., *Interleukin 12p40 is required for dendritic cell migration and T cell priming*

after Mycobacterium tuberculosis infection. J Exp Med, 2006. **203**(7): p. 1805-15.

57.   Cua, D.J., et al., *Interleukin-23 rather than interleukin-12 is the critical cytokine for autoimmune*

inflammation of the brain. Nature, 2003. **421**(6924): p. 744-8.

58.   Mondal, S., A. Roy, and K. Pahan, *Functional blocking monoclonal antibodies against IL-12p40*

homodimer inhibit adoptive transfer of experimental allergic encephalomyelitis. J Immunol, 2009.

182(8): p. 5013-23.

59.   Segal, B.M., et al., *Repeated subcutaneous injections of IL12/23 p40 neutralising antibody,*

ustekinumab, in patients with relapsing-remitting multiple sclerosis: a phase II, double-blind, placebo-

controlled, randomised, dose-ranging study. Lancet Neurol, 2008. **7**(9): p. 796-804.

60.   Vollmer, T.L., et al., *A phase 2, 24-week, randomized, placebo-controlled, double-blind study*

examining the efficacy and safety of an anti-interleukin-12 and -23 monoclonal antibody in patients with

relapsing-remitting or secondary progressive multiple sclerosis. Mult Scler, 2011. **17**(2): p. 181-91.

61.   Gordon, K.B., et al., *A phase III, randomized, controlled trial of the fully human IL-12/23 mAb*

briakinumab in moderate-to-severe psoriasis. J Invest Dermatol, 2012. **132**(2): p. 304-14.

62.   Chitnis, T., Singhal, T., Zuraswski, J., Saraceno, T.J., Gopalakrishnan, N., Cain, L., LaBarre, B.,

King, D., Bermark, R.W., Maxfield, A.Z., Cicero, S., Pan, H., Dubey, S., Vaquerano, S., Hansel, C.,

Healy, B.C., Saxena, S., Lokhande, H.A., Bharnoori, M., Madill, E.S., Sheth, M.S., Rodin, R., Ye, J.,

Clementi, N., Clementi, W.A., and Weiner, H.L., *Nasal Foralumab for the Treatment of Progression*

Independent of Relapses in Patients With Nonactive Secondary Progressive Multiple Sclerosis.

Neurology Neuroimmunology, and Neuroinflammation, 2026. **13**(2): p. e200543.

63.   Blanquart, E., et al., *Targeting androgen signaling in ILC2s protects from IL-33-driven lung*

inflammation, independently of KLRG1. J Allergy Clin Immunol, 2022. **149**(1): p. 237-251 e12.

64.   Bal, S.M., et al., *IL-1beta, IL-4 and IL-12 control the fate of group 2 innate lymphoid cells in*

human airway inflammation in the lungs. Nat Immunol, 2016. **17**(6): p. 636-45.

65.   Blackmore, S., et al., *Influenza infection triggers disease in a genetic model of experimental*

autoimmune encephalomyelitis. Proc Natl Acad Sci U S A, 2017. **114**(30): p. E6107-E6116.

66.   Wilcoxen, S.C., et al., *Gender-dependent IL-12 secretion by APC is regulated by IL-10.* J

Immunol, 2000. **164**(12): p. 6237-43.

67.   Douin-Echinard, V., et al., *Estrogen receptor alpha, but not beta, is required for optimal dendritic*

cell differentiation and [corrected] CD40-induced cytokine production. J Immunol, 2008. **180**(6): p.

3661-9.

68.   Zhang, M.A., et al., *Peroxisome proliferator-activated receptor (PPAR)alpha and -gamma*

regulate IFNgamma and IL-17A production by human T cells in a sex-specific way. Proc Natl Acad Sci

U S A, 2012. **109**(24): p. 9505-10.

69.   Kadel, S., et al., *A Major Population of Functional KLRG1(-) ILC2s in Female Lungs Contributes*

to a Sex Bias in ILC2 Numbers. Immunohorizons, 2018. **2**(2): p. 74-86.

70.   Laffont, S., et al., *Androgen signaling negatively controls group 2 innate lymphoid cells.* J Exp

Med, 2017. **214**(6): p. 1581-1592.

71.   Doz, E., et al., *Cigarette smoke-induced pulmonary inflammation is TLR4/MyD88 and IL-*

1R1/MyD88 signaling dependent. J Immunol, 2008. **180**(2): p. 1169-78.

72.   Zhang, P., et al., *HMGB1 mediates Aspergillus fumigatus-induced inflammatory response in*

alveolar macrophages of COPD mice via activating MyD88/NF-kappaB and syk/PI3K signalings. Int

Immunopharmacol, 2017. **53**: p. 125-132.

73.   Kronzer, V.L., et al., *Lifestyle and Clinical Risk Factors for Incident Rheumatoid Arthritis-*

associated Interstitial Lung Disease. J Rheumatol, 2021. **48**(5): p. 656-663.

74.   Egesten, A., et al., *Increased prevalence of multiple sclerosis among COPD patients and their*

first-degree relatives: a population-based study. Lung, 2008. **186**(3): p. 173-178.

75.   Marrie, R.A., et al., *Multiple sclerosis and antecedent infections: a case-control study.*

Neurology, 2000. **54**(12): p. 2307-10.

76.   Buljevac, D., et al., *Prospective study on the relationship between infections and multiple*

sclerosis exacerbations. Brain, 2002. **125**(Pt 5): p. 952-60.

77.   Kriesel, J.D., et al., *Multiple sclerosis attacks are associated with picornavirus infections.* Mult

Scler, 2004. **10**(2): p. 145-8.

78.   Sibley, W.A., C.R. Bamford, and K. Clark, *Clinical viral infections and multiple sclerosis.*

Lancet, 1985. **1**(8441): p. 1313-5.

79.   Panitch, H.S., *Influence of infection on exacerbations of multiple sclerosis.* Ann Neurol, 1994. **36**

Suppl(Suppl): p. S25-8.

80.   Edwards, S., et al., *Clinical relapses and disease activity on magnetic resonance imaging*

associated with viral upper respiratory tract infections in multiple sclerosis. J Neurol Neurosurg

Psychiatry, 1998. **64**(6): p. 736-41.

81.   Tremlett, H., et al., *Monthly ambient sunlight, infections and relapse rates in multiple sclerosis.*

Neuroepidemiology, 2008. **31**(4): p. 271-9.

82.   Chen, Q., et al., *Influenza virus infection exacerbates experimental autoimmune*

encephalomyelitis disease by promoting type I T cells infiltration into central nervous system. J

Autoimmun, 2017. **77**: p. 1-10.

83.   Hosang, L., et al., *The lung microbiome regulates brain autoimmunity.* Nature, 2022. **603**(7899):

p. 138-144.

84.   Saito, E., et al., *Modulating lung immune cells by pulmonary delivery of antigen-specific*

nanoparticles to treat autoimmune disease. Sci Adv, 2020. **6**(42).

85.   Enzmann, G., et al., *The Genetic Background of Mice Influences the Effects of Cigarette Smoke*

on Onset and Severity of Experimental Autoimmune Encephalomyelitis. Int J Mol Sci, 2019. **20**(6).

86.   Rubin, R.L., et al., *Effect of cigarette smoke on autoimmunity in murine and human systemic*

lupus erythematosus. Toxicol Sci, 2005. **87**(1): p. 86-96.

87.   Muscat, J.E., et al., *Time to first cigarette after waking predicts cotinine levels.* Cancer Epidemiol

Biomarkers Prev, 2009. **18**(12): p. 3415-20.

88.   Ucciferri, C.C., et al., *Scoring Central Nervous System Inflammation, Demyelination, and Axon*

Injury in Experimental Autoimmune Encephalomyelitis. J Vis Exp, 2024(204).

89.   Cordeiro, B., et al., *Obesity intensifies sex-specific interferon signaling to selectively worsen*

central nervous system autoimmunity in females. Cell Metab, 2024. **36**(10): p. p2298-2314.

8. Figure Legends

Figure 1. Chronic exposure of mice to cigarette smoke (CS) enhances MOGp35-55-specific Th1

responses. Ambient air (AA)- or CS-exposed male C57BL6/J mice were immunized with MOG p35-

55/CFA and the recall cytokine and proliferative responses of cells were assessed 9 days later in spleen

and draining lymph node (dLN) mononuclear cells. (A-B) Levels of cytokines in supernatants of MOG_{35-}

$_{55}$-stimulated cell cultures from spleens (A) and dLNs (B) measured using a multiplex assay. (C-E)

Division of splenic $CD44^{high}CD4^+$ T cells after stimulation in the presence/absence of MOG p35-55 *ex*

vivo as assessed by CFSE dilution. Representative plots of dividing $CD4^+$ T cells (C), and frequency (D)

and total numbers (E) of dividing $CD44^{high}CD4^+$ T cells. Data in A-B are mean $\pm$ SEM of triplicate wells

of cultures from pooled mice and representative of three independent experiments in males and two

independent experiments in females. Data points in C are individual mice from one experiment

representative of 3 proliferation assays performed. *: p≤0.05 between groups as determined by Mann-

Whitney (A,B) or 2-way ANOVA and Bonferroni post hoc test (C,D).

Figure 2. Cigarette smoke (CS) exposure in males decreases the severity of active EAE, yet

promotes spontaneous autoimmunity in MOG T cell receptor transgenic mice. (A) Schematic of

experiments: C57BL/6J mice were exposed to 8-10 weeks of ambient air (AA) or CS prior to active EAE

induction by immunization with pMOG/CFA plus pertussis toxin. Alternatively, 2D2 mice were exposed

to AA or CS for up to 8 weeks or until clinical symptoms developed. Black denotes males and red denotes

females. (B) EAE clinical scores over time, (C) Cumulative EAE scores, (D) Peak EAE scores, and (E)

Day of EAE onset. (F-G) Histological scoring of EAE was done at endpoint by counting the density of

$CD45^+$ cells/mm^2 of white matter (WM) (F) or percent demyelination (G) at the level of the thoracic

spine in AA- or CS-exposed EAE mice. (H) Frequency of 2D2 mice developing CNS autoimmunity after

AA- or CS-exposure. (I) Representative CD45 staining showing sparse immune cell infiltration in the

spinal cord, the dorsal root ganglion and spinal nerve roots of CS-exposed 2D2 mice. Clinical data in (H)

are from two consecutive cohorts each with (n=9-21/group/sex). *: p≤0.05 between groups as determined

by two-way ANOVA and Bonferroni post hoc test (B), by two-tailed Mann-Whitney U test (B-G), or by

χ2 test (H).

Figure 3. **Myelin-reactive T cells are stalled and hyperactivated in cigarette smoke (CS)-exposed**

lungs. Passive EAE was induced in male C57BL/6J mice that had been exposed to ambient air (AA) or

CS, by transferring IL-23-polarized, MOG_{35-55}-reactive female T cells from unexposed donor mice. (A)

EAE scores in AA- and CS-exposed recipients. (B) Representative flow cytometry plots of $CD45.1^+$ T

helper cells in the CNS compartment in established EAE (*grey arrows* in A). (C-D) Total numbers of

mononuclear cells (C) and of $CD45.1^+CD4^+$ T cells (D); (E-F) frequencies of MOG_{35-55}-reactive

CD45.1$^+$CD4$^+$CD44^{high} T cells producing IL-17 (E) or co-producing IL17 and GM-CSF (F). (G) Shows

representative FACs plots of CD4$^+$ T cells in the lung. (H) Total numbers of IL17$^+$, IL17$^+$IFN-γ$^+$, and

IFN-γ$^+$ CD4$^+$ T cells in lungs. (I) IL-17 and GM-CSF median fluorescence intensities (MFIs) of lung

MOG$_{35-55}$–reactive IL17$^+$IFN-γ$^-$ CD4$^+$ T cells. (J) CD25, CD69 and PD-1 expression by donor

CD45.1$^+$CD4$^+$CD44^{high} T cells in the lungs. (K) Immunolabeling of donor CD45.1$^+$ cells in the lungs.

Data in A are presented as mean ± SEM of individual mice from one representative experiment of at least

four performed in males. Data in C-F are mean ± SEM of individual mice from one experiment of two

that were performed, one conducted at disease onset and one at peak disease. Data investigating cytokines

and activation markers in the lung are mean + SEM of individual mice and are representative of 4

independent experiments. *: p≤0.05 between groups as determined by two-way ANOVA and Bonferroni

post hoc test (H) or by two-tailed Mann-Whitney U test (A,C-F, I-J).

Figure 4. Cigarette smoke (CS) increases pro-inflammatory immune pathways in human lungs and

cytokine and chemokine levels in murine lungs. (A) GO term enrichment analysis of microarray-

derived differentially expressed genes (DEGs) from human lung tissue of smoking versus non-smoking

patients [35]. Significantly enriched immune pathways, and their associated DEGs, were identified using

a hypergeometric test with Benjamini–Hochberg correction (FDR > 0.05), with representative immune

response and specific cytokine signaling pathways displayed on the heatmap. B) Dotplot of immune

response-related MSigDB Hallmark and Reactome gene sets significantly enriched in an exploratory list

of microarray-derived DEGs from the human lung tissue of smoking versus non-smoking patients [35].

Significance was determined using the Monte-Carlo method with Benjamini–Hochberg correction (FDR

> 0.05). (C) Lysates were prepared from the lungs for AA- or CS-exposed mice that had been immunized

with pMOG/CFA. Lung were harvested at d9 post-immunization, were snap frozen in liquid nitrogen,

and lungs were homogenized in TPER buffer. Analytes were measured using a multiplex assay and

individual ELISA kits. Shown is a heatmap of pg/ml levels of each analyte normalized for total lung

protein measured. Black asterisks indicate a difference of CS-group from AA-group in males. Red

asterisks indicate a difference of the CS-group from the AA-group in females. Blue asterisks indicate an

interaction. Data were analyzed by 2-way ANOVA followed by a Bonferroni post-hoc test. Values

represent tissues collected from individual mice from 2 experiments.

Figure 5. Cigarette smoke (CS) increases the activation of antigen-presenting cells (APCs) in the

lung and lung CD11c$^+$ cells from CS-exposed mice have a greater capacity to prime myelin-reactive

Th cells. Male C57BL/6J mice were exposed to ambient air (AA) or CS for 8-10 weeks and then were

transferred with MOG p35-55 reactive T cells. Lungs were harvested at a time when AA-exposed mice

had developed EAE and CS-exposed mice were still asymptomatic. (A) Representative FACS plots of

the expression of activation markers (CD86, MHC-II) in APCs isolated from male C57BL/6J mice

exposed to ambient air (AA) or CS. (B-F) Dendritic cells (DCs) isolated from lungs of AA- and CS-

exposed male C57BL/6J mice were cultured in presence of MOG$_{35-55}$ (0-10 µg/mL) with myelin-reactive

CD4$^+$ T cells isolated from spleens of 2D2 male mice. Proliferation of 2D2 CD4$^+$ T cells (B), and

expression of CD25 (C), CD69 (D), IFN-γ (E) and IL-17 (F) on memory 2D2 CD4$^+$ T cells.

Figure 6. IL-12p40 is a mediator of the effect of CS-exposure in enhancing Th17 and GM-CSF

production by myelin-reactive T helper cells. (A-D) AA- and CS-exposed lung DCs were co-cultured

with splenic myelin-reactive 2D2 CD4$^+$ T cells in presence/absence of MOG$_{35-55}$ (5 µg/mL) with either

blocking anti-IL12p40 or isotype control Ab. Proliferation of 2D2 CD4$^+$ T cells (A), expression of CD25

(B), IFN-γ (C) and IL-17 (D) in 2D2 CD44^{hi}CD4$^+$ T cells. (E-K) Passive EAE was induced by

transferring IL-23-polarized, MOG$_{35-55}$-reactive T cells from unexposed donor CD45.1$^+$ mice into AA-

and CS-exposed male C57BL/6J (CD45.2$^+$) mice; 2 days prior to, and 1 day after the adoptive transfer.

Recipient mice received blocking anti-IL12p40 or isotype control Ab intranasally (125 µg/animal/dose).

(E) Representative cytokine staining in the donor CD45.1$^+$ CD4$^+$ T cells. (F-H) Numbers of total CD45

cells (F), donor CD4$^+$ T cells (G) and MOG$_{35-55}$-reactive donor CD44^{high}CD4$^+$ T cells producing IL-17

(H). (I-J) Frequencies of donor CD4$^+$ T cells expressing and IL-17 (I) with GM-CSF (J). (K) Expression

of CD25 on the donor CD4$^+$ T cells. Data presented are mean ± SEM of triplicates of one experiment

(A-D) or are individual mice (F-K). *: p≤0.05 between groups as determined by two-way ANOVA and

Bonferroni post hoc test.

Supplementary Figure 1. Exposure of mice to CS over eight weeks induces immune cell infiltration

in the lung. (A-D) Sections of lungs of AA- and CS-exposed males were stained for CD45. Arrows

show immune cell clusters located close to blood vessels and bronchioles. (B-C) Shown are the number

of CD45$^+$ cells (B) and the mean area of clusters of CD45$^+$ cells in mm^2 (C). (D-G) Lung mononuclear

cells were isolated from AA- and CS-exposed and were analyzed by flow cytometry. (D-E) The total

number of live CD45 leukocytes isolated from the lungs. (F) Gating strategy (pre-gated on live singlets)

used to identify lung immune cell subsets. (G) Heat map of the numbers of the different immune

populations in individual mice (n=5 mice per group). Data in B-E are presented as mean ± SEM. Data

are from one representative experiment of at least three that were performed. *: p≤0.05 between groups

as determined by two-tailed Mann-Whitney U test (B-E) or 2-way ANOVA followed by Bonferroni post-

hoc test (G). In (G), p≤0.05 between male groups is shown with black asterisks, and between female

groups with red asterisks.

Supplementary Figure 2. B and T cells accumulate in perivascular and peri-bronchial clusters in

cigarette smoke (CS)-exposed lungs. Dual CD3 (DAB) and B220 (Vector Blue) staining in lung

sections of ambient air (AA) and CS-exposed mice. (A) Presence of scattered T cells in the AA-exposed

lung. (B-D) Examples of pathology in the CS-exposed lungs. (B) Shows a larger and smaller

lymphocytic infiltrate containing T cells and B cells (arrows). (C) Shows a less organized lymphocytic

structure around the airway with T and B cells present (arrows). (D) Shows an example of T cell clusters

with no B cells present.

Supplementary Figure 3. Effect of CS exposure on pMOG-reactive Th cytokine responses in female

mice during active EAE. A-B) Ambient air (AA)- or CS-exposed female C57Bl/6J mice were

immunized with MOG$_{35-55}$ to assess the *in vitro* cytokine and proliferative recall responses of cells

isolated from spleen and draining lymph nodes (dLNs) 9 days after immunization. Levels of Th1, Th17

and Th2 cytokines in supernatants of MOG$_{35-55}$-stimulated (0-10 µg/mL) cell cultures from spleens (A)

and dLNs (B). Data are mean ± SEM of triplicate wells of cultures from pooled mice from one experiment

that is representative of two performed. *: p≤0.05 between groups as determined by Mann-Whitney test.

Supplementary Figure 4. EAE is still attenuated in male mice if CS-exposures are stopped prior to

the induction of EAE. Male C57BL/6J mice were exposed to 8-10 weeks of AA or CS, followed of a

4.5 day washout period before active EAE induction according to the schematic in (A). EAE scores (B)

and cumulative EAE scores (C). Data are presented as mean ± SEM of individual mice from one

experiment. *: p≤0.05 between groups as determined by two-way ANOVA and Bonferroni post hoc test

(B) or by two-tailed Mann-Whitney U test (C).

Supplementary Figure 5. CS also induces a hyperactivation of female Th cells in passive transfer

EAE, but does not activate the host cells. (A) Passive EAE was induced in male (black) or female

(red) C57BL/6J mice that had been exposed to ambient air (AA) and CS, by transferring IL-23-polarized,

MOG$_{35-55}$-reactive female T cells from unexposed donor mice. Mononuclear cells were isolated from the

lungs and the spleen at a time when EAE was delayed in the CS group. (A) Numbers of MOG$_{35-55}$-

reactive T cells producing IL17, co-producing IL17 and IFN-γ, or producing IFN-γ in spleens of male

mice during passive transfer EAE as assessed by flow cytometry after stimulation *in vitro* with MOG$_{35-}$

$_{55}$ for 12 hours, with GolgiStop added in the last 6 h. (B-D) Results from an experiment done in females

mice. (B) The numberof donor pMOG-reactive T cells detected in the AA- and CS-exposed female lungs

in passive EAE. (C) The expressions CD25 and CD69 by median fluorescence intensity (MFI) in donor

T cells in the lungs of female recipients. (D) Shows the clinical scores of mice in a companion experiment

where females were followed for clinical signs. (E) Shows expressions of CD25 and CD69 in recipient

T cells (CD45.1$^-$ CD4$^+$) in the lungs of an experiment done in males. Data in A and E are mean + SEM

of individual mice in one experiment that is representative of 4 independent experiments that were

performed. Data points in B-D are individual mice from one experiment of two that were performed. *:

p≤0.05 between groups as determined by two-way ANOVA and Bonferroni post hoc test (A) or by two-

tailed Mann-Whitney U test (B-D).

Supplementary Figure 6. Cigarette smoke (CS) enhances cytokine production by 2D2 CD4$^+$ T cells

in the lungs. Male 2D2 mice aged 6-7 weeks were exposed to ambient air (AA) or CS. Mice were

followed for clinical signs and flow cytometry was performed at endpoint to examine the profile of 2D2

CD4$^+$ T cells in the lung and spleen. A) Total number of cells in lungs. B) Total number of myelin-

reactive, transgenic TCR-expressing, 2D2 CD4$^+$ T cells in the lungs. C) Distribution of naïve (CD44$^-$

CD62L$^+$), central memory (CD44$^+$CD62L$^+$) and effector memory (CD44$^+$CD62L$^-$) in myelin-reactive

2D2 CD4$^+$ T cells in the lungs in the AA- and CS-exposed mice. D-E) Median fluorescence intensity

(MFI) of CD25 (D) and CD69 (E) in memory (CD44^{high}) and naïve (CD44^{low}) myelin-reactive 2D2 CD4$^+$

T cells in the lungs. F) MFI of IFN-γ and IL-17 in lung cytokine-producing, memory 2D2 CD4$^+$ T cells

cultured *in vitro* with/without MOG$_{35-55}$. G) Representative FACS histograms of IFN-γ and IL-17

production by lung cytokine-producing, memory 2D2 CD4$^+$ T cells. G) Total number of cells in spleen.

H) Total number of myelin-reactive, transgenic TCR-expressing, 2D2 CD4$^+$ T cells in the spleen. I)

Distribution of naïve (CD44$^-$CD62L$^+$), central memory (CD44$^+$CD62L$^+$) and effector memory

(CD44$^+$CD62L$^-$) phenotypes in myelin-reactive 2D2 CD4$^+$ T cells in the spleen. J) Median fluorescence

intensity (MFI) of CD69 in memory (CD44^{high}) myelin-reactive 2D2 CD4$^+$ T cells in the spleen. K-L)

MFI of IFN-γ and IL-17 in splenic cytokine-producing, memory 2D2 CD4$^+$ T cells cultured *in vitro*

with/without MOG$_{35-55}$. Data are presented as mean ± SEM of individual mice (n=8-9/group) from one

experiment. *: p≤0.05 between groups as determined by two-tailed Mann-Whitney U test (A-B, C-D, E,

G-L), χ2 test (C, I), and two-way ANOVA and Bonferroni post hoc test (F).

Supplementary Figure 7. Cigarette smoke (CS) increases the activation of antigen presenting cells

(APCs) in the lung. Female C57BL/6J mice were exposed to ambient air (AA) or CS for 8-10 weeks

and then were transferred with MOG p35-55 reactive T cells. Lungs were harvested at a time when AA-

exposed mice had developed EAE and CS-exposed mice were still asymptomatic. Representative FACS

plots of the expression of activation markers (CD86, MHC-II) in APCs isolated from female C57BL/6J

mice exposed to AA or CS.

Supplementary Figure 8. Blockade of lung IL-12p40 production by intranasal vs intratracheal

administration of a blocking anti-IL-12p40 antibody. (A) Blocking anti-IL12p40 Ab (0-250

μg/animal) was administered either intranasally or intratracheally to male C57BL/6J mice; three days

after the administration, IL-12p40 levels were measured in tissue lysates of lungs, small intestine, and

spleen using an ELISA kit that used different antibodies from the ones administered. Data are presented

as mean ± SEM of individual mice from one experiment *: p≤0.05 between groups as determined by

two-way ANOVA and Bonferroni post hoc test; asterisks show statistical significance between 0 and the

marked concentration of Ab for intranasal (black asterisks) and intratracheal (grey asterisks)

administration.

A

Male or female (6 weeks)

↓ | 8 weeks | MOG₃₅₋₅₅/CFA+PTx ↓ Clinical scoring

CS or AA (2x daily, 5 days/week)

Male or female 2D2 (6 weeks)

Clinical scoring

CS or AA (2x daily, 5 days/week)

△ AA males   ▲ CS males   ○ AA females   ● CS females

I

| Dorsal spinal cord | Dorsal root ganglion (DRG) | | Nerve |

50 μm

CD45 Hematoxylin

CNS mononuclear cells

A Immune GO biological processes in smoking versus non-smoking patients (Z-scores)

B Immune programs enriched in smoking patients
MSigDB Hallmark and Reactome GSEA