Angiogenic CD8 T cells from PWH induce Granzymes-dependent PAR1 activation promoting endothelial inflammation

Tong Li[1], Chinmayee Mehta[1], Cooper James[1], Jinya Chen[1], Hui Chen[1], Gerard P Ahern[2], Ziang Zhu[1], Maitane Faus Cid[1], Maryam Abdussamad[1], Shiwen Yu[1], Jai Kumar[3], Princy Kumar[3], Colleen Hadigan[4], Kyle DiVito[5], Dorian B McGavern[6], Leonid Pobezinsky[7], and Marta Catalfamo[1]

[1]Department of Microbiology and Immunology, Georgetown University School of Medicine. Washington DC, USA

[2]Department of Pharmacology and Physiology, Georgetown University School of Medicine. Washington DC, USA

[3]Division of Infectious Diseases and Travel Medicine, Georgetown University School of Medicine. Washington DC, USA

[4]Laboratory of Immunoregulation, National Institute of Allergy and Infectious Diseases, National Institutes of Health, Bethesda, MD 20892, USA

[5]Department of Biochemistry, Molecular and Cellular Biology, Georgetown University School of Medicine. Washington DC, USA

[6]Viral Immunology and Intravital Imaging Section, National Institute of Neurological Disorders and Stroke, National Institutes of Health, Bethesda, Maryland, USA

[7]Department of Veterinary and Animal Science, University of Massachusetts, Amherst, MA, USA

Abstract

In people with HIV (PWH), T cell immune activation and endothelial inflammation are

contributors of the increased cardiovascular risk but the mechanisms remain poorly understood. In

this study, we hypothesized that immune activation alters angiogenetic CD8 T cell (T_{ang}) function

driving endothelial chronic inflammation. Endothelial cell repair is partially mediated by

angiogenic T (T_{ang}) cells which facilitate proliferation and differentiation of bone marrow derived

and resident endothelial progenitor cells at site of vascular injury. The role of CD8 T_{ang} cells in

the setting of viral infection, including chronic HIV infection is largely unknown. In this study, we

identified a subset of CD8 T cells expressing $CD31^{high}$ that exhibited a long-lived memory

precursor phenotype ($CD127^+KLRG1^-$) and secreted the pro-angiogenic cytokine vascular

endothelial growth factor (VEGF) in mice infected with acute LCMV. Furthermore, in PWH, the

frequencies of circulating CD8 T_{ang} cells were reduced and showed an activated phenotype and

expression of granzymes. $GZMA^+GZMB^+CD8$ T_{ang} cells correlated with atherosclerotic

cardiovascular disease (ASCVD) risk. *In vitro*, granzyme dependent PAR1 activation led to

calcium mobilization and secretion of pro-inflammatory cytokines IL-6, IL-8 and Angiopoietin-2

by primary human endothelial cells. Altogether, these findings suggest that HIV driven activation

of CD8 T_{ang} cells promotes endothelial inflammation and chronic stimulation of this pathway may

contribute to the increased risk of cardiovascular disease in PWH.

Introduction

Immune activation remains the hallmark of HIV infection, despite successfully suppressed

viremia by antiretroviral therapy (ART). HIV-associated inflammation and immune activation

drives cardiovascular risk, independent of traditional risk factors, duration of antiretroviral

treatment and CD4 counts (1-15). Compared to the general population, People with HIV (PWH)

have twice the risk of cardiovascular disease and standard cardiovascular risk prediction scores

underestimate cardiovascular risk in this population   (16-18). The Strategic Timing of

AntiRetroviral Treatment (START) trial reported that the risk of AIDS and Serious non-AIDS

associated events (SNAEs) in early versus delayed treatment initiation was reduced, however,

excess risk remains for the most common non-AIDS defining illnesses, cardiovascular disease

(CVD) and cancer (19).

Atherosclerosis, the underlying pathology of cardiovascular events, is characterized by lipid

deposition, chronic inflammation, endothelial dysfunction; where cells from the innate and

adaptive immune system participate in the pathology of the disease (20). In PWH, CD8 T cell

immune activation is associated with carotid intima-media thickness and coronary plaque

suggesting an interplay between CD8 T cell activation and endothelial inflammation and

dysfunction (10, 11, 21-23). The molecular pathways by which HIV contributes to increased risk

of CVD remain largely unknown, which makes the development of more targeted therapies

difficult.

During systemic infections such as HIV, endothelial activation and injury can be mediated

by viral proteins and the host response (24, 25). Endothelial repair, is mediated in part by

angiogenic T cells (T_{ang}) that support proliferation and differentiation of bone marrow derived and

resident endothelial progenitor cells/endothelial colony forming cells (EPC/ECFC) at the site of

endothelial injury (26-28). In humans, T_{ang} cells have been defined by co-expression of CD31

(PECAM-1, Platelet endothelial cell adhesion molecule) and CXCR4 (stromal-cell-derived factor

receptor) (26). In addition, reduced circulating frequencies have been associated with age and

increased cardiovascular risk (26, 29, 30). However, their role in the setting of viral infections

including HIV has not been investigated.

Under physiological conditions, vascular permeability induced by trauma or infection triggers

activation of the coagulation cascade (leading to thrombin generation and clot formation) and a

host inflammatory response that promotes vascular and tissue repair. Thrombin, the main serine

protease of the coagulation cascade, connects the coagulation cascade, platelets and the host

inflammatory response by signaling through its receptor the Protease Activated Receptor 1 (PAR1)

expressed on immune cells. PAR1 is one of the four members of the seven transmembrane domain

G protein-coupled receptor family (PAR1, PAR2, PAR3 and PAR4) (31). PAR's activation is

mediated by proteolytic cleavage of the N-terminus that uncovers a tethered ligand (32, 33). In

humans, PAR1, PAR3 and PAR4 are activated by thrombin, whereas in mice PAR3 and PAR4 are

the main receptor for thrombin. In addition, PAR2 is activated by trypsin, tryptase and other

coagulation factors (32, 34-37).

CD8 T cells express PAR1, and its activation by thrombin enhances T cell function (32, 38-

41). These data provided evidence that PAR1 signaling mediates the cross-talk between the

coagulation system, endothelium and the adaptive immune system (CD8 T cells) at sites of

endothelial injury.

Thrombin-dependent activation of PAR1 occurs by proteolytic cleavage of the extracellular

domain at the Arg^{41}-Leu^{42} (32, 42, 43). In addition, other proteases including granzymes (A, B and

K) activate PAR1 although the cleavage site is not well defined. Moreover, granzyme-dependent

activation of PAR1 induces downstream signaling pathways that differ from thrombin leading to

a non-canonical activation or "biased signal" (43-50). Extracellular granzymes A, B and K have

been shown to induce inflammation by activating PAR1 in multiple cell types including neurons,

fibroblast and endothelial cells (44, 46-48, 50, 51).

In this study, we evaluated whether T_{ang} cells are involved in endothelial repair in response to

immune mediated injury during acute Lymphocytic Choriomeningitis virus (LCMV) infection.

We found that CD31^{high} expressing CD8 T_{ang} cells increased after viral control, have a memory

precursor phenotype (KLRG1$^-$CD127$^+$) and secrete vascular endothelial growth factor (VEGF)

suggesting their contribution in endothelial repair. Moreover, we found that in the setting of

chronic HIV infection, immune activation drives a proinflammatory phenotype of CD8 T_{ang}

expressing granzymes. The frequencies of GZMA$^+$GZMB$^+$CD8 T_{ang} are associated with the

atherosclerotic cardiovascular disease (ASCVD) risk. In addition, plasma levels of GZMA

correlated with endothelial inflammatory markers including sVCAM1 and IL-6. *In vitro* GZMs

activate PAR1 favoring the secretion of pro-inflammatory cytokines over the pro-angiogenic

cytokines. This evidence supports the notion that HIV driven immune activation alters CD8 T_{ang}

cell function and chronic stimulation of this pathway may contribute endothelial inflammation and

the increased risk of cardiovascular disease.

Material and Methods

Study Participants

Participants were studied under a MedStar Georgetown University Hospital and NIAID

Institutional Review Board approved HIV clinical research studies. Characteristics of the study groups

of PWoH (n= 17) and PWH (n= 25) are described in Table 1. All study participants signed a written

informed consent for the collection of samples.

C57BL/6 f2r$^{fl/fl}$ Mice

The mouse carrying a conditional floxed (fl) PAR1 (C57BL/6 F2r$^{fl/fl}$) was constructed with

Ingenious Targeting Laboratories (NY, USA). Exon 2 of F2r gene was conditionally targeted and it is

fused in frame with p2A-Tomato reporter cassette. F2r and tdTomato genes are inactivated upon Cre

mediated deletion between the LoxP elements. A FRT/LoxP selection ivNeo marker as homology

arms were included in the final targeting vector.

Targeted iTL IN2 (C57BL/6) embryonic stem cells were microinjected into Balb/c blastocysts.

Resulting chimeras with a high percentage black coat color were mated to C57BL/6 FLP mice to

generate Somatic Neo Deleted mice. Confirmed somatic neomycin cassette deleted mice were bred

with wild-type C57BL/6N mice to generate germline neomycin cassette deleted floxed PAR1 mice

C57BL/6 f2r$^{fl/fl}$.

All mice were bred and maintained under specific pathogen-free conditions at Georgetown

University Division of Comparative Medicine facility. All experimental procedures were conducted

in accordance with the guidelines and an approved protocol by Institution Animal Care and Use

Committee at Georgetown University. Mice between 8 to 12 weeks old were used for experiments.

RT-qPCR analysis

qPCR was performed to determined mRNA expression of PARs in resting murine cells. RNA was

obtained from magnetic isolated B, CD4 and CD8 T cells (Miltenyi Biotech, MD). RNA was extracted

and purified using RNeasy Mini kit (Qiagen, MD). RNA was reverse transcribed into cDNA using

QuantiNova Reverse Transcription kit (Qiagen, Germany). qPCR performed with QuantiNova SYBR

Green PCR kit in CFX Opus Real-Time PCR system (Bio-Rad, CA). Primers used: murine *F2r*

(NM_010169.3):   forward,   5′-ATGAAAGCGTCCTGCTGGAG-3′;   reverse,   5′-

GGACGTTCAGAGGAAGGCTG-3′;   murine   *F2rl1*   (NM_007974.4):   forward,   5′-

GCTGGGAGGTATCACCCTTCT-3′;   reverse,   5′-CGCAGAGAACTCATCGATGGA-3′;   murine

F2rl2   (NM_010170.4):   forward,   5′-TGCCAGTCACTGTTTGCCAA-3′;   reverse,   5′-

CTCGGGACACTCCGCTTTTAT-3′;   murine   *F2rl3*   (NM_007975.3):   forward,   5′-

GACCCCCAGCATCTACGATG-3′; reverse, 5′-CAGCAGCAGTGCTTGAGAGCT-3′.

Housekeeping gene HPRT was used: murine HPRT (NM_013556): forward, 5′ -

GCGATGATGAACCAGGTTATGA- 3′; reverse, 5′ - ACAATGTGATGGCCTCCCAT - 3′

LCMV Glycoprotein: forward, 5′- CATTCACCTGGACTTTGTCAGACTC -3′; reverse, 5′-

GCAACTGCTGTGTTCCCGAAA -3′ (52, 53).

Ct of each sample obtained from the amplification curve was normalized to HPRT

and calculated ΔCt (Ct gene of interest – Ct HPRT). Results were expressed as $2^{-\Delta Ct}$ value.

LCMV infection

C57BL/6-F2r$^{fl/fl}$ mice were intravenously infected with 2×10^5 pfu LCMV Armstrong strain. At

days 6 and 21 post infection (p.i.), mice were anesthetized and perfused with cold HBSS before organs

(lung and spleen) were collected. *In vivo* CD8 T cell depletion was performed in LCMV infected mice

at day 15 p.i.. Mice were administered (i.p.) with 300 μg of anti-mouse CD8α mAbs (YTS 169.4,

BioCell, MA) or IgG2b isotype control (LTF-2, BioCell, MA). CD8 T cell depletion was confirmed

by flow cytometry in lung and spleen.

Cell suspensions from the spleens were obtained by smashing the spleens with a syringe plunger

in PBS and red blood cells were lysed with ACK (Gibco, MA) for 3 min on ice. After cell lysis cell

suspension was passed through a 70 μm mesh, and cell count and viability was assessed.

Cell suspension from the lungs were obtained by enzymatic digestion. Lungs were cut into small

pieces and digested using Lung Dissociation kit (Miltenyi Biotech, MD). The total cell suspensions

were passed through a 70 μm mesh, and cell count and viability were assessed. The total cell

suspension was used to analyze endothelial cells in the lung. For T cell analysis immune cell

enrichment was performed using 35% Percoll gradient. Total lung cell suspensions were centrifuged

and the pellet resuspended in 12 ml of 35% Percoll (Sigma, MA) and centrifuged for 15 min without

break at 1800 rpm (757g) at 7 °C. Immune cell suspensions were passed through a 70 μm mesh, and

cell count and viability were assessed.

Flow cytometry:

Phenotype lymph organs and endothelial cell phenotype: 2×10^6 of cells from spleen, lymph nodes

and lung cells (pre-Percoll) were stained with LIVE/DEAD (Invitrogen, CA) for 30 minutes at 4ºC.

Cells were incubated with a cocktail of mAbs (Table S1, panel-1 and panel-2) for 20 minutes at 4ºC.

Cell were washed with staining buffer (HBSS with 0.01% of NaN_3 and 0.01% of BSA) twice and

acquired with a BD Symphony (BD Biosciences, CA). Cells were analyzed with FlowJo (BD

Biosciences, CA).

Cytokine secretion assay: $2x10^6$ splenic and immune lung cells (post-Percoll) were stimulated *in*

vitro with 1 μg/mL of LCMVGp33-41 peptide (KAVYNFATC, Anaspec, CA) in the presence of 5

μg/mL of Brefeldin A (Sigma, MA) and 50 U/mL of rhIL-2 (National Cancer Institution, MD) for 4

hours at 37°C and 5% CO_2. After stimulation, cells were stained with LIVE/DEAD (Invitrogen, CA)

for 30 minutes at 4°C. Cells were washed twice and incubated with surface mAbs cocktail (Table S1,

panel-3) for 20 min at 4°C. Cells were fixed and permeabilized with BD Cytofix/Cytoperm solution

(BD Bioscience, CA) and incubated with the mAbs cocktail for intracellular markers for 20 min at

4°C (Table S2 panel-2). Samples were acquired in a BD Symphony and analyzed with FlowJo (BD

Biosciences, CA).

MHC class I pentamer staining: $1x10^6$ splenic and lung immune cells from infected mice were

stained with H-2D^b restricted LCMVGP33-41aa pentamer (H-2D^b-KAVYNFATC, Proimmune, UK)

at room temperature (22 °C) for 10 min followed by incubation with surface mAbs cocktail (Table S1)

for 20 min at 4°C. Cell suspensions were acquired in a BD Symphony (BD Biosciences, CA) and

analyzed using FlowJo.

CD8 T$_{ang}$ sorting and cytokine secretion: T cells were isolated from the lungs as described above.

Lung cell suspensions from naïve and infected mice at day 21 p.i. were stained with a cocktail of mAbs

for 20 minutes at 4°C (Table S1) and live cells were selected using Sytox Blue staining before sorting

(Biolegend, CA). CD31^{high} and CD31^{neg} CD8 T cells were sorted on a FACSAria (BD Bioscience,

CA). T cells were stimulated at a concentration of 300,000 cells/mL in plates with cross-linked anti-

CD3 and anti-CD28 mAbs (BD Bioscience, CA). After 3 days of culture, supernatants were collected

and cytokines (M-CSF, PDGF-A, GM-CSF, IL-2, IL-4, IL-5, IL-13, IL-17A, IL-6, IL-22, IL-9 and

IL-10) were measured using LEGENDplex (Biolegend, CA) following manufacturer's instructions.

Briefly, 25 μL of sample in duplicates were incubated with capture mAbs coated beads for 2 hours

followed by incubation with detection mAbs for 1 hour at room temperature in a shaker. Each sample

was incubated with streptavidin-PE for 30 min and acquired with a BD Symphony (BD Biosciences,

CA). Results were analyzed using cloud-based software Qognit (Qognit Inc, CA).

Flow cytometry human peripheral blood mononuclear cells

Whole blood from PWoH (n= 17) and PWH (n= 25) were collected in BD Vacutainer blood

collection tube containing EDTA (BD Bioscience, CA). Whole blood was diluted with PBS with 2%

FBS (1:1) and loaded on the top of Ficoll-Paque (Cytiva, MA) followed by a centrifugation at 400xg

for 30 min at room temperature (25°C). Peripheral blood mononuclear cells (PBMCs) were collected

and stained with LIVE/DEAD (Invitrogen, CA) for 30 minutes at 4°C. Cells were incubated with

1μg/mL human IgG (Sigma, MO) for 10 minutes at 4°C to block Fc receptors followed by incubation

with a monoclonal antibodies (mAbs) cocktail (Table S2, panel-1) for 30 minutes at 4°C. Samples

were acquired with a BD Symphony (BD Biosciences, CA) and analyzed using FlowJo software.

Measurements of cytokines and biomarkers of endothelial inflammation in plasma

Plasma were obtained from whole blood collected in BD Vacutainer blood collection tube

containing EDTA (BD Bioscience, CA) from PWH (n= 25) and stored at -80°C until use. Cytokines,

biomarkers of endothelial inflammation and granzymes (VCAM-1, IL-6, TNFα, IFNγ, sFasL, IL-2,

granzyme A and granzyme B) were measured using LEGENDplex (Biolegend, CA) following

manufacturer's instructions as described above.

Unsupervised clustering analysis

Multidimensional analysis was performed using Phenograph algorithm (FlowJo, CA). 10,000 of

manually gated lung endothelial cells (CD326$^-$CD45$^-$CD31$^+$) were concatenated from naïve mice and

infected mice at days 6 and 21 p.i.. Samples acquired in different days were normalized using Mutual

Nearest Neighbors (MNN) (FlowJo, CA) (54). Unsupervised clusters were analyzed with Phenograph

(k= 100) based on surface markers (CD31, PDL1, tdTomato, CD34, CD309, VCAM-1, CD105,

CD90.2 and CD201). Mean fluorescence intensity of each marker for all the clusters was presented in

heatmap and median frequency of each cluster from naïve and infected mice were expressed as bubble

plots.

Co-culture of primary human umbilical vein endothelial cells (HUVEC) and CD8 T_{ang} cells

Frozen PBMCs from PWoH (n= 10) and PWH (n= 10) were thawed and rested for 2 hours. Cells

were incubated with 1μg/mL human IgG (Sigma, MO) for 10 minutes at 4°C to block Fc receptors

followed by a cocktail of mAbs incubation for 30 minutes at 4°C (Table S2, panel-2). Live cells were

selected using Sytox Blue staining before sorting (Biolegend, CA). CD8 T_{ang} cells were sorted on a

FACSAria (BD, CA) and centrifuged at 1000 rpm for 30 minutes. Sorted CD8 T_{ang} cells were activated

with anti-CD3 and anti-CD28 mAbs coated beads (Miltenyi Biotech, MD) for 3 days and expanded in

the presence of 50 U/mL of recombinant human (rh) IL-2 (National Cancer Institution, MD) for 14

days. 5,000 HUVEC and 100,000 of CD8 T_{ang} cells were resuspended in endothelial cell growth basal

media (Lonza, CA) and seeded in the 96 well plate in the presence or absence of anti-CD3/anti-CD28

mAbs coated beads (Miltenyi Biotech, MD) and incubated at 37°C and 5% CO_2 overnight. As a

control, HUVEC alone was stimulated with 20 ng/mL of rhIFNγ (PeproTech, NJ) and 0.5 ng/mL of

rhTNFα (R&D System, MN) overnight. Cytokines in supernatant were measured using LEGENDplex

(Biolegend, CA) as described above.

Tube formation assay

Frozen PBMCs from PWH (n= 3) and PWoH (n= 2) were thawed and rested for 2 hours and sorted

as described above. Cells were incubated with 1μg/mL human IgG (Sigma, MO) for 10 minutes at

$4^{o}C$ to block Fc receptors followed by a cocktail of mAbs incubation for 30 minutes at $4^{o}C$ (Table S2,

panel 2). Live cells were selected using Sytox Blue staining before sorting (Biolegend, CA). CD8 T_{ang}

memory cells (CD45RA$^-$CD27$^+$) were sorted as described above. For HIV-Gag-specific CD8 T cells,

PBMCs from PWH were stimulated with HIV$_{gag}$ peptide pool (2 µg/mL, NIH AIDS Reagent Program)

overnight and CD137$^+$CD8 T cells were sorted. Sorted cells were washed twice with cold PBS and

labeled with 1.5 µM of CSFE (Invitrogen, CA) for 8 min at $37\,^{o}C$ followed by quenching with cold

FBS for 5 min on ice. CFSE labeled sorted cells were rested overnight. 10 µL of basement membrane

extract (Cultrex reduced growth factor, R&D System, MN) was applied in the inner well of 15 well

µ-slide chambered coverslip (ibidi, WI) and incubated for 20 min at $37°C$ for gelation. HUVEC and

CFSE labeled CD8 T_{ang} cells were resuspend in endothelial cell growth basal media (Lonza, CA) at

2:1 endothelial cell to T cell ratio and seeded on the top of gel in each well. Chambered coverslips

(Ibidi, WI) were incubated in a stage-top cell incubator at $37°C$ and 5% CO_2 and evaluated using Leica

SP8 live cell imaging (Leica, IL) at 4, 6, 8, 10 and 12 hours at 10X magnification. Analysis was

analyzed using image analysis platform Wimasis (Wimasis GmbH, Germany).

Calcium mobilization assay

60,000 HUVEC were cultured on a coverslip in endothelial cell growth media (Lonza, CA) for 8

hours and starved with endothelial cell growth basal media-2 (EBM-2) containing 1% PBS overnight.

Cells were loaded for 40 min with 1 µM Fluo-4-AM (Invitrogen, Thermo Fisher Scientific) in a buffer

solution containing 140 mM NaCl, 4 mM KCl, 1 mM $MgCl_2$, 1.2 mM $CaCl_2$, 10 mM HEPES, and 5

mM glucose (pH 7.3). Cells were imaged with 20X objective using a Nikon TE2000 microscope with

an excitation filter of 480 ± 15 nm and an emission filter of 535 ± 25 nm. The images were captured

by a Retiga 3000 digital camera (QImaging) and analysis was performed offline using ImageJ.

Granzyme- dependent PAR1 activation

100,000 HUVEC were seeded a 12 well plate in endothelial cell growth media supplemented with

EGM-2 bullet kit overnight. Next day cells were starved in EBM-2 containing 1% FBS for 6 hours.

HUVEC were incubated pre-incubated with 10 μg/mL of ATAP-2 (Invitrogen, CA) or 300 nM of

PAR1 antagonist SCH530348 diluted in EBM-2 containing 1% FBS for 20 min at room temperature

(25°C). After the preincubation period, supernatants were removed from the wells, and HUVEC were

stimulated with 100 nM of recombinant human granzyme A (rhGZMA, Kamiya Biomedical, WA)

and recombinant human granzyme K (rhGZMK, Kamiya Biomedical, Japan) with fresh inhibitors at

the same concentration. As the positive control, HUVEC were incubated with 100 μM of PAR1

agonist TFLLR-NH2 (Tocris Bioscience, UK) and control peptide RLLFT-NH2 (Tocris Bioscience,

UK). After overnight incubation at 37°C and 5% CO_2 supernatants were collected and cytokines were

measured using LEGENDplex (Biolegend, CA) as described above.

Statistical analysis

Comparisons were performed using non-parametric Mann-Whitney test or Wilcoxon test and

p value $\leq$ 0.05 was considered significant. Correlations were performed using nonparametric

Spearman correlation and *p* value $\leq$ 0.01 was considered significant.

Results

PAR1 expression is modulated in virus-specific CD8 T lymphocytes during acute LCMV

infection

At the site of vascular injury mediated by trauma or infection, the cross-talk between the

coagulation cascade, platelets, endothelial cells and CD8 T cells is mediated by thrombin

dependent activation of PAR1 (32, 38, 40, 55). Particularly, PAR1 signaling is involved in CD8 T

cell trafficking and function (40, 41). However, whether PAR1 is modulated in CD8 T cells

during the immune response remains to be defined.

We generated a PAR1 reporter mouse that allows us to evaluate PAR1 expression in CD8 T

cells during the course of a systemic viral infection and better understand the interplay between

CD8 T cells and endothelial cells. C57BL/6 F2r$^{fl/fl}$ mouse carrying floxed Exon 2 of *F2r* gene was

conditionally targeted as well as fused in frame with p2A-Tomato (tdT) reporter cassette (as

described in Material and Methods section and Figure S1A).

C57BL/6 F2r$^{fl/fl}$ mice showed similar cell numbers in lymph nodes and spleen compared with

the wild type C57BL/6 (Figure S1B). The frequencies and cell count of innate cells including

neutrophils, eosinophils, monocytes, macrophages and dendritic cell subsets were similar between

the C57BL/6 F2r$^{fl/fl}$ and the C57BL/6 wild type mice (Figure S1C and D). Similarly, no significant

differences between C57BL/6 F2r$^{fl/fl}$ and wild type mice were observed in the frequencies and cell

numbers of lymphocytes subsets, B, NK and T cells in lymphoid organs (Figure S1E).

PAR1 expression, tdTomato^{+} (tdT^{+}) cells, included CD4 and CD8 T cells and NK cells and it

was negligible in B cells (Figure 1A, B and C). Similar to our previous observations in human T

cells, CD8 T cells expressed higher levels of PAR1 compared with CD4 T cells. Among

lymphocytes, NK expressed the highest level of PAR1 at steady state conditions (Figure 1C) (41).

mRNA of *F2r* expression in tdT expressing cells was further confirmed by qPCR in magnetic

cell isolated CD4 and CD8 T cells and B cells. CD8 T cells expressed higher levels of mRNA of

F2r than CD4 T cells (Figure 1D). In addition, both CD4 and CD8 T cells expressed mRNA *F2rl1*

(PAR2) but not *F2rL2* (PAR3) and *F2rl3* (PAR4) (data not shown)(41). PAR1 protein expression

in CD4 and CD8 T cells was further confirmed by flow cytometry in tdT$^+$ T cells (Figure S2).

These data show that at steady state conditions, PAR1 is expressed in T and NK but not B

lymphocytes.

We next investigated the expression of PAR1 in virus-specific CD8 T cells during a systemic

acute viral infection with the Lymphocytic Choriomeningitis virus Armstrong strain (LCMVArm).

This infection model recapitulates features of human viral infections, the acute form, LCMV

Armstrong (Arm) is cleared 8-10 days post infection (p.i) and induces the generation of LCMV-

specific memory CD8 T cells (56-58). LCMV is a non-cytolytic virus and the pathology induced

is mediated by the host immune response (59, 60).

Using this model, we aimed to determine the immune mediated endothelial injury in the lung,

and address whether T cells are involved in endothelial repair after viral control. We studied two

time points of the LCMVArm infection, day 6 p.i (effector phase) and day 21 p.i (contraction phase

and early memory development) (58, 61).

LCMVGp33-specific CD8 T cells secreted IFNγ, TNFα and showed cytotoxic potential (as

measured by CD107a surface expression) upon *in vitro* stimulation with LCMVGp33-41aa peptide

(Figure S3A and B). LCMVGp33-specific CD8 T cells peaked by day 6 p.i in both spleen and

lung and declined by day 21 in the spleen when the virus is controlled (Figure S3A and B). In

contrast, in the lung, LCMVGp33-specific CD8 T cells remain higher consistent with development

of memory cells (Figure S3A and B).

Expression of PAR1 was monitored by tdT$^+$ cells in total CD8 T cells and virus-specific CD8

T cells using pentamer staining (H2-D^{b}LCMVGp33-41aa) (Figure S3C and Figure 1E

respectively). At day 6 p.i. and compared to total CD8 T cells from the naïve (dashed line) mice,

expression of PAR1 (tdT$^+$) in LCMVGp33-specific CD8 T cells was reduced, and it was

significantly upregulated by day 21 p.i. (Figure 1E and F). Similar observations were noted in total

CD8 T cells in both spleen and lung (Figure S3C). PAR1 transcriptional regulation in

LCMVGp33-specific CD8 T cells may be induced by *in vivo* granzyme activation during

degranulation (Figure S3B) (41). Accordantly, tdT expression by CD107a$^+$CD8 T cells showed

lower expression of granzyme A compared to CD107a$^-$CD8 T cells (Figure S3D and E). In

addition, tdT$^+$CD8 T cells negatively correlated with the frequency of GZMA$^+$CD8 T cells, R= -

0.61, *p*= 0.02.

Altogether these data show that PAR1 is expressed at distinct stages of CD8 T cell

differentiation. *In vivo*, PAR1 expression is downregulated at the effector phase and upregulated

after viral control and memory development.

Angiogenic CD31^{high}CD8 T (T$_{ang}$) cells promote endothelial recovery during acute viral

infection

T$_{ang}$ cells have been originally described in the center of the endothelial progenitor cell colonies

isolated from peripheral mononuclear cells (PBMCs). In *in vivo* experiments, T$_{ang}$ cells support

new vessel formation (26). However, whether T$_{ang}$ cells are virus-specific CD8 T cells that acquire

these properties once the infection is controlled and are involved in endothelial repair remains

poorly understood.

We hypothesized that systemic LCMVArm infection induces immune mediated endothelial

injury in the lung, and T_{ang} cells play a role in repair after viral control. We found that at day 6 p.i.,

when LCMV-specific CD8 T cells are recruited into the lung, there was a significant reduction in

the frequency and cell count of lung endothelial cells (ECs) (Figure 1G). ECs recovered to baseline

levels by day 21 p.i. (Figure 1G). In contrast to CD8 T cells, expression of PAR1 in endothelial

cells increased over the course of infection suggesting that PAR1 can be targeted by different

proteases during the course of the infection (Figure 1H). These data show that immune mediated

endothelial cell injury occurs in lung during a systemic viral infection and recovers upon viral

control.

We next determined if CD8 T cells are involved in the recovery of lung ECs. LCMV infected

mice were either treated with anti-CD8 mAb and IgG control at day 15 p.i. when the infection is

controlled. CD8 T cells were efficiently depleted in the lung by day 21 p.i. (Figure 2A). Compared

to the mice treated with IgG control, CD8 T cell depleted mice showed a delayed recovery of EC

count suggesting that CD8 T cells are involved in this process (Figure 2B). At this time point of

infection, we did not detect LCMV Glycoprotein by qPCR in both spleen and lung suggesting viral

control (Figure S4A) (62).

We next tested the hypothesis that the delayed recovery of lung ECs is associated with the loss

of CD8 T_{ang} cells during depletion. T_{ang} cells have been defined by expression of CD31 (26). In

the lung, during the course of the LCMV infection, a small subset of CD8 T cells expressed

CD31^{high} and CD309 (vascular endothelial growth factor receptor 2 (VEGFR2)) and although the

frequency showed an increased trend by day 21 p.i., did not rich statistical significance (Figure

2C). In contrast, CD31^{high}CD8 T cell count was significant increased suggesting an expansion of

this subset during the recovery phase of lung ECs (Figure 2C). The expression of PAR1 of the

CD31^{high} and CD31^{low} CD8 T cells followed the similar kinetics to that observed in LCMVGp33-

specific CD8 T cells (Figure S4B and Figure 1F).

We then investigated whether CD31^{high}CD8 T cells were LCMV-specific and secrete

cytokines upon *in vitro* stimulation with LCMVGp33 peptide (Figure 2D). At day 21 p.i. and

compared to CD31^{neg}CD8 T cells, CD31^{high}CD8 T$_{ang}$ cells secrete similar levels of IFNγ, TNFα

and have similar cytotoxic potential (CD107a). Of note, the frequency of CD31^{high}LCMVGp33-

specific CD8 T cells single IFNγ producers were reduced compared to the CD31^{neg} (Figure 3D).

Interestingly, LCMVGp33-specific CD31^{high}CD8 T cells were enriched in the memory

precursor effector T cells (KLRG1^{-}CD127^{+}), which have survival advantage during the

contraction phase and possesses plasticity to give rise to multiple memory subsets (Figure 2E)

(63-66). As expected, at day 21 p.i., a reduction of double positive effectors (KLRG1^{+}CD127^{+})

and short-live effector KLRG1^{+}CD127^{-} and KLRG1^{-}CD127^{-} CD8 T cells was observed at day

21 p.i. (Figure 2E).

In addition, we evaluated the ability of lung CD31^{high}CD8 T cells to secrete pro-angiogenic

cytokines and T cell helper cytokines. We sorted CD31^{high} and CD31^{neg} bulk CD8 T cells from

lung of LCMV mice infected at day 21 p.i., and CD31^{high} bulk CD8 T cells from lung of naïve

mice as control. Sorted subsets were TCR stimulated with cross-linked CD3 and CD28 mAbs for

3 days and cytokines were measured in the supernatants (Figure 2F and Figure S4C). CD31^{high}

but not CD31^{neg}CD8 T cells secreted vascular endothelial growth factor (VEGF), a cytokine that

induces proliferation of ECs suggesting a pro-angiogenic function (67, 68). In addition, compared

to the CD31^{neg}, CD31^{high}CD8 T cells showed an increased trend, although not significant, in the

ability to secrete granulocyte-macrophage-colony stimulating factor (GM-CSF), known to induce

EC proliferation and mobilize endothelial progenitor cells from the bone marrow (Figure S4C)

(69, 70). Lastly, CD31^{high}CD8 T cells showed the ability to secrete of IL-2 and secrete low levels

of Th2, Th17, Th22 and Th9 associated cytokines (Figure S4C).

These data suggest that angiogenic CD31^{high}CD8 T$_{ang}$ cells are virus specific with an enriched

memory precursor phenotype. In addition, CD31^{high}CD8 T$_{ang}$ secrete pro-angiogenic cytokines

contributing to lung EC recovery after viral control.

Endothelial recovery is associated to CD31^{high}CD8 T$_{ang}$ cells

We next investigated the relationships between CD8 T$_{ang}$ cells and lung ECs during the course

of the LCMV infection. ECs are heterogenous and play important roles by supporting immune cell

trafficking and regulating inflammation during immunity (71-73). Diverse phenotypes with

distinct functions including "immune cell-like" and "developmental/regenerative" involved in

immunity and repair respectively have been described (74, 75).

We hypothesized that during the expansion of CD31^{high}CD8 T$_{ang}$ cells and recovery EC phase

ECs will be enriched with a regenerative phenotype. We determined the expression of markers

associated with inflammation (PD-L1, VCAM1) and pro-angiogenic/repair (CD34, CD309,

CD105) (60, 74, 76, 77). CD309 is a receptor for the VEGF, an essential factor for EC function

during angiogenesis (67). CD105 (Endoglin) binds to different components of the TGF-β

superfamily and also regulates endothelial proliferation, migration, and vascular homeostasis (78,

79). In addition, CD90.2 was used as maker for lymphatic endothelial cells (80).

We performed an unsupervised cluster analysis using PhenoGraph to classify high-parameter

single-cell data into phenotypically distinct clusters (81). We concatenated 10,000 events per

mouse of CD326$^-$CD45$^-$CD31$^+$ lung ECs from Naïve and LCMV infected mice from day 6 and 21

p.i. to analyze the dynamics of endothelial cells (Figure 3A).

We found that the frequency of lung EC clusters (CL) 1, 2, 3, and small CL 7, and 8

significantly increased at day 6.p.i (Figure 3A, bubble plots and Figure S5). These CLs expressed

higher levels of PD-L1 and distinct levels of CD31 and PAR1 (Figure 3A). Expression of PD-L1

suggest that endothelial cells interact with immune cells modulating immune mediated pathology

through PD1-PD-L1 interaction (82).

The CL4, 5, 6, and 13 were present in the naïve mice and at day 21 p.i. (Figure 3A and Figure

S5). These clusters expressed markers associated with EC proliferation including CD105

(Endoglin), and CD309 (VEGFR2), A small EC subset (CL14), expressed high levels of CD105

and was expanded by day 21 p.i. suggesting a "regenerative" phenotype (Figure S5). In addition,

CL10 and CL11 remained significantly reduced compared to the naïve mice. CL9 correspond to

CD90.2$^+$ lymphatic endothelial cells and its frequency remained constant through the course of the

infection.

These data suggested that ECs are heterogenous and dynamic during viral infection. The ECs

in CL6 expressed CD201 (endothelial protein C receptor, EPCR), a marker associated with

resident vascular endothelial cell progenitors and activated protein C (APC) dependent activation

of PAR1 during angiogenesis (75, 83, 84). More importantly, CL6 expressed higher levels of

CD105, CD309 and CD34 suggesting a proliferative and regenerative phenotype (76).

Interestingly, CL6 was positively associated (R= 0.66, $p<$ 0.01) with the frequency of

CD31^{high}CD8 T$_{ang}$ cells suggesting potential interaction during vascular remodeling (Figure 3B).

Altogether these data suggest that CD31^{high}CD8 T$_{ang}$ cells play a role in endothelial remodeling

during infection.

Circulating angiogenic CD8 T (T$_{ang}$) cells from PWH have pro-inflammatory phenotype.

444  The data above shows that during a systemic viral infection the host immune response mediates

445 endothelial injury and CD31^{high}CD8 T_{ang} cells play a role in endothelial repair by secreting pro-

446 angiogenic cytokines involved in endothelial proliferation and remodeling.

447  The role of T_{ang} cells in the setting of human chronic viral infections such as HIV remains to

448 be defined. Previously, we reported that circulating T_{ang} cells from PWH express the vascular

449 homing receptor CX3CR1 suggesting their recruitment to the sites of endothelial inflammation

450 (85). Given that immune activation and systemic inflammation are the main features of chronic

451 HIV infection, we hypothesized that this environment alters T_{ang} cell function promoting a pro-

452 inflammatory function that contributes to endothelial inflammation.

453  To address this question, we investigated circulating T_{ang} cells in PBMCs from PWH (n= 25)

454 and uninfected controls People without HIV (PWoH, n=17). PWH had a median CD4 count 663

455 (532-861.5) cells/µL and HIV RNA viral load < 20 copies/mL (n= 20) and 50-110 copies/mL (n=

456 4). Study participants had a median ASCVD (Atherosclerosis Cardiovascular Disease) risk score

457 of 7.9 (IQR: 5.0-18.6) (Table 1).

458  In humans, T_{ang} cells are defined as surface expression of CD31/PECAM-1 (Platelet

459 Endothelial Cell Adhesion Molecule) and the chemokine receptor CXCR4 (receptor for the

460 Stromal Cell Derived Factor-1) *(26, 86)*. The flow cytometry analysis showed that in PWH there

461 is a significantly reduction in the frequencies of circulating CD8 T_{ang} cells (*p*< 0.001). A trend

462 although not statistically significant was observed in CD4 T_{ang} cells as well (Figure 4A and B).

463  To evaluate the impact of HIV driven immune activation, we studied expression of markers

464 associated with adhesion, homing and trafficking during vascular inflammation including the

465 Protease Activated Receptor 1 (PAR1) which is involved in the recruitment of T cells at the site

466 of vascular injury (38, 40, 41). In addition, we investigated the vascular homing receptor CX3CR1

that binds to CX3CL1 (fractalkine) expressed by inflamed endothelial cells, and the integrin

CD49f (Integrin alpha-6) that binds to laminin, a component of the vascular basement membrane

(38-40, 87).

In PWH, both CD4 T_{ang} and CD8 T_{ang} cells showed significantly increased expression of

PAR1, CX3CR1 and CD49f compared to PWoH suggesting an increased potential for recruitment

and trafficking to the sites of vascular inflammation (Figure 4C and D). In addition to the increased

trafficking potential, both CD4 T_{ang} and CD8 T_{ang} cells from PWH showed an increased expression

of effector molecules Granzyme A (GZMA) and Granzyme B (GZMB) when compared to PWoH

(Figure 4E).

All together these data suggest that PWH and virally suppressed by ART have a significant

reduction in the frequencies of CD8 but not CD4 T_{ang} cells. T_{ang} cells showed a differentiated

phenotype with cytotoxic potential and increased potential to migrate to the sites of vascular

inflammation.

Frequencies of activated CD8 T_{ang} cells are associated with cardiovascular risk in PWH

In PWH, immune activation, systemic inflammation and endothelial dysfunction have been

identified as the main drivers of cardiovascular events (4, 9-11, 88). The data above suggest that

in the context of HIV infection T_{ang} cells have a pro-inflammatory phenotype. We next investigated

the relationship between activation markers expressed by T_{ang} cells and the Atherosclerotic

Cardiovascular Disease (ASCVD) risk score (Figure 4F, G and H).

The frequencies of circulating total CD4 and CD8 T_{ang} cells did not show an association with

ASCVD (data not shown). In contrast, we observed a positive correlation between ASCVD and

the frequency of cells expressing activation markers including homing and trafficking

CX3CR1$^+$PAR1$^+$CD8 T$_{ang}$ and CD49f$^+$PAR1$^+$CD8 T$_{ang}$ (R= 0.50, p= 0.04 and R= 0.58, p= 0.02

respectively) (Figure 4F and G). Similarly, GZMA$^+$GZMB$^+$CD8 T$_{ang}$ cells showed a positive

association with the ASCVD risk score R= 0.65, p< 0.01 (Figure 4H). Of note is that expression

of these markers in CD4 T$_{angs}$ cells did not correlate with ASCVD risk (Figure 4F, G and H). These

observations suggest that in PWH immune activation of CD8 T$_{ang}$ cells drives a pro-inflammatory

phenotype and may play a role in sustaining endothelial inflammation.

To better address the potential contribution of activated CD8 T$_{ang}$ cells on endothelial

inflammation, we further studied the relationship between plasma levels of biomarkers of

endothelial inflammation and immune cell trafficking including soluble Vascular and Intracellular

Adhesion Molecule-1 (sVCAM-1 and sICAM-1 respectively), IL-6; and T cell-derived cytokines

and effector molecules including TNFα, IFNγ, IL-2, sFASL, GZMA and GZMB (Figure 4I and

Figure S6A). We found that plasma levels of GZMA but not GZMB were positively associated

with IL-6 (R= 0.62, p< 0.001). sVCAM-1 showed a positive trend (R= 0.449, p= 0.024) but not

with sICAM-1 (Figure 4I). As expected, plasma levels of cytokines (TNFα, IFNγ, sFASL, IL-2)

positively correlated with the plasma levels of granzymes (Figure S6A).

Altogether the data shows that chronic HIV driven immune activation increase trafficking and

pro-inflammatory properties of CD8 T$_{ang}$ cells. Moreover, the association with ASCVD suggest

that CD8 T$_{ang}$ cells may contribute to endothelial inflammation, dysfunction and disease risk.

***In vitro*, CD8 T$_{ang}$ cells from PWH showed enhanced ability to induced sICAM-1 and**

VCAM-1 by endothelial cells.

The above results suggest that chronic HIV infection alters CD8 T$_{ang}$ cells skewing their

function to pro-inflammatory. Because activated CD8 but not CD4 T$_{ang}$ cells showed association

with cardiovascular risk in the following sections of the manuscript we focused our study on CD8

T_{ang} cells.

To address the potential mechanisms involved by which CD8 T_{ang} cells drive endothelial

inflammation, we first determined the effects of *in vitro* activated CD8 T_{ang} cells on primary

Human Umbilical Vein Endothelial cells (HUVEC).　In the supernatants of the co-cultures, we

measured adhesion molecules including sICAM-1 and sVCAM-1, pro-inflammatory cytokines IL-

6, Angiopoetin-2 (Ang-2) and IL-8; and pro-angiogenic cytokines, basic Fibroblast Growth Factor

(bFGF) and Vascular Endothelial Growth Factors (VEGF) (89-91).

CD8 T_{ang} cells from PWH (n= 10) and PWoH (n= 10) were sorted based on the expression of

CD31, CXCR4 and memory phenotype. CD8 T_{ang} cells were *in vitro* expanded with anti-

CD3/CD28 mAbs and rhIL-2 (50 U/ml) for 14 days. Expanded CD8 T_{ang} cells were co-cultured

with HUVEC at 20:1 CD8 T_{ang} to HUVEC ratio and stimulated in media (control) and with anti-

CD3/CD28 mAbs coated beads overnight.

CD8 T_{ang} co-cultured with HUVEC cells in media alone showed negligible levels of shedding

of sICAM-1 and sVCAM-1. In contrast, when CD8 T_{ang} cells were TCR stimulated a significant

increase of sICAM-1 and sVCAM-1 was detected in the supernatant of the cultures (Figure 5A).

The effects mediated by CD8 T_{ang} cells from PWH were significantly higher than those observed

with CD8 T_{ang} cells from PWoH (Figure 5A). Moreover, stimulation of CD8 T_{ang} from PWH led

to an increased trend, although did not reach statistical significance, in the production of pro-

inflammatory and pro-angiogenic cytokines IL-6, and bFGF respectively. No differences were

observed in the secretion of IL-8 Angiopoietin-2 (Ang-2) and VEGF. (Figure 5B). The

combination of cytokines IFNγ (10 ng/mL) and TNFα (0.5 ng/mL) used as positive control in

HUVEC cultured alone, showed no statistically significant effect on the shedding of sVCAM-1,

sICAM-1. In addition, low IL-6 secretion was observed suggesting that CD8 T_{ang} cell-HUVEC

contact dependent mechanism is involved in mediating shedding of adhesion molecules and IL-6

secretion by endothelial cells (Figure S6B and Figure 5B). Similar levels of Ang-2, IL-8/CXCL8

and the pro-angiogenic cytokines VEGF and FGFb were detected when endothelial cells were

stimulated with combination IFNγ and TNFα (Figure S6B).

These observations suggest that secretion of cytokines and effector molecules from activated

CD8 T_{ang} cells can lead to endothelial activation promoting trafficking of immune cells and

secretion of pro-inflammatory and pro-angiogenic cytokines by endothelial cells.

We next performed a tube forming assay in which endothelial cells cultured in an extracellular

matrix form capillary-like structures as a measurement of angiogenesis to better evaluate the

interactions between CD8 T_{ang}-endothelial cells (92). CD8 T_{angs} cells were sorted from PBMCs,

labeled with CFSE and cultured with HUVEC at 1:2 T cell to HUVEC ratio (Figure 5C and D).

CD8 T_{ang} cells overtime migrate and adhere to the endothelial cells as the tubes formed

showing a mesh of capillary-like structures and $CFSE^+CD8$ T_{ang}, suggesting a close interaction

with endothelial cells in the extracellular matrix (Figure 5C and D). Overtime, $CSFE^+CD8$ T_{ang}

from PWH and sorted HIV_{Gag} specific CD8 T cells showed a trend of slowing the growth of the

tube (loop area), although these responses were heterogenous among the donors (Figure 5E). In

contrast, CD8 T_{ang} from PWoH showed a trend promoting bigger loop area overtime.

Altogether, these data imply a cross-talk between CD8 T_{ang} cells and endothelial cells. CD8

T_{ang} cells from PWH promote endothelial activation (ICAM1 and VCAM1), and induced secretion

of pro-inflammatory and pro-angiogenic cytokines influencing endothelial cell function.

Activated CD8 T$_{ang}$ cells mediate endothelial cell activation through a granzyme-dependent PAR1 activation

We next investigated the mechanisms by which CD8 T$_{ang}$ cells activate endothelial cells. PAR1 is expressed by endothelial cells and its signaling mediates important functions in maintaining barrier function (32). In addition, PAR1 connects endothelial cells and CD8 T cells at the site of vascular injury (38, 40, 41). The above data shows that CD8 T$_{ang}$ cells express PAR1 and endothelial homing markers, and GZMA$^+$GZMB$^+$CD8 T$_{ang}$ cells are associated with ASCVD (Figure 4). These observations and the evidence that in addition to thrombin, other proteases including granzymes (A, B and K) activate PAR1, led us to the hypothesis that granzymes are involved in endothelial activation in the context of HIV infection (93-95).

GZMs were first thought to be mainly involved in cytotoxic function, however non-cytotoxic and extracellular biological activities including regulation of vascular integrity, angiogenesis, extracellular matrix remodeling have been reported (93, 96, 97). In contrast to thrombin, granzymes cleavage site and the downstream signaling pathways differ from thrombin leading to a non-canonical activation of PAR1 called biased signal (43-50, 98, 99).

We next studied the effects of granzymes in endothelial cells. To isolate the effects of granzymes from cytokines secreted by CD8 T$_{ang}$ cells we used recombinant human (rh) GZMA and GZMK. GZMA and GZMK have tryptase activity and preferentially cleave after basic residues (Arg and Lys) which is similar to thrombin. GZMB is an aspartase and cleaves after Asp (49, 50, 98, 99).

We determined the effects of GZs dependent PAR1 activation on calcium mobilization. HUVEC were cultured *in vitro* and activated either with rhGZMA (100nM) and Thrombin (3nM). Compared to thrombin, GZMA induced a calcium mobilization with slower kinetics and sustained

overtime (Figure 6B). These effects were not observed on GZMK and GZMB (data not shown).

We then investigated the downstream effects induced by GZM-dependent activation of PAR1 at

later time points of stimulation and measure cytokine secretion by endothelial cells (Figure 6C).

HUVEC were activated with either rhGZMA and rhGZMK (100 nM) in the absence or

presence of PAR1 antagonist SCH530348 which inhibit binding of the tethered ligand (Figure 6A).

In addition, we used the mAb ATAP-2 that blocks PAR1 cleavage by binding to an epitope within

the PAR1 tethered ligand domain, Figure 6A (43, 100-102). As positive control, we used an agonist

peptide specific for PAR1 (mimics the tethered ligand) and mediates activation of PAR1 in a

cleavage independent manner (Figure 6A).

We found that rhGZMA and rhGZMK induced secretion of pro-inflammatory cytokines IL-

6, IL-8/CXCL8 and Ang-2 (Figure 6 and Figure S6C). Moreover, GZA and GZK induced pro-

angiogenic cytokine VEGF, albeit at lower levels (Figure S7). The antagonist SCH530348

(vorapaxar) partially blocked PAR1 activation by rhGZMA and consequently the secretion of IL-

6 and IL-8 and Ang-2 (Figure 6C, D and E). The effects mediated by the ATAP-2 mAb showed

inhibition Ang-2 but not IL-6 and IL-8.

In addition, rhGZMK dependent PAR1 activation by the antagonist SCH530348 inhibited

secretion of IL-6 and IL-8/CXCL8 but not Ang-2 secretion by endothelial cells (Figure 6C, D and

E). In contrast, ATAP-2 mAb blocked GZMK dependent PAR1 activation and the secretion of

IL-6, IL-8/CXCL8 and Ang-2 (Figure 6C and S7). As expected, stimulation with PAR1 agonist

peptide (cleavage independent activation) induced secretion of IL-6 and IL-8, however it did not

induce a significant increase in secretion of Ang-2 (Figure 6H and S6C).

All together these data suggest that CD8 T$_{ang}$ cells mediate endothelial cell activation in part

through GZMA and GZMK dependent PAR1 activation leading to the secretion of pro-

inflammatory cytokines.

Discussion

Chronic CD8 T cell activation is associated with increased cardiovascular disease risk in PWH, however the mechanisms remain to be defined (10, 11, 18, 22, 23). Because endothelial inflammation and dysfunction are the main features of cardiovascular disease, in the present study we investigated a physiological mechanism in which the immune system (with a focus on CD8 T cells) intersects with the homeostasis of the endothelium such as the vascular repair mechanism in the setting of a systemic viral infection.

In this study, using a PAR1 reporter mouse we established the dynamics of PAR1 expression in virus-specific CD8 T cells during the course of an acute viral infection with LCMV. We showed that PAR1 expression in LCMV-specific CD8 T cells is downregulated during the effector phase and its expression is upregulated during the contraction phase and early memory development (day 21 p.i). In addition, we determined that a subset of CD8 T cells expressing CD31^{high} play a role endothelial repair providing new evidence of the important role of CD8 T_{ang} cells in vascular repair and remodeling in the setting of a viral infection.

T_{ang} cells were originally discovered in the center of endothelial progenitor cell colonies from PBMCs by Hur J. et al. (26). Vascular repair is mediated in part by T_{ang} cells that support proliferation and differentiation of bone marrow derived and resident endothelial progenitor cells/endothelial colony forming cells (EPC/ECFC) at the site of injury (26-28). However, the type CD4 and CD8 of T cells involved in angiogenic function and whether they are virus-specific T cells in the setting of infection has not been elucidated.

In the present study, we investigated endothelial injury and repair during acute LCMV infection. We found that expansion of CD31^{high}CD8 T cells occurs concomitant with the EC recovery and TCR stimulation of CD31^{high}CD8 T cells showed secretion of pro-angiogenic

cytokines VEGF and GMCSF. These cytokines are associated with endothelial cell proliferation,

survival and bone marrow mobilization of endothelial progenitors (67-69, 103). Accordantly,

previous reports showed that administration of GMCSF and granulocyte colony stimulating factor

(G-CSF) induced mobilization of pro-angiogenic cells in patients with acute myocardial infarction

(69, 104). Moreover, mobilization endothelial progenitor cells by GMCSF enhanced endothelial

regeneration and reduced vascular inflammation after intravascular radiation (105, 106). Our study

shows that CD31^{high}CD8 T$_{ang}$ cells may regulate angiogenesis by promoting the recruitment of

endothelial progenitor cells, and other cells including myeloid cells to the site of vascular

remodeling as depletion of total CD8 T cells delay the recovery of ECs.

Particularly, CD31^{high}CD8 T$_{ang}$ cells expressed CD309 (VEGFR2) and secrete pro-angiogenic

and Th1 cytokines. Similar to these findings, a study reported that hypoxia upregulates expression

of CD309 (VEGFR2) and VEGF signaling, which in turn increaseed expression of IFNγ and

inhibited IL-10 (68). In our model of a systemic viral infection with LCMV, immune mediated

pathology may induce hypoxia driving expression of CD309 and secretion of VEGF by CD8 T$_{ang}$

cells. It is also possible VEGF signal promotes the acquisition of a pro-angiogenic property by

CD8 T$_{ang}$ cells and warrants further investigation.

CD8 T cells are predominantly viewed as main players in immunity against viruses and cancer

due to their cytotoxic function, however, emerging evidence suggest that they play a role in tissue

regeneration and remodeling. CD31^{high}CD8 T$_{ang}$ cells were LCMV-specific and secrete VEGF,

IFNγ and TNFα and degranulate upon *in vitro* stimulation LCMVGp33 suggesting that may be a

function of memory cells after viral control. More importantly, CD31^{high}CD8 T$_{ang}$ cells showed a

memory precursor phenotype (KLGRG1^{-}CD127^{+}) suggesting a survival advantage during the

contraction phase and a proliferative potential since they produce IL-2 (63-66).

Expression of CD31 in T cells has been associated with T cell trafficking and studies have

shown that CD31 attenuates TCR signaling in T cells as contains two immunoreceptor tyrosine-

based inhibitory motif (ITIM) units in the cytoplasmic domain (107-109). It is possible that CD31

expression in CD8 T_{ang} cells allows close interactions with endothelial cells preventing their

potential cytotoxic activity on endothelial cells.

CD8 T cells with tissue remodeling properties that secrete IFNγ, TNFα and epidermal growth

factor receptor (EGFR)-ligand amphiregulin (AREG) have been described highlighting CD8 T cell

function beyond cytotoxicity (110). Moreover, in pathological conditions such as retinal vascular

diseases, VEGF producing CD8 T cells drive inflammation and abnormal angiogenesis suggesting

that dysregulated T_{ang} cell function may contribute to the pathology of disease (111).

We also investigated changes in lung endothelial cells. ECs are heterogenous and play distinct

roles during immune response by supporting trafficking of immune cells, immune functions

(immune cell-like ECs) and repair (regenerative ECs) (73, 112). Particularly, our study shows that

at the peak of CD8 T cell recruitment into the lung, ECs expressed high levels of PD-L1 likely

induced by cytokines produced by LCMV-specific CD8 T cells regulating the immune mediated-

pathology including endothelial injury (113). The "immune-EC" phenotype returns to basal levels

when the virus is controlled and ECs expressing receptors associated with cell proliferation and

survival (CD34, CD309, CD105, CD201) are more predominant.

Studies of lung endothelial cells in response to acute lung injury by LPS and influenza infection

using single cell RNAseq, revealed distinct functions of ECs "immune-ECs" and "regenerative-

ECs" the later involved in vascular remodeling suggesting an important cross-talk between ECs,

host immunity and endothelial repair (74, 76). In contrast to these studies that used an acute

respiratory lung viral infection or injury, we used a systemic virus to evaluate vascular injury and

investigate potential mechanisms associated with chronic HIV infection and endothelial

inflammation. Similar to our studies, the authors observed a trend of reduced EC count at day 7

p.i, (76). Although this study did not investigate the potential mechanisms driving proliferation of

ECs in the reconstitution phase (between day 14 to 21 p.i), they described a proliferating

population of CD34+ECs with regenerative capacity (76). We found that CD8 T_{ang} cells play a role

in this phase and are associated with population of ECs (cluster 6) that express CD34 and have the

potential to proliferate in response to pro-angiogenic cytokines through CD309 and CD105

signaling. CD309, the receptor for VEGF, is a critical factor for endothelial cell proliferation,

migration, and survival, and plays important role during vascular remodeling (114). In addition,

CD105, a marker for endothelial cell proliferation, is expressed on vascular endothelium during

angiogenesis and prevents endothelial cell apoptosis induced by TGFβ (115, 116). Moreover, this

EC population express of endothelial protein C receptor (CD201) that has been proposed as a

marker of endothelial progenitor cells with new vessel forming properties (83, 117, 118).

T_{ang} cells are key players in maintaining endothelial homeostasis and alterations in vascular

repair mechanisms may contribute to endothelial dysfunction and increase cardiovascular disease

risk (119). In fact, reduced proportion of circulating T_{angs} has been associated with age and

increased cardiovascular risk (26, 29, 30). Patients with rheumatoid arthritis and inflammatory

diseases that suffered a cardiovascular event complication had lower circulating T_{ang} cells (30, 86,

120-124).

The role of T_{ang} cells in the setting of chronic viral infections including HIV is largely

unknown. HIV-associated T cell immune activation and inflammation drives cardiovascular risk,

independently of traditional risk factors, duration of antiretroviral treatment and CD4 counts (1-

15, 125-127). In this study, we hypothesized that HIV-driven CD8 T cell immune activation has a

detrimental effect on T_{ang} cell function, compromising their ability to promote endothelial repair,

and chronic interference of this pathway results in vascular inflammation/damage and increased

risk of disease.

PAR1 signaling mediates the cross-talk between the coagulation system, endothelium and

adaptive immune system (CD8 T cells) (38-41). We found that compared with uninfected controls

(PWoH), CD8 T_{ang} cells from PWH had pro-inflammatory phenotype, express PAR1, and the

adhesion and homing receptors CD49f and CX3CR1 respectively (39, 87, 128). In addition, CD8

T_{ang} cells had cytotoxic potential and expressed GZMA and GZMB. The activated phenotype of

CD8 T_{ang} cells but not CD4 T_{ang} was associated with the ASCVD suggesting a relationship between

endothelial inflammation and CD8 T_{ang} cell immune activation.

PAR1 is the main receptor of thrombin, however, other serine proteases including granzymes

activate PAR1 promoting inflammation (44, 47, 50, 129-131). Granzymes (GZMA, GZMB,

GZMK), can activate PAR1 and induce a "biased" or non-canonical signal distinct from thrombin

(93-95). GZMs were first thought to be involved in cytotoxic function and recent evidence

suggests non-cytotoxic and extracellular biological activities including regulation of vascular

integrity, angiogenesis, extracellular matrix remodeling and trans-endothelial migration (32, 46-

48, 50, 93, 94, 96, 97, 132). The present study shows that, *in vitro*, CD8 T_{ang} cells induced

activation of ECs via granzyme induced production of pro-inflammatory cytokines. In addition,

granzyme dependent PAR1 activation provides a potential mechanism by which HIV alters pro-

angiogenic function and the chronic stimulation of this pathway may lead endothelial

inflammation and dysfunction.

Granzymes are emerging as potential contributors of inflammation in cardiovascular diseases

(93, 94, 133-136). Recent study reported a subset of CD8 T cells expression GZMK and GZMB

with proinflammatory (rather than cytotoxic) function in chronically inflamed human tissues

supporting their potential role in inflammation (97). Moreoever, increased levels of GZMs in

biological fluids and tissues has been reported in inflammatory conditions including cardiovascular

disease, autoimmunity, chronic obstructive pulmonary disease, and cancer (20, 93, 97, 134, 135,

137-146). Supporting these observations, we found an association between plasma GZMA leves

and endothelial inflammation markers including sVCAM1 and IL-6 was observed in PWH

suggesting that ongoing endothelial inflammation in the setting of suppressed viremia may be

driven by T_{ang} cell activation.

Alltoghether these data highlight a novel role of CD8 T_{ang} cells in vascular remodeling, and

granzyme-PAR1 dependent pathway promoting endothelial inglammation. CD8 T_{ang} cells and

GZMA dependent PAR1 activation could be a target of intervetion to reduce endothelial

inflammation and injury in PWH potentially mitigating the risk of cardiovascular disease during

chronic HIV infection.

Acknowledgements

Research reported in this publication was supported by the National Institute of Allergy and Infectious Diseases of the National Institutes of Health under award number NIH R01AI145549 and R21AI186960. We thank the District of Columbia Center for AIDS Research, an NIH funded program (P30AI117970) which is supported by the following NIH Co-Funding and Participating Institutes and Centers: NIAID, NCI, NICHD, NHLBI, NIDA, NIMH, NIA, NIDDK, NIMHD, NIDCR, NINR, FIC and OAR. The content of this publication does not necessarily reflect the views or policies of the Department of Health and Human Services, nor does mention of trade names, commercial products, or organizations imply endorsement by the U.S. Government.

Authors' contributions

MC conceptualized the study and supervise data analysis. TL, CM, CJ performed and analyzed mice experiments, JC, ZZ, MFC, MA, SY, performed human experiments, JK, PK, HC recruited patients, KDV, GPA advise and performed endothelial assay, DBG and LP advice mice infection model. MC, TL and LP wrote the manuscript.

750   **References**

751   1.   Baker JV, Sharma S, Grund B, Rupert A, Metcalf JA, Schechter M, et al. Systemic
752   Inflammation, Coagulation, and Clinical Risk in the START Trial. *Open Forum Infect*
753   *Dis.* 2017;4(4):ofx262.
754   2.   Molina JM, Grund B, Gordin F, Williams I, Schechter M, Losso M, et al. Which HIV-
755   infected adults with high CD4 T-cell counts benefit most from immediate initiation of
756   antiretroviral therapy? A post-hoc subgroup analysis of the START trial. *Lancet HIV.*
757   2018.
758   3.   Hadigan C, Paules CI, and Fauci AS. Association Between Human Immunodeficiency
759   Virus Infection and Cardiovascular Diseases: Finding a Solution to Double Jeopardy.
760   *JAMA Cardiol.* 2017;2(2):123-4.
761   4.   Hsue PY, Deeks SG, and Hunt PW. Immunologic basis of cardiovascular disease in HIV-
762   infected adults. *The Journal of infectious diseases.* 2012;205 Suppl 3:S375-82.
763   5.   Sinha A, Ma Y, Scherzer R, Hur S, Li D, Ganz P, et al. Role of T-Cell Dysfunction,
764   Inflammation, and Coagulation in Microvascular Disease in HIV. *Journal of the*
765   *American Heart Association.* 2016;5(12).
766   6.   Lipshultz SE, Miller TL, Wilkinson JD, Scott GB, Somarriba G, Cochran TR, et al.
767   Cardiac effects in perinatally HIV-infected and HIV-exposed but uninfected children and
768   adolescents: a view from the United States of America. *J Int AIDS Soc.* 2013;16:18597.
769   7.   Miller TL, Somarriba G, Orav EJ, Mendez AJ, Neri D, Schaefer N, et al. Biomarkers of
770   vascular dysfunction in children infected with human immunodeficiency virus-1. *J*
771   *Acquir Immune Defic Syndr.* 2010;55(2):182-8.
772   8.   Grund B, Baker JV, Deeks SG, Wolfson J, Wentworth D, Cozzi-Lepri A, et al. Relevance
773   of Interleukin-6 and D-Dimer for Serious Non-AIDS Morbidity and Death among HIV-
774   Positive Adults on Suppressive Antiretroviral Therapy. *PLoS One.* 2016;11(5):e0155100.
775   9.   Freiberg MS, Chang CC, Kuller LH, Skanderson M, Lowy E, Kraemer KL, et al. HIV
776   infection and the risk of acute myocardial infarction. *JAMA internal medicine.*
777   2013;173(8):614-22.
778   10.   Longenecker CT, Funderburg NT, Jiang Y, Debanne S, Storer N, Labbato DE, et al.
779   Markers of inflammation and CD8 T-cell activation, but not monocyte activation, are
780   associated with subclinical carotid artery disease in HIV-infected individuals. *HIV Med.*
781   2013;14(6):385-90.
782   11.   Mattingly AS, Unsal AB, Purdy JB, Gharib AM, Rupert A, Kovacs JA, et al. T-cell
783   Activation and E-selectin Are Associated With Coronary Plaque in HIV-infected Young
784   Adults. *Pediatr Infect Dis J.* 2017;36(1):63-5.
785   12.   Serrano-Villar S, Sainz T, Lee SA, Hunt PW, Sinclair E, Shacklett BL, et al. HIV-
786   infected individuals with low CD4/CD8 ratio despite effective antiretroviral therapy
787   exhibit altered T cell subsets, heightened CD8+ T cell activation, and increased risk of
788   non-AIDS morbidity and mortality. *PLoS Pathog.* 2014;10(5):e1004078.
789   13.   Stein JH, and Hsue PY. Inflammation, immune activation, and CVD risk in individuals
790   with HIV infection. *JAMA.* 2012;308(4):405-6.
791   14.   Longenecker CT, Sullivan C, and Baker JV. Immune activation and cardiovascular
792   disease in chronic HIV infection. *Curr Opin HIV AIDS.* 2016;11(2):216-25.
793   15.   Gharib AM, and Hadigan C. Imaging to End Points: Cardiovascular Disease Risk
794   Assessment in HIV. *Circulation Cardiovascular imaging.* 2017;10(10).

16.   Shah ASV, Stelzle D, Lee KK, Beck EJ, Alam S, Clifford S, et al. Global Burden of
Atherosclerotic Cardiovascular Disease in People Living with the Human
Immunodeficiency Virus: A Systematic Review and Meta-Analysis. *Circulation.* 2018.
17.   Triant VA, Perez J, Regan S, Massaro JM, Meigs JB, Grinspoon SK, et al.
Cardiovascular Risk Prediction Functions Underestimate Risk in HIV Infection.
Circulation. 2018;137(21):2203-14.
18.   Shah ASV, Stelzle D, Lee KK, Beck EJ, Alam S, Clifford S, et al. Global Burden of
Atherosclerotic Cardiovascular Disease in People Living With HIV: Systematic Review
and Meta-Analysis. *Circulation.* 2018;138(11):1100-12.
19.   Lundgren JD, Babiker AG, Sharma S, Grund B, Phillips AN, Matthews G, et al. Long-
Term Benefits from Early Antiretroviral Therapy Initiation in HIV Infection. *NEJM Evid.*
2023;2(3).
20.   Schafer S, and Zernecke A. CD8(+) T Cells in Atherosclerosis. *Cells.* 2020;10(1).
21.   Hsue PY, Deeks SG, and Hunt PW. Immunologic basis of cardiovascular disease in HIV-
infected adults. *The Journal of infectious diseases.* 2012;205 Suppl 3(Suppl 3):S375-82.
22.   Kaplan RC, Sinclair E, Landay AL, Lurain N, Sharrett AR, Gange SJ, et al. T cell
activation and senescence predict subclinical carotid artery disease in HIV-infected
women. *The Journal of infectious diseases.* 2011;203(4):452-63.
23.   Karim R, Mack WJ, Kono N, Tien PC, Anastos K, Lazar J, et al. T-cell activation, both
pre- and post-HAART levels, correlates with carotid artery stiffness over 6.5 years
among HIV-infected women in the WIHS. *J Acquir Immune Defic Syndr.*
2014;67(3):349-56.
24.   Keller TT, Mairuhu AT, de Kruif MD, Klein SK, Gerdes VE, ten Cate H, et al. Infections
and endothelial cells. *Cardiovascular research.* 2003;60(1):40-8.
25.   Wang Y, Brichacek B, Dubrovsky L, Pushkarsky T, Korolowicz K, Rodriguez O, et al.
Increased atherosclerosis in HIV-infected humanized mice is caused by a single viral
protein, Nef. *The Journal of infectious diseases.* 2025.
26.   Hur J, Yang HM, Yoon CH, Lee CS, Park KW, Kim JH, et al. Identification of a novel
role of T cells in postnatal vasculogenesis: characterization of endothelial progenitor cell
colonies. *Circulation.* 2007;116(15):1671-82.
27.   Asahara T, Murohara T, Sullivan A, Silver M, van der Zee R, Li T, et al. Isolation of
putative progenitor endothelial cells for angiogenesis. *Science (New York, NY.*
1997;275(5302):964-7.
28.   Yoder MC. Endothelial stem and progenitor cells (stem cells): (2017 Grover Conference
Series). *Pulm Circ.* 2018;8(1):2045893217743950.
29.   Rouhl RP, Mertens AE, van Oostenbrugge RJ, Damoiseaux JG, Debrus-Palmans LL,
Henskens LH, et al. Angiogenic T-cells and putative endothelial progenitor cells in
hypertension-related cerebral small vessel disease. *Stroke; a journal of cerebral*
circulation. 2012;43(1):256-8.
30.   Rodriguez-Carrio J, Alperi-Lopez M, Lopez P, Alonso-Castro S, Ballina-Garcia FJ, and
Suarez A. Angiogenic T cells are decreased in rheumatoid arthritis patients. *Ann Rheum*
Dis. 2015;74(5):921-7.
31.   Coughlin SR. How the protease thrombin talks to cells. *Proceedings of the National*
Academy of Sciences of the United States of America. 1999;96(20):11023-7.
32.   Coughlin SR. Thrombin signalling and protease-activated receptors. *Nature.*
2000;407(6801):258-64.

33. Soh UJ, Dores MR, Chen B, and Trejo J. Signal transduction by protease-activated
receptors. *British journal of pharmacology.* 2010;160(2):191-203.
34. Sambrano GR, Weiss EJ, Zheng YW, Huang W, and Coughlin SR. Role of thrombin
signalling in platelets in haemostasis and thrombosis. *Nature.* 2001;413(6851):74-8.
35. Kahn ML, Zheng YW, Huang W, Bigornia V, Zeng D, Moff S, et al. A dual thrombin
receptor system for platelet activation. *Nature.* 1998;394(6694):690-4.
36. Ishihara H, Zeng D, Connolly AJ, Tam C, and Coughlin SR. Antibodies to protease-
activated receptor 3 inhibit activation of mouse platelets by thrombin. *Blood.*
1998;91(11):4152-7.
37. Xu WF, Andersen H, Whitmore TE, Presnell SR, Yee DP, Ching A, et al. Cloning and
characterization of human protease-activated receptor 4. *Proceedings of the National*
Academy of Sciences of the United States of America. 1998;95(12):6642-6.
38. Green SA, Smith M, Hasley RB, Stephany D, Harned A, Nagashima K, et al. Activated
platelet-T-cell conjugates in peripheral blood of patients with HIV infection: coupling
coagulation/inflammation and T cells. *AIDS (London, England).* 2015;29(11):1297-308.
39. Mudd JC, Panigrahi S, Kyi B, Moon SH, Manion MM, Younes SA, et al. Inflammatory
Function of CX3CR1+ CD8+ T Cells in Treated HIV Infection Is Modulated by Platelet
Interactions. *The Journal of infectious diseases.* 2016;214(12):1808-16.
40. Hurley A, Smith M, Karpova T, Hasley RB, Belkina N, Shaw S, et al. Enhanced effector
function of CD8(+) T cells from healthy controls and HIV-infected patients occurs
through thrombin activation of protease-activated receptor 1. *The Journal of infectious*
diseases. 2013;207(4):638-50.
41. Chen H, Smith M, Herz J, Li T, Hasley R, Le Saout C, et al. The role of protease-
activated receptor 1 signaling in CD8 T cell effector functions. *iScience.*
2021;24(11):103387.
42. Hollenberg MD. PARs in the stars: proteinase-activated receptors and astrocyte function.
Focus on "Thrombin (PAR-1)-induced proliferation in astrocytes via MAPK involves
multiple signaling pathways". *American journal of physiology Cell physiology.*
2002;283(5):C1347-50.
43. Ramachandran R, Noorbakhsh F, Defea K, and Hollenberg MD. Targeting proteinase-
activated receptors: therapeutic potential and challenges. *Nature reviews Drug discovery.*
2012;11(1):69-86.
44. Wang T, Lee MH, Choi E, Pardo-Villamizar CA, Lee SB, Yang IH, et al. Granzyme B-
induced neurotoxicity is mediated via activation of PAR-1 receptor and Kv1.3 channel.
PLoS One. 2012;7(8):e43950.
45. Zhao P, Metcalf M, and Bunnett NW. Biased signaling of protease-activated receptors.
Frontiers in endocrinology. 2014;5:67.
46. Lee PR, Johnson TP, Gnanapavan S, Giovannoni G, Wang T, Steiner JP, et al. Protease-
activated receptor-1 activation by granzyme B causes neurotoxicity that is augmented by
interleukin-1beta. *J Neuroinflammation.* 2017;14(1):131.
47. Cooper DM, Pechkovsky DV, Hackett TL, Knight DA, and Granville DJ. Granzyme K
activates protease-activated receptor-1. *PLoS One.* 2011;6(6):e21484.
48. Sharma M, Merkulova Y, Raithatha S, Parkinson LG, Shen Y, Cooper D, et al.
Extracellular granzyme K mediates endothelial activation through the cleavage of
protease-activated receptor-1. *The FEBS journal.* 2016;283(9):1734-47.

49.   Bovenschen N, Quadir R, van den Berg AL, Brenkman AB, Vandenberghe I, Devreese
887   B, et al. Granzyme K displays highly restricted substrate specificity that only partially
888   overlaps with granzyme A. *J Biol Chem.* 2009;284(6):3504-12.

889   50.   Suidan HS, Bouvier J, Schaerer E, Stone SR, Monard D, and Tschopp J. Granzyme A
890   released upon stimulation of cytotoxic T lymphocytes activates the thrombin receptor on
891   neuronal cells and astrocytes. *Proc Natl Acad Sci U S A.* 1994;91(17):8112-6.

892   51.   Kaiserman D, Zhao P, Rowe CL, Leong A, Barlow N, Joeckel LT, et al. Granzyme K
893   initiates IL-6 and IL-8 release from epithelial cells by activating protease-activated
894   receptor 2. *PLoS One.* 2022;17(7):e0270584.

895   52.   !!! INVALID CITATION !!! (52).

896   53.   Harker JA, Wong KA, Dallari S, Bao P, Dolgoter A, Jo Y, et al. Interleukin-27R
897   Signaling Mediates Early Viral Containment and Impacts Innate and Adaptive Immunity
898   after Chronic Lymphocytic Choriomeningitis Virus Infection. *Journal of virology.*
899   2018;92(12).

900   54.   Haghverdi L, Lun ATL, Morgan MD, and Marioni JC. Batch effects in single-cell RNA-
901   sequencing data are corrected by matching mutual nearest neighbors. *Nat Biotechnol.*
902   2018;36(5):421-7.

903   55.   Lopez ML, Soriano-Sarabia N, Bruges G, Marquez ME, Preissner KT, Schmitz ML, et al.
904   Expression pattern of protease activated receptors in lymphoid cells. *Cell Immunol.*
905   2014;288(1-2):47-52.

906   56.   Anderson KG, Sung H, Skon CN, Lefrancois L, Deisinger A, Vezys V, et al. Cutting
907   edge: intravascular staining redefines lung CD8 T cell responses. *J Immunol.*
908   2012;189(6):2702-6.

909   57.   Masopust D, Vezys V, Marzo AL, and Lefrancois L. Preferential localization of effector
910   memory cells in nonlymphoid tissue. *Science (New York, NY.* 2001;291(5512):2413-7.

911   58.   Purton JF, Martin CE, and Surh CD. Enhancing T cell memory: IL-7 as an adjuvant to
912   boost memory T-cell generation. *Immunol Cell Biol.* 2008;86(5):385-6.

913   59.   Kang SS, and McGavern DB. Lymphocytic choriomeningitis infection of the central
914   nervous system. *Frontiers in bioscience : a journal and virtual library.* 2008;13:4529-43.

915   60.   Baccala R, Welch MJ, Gonzalez-Quintial R, Walsh KB, Teijaro JR, Nguyen A, et al.
916   Type I interferon is a therapeutic target for virus-induced lethal vascular damage.
917   *Proceedings of the National Academy of Sciences of the United States of America.*
918   2014;111(24):8925-30.

919   61.   Wherry EJ, Teichgraber V, Becker TC, Masopust D, Kaech SM, Antia R, et al. Lineage
920   relationship and protective immunity of memory CD8 T cell subsets. *Nature
921   immunology.* 2003;4(3):225-34.

922   62.   Wherry EJ, Blattman JN, Murali-Krishna K, van der Most R, and Ahmed R. Viral
923   persistence alters CD8 T-cell immunodominance and tissue distribution and results in
924   distinct stages of functional impairment. *Journal of virology.* 2003;77(8):4911-27.

925   63.   Joshi NS, Cui W, Chandele A, Lee HK, Urso DR, Hagman J, et al. Inflammation directs
926   memory precursor and short-lived effector CD8(+) T cell fates via the graded expression
927   of T-bet transcription factor. *Immunity.* 2007;27(2):281-95.

928   64.   Sarkar S, Kalia V, Haining WN, Konieczny BT, Subramaniam S, and Ahmed R.
929   Functional and genomic profiling of effector CD8 T cell subsets with distinct memory
930   fates. *The Journal of experimental medicine.* 2008;205(3):625-40.

65. Chang JT, Wherry EJ, and Goldrath AW. Molecular regulation of effector and memory T
cell differentiation. *Nature immunology.* 2014;15(12):1104-15.

66. Herndler-Brandstetter D, Ishigame H, Shinnakasu R, Plajer V, Stecher C, Zhao J, et al.
KLRG1(+) Effector CD8(+) T Cells Lose KLRG1, Differentiate into All Memory T Cell
Lineages, and Convey Enhanced Protective Immunity. *Immunity.* 2018;48(4):716-29 e8.

67. Perez-Gutierrez L, and Ferrara N. Biology and therapeutic targeting of vascular
endothelial growth factor A. *Nat Rev Mol Cell Biol.* 2023;24(11):816-34.

68. Mor F, Quintana FJ, and Cohen IR. Angiogenesis-inflammation cross-talk: vascular
endothelial growth factor is secreted by activated T cells and induces Th1 polarization. *J*
Immunol. 2004;172(7):4618-23.

69. Wang QR, Wang F, Zhu WB, Lei J, Huang YH, Wang BH, et al. GM-CSF accelerates
proliferation of endothelial progenitor cells from murine bone marrow mononuclear cells
in vitro. *Cytokine.* 2009;45(3):174-8.

70. Bussolino F, Wang JM, Defilippi P, Turrini F, Sanavio F, Edgell CJ, et al. Granulocyte-
and granulocyte-macrophage-colony stimulating factors induce human endothelial cells
to migrate and proliferate. *Nature.* 1989;337(6206):471-3.

71. Biedermann BC. Vascular endothelium: checkpoint for inflammation and immunity.
News in physiological sciences : an international journal of physiology produced jointly
by the International Union of Physiological Sciences and the American Physiological
Society. 2001;16:84-8.

72. Danese S, Dejana E, and Fiocchi C. Immune regulation by microvascular endothelial
cells: directing innate and adaptive immunity, coagulation, and inflammation. *J Immunol.*
2007;178(10):6017-22.

73. Amersfoort J, Eelen G, and Carmeliet P. Immunomodulation by endothelial cells -
partnering up with the immune system? *Nature reviews.* 2022;22(9):576-88.

74. Zhang L, Gao S, White Z, Dai Y, Malik AB, and Rehman J. Single-cell transcriptomic
profiling of lung endothelial cells identifies dynamic inflammatory and regenerative
subpopulations. *JCI Insight.* 2022;7(11).

75. Godoy RS, Cober ND, Cook DP, McCourt E, Deng Y, Wang L, et al. Single-cell
transcriptomic atlas of lung microvascular regeneration after targeted endothelial cell
ablation. *Elife.* 2023;12.

76. Niethamer TK, Stabler CT, Leach JP, Zepp JA, Morley MP, Babu A, et al. Defining the
role of pulmonary endothelial cell heterogeneity in the response to acute lung injury.
Elife. 2020;9.

77. Rodor J, Chen SH, Scanlon JP, Monteiro JP, Caudrillier A, Sweta S, et al. Single-cell
RNA sequencing profiling of mouse endothelial cells in response to pulmonary arterial
hypertension. *Cardiovasc Res.* 2022;118(11):2519-34.

78. Jerkic M, Rivas-Elena JV, Prieto M, Carron R, Sanz-Rodriguez F, Perez-Barriocanal F, et
al. Endoglin regulates nitric oxide-dependent vasodilatation. *FASEB journal : official*
publication of the Federation of American Societies for Experimental Biology.
2004;18(3):609-11.

79. Toporsian M, Gros R, Kabir MG, Vera S, Govindaraju K, Eidelman DH, et al. A role for
endoglin in coupling eNOS activity and regulating vascular tone revealed in hereditary
hemorrhagic telangiectasia. *Circulation research.* 2005;96(6):684-92.

80. Kretschmer S, Dethlefsen I, Hagner-Benes S, Marsh LM, Garn H, and Konig P.
Visualization of intrapulmonary lymph vessels in healthy and inflamed murine lung using
CD90/Thy-1 as a marker. *PLoS One.* 2013;8(2):e55201.
81. Levine JH, Simonds EF, Bendall SC, Davis KL, Amir el AD, Tadmor MD, et al. Data-
Driven Phenotypic Dissection of AML Reveals Progenitor-like Cells that Correlate with
Prognosis. *Cell.* 2015;162(1):184-97.
82. Li Q, Wei S, Li Y, Wu F, Qin X, Li Z, et al. Blocking of programmed cell death-ligand 1
(PD-L1) expressed on endothelial cells promoted the recruitment of CD8(+)IFN-
gamma(+) T cells in atherosclerosis. *Inflammation research : official journal of the*
European Histamine Research Society [et al]. 2023;72(4):783-96.
83. Yu QC, Song W, Wang D, and Zeng YA. Identification of blood vascular endothelial
stem cells by the expression of protein C receptor. *Cell research.* 2016;26(10):1079-98.
84. Thiyagarajan M, Cheng T, and Zlokovic BV. Endothelial cell protein C receptor: role
beyond endothelium? *Circulation research.* 2007;100(2):155-7.
85. Zhu Z, Li T, Chen J, Kumar J, Kumar P, Qin J, et al. The Role of Inflammation and
Immune Activation on Circulating Endothelial Progenitor Cells in Chronic HIV
Infection. *Frontiers in immunology.* 2021;12:663412.
86. Kushner EJ, MacEneaney OJ, Morgan RG, Van Engelenburg AM, Van Guilder GP, and
DeSouza CA. CD31+ T cells represent a functionally distinct vascular T cell phenotype.
Blood Cells Mol Dis. 2010;44(2):74-8.
87. Pilling JE, Galvin A, Robins AM, Sewell HF, and Mahida YR. Expression of alpha5
(CD49e) and alpha6 (CD49f) integrin subunits on T cells in the circulation and the
lamina propria of normal and inflammatory bowel disease colonic mucosa. *Scandinavian*
journal of immunology. 1998;48(4):425-8.
88. Gimbrone MA, Jr., and Garcia-Cardena G. Endothelial Cell Dysfunction and the
Pathobiology of Atherosclerosis. *Circulation research.* 2016;118(4):620-36.
89. Singh V, Kaur R, Kumari P, Pasricha C, and Singh R. ICAM-1 and VCAM-1:
Gatekeepers in various inflammatory and cardiovascular disorders. *Clin Chim Acta.*
2023;548:117487.
90. Groten SA, Smit ER, Janssen EFJ, van den Eshof BL, van Alphen FPJ, van der Zwaan C,
et al. Multi-omics delineation of cytokine-induced endothelial inflammatory states.
Commun Biol. 2023;6(1):525.
91. Nakamichi M, Akishima-Fukasawa Y, Fujisawa C, Mikami T, Onishi K, and Akasaka Y.
Basic Fibroblast Growth Factor Induces Angiogenic Properties of Fibrocytes to Stimulate
Vascular Formation during Wound Healing. *The American journal of pathology.*
2016;186(12):3203-16.
92. Arnaoutova I, and Kleinman HK. In vitro angiogenesis: endothelial cell tube formation
on gelled basement membrane extract. *Nature protocols.* 2010;5(4):628-35.
93. Zeglinski MR, and Granville DJ. Granzymes in cardiovascular injury and disease.
Cellular signalling. 2020;76:109804.
94. Chamberlain CM, and Granville DJ. The role of Granzyme B in atheromatous diseases.
Canadian journal of physiology and pharmacology. 2007;85(1):89-95.
95. Choy JC, McDonald PC, Suarez AC, Hung VH, Wilson JE, McManus BM, et al.
Granzyme B in atherosclerosis and transplant vascular disease: association with cell
death and atherosclerotic disease severity. *Modern pathology : an official journal of the*
United States and Canadian Academy of Pathology, Inc. 2003;16(5):460-70.

96.   Arias M, Martinez-Lostao L, Santiago L, Ferrandez A, Granville DJ, and Pardo J. The
Untold Story of Granzymes in Oncoimmunology: Novel Opportunities with Old
Acquaintances. *Trends Cancer.* 2017;3(6):407-22.

97.   Jonsson AH, Zhang F, Dunlap G, Gomez-Rivas E, Watts GFM, Faust HJ, et al.
Granzyme K(+) CD8 T cells form a core population in inflamed human tissue. *Sci Transl*
Med. 2022;14(649):eabo0686.

98.   Mahrus S, and Craik CS. Selective chemical functional probes of granzymes A and B
reveal granzyme B is a major effector of natural killer cell-mediated lysis of target cells.
Chem Biol. 2005;12(5):567-77.

99.   Plasman K, Demol H, Bird PI, Gevaert K, and Van Damme P. Substrate specificities of
the granzyme tryptases A and K. *J Proteome Res.* 2014;13(12):6067-77.

100.   Zhang C, Srinivasan Y, Arlow DH, Fung JJ, Palmer D, Zheng Y, et al. High-resolution
crystal structure of human protease-activated receptor 1. *Nature.* 2012;492(7429):387-92.

101.   Brass LF, Pizarro S, Ahuja M, Belmonte E, Blanchard N, Stadel JM, et al. Changes in the
structure and function of the human thrombin receptor during receptor activation,
internalization, and recycling. *J Biol Chem.* 1994;269(4):2943-52.

102.   O'Brien PJ, Prevost N, Molino M, Hollinger MK, Woolkalis MJ, Woulfe DS, et al.
Thrombin responses in human endothelial cells. Contributions from receptors other than
PAR1 include the transactivation of PAR2 by thrombin-cleaved PAR1. *J Biol Chem.*
2000;275(18):13502-9.

103.   Qiu C, Xie Q, Zhang D, Chen Q, Hu J, and Xu L. GM-CSF induces cyclin D1 expression
and proliferation of endothelial progenitor cells via PI3K and MAPK signaling. *Cell*
Physiol Biochem. 2014;33(3):784-95.

104.   Bruno S, Bussolati B, Scacciatella P, Marra S, Sanavio F, Tarella C, et al. Combined
administration of G-CSF and GM-CSF stimulates monocyte-derived pro-angiogenic cells
in patients with acute myocardial infarction. *Cytokine.* 2006;34(1-2):56-65.

105.   Cho HJ, Kim HS, Lee MM, Kim DH, Yang HJ, Hur J, et al. Mobilized endothelial
progenitor cells by granulocyte-macrophage colony-stimulating factor accelerate
reendothelialization and reduce vascular inflammation after intravascular radiation.
Circulation. 2003;108(23):2918-25.

106.   Stabile E, Kinnaird T, la Sala A, Hanson SK, Watkins C, Campia U, et al. CD8+ T
lymphocytes regulate the arteriogenic response to ischemia by infiltrating the site of
collateral vessel development and recruiting CD4+ mononuclear cells through the
expression of interleukin-16. *Circulation.* 2006;113(1):118-24.

107.   Newman PJ. Switched at birth: a new family for PECAM-1. *The Journal of clinical*
investigation. 1999;103(1):5-9.

108.   Fornasa G, Groyer E, Clement M, Dimitrov J, Compain C, Gaston AT, et al. TCR
stimulation drives cleavage and shedding of the ITIM receptor CD31. *J Immunol.*
2010;184(10):5485-92.

109.   Ma L, Mauro C, Cornish GH, Chai JG, Coe D, Fu H, et al. Ig gene-like molecule CD31
plays a nonredundant role in the regulation of T-cell immunity and tolerance.
Proceedings of the National Academy of Sciences of the United States of America.
2010;107(45):19461-6.

110.   Delacher M, Schmidleithner L, Simon M, Stuve P, Sanderink L, Hotz-Wagenblatt A, et
al. The effector program of human CD8 T cells supports tissue remodeling. *The Journal*
of experimental medicine. 2024;221(2).

111. Deliyanti D, Figgett WA, Gebhardt T, Trapani JA, Mackay F, and Wilkinson-Berka JL.
CD8(+) T Cells Promote Pathological Angiogenesis in Ocular Neovascular Disease.
Arterioscler Thromb Vasc Biol. 2023;43(4):522-36.

112. Pober JS, and Sessa WC. Evolving functions of endothelial cells in inflammation. *Nature*
reviews. 2007;7(10):803-15.

113. Frebel H, Nindl V, Schuepbach RA, Braunschweiler T, Richter K, Vogel J, et al.
Programmed death 1 protects from fatal circulatory failure during systemic virus
infection of mice. *The Journal of experimental medicine.* 2012;209(13):2485-99.

114. Lee C, Kim MJ, Kumar A, Lee HW, Yang Y, and Kim Y. Vascular endothelial growth
factor signaling in health and disease: from molecular mechanisms to therapeutic
perspectives. *Signal Transduct Target Ther.* 2025;10(1):170.

115. Li C, Issa R, Kumar P, Hampson IN, Lopez-Novoa JM, Bernabeu C, et al. CD105
prevents apoptosis in hypoxic endothelial cells. *Journal of cell science.* 2003;116(Pt
13):2677-85.

116. Sharma S, Ehrlich M, Zhang M, Blobe GC, and Henis YI. NRP1 interacts with endoglin
and VEGFR2 to modulate VEGF signaling and endothelial cell sprouting. *Commun Biol.*
2024;7(1):112.

117. Cissy Yu Q, Bai L, Chen Y, Chen Y, Peng G, Wang D, et al. Embryonic vascular
establishment requires protein C receptor-expressing endothelial progenitors.
Development. 2022;149(12).

118. Lin Y, Banno K, Gil CH, Myslinski J, Hato T, Shelley WC, et al. Origin, prospective
identification, and function of circulating endothelial colony-forming cells in mice and
humans. *JCI Insight.* 2023;8(5).

119. Weil BR, Kushner EJ, Diehl KJ, Greiner JJ, Stauffer BL, and Desouza CA. CD31+ T
cells, endothelial function and cardiovascular risk. *Heart Lung Circ.* 2011;20(10):659-62.

120. Lopez P, Rodriguez-Carrio J, Martinez-Zapico A, Caminal-Montero L, and Suarez A.
Senescent profile of angiogenic T cells from systemic lupus erythematosus patients. *J*
Leukoc Biol. 2016;99(3):405-12.

121. de Boer SA, Reijrink M, Abdulahad WH, Hoekstra ES, Slart R, Heerspink HJL, et al.
Angiogenic T cells are decreased in people with type 2 diabetes mellitus and recruited by
the dipeptidyl peptidase-4 inhibitor Linagliptin: A subanalysis from a randomized,
placebo-controlled trial (RELEASE study). *Diabetes Obes Metab.* 2020;22(7):1220-5.

122. Kul A, Ozturk N, Kurt AK, and Arslan Y. Detection of Angiogenic T Cells and
Endothelial Progenitor Cells in Behcet Disease and Determination of Their Relationship
with Disease Activity. *Life (Basel).* 2023;13(6).

123. Manetti M, Pratesi S, Romano E, Bellando-Randone S, Rosa I, Guiducci S, et al.
Angiogenic T cell expansion correlates with severity of peripheral vascular damage in
systemic sclerosis. *PLoS One.* 2017;12(8):e0183102.

124. Zhang G, Liu Y, Qiu Y, Zhang J, Sun J, Zhou Z, et al. Circulating senescent angiogenic T
cells are linked with endothelial dysfunction and systemic inflammation in hypertension.
J Hypertens. 2021;39(5):970-8.

125. Deeks SG, Lewin SR, and Havlir DV. The end of AIDS: HIV infection as a chronic
disease. *Lancet.* 2013;382(9903):1525-33.

126. Triant VA. Cardiovascular disease and HIV infection. *Curr HIV/AIDS Rep.*
2013;10(3):199-206.

127. Nou E, Lo J, and Grinspoon SK. Inflammation, immune activation, and cardiovascular disease in HIV. *AIDS (London, England)*. 2016;30(10):1495-509.

128. Panigrahi S, Chen B, Fang M, Potashnikova D, Komissarov AA, Lebedeva A, et al. CX3CL1 and IL-15 Promote CD8 T cell chemoattraction in HIV and in atherosclerosis. *PLoS Pathog*. 2020;16(9):e1008885.

129. Ossovskaya VS, and Bunnett NW. Protease-activated receptors: contribution to physiology and disease. *Physiological reviews*. 2004;84(2):579-621.

130. van Eck JA, Shan L, Meeldijk J, Hack CE, and Bovenschen N. A novel proinflammatory role for granzyme A. *Cell Death Dis*. 2017;8(2):e2630.

131. Wensink AC, Hack CE, and Bovenschen N. Granzymes regulate proinflammatory cytokine responses. *J Immunol*. 2015;194(2):491-7.

132. Richardson KC, Jung K, Pardo J, Turner CT, and Granville DJ. Noncytotoxic Roles of Granzymes in Health and Disease. *Physiology (Bethesda)*. 2022;37(6):323-48.

133. Ikemoto T, Hojo Y, Kondo H, Takahashi N, Hirose M, Nishimura Y, et al. Plasma granzyme B as a predicting factor of coronary artery disease--clinical significance in patients with chronic renal failure. *Journal of cardiology*. 2009;54(3):409-15.

134. Kondo H, Hojo Y, Tsuru R, Nishimura Y, Shimizu H, Takahashi N, et al. Elevation of plasma granzyme B levels after acute myocardial infarction. *Circulation journal : official journal of the Japanese Circulation Society*. 2009;73(3):503-7.

135. Skjelland M, Michelsen AE, Krohg-Sorensen K, Tennoe B, Dahl A, Bakke S, et al. Plasma levels of granzyme B are increased in patients with lipid-rich carotid plaques as determined by echogenicity. *Atherosclerosis*. 2007;195(2):e142-6.

136. Hendel A, Hiebert PR, Boivin WA, Williams SJ, and Granville DJ. Granzymes in age-related cardiovascular and pulmonary diseases. *Cell death and differentiation*. 2010;17(4):596-606.

137. Xu T, Zhu HX, You X, Ma JF, Li X, Luo PY, et al. Single-cell profiling reveals pathogenic role and differentiation trajectory of granzyme K+CD8+ T cells in primary Sjogren's syndrome. *JCI Insight*. 2023;8(8).

138. Tsuru R, Kondo H, Hojo Y, Gama M, Mizuno O, Katsuki T, et al. Increased granzyme B production from peripheral blood mononuclear cells in patients with acute coronary syndrome. *Heart*. 2008;94(3):305-10.

139. Hiebert PR, Boivin WA, Abraham T, Pazooki S, Zhao H, and Granville DJ. Granzyme B contributes to extracellular matrix remodeling and skin aging in apolipoprotein E knockout mice. *Experimental gerontology*. 2011;46(6):489-99.

140. Hansson GK, Holm J, and Jonasson L. Detection of activated T lymphocytes in the human atherosclerotic plaque. *The American journal of pathology*. 1989;135(1):169-75.

141. Kyaw T, Winship A, Tay C, Kanellakis P, Hosseini H, Cao A, et al. Cytotoxic and proinflammatory CD8+ T lymphocytes promote development of vulnerable atherosclerotic plaques in apoE-deficient mice. *Circulation*. 2013;127(9):1028-39.

142. Cochain C, Koch M, Chaudhari SM, Busch M, Pelisek J, Boon L, et al. CD8+ T Cells Regulate Monopoiesis and Circulating Ly6C-high Monocyte Levels in Atherosclerosis in Mice. *Circulation research*. 2015;117(3):244-53.

143. Olofsson PS, Soderstrom LA, Wagsater D, Sheikine Y, Ocaya P, Lang F, et al. CD137 is expressed in human atherosclerosis and promotes development of plaque inflammation in hypercholesterolemic mice. *Circulation*. 2008;117(10):1292-301.

1157   144.   Kolbus D, Ramos OH, Berg KE, Persson J, Wigren M, Bjorkbacka H, et al. CD8+ T cell
1158   activation predominate early immune responses to hypercholesterolemia in Apoe(-)(/)(-)
1159   mice. *BMC Immunol.* 2010;11:58.
1160   145.   Kolbus D, Ljungcrantz I, Andersson L, Hedblad B, Fredrikson GN, Bjorkbacka H, et al.
1161   Association between CD8+ T-cell subsets and cardiovascular disease. *J Intern Med.*
1162   2013;274(1):41-51.
1163   146.   Tyrrell DJ, Wragg KM, Chen J, Wang H, Song J, Blin MG, et al. Clonally expanded
1164   memory CD8(+) T cells accumulate in atherosclerotic plaques and are pro-atherogenic in
1165   aged mice. *Nat Aging.* 2023;3(12):1576-90.
1166
1167

Figure Legends

Figure 1. PAR1 expression is modulated in LCMVGp33-specific CD8 T lymphocytes during acute LCMV infection.

In vivo expression of PAR1 in splenic cells from naïve C57BL/6-F2r$^{fl/fl}$ (n= 5) and C57BL/6 (n= 5) mice analyzed by flow cytometry. **(A)** Cells were analyzed after gating on live cells, and singlets and CD45$^+$ lymphocytes. Representative contour plots gating cell types: NK cells (CD3$^-$NK1.1$^+$), B cells (MHCII$^+$B220$^+$), CD4 (CD3$^+$CD4$^+$) and CD8 T cells (CD3$^+$CD8$^+$). **(B)** Representative histogram of tdTomato (PAR1) expression in CD4 and CD8 T cells, B cells and NK cells from C57BL/6 mice (solid grey) and C57BL/6-F2r$^{fl/fl}$ mice (black line). **(C)** Median Fluorescence Intensity (MFI) of tdTomato expression of total splenic cells from C57BL/6 mice (SP) and B cells, NK and CD4 and CD8 T cells and NK cells from C57BL/6-F2r$^{fl/fl}$ mice. **(D)** Relative *F2r* mRNA expression in sorted B cells and purified by magnetic isolation of CD4 and CD8 T cells from C57BL/6-F2r$^{fl/fl}$ mice (n= 6). **(E)** C57BL/6-F2r$^{fl/fl}$ mice were infected with LCMVArm strain (2 x 10^5 pfu, i.v). Splenic and lung cells were assessed at day 6 and 21 p.i. by flow cytometry. **(F)** Representative histogram of tdTomato (tdT) expression in total CD8 T cells from naïve (N) and infected mice at day 6 and 21 p.i. (solid gray). Overlayed, tdTomato expression in LCMVGp33-specific CD8 T cells detected by H2-D^{b}LCMVGp33 pentamer staining (red line). **(F)** Median Fluorescence Intensity (MFI) of tdTomato expression in LCMVGp33-specific CD8 T cells detected by H2-D^{b}LCMVGp33 pentamer staining. Dashed line represents level expression of total naïve CD8 T cells. **(G)** Representative plot of lung endothelial cells (EC). After excluding dead cells and doublets, ECs were gated as CD326$^-$CD45$^-$CD31$^+$. Frequency and cell count of lung endothelial cells from naïve (N) and infected mice at days 6 and 21 p.i. Frequencies of ECs are

expressed as frequency of total live cells. **(H)** Median Fluorescence Intensity (MFI) of tdTomato

expression in lung ECs from naïve (N) and infected mice (day 6 and day 21p.i.).

The bar graph is represented by box and whisker showing the median value with first and third

quartiles in the box, with whiskers extending to the minimum and maximum values. The data in

line graphs represent mean±SEM in each time point. Data are pooled from three independent

experiments with 5-9 naïve animals (C, E and F) and 8-17 per time point (F and H). Statistical

analysis was performed using non-parametric Mann-Whitney test. *P* value < 0.05 was considered

significant.

1198

Figure 2. Angiogenic CD8 T cells promote endothelial recovery during acute viral infection.

(A) Schematic presentation of CD8 T cell depletion. C57BL/6-F2r$^{fl/fl}$ mice were infected with

LCMVArm strain (2 x 10^5 pfu, i.v). At day 15 p.i mice were administered anti-CD8α mAb and

IgG2b isotype for control mice. Frequency and cell count of CD8 T cells at day 21p.i in CD8 T

cell depleted mice and IgG treated control mice assessed by flow cytometry. **(B)** Lung endothelial

cell frequency and count from CD8 T cell depleted and control mice at day 21 p.i. Frequencies of

ECs are expressed as frequency of total live cells. **(C)** Representative contour plot of CD31^{high} and

CD309 expression by lung CD8 T cells from naïve mice and infected mice. Frequency and cell

count of CD31^{high}CD8 T cells from naïve (N) mice and infected mice at day 6 and 21 p.i. **(D)** Lung

CD31^{high} and CD31^{neg} CD8 T cells, representative contour plot of secretion of IFNγ, TNFα

secretion and CD107a upon *in vitro* stimulation with LCMVGp33-44 peptide. DMSO was used as

control. **(E)** Representative contour plot of KLRG1 and CD127 expression by LCMVGp33-

specific CD8 T cells. Frequency of KLRG1$^-$CD127$^+$ (R1), KLRG1$^+$CD127$^+$ (R2) and

KLRG1$^+$CD127$^-$ and KLRG1$^-$CD127$^-$ (R3+R4) LCMVGp33 specific CD8 T cells from lung at

days 6 and 21 p.i.. **(F)** Lung CD31^{high} and CD31^{neg} bulk CD8 T cells were sorted from naïve mice

and infected mice at 21 p.i. and stimulated *in vitro* with anti-CD3 and anti-CD28 mAbs coated

plate for 3 days. Detection of VEGF in the supernatants of the culture. The bar graphs are

represented by box and whisker showing the median value with first and third quartiles in the box,

with whiskers extending to the minimum and maximum values. The data in line graphs represent

mean±SEM in each time point. Data are pooled from two independent experiments with 8 naïve

animals (A and B) and 7-16 per time point (C, D and E). Sorted cells from 3 naïve and 6 infected

mice per time point (G). Statistical analysis was performed using non-parametric Mann-Whitney

test. *P* value < 0.05 was considered significant.

1222

Figure 3. CD31^{high}CD8 T cells are associated with endothelial that express a regenerative

phenotype.

(A) Unsupervised clustering analysis of lung endothelial cells using phonograph (k= 100). The

heatmap represent mean fluorescence intensity of CD31, PDL1, tdTomato, CD34, CD309,

VCAM-1, CD105, CD90.2 and CD201 expression in ECs. Bubble plots represent the mean

frequency of each cluster from naïve and infected mice at day 6 and 21 p.i. **(B)** Relationship

between frequency of Cluster 6 and lung CD31^{high}CD8 T cells. Data are pooled from two

independent experiments with 7 naïve (A and B) and 8-10 per time point animals (A and B).

Correlation were performed using non-parametric Spearman correlation and *p* value ⩽ 0.01 was

considered significant.

1233
Figure 4. Pro-inflammatory phenotype of CD8 T$_{ang}$ correlates to ASCVD risk in PWH.

Circulating T$_{ang}$ were analyzed in PBMCs from PWH (n= 25, red symbols) and PWoH (n= 17,

blue symbols) by flowcytometry. After gating on live cells and singlets, CD4 and CD8 T cells

were analyzed based on expression of CD31 and CXCR4: **(A)** Representative contour plots for

gating angiogenic T cells, CD4 T_{ang} (CD31$^+$CXCR4$^+$) and CD8 T_{ang} (CD31$^+$CXCR4$^+$). **(B)**

Frequency of CD4 T_{ang} and CD8 T_{ang} from PWH and PWoH. CD4 T_{ang} and CD8 T_{ang} expression

of: **(C)** CX3CR1 and PAR1 **(D)** CD49f and PAR1 **(E)** granzyme A (GZMA) and B (GZMB). **F,**

G and H. Relationship between expression of CD4 and CD8 T_{ang} cells and atherosclerotic

cardiovascular disease (ASCVD) risk score in PWH. **(I)** Relationship between expression of

plasma levels of sVCAM-1, sICAM-1, IL-6 and granzyme A in PWH (n= 25). The bar graph is

represented by box and whisker showing the median value with first and third quartiles in the box,

with whiskers extending to the minimum and maximum values. Statistical analysis was performed

using non-parametric Mann-Whitney test. *P* value < 0.05 was considered significant. Correlations

were performed using non-parametric Spearman correlation and *p* value $\leqslant$ 0.01 was considered

significant.

1249

Figure 5. *In vitro*, CD8 T_{ang} cells from PWH induced shedding of sICAM-1 and sVCAM-1 by

endothelial cells.

CD8 T_{ang} from PWH (n= 10, red symbols) and PWoH (n= 10, blue symbols) were sorted and

stimulated with CD3 and CD28 mAbs coated beads for 3 days and expanded in the presence of

IL-2 for an additional 14 days. Activated CD8 T_{ang} cells were co-cultured with primary human

umbilical vein endothelial cells HUVEC at 2 to 1 CD8 T_{ang} to HUVEC ratio for 24 hours.

Supernatants were collected and measured: **(A)** Soluble vascular cell adhesion molecule 1

(sVCAM1) and intracellular adhesion molecule 1 (sICAM1). **(B)** Cytokines IL-6, Angiopoietin-2

(Ang-2), fibroblast growth factor (bFGF), vascular endothelial growth factor (VEGF) and IL-

8/CXCL8. **(C)** Sorted CD8 T_{ang} memory cells (CD45RA$^-$CD27$^+$) from PWH (n= 3, opened red

symbols), PWoH (n= 2, blue symbols), and HIVGag-specific CD8 T cells from PWH (n= 2, solid

red symbols) were labeled with CFSE and cultured with HUVEC at a 2 to 1 ratio in Ibidi chambers

for up to 12 hours. As control, HUVEC were cultured alone (black symbol). Representative phase

contrast images at 2, 4, 6, 8, 10 and 12 hours with masks of tube (black line), branching point

(white point), loop (labeled with number) and CD8 T_{ang} cells (red dots). **(D)** Representative

fluorescent images of T_{ang} cells and HUVEC. **(E)** Fold change of median loop area of individual

donors from PWH and PWoH and HIVGag-specific CD8 T cells. **(F)** Fold change of median loop

area of pooled all donors. HUVEC shows mean±SEM of pooled experiments with all donors. The

bar graph is represented by box and whisker showing the median value with first and third quartiles

in the box, with whiskers extending to the minimum and maximum values. Line graphs represent

the average of all donors. Statistical analysis was performed using non-parametric Mann-Whitney

test. *P* value < 0.05 was considered significant.

1272

Figure 6. Activated CD8 T_{ang} cells mediate endothelial cell activation through a granzyme-

dependent PAR1 activation

(A) Diagram illustrates the molecular mechanism of cleavage dependent PAR1 activation via

serine proteases. Inhibition of PAR1 activation by ATAP-2 mAb and antagonist SCH530348

(vorapaxar). Peptide agonist (mimics the tethered ligand) and cleavage independent PAR1

activation. The figure was generated with Biorender. **(B)** Image of Granzyme A induced

intracellular calcium mobilization in HUVEC. Intracellular calcium mobilization kinetics in

HUVEC stimulated with granzyme A and thrombin as control. Each line represents the kinetics

overtime of individual cells (lower panel). **(C, D and E)** HUVEC were stimulated with 100 nM of

rh GZMA and GZMK overnight in the presence or absence of 10 µg/mL of ATAP-2 mAb or 300

nM of PAR1 antagonist SCH530348. (**F, G** and **H**) HUVEC were stimulated with 100 μM of

PAR1 agonist peptide (TFLLR-NH2) and peptide control (RLLFT-NH2) as positive and negative

controls respectively. IL-6, IL-8/CXCL8 and Angiopoietin-2 were measured in the supernatants

of the cultures. The bar graph is represented by box and whisker showing the median value with

first and third quartiles in the box, with whiskers extending to the minimum and maximum values.

Statistical analysis was performed using non-parametric Mann-Whitney test. *P* value <0.05 was

considered significant.

1290

Table 1. Study groups characteristics

Demographics	PWH (n= 25)	PWOH (n= 17)
Age, yr, median (IQR)	52 (42-57.5)	48 (34-60)
Gender n (%)		
Male	22 (88)	8 (47.1)
Female	3 (12)	9 (52.9)
Ethnicity n (%)		
White	8 (32)	11 (64.8)
African American	12 (48)	3 (17.6)
Other	5 (20)	3 (17.6)
Clinical Characteristics		
Smoker ever, n (%)	8 (32)	N/A
Diabetes, n (%)	9 (36)	N/A
ASCVD risk estimator, median (IQR)[1]	7.9 (5.0-18.6)	N/A
Clinical Laboratory, median (IQR)		
Cholesterol, mg/dL[2]	168 (147-194)	N/A
HDL, mg/dL[3]	47 (41-56.3)	N/A
LDL, mg/dL[3]	99.5 (78.5-117.5)	N/A
HIV Characteristics, median (IQR)		
Viral Load copies/mL[4]	<20 (20-20)	N/A
CD4 count, cells/μL	663 (532-861.5)	N/A
ART treatment		
NRTIs, n (%)	19 (76)	N/A
NNRTIs, n (%)	4 (16)	N/A
PIs, n (%)	4 (16)	N/A
INSTIs, n (%)	14 (56)	N/A
Serum Biomarkers, median (IQR)		
sVCAM-1, pg/mL	288.5 (188.6-366.6)	
IL-6	3.7 (1.2-4.6)	
Granzyme A	256.1 (111.1-514)	
Granzyme B	920.8 (523.4-1566)	
IFNγ	289.8 (87.0-1012)	
TNFα	2.5 (1.1-4.6)	
sFASL	9.8 (3.6-24.1)	

[1] 5 out of 25 participants didn't have ASCVD risk estimator. [2] 10 out of 25 participants didn't have total cholesterol. [3] 11 out of 25 participants didn't have HDL and LDL. [4] 4 out of 25 participants have viral load 110, 70, 60 and 50 HIV RNA copies/mL respectively.

Figure 1

Figure 2

Figure 3

(A)

(B)

Figure 4

Figure 5

Figure 6