High Cross Pollination Frequency in Rice Landraces in Field Condition

Debal Deb*, Debdulal Bhattacharya, Mahendra Nauri

Centre for Interdisciplinary Studies
Kolkata

* Corresponding author : debdebal@gmail.com

Key words : Landraces, Rice, Cross Pollination, Flower opening time, Flower exposure duration

Introduction

The flower opening time (FOT) and the flower exposure duration (FED, measured as the period between flower opening time and closing time) are crucially important for the success of pollination and fertilization in rice plants. Although several studies have reported the range of FOT and flower closing time (FCT) of different landraces[1-9] (**Table 1**), none of them documented the FED of the pollen parent (PP) and ovary parent (OP) landraces in the crossing experiments.

Table 1. Published Records of Earliest and Latest FOT for Rice Cultivars from Different Countries.

Country of Origin	*Indica*		*Japonica*		Author Reference
	Earliest FOT	Last FOT	Earliest FOT	Last FOT	
India	07:00	12:00			1
	09:53	11:16			2
Long day	8:55	11:36			3; 4
Short day	9:18	12:19	9:37	12:10	3; 4
Indonesia	09:00	13:00	11:01	-	5
	10:44	12:00			2
Nepal			10:51	11:30	2
Myanmar	07:00	09:00			6
	10:38	11:31			2
China	09:01	10:47	11:44	12:02	2
Japan	10:08	11:42	10:19	12:35	2
The Philippines	09:00	11:30			7
	09:55	11:15			2
USA	08:00	17:00	10:10	11:42	8; 2
			10:14	13:00	9

The most crucial period for successful cross pollination is the pollen recipient flower's exposure duration, calculated as the difference between the PP's FCT and OP's FOT, or alternatively, the OP's FCT and PP's FOT, which is denoted here as the 'effective exposure duration' (EED). The CPF is likely to be proportionately high only when EED > 0. In other words, if the flowers of the ovary parent (OP) close even a second before the FOT of the pollen parent (PP), the FED overlap between the two parents is zero, and consequently, there would be no cross pollination. Oblivious to this overlap of the crossing pair's exposure duration (or EED as defined here), all previous experimental studies [1-9] as well as authentic reference books

[10,11] inform that cross pollination frequency (CPF) in cultivated rice (*Oryza sativa*) rarely exceeds 2%. This inference of low CPF is also reported from experiments involving crosses between cultivated and transgenic rice cultivars [12-15]. The only report of over 9% CPF, in one instance, came from an experiment by Bae et al. [16], yet the overall CPF did not exceed 1% "in the field condition" (ref. 16, p. 168). We surmise that in all these experiments [1-9, 12-16], near-zero overlap of the FED between the PP and OP might be the only factor resulting in extremely low CPF reported in those experiments, and therefore the received wisdom of extremely low CPF in cultivated rice is a severe underestimation of the actual value in real life conditions [17]. Our 2019 study (published in 2021, ref. 17) reported for the first time that CPF exceeded 80% in a pair of rice landraces, whose respective FEDs had significantly overlapped.

In continuation of our previous experiment in cross pollination conducted in 2019, we conducted similar experiments in subsequent years using 19 pairs of landraces during winter of 2020 and 8 pairs during summer of 2021. The range of flowering periods of the landraces in each pair were overlapping. We report here the results of the new sets of experiment to test our conjecture that the CPF in cultivated rice is usually very high unless the overlap between the FEDs of the crossing pairs is zero or near-zero.

Materials and Methods

Study site

The experiment was conducted at Basudha conservation farm (19°42′32.0″N, 83°28′8.4″E), located in Rayagada district, Odisha, India, during *aman* season (June–December) of the year 2020 and *boro* (Januiary – March) of 2021. The uniformly sandy loam soil of the farm was enriched with decomposed farm yard manure, farm-grown phosphate solubilizing microbial (PSM) inocula, and leaf mulching, and received no synthetic agrochemicals for the past 25 years.

Selection of landraces

The experiment was conducted with rice landraces in two seasons, first with the short-day (*aman*) season landraces, flowering in winter (October-November 2020); and subsequently, with the photoperiod-insensitive landraces flowering in long day (*boro*) season (April-May 2021).

Based on the previously recorded booting dates and unmistakable contrasting morphological characters of the landraces[27], we selected a total of 26 pure lines of cultivars from Basudha's repertoire of 1480 landraces, based on matching anthesis dates and FOT, and contrasting morphological characters (**Table 2**). The PP and the OP landraces were selected in order that they differed in at least 1 morphological trait. Among the 26 landraces, 16 were used as pollen parents and 19 as ovary parents for the crosses (**Suppl. Table S1**), comprising 19 parental pairs during *aman* and 8 pairs during *boro* season (**Suppl. Table S1**), totalling 25 pairs including 5 repeats.

Table 2: Countries of Origin and Heritable Morphological Characters[27],[28] of the Landraces Used in Pair-wise Cross-Pollination Experiments.

Landrace Code	Origin	Pollen- or Ovary Parent	Flag Leaf Attitude	Awn: Max Length (mm)	Hull Colour	Bran Colour	Kernel: Notched Belly %
AA19	India	PP & OP	Droopy	0	Straw	Brown	0
B103	Sri Lanka	PP & OP	Semi Erect	0	Straw	Light Brown	0
C34	India	PP	Erect	14	Black	Black	0
C44	India	OP	Semi Erect	0	Brown furrows	Brown	0
DD16	India	OP	Erect	25	Gold furrows	White	0
G25	India	OP	Horizontal	0	Purple furrows	White	0
H01	India	PP & OP	Horizontal	0	Gold furrows	White	0
H19	India	OP	Erect	0	Gold furrows	White	0
H40	Sri Lanka	OP	Erect	0	Brown furrows	Light Brown	0
J33	India	PP & OP	Horizontal	0	Gold furrows	White	0
K131	India	PP	Droopy	37	Gold furrows	White	>80%
K169	India	OP	Horizontal	54	Gold furrows	Light Brown	0
L59	India	PP	Horizontal	0	Brown furrows	White	0
M120	India	PP	Erect	30	Gold furrows	White	0
M121	India	PP	Semi Erect	0	Brown (tawny)	White	0
M84	Philippines	OP	Erect	0	Straw	Light Brown	0
M63	India	OP	Erect	0	Straw	White	0
M91	India	PP & OP	Erect	0	Brown (tawny)	White	0
M94	India	PP & OP	Semi Erect	3	Brown (tawny)	Brown	0
N01	India	PP & OP	Semi Erect	2	Straw	Light Brown	0
Q11	India	PP & OP	Semi Erect	11	Straw	White	0
S72	India	PP & OP	Erect	0	Straw	White	0
S87	India	OP	Horizontal	6	Straw	White	0
S20	India	OP	Erect	0	Straw	White	0
T05	India	PP	Semi Erect	0	Purple furrows	Black	0
V17	India	PP	Semi Erect	62	Black furrows	White	>80%

Selection of morphological characters

Each of the 26 landraces had already been assessed for 56 morphological characters to maintain genetic purity over the past 26 years (http://cintdis.org/basudha). We selected 5 visually distinctive morphological characters so that the PP and the OP of each pair differed in at least one of these 5 characters. Most of these characters were previously employed in crossing experiments by many authors[11-18].

Fig. 1. Planting Design of Two Experimental Plots with a Single Ovary Parent (OP, *in red circle*), Surrounded by 24 Pollen Parents (PP), Planted in a Density of 16 m^{-2}. Plot A1 and Plot A2 Represent Different Pairs of PP and OP, planted during *Aman* season. Plot B1 and Plot B2 Represent Two Reciprocal Pairs (the OP in Plot B1 is the PP in Plot B2) Planted During *Boro* Season.

Experimental design

The experimental design of the crossing pairs was identical to our previous experiment[26]. Experiments were set up on Basudha farm in two cultivation seasons – *Aman* (sown in June 2020, harvested in December 2020), and *Boro* (sown in February 2021, harvested in May 2021). During *Aman* 2020, 13 pairs of landraces

were Selected, and 5 of the 13 pairs were replicated in two plots each. Thus the total number of plots was 18. In each plot, a single seedling of a landrace, demarcated as the pollen recipient ovary parent (OP), was planted in the centre, around which 24 seedlings of a different landrace as the pollen parent (PP) were planted in a 5 x 5 matrix with 25 cm-spacing (**Fig. 1**, Plot A1 and Plot A2). The central position of the OP plant ensured the possible reception of pollen from all the surrounding PP plants, regardless of wind direction and speed. All seedlings were uniformly 14-day old when transplanted.

During *boro* season, 4 pairs of landraces were selected for reciprocal crossing, to examine whether the pollen flow from PP to OP remains effectively the same when the role of the parents are swapped. The arrangement of the each pair was reciprocally reversed in a corresponding neighbouring plot, such that each PP in a plot was planted as OP, and *vice versa*, in another plot (**Fig. 1**, Plot B1 and Plot B2). The total number of experimental plots with different pairs of these landraces, flowering during summer 2021, was 8.

At the booting stage, a thin polyethylene sheet was vertically placed as a barrier to pollen flux between the neighbouring plots. The height of the barrier was 1.5 m, always above the top panicle level in each plot.

Recording of FOT, FCT, FED and EED of the parental pairs

After the anthesis of each landrace, we recorded the first flower opening time (FOT) and the last floret closing time (FCT) in each varietal plot. The methods are given in detail in ref. [4]. The longest duration of flowers remaining open (= flower opening duration, FED) was the interval between the FOT and the FCT of the last floret on a given varietal plot. As the transfer of pollen is possible only during the overlap of FED of OP and PP, we calculated the effective exposure duration (EED) of the OP florets, or the time widest window available for the OP's florets receiving pollen. For any period during which both PP and OP florets remain open simultaneously, EED > 0, and EED = 0 otherwise.

Collection of F1 progeny seeds

After maturity of seeds, panicles of the F1 progeny were harvested from the single OP plant from each plot. The OP panicles of *Aman* season were harvested in January 2021, and the OP panicles from *Boro* season were harvested in April 2021.

Cultivation of F1 plants for assessment of gene flow frequency

All the F1 seeds from each panicle with labels bearing the parental pairs and corresponding plot numbers were sown in separate nursery beds in June 2021. After germination, 14- to 16-d old seedlings were transplanted in July 2021 in two rows in separate demarcated plots, where the soil had been prepared following standard organic method, with no synthetic agrochemical inputs.

Characterisation of the F1 offspring and CPF estimation

Every plant of the F1 progeny, germinated from viable F1 seeds, was visually examined for the selected morphological characters at maturity, namely, flag leaf angle, and the presence of awn. All panicles of the F1 plants were harvested after maturity in January 2022, and their seeds were assessed for hull (lemma and palea) colour, bran (pericarp) colour and the frequency of 'notched belly' trait (**Table 2**), to discern if any trait was inherited from the respective PP of the progeny.

Assessment of gene flow frequency

The detection of at least one of the 5 contrasting characters of the PP in the F1 progeny would indicate the success of cross pollination. Since all these characters are a result of interactions of multiple genes, crosses between two landraces with contrasting traits is likely to produce offspring with variable quantifiable morphotypes. In particular, at least one of the offspring of a long-awned PP crossed with an awnless OP may possess smaller awns on most grains. In this case, the presence of awn would be a clear evidence of inheritance of the PP trait, regardless of the exact length of the awn in the offspring.

The cross pollination frequency (CPF) is a measure of the inheritance of at least 1 morphological character that is absent in OP, and was calculated as

$$V = 1 - P\,|_{F1 = OP} \qquad (eq.\ 1)$$

where V is the frequency of inheritance of non-OP characters in the viable progeny (germinated from non-sterile seeds), and P is the fraction of the viable F1 offspring inheriting all OP characters. However, considering that inter-varietal cross pollination often results in variable proportions of F1 sterility[29, 30,31], the equation 1 was modified to include the sterile seeds for estimation of the total transmission of PP genes to the offspring, including the sterile seeds:

$$CPF = (N1.V + N2\ S)/(N1+N2) \qquad (eq.\ 2)$$

where N1 is the number of viable offspring, N2 is the number of sterile seeds, and S is the F1 sterility %. When all F1 seeds are sterile, CPF = S. The background frequency of sterile grains in all the landraces selected for our experiments is 0% - 2%; therefore, any grain sterility significantly exceeding this background maximum rate of 2% (that is, $S > 2\%$) was inferred to be a result of CPF.

Results

Phenological details of flowering in the PP and OP landraces in all experimental plots, as well as the range of FED and the EED for each pair of crosses are given in **Fig. 2**. Using a total of 16 landraces as PP and 18 landraces PP, we planted selected pairs in a total of 26 experimental plots (including 5 replicates), of which 18 plots were established during *Aman* and 8 plots during *Boro* season (**Suppl. Table S1**). However, the entire progeny of one of these crosses (Q11 x S87 in plot A11) was lost due to insect pest attack before

harvest. Therefore, we were unable to estimate the CPF from this pair. Furthermore, The F1 progeny of three crosses (plot A11, plot B2 and plot B4) produced 100% sterile grains (**Suppl. Table S2**).

Fig. 2. The Range of FED (green bar), Beginning from FOT and Ending at FCT, with the Overlapping Period (blue) Indicating Effective Exposure Duration (EED) for Each Pair of Landraces. The column on the right margin shows the EED (in minutes) for each pair.

High F1 sterility in some crosses

The data presented in **Fig. 2** reveal that the EED of all pairs of landraces in this experiment ranged from 23 min (Plot A12a and Plot A12b) to 76 min (Plot A8). The mean of EED was 50.3 min during *Aman* season, and 56.8 min during *Boro* season. Each OP was expected to yield a progeny with a non-zero probability of inheritance of at least 1 trait from the corresponding PP. Most of the F1 progeny of the crosses engendered sterile seeds.

The crosses between the landraces in plot A13 (T05 x K169), plot B2 (B103 x AA19) and plot B4 (H01 x M94) produced 100% sterile grains. Some of the other crosses also yielded a high proportion of sterile grains: the crosses between the landraces grown in plot A1, plot A7a, plot A7b, plot A8, plot A12 a, plot A12 b, plot B6 and plot B7 produced >60% sterile grains borne on the respective OP's panicle (**Suppl. Table S2**). By contrast, plot A9 (M121 x Q11) produced no sterile seeds, and the crosses in plot A5 (V16 x M84), plot A6a and plot A6b (V17 x S20) respectively yielded 3.2%, 0.2% and 1.2% sterile seeds (in excess of the max. background rate of 2%). When the proportions of F1 sterility were plotted against EED for all crosses, the slope of regression was significant at 99% confidence level (**Fig. 3**).

Fig. 3: Regression of % Sterility of Grains in the Progeny of Crossed Parents against EED (min). Slope of regression = —1.2, *df* = 25, R^2 = 0.32, *t* = 3.33, *p* < 0.002.

High frequency of gene flow in all varietal pairs

The frequency of inheritance of PP traits in each progeny from the experimental plots assigned to the corresponding parental pairs (**Suppl. Table S2**) was estimated using equation 2. Excluding the progeny of 2 pairs destroyed by pest attack, 16 progeny lines was observed to have inherited > 60% of PP characters. No

PP character was visible in the viable offspring in plot 1A (C34 x H19), implying zero inheritance in the viable offspring, although 80.4% of the total F1 seeds was sterile.

A sufficiently wide time window is logically conducive to successful cross pollination. Thus, the CPF is expected to be associated with the length of EED. When the proportion of inheritance of PP traits is plotted against EED, a definitive trend of CPF escalating with EED is evident, which can best described as a strong ($p < 0.00001$) 3rd degree polynomial relationship: % Inheritance = 177.7 — 0.002 EED^3 + 0.22 EED^2 - 7.93 EED (**Fig. 4**).

Fig. 4: The Association of the Frequency of PP Trait Inheritance with EED of the OP, Best Described as a Polynomial Function ($df = 25$, $R^2 = 0.55$, $t = 5.37$, $p < 0.00001$).

Discussion and Conclusion

In the trivial case of zero- or near-zero overlap of FED between the PP and OP (EED ≈ 0), CPF ≈ 0; and this justifies the reports of zero or near-zero CPF in most of the previous cross pollination experiments (ref. **1-9, 12-15**). In our previous experiment[26], one of the two parental pairs had an EED of ca. 10 min, and showed CPF = 0. In that same experiment, the FED overlap of 80 min was associated with CPF >80% — a result that is corroboration in the present study. Because pollen remains viable for no longer than 9 min after anther dehiscence, the long duration of FED overlap does not imply extension of pollen longevity. Rather, it implies that the ovary of an open floret remains receptive to viable pollen released from successively opening multiple florets borne on other panicles.

Two consequences of inter-varietal cross pollination

Cross pollination results in the transmission of the PP genes to the F1 progeny, but the inheritance does not follow simple Mendelian pattern. Because most of the rice characters are polygenic, different combinations of the genes from the pollen with the genes in the ovary produces an array of phenotypic descriptors. In

addition, a large number of pleiotropic genes linked different biological pathways, governing heading date, grain length, plant height, flag leaf angle, and other characters[33, 34].

One of the 5 characters selected in this study is the 'flag leaf angle' phenotype, which is expressed in 4 categories, namely, erect, semi-erect, horizontal and droopy[28]. The cross between M94 with semi-erect flag leaf (as PP) and H01 with horizontal flag leaf (as OP) produced a viable progeny with erect (15%) and semi-erect (18%), as well as horizontal flag leaf (67%). Both the erect and semi-erect characters are non-OP characters, although the PP is a pure line with entirely semi-erect flag leaf. In the same pair, the pericarp colour of 55% of F1 seeds was light brown, although the rice kernels of both OP and PP had white bran. Likewise, the cross between L59 and G25 (plot A14), both parents characterised by horizontal flag leaf and white bran, produced 66% of the progeny with semi-erect flag leaf and 56% with light brown pericarp. Such (apparently unpredictable) departures from the parental characters in the offspring is because flag leaf angle and bran colour are complex phenotypes under polygenic control, with various associations of different genes on different chromosomes, each with multiple alleles[35-37]. Low angles of flag leaf (erect and semi-erect) are partially dominant[38], and light brown bran colour is governed by a dominant gene in the absence of a complementary gene[39, 40]. The combinations of the multiple genes and their alleles in the zygote may result in the expression of an assortment of colour and flag leaf angles not exactly matching the character of either parent.

A second consequence of cross pollination is a range of seed sterility frequency, orchestrated by genetic interactions between the parental gametes F1 sterility, as reported in several studies (summarized in ref. [29] and ref. [30]). Low F1 fertility has been recorded mostly in *Oryza sativa* ssp. *indica* x *O. Sativa* ssp. *japonica* crosses, but also in some inter-varietal *indica* x *indica* crosses,[30],[31] where the sterility genes determined by allelic interaction seem to be of wide occurrence. Different combinations of the *HSA1a* and *HSA1b* genes in *indica* rice are currently known to confer sterility in F1 and F2 hybrids .[30], [32]

The reciprocal crossing trials in 8 plots show that at least two pairs yielded a strikingly asymmetric result: In plot B1, the number of fertile grains of the progeny was 54, whereas the progeny of the reciprocal cross in Plot B2 produced no viable offspring (100% F1 sterility). Likewise, the cross between the pair grown in Plot B3 produced 66 offspring, whereas the reciprocal cross between the same landraces grown in Plot B4 produced only sterile grains (**Table 3**). Asymmetry in F1 sterility was also found in other crosses when a landrace is the PP, compared with the same landrace as OP. When Q11 was the PP in three crosses (plot 10a, plot 10b and plot 11), F1 sterility ranged from 43% to 57%, whereas with Q11 as the OP (plot 9), F1 sterility was 0% (**Suppl. Table S2**).

Our observation of the two consequences of cross pollination in rice — F1 sterility (in excess of the background rate of 2%) and inheritance of a combination of assorted phenotypes the viable offspring — is summarised in a flowchart (**Fig. 5**), without allusion to biochemical and molecular genetic explanations, which are beyond the scope of this paper. This flowchart highlights the importance of FED overlap as well as the parental genotypes for cross pollination success.

The implication of F1 sterility in CPF estimation

Because grain sterility is a consequence of cross pollination itself, all the flowers of the OP plants in both Plot B2 and Plot B4 were cross pollinated (CPF =100). Taken together, our study presents a convincing empirical evidence of considerably high CPF between *indica* landraces, as long as the overlap of FED between PP and OP flowers is of sufficiently long (>20 min) duration. The high frequency of CPF and gene flow reported here is in sharp contradiction to the prevalent knowledge of 'less than 2% CPF' presented in all biology text books and official rice biology reports. The high frequency of inheritance of PP traits by the progeny documented in this study establishes the factual possibility of high CPF between landraces in field conditions, corroborating our earlier finding [26].

Fig. 5. Flowchart Describing the Phenomenological Observations of Rice Phenology, Cross Pollination, and its Consequences in Field Condition.

The results of this study seem to be conclusive of the actuality of naturally high CPF in rice under circumstances of considerable FED overlap between parental cultivars, and indicates the necessity of further exploration into the biochemical pathways and genetic interactions of sterile grain production and assortments of polygenic traits in the offspring from crosses between different varietal genotypes. The latter

investigation may consolidate our contention that contrary to the the extremely low CPF that is generally accepted in the mainstream rice literature over a century, CPF in field conditions is much higher, depending on FED overlap between neighbouring cultivars. These findings are of profound importance in successful hybridization efforts (by selecting cultivars with wider EED); and conversely, in efforts to maintain genetic purity of rice landraces (by planting multiple landraces next to one another with non-overlapping EEDs). The rice breeders' and farmers' concern of unintended genetic contamination of their crops from transgenic (such as herbicide tolerant) rice pollen flow can also be addressed by de-synchronising the FOT and FED of the OP from that of the transgenic PP cultivars.

References

1 Sharngapani SG 1928. A few observations of paddy (*Oryza sativa*) crossing. *Agriculture Journal of India*. 19, 48-50.

2. Kobayasi K, Masui H, Atsuta Y, Matsui T, Yoshimoto M, Hasegawa, T. 2009. Flower opening time in rice —Cultivar difference and effect of weather factors. *Proceedings of the MARCO Symposium*; Tsukuba, Japan. 5–7 October 2009. Available from: https://www.naro.affrc.go.jp/archive/niaes/marco/marco2009/english/W2-12_Kobayasi_Kazuhiro_P.pdf

3. Deb, D., 2022, 'Rice flower opening time and flower opening duration data', https://doi.org/10.7910/DVN/KYW1UC, Harvard Dataverse, V10

4. Deb D, Joshi NV, Bhattacharya D, Nauri M, Ganguly R 2023. Diversity of flower opening time and duration in rice (*Oryza sativa* ssp. *indica*) landraces of South and Southeast Asia in different cultivation seasons. *J. Bot Res* 5(3): 36-48. doi: 10.30564/jbr.v5i3.5724

5. van der Stok, J.E., 1910. Onderzorkingen omtrent rijst en tweede gewassen (Dutch) [Experiments with rice and secondary crops]. G. Kolff. Co.: Batavia.

6. Thompstone E 1915. Some observations on Upper Burma paddy (grown under irrigation). The Agricultural Journal of India. 10: 26-5.

7. Torres, J.P., 1923. Some notes on rice hybridization work. *Philippine Agric. Rev.*16: 46-48.

8. Jones JW 1924. Observations on the time of blooming of rice flowers. *J. Amer. Soc. Agron.* 16 (10), 665-670.

9. Pope M N. 1916. The mode of pollination in some farm crops. Journal of the American Society of Agronomy. 8(4), 209-227.

10. Beachell H M, Adair C R, Jodon N E, Davis L. L, Jones J W 1938. Extent of natural crossing in rice. *J. Am. Soc. Agron.* 30: 743–653.

11. da Silva E F, da Silva LM, Montalván R 2005. Crossing rate and distance in upland rice. *Bragantia* 64(2): 197–201.

12. Coste, R 1969. *El Arroz: Colección Agricultura Tropical*, Editorial Blume, Barcelona, Spain, 1969, 2nd edn, pp. 35–72.

13. Endo T Sato H, Yamaguchi M., Kataoka T, Nakagomi K, Ito T, Mori K. 2009. Estimate of outcrossing rates in a rice plant (*Oryza sativa* L.) under field conditions using a purple grain rice cultivar. *Breed. Sci.* 59: 195-202.

14. Jodon N E 1959. Occurrence and importance of natural crossing in rice. *Rice J.:* 62(8): 8–10.

15. Reaño R, Pham JL 1998. Does cross-pollination occur during seed regeneration at the International Rice Genebank? *IRRN* 23: 5–6.

16. Robert EH, Craufurd RQ, Cochet FL 1961. Estimation of percentage of natural cross-pollination: experiment on rice. *Nature* 190: 1084–1085.

17. Sahadevan P C, Namboodiri K. M N 1963. Natural crossing in rice. *Proc. Indian Acad. Sci*. Sect. B. 58: 176–185.

18. Somaratne L H M Y K, Abayawickrama A S M T, Wickramasinghe I P, Samarasinghe W L G 2012. Estimating outcrossing rate of Bg 379-2 using morphological markers and confirmation by molecular markers. *Rice Sci*. 19(2): 166–168.

19. Moldenhauer K A K, Gibbons J H 2003. Rice morphology and development, pp. 103–127. In *Rice: Origin, History, Technology and Production* (eds Smith, C. W. and Dilday, R. H.), Wiley, Hoboken, NJ, USA.

20. OGTR 2005. *The Biology and Ecology of Rice (Oryza sativa L.) in Australia*, Office of the Gene Technology Regulator, Woden, Australia.

21. Langevin S A, Clay K, Grace J B 1990. The incidence and effect of hybridization between cultivated rice and its related weedy red rice (*Oryza sativa* L.). *Evolution* 44: 1000–1008.

22. Rong J. et al. 2007. Dramatic reduction of crop-to-crop gene flow within a short distance from transgenic rice fields. *New Phytol*. 173: 346–353.

23. Song Z P, Lu B.-R, Zhu Y G, Chen J K. 2003. Gene flow from cultivated rice to the wild species *Oryza rufipogon* under experimental field conditions. *New Phytol*. 157: 657–665.

24. Messeguer J, Marfà V, Català M M, Guiderdoni E, Melé E A. 2004. Field study of pollen-mediated gene flow from Mediterranean GM rice to conventional rice and the red rice weed. *Mol. Breed*. 13: 103–112.

25. Bae H.-K. et al., 2013. Evaluation of gene flow from GM to non-GM rice. *Plant Breed. Biotech.* 1(2): 162–170.

26. Deb D, Bhattacharya D 2021. Unusually high frequency of cross-pollination between rice landraces Shiuli and Khara with coincident flower-opening times. *Curr. Sci.* 121(1): 121-126. doi: 10.18520/cs/v121/i1/121-126

27. Deb D. 2005. *Seeds of Tradition, Seeds of Future*. Research Foundation for Science Technology & Ecolohgy: New Delhi.

28. Bioversity International, International Rice Res. Institute and the Africa Rice Center. *Descriptors for Wild and Cultivated Rice* (*Oryza* spp.). Bioversity International: Rome, Italy, 2007.

29. Miller R L 1958. The Inheritance of Sterility in an Intervarietal Hybrid of Rice, *Oryza Sativa* L. *LSU Historical Dissertations and Theses*. 498. https://repository.lsu.edu/gradschool_disstheses/498
DOI: 10.31390/gradschool_disstheses.498

30. Zhang Y, Wang J, Pu Q, Yang Y, Lv Y, Zhou J, Li J, Deng X, Wang M and Tao D. 2022. Understanding the nature of hybrid sterility and divergence of Asian cultivated rice. *Front. Plant Sci.* 13:908342. doi: 10.3389/fpls.2022.908342

31. Engle L M, Chang T T, Ramirez D A. 1969. The cytogenetics of sterility in F1 hybrids of *indica × indica* and *indica × javanica* varieties of rice (*Oryza sativa* L.). *Philipp. Agric*. 53: 289–307.

32. Kubo T, Takashi T, Ashikari M, Yoshimura A, Kurata N. 2016. Two tightly linked genes at the hsa1 locus cause both F1 and F2 hybrid sterility in rice. Mol. Plant. 9: 221–232. http://dx.doi.org/10.1016/j.molp.2015.09.014

33 Yao W, Li G, Yu Y, Ouyang Y 2018. funRiceGenes dataset for comprehensive understanding and application of rice functional genes. *GigaScience* 7: 1–9.

34. Li Z, Liu J, Wang X, Wang J, Ye J, Xu S, Zhang Y, Hu D, Zhang M, Xu Q, et al. *LG5*, a Novel Allele of *EUI1*, Regulates Grain Size and Flag Leaf Angle in Rice. *Plants*. 2023; 12(3):675. https://doi.org/10.3390/plants12030675

35. Zhang B, Ye W, Ren D, Tian P, Peng Y, Gao Y, Ruan B, Wang L, Zhang G, Guo L, Qian Q, Gao Z. 2015. Genetic analysis of flag leaf size and candidate genes determination of a major QTL for flag leaf width in rice. 2015. *Rice* 8: 2; DOI 10.1186/s12284-014-0039-9

36. Jiang J, Zhang Y, Li Y, Hu C, Xu L, Zhang Y, Wang D, Hong D and Dang X. 2022. An analysis of natural variation reveals that *OsFLA2* controls flag leaf angle in rice (*Oryza sativa* L.). *Front. Plant Sci.* 13: 906912. doi: 10.3389/fpls.2022.906912

37. Singh, G., Jyoti, S.D., Uppalanchi, P. *et al.* Genomic regions associated with flag leaf and panicle architecture in rice (*Oryza sativa* L.). *BMC Genomics* **25**, 1200 (2024). https://doi.org/10.1186/s12864-024-11037-z

38. Shen, F C. 1983. Inheritance of length, width, angle and ratio of dry weight over area of flag leaf in rice (*Oryza sativa* L.). *Guizhou Agric. Sci*. 6: 18–25.

39. Lee KE, Rahman M M, Kim JB, Kang SG. 2018. Genetic analysis of complementary gene interactions of *Pb* and *Pp* Genes for the purple pericarp trait in rice. *Journal of Life Science* 28(4): 398~407.

40. Yang W, Chen L, Zhao J, Wang J, Li W, Yang T, Dong J, Ma Y, Zhou L, Chen J, Wu W, Zhang S, Liu B. 2022. Genome-wide association study of pericarp color in rice using different germplasm and phenotyping methods reveals different genetic architectures. *Front. Plant Sci.* 13: 841191. doi: 10.3389/fpls.2022.841191.

41. Deb D. 2025. Phenological Data of Rice Cross Pollination - 2022. Harvard Dataverse, V2; https://doi.org/10.7910/DVN/TZCNYR.

Acknowledgements

We are immensely grateful to Prof. N V Joshi of Indian Institute of Science, Bangalore for his meticulous scrutiny of data and critical comments on earlier drafts of the mss. We are indebted to Sri Avik Saha for all logistic support to our work on Basudha farm and laboratory.

Funding

This work did not receive any institutional funding support.

Data availability

In addition to the data provided in Supplementary Tables, phenological details of 26 rice landraces are available on Harvard Dataverse [41]. Any other data will be freely available on reasonable request.

Author Contributions

Conception, experimental design, data curation, analysis, m/s writing and editing: D.D.
Field observation, data generation, documentation and validation: D.B. and M.N.

Conflict of interest

Authors declare that there is no conflict of interest.