TITLE: Size-stabilized, Hypoallometric, Genitalia Determined for Male Black Soldier Fly, *Hermetia*

illucens (Diptera: Stratiomyidae)

RUNNING HEAD: Hypoallometric Genitalia in Black Soldier Fly

AUTHORS: *Lisa N. Rollinson[1,3], *Noah B. Lemke[1], Scott Crawford[2], James B. Woolley[1], and Jeffery

K. Tomberlin[1]

* LNR and NBL are co-first authors.

AFFILIATIONS/ADDRESS

[1]Department of Entomology, Texas A&M University, 2475 TAMU, College Station, TX, USA

[2]Statistics Consulting Center, Texas A&M University, 3143 TAMU, College Station, TX, USA

[3]To whom correspondence should be addressed: Email: lisanrollinson@gmail.com

KEYWORDS:

Allometry, Reproduction, Morphology, Breeding

LAY SUMMARY:

Adult male black soldier fly shown to have similar sized genitals despite differences in body size.

SHORT SUMMARY (80 Words):

The black soldier fly, *Hermetia illucens,* is an economically important insect mass-reared throughout the

globe; however, a large knowledge gap exists in terms of its reproductive physiology. This study examined

the relationship of male genitalia to body size, finding a 10% increase in body size corresponded with a

1.8% increase in genitalia size, meaning the structures are hypoallometric. This finding is important because

it indicates little adverse consequences for selecting larger flies, as differently-sized individuals should be

able to copulate.

ABSTRACT:

The black soldier fly, *Hermetia illucens*, is mass-reared throughout the world to convert organic waste into ingredients for pet and livestock feed, as well as frass and other biological products. To promote the optimization of breeding regimes carried out by industrial black soldier fly production plants, it is critical to better understand adult reproductive physiology and its potential relationship with fertile egg production. However, in this species, little is known about how reproductive morphology changes with respect to increases in body size, nor the potential impacts selection can have on the form of reproductive structures. Hence, this study investigated the nutritional static allometric relationship between the external genitalia of adult male black soldier flies and their body size. Size differences were induced top-down by varying larval rearing densities, which has a downstream effect on adult body size. For each rearing density, a random sample of 30 individuals was selected, and measurements taken three times each for the head, thorax, parameral sheath, and gonostylus. Fitting a generalized linear log-log model to the data revealed that for every 10.0% increase in body size (thorax length), genital length (parameral shealth) only increased by 1.8%. The resulting allometric slope of genitalia to thorax size was 0.195, indicating a pattern of hypoallometry. The presence of hypoallometric genitalia in a domestic population indicates individuals can copulate irregardless of differences in body size, which is consistent with most other insects. Moreover, it is likely that the ancestral population of black soldier flies historically underwent selection to maintain a single genitalia-size independent of body size that continues to persist in captivity. To confirm, future work should investigate the direct impacts of hypoallometric genitalia on fitness, especially in flies which have been genetically edited or artificially selected to be increasingly large.

INTRODUCTION

The black soldier fly (BSF), *Hermetia illucens* (L. 1758) (Diptera: Stratiomyidae), is native to the southern United States as well as the American tropics (Kaya et al., 2021). It is bred industrially throughout the world for applications in animal feed and organic waste bioconversion. Over other insects, it is prized (Tomberlin & van Huis, 2020) not only for its ease of cultivation, but because its larvae can feed on most organic substrates and quickly bio-convert them into insect biomass and frass. Consider that in just a few weeks, BSF larvae can grow up to 10,000-times their body size, eating between 12.5 and 200-mg/day (Supriyatna et al., 2018; Dzepe et al., 2020). However, individual larvae are small, and so millions, if not billions, are needed to reach industrial targets, which can be as high as 60,000 metric tons of protein produced from waste conversion per year, or about 165 metric tons per day (C&E News, American Chemical Society). Companies have rallied around the promise of BSF in shaping future sustainable agricultural practices, and now billions of larvae are collectively produced per year across the globe (Barrett et al., 2023). But despite the fact that acquiring larvae relies on fertile egg production of the breeding colony, the reproductive biology of adult BSF remains understudied, especially compared to their larval counterparts (Lemke et al., 2023; Meneguz et al., 2023), as well as other mass-reared flies (Boller, 1972; Chambers, 1977; Scott et al., 2017). While the situation is not as critical as a decade ago (Čičková et al., 2015; Pastor et al., 2015), increased consistency and efficiency in breeding operations are both needed to reduce operational costs and drive down the final cost of industrial products (e.g., whole, dried, pulverized larvae, etc.). Therefore, this morphological study was designed as a first step to document the allometric relationships of male BSF genitalia of different size classes to aid ongoing optimization efforts of industrial breeding programs.

Historically, the focus of BSF-breeding programs has prioritized industrially relevant traits in larvae, such as increased larval mass, feed conversion ratios, or macronutrient profiles. Each of these has been accomplished both through artificial selection or genetic modification (Zhou et al., 2013; Zhan et al., 2020; Sandrock et al., 2022; Generalovic, 2023). Recent examples include an increase in size (Zhan et al.,

2020; Generalovic, 2023), differences in feed conversion rates (Zhou et al., 2013; Sandrock et al., 2022;

Generalovic et al.), and increased larval nutrients such as carotenoids (Gunther et al., 2024) in industrial or

research colony strains. Given these successes, greater attention should now be directed towards adult traits.

In other mass-reared insect species, such those reared for Sterile Insect Technique (SIT) (e.g.,

mosquitoes (Diptera: Culicidae), common fruit flies (Diptera: Drosophilidae), screwworm flies (Diptera:

Calliphoridae), and true fruit flies (Diptera: Tephritidae)), high-priority traits include size, longevity, flight

ability, and reproductive competitiveness (Lance & McInnis, 2021). In practice, however, translating such

traits 1:1 to BSF is not necessarily needed and may in fact be contrary to industrial aims; for example,

increased longevity would cause pre- and post-reproductives to overlap in cages using continuous release

(Lemke et al. 2025). The ultimate fate of mass-reared BSF is not wild release, meaning artificial selection

that reduces mating competitiveness in the wild does not need to be curbed. In practice this means,

compared to SIT species, there are fewer adult parameters relevant for BSF production. Even so, with little

species-specific knowledge of adult morphology, mating behavior, and sexual selection in BSF, it is

difficult to know *a priori* how these interact to increase fertile egg production before they are first

documented.

Certainly, increased larval size can translate to increased adult size and fecundity (Gobbi et al.,

2013), but the interplay of adult morphology and life history traits with other industrial parameters is

complex and not necessarily straightforward. Industrial producers typically operate by breeding large

numbers of adults, often in cages as large as 6 m long (S. Salari, 2024). Densities within these cages can

reach as high as 20,000 flies/m³ (S. Salari, 2024). With such high densities, there are too many flies for the

human eye to track, and densities even exceed the capabilities of computer vision or automated tracking

systems, since classification models can break down whenever individuals leave and re-enter the camera

frame (Nawoya et al., 2024). Previous studies, at greatly reduced spatio-temporal resolution, have examined

stocking density (Hoc et al., 2019), heterogeneity of cohort size (Jones & Tomberlin, 2021) and age

(Dickerson et al., 2024), as well as abiotic conditions (Li et al., 2024) and habitat design (Lemke et al.,

2024). To date, no studies have investigated the scaling of the external genitalia, which is almost certainly

linked to pre- and post-copulatory processes and ultimately fitness.

Male adult BSF have external genitalia which grasp the female, allowing them to attach for ~30

min during mating (Giunti et al., 2018; Masse et al., 2023; Manas et al., 2025). Specifically, lateral

extensions of the gonocoxite, which extend past the gonostylus, bend medially to grasp the female (Personal

communication, Martin Hauser). The parameral sheath then slots into the female and the three aedeagal

wands extend from the male's parameral sheath to release accessory fluid and sperm (personal

observations). In addition, any of these male genital structures could play a role in mating success via

postcopulatory female choice, since females might prefer certain shapes of genitals, which is the case for

other flies (e.g., crane flies (Diptera: Tipulidae) (Eberhard & Gelhaus, 2009)), and these in turn might be

affected by adult body size.

Black soldier fly genitalia structures may scale disproportionately with body size because the same

is true for other anatomical structures. For example, BSF eye length, ommatidia density, and wing centroids

do not scale proportionally with body size (Birrell, 2018). In addition, in laboratory settings, large flies, that

have presumably received more nutrition as larvae, were found to have proportionally smaller eyes than

those that received less nutrition as larvae because their eye size increased less than their body size (Birrell,

2018). Females exhibited larger antennae, wing centroids, and eyes, which may indicate sexual selection

acting on these traits (Birrell, 2018). Put simply, large BSF are not scale models of small BSF.

In terms of reproductive structures, BSF genitalia are likely to be size-stabilized, meaning selection

by females results in approximately one size of male genitalia irrespective of body size. Genital size

stabilization is exhibited by most other insects and has been documented in the vast majority of analysed

insect species (Eberhard, 2009). However, this pattern has yet to be confirmed in BSF. For these other

insects, when larval nutrition is increased, adult body size in turn increases, and indeed this is also the case

in BSF (Jones & Tomberlin, 2019). However, the rate at which genitalia size increases is typically much

slower than for body size (Eberhard, 2009). This trend of nutritionally mediated static hypoallometry

(NMSH) is widespread, but not ubiquitous, in insects, and indeed three alternatives exist. When compared

to a smaller conspecific, an individual might have proportionally smaller, similar, or larger genitalia, which

are referred to as hypo-, iso-, or hyper allometry, and indicated when a log-log regression is fit to the data

and is either less than 1, statistically equal to 1, or greater than 1; respectively (Shingleton, 2010). Genitalia

with low allometric slopes have been theorized to increase the total number of females with which a given

male can mate and stimulate effectively (Eberhard, 2009). By contrast, hyperallometric genitalia are much

rarer but may still play a role in sexual selection. For example, male scorpionflies (Mecoptera), use their

genitalia directly in battles with rival males (Johnson, 1995; Eberhard, 2009).

The objective of this study was to investigate the NMSA relationship of male BSF genitalia using

morphometrics. Following the convention that isometry is used as a null hypothesis (Eberhard, 2009), we

present the null that male external genitalia in the BSF will scale in direct proportion (log-log slope=1) with

body size. However, our alternative hypothesis, we expect a hypoallometric deviation to occur (log-log

slope<1.00), following the general trend found in other insects (Eberhard, 2009). More broadly, this is a

step to understand how breeding amongst different sized flies might be maintained in the midst of industrial

selection pressures.

METHODS

Because the goal was to analyse genital allometry across a range of adult body sizes, and body size

is inversely proportional to the degree of larval competition in BSF (Jones & Tomberlin, 2019), larvae were

reared at three densities of 5,000, 7,500, and 10,000 per 10.5 kg of substrate. Critically, these sizes are

reasonable to have been sampled haphazardly from colony flies.

Rearing Methods - Eggs were harvested from a BSF colony established at the Forensic Laboratory for

Investigative Entomological Sciences (FLIES) at Texas A&M University. Larvae were reared on 70%

moisture Gainesville (Producers Cooperative Association, Bryan, TX, USA) diet (Hogsette, 1992)

according to the FLIES colony SOP (see: Supplementary Info) until they reached 8-d-old. After this, larvae

were split into different density treatments (n = 3 levels, n = 2 replicates) of approximately 5,000, 7,500,

and 10,000 larvae (gravimetrically estimated). Aliquots of larvae were placed directly atop 7.5 kg

Gainesville feed at 70% moisture in 27 L pans (Sterilite Corporation, Townsend, MA, USA). An additional

0.75 kg of dry Gainesville was placed around the edge of the pans to prevent escape. For the remainder of

the rearing-cycle, pans were housed in a walk-in incubator maintained at approximately 25°C, 50% relative

humidity, and 14:10 L:D photoperiod (fluorescent lighting).

Contents of each pan were hand-mixed on days 11 and 12 to vent heat, prevent anoxic composting,

and give larvae better access to the undigested portions of the substrate. An additional kilogram of 70%

moisture Gainesville diet was supplemented to pans on days 15, 16, and 18. On day 26, when at least 40%

of the larvae began pupation (Bosch et al., 2020), the two pans of larvae per density-level were combined

prior to being sieved. Then, 150 g aliquots from each respective density-level were transferred to cylindrical

plastic containers (13.5x13.2 cm: d × h) to pupate (Dickerson et al., 2024; Lemke et al., 2024). 150 g dry

Gainesville was used as bedding material, and the tops of the containers were secured with Tulle fabric

(Wal-Mart Stores Inc., Little Rock, AR, USA).

Newly emerged adults (unsexed) of each density level were collected daily from cylindrical

pupation containers and placed in BugDorm-1 rearing-cages (30 × 30 × 30 cm) (InsectaBio, Riverside, CA,

USA) housed within the same walk-in incubator for a minimum of 12 h, but not more than 36 h, so their

exoskeleton could sclerotize prior to being euthanized. During these procedures, mating was never observed

between flies within the Bug-Dorms, which aligns with established notions that mating occurs later in the

reproductive ontogeny (Tomberlin & Sheppard, 2002; Kobelski et al., 2024). The flies that emerged on

days 37-40, were then frozen in a -20°C freezer and transferred to 0.95-liter freezer bags for long-term

storage. Flies from each treatment level were combined in a 3.79-liter freezer bag and commingled.

Sample Preparation – Specimens were sorted based on external genitalia, and females discarded. Male

specimens from each treatment were selected on the basis that they did not have damage to their heads or

bodies. The sclerotized portion of the genitalia were removed whole and dry-mounted between two glass

slides. Unscleritorized tissue was removed with HTS 144S7 4.5" straight squeeze scissors (Hobby Tool

Supply USA LLC, Fallbrook CA, USA). The epandrium was also removed to improve the visibility of the

parameral sheath and gonostylus against the flat plane of the microscope slide. Dissected genitalia were

secured in square wells measuring approximately 4×4 mm created between two Corning 2947 Micro

Slides (Corning Glass Works, Corning, NY, USA) and two layers of Duck Brand masking tape (Shurtape

Technologies, LLC, Hickory, NC, USA). This allowed the genitalia to be stored without being crushed

(Supplementary Figure 1).

The head was removed from the thorax by cutting through the membranous neck. Each head was

mounted through the center, ventral to the antennae, using a #2 Physus pin (Making it Work Inc.,

Franklintown, PA, USA). Each thorax was pinned according to conventional standards, i.e., by placing a

#2 pin through the right side.

Structures- Because the parameral sheath is sclerotized, easily visible, and in direct contact with the female,

it was chosen as a good proxy for genital size. The gonostylus was chosen as a secondary proxy for genital

size because it less visible but is in contact with more of the genitalia and may help maintain its shape. This

investigation was limited to male external genitalia because female external genitalia are unsclerotized

(Munsch-Masset et al., 2023) and difficult to measure accurately; this is common in genital allometry

investigations.

Measurements- The length of the head, thorax, and parameral sheath were determined using a Leica M205

FA with a DFC 450 C lens operated with Leica LAS X software (Version 3.7.24914.5). These structures

were selected because they are sclerotized in BSF, and therefore putatively less prone to distortion from

mounting (personal observation, LNR). To account for error, each structure was measured on three separate

occasions. Between repeated measurements, the pinned head and thorax were moved from the specimen

stage and repositioned. Likewise, the genitalia structures were also moved from the specimen stage,

although they remained mounted in the wells of glass slides.

The length of each structure was automatically calculated using the integrated measure function

included in LAS X (Version 3.7.24914.5) by superimposing a line over the digital magnifications. How

these lines were drawn in reference to other anatomical reference points was defined by Birrell (2018),

though we note that in their work head length and thorax length are instead referred to as eye length and

pronotum length [sic].

Per Birrell (2018), head length was defined as the distance from the dorsal to ventral boundary of

the inner margin of the right compound eye, and thorax length was taken from the anterior of the pronotum

to the posterior of the scutellum. The parameral sheath was measured medially, perpendicular to its

intersection into the rest of the genitalic structure. The length of the gonostylus was taken from the distal

tip to a line perpendicular to the base of the parameral sheath (Figure 1).

Statistical Analyses - Raw data was cleaned in MS Excel (Version 2308 Build 16731.20170) by LNR to

indicate damaged specimens (n = 4) and later analyzed using R Studio (Version 4.2.2) by SC at the Texas

A&M Statistical Consulting Center. For each of the three rearing density levels, 30 flies were measured,

less the damaged specimens, yielding 86 total replicates (5,000 n = 28, 7,500 n = 29, 10,000 n = 29) (Table

1).

Because each fly replicate was re-measured three times, a model needed to be devised that could

account for measurement error. A generalized linear log-log model with random effect for 'fly' using

backwards variable selection to model *P*-values based on Satterthwaite degrees of freedom with Kenward-

Roger approximation was chosen (which we will abbreviate as a 'random effects model'). Alpha for the

analyses was set at 0.05.

Model Selection- Although preliminary results comparing untransformed data to a log-log transformation

were similar, the latter was preferred to be consistent with allometric Power-Scaling Laws seen in the

general literature (Shingleton et al., 2007). Preliminary modelling also revealed that thorax and head lengths

were highly positively correlated (R^2 = 0.90), and statistically significant ($P = 2.2*10^{-16}$), with a quadratic

interaction between increasing length of parameral sheath and larval density. Still, larval density was

removed from the final model because it is an expected artifact of nutrition affecting body size (Jones &

Tomberlin, 2019). To simplify the model further, the random effects model only considered the effects of

thorax on genitalia (parameral sheath, gonostylus). Head size was omitted because thorax length is a more

standard proxy for body size.

Dimensional Reduction - In addition to the random effects model, principal component analysis (PCA) was

performed (Borcard et al., 2011) on the untransformed data to explore how thorax, head, sheath, gonostylus,

and larval density were related to one another in multidimensional space. (Supplementary Figure 2) Because

much of the variability could be explained by the first two principal components (>75%), only these were

retained to make model interpretation tractable (Supplementary Figure 3; Supplementary Table 1).

RESULTS

Chosen Model- The generalized linear log-log model with backwards variable selection found that the slope

of the logarithmic line, α, was highly significantly different from 1 (two-tailed $|P|$< 0.006, df = 193.74).

Therefore, the null hypothesis of isometry (slope=1) was rejected (Table 2).

The random effects model produced the following equation:

$$\ln(Sheath)=-7.61413+0.19468\ln(Thorax)$$

On a log-log plot, this equation appeared as a straight line with a slope (α) of 0.19468 and a y-intercept of

-7.61413 (Figure 2).

A 10% increase in the size of the thorax was shown to translate to a 1.87% increase in sheath size.

Some sample projections are provided in Table 3. Study data support a 10% increase in size. Values greater

than this were extrapolated for achievable increases on body size based on industry press releases (Varol,

2023) and artificial selection studies (Generalovic Thesis).

Principal Component Analysis (PCA) indicated most of the variance in density, thorax, and head

were explained by principal component one, while the variance for gonostylus and sheath are almost solely

explained by principal component two. Sheath length had a slight positive association with principal

component two, indicating its correlation with head and thorax length. In addition, PCA indicated that

thorax and head are closely related, with an inverse relationship to density.

DISCUSSION

The aim of this study was to document the nutritionally mediated static allometry of adult male

BSF external genitalia. The data collected indicate that the log-log relationship between adult BSF thorax

length and male parameral sheath length significantly deviated from 1.0. Thus, the null hypothesis of

isometry was rejected, and the alternative confirmed: male BSF genitalia were found to be hypoallometric

with respect to body size ($P = 0.005$), which aligns with patterns found in most other insects studied thus

far. The models developed in this study specifically indicate a nutritionally mediated 10% increase in thorax

length, the proxy for body size, correlated with just a 1.87% increase in parameral sheath length, the proxy

for genitalia size. Effectively, this means both large and small BSF from the same research colony have

similarly sized genitalia relative to their body size. These findings affirm that while many aspects of adult

BSF reproduction are size-dependent, external male genital morphology is relatively size-independent; this

could be relevant to industrial producers seeking to simplify their selection regimes by reducing the number

of parameters being considered for their breeding programs. Still, although only a 5.87% proportional

increase in genitalia relative to body size was observed (10% thorax divided by 1.87% parameral sheath),

insects may be sensitive to even small changes, being small themselves. Future studies linking genital size

increases directly with fitness are needed to confirm.

Indeed, in other insect species, the size and shape of genitalia has been demonstrated to affect

reproductive success. Although sexual selection on genitalia is difficult to directly quantify (Eberhard,

2011), larger genital size has been positively correlated with paternity rates and negatively associated with

sperm rejection in tortoise beetles (Coleoptera: Chrysomelidae) (Rodríguez et al., 2004). In addition,

reducing male genitalia size through ablation was shown to alternatively reduce reproductive success in a

seed bug (Hemiptera: Lygaeidae) (Dougherty et al., 2015), increases female sperm rejection in a tortoise

beetle (Coleoptera: Chrysomelidae) (Rodriguez, 1995), and can reduce copulatory success in fruit flies

(Diptera: Drosophilidae) (Grieshop & Polak, 2012). Likewise, altering the scaling of insect genitalia

through genetic modifications can also negatively impact reproductive success. For instance, *Drosophila*

melanogaster Meigan 1830 (Diptera: Drosophilidae) genitalia size have been experimentally reduced by

genetically editing the flies to make Forkhead box O (FOXO) activity, which slows growth when nutrition

is low, constitutively active. Males with posterior genital lobes that were smaller were less likely to copulate

and produced fewer offspring (Dreyer & Shingleton, 2011). These studies hint that the maintenance of

appropriately sized hypoallometric genitalia could be important to the reproductive success of many insects,

possibly including black soldier flies.

Extrapolating further, the hypoallometic genitalia of BSF may be a result of similar selective

pressures as in other insects and may be similarly important in reproductive success. In insects broadly

(Eberhard, 2009) past selection events have acted on genital structures to maintain hypoallometric

relationships, and the same may be occurring in BSF, though this is unconfirmed. Theory indicates that

over evolutionary time, a single optimal genital size could have been achieved through three alternative

routes: (1) stabilizing selection (Eberhard et al., 1998), (2) two opposing forms of directional selection

(Dreyer & Shingleton, 2011), or (3) different strengths of selection on different sizes of flies (Eberhard et

al., 2009). In terms of reproductive ecology, adult female BSF may have had exhibited either a choice for

a single preferred size of male genitalia (Eberhard et al., 1998) or two co-occurring mating strategies that

selected for smaller genitalia in large flies and larger genitalia in small flies (Eberhard et al., 2009; Dreyer

& Shingleton, 2011). Preserved over time, either of these cases could result in a single preferred size of

male genitalia and thus a hypoallometric relationship, as observed in the flies of our study. In other species,

this choice for genital size has been postulated to be driven by increased stimulation and is not necessarily

based on a precise mechanical 'locking' fit because the female genitalia are flexible and yielding (Eberhard

et al., 1998; Munsch-Masset et al., 2023).

LIMITATIONS

This study was limited to linear measurements of external adult male genitalia in a single population of BSF and offers a single, preliminary data set to guide future research. Linear measurements of the thorax, head, and genitalia may not fully explain the three-dimensional changes in morphology and may be subject to measurement error, but the use of multiple structures and a model incorporating repeated measurements was used to partially mitigate this. As in many genital allometry studies, female structures were not measured due to their non-sclerotized nature, although attempts were made. Additionally, the nature of purely morphological data limits predictions of sexual selection and reproductive success related to BSF genital size, especially across populations and strains. The allometric slope of male genitalia has also been shown to differ across populations (Kawano, 2002; Bertin & Fairbairn, 2007; Eberhard et al., 2009) and may do so in BSF. The exact genetic provenance of this population is not fully known, although the EVO Conversion Systems strain descended from the original Sheppard strain, which has served for numerous populations globally.

As in many genital allometry studies, female structures were not measured due to their non-sclerotized nature, although attempts were made. The nature of purely morphological data limits predictions of sexual selection and reproductive success related to BSF genital size, especially across populations and strains. The allometric slope of male genitalia has also been shown to differ across populations (Kawano, 2002; Bertin & Fairbairn, 2007; Eberhard et al., 2009) and may do so in BSF. The shifts in mating systems that BSF have undergone in industry may have shifted the exact allometric relationship, however determining differences between populations is beyond the breadth of the current study. Given the ubiquity of hypoallometry in insects (Eberhard et al., 1998), theories of selection maintaining hypoallometric genitalia (Eberhard et al., 1998), and the mechanisms by which sexual selection acts on genital size in other insects (Dreyer & Shingleton, 2019), it is probable that this hypoallometry is consistent across different BSF strains.

FUTURE DIRECTIONS

This investigation acts as preliminary research documenting the allometric relationships in male

BSF external genitalia. Future studies should investigate the direct relationship between size-stabilized

genitalia on fitness by examining how increases in body size and corresponding increases in genital

parameters act to influence mating success and fertile egg production. Differences in adult mass of just 50%

have been shown to increase hatch variability by 40 times (Jones & Tomberlin, 2021); however, the reason

for this variability has yet to be determined, and could be due to physiological or behavioral mechanisms,

or some combination of the two. Examining the behavior, egg production, and hatch rate of individual

mating pairs of different sizes could elucidate the reason for hatch variability in heterogeneous populations.

In addition, future research should consider using genetically modified strains, so that the results

can be better generalized to industry, where improvement through traditional selection and genome

modification can increase mean pupal body size by 26% (Generalovic Thesis) and 50% (Zhan et al., 2020;

Varol, 2023) respectively, with similar numbers being reported in industry press releases. Genetic editing,

especially of insulin pathways, could theoretically impact allometric relationships as well, and impact

mating success. However, over time natural selection could reinforce effective genitalia sizes and allometric

relationships in the edited population. Hence, understanding how artificial selection and gene-editing

interplay to affect adult morphology and reproductive success will be important for understanding the

dynamics of sexual selection during mass-breeding.

ACKNOWLEDGEMENTS

This material is based upon work supported by the National Science Foundation Graduate Research

Fellowship under Grant No. 1746932. Any opinion, findings, and conclusions or recommendations

expressed in this material are those of the authors and do not necessarily reflect the views of the National

Science Foundation.

The authors would like to thank the Forensic Laboratory for Investigative Entomological Sciences

(FLIES) Facility and the Texas A&M Department of Entomology for funding. In addition, the authors

would like to thank J. Emmanuel Mendoza for his perspectives and assistance, Davíd Salazar for his

support, and Dr. Chelsea Miranda for her assistance in rearing neonates.

CONFLICT OF INTEREST

Material for use in this project was purchased from EVO Conversion Systems LLC, a company with which

Dr. Tomberlin has a significant financial interest. This conflict of interest is managed by a plan submitted

to and approved by Texas A&M University and Texas A&M AgriLife.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable

request.

REFERENCES

Barrett M, Chia S, Fischer B & Tomberlin J K. (2023) Welfare considerations for farming black

soldier flies, *Hermetia illucens* (Diptera: Stratiomyidae): a model for the insects as food

and feed industry. *Journal of Insects as Food and Feed* 9:119–148.

https://doi.org/10.3920/JIFF2022.0041

Bertin A & Fairbairn D J (2007) The form of sexual selection on male genitalia cannot be

inferred from within-population variance and allometry? A case study in *Aquarius*

remgis. *Evolution* 61:825–837. https://doi.org/10.1111/j.1558-5646.2007.00074.x

Birrell N (2018) Sensory Morphology, Investment and Plasticity in the Black Soldier Fly

(*Hermetia illucens*). Thesis. University of Aukland.

Boller E (1972) Behavioral aspects of mass-rearing of insects. *Entomophaga* 17:9–25.

https://doi.org/10.1007/BF02371070

Borcard D, Gillet F & Legendre P (2011) Numerical ecology with R. *Springer*, New York, NY.

Bosch G, Oonincx DGAB, Jordan HR, Zhang J, van Loon JJA, van Huis A & Tomberlin JK

(2020) Standardisation of quantitative resource conversion studies with black soldier fly

larvae. Journal of Insects as Food and Feed 6:95–109.

https://doi.org/10.3920/JIFF2019.0004

Chambers DL (1977) Quality control in mass rearing. *Annual Review of Entomology* 22:289–

308. https://doi.org/10.1146/annurev.en.22.010177.001445

Čičková H, Newton GL, Lacy RC & Kozánek M (2015) The use of fly larvae for organic waste

treatment. *Waste Management* 35:68–80. https://doi.org/10.1016/j.wasman.2014.09.026

Dickerson AJ, Lemke NB, Li C & Tomberlin JK (2024) Impact of age on the reproductive

output of *Hermetia illucens* (Diptera: Stratiomyidae). J*ournal of Economic*

Entomology:toae107. https://doi.org/10.1093/jee/toae107

Dougherty LR, Rahman IA, Burdfield-Steel ER, Greenway EV (Ginny) & Shuker DM (2015)

Experimental reduction of intromittent organ length reduces male reproductive success in

a bug. *Proceedings of the Royal Society B: Biological Sciences* 282:20150724.

https://doi.org/10.1098/rspb.2015.0724

Dreyer AP & Shingleton AW (2011) The effect of genetic and environmental variation on

genital size in male *Drosophila*: canalized but developmentally unstable. PLOS ONE

6:e28278. https://doi.org/10.1371/journal.pone.0028278

Dreyer AP & Shingleton AW (2019) Insulin-insensitivity of male genitalia maintains

reproductive success in *Drosophila*. Biology Letters 15:20190057.

https://doi.org/10.1098/rsbl.2019.0057

Dzepe D, Nana P, Fotso A, Tchuinkam T & Djouaka R (2020) Influence of larval density,

substrate moisture content and feedstock ratio on life history traits of black soldier fly

larvae. *Biosaintifika* 10: 448–454. https://doi.org/10.3920/JIFF2019.0034

Eberhard WG (2009) Static allometry and animal genitalia. *Evolution* 63:48–66.

https://doi.org/10.1111/j.1558-5646.2008.00528.x

Eberhard WG (2011) Experiments with genitalia: a commentary. *Trends in Ecology & Evolution*

26:17–21. https://doi.org/10.1016/j.tree.2010.10.009

Eberhard WG & Gelhaus JK (2009) Genitalic stridulation during copulation in a species of crane

fly, *Tipula* (*Bellardina*) sp. (Diptera: Tipulidae). *Rev. Biol. Trop.* 57.

Eberhard WG, Huber BA, S. RLR, Briceño RD, Salas I & Rodriguez V (1998) One size fits all?

Relationships between the size and degree of variation in genitalia and other body parts in

twenty species of insects and spiders. *Evolution* 52:415–431.

https://doi.org/10.1111/j.1558-5646.1998.tb01642.x

Eberhard W, Rodriguez RL & Polihronakis M (2009) Pitfalls in understanding the functional

significance of genital allometry. *Journal of Evolutionary Biology* 22:435–445.

https://doi.org/10.1111/j.1420-9101.2008.01654.x

Generalovic TN (2023) Population genomics and domestication of the Black Soldier Fly

(*Hermetia illucens* L.). Thesis. Apollo - University of Cambridge Repository.

doi:10.17863/CAM.103274.

Generalovic TN, Sandrock C, Leonard S, Schuldiner-Harpaz T, Pipan M, Welch JJ & Jiggins

CD Variation in strain performance and estimates of heritability of body size indicate

considerable potential for genetic improvement of the black soldier fly (*Hermetia*

illucens). *Entomologia Experimentalis et Applicata* 173:558–574.

https://doi.org/10.1111/eea.13565

Giunti G, Campolo O, Laudani F & Palmeri V (2018) Male courtship behaviour and potential for

female mate choice in the black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae).

Entomologia Generalis 38:29–46. https://doi.org/10.1127/entomologia/2018/0657

Gobbi P, Martínez-Sánchez A & Rojo S (2013) The effects of larval diet on adult life-history

traits of the black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *European*

Journal of Entomology 110:461–468. https://doi.org/10.14411/eje.2013.061

Grieshop K & Polak M (2012) The precopulatory function of male genital spines in *drosophila*

ananassae[doleschall] (diptera: drosophilidae) revealed by laser surgery. *Evolution*

66:2637–2645. https://doi.org/10.1111/j.1558-5646.2012.01638.x

Gunther D, Alford R, Johnson J, Neilsen P, Zhang L, Harrell R & Day C (2024) Transgenic

black soldier flies for production of carotenoids. *Insect Biochemistry and Molecular*

Biology 168:104110. https://doi.org/10.1016/j.ibmb.2024.104110

Hoc B, Noël G, Carpentier J, Francis F & Megido RC (2019) Optimization of black soldier fly

(*Hermetia illucens*) artificial reproduction. *PloS one* 14:e0216160.

https://doi.org/10.1371/journal.pone.0216160

Hogsette JA (1992) New diets for production of house flies and stable flies (Diptera: Muscidae)

in the laboratory. *Journal of Economic Entomology* 85:2291–2294.

https://doi.org/10.1093/jee/85.6.2291

Johnson NF (1995) Variation in male genitalia of *Merope tuber* Newman (Mecoptera:

Meropeidae). *Journal of the Kansas Entomological Society* 68:224–233.

Jones BM & Tomberlin JK (2019) Impact of larval competition on life-history traits of the black

soldier fly (Diptera: Stratiomyidae). *Annals of the Entomological Society of America*

112:505–510. https://doi.org/10.1093/aesa/saz014

Jones BM & Tomberlin JK (2021) Effects of adult body size on mating success of the black

soldier fly, *Hermetia illucens* (L.) (Diptera: Stratiomyidae). *Journal of Insects as Food*

and Feed 7:5–20. https://doi.org/10.3920/JIFF2020.0001

Kawano K (2002) Character displacement in giant rhinoceros beetles. *The American Naturalist*

159:255–271. https://doi.org/10.1086/338512

Kaya C, Generalovic TN, Ståhls G, Hauser M, Samayoa AC, Nunes-Silva CG, Roxburgh H,

Wohlfahrt J, Ewusie EA, Kenis M, Hanboonsong Y, Orozco J, Carrejo N, Nakamura S,

Gasco L, Rojo S, Tanga CM, Meier R, Rhode C, Picard CJ, Jiggins CD, Leiber F,

Tomberlin JK, Hasselmann M, Blanckenhorn WU, Kapun M & Sandrock C (2021)

Global population genetic structure and demographic trajectories of the black soldier fly,

Hermetia illucens. *BMC Biology* 19:94. https://doi.org/10.1186/s12915-021-01029-w

Kobelski A, Hempel A-J, Padmanabha M, Klüber P, Wille L-C & Streif S (2024) Model-based

process optimization of black soldier fly egg production. *Frontiers in Bioengineering and*

Biotechnology 12. https://doi.org/10.3389/fbioe.2024.1404776

Lance DR & McInnis DO (2021) Biological Basis of the Sterile Insect Technique. Sterile Insect

Technique. 2nd edn. *CRC Press*.

Lemke N, Dickerson A & Jeffery T (2023) No neonates without adults: A review of adult black

soldier fly biology, *Hermetia illucens* (Diptera: Stratiomyidae). *BioEssays* 45:2200162

Lemke NB, Rollison LN & Tomberlin JK (2024) Sex-Specific Perching: Monitoring of artificial

plants reveals dynamic female-biased perching behavior in the black soldier fly,

Hermetia illucens (Diptera: Stratiomyidae). *Insects*

15:770.https://doi.org/10.3390/insects15100770

Li C, Addeo NF, Rusch TW, Chappell TM, Tarone AM & Tomberlin JK (2024) Age and

calorific restriction impact immature black soldier fly (Diptera: Stratiomyidae) thermal

tolerance and preference. *Journal of Insects as Food and Feed* 11:259–272.

Manas F, Labrousse C & Bressac C (2025) Plastic responses in sperm expenditure to sperm

competition risk in black soldier fly (*Hermetia illucens*, Diptera) males. J*ournal of Insect*

Physiology 161:104751. https://doi.org/10.1016/j.jinsphys.2025.104751

Masse M, Paul Serge M, Loudh M, Yede, Libong N, Glwadys S, Totuom M, Sidoine C, Bilong

Bilong C, Felix C, Loudh G, Yede N, Libong S, Glwadys M, Totuom C, Sidoine B, &

Bilong (2023) Differences in life history traits and morphology in wild vs. domesticated

populations of black soldier fly, *Hermetia illucens* (Diptera: Stratiomyidae). *Animal and*

Veterinary Sciences 11:71–79. https://doi.org/10.11648/j.avs.20231103.13

Melody M. Bomgardner. (2020). ADM brings insect protein to the US. *C&EN Global*

Enterprise 98: 13–13. https://doi.org/10.1021/cen-09846-buscon3

Meneguz M, Miranda CD, Cammack JA & Tomberlin JK (2023) Adult behaviour as the next

frontier for optimising industrial production of the black soldier fly *Hermetia illucens*

(L.) (Diptera: Stratiomyidae). *Journal of Insects as Food and Feed* 9:399–414.

https://doi.org/10.3920/JIFF2022.0055

Munsch-Masset P, Labrousse C, Beaugeard L & Bressac C (2023) The reproductive tract of the

black soldier fly (*Hermetia illucens*) is highly differentiated and suggests adaptations to

sexual selection. *Entomologia Experimentalis et Applicata* 171:857–866.

https://doi.org/10.1111/eea.13358

Nawoya S, Ssemakula F, Akol R, Geissmann Q, Karstoft H, Bjerge K, Mwikirize C, Katumba A

& Gebreyesus G (2024) Computer vision and deep learning in insects for food and feed

production: A review. *Computers and Electronics in Agriculture* 216:108503.

https://doi.org/10.1016/j.compag.2023.108503

Pastor B, Velasquez Y, Gobbi P & Rojo S (2015) Conversion of organic wastes into fly larval

biomass: bottlenecks and challenges. *Journal of Insects as Food and Feed* 1:179–194.

https://doi.org/10.3920/JIFF2014.0024

Rodriguez V (1995) Relation of flagellum length to reproductive success in male *Chelymorpha*

alternans Boheman (Coleoptera: Chrysomelidae: Cassidinae). *The Coleopterists Bulletin*

49:201-205

Rodríguez V, Windsor DM & Eberhard WG (2004) Tortoise beetle genitalia and demonstrations

of a sexually selected advantage for flagellum length in *Chelymorpha alternans*

(Chrysomelidae, Cassidini, Stolaini). *Brill*. https://doi.org/10.1163/9789004475335_063

Salari S P, de Goede M L (2024) Study into *Hermetia illucens* behaviour in large artificial

mating chambers and possible optimizations. *Journal of Insects as Food and Feed* 10:1-

353. https://doi.org/10.1163/23524588-20241013

Sandrock C, Leupi S, Wohlfahrt J, Kaya C, Heuel M, Terranova M, Blanckenhorn WU,

Windisch W, Kreuzer M & Leiber F (2022) Genotype-by-diet interactions for larval

performance and body composition traits in the black soldier fly, *Hermetia illucens*.

Insects 13:424. https://doi.org/10.3390/insects13050424

Scott MJ, Concha C, Welch JB, Phillips PL & Skoda SR (2017) Review of research advances in

the screwworm eradication program over the past 25 years. *Entomologia Experimentalis*

et Applicata 164:226–236.https://doi.org/10.1111/eea.12607

Shingleton AW (2010) Allometry: The Study of Biological Scaling. *Nature Education*

Knowledge 3:2.

Shingleton AW, Frankino WA, Flatt T, Nijhout HF & Emlen DouglasJ (2007) Size and shape:

the developmental regulation of static allometry in insects. *BioEssays* 29:536–548.

https://doi.org/10.1002/bies.20584

Supriyatna A, Kurrahman OT, Cahyanto T, Yuliawati A & Kulsum Y (2018) The potency of

black soldier larvae (*Hermetia illucens* L.) as a source of protein for livestock feed.

Biosaintifika: Journal of Biology & Biology Education 10:449–455.

https://doi.org/10.15294/biosaintifika.v10i2.14422

Tomberlin J k. & van Huis A (2020) Black soldier fly from pest to 'crown jewel' of the insects

as feed industry: an historical perspective. *Journal of Insects as Food and Feed* 6:1–4.

https://doi.org/10.3920/JIFF2020.0003

Tomberlin JK & Sheppard DC (2002) Factors Influencing Mating and Oviposition of Black

Soldier Flies (Diptera: Stratiomyidae) in a Colony. *Journal of Entomological Science*

37:345–352. https://doi.org/10.18474/0749-8004-37.4.345

Varol B (2023) FreezeM launches enhanced genetic line of Black Soldier Flies. *Feed & Additive*

Magazine.

Zhan S, Fang G, Cai M, Kou Z, Xu J, Cao Y, Bai L, Zhang Y, Jiang Y, Luo X, Xu J, Xu X,

Zheng L, Yu Z, Yang H, Zhang Z, Wang S, Tomberlin JK, Zhang J & Huang Y (2020)

Genomic landscape and genetic manipulation of the black soldier fly *Hermetia illucens*, a

natural waste recycler. *Cell Research* 30:50–60. https://doi.org/10.1038/s41422-019-

0252-6

Zhou F, Tomberlin JK, Zheng L, Yu Z & Zhang J (2013) Developmental and waste reduction

plasticity of three black soldier fly strains (Diptera: Stratiomyidae) raised on different

livestock manures. *Journal of Medical Entomology* 50:1224–1230.

https://doi.org/10.1603/ME13021

Figure 1. Photo of male black soldier fly genitalia with epandrium already removed to increase visibility to
the structures below. Colored measurement lines incidate the parameral sheath (pm sh) in blue and
gonostylus (gonst) in red.

Figure 2. Black soldier fly sheath length plotted against thorax length, with both axes log transformed. Data points are colored according to the larval rearing density (i.e., 10 000, 7 500, or 5 000 larvae/pan), and each point represents the average for each individual specimen (n = 30), which was measured three times.

TABLES

Table 1. Black soldier fly body and genitalia measurements (mm). Thorax and head lengths were used as proxies for body size, whereas the gonostylus and shealth were measured as proxies of genital size. Presented below are the group means ± SD and ranges (min, max) for these measurements according to each density that specimens were reared at (10 000, 7 500, or 5000 larvae per container). Treatment groups were pooled together to yield the grand mean and total range.

	10,000		7,500		5,000		Combined	
	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)
Thorax	5.012 ± 0.313	(4.278, 5.856)	5.262 ± 0.34	(4.433, 5.86)	5.678 ± 0.304	(5.067, 6.328)	5.313 ± 0.42	(4.278, 6.328)
Head	2.072 ± 0.094	(1.871, 2.318)	2.147 ± 0.098	(1.954, 2.304)	2.266 ± 0.077	(2.106, 2.433)	2.16 ± 0.12	(1.871, 2.433)
Gonostylus	0.391 ± 0.024	(0.345, 0.454)	0.401 ± 0.023	(0.349, 0.448)	0.388 ± 0.032	(0.315, 0.438)	0.393 ± 0.027	(0.315, 0.454)
Sheath	0.174 ± 0.011	(0.142, 0.199)	0.182 ± 0.011	(0.161, 0.207)	0.179 ± 0.015	(0.152, 0.221)	0.178 ± 0.013	(0.142, 0.221)

Table 2. Coefficients table. A generalized linear log-log model with backwards variable selection relating increases in adult male BSF genital size (parameral sheath) to increases in body size (Thorax).

	Estimate	Std. Error	df	T value	Pr(>|t|)
(Intercept)	-7.61413	0.36231	193.69667	-21.016	< 0.0000 ***
log(Thorax)	0.19468	0.06912	193.74983	2.817	0.00536 **

Table 3. Estimated projections for the absolute increase in male genital (sheath) size with respect to increases in body (thorax) size in the black soldier fly. Body size increases of 10% or less are interpolated

551   from this study's data. Beyond this range, the projected size increase to genitalia is extrapolated from other

552   studies or claims made by industry press releases.

Increase in Body Size	Reference	Increase in Genital Size
+5%	Interpolation	0.95%
+10%	Interpolation	1.87%
+25%	Artificial Selection (Generalovic Thesis)	4.44%
+50%	FreezeM Genetic Line (Varol, 2023)	8.21%
+100%	Extrapolation	14.4%

SUPPLEMENTARY INFO

FLIES Colony SOP

Oviposition traps of fluted cardboard were collected from the colony and transferred into a mason jar

covered by two layers of paper towel to allow airflow but prevent escape. During this time, the eggs hatched,

and neonates migrated into the bottom of the mason jar. Then, 0.5 g of neonates were transferred into each

of three square plastic containers. The neonates were fed with 500 g Gainesville feed (Hogsette 1992)

formulated at 70% moisture. A border with 50 g of dry feed was used to prevent larvae from escaping. The

black soldier fly larvae were reared in a walk-in incubator at approximately 27 °C.

Supplemental Figure 1. Genitalia secured between microslides in wells made of two layers of Duck Brand

masking tape.

Supplemental Figure 2. Principal component analysis (PCA) performed with the variables of thorax, head,

sheath, gonostylus, and larval density

Supplemental Figure 3. Scree plot demonstrating that principal components and two explain a sufficient

amount of the variability (>75%).

Supplemental Table 1. Principal Component Analysis

	PC1	PC2

Thorax	0.58361004	0.1027914
Head	0.58406062	0.0847625
Sheath	0.26451689	-0.6319838
Gonostylus	0.01790199	-0.7499736
Density	-0.49797876	-0.1427769