TITLE: Size-stabilized, Hypoallometric, Genitalia Determined for Male Black Soldier Fly, *Hermetia*

illucens (Diptera: Stratiomyidae)

RUNNING HEAD: Hypoallometric Genitalia in Black Soldier Fly

AUTHORS: *Lisa N. Rollinson[1,3], *Noah B. Lemke[1], Scott Crawford[2], James B. Woolley[1], and Jeffery

K. Tomberlin[1]

* LNR and NBL are co-first authors.

AFFILIATIONS/ADDRESS

[1]Department of Entomology, Texas A&M University, 2475 TAMU, College Station, TX, USA

[2]Statistics Consulting Center, Texas A&M University, 3143 TAMU, College Station, TX, USA

[3]To whom correspondence should be addressed: Email: lisanrollinson@gmail.com

KEYWORDS:

Allometry, Reproduction, Morphology, Breeding

LAY SUMMARY:

Adult male black soldier fly shown to have similar sized genitals despite differences in body size.

SHORT SUMMARY (80 Words):

The black soldier fly, *Hermetia illucens,* is an economically important insect mass-reared throughout the

globe; however, a large knowledge gap exists in terms of its reproductive anatomy and physiology. This

study examined the relationship of male genitalia to body size, finding a 10% increase in body size

corresponded with a 1.8% increase in genitalia size, meaning the structures are hypoallometric. This finding

is important because it indicates large and small flies have similar sized genitalia, which may allow

differently-sized individuals to copulate.

ABSTRACT:

The black soldier fly, *Hermetia illucens*, is mass-reared throughout the world to convert organic waste into

ingredients for pet and livestock feed, as well as frass and other biological products. To promote the

optimization of breeding regimes carried out by industrial black soldier fly operations, it is critical to better

understand adult reproductive anatomy and its potential relationship with fertile egg production. However,

in this species, little is known about how reproductive anatomy changes with respect to increases in body

size. Hence, this study investigated the nutritional static allometric relationship between the external

genitalia of adult male black soldier flies and their body size. Size differences were induced top-down by

increasing the larval rearing density, which has a negative effect on adult body size. For each of 3 rearing

densities, a random sample of 30 adults was selected, and measurements taken three times each for the

head, thorax, parameral sheath, and gonostylus. Fitting a generalized linear log-log model to the data

revealed that for every 10.0% increase in body size (thorax length), genital length (parameral shealth) only

increased by 1.8%. The resulting allometric slope of genitalia to thorax size was 0.195, indicating a pattern

of hypoallometry. The presence of hypoallometric genitalia in a domestic population suggests individuals

should be able to copulate regardless of differences in body size, which is consistent with most other insects.

Moreover, this finding implies that black soldier fly genitalia were hypoallometric prior to their

domestication and continues to persist within captive black soldier fly populations. To confirm, future work

should investigate the direct impacts of hypoallometric genitalia on fitness, especially in flies which have

been genetically edited or artificially selected to be increasingly large.

 INTRODUCTION

The black soldier fly (BSF), *Hermetia illucens* (L. 1758) (Diptera: Stratiomyidae), is native to the

Neotropics and has since been naturalized or introduced across most regions of the globe (Kaya et al. 2021).

It is bred industrially throughout the world for applications in animal feed and organic waste bioconversion

due to its ease of cultivation (Tomberlin and van Huis 2020), but also because BSF larvae can feed on most

organic substrates and quickly bio-convert them into biomass and frass. In just a few weeks, BSF larvae

can grow up to 10,000-times their body size. However, individual larvae are small, and so to reach industrial

targets, which can be as high as 60,000 tons of protein produced per year (= 165 metric tons per day) (C&E

News, American Chemical Society), millions if not billions of flies need to be bred by such operations.

As such, trillions of BSF larvae are collectively produced by the industry each year (Barrett and

Fischer 2023), and yet the reproductive biology of adult BSF remains understudied (Lemke et al. 2023,

Meneguz et al. 2023), especially compared to studies of larvae (Lemke and Puniamoorthy 2025), as well

as other mass-reared flies such as mosquitoes (Culicidae), true fruit flies (Tephritidae), and common fruit

flies (Drosophilidae), which have been studied for decades (Boller 1972, Chambers 1977, Scott et al. 2017).

For BSF, increased consistency and efficiency in breeding operations are both needed to reduce operational

costs so that valorised products (e.g., whole larvae, protein meal, larvae puree, lipids, frass, etc.) can better

compete in the marketplace. To keep operational costs low, BSF in practice often need to be reared on

nutritionally poor and/or heterogenous substrates, but this can produce a high degree of variability in body

size and possibly further increase the heterogeneity within breeding populations (due to overlapping cohorts

causing adults to have different ages, reproductive statuses, conditions, etc.) (Lemke et al. 2025).

Of course, variation in body-size may also affect reproductive success (Jones and Tomberlin 2021,

Julita et al. 2025) and egg production (Gobbi et al. 2013). A study found BSF breeding cohorts of mixed-

sizes had 2-3 times more failed mating attempts than same-sized cohorts, and a range of 48-343% more

eggs produced, though precise mechanisms are unknown (Jones and Tomberlin 2021). Recent work has

highlighted the potential for a physiological mechanisms (i.e., sperm competition, seminal fluid transfers)

(Munsch-Masset et al. 2023, Manas et al. 2024); however it is also possible that external genital anatomy

plays a role in structuring reproductive success, since male genitalia not only provide a physical stimulus

for females selection to act on (Eberhard and Gelhaus 2009), but also must theoretically be similar-sized to

female genitalia for the efficient transfer of gametes.

Form follows function, and male BSF possess external genitalia structures which allow them to

grasp the female, allowing them to remain attached for ~30 min during mating (Giunti et al. 2018, Masse

et al. 2023, Manas et al. 2025). Specifically, lateral extensions of the gonocoxite, which extend past the

gonostylus, bend medially to grasp the female (Personal communication, Martin Hauser). These structures

allow the male to remain linked to the female while switching from a 'from-behind' mount to an 'end-to-

end' position (Chiabotto et al. 2024, Lemke and Puniamoorthy 2025) potentially while in flight (personal

observation). The parameral sheath then slots into the female and the three aedeagal wands extend from the

male's parameral sheath to release accessory fluid and sperm (personal observations).

Of course, as BSF body size varies, genitalia structures may also scale disproportionately. The same

is true for other anatomical structures; for example, BSF eye length, ommatidia density, and wing centroids

do not scale proportionally with body size (Birrell 2018). In addition, lab-reared large BSF, that presumably

received more nutrition as larvae, were found to have proportionally smaller eyes than those that received

less nutrition as larvae because their eye size increased less than their body size (Birrell 2018). Females

exhibited larger antennae, wing centroids, and eyes, which may indicate sexual selection acting on these

traits (Birrell 2018).

Put simply, large BSF are not scale models of small BSF. When larval nutrition is increased, adult

body size increases in turn (Jones and Tomberlin 2019); however, typically the rate at which genitalia size

increases in organisms is typically much slower than for their body size (Eberhard 2009). This trend of

nutritionally-mediated static hypoallometry (NMSH) is widespread across the Insecta and Arachnida

(Eberhard 2009), but not ubiquitous, and indeed three alternatives exist: Larger individuals might have

proportionally smaller, similar, or larger genitalia than smaller conspecifics, which are referred to as hypo-,

iso-, or hyper allometry, and indicated when a log-log regression is fit to the data and is either less than 1,

statistically equal to 1, or greater than 1; respectively (Shingleton 2010). Genitalia with low allometric

slopes have been theorized to increase the total number of females with which a given male can mate and

stimulate effectively (Eberhard 2009). By contrast, hyperallometric genitalia are much rarer but may still

play a role in sexual selection. For example, male scorpionflies (Mecoptera), use their genitalia directly in

battles with rival males (Johnson 1995, Eberhard 2009).

The objective of this study was thus to investigate the NMSA relationship of male BSF genitalia

using morphometrics. Following the convention that isometry is used as a null hypothesis (Eberhard 2009),

we predict male external genitalia in the BSF will scale in direct proportion (log-log slope=1) with body

size. However, our alternative hypothesis, we expect a hypoallometric deviation to occur (log-log

slope<1.00), following the general trend found in other insects (Eberhard 2009). More broadly, this is a

first step to understand how breeding amongst different sized flies might influence reproductive success

amidst a plethora of selection pressures.

113   METHODS

Research was conducted at the Forensic Laboratory for Investigative Entomological Sciences

(FLIES) at Texas A&M University (College Station, Texas), where a BSF production colony is regularly

maintained, from which neonate larvae were acquired for our study. The BSF maintained at FLIES are

descended from the 'Sheppard Strain', originally maintained laboratory at the Coastal Plain Experiment

Station of the University of Georgia, Tifton, GA (Sheppard et al. 2002).

Because adult body size is inversely proportional to the degree of larval competition in BSF (Jones

and Tomberlin 2019), larvae were reared at three densities of 5,000, 7,500, and 10,000 per 10.5 kg of

substrate (Gainesville 'Housefly' diet (Hogsette 1992) acquired from Producers Cooperative Association

Bryan, TX, USA) prepared at 70% moisture. This produced a continuum of adults from three overlapping

size-classes, which was well within the range that could normally be expected to normally occur in our

production colony due to variations in nutrition.

Newly emerged adults (unsexed) reared from each density level were collected daily from their

pupation containers and placed temporarily in BugDorm-1 rearing-cages (30 × 30 × 30 cm) (InsectaBio,

Riverside, CA, USA) and kept inside a walk-in incubator maintained at approximately 25 °C, 50% relative

humidity, and 14:10 L:D photoperiod (fluorescent lighting). All adults used for this study housed within

the incubator for a minimum of 12 h, but not more than 36 h, so their exoskeleton could sclerotize prior to

being euthanized. During these procedures, mating was never observed between flies within the Bug-

Dorms, which aligns with established notions that mating occurs later in the reproductive ontogeny

(Tomberlin and Sheppard 2002, Kobelski et al. 2024) and in the presence of UV-light (Oonincx et al. 2016)

The flies were then frozen in a -20°C freezer and transferred to 0.95-liter freezer bags for long-term storage.

Flies from each treatment level were combined in a 3.79-liter freezer bag and commingled to randomize

them.

Sample Preparation – Specimens were sorted based on external genitalia, and females discarded.

Male specimens (n = 30) from each treatment were selected on the basis that they did not have damage to

their heads or bodies. The sclerotized portion of the genitalia were removed whole and dry-mounted

between two glass slides. Unsclerotized tissue was removed with HTS 144S7 4.5" straight squeeze scissors

(Hobby Tool Supply USA LLC, Fallbrook CA, USA). The epandrium was also removed to improve the

visibility of the parameral sheath and gonostylus against the flat plane of the microscope slide. Dissected

genitalia were secured in square wells measuring approximately 4 × 4 mm created between two Corning

2947 Micro Slides (Corning Glass Works, Corning, NY, USA) and two layers of Duck Brand masking tape

(Shurtape Technologies, LLC, Hickory, NC, USA). This allowed the genitalia to be stored without being

crushed (Supplementary Figure 1). The head was removed from the thorax by cutting through the

membranous neck. Each head was mounted through the center, ventral to the antennae, using a #2 Physus

pin (Making it Work Inc., Franklintown, PA, USA). Each thorax was pinned according to conventional

standards, i.e., by placing a #2 pin through the right side.

Structures- Because the parameral sheath is sclerotized, easily visible, and in direct contact with

the female, it was chosen as our proxy for genital size. The gonostylus was chosen as a secondary proxy

for genital size because it less visible but is in contact with more of the genitalia and may help maintain its

shape. This investigation was limited to male external genitalia because female external genitalia are

unsclerotized (Munsch-Masset et al. 2023) and difficult to measure accurately; this is common in genital

allometry investigations for which often only information is known about male structures (Eberhard et al.

1998).

Measurements- For each specimen, four measurements were taken: the length of the head, thorax,

and parameral sheath. These were determined using a Leica M205 FA with a DFC 450 C camera operated

with Leica LAS X software (Version 3.7.24914.5). These structures were selected because they are

sclerotized in BSF, and therefore putatively less prone to distortion from mounting (personal observation,

LNR). To account for error, each structure was measured on three separate occasions (i.e., 30 replicates x

4 measures x 3 repeated measures). Between sessions of repeated measurements, the pinned head and thorax

were moved off the specimen stage, replaced, and repositioned. The same was true for the glass slides

containing genitalia. The length of each structure was automatically calculated using the integrated measure

function included in LAS X by superimposing a line over the digital magnifications. How these lines were

drawn in reference to other anatomical reference points was defined by Birrell (Birrell 2018), though we

note that in their work head length and thorax length are instead referred to as eye length and pronotum

length [sic]. Per Birrell (2018), head length was defined as the distance from the dorsal to ventral boundary

of the inner margin of the right compound eye, and thorax length was taken from the anterior of the

pronotum to the posterior of the scutellum. The parameral sheath was measured medially, perpendicular to

its intersection into the rest of the genitalic structure. The length of the gonostylus was taken from the distal

tip to a line perpendicular to the base of the parameral sheath (Figure 1).

Statistical Analyses - Raw data was cleaned in MS Excel (Version 2308 Build 16731.20170) by

LNR to indicate damaged specimens (n = 4) and later analyzed using R Studio (Version 4.2.2) by SC at the

Texas A&M Statistical Consulting Center. For each of the three rearing density levels, 30 BSF were

measured, less the damaged specimens (5,000 n = 28, 7,500 n = 29, 10,000 n = 29) (Table 1).

Model Selection- Because each fly replicate was remeasured three times, a model accounting for

measurement error was developed. A generalized linear log-log model with random effect for 'fly' using

backwards variable selection to model *P*-values based on Satterthwaite degrees of freedom with Kenward-

Roger approximation was chosen (which we will abbreviate as a 'random effects model'). Alpha for the

analyses was set at 0.05. Although preliminary results comparing untransformed data to a log-log

transformation were similar, the log-log transformed data was preferred to be consistent with the convention

for describing allometric Power-Scaling Laws (Shingleton et al. 2007). Preliminary modelling also revealed

that thorax and head lengths were highly positively correlated ($R^2 = 0.90$), and statistically significant ($P =$

$2.2*10^{-16}$) (although there was a quadratic interaction between increasing length of parameral sheath and

larval density). Larval density (i.e., the treatment effect) was removed from the final model because it was

an expected artifact of nutrition affecting body size (Jones and Tomberlin 2019). To further simplify the

model further, we only considered the effects of thorax on genitalia (parameral sheath, gonostylus). Both

head and thorax length were correlated to genitalia size, and thorax length was chosen as the proxy for body

size because head size is sexually dimorphic (Birrell 2018). Lastly, because artificial selection can further

increase the body size of BSF beyond what we could artificially produce, the random effects model was

used to predict potential increases in genitalia size for BSF body-size increases up to 100%.

Dimensional Reduction - In addition to the random effects model, principal component analysis

(PCA) was performed (Borcard et al. 2011) on the untransformed data to explore how thorax, head, sheath,

gonostylus, and larval density were related to one another in multidimensional space (Supplementary Figure

2). Because much of the variability could be explained by the first two principal components (>75%), only

these were retained to make model interpretation tractable (Supplementary Figure 3; Supplementary Table

1).

RESULTS

Summary Statistics. The BSF we measured in our study across all 3 rearing densities/size-classes

had an average thorax measuring 5.313 ± 0.42 mm (Mean + SD). The smallest individual thorax length

measured was 4.278 mm and the largest thorax length was measured to be 6.328 mm. Likewise, head

measurements were 2.16 ± 0.12 mm on average, with a range of 1.871-2.433 mm. Gonostylus

measurements were 0.393 ± 0.027 mm on average and ranged from 0.315-0.454 mm. Finally, sheath

measured 0.178 ± 0.013 mm on average and ranged from 0.142-0.221 mm. Statistics for individual

treatment groups are also given in Table 1.

Random effects model. The random effects model found that the slope of the logarithmic line, α,

was highly significantly different from 1 (two-tailed $|P| < 0.006$, df = 193.74), and produced the following

equation: $\ln(Sheath) = 7.61413 + 0.19468 \ln(Thorax)$, which was then transformed to a log-log

relationship following the convention for modelling allometric relationships (Figure 2). This model

revealed that a 10% increase in the size of the thorax translated to just a 1.87% increase in sheath size. For

these reasons, the null hypothesis of isometry (slope=1) was rejected (Table 2). The model also predicts a

14.4% increase in sheath size when increases in BSF body size were extrapolated out to 100% (Table 3).

Dimensional reduction - Principal Component Analysis (PCA) indicated most of the variance in

density, thorax, and head were explained by principal component one, while the variance for gonostylus

and sheath are almost solely explained by principal component two. Sheath length had a slight positive

association with principal component two, indicating its correlation with head and thorax length. In

addition, PCA indicated that thorax and head are closely related, with an inverse relationship to density.

DISCUSSION

The aim of this study was to document the nutritionally mediated static allometry of adult male

BSF external genitalia. The data collected indicate that the log-log relationship between adult BSF thorax

length and male parameral sheath length significantly deviated from a slope of 1.0. Thus, the null hypothesis

of isometry was rejected, and the alternative confirmed: male BSF genitalia were found to be

hypoallometric with respect to body size ($P = 0.005$), which aligns with patterns found in most insects

studied this far, including flies, beetles, and ants (Eberhard 2009). The model developed in this study

specifically indicates a nutritionally mediated 10% increase in thorax length, the proxy for body size, is

correlated with just a 1.87% increase in parameral sheath length, the proxy for genitalia size. Effectively,

this means both large and small BSF from the same research colony have similarly sized genitalia. These

findings affirm that while many aspects of adult BSF reproduction are size-dependent (e.g., reproductive

success, egg production, sperm competition), external male genital morphology is relatively size-

independent, and indeed is confirmed also by the PCOA we conducted. These findings are relevant to

industrial producers seeking to simplify their selection regimes by reducing the number of parameters being

considered for their breeding programs, and increase the confidence that despite body size variations, the

size of male genitalia are largely unaffected (and thus should putatively allow effective reproduction across

size-classes, as has previously been demonstrated (Jones and Tomberlin 2021). Still, although only a 5.87%

proportional increase in genitalia relative to body size was observed (10% increase in thorax relative to a

1.87% increase in parameral sheath), insects may be sensitive to even small changes, being small

themselves (and indeed our model predicts a 14.4% increase in genital size when body size doubles, which

is not an insignificant amount). Future studies linking genital size increases directly with fitness are needed

to confirm this relationship.

Indeed, across the Insecta, the size and shape of genitalia have been demonstrated to affect

reproductive success. Although sexual selection on genitalia is difficult to directly quantify (Eberhard

2011), larger genital size has been positively correlated with paternity rates and negatively associated with

sperm rejection in tortoise beetles (Coleoptera: Chrysomelidae) (Rodríguez et al. 2004). In addition,

reducing male genitalia size through ablation was shown to alternatively reduce reproductive success in a

seed bug (Hemiptera: Lygaeidae) (Dougherty et al. 2015), increases female sperm rejection in a tortoise

beetle (Coleoptera: Chrysomelidae) (Rodriguez 1995), and can reduce copulatory success in fruit flies

(Diptera: Drosophilidae) (Grieshop and Polak 2012). Likewise, altering the scaling of insect genitalia

through genetic modifications can also negatively impact reproductive success. For instance, *Drosophila*

melanogaster Meigan 1830 (Diptera: Drosophilidae) genitalia size have been experimentally reduced by

genetically editing the flies to make Forkhead box O (FOXO) activity, which slows growth when nutrition

is low, constitutively active. Males with posterior genital lobes that were smaller were less likely to copulate

and produced fewer offspring (Dreyer and Shingleton 2011). These studies hint that the maintenance of

appropriately sized hypoallometric genitalia could be important to the reproductive success of many insects

(Eberhard 2009), possibly including black soldier flies.

As for how hypoallometric genitalia have been maintained in BSF, theory indicates that over

evolutionary time, a single optimal genital size could have been achieved through three alternative routes:

(1) stabilizing selection (Eberhard et al. 1998), (2) two opposing forms of directional selection (Dreyer and

Shingleton 2011), or (3) different strengths of selection on different sizes of flies (Eberhard et al. 2009). In

terms of reproductive ecology, adult female BSF may have had theoretically exhibited either a choice for a

single preferred size of male genitalia (Eberhard et al. 1998) or two co-occurring mating strategies that

selected for smaller genitalia in large flies and larger genitalia in small flies (Eberhard et al. 2009, Dreyer

and Shingleton 2011). Preserved over time, either of these cases could result in a single preferred size of

male genitalia and thus a hypoallometric relationship. In other species (e.g., crane flies (Diptera:

Tipuliidae)), this choice for genital size has been postulated to be driven by increased stimulation and is not

necessarily based on a precise mechanical 'locking' fit because the female genitalia are flexible and yielding

(Eberhard et al. 1998, Munsch-Masset et al. 2023).

Limitations and Future Work. This study was limited to linear measurements of external adult male

genitalia in a single population of BSF. Linear measurements of the thorax, head, and genitalia may not

fully explain the three-dimensional changes in morphology and may be subject to measurement error, but

the use of multiple structures and a random-effects model incorporating repeated measurements was used

to mitigate this. As in many genital allometry studies, female structures were not measured due to their non-

sclerotized nature, although attempts were made. Additionally, since measuring the genitalia is a destructive

process, we did not measure reproductive success related to BSF genital size. Thus, the door is open for

future studies to confirm whether a size-mismatch between male and female BSF genitalia is possible (or

whether there are negative fitness consequences that result). Of course, we also only took measurements of

BSF from a single population; and indeed, future work can also conduct morphometrics across global BSF

populations to document the pan-phenome. However, given the ubiquity of hypoallometry in insects and

spiders (Eberhard et al. 1998), theories of selection maintaining hypoallometric genitalia (Eberhard et al.

1998), and the mechanisms by which sexual selection acts on genital size in other insects (Dreyer and

Shingleton 2019), it is probable that hypoallometric genitalia remains a consistent feature across different

BSF strains. Lastly, since artificial selection can result in increased BSF body size, i.e., an increase in pupal

surface area by 15% (Generalovic 2023) and BSF weight 25% (Facchini et al. 2022), while gene editing

can also increase larval mass by 50% (Zhan et al. 2020, Varol 2023), with similar numbers being reported

by industry, future work should confirm that the allometric slopes remain constant under these increases.

ACKNOWLEDGEMENTS

This material is based upon work supported by the National Science Foundation Graduate Research

Fellowship under Grant No. 1746932. Any opinion, findings, and conclusions or recommendations

expressed in this material are those of the authors and do not necessarily reflect the views of the National

Science Foundation.

The authors would like to thank the Forensic Laboratory for Investigative Entomological Sciences

(FLIES) Facility and the Texas A&M Department of Entomology for funding. In addition, the authors

would like to thank J. Emmanuel Mendoza for his perspectives and assistance, Davíd Salazar for his

support, and Dr. Chelsea Miranda for her assistance in rearing neonates.

CONFLICT OF INTEREST

Material for use in this project was purchased from EVO Conversion Systems LLC, a company

with which Dr. Tomberlin has a significant financial interest. This conflict of interest is managed by a plan

submitted to and approved by Texas A&M University and Texas A&M AgriLife.

REFERENCES

Barrett M, Fischer B. 2023. Challenges in farmed insect welfare: Beyond the question of sentience. Anim. welf. 32:e4. https://doi.org/10.1017/awf.2022.5

Birrell N. 2018. Sensory Morphology, Investment and Plasticity in the Black Soldier Fly (*Hermetia illucens*) [Thesis]. University of Aukland. [accessed 2022 Aug 27]. https://researchspace.auckland.ac.nz/handle/2292/37256.

Boller E. 1972. Behavioral aspects of mass-rearing of insects. Entomophaga. 17(1):9–25. https://doi.org/10.1007/BF02371070

Borcard D, Gillet F, Legendre P. 2011. Numerical ecology with R. New York, NY: Springer.

Bosch G, Oonincx DGAB, Jordan HR, et al. 2020. Standardisation of quantitative resource conversion studies with black soldier fly larvae. Journal of Insects as Food and Feed. 6(2):95–109. https://doi.org/10.3920/JIFF2019.0004

Chambers DL. 1977. Quality Control in Mass Rearing. Annu. Rev. Entomol. 22(1):289–308. https://doi.org/10.1146/annurev.en.22.010177.001445

Chiabotto C, Grosso F, Doretto A, et al. 2024. Observation of mating behavior using marked flies of black soldier fly (Hermetia illucens) under sunlight condition. Journal of Insects as Food and Feed. 10(12):2017–2029.

Dickerson AJ, Lemke NB, Li C, et al. 2024 May 22. Impact of age on the reproductive output of *Hermetia illucens* (Diptera: Stratiomyidae). Journal of Economic Entomology.:1–10. https://doi.org/10.1093/jee/toae107

Dougherty LR, Rahman IA, Burdfield-Steel ER, et al. 2015. Experimental reduction of intromittent organ length reduces male reproductive success in a bug. Proceedings of the Royal Society B: Biological Sciences. 282(1808):20150724. https://doi.org/10.1098/rspb.2015.0724

Dreyer AP, Shingleton AW. 2011. The Effect of Genetic and Environmental Variation on Genital Size in Male *Drosophila*: Canalized but Developmentally Unstable. PLOS ONE. 6(12):e28278. https://doi.org/10.1371/journal.pone.0028278

Dreyer AP, Shingleton AW. 2019. Insulin-insensitivity of male genitalia maintains reproductive success in *Drosophila*. Biology Letters. 15(5):20190057. https://doi.org/10.1098/rsbl.2019.0057

Eberhard W, Rodriguez RL, Polihronakis M. 2009. Pitfalls in understanding the functional significance of genital allometry. Journal of Evolutionary Biology. 22(3):435–445. https://doi.org/10.1111/j.1420-9101.2008.01654.x

Eberhard WG. 2009. Static allometry and animal genitalia. Evolution. 63(1):48–66. https://doi.org/10.1111/j.1558-5646.2008.00528.x

Eberhard WG. 2011. Experiments with genitalia: a commentary. Trends in Ecology & Evolution. 26(1):17–21. https://doi.org/10.1016/j.tree.2010.10.009

Eberhard WG, Gelhaus JK. 2009. Genitalic stridulation during copulation in a species of crane fly, Tipula
(Bellardina) sp. (Diptera: Tipulidae). Rev. Biol. Trop. 57.

Eberhard WG, Huber BA, S. RLR, et al. 1998. One Size Fits All? Relationships Between the Size and
Degree of Variation in Genitalia and Other Body Parts in Twenty Species of Insects and Spiders.
Evolution. 52(2):415–431. https://doi.org/10.1111/j.1558-5646.1998.tb01642.x

Facchini E, Shrestha K, van den Boer E, et al. 2022. Long-term artificial selection for increased larval
body weight of *Hermetia illucens* in industrial settings. Frontiers in Genetics. 13. [accessed 2023 Nov
27]. https://www.frontiersin.org/articles/10.3389/fgene.2022.865490.

Generalovic TN. 2023. Population genomics and domestication of the Black Soldier Fly (Hermetia
illucens L.) [PhD Thesis]. University of Cambridge. https://doi.org/10.17863/CAM.103274.

Giunti G, Campolo O, Laudani F, et al. 2018. Male courtship behaviour and potential for female mate
choice in the black soldier fly *Hermetia illucens* L. (Diptera: Stratiomyidae). Entomologia Generalis.
38:29–46. https://doi.org/10.1127/entomologia/2018/0657

Gobbi P, Martínez-Sánchez A, Rojo S. 2013. The effects of larval diet on adult life-history traits of the
black soldier fly, Hermetia illucens (Diptera: Stratiomyidae). Eur. J. Entomol. 110(3):461–468.
https://doi.org/10.14411/eje.2013.061

Grieshop K, Polak M. 2012. The precopulatory function of male genital spines in *drosophila
ananassae*[doleschall] (diptera: drosophilidae) revealed by laser surgery. Evolution. 66(8):2637–2645.
https://doi.org/10.1111/j.1558-5646.2012.01638.x

Hogsette JA. 1992. New Diets for Production of House Flies and Stable Flies (Diptera: Muscidae) in the
Laboratory. Journal of Economic Entomology. 85(6):2291–2294. https://doi.org/10.1093/jee/85.6.2291

Johnson NF. 1995. Variation in Male Genitalia of *Merope tuber* Newman (Mecoptera: Meropeidae).
Journal of the Kansas Entomological Society. 68(2):224–233.

Jones BM, Tomberlin JK. 2019. Impact of Larval Competition on Life-History Traits of the Black Soldier
Fly (Diptera: Stratiomyidae). Annals of the Entomological Society of America. 112(6):505–510.
https://doi.org/10.1093/aesa/saz014

Jones BM, Tomberlin JK. 2021. Effects of adult body size on mating success of the black soldier fly,
Hermetia illucens (L.) (Diptera: Stratiomyidae). Journal of Insects as Food and Feed. 7(1):5–20.
https://doi.org/10.3920/JIFF2020.0001

Julita U, Permana A, Supriatna A, et al. 2025. Impact of Male and Female Body Size on Mating
Performance and Reproductive Success in Black Soldier Fly, Hermetia illucens L. (Diptera:
Stratiomyidae). J. Entomol. Res. Soc.:1. https://doi.org/10.51963/jers.v27i1.2684

Kaya C, Generalovic TN, Ståhls G, et al. 2021. Global population genetic structure and demographic
trajectories of the black soldier fly, *Hermetia illucens*. BMC Biology. 19(1):94.
https://doi.org/10.1186/s12915-021-01029-w

Kobelski A, Hempel A-J, Padmanabha M, et al. 2024. Model-based process optimization of black soldier
fly egg production. Front. Bioeng. Biotechnol. 12. https://doi.org/10.3389/fbioe.2024.1404776

Lemke N, Puniamoorthy N. 2025. Beyond the Black Box: Reproductive Strategies of the Black Soldier Fly as a Model for Bridging Evolutionary Biology and Applied Entomology.

Lemke NB, Dickerson AJ, Tomberlin JK. 2023. No neonates without adults. BioEssays. 45(1):2200162. https://doi.org/10.1002/bies.202200162

Lemke NB, Li C, Dickerson AJ, et al. 2025. Heterogeny in cages: Age-structure and attractant availability impacts fertile egg production in the black soldier fly, *Hermetia illucens*. Journal of Insects as Food and Feed. Online. https://doi.org/10.1163/23524588-bja10275

Lemke NB, Rollison LN, Tomberlin JK. 2024. Sex-Specific Perching: Monitoring of Artificial Plants Reveals Dynamic Female-Biased Perching Behavior in the Black Soldier Fly, *Hermetia illucens* (Diptera: Stratiomyidae). Insects. 15(10):770. https://doi.org/10.3390/insects15100770

Manas F, Labrousse C, Bressac C. 2025. Plastic responses in sperm expenditure to sperm competition risk in black soldier fly (*Hermetia illucens*, Diptera) males. Journal of Insect Physiology. 161:104751. https://doi.org/10.1016/j.jinsphys.2025.104751

Manas F, Piterois H, Labrousse C, et al. 2024. Gone but not forgotten: dynamics of sperm storage and potential ejaculate digestion in the black soldier fly Hermetia illucens. :2024.05.31.596011. https://doi.org/10.1101/2024.05.31.596011

Masse M, Paul Serge M, Loudh M, et al. 2023 June 6. Differences in Life History Traits and Morphology in Wild vs. Domesticated Populations of Black Soldier Fly, *Hermetia illucens* (Diptera: Stratiomyidae). :71–79. https://doi.org/10.11648/j.avs.20231103.13

Meneguz M, Miranda CD, Cammack JA, et al. 2023. Adult behaviour as the next frontier for optimising industrial production of the black soldier fly Hermetia illucens (L.) (Diptera: Stratiomyidae). Journal of Insects as Food and Feed. 9(4):399–414. https://doi.org/10.3920/JIFF2022.0055

Munsch-Masset P, Labrousse C, Beaugeard L, et al. 2023. The reproductive tract of the black soldier fly (*Hermetia illucens*) is highly differentiated and suggests adaptations to sexual selection. Entomologia Experimentalis et Applicata. 171(11):857–866. https://doi.org/10.1111/eea.13358

Oonincx DGAB, Volk N, Diehl JJE, et al. 2016. Photoreceptor spectral sensitivity of the compound eyes of black soldier fly (Hermetia illucens) informing the design of LED-based illumination to enhance indoor reproduction. Journal of Insect Physiology. 95:133–139. https://doi.org/10.1016/j.jinsphys.2016.10.006

Rodriguez V. 1995. Relation of Flagellum Length to Reproductive Success in Male *Chelymorpha alternans* Boheman (Coleoptera: Chrysomelidae: Cassidinae).

Rodríguez V, Windsor DM, Eberhard WG. 2004. Tortoise beetle genitalia and demonstrations of a sexually selected advantage for flagellum length in *Chelymorpha alternans* (Chrysomelidae, Cassidini, Stolaini). Brill. https://doi.org/10.1163/9789004475335_063

Scott MJ, Concha C, Welch JB, et al. 2017. Review of research advances in the screwworm eradication program over the past 25 years. Entomologia Experimentalis et Applicata. 164(3):226–236. https://doi.org/10.1111/eea.12607

Sheppard C, Tomberlin J, Joyce J, et al. 2002. Rearing methods for the black soldier fly (Diptera:
Stratiomyidae). Journal of Medical Entomology. 39(4):695–698. https://doi.org/10.1603/0022-2585-
39.4.695

Shingleton AW. 2010. Allometry: The Study of Biological Scaling. Nature Education Knowledge.
3(10):2.

Shingleton AW, Frankino WA, Flatt T, et al. 2007. Size and shape: the developmental regulation of static
allometry in insects. BioEssays. 29(6):536–548. https://doi.org/10.1002/bies.20584

Tomberlin J k., van Huis A. 2020. Black soldier fly from pest to 'crown jewel' of the insects as feed
industry: an historical perspective. Journal of Insects as Food and Feed. 6(1):1–4.
https://doi.org/10.3920/JIFF2020.0003

Tomberlin JK, Sheppard DC. 2002. Factors Influencing Mating and Oviposition of Black Soldier Flies
(Diptera: Stratiomyidae) in a Colony. Journal of Entomological Science. 37(4):345–352.
https://doi.org/10.18474/0749-8004-37.4.345

Varol B. 2023 Apr 26. FreezeM launches enhanced genetic line of Black Soldier Flies. Feed & Additive
Magazine. [accessed 2024 Apr 23]. https://www.feedandadditive.com/freezem-launches-enhanced-
genetic-line-of-black-soldier-flies/.

Zhan S, Fang G, Cai M, et al. 2020. Genomic landscape and genetic manipulation of the black soldier fly
Hermetia illucens, a natural waste recycler. Cell Res. 30(1):50–60. https://doi.org/10.1038/s41422-019-
0252-6

FIGURES

Figure 1. Photo of male black soldier fly genitalia with epandrium already removed to increase visibility to

the structures below. Colored measurement lines incidate the parameral sheath (pm sh) in blue and

gonostylus (gonst) in red.

Figure 2. Black soldier fly sheath length plotted against thorax length, with both axes log transformed. Data

points are colored according to the larval rearing density (i.e., 10 000, 7 500, or 5 000 larvae/pan), and each

point represents the average for each individual specimen (n = 30), which was measured three times.

TABLES

Table 1. Black soldier fly body and genitalia measurements (mm). Thorax and head lengths were used as

proxies for body size, whereas the gonostylus and shealth were measured as proxies of genital size.

Presented below are the group means ± SD and ranges (min, max) for these measurements according to

each density that specimens were reared at (10 000, 7 500, or 5000 larvae per container). Treatment groups

were pooled together to yield the grand mean and total range.

	10,000		7,500		5,000		Combined	
	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)	Mean ± SD	(Min,Max)
Thorax	5.012 ± 0.313	(4.278, 5.856)	5.262 ± 0.34	(4.433, 5.86)	5.678 ± 0.304	(5.067, 6.328)	5.313 ± 0.42	(4.278, 6.328)
Head	2.072 ± 0.094	(1.871, 2.318)	2.147 ± 0.098	(1.954, 2.304)	2.266 ± 0.077	(2.106, 2.433)	2.16 ± 0.12	(1.871, 2.433)
Gonostylus	0.391 ± 0.024	(0.345, 0.454)	0.401 ± 0.023	(0.349, 0.448)	0.388 ± 0.032	(0.315, 0.438)	0.393 ± 0.027	(0.315, 0.454)
Sheath	0.174 ± 0.011	(0.142, 0.199)	0.182 ± 0.011	(0.161, 0.207)	0.179 ± 0.015	(0.152, 0.221)	0.178 ± 0.013	(0.142, 0.221)

458 Table 2. Coefficients table. A generalized linear log-log model with backwards variable selection relating

459 increases in adult male BSF genital size (parameral sheath) to increases in body size (Thorax).

	Estimate	Std. Error	df	T value	Pr(>|t|)
(Intercept)	-7.61413	0.36231	193.69667	-21.016	< 0.0000 ***
log(Thorax)	0.19468	0.06912	193.74983	2.817	0.00536 **

461 Table 3. Estimated projections for the absolute increase in male genital (sheath) size with respect to

462 increases in body (thorax) size in the black soldier fly. Body size increases of 10% or less are interpolated

463   from this study's data. Beyond this range, the projected size increase to genitalia is extrapolated from other

464   studies or claims made by industry press releases.

Increase in Body Size	Reference	Increase in Genital Size
+5%	Interpolation	0.95%
+10%	Interpolation	1.87%
+25%	Extrapolation - Facchini 2022 increased BSF dry weight by 25%	4.44%
+50%	Extrapolation - FreezeM Genetic Line increased BSF larval *mass* by 50% (Varol 2023)	8.21%
+100%	Extrapolation	14.4%

SUPPLEMENTARY INFO

Rearing Methods

Rearing Methods - Eggs were harvested from a BSF colony established at the Forensic Laboratory

for Investigative Entomological Sciences (FLIES) at Texas A&M University. Larvae from this colony are

descended from the 'Sheppard Strain' originally maintained laboratory at the Coastal Plain Experiment

Station of the University of Georgia, Tifton, GA (Sheppard et al. 2002). Larvae were reared on 70%

moisture Gainesville (Producers Cooperative Association, Bryan, TX, USA) diet (Hogsette 1992)

according to the FLIES colony SOP (see: Supplementary Info) until they reached 8-d-old. After this, larvae

were split into different density treatments (n = 3 levels, with 2 duplicate pans for redundancy) of

approximately 5,000, 7,500, and 10,000 larvae (gravimetrically estimated). Aliquots of larvae were placed

directly atop 7.5 kg Gainesville feed at 70% moisture in 27 L pans (Sterilite Corporation, Townsend, MA,

USA). An additional 0.75 kg of dry Gainesville was placed around the edge of the pans to prevent escape.

For the remainder of the rearing-cycle, pans were housed in a walk-in incubator maintained at

approximately 25°C, 50% relative humidity, and 14:10 L:D photoperiod (fluorescent lighting).

Contents of each pan were hand-mixed on days 11 and 12 to vent heat, prevent anoxic composting,

and give larvae better access to the undigested portions of the substrate. An additional kilogram of 70%

moisture Gainesville diet was supplemented to pans on days 15, 16, and 18. On day 26, when at least 40%

of the larvae began pupation (Bosch et al. 2020), the two pans of larvae per density-level were combined

prior to being sieved. Then, 150 g aliquots from each respective density-level were transferred to cylindrical

plastic containers (13.5x13.2 cm: d × h) to pupate (Dickerson et al. 2024 May 22, Lemke et al. 2024). 150

g dry Gainesville was used as bedding material, and the tops of the containers were secured with Tulle

fabric (Wal-Mart Stores Inc., Little Rock, AR, USA).

Oviposition traps of fluted cardboard were collected from the colony and transferred into a mason jar covered by two layers of paper towel to allow airflow but prevent escape. During this time, the eggs hatched, and neonates migrated into the bottom of the mason jar. Then, 0.5 g of neonates were transferred into each of three square plastic containers. The neonates were fed with 500 g Gainesville feed (Hogsette 1992) formulated at 70% moisture. A border with 50 g of dry feed was used to prevent larvae from escaping. The black soldier fly larvae were reared in a walk-in incubator at approximately 27 °C.

Supplemental Figure 1. Genitalia secured between microslides in wells made of two layers of Duck Brand masking tape.

Supplemental Figure 2. Principal component analysis (PCA) performed with the variables of thorax, head, sheath, gonostylus, and larval density

507 Supplemental Figure 3. Scree plot demonstrating that principal components and two explain a sufficient

508 amount of the variability (>75%).

509
510

511 Supplemental Table 1. Principal component analysis loadings

	PC1	PC2
Thorax	0.58361004	0.1027914
Head	0.58406062	0.0847625
Sheath	0.26451689	-0.6319838
Gonostylus	0.01790199	-0.7499736

| Density | -0.49797876 | -0.1427769 |