Human Beta Oscillations Reflect Magnitude and Fidelity of Priority Shifts in Working Memory

Nicholas E. Myers[1,2], Mark G. Stokes[3], Paul S. Muhle-Karbe[4,5]
1. School of Psychology, University of Nottingham
2. Centre for Neurotechnology, Neuromodulation and Neurotherapeutics, University of Nottingham
3. Department of Experimental Psychology, University of Oxford
4. School of Psychology, University of Birmingham
5. Centre for Human Brain Health, University of Birmingham

Corresponding author: nicholas.myers@nottingham.ac.uk

Acknowledgments

This work was supported by funding from the Wellcome Trust (Award 201409/Z/16/Z/ to N.E.M. and Award 210849/Z/18/Z to P.S.M.K), University College, Oxford (N.E.M.), and Linacre College, Oxford (P.S.M.K.), the Biotechnology and Biological Sciences Research Council (Award BB/M010732/1 to M.G.S.), the James S. McDonnell Foundation (Award 220020405 to M.G.S.), and the National Institute for Health Research Oxford Health Biomedical Research Centre. The Wellcome Centre for Integrative Neuroimaging is supported by core funding from the Wellcome Trust (Grant 203139/Z/16/Z). This article is dedicated to Mark G. Stokes, who passed away in 2023.

Abstract

Flexible prioritisation in working memory (WM) is supported by neural oscillations in frontal and sensory brain areas, but the roles of different oscillations remain poorly understood. Recordings in humans suggest an interplay between prefrontal slow frequency (2-8Hz) and posterior alpha-band (10Hz) oscillations regulating top-down control and retrieval of WM representations, respectively. Complementary work, primarily in non-human primates, suggests an additional role for beta (15-30Hz) oscillations in clearing or inhibiting stimuli from entering WM. Here we investigated the role of neural oscillations in prioritising WM content using electroencephalography (EEG) as participants (humans of any sex) performed a task requiring frequent priority switches between two memorized oriented bars. Behavioural performance revealed switch costs, which scaled with the angular distance between the two items, suggesting that priority shifts are modulated by shift magnitude. Time-frequency analyses revealed increased frontal theta (4–8Hz) and decreased central-parietal beta (15–25Hz) power during switches. Crucially, only beta power scaled with the magnitude of the priority shift and predicted the fidelity of neural decoding of the newly prioritized item during subsequent recall. Theta power, by contrast, was elevated on switch trials but did not vary with update magnitude or decoding strength, suggesting a more general role in signaling control demands. Our findings highlight a particular and previously overlooked role for beta-band oscillations in the flexible prioritisation of WM content.

Significance Statement

Working memory permits flexible switching between mental representations, so we can focus on what is most relevant at the moment. Different brain rhythms in frontal control and sensory memory storage areas orchestrate switches but their respective roles remain unclear. Here, using EEG, we show that power reductions of ~20Hz oscillations over central-parietal regions, usually associated with the motor system, closely track the magnitude of the required switch and the fidelity of the prioritized memory. In contrast, slower 4-8Hz (theta-band) activity over frontal regions increases during priority switches but tracks neither magnitude nor fidelity. Our findings suggest a unique function for central-parietal beta oscillations in the flexible control of working memory.

Keywords: working memory, attention, task switching, beta oscillations, theta oscillations, multivariate pattern analysis, EEG

Introduction

Working memory (WM) supports the maintenance of task-relevant information, but its capacity is severely limited. Focusing attention on individual items in WM permits effective use of this limited capacity, allowing us to remember more quickly and accurately (Griffin & Nobre, 2003; Souza & Oberauer, 2016; Van Ede & Nobre, 2023). Multiple-state models (Cowan, 2000; Oberauer, 2002) propose that high-priority items are stored in a privileged state that guides perception and action, while other items are concurrently held in a de-prioritised accessory state without driving behaviour (Olivers et al., 2011).

Neurally, prioritised and de-prioritised WM items may be represented in different brain areas (Christophel et al., 2018) and representational formats (El-Gaby et al., 2024; Iamshchinina et al., 2021; Paluch et al., 2025; Panichello & Buschman, 2021; van Loon et al., 2018; Yu et al., 2020). Remarkably, items can be flexibly moved in and out of the focus of attention (Muhle-Karbe et al., 2021; Rerko et al., 2014; Rerko & Oberauer, 2013). Yet the neural mechanisms that underpin priority switches remain to be delineated. This is the focus of the current study.

Prefrontal neural synchronisation for controlled access to items stored in sensory brain areas (Wallis et al., 2015) may represent such a mechanism. More specifically, low-frequency oscillations in the delta and theta frequency ranges (2-8Hz) originating in frontal cortex may play a role in orchestrating top-down control over WM representations stored in sensory or medial temporal areas (Berger et al., 2019; Johnson et al., 2018; Liebe et al., 2012; Roux & Uhlhaas, 2014). Retrieval of these sensory WM representations may occur via spatiotopic desynchronisation of alpha-band oscillations (de Vries et al., 2017, 2018; Myers, Walther, et al., 2015; Poch et al., 2014; Riddle et al., 2020, 2024; van Ede et al., 2016).

Parallel work has emphasised the role of beta oscillations (15-30Hz) in WM (Liljefors et al., 2024; Lundqvist et al., 2016; Miller et al., 2018; Spitzer & Haegens, 2017) but their role in prioritisation is less clear, as priority switches likely require multiple sub-processes that could be linked to beta: a shift in beta synchrony could signal a change in task context (Engel & Fries, 2010) or facilitate removal of the previously prioritised memory (Lundqvist et al., 2018). Given the role of beta in coordinating cortical-basal ganglia loops (Brittain & Brown, 2014; Jenkinson & Brown, 2011), which have been implicated in gating information in and out of WM (Chatham et al., 2014; Chatham & Badre, 2015; D'Ardenne et al., 2012), they may alternatively play a role in selecting a new memory into the prioritised state.

Prioritisation in WM is typically investigated with the retro-cue paradigm, where participants initially encode multiple memory items until a subsequent cue signals which item will be task-relevant for a forthcoming decision (Griffin & Nobre, 2003). This procedure typically incurs only a single priority shift and therefore confounds changes in priority states with changes in memory load,

as the initially prioritised item can be dropped after a switch. We recently
developed a paradigm that avoids this issue by requiring frequent priority
switches between the same two items, so that neither can be dropped from
memory. Using this paradigm, we have previously shown that both prioritised
and de-prioritised memory items can be decoded from
electroencephalography (EEG) recordings and that they track different
aspects of memory-guided behaviour (Muhle-Karbe et al., 2021).
121
Here, we sought to leverage this dataset to characterise the role of theta and
beta oscillations during priority switches. We related both oscillations to the
magnitude of priority switches and their fidelity by harnessing single-trial
decoding of prioritised and de-prioritised memory items. Priority switches
increased frontal theta and decreased central-parietal alpha and beta power.
We found that beta (but not theta or alpha) oscillations correlated with the
magnitude of priority switches and the decoding strength of newly prioritised
memory items, but not the previously prioritised item, suggesting that beta
oscillations are primarily involved in prioritising new items.
131

Materials and Methods

Participants

We reanalysed data from a previously published study (Muhle-Karbe et al., 2021). 43 adults participated in total. We combined data from two closely related variants of the same visual WM task (experiments 1 and 2). Our previous study showed identical results across both variants. We therefore collapsed all analyses across both experiments. Twenty adults participated in experiment 1 (mean age = 28.1, age range = 18-37, 10 female, 10 male, 1 left-handed, 19 right-handed) and 30 participated in experiment 2 (mean age = 26.8, age range = 19-41, 14 female, 16 male, 2 left-handed, 28 right-handed). Seven participants took part in both experiments, yielding a final sample of 43 participants. For participants who completed both experiments, we analysed data separately within each experiment before averaging the results across the two experiments. All participants reported normal or corrected-to-normal vision and were compensated for their participation. The study was approved by the Central University Research Ethics Committee of the University of Oxford.

Apparatus

Stimuli were presented using the Psychophysics Toolbox (Brainard, 1997) on a 22" monitor with a refresh rate of 100Hz, viewed at a distance of approximately 60cm in an electrically shielded, sound-attenuated booth with dim background lighting. Stimuli appeared on a grey background ([127 127 127]). Participants responded with left and right index fingers on the "B" and "Y" buttons of a QWERTY keyboard (placement counterbalanced across participants).

EEG Acquisition

EEG data were collected with a NeuroScan SynAmps RT Amplifier and Curry 7 software. 61 EEG channels were placed following the extended 10-20 positioning system. Channel impedances were kept below 5 kOhm. In both studies, eye movements were recorded via electrooculography (EOG) using electrodes placed above and below the left eye and to the outside of each eye. We also recorded activity in the first dorsal interosseus muscle of the left and right hand via electromyography (EMG). Additional electrodes were placed on the left and right mastoids, which were averaged and used for offline re-referencing. The ground electrode was placed on the left elbow. Data were recorded at 1000Hz.

Eye-tracking

In some participants (experiment 2, 30 participants) we additionally recorded eye movements using a remote infrared eye-tracker (SR Research, Eyelink 1000) sampling both eyes at 1000Hz. These data were used to confirm that participants maintained close fixation and that EEG-based decoding of memory contents could not be explained by systematic deviations in fixation location (see Muhle-Karbe et al., 2021, for control analyses). Eye-tracking data were not further analysed here.

Task Design

The experiment used a 2-item visual working memory task requiring frequent
priority switches (Figure 1a). Task blocks consisted of 16 trials. At the
beginning of each block, participants encoded in WM two orientated bars (6°
visual angle in length and 0.25° in width, presented at the same location as
subsequent stimuli) in blue ([25.5 25.5 204]) and yellow ([204 204 25.5]). These
bars served as memory items for the remainder of the block. The two
orientations were drawn randomly from a set of 16 possible orientations
(spaced evenly at 11.25° intervals from 2.8125° to 171.5625°) with the
constraint that the two items could not be identical or exactly orthogonal. As
a result, the absolute angular distance ('item distance') between them ranged
from 11.25° to 78.75°. One bar colour was associated with a high-pitch tone
and the other with a low-pitch tone (mapping counterbalanced across
participants), with the two tones serving as retrieval cues for the respective
bars later in the block. Participants had unlimited time to encode the two bars
and initiated the block via button press.

Within the block, each trial started with a presentation of an auditory cue (pure
sinusoidal tones, low tone: 440Hz, high tone: 880Hz, duration: 100ms
including a 10ms ramp-up and 10ms ramp-down). The tone signalled which
memory item should be used for a forthcoming perceptual decision (cued
item), while the other item was maintained for later use in the block (uncued
item). The cue was followed by a 700ms delay period within which a black
fixation dot was presented centrally on the screen (0.15° diameter). A randomly
oriented Gabor patch was presented (orientation drawn from 16 possible
orientations, spaced evenly at 11.25° intervals from 8.4375° to 177.1875°;
patches had 6° diameter, 50% contrast, 1.75 cycles/°, random phase, and a
Gaussian envelope with 1.2° SD). Participants were given a maximum of
4,000ms to judge the probe via button press as clockwise or counter-
clockwise relative to the cued memory orientation. Probes were presented for
100ms and replaced by a fixation dot for the remainder of the response period.
In experiment 1, the probe was presented centrally on the screen on all trials.
In experiment 2, the probe was presented laterally at a distance of 6° from the
screen centre. The side of probe presentation (left vs. right) alternated
predictably across blocks. A noise patch (Gaussian smoothed random white
noise using a kernel with 0.13° SD, convolved with a Gaussian envelope with
1.2° SD) that matched the probe stimulus in luminance, size, contrast, and
eccentricity, was presented on the side of the screen at which no probe
appeared. Clockwise and counter-clockwise decisions were indicated with the
right and left index fingers respectively. The response period was followed by
a variable inter-trial interval (400-900ms, drawn from a truncated exponential
distribution with mean 550ms).

The next trial cued either the other item (Switch trial) or the same item again
(Repeat trial). The likelihood of a cue repeat on each trial was drawn from a
modified exponential distribution with a minimum of 1 repeat and a maximum
of 7 repeats after each Switch trial. This resulted in comparable trial numbers
for Switch trials (464±9, mean±s.e.m.), the first Repeat trial after a switch
(Repeat 1, 553±9), and subsequent Repeat trials (Repeats 2-7, abbreviated

Repeat 2+, 646±10). Participants received feedback only at the end of each
block when their mean accuracy and response time were presented.
Accordingly, they had to maintain precise representations of both memory
items for the whole duration of the block, as they could not rely on trial-wise
feedback to infer the orientation of an item if it was forgotten. Overall,
participants completed 128 blocks, resulting in a total of 2048 trials and lasting
approximately 2h.
236
Behavioural Analysis
We were primarily interested in the effect of priority switches, i.e., trials in
which the item cued as relevant by the auditory cue was not the same as the
item cued on the preceding trial (Switch trials). We compared Switch trials
with Repeat trials, i.e., trials in which the same item was cued as on the
preceding trial. We excluded the first trial in each block because it was
neither a switch nor a repeat trial. We then calculated mean accuracy and
reaction time on switch trials, the first repeat trial (Repeat 1), and any
subsequent repeat trials (Repeat 2+). Reaction times were calculated as the
median of all correct responses to a given condition. Since accuracy on
Repeat 1 and Repeat 2+ trials was largely the same (Fig. 1a), for the
remainder of the manuscript we focused on comparing Switch with Repeat 1
trials. Including Repeat 2+ trials in any analyses did not materially affect our
results.
251
252   We next fit a psychometric curve to the behavioural data to separately
253   estimate the memory precision (inverse of the slope, σ) and the lapse rate (λ)
254   using the response function:
255
256    $$P(\text{response} = \text{clockwise} \mid \theta) = (1-\lambda)*\Phi(\theta,\mu,\sigma) + \lambda$$
257
where θ is the angular difference between the probe and cued memory
stimulus, λ is the lapse rate, μ is the response bias (fixed to 0), σ is the
inverse of the memory precision, and Φ is the cumulative density function of
the normal distribution.
Precision ($1/\sigma$) and lapse rate (λ) were fit using maximum likelihood
estimation separately for Switch and Repeat trials. Estimated parameters
were compared between conditions using pairwise t-tests.
265
266   We extended the psychometric model to include erroneous comparisons to
267   the uncued item ('swap errors'):
268
269    $$P(\text{response} = \text{clockwise} \mid \theta_{cued}, \theta_{uncued}) = (1-p_{swap}-\lambda)*\Phi(\theta_{cued},\mu,\sigma) +$$
270    $$p_{swap}*\Phi(\theta_{uncued},\mu,\sigma) + \lambda;$$
271
272   where θ_{cued} is the angular difference of the probe orientation relative to the
273   cued WM item, θ_{uncued} is the angular difference of the probe orientation
274   relative to the uncued WM item, and p_{swap} is the swap rate.
275

276 To examine the effect of item distance, we fit a linear model of item distance
277 (with values ranging from 11.25º-78.75º) to accuracy and reaction time data
278 on Switch and Repeat trials separately. Regression coefficients were then
279 compared against zero and each other using one-sample or paired-samples
280 t-tests.
281

EEG Preprocessing
EEG data were re-referenced to the average of both mastoids. EEG, EOG, and
EMG data were down-sampled to 250Hz and bandpass-filtered between
0.1Hz and 45Hz. EEG channels with excessive noise were identified through
visual inspection and replaced via interpolation using a weighted average of
the surrounding electrodes. The continuous data were epoched from 750ms
before to 2500ms after the onset of the cue on each trial. Each trial was
inspected visually for blinks, eye-movements, and non-stereotyped artefacts.
Trials were rejected if they contained any of those artefacts during the cue-
probe delay or probe period. Stereotyped artefacts (from blinks, saccades, or
muscle artefacts) outside those periods were subsequently removed via
independent component analysis. Unless stated otherwise, the data were
baseline-corrected for the decoding analyses using the average signal from
the time window of 200 to 50 ms before cue onset.
296

EEG Decoding
We conducted multivariate pattern analysis to characterise the neural
representations of the cued and uncued WM item. We followed a previously
established approach (Muhle-Karbe et al., 2021; Wolff, Jochim, et al., 2020;
Wolff, Kandemir, et al., 2020) that used the signal at multiple posterior EEG
sensors, pooled across multiple time points, to decode remembered
orientations. This approach exploits the dynamic temporal structure of event-
related potentials by pooling multivariate information in time, capturing
information encoded in spatial activation patterns and in the temporal
unfolding of these patterns to achieve greater decoding sensitivity (Wolff,
Jochim, et al., 2020).
308

In keeping with this previous work, decoding analyses were conducted only
within posterior EEG sensors (P7, P5, P3, P1, P2, P4, P6, P8, PO7, PO5, PO3,
POz, PO4, PO8, O1, Oz, O2). We selected these channels to be consistent
with previous studies and because orientation signals are typically expressed
most strongly in posterior regions (Cichy et al., 2015; Myers, Rohenkohl, et al.,
2015; Wolff, Jochim, et al., 2020). We pooled the signal across a 450ms
backward-looking sliding window (i.e., decoding at 800ms after cue onset
used data from 350ms-800ms), downsampled to 83.3Hz (i.e., one timepoint
every 12ms), yielding 37 timepoints from each of the 17 EEG sensors, or 629
features in total. Data were demeaned across timepoints within each sensor
(to reduce any potential carryover of decodable signal from other time
windows, although this step seemed to have a minimal effect in our data, see
Fig. S5 top vs. bottom row). Our previous work showed that WM decoding
peaked around the time of the decision, approximately 200-400ms after probe
onset. We therefore focused on the post-probe window for analysis, repeating

the decoding analysis in a sliding window from 800-1,400ms relative to cue
onset (or 0-600ms relative to probe onset), in 50ms steps.
326
We computed Mahalanobis distances between the patterns of sensor activity
that were evoked by different stimulus orientations and measured the extent
to which these distances reflected the underlying circular orientation space.
We used a 13-fold cross-validation procedure. On each fold, data from 10 task
blocks served as test data and all the remaining data served as training data.
From the training data we calculated the average response to each of the 16
orientations, yielding a $N_{orientations}$-by-$N_{features}$ (16x629) training data matrix M_{train}.
Mahalanobis distances between the 16 average patterns in the training data
and each test trial i (M_i, 1x629) were then computed, using the noise
covariance from the training data. Noise covariance was calculated on the
residual data after subtracting orientation-specific mean activity from each
trial, using a shrinkage estimator (Ledoit & Wolf, 2004). Smaller values
indicated a smaller Mahalanobis distance, i.e., greater similarity between the
test trial and a given orientation from the training set. For each test trial, this
yielded a vector D_i of 16 distances. D_i was circularly shifted to center the vector
on the test trial orientation θ_i and inverted so larger values indicated greater
pattern similarity, yielding a 'tuning curve' that peaks at the centre of the curve
when the orientation is decodable. To calculate a trial-wise index of decoding
we computed a summary measure that estimated how closely the tuning curve
resembled a cosine peaking at the test trial orientation. This index was
calculated as the cosine vector mean of the 16-dimensional tuning curve by
multiplying each distance with the cosine of the orientation difference to the
test trial (Sprague et al., 2016; Wolff et al., 2017):

$$decoding_i = \sum_{j=1}^{16} cos(2\theta_j)D_{i,j}$$

where θ_j is the orientation offset of the jth point in the tuning curve (ranging
from -78.75° to +90°) and $D_{i,j}$ is the Mahalanobis distance between test trial i
and the training set average for the orientation at θ_j relative to θ_i. Angles were
doubled in the equation to account for the 180°-symmetry of the orientation
stimuli. Because the cosine is zero-centred, if the tuning curve is flat this
operation yields a mean decoding score of 0. If decoding peaks at the centre
of the curve (at θ_j=0°), this yields a positive decoding score. An inverted curve
(peak near 90°) would yield a negative decoding score. The absolute value of
the decoding score is difficult to interpret because it depends on the voltage
change over time, the number of sensors, and the number of timepoints. The
analysis was run separately for the cued and the uncued orientation. In
experiment 2, decoding analyses were conducted separately for blocks with
probe presentation on the left and right side, with training and test data
matched by side. Training data consisted of all available trials, with decoding
accuracy of the test data split afterwards into Switch and Repeat trials.
366
EEG Time-Frequency Analysis
Preprocessed EEG data were transformed with a Spatial Laplacian filter to
reduce spatial spread of neural sources across the scalp. We used the

FieldTrip function *ft_scalpcurrentdensity* (using default conductivity (0.33) and
lambda (10^{-5}) values and a degree of 10 as in (de Vries et al., 2018)). We then
calculated the time-frequency transform in sliding windows from -0.75s
before cue onset to 2.5s after cue onset (in 40ms steps). At each timepoint,
we calculated power at frequencies between 2 and 40Hz (in 1-Hz steps)
using Hanning tapers (using a time window corresponding to 5 cycles of the
frequency of interest centered on the current timepoint). We then log-
transformed power (10*log10) and baseline-corrected using power from 0.75-
0.25s before cue onset.
379
Statistical Testing
Unless otherwise noted, we used multiple linear regression to establish
relationships between oscillatory power and other task variables (item
distance, WM decoding), separately for Switch and Repeat trials. Regressors
and dependent variables were z-scored prior to regression. We tested for
significance using one-sample t-tests against 0 (at each time point, frequency,
or time-frequency combination) and corrected for multiple comparisons via
cluster-based permutation testing using 10,000 permutations (Maris &
Oostenveld, 2007).
389

390 **Results**
391
WM Priority Switches Incur a Temporary Behavioural Cost
43 Participants performed a visual WM task designed to induce frequent
priority switches while we recorded their brain activity through
electroencephalographic (EEG) recordings (see Methods for details). The task
consisted of 128 mini-blocks each of which contained 16 trials. Participants
first encoded two randomly oriented bars into WM and maintained both for
the duration of the mini-block. Each bar was associated with a different
auditory cue (see Fig. 1a). Every trial began with an auditory cue indicating
which of the two items was relevant (Fig. 1b). Participants therefore
prioritised the cued stimulus and deprioritised the other (uncued) stimulus.
After a 700ms cue-probe interval a randomly oriented probe stimulus
appeared and was judged as clockwise or counterclockwise relative to the
cued WM orientation. The following trial could then either repeat the same
cue (Fig. 1c, Repeat trials, max. 7 repeats in a row) or switch to the other
auditory cue (Switch trials), requiring prioritisation of the previously uncued
item. Multiple switches occurred in each block, so participants had to keep
both stimuli in WM.
409
Accuracy on Switch trials (78.3%±0.9% s.e.m., Fig. 1d) was modestly but
significantly lower than on the first Repeat trial (80.4±0.7%, t_{42}=-4.41,
p=7.01*10^{-5}, Cohen's d=-0.67) and lower than on subsequent repeat trials
(80.0±0.7%, t_{42}=-4.06, p=2.08*10^{-4}, d=-0.62). The first Repeat trial did not
differ from subsequent repeats (t_{42}=1.11, p=0.275, d=0.17). Switch costs
were driven by an increase in guess rate on Switch trials (9.2±1.1%, Fig. 1f)
compared to all Repeat trials (7.0±0.7%, t_{42}=2.41, p=0.021, d=0.37). By
contrast, precision remained unchanged (Switch: 18.2±1.2°, Repeat:
17.6±0.7°, t_{42}=0.43, p=0.67, d=0.07).
419
Similarly, median reaction time on correct trials was significantly slower on
Switch trials (693±23ms, Fig. 1e) compared to either the first Repeat trial
(616±18ms, t_{42}=9.19, p=1.30*10^{-11}, d=1.40) or subsequent Repeat trials
(604±16ms, t_{42}=9.91, p=1.47*10^{-12}, d=1.51). Reaction time on the first Repeat
trial was also significantly slower than on subsequent Repeats (t_{42}=3.71,
p=6.08*10^{-4}, d=0.57).
426
Priority Switch Costs Scale with Update Magnitude
We next explored whether the similarity between the two items held in WM
affected behaviour (Fig. 1g-h). As the angular distance between the cued and
uncued item ('item distance', ranging from 11-79° in 11.25° steps) increased,
accuracy decreased on both Switch trials (decrease in accuracy of
0.97±0.17% per 10 degrees, t-test on slope: t_{42}=-5.71, p=1.04*10^{-6}, d=-0.87)
and Repeat trials (0.46±0.11% per 10 degrees, t_{42}=-4.16, p=1.54*10^{-4}, d=-
0.63). However, the effect of item distance was stronger on Switch trials (t_{42}=-
3.00, p=0.0045, d=-0.46). After regressing out item distance, there was no
longer a significant switch cost (t_{42}=0.73, p=0.47, d=0.11), indicating that
switch costs were driven primarily by trials requiring a large update from the

previous to the currently cued item. Consistent with this, blocks with similar
WM item orientations (item distance <25°) showed no switch cost on
accuracy ($t_{42} \leq 0.29$, $p \geq 0.77$, $d \leq 0.04$).
441
We found complementary effects on reaction time (Fig. 1e), which increased
significantly with item distance on Switch trials (22.5±2.6ms per 10 degrees,
$t_{42}=8.64$, $p=7.24*10^{-11}$, $d=1.32$) and, to a lesser degree, on Repeat trials
(5.9±0.9ms per 10 degrees, $t_{42}=6.60$, $p=5.4*10^{-8}$, $d=1.01$). Again, the effect of
item distance was significantly larger on Switch trials ($t_{42}=7.69$, $p=1.5*10^{-9}$,
$d=1.17$). However, even after regressing out the effect a significant difference
between Switch and Repeat trials remained ($t_{42}=2.31$, $p=0.026$, $d=0.35$).
Therefore, reaction time switch costs were present even in blocks with very
similar WM items (item distance 11°: $t_{42}=-3.89$, $p=3.53*10^{-4}$, $d=0.59$).
451
The item-distance effect could have been driven in part by occasional failures
to update the cued item on Switch trials, leading to responses relative to the
uncued item (sometimes referred to as a swap error, Bays et al., 2009). When
we estimated the swap rate separately from the lapse rate we indeed found
that both increased on Switch trials (Suppl. Fig. S1d-e). Nevertheless, when
we accounted for potential swap error effects, there was still an independent
significant effect of item distance on both accuracy and reaction time (Suppl.
Fig. S1f-g).

Figure 1. *Shifting priority between WM contents incurs a temporary switch*
cost. **A.** *Each block began with encoding of two oriented bars. Each bar was*
associated with an auditory cue. **B.** *On each trial, a cue indicating which WM*
item was relevant was followed by a probe. Participants judged the probe as

clockwise or counterclockwise relative to the cued memory orientation. **C.**
Successive trials could cue either the same or different items, requiring
frequent priority switches and maintenance of both items throughout the block.
D. *Accuracy was lower on Switch trials than the first Repeat trial after a switch*
or subsequent Repeat trials (max. 7 repetitions). Coloured horizontal lines show
median, black horizontal lines show mean. Dots are individual observers. **E.**
Reaction time was slowest on Switch trials and slightly slower on the first
Repeat trial. Empty black circles (both Repeat conditions) represent one outlier.
F. *Psychometric curves fit to accuracy had comparable slopes but higher*
guess rates on Switch trials. Dots are data, with error bars indicating standard
error of the mean (s.e.m.). Line is the mean psychometric curve fit (shaded area
around mean shows s.e.m.). Horizontal dashed lines are the mean asymptotic
performance ('guess rate') for each condition (blue: Switch, grey: Repeat). **G.**
The similarity between WM orientations ('Item Distance') affected accuracy,
especially on Switch trials, and accounted for the switch cost. Lines are mean
of linear fits to each observer and condition (shaded area shows s.e.m.). **H.**
Item distance increased reaction time more on Switch than Repeat trials. ***:
p<0.001, **: p<0.01, *: p<0.05, n.s.: not significant.
483

Priority Switches Lead to Temporary Decreases in Beta-band Power and Sustained Increases in Theta-band Power

We next examined the neural oscillatory correlates of priority switches in WM. We focused on oscillations because previous work has implicated frontal slow oscillations in the delta/theta-band and posterior alpha-band power reductions (de Vries et al., 2018; Poch et al., 2014; Riddle et al., 2020; Wallis et al., 2015) in switching attention between items held in WM. Here, we observed increased frontal theta power and decreased posterior alpha power, but additionally found a pronounced reduction of power in the beta-band (15-25Hz).

Beta-band power was significantly lower on Switch trials compared to Repeat trials (Fig. 2a). The effect was strongest in a frequency-limited band (approximately 15-25Hz) and peaked in a brief period shortly after the onset of the cue (roughly 400-800ms after cue onset) but before the onset of the memory probe (at 800ms after cue onset). The effect (i.e., decreased power) was most pronounced at central-parietal sites (Fig. 2b) and all subsequent analyses focused on these channels ('Central Channels': C1,Cz,C2,CP1,CPz,CP2,P1,Pz,P2). The power decrease on Switch trials at central channels extended to the Alpha band (8-14Hz) and to higher frequencies (25-40Hz, sig. cluster 7-40Hz, $p<10^{-4}$ after cluster-based permutation test) but showed a pronounced peak in the beta-band (Fig. 2c) indicating that beta oscillations are a distinct phenomenon. The effect was significant throughout the trial (from 320ms after cue onset) but strongest in the latter half of the cue period (Fig. 2d).

In parallel, we observed a significant increase in theta (4-8Hz) power on Switch trials (Fig. 2e) that peaked at frontal midline channels (Fig. 2f, 'Frontal Channels': AF3,AFz,AF4,F1,Fz,F2). The effect was limited to the theta and alpha bands (2-15Hz, corrected p=0.001, Fig. 2g) but accompanied by a reduction in higher-frequency power (24-36Hz, p=0.013, Fig. 2g). The increase in theta-band power persisted throughout the Switch trial (Fig. 2h). Finally, power in the alpha-band (8-14Hz) decreased at posterior channels (O1,Oz,O2,PO7,PO3,POz,PO4,PO8, see Supplementary Fig. S2a). We observed the same effects when comparing Switch trials to later Repeat trials (Repeat2+, Fig. S3).

521
Figure 2. *Priority switches led to a temporary decrease in beta-band (15-25Hz) and an increase in theta-band (4-8Hz) power. **A.** Average difference in power between Switch and Repeat trials at central channels. Between the cue and probe onset, power reduction on Switch trials was strongest in the 15-25Hz range but extended from ~7 to 40Hz (shading indicates significant cluster from permutation test). Time-frequency points are plotted with saturated colours if they belong to a significant cluster (permutation test, p<0.05). **B.** Beta power reduction on Switch trials was strongest at central-parietal channels. **C.** In the late cue period (400-800ms after cue onset), power at central channels peaked in the alpha and beta bands and was lower on Switch trials from 7-40Hz. Shaded area around mean indicates s.e.m., black bar denotes significant difference between Switch and Repeat trials after cluster-based permutation testing. **D.** Central 15-25Hz power was lower throughout Switch trials but strongest in the late cue period. **E-H.** 4-8Hz power at frontal channels was stronger on Switch trials.*
537

Beta (but not Theta) Power Scales With Magnitude of Priority Switches

Given the pronounced effect of item distance on behaviour, particularly on Switch trials, we tested if beta- and theta-band effects were also sensitive to item distance. Indeed, beta-band power scaled with item distance (Fig. 3a, $F_{6,252}=2.303$, p=0.035, partial $\eta^2=0.009$), with opposing effects on Switch and Repeat trials (interaction Switch effect with Item Distance effect $F_{6,252}=8.712$, p=$1.25*10^{-8}$, partial $\eta^2=0.033$). On Switch trials, larger priority switches (i.e., larger item distances) led to lower beta-band power (t-test on linear fit, $t_{42}=-5.74$, p=$9.54*10^{-7}$, d=-0.88). On Repeat trials, the effect was reversed, with a (smaller) increase in beta power for larger item distances ($t_{42}=3.12$, p=0.0033, d=0.48), leading to a significant difference in slope between Switch and Repeat trials ($t_{42}=-5.86$, p=$6.56*10^{-7}$, d=-0.89). Even after regressing out the effect of item distance, beta power was still lower on Switch trials ($t_{42}=-3.42$, p=0.0014, d=-0.52). The effect of item distance was confined to the beta band (Fig. 3b, with band-limited sig clusters on Switch trials 15-33Hz, p<10^{-4}, on Repeat trials 17-27Hz, p=0.0062, Switch vs. Repeat 15-33Hz, p<10^{-4}, Fig. 3e) and to central-parietal channels (Fig. 3c), emerging approximately 200ms after cue onset (Fig. 3d, sig. effect on Switch trials 162-782ms, corrected p<10^{-4}, effect on Repeat trials 482-782ms, p=0.0048, difference Switch vs. Repeat 242-782ms, p<10^{-4}). Therefore, the beta-band power decrease observed on Switch trials scales with the magnitude of the update between the previous and currently cued item.

By contrast, frontal theta power was largely insensitive to item distance (Fig. 3f, main effect of item distance $F_{6,252}=0.343$, p=0.913, partial $\eta^2=0.001$, interaction Switch by item distance $F_{6,252}=0.867$, p=0.519, partial $\eta^2=0.0034$). Linear fits showed no effect of item distance in either Switch ($t_{42}=1.21$, p=0.233, d=0.18) or Repeat trials ($t_{42}=-0.57$, p=0.574, d=-0.09). Theta power remained higher on Switch trials after regressing out item distance ($t_{42}=3.24$, p=0.0024, d=0.49). We found no significant effect at frontal channels when analysed separately by timepoint and frequency (Fig. 3g) or averaged over the late cue period (Fig. 3j). We also saw no clear theta effects at other channels (Fig. 3h). When we examined the effect separately for each time point in the trial, we did find a small effect of item distance on Switch trials immediately following the cue onset (Fig. 3i, effect on Switch trials from 122-422ms, p=0.03, difference between Switch vs. Repeat effects 42-382ms, p=0.018). The early and brief effect suggests that this may reflect a phasic event-related response to the cue rather than the sustained theta oscillation increase observed on Switch trials more generally (Fig. 2e). We found no effect of posterior alpha power on item distance on Switch trials, with only a small effect on Repeat trials (Fig. S2b) that could have been driven by the spatially and spectrally adjacent beta-band effect. In sum, unlike the graded beta-band effect increasing with item distance, oscillatory theta power increases on Switch trials appear to be largely an all-or-none phenomenon.

Figure 3. *Central Beta Power Scales with Item Distance.* **A.** *Beta (15-25Hz) power at central channels (400-800ms post-cue), separated by Switch (blue) vs. Repeat (grey) trials and by angular distance between WM items. A large item distance required a larger-magnitude update of the prioritised WM representation on Switch trials. Dots show average power (error bars show s.e.m.), line shows linear fit (shaded area is s.e.m.). Inset shows regression coefficients for each participant.* **B.** *The effect of item distance on power (regression coefficient on Switches vs. coefficient on Repeats) was resolved over time and frequency and was significant primarily between 15 and 30Hz (area with saturated colour indicates significant cluster from permutation test, p<0.05 corrected).* **C.** *The effect peaked at central-parietal channels (compare Fig. 2b).* **D.** *Regression coefficient resolved by time (average of 15-25Hz power at central channels), separately for Switch and Repeat trials. Bars indicate significant effects after correction (blue: Switch vs. 0, grey: Repeat vs. 0, black; Switch vs. Repeat).* **E.** *Effect resolved by frequency (Central channels, 400-800ms after cue onset).* **F-J.** Frontal 4-8Hz power showed no comparable effects, except for a brief correlation on Switch trials immediately after cue onset (panel I).

Beta Power After Priority Switches Predicts Decoding of Cued WM Item
Since the magnitude of beta power reduction after priority switches scaled
with the magnitude of the update, we reasoned that it might be involved in
retrieving or shifting attention to the newly cued WM item. We therefore
expected that beta power should predict the strength of WM decoding. To
this end, we calculated a trial-by-trial orientation decoding score around the
time of decision-making after the onset of the probe (200-400ms following
probe onset, i.e., 1,000-1,200ms after cue onset). Decoding in this window
was maximal (Fig. 4a inset, see also Muhle-Karbe et al., 2021) and
comparable on Switch and Repeat trials (Switch trials, significant decoding
800-1,400ms after cue onset, e.g., 0-600ms after probe onset, corrected
$p<10^{-4}$, Repeat trials: significant decoding 900-1,400ms, $p<10^{-4}$, no sig.
difference between Switch and Repeat, smallest corrected p=0.088).
616
We then related this to trial-by-trial beta-band power. When we binned all
Switch trials by beta power into 8 non-overlapping bins, we saw a clear
relationship to decoding: the lower the beta-band power, the higher the
decoding (Fig. 4a, linear regression across 8 bins, t_{42}=-3.93, p=0.0003, d=-
0.60). This was not the case on Repeat trials (t_{42}=0.14, p=0.89, d=0.02),
leading to a significant difference between slopes (t_{42}=-2.68, p=0.011, d=-
0.41). Only beta power after the switch cue and shortly before probe onset
predicted decoding (Fig. 4b, sig. cluster 492-772ms, corrected p=0.0029,
difference to Repeat trials 532-772ms, p=0.0096). Beta power in the pre-
probe interval (400-800ms) predicted decoding throughout the post-probe
interval (Fig. 4c, 950-1,400ms after cue onset, p=0.0001, difference to repeat
trials 1,000-1,400ms, p=0.0033). Decoding before probe onset was much
weaker and not correlated with beta power (Fig. S5). At central channels,
only alpha- and beta-band power predicted decoding (Fig. 4d,e, 10-24Hz,
p=0.001, sig. difference to repeat trials 16-25Hz, p=0.0172).
632
Because of the relationship between item distance and both beta power and
accuracy, these effects could have been driven by differences in item
distance. To account for this possibility, we conducted a multiple regression
analysis with (z-scored) beta power, z-scored item distance and their
interaction as predictors and WM decoding as the dependent variable. The
effect of beta power on decoding remained significant for Switch trials (Fig.
4f, t_{42}=-4.420, p=6.82*10^{-5}, d=-0.67) and non-significant for Repeat trials
(t_{42}=0.672, p=0.506, Switch vs. Repeat: t_{42}=3.428, p=0.0014, d=-0.52) and we
saw no difference in the effect of item distance on decoding between Switch
and Repeat trials (t_{42}=1.03, p=0.31 for model including 15-25Hz power,
t_{42}=0.79, p=0.43 for model including 4-8Hz power), and there was no
significant interaction between beta power and item distance on decoding
(p>0.22).
646
We saw no comparable effect of theta power on Switch trials. In fact, there
appeared to be only a modest effect of theta power on decoding on Repeat
trials. This effect trended into an unexpected direction, with higher theta
power predicting lower decoding (Fig. 4g,h, sig. cluster 532-772ms, p=0.044,

also after correcting for item distance, Fig. 4l). This effect occurred in a
similar time range (Fig. 4i) as the beta power effect but appeared to be
statistically weaker and showed no clear peak in the frequency spectrum
(Fig. 4j) or strong effects at non-frontal channels (Fig. 4k). More importantly, it
appeared to depend on the choice of baseline and may have been driven by
fluctuations in pre-trial theta power. When we conducted the analysis without
baseline-correction (Fig. S4b) or after regressing baseline power out of the
signal (Fig. S4d), there was no longer a significant relationship between theta
power and decoding. Instead, increased theta power on switch trials
predicted increased decoding of the cued item late in the trial, mirroring the
beta power effect (Fig. S4b,d, third panel). Similarly, alpha power had no
effect on decoding of the cued item with or without baseline correction (Fig.
S2c,f). The beta power effect (Fig. S4a,c), on the other hand, remained
unchanged and significant.
665

Figure 4. *15-25Hz power on priority switch trials predicts decoding of the*
newly prioritised item. **A.** *Switch trials binned by 15-25Hz power (400-800ms,*
central channels) show a linear relationship with cued orientation decoding.
This is not the case on repeat trials. Dots and error bars show mean±s.e.m. of
the data. Line and shaded area shows mean±s.e.m. of linear fit. Inset:
Timecourse of cued orientation decoding strength (a.u.) from probe onset
(0.8s) to 600ms after cue onset (1.4), separately for Switch and Repeat trials.
B. *We regressed beta power across the timecourse of the trial against post-*
probe decoding (1-1.2s). Only beta power in the late cue interval predicted
later decoding (Switch trials only), with no effect on Repeat trials. **C.** *Switch*
trial beta power in the late cue period (400–800ms) predicted decoding
throughout the post-probe period, from 150ms after probe onset (0.95s) to
the end of the probe epoch (1.4s, 600ms after probe onset). **D.** *Cue interval*
power predicted decoding in a frequency range from 10-24Hz. **E.**
Topography of channel-wise power predicting decoding, peaking at central-
parietal channels. **F.** *Regression coefficient of beta power on decoding from a*
multiple regression model including item distance and power*item distance
interaction. Coloured horizontal lines show median, black horizontal lines

show mean. Dots are individual observers. **G-L.** *Frontal theta power does not*
show this effect, but higher theta power predicts lower decoding on repeat
trials (but see analysis without baseline correction, Fig. S4).
688
In sum, beta power had a significant effect on decoding only on switch trials,
suggesting a role in prioritising the newly relevant WM item. Conversely, an
insufficient reduction in beta power might reflect insufficient or ineffective
deprioritisation of the uncued item and thus predict its decoding remaining
high. To test this possibility, we repeated the analysis but with trial-by-trial
decoding of the uncued item. Decoding of uncued orientations was highly
significant on both Switch (1-1.4s after cue onset, corrected p=0.0001) and
Repeat trials (1.05-1.4s, p=0.0028, Switch>Repeat 1.25-1.4s, p=0.0295, Fig.
5a, left panel inset). Nevertheless, beta power had no effect on decoding
(main effect: $F_{6.29,264.14}=0.72$, p=0.64, Switch trials: $t_{42}=0.66$, p=0.51, Repeat
trials: $t_{42}=0.78$, p=0.44, Fig. 5a). This remained true after regressing out the
effect of item distance (all p>0.51, Fig. 5a, right panel). Theta power also did
not predict decoding of the uncued item (main effect: $F_{5.91,248.24}=0.319$,
p=0.92, Switch trials: $t_{42}=-1.06$, p=0.30, Repeat trials: $t_{42}=-0.56$, p=0.58, Fig.
5b). Regressing out the effect of item distance did not change this result (all
p>0.20).
705
A related possibility is that insufficient beta desynchronisation on Switches
leads to lingering activation of the uncued item in the representational format
of a cued item (i.e., in a prioritised state). To test this, we trained a decoder
on cued WM items and tested it on uncued items. We again found no
significant effect. Cross-decoding of uncued orientations (training on cued
orientations) was significant on both Switch (0.95-1.4s after cue onset,
corrected p=0.0001) and Repeat trials (0.95-1.4s, p=0.0002, Switch>Repeat
n.s. Fig. 5c, left panel inset). Beta power had no effect on decoding (main
effect: $F_{6.15,258.32}=1.78$, p=0.10, Switch trials: $t_{42}=0.05$, p=0.96, Repeat trials:
$t_{42}=-1.26$, p=0.21, Fig. 5c). This remained true after regressing out the effect
of item distance (all p>0.25, Fig. 5c, right panel). Theta power also did not
predict decoding of uncued orientations (main effect: $F_{6.14,258.03}=0.418$,
p=0.87, Switch trials: $t_{42}=-0.95$, p=0.35, Repeat trials: $t_{42}=0.80$, p=0.43, Fig.
5d). Regressing out the effect of item distance did not change this result (all
p>0.41, Fig. 5d, right panel). Alpha power again had no effect on decoding of
the uncued item (Fig. S2d,e).
722
723 We conclude that beta power reduction affects only the activation of a newly
724 cued item but not the removal of a previously relevant item from the
725 prioritised representational state.
726

Figure 5. *15-25Hz and 4-8Hz power on priority switch trials do not predict decoding of the uncued item.* **A.** *No relationship between 15-25Hz power and decoding of the uncued item.* **B.** *Similarly, 4-8Hz power had no effect.* **C-D.** *Crossdecoding the uncued item (train on cued, test on uncued) also showed no effect. Conventions as in Fig. 4.*

Discussion

Our study examined the role of neural oscillations in priority switches in working memory. Previous studies had identified predominantly theta and alpha oscillations, with relatively less emphasis on beta-band oscillations, but their respective contributions have remained unclear. Here we identified a central role for transient reductions in beta-band power in two aspects of switching priority in working memory: the magnitude of the representational shift and the fidelity of the update, as measured by the decoding strength of the newly prioritised item. By contrast, while priority switches also affected theta and alpha power, neither of those oscillations strongly correlated with update magnitude or fidelity.

At the behavioural level, priority switches incurred a temporary cost that manifested in reduced accuracy and increased reaction time, consistent with classic work on task switching (Koch et al., 2018; Monsell, 2003). The reduction in accuracy was linked to an increase in guess rate but no change in precision, suggesting that switch costs may have arisen because of occasional failures to prioritise but not because the representation itself was noisier. In line with this interpretation, switches also increased the rate of swap errors. Interestingly, the cost of switching priorities between memories scaled linearly with the angular distance between the two items. This was a surprisingly strong effect that explained the entirety of the switch cost on accuracy (although swap errors may have contributed to part of this effect). The dependence of switch costs on memory similarity suggests that updating the focus of attention is a graded operation, akin to mental rotation (Shepard & Metzler, 1971). This finding appears to be at odds with models of WM as a storage system for arbitrary content (Bocincova et al., 2022; Bouchacourt & Buschman, 2019; Manohar et al., 2019) since it suggests that control operations in WM reflect the representational space of features in WM. A comparable similarity effect has recently been reported for switch costs between different task rules (Bustos et al., 2024). Cognitive control operations more generally may therefore depend on the representational geometry of task-relevant features, consistent with reports of map-like representations for other cognitive functions (Bellmund et al., 2018).

At the neural level, we observed a robust beta-band (approx. 15-25Hz) power reduction following switch cues, particularly at central-parietal electrodes. Power reduction scaled with item distance, suggesting that beta oscillations reflect the magnitude of the required switch. Notably, this effect was specific to Switch trials and absent (or modestly reversed) on Repeat trials, while behaviour on Repeat trials still showed an effect of item distance. This specificity to switch trials suggests a role for beta in the switching process, rather than coding for item distance per se. It is unclear why switches between more dissimilar items require a greater reduction in beta amplitude. One possibility is that beta tracks the duration of the switching process, and switching to a more dissimilar item takes longer (as with mental rotation). More dissimilar Items could also be encoded in more spatially distant populations, possibly requiring stronger beta reduction (Lundqvist et al.,

782   2023) during a switch. We stress, however, that these are speculative
783   explanations that cannot be resolved with the current data.
784
Importantly, beta power on priority switches also predicted the fidelity of the
updated representation. On Switch trials, lower beta power was associated
with stronger decoding of the newly prioritised item later in the decision
phase. The correlation was specific to beta frequencies, to the cued (but not
the uncued) item, and occurred only on Switch trials. This suggests that beta
oscillations are involved in the selection and prioritisation of relevant memory
items rather than the removal of de-prioritised items. This conclusion appears
to contrast with previous reports of beta oscillations playing a key role in
clearing items from WM when they are no longer needed (Lundqvist et al.,
2018). In our paradigm, however, items were not fully removed from memory,
as uncued items could still be probed later in the block. Beta oscillations may
therefore play distinct roles in updating and clearing WM (Lundqvist et al.,
2023, 2024). Beta-band desynchronisation has previously been implicated in
increasing the coding capacity of neural populations (Hanslmayr et al., 2012)
to aid reinstatement of long-term memories (Griffiths et al., 2019), consistent
with the proposed role in updating in our task. We also observed a temporal
progression of events, with beta desynchronisation occurring before probe
onset but affecting decoding after probe onset. While it is possible that this
was simply because stronger decoding after the probe made it easier to find
a significant correlation, it may hint at a temporal sequence in which a
reduction in beta power initiates a preparatory cascade of events from
identifying and accessing the cued item to then prioritising it before this is
reflected in improved decoding when the cued item is needed for decision-
making. We speculate that reductions in beta bursts may not only serve to
prepare for upcoming external updates of WM (Liljefors et al., 2024) but also
for internal updates within the focus of attention. Consistent with this view, a
related line of work has suggested that beta synchronisation helps in the
selection and downstream routing of behaviourally relevant neural ensembles
(Rassi et al., 2023; Salazar et al., 2012; Spitzer & Haegens, 2017).
814
While the limited spatial resolution of EEG prevents us from drawing strong
conclusions about anatomical sources of the beta-band effect, its well-
documented involvement in regulating activity between cortex and the basal
ganglia (Brittain & Brown, 2014; Engel & Fries, 2010; Jenkinson & Brown,
2011; Schmidt et al., 2019) suggests beta synchrony in corticostriatal loops
could orchestrate WM prioritisation. Along these lines, the basal ganglia have
been proposed as a source of selecting from WM for the next action
(Chatham et al., 2014; Hazy et al., 2006). Other work has suggested a direct
role for beta desynchronisation in the motor system in WM tasks which
permit preparation of an action before recall (Boettcher et al., 2021;
Schneider et al., 2017). This is unlikely in our case because participants could
not anticipate the correct response (probe rotated
clockwise/counterclockwise) at the time of the peak beta effect (before probe
onset). More broadly, surprise at the less frequent Switch trials (approx. 30%
of trials) could have led to inhibition of the motor system (Wessel & Aron,

2013, 2017), though it is unclear why such an effect would scale with update
magnitude or predict the strength of decoding. In sum, while several
processes likely occur during prioritisation, the specificity of the beta-band
effects in our study are most consistent with a role in selecting and
prioritising new information.
835
Theta oscillations, while elevated on Switch trials, did not scale with item
distance and showed no consistent relationship with decoding strength. This
suggests that theta may reflect a more general, possibly all-or-none, process
signaling control demands rather than the implementation of specific
updates. This interpretation is consistent with the apparent role of delta/theta
in regulating posterior alpha desynchronisation when space-based selection
of WM contents is possible (de Vries et al., 2020). It also aligns with a
longstanding proposal that midline frontal theta signals cognitive control
demands more generally (Cavanagh & Frank, 2014) rather than executing
control.
846
Alpha oscillations, like beta, showed a significant reduction on Switch trials in
our study. However, as with theta, we did not observe any correlations with
update magnitude or decoding. Spatially lateralised alpha oscillations have
previously been linked to successful prioritisation of cued WM contents
(Boettcher et al., 2021; Myers, Walther, et al., 2015; Poch et al., 2014; Wallis
et al., 2015). This discrepancy may be due to the non-spatial nature of our
task, where items were cued via auditory tones rather than by spatial
location. Future work could untangle the contributions of different
prioritisation mechanisms by comparing spatial with non-spatial selection.
856
Taken together, our results highlight a previously underappreciated role for
beta oscillations in the dynamic reorganisation of WM contents. Beta power
reduction appears to facilitate the transition of items into the prioritised state,
with the degree of reduction reflecting both the magnitude of the update and
the quality of the resulting representation. We saw no such relationship with
the uncued item, indicating that beta in our task was not directly correlated
with deprioritising previously cued information. However, swap errors were
more frequent after priority switches, indicating occasional failures to update
the focus of attention, and beta could be relevant to this process or in
suppressing the uncued item more generally. While recent work has begun to
address the neural mechanisms of swap errors in working memory (Alleman
et al., 2024), the role of neural oscillations in this process is unknown and
appears to be a fruitful avenue for further research. These findings more
broadly highlight the potential relationship between neural oscillations and
interference-free maintenance of multiple task representations (Fang et al.,
2025). Future work could aim to further dissociate the causal role and
specific subprocesses of priority switching supported by beta-band
oscillations by examining prioritisation efficacy when beta oscillations are
entrained artificially or when they are affected by neurological conditions
such as Parkinson's disease.
877

References

Alleman, M., Panichello, M., Buschman, T. J., & Johnston, W. J. (2024). The neural basis of swap errors in working memory. *Proceedings of the National Academy of Sciences*, *121*(33), e2401032121.

Bays, P. M., Catalao, R. F., & Husain, M. (2009). The precision of visual working memory is set by allocation of a shared resource. *Journal of Vision*, *9*(10), 7, 1-11.

Bellmund, J. L. S., Gärdenfors, P., Moser, E. I., & Doeller, C. F. (2018). Navigating cognition: Spatial codes for human thinking. *Science*, *362*(6415). https://doi.org/10.1126/science.aat6766.

Berger, B., Griesmayr, B., Minarik, T., Biel, A. L., Pinal, D., Sterr, A., & Sauseng, P. (2019). Dynamic regulation of interregional cortical communication by slow brain oscillations during working memory. *Nature Communications*, *10*(1), 4242. https://doi.org/10.1038/s41467-019-12057-0

Bocincova, A., Buschman, T. J., Stokes, M. G., & Manohar, S. G. (2022). Neural signature of flexible coding in prefrontal cortex. *Proceedings of the National Academy of Sciences*, *119*(40), e2200400119. https://doi.org/10.1073/pnas.2200400119

Boettcher, S. E. P., Gresch, D., Nobre, A. C., & Van Ede, F. (2021). Output planning at the input stage in visual working memory. *Science Advances*, *7*(13). https://doi.org/10.1126/sciadv.abe8212

Bouchacourt, F., & Buschman, T. J. (2019). A Flexible Model of Working Memory. *Neuron*, *103*(1), 147-160.e8.

Brainard, D. H. (1997). The Psychophysics Toolbox. *Spatial Vision*, *10*(4), 433–436.

Brittain, J.-S., & Brown, P. (2014). Oscillations and the basal ganglia: Motor control and beyond. *Neuroimage*, *85*, 637–647.

Bustos, B., Mordkoff, J. T., Hazeltine, E., & Jiang, J. (2024). Task switch costs scale with dissimilarity between task rules. *Journal of Experimental Psychology: General*, *153*(7), 1873.

Cavanagh, J. F., & Frank, M. J. (2014). Frontal theta as a mechanism for cognitive control. *Trends in Cognitive Sciences*, *18*(8), 414–421.

Chatham, C. H., & Badre, D. (2015). Multiple gates on working memory. *Current Opinion in Behavioral Sciences*, *1*, 23–31.

Chatham, C. H., Frank, M. J., & Badre, D. (2014). Corticostriatal Output Gating during Selection from Working Memory. *Neuron*, *81*(4), 930–942.

Christophel, T. B., Iamshchinina, P., Yan, C., Allefeld, C., & Haynes, J.-D. (2018). Cortical specialization for attended versus unattended working memory. *Nature Neuroscience*, *21*(4), 494–496.

Cichy, R. M., Ramirez, F. M., & Pantazis, D. (2015). Can visual information encoded in cortical columns be decoded from magnetoencephalography data in humans? *NeuroImage*, *121*, 193–204.

Cowan, N. (2000). The magical number 4 in short-term memory: A reconsideration of mental storage capacity. *Behavioral and Brain Sciences*, *24*(1), 87-114-discussion 114-85.

D'Ardenne, K., Eshel, N., Luka, J., Lenartowicz, A., Nystrom, L. E., & Cohen,
J. D. (2012). Role of prefrontal cortex and the midbrain dopamine
system in working memory updating. *Proceedings of the National*
Academy of Sciences, 19900–19909.
de Vries, I. E. J., Slagter, H. A., & Olivers, C. N. L. (2020). Oscillatory Control
over Representational States in Working Memory. *Trends in Cognitive*
Sciences, *24*(2), 150–162.
de Vries, I. E. J., van Driel, J., Karacaoglu, M., & Olivers, C. N. L. (2018).
Priority Switches in Visual Working Memory are Supported by Frontal
Delta and Posterior Alpha Interactions. *Cerebral Cortex (New York,*
N.Y.: 1991), *28*(11), 4090–4104.
de Vries, I. E. J., van Driel, J., & Olivers, C. N. L. (2017). Posterior α EEG
Dynamics Dissociate Current from Future Goals in Working Memory-
Guided Visual Search. *Journal of Neuroscience*, *37*(6), 1591–1603.
El-Gaby, M., Harris, A. L., Whittington, J. C., Dorrell, W., Bhomick, A., Walton,
M. E., Akam, T., & Behrens, T. E. (2024). A cellular basis for mapping
behavioural structure. *Nature*, *636*(8043), 671–680.
Engel, A. K., & Fries, P. (2010). Beta-band oscillations—Signalling the status
quo? *Current Opinion in Neurobiology*, *20*(2), 156–165.
Fang, W., Jiang, X., Chen, J., Zhang, C., & Wang, L. (2025). Oscillatory
control over representational geometry of sequence working memory in
macaque frontal cortex. *Current Biology*, *35*(7), 1495–1507.
Griffin, I. C., & Nobre, A. C. (2003). Orienting attention to locations in internal
representations. *Journal of Cognitive Neuroscience*, *15*(8), 1176–1194.
Griffiths, B. J., Mayhew, S. D., Mullinger, K. J., Jorge, J., Charest, I., Wimber,
M., & Hanslmayr, S. (2019). Alpha/beta power decreases track the
fidelity of stimulus-specific information. *Elife*, *8*, e49562.
Hanslmayr, S., Staudigl, T., & Fellner, M.-C. (2012). Oscillatory power
decreases and long-term memory: The information via
desynchronization hypothesis. *Frontiers in Human Neuroscience*, *6*,
74.
Hazy, T. E., Frank, M. J., & Reilly, R. O. (2006). Banishing the homunculus:
Making working memory work. *Neuroscience, 139*(1), 105-118.
Iamshchinina, P., Christophel, T. B., Gayet, S., & Rademaker, R. L. (2021).
Essential considerations for exploring visual working memory storage
in the human brain. *Visual Cognition*, *29*(7), 425–436.
Jenkinson, N., & Brown, P. (2011). New insights into the relationship between
dopamine, beta oscillations and motor function. *Trends in*
Neurosciences, *34*(12), 611–618.
Johnson, E. L., Adams, J. N., Solbakk, A.-K., Endestad, T., Larsson, P. G.,
Ivanovic, J., Meling, T. R., Lin, J. J., & Knight, R. T. (2018). Dynamic
frontotemporal systems process space and time in working memory.
PLoS Biology, *16*(3), e2004274.
Koch, I., Poljac, E., Müller, H., & Kiesel, A. (2018). Cognitive structure,
flexibility, and plasticity in human multitasking—An integrative review of
dual-task and task-switching research. *Psychological Bulletin*, *144*(6),
557–583.
Ledoit, O., & Wolf, M. (2004). A well-conditioned estimator for large-

dimensional covariance matrices. *Journal of Multivariate Analysis*,
88(2), 365–411.

Liebe, S., Hoerzer, G. M., Logothetis, N. K., & Rainer, G. (2012). Theta
coupling between V4 and prefrontal cortex predicts visual short-term
memory performance. *Nature Neuroscience*, *15*(3), 456–462.

Liljefors, J., Almeida, R., Rane, G., Lundström, J. N., Herman, P., &
Lundqvist, M. (2024). Distinct functions for beta and alpha bursts in
gating of human working memory. *Nature Communications*, *15*(1),
8950.

Lundqvist, M., Brincat, S. L., Rose, J., Warden, M. R., Buschman, T. J., Miller,
E. K., & Herman, P. (2023). Working memory control dynamics follow
principles of spatial computing. *Nature Communications*, *14*(1), 1429.

Lundqvist, M., Herman, P., Warden, M. R., Brincat, S. L., & Miller, E. K.
(2018). Gamma and beta bursts during working memory readout
suggest roles in its volitional control. *Nature Communications*, 1–12.

Lundqvist, M., Miller, E. K., Nordmark, J., Liljefors, J., & Herman, P. (2024).
Beta: Bursts of cognition. *Trends in Cognitive Sciences*, *28*(7), 662–
676. https://doi.org/10.1016/j.tics.2024.03.010

Lundqvist, M., Rose, J., Herman, P., Brincat, S. L., Buschman, T. J., & Miller,
E. K. (2016). Gamma and Beta Bursts Underlie Working Memory.
Neuron, *90*(1), 152–164.

Manohar, S. G., Zokaei, N., Fallon, S. J., Vogels, T. P., & Husain, M. (2019).
Neural mechanisms of attending to items in working memory.
Neuroscience & Biobehavioral Reviews, *101*, 1–12.

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG-
and MEG-data. *Journal of Neuroscience Methods*, *164*(1), 177–190.

Miller, E. K., Lundqvist, M., & Bastos, A. M. (2018). Working Memory 2.0.
Neuron, *100*(2), 463–475.

Monsell, S. (2003). Task switching. *Trends in Cognitive Sciences*, *7*(3), 134–
140.

Muhle-Karbe, P. S., Myers, N. E., & Stokes, M. G. (2021). A Hierarchy of
Functional States in Working Memory. *The Journal of Neuroscience*,
41(20), 4461–4475. https://doi.org/10.1523/JNEUROSCI.3104-20.2021

Myers, N. E., Rohenkohl, G., Wyart, V., Woolrich, M. W., Nobre, A. C., &
Stokes, M. G. (2015). Testing sensory evidence against mnemonic
templates. *eLife*, *4*, e09000. https://doi.org/10.7554/eLife.09000

Myers, N. E., Walther, L., Wallis, G., Stokes, M. G., & Nobre, A. C. (2015).
Temporal Dynamics of Attention during Encoding versus Maintenance
of Working Memory: Complementary Views from Event-related
Potentials and Alpha-band Oscillations. *Journal of Cognitive
Neuroscience*, *27*(3), 492–508.

Oberauer, K. (2002). Access to information in working memory: Exploring the
focus of attention. *Journal of Experimental Psychology: Learning,
Memory, and Cognition*, *28*(3), 411–421.

Olivers, C. N. L., Peters, J., Houtkamp, R., & Roelfsema, P. R. (2011).
Different states in visual working memory: When it guides attention and
when it does not. *Trends in Cognitive Sciences*, *15*(7), 327–334.

Paluch, K., Magnuski, M., Średniawa, W., Ivanovski, D., Rysz, A., Służewska-

Niedźwiedź, M., Pasterski, T., Fortuna, W., Smarzewska, K., &
Reinacher, P. C. (2025). Unattended working memory items are coded
by persistent activity in human medial temporal lobe neurons. *Nature*
Human Behaviour, 1–15.
Panichello, M. F., & Buschman, T. J. (2021). Shared mechanisms underlie the
control of working memory and attention. *Nature, 592*(7855), 601–605.
Poch, C., Campo, P., & Barnes, G. R. (2014). Modulation of alpha and
gamma oscillations related to retrospectively orienting attention within
working memory. *European Journal of Neuroscience, 40*, 2399–2405.
Rassi, E., Zhang, Y., Mendoza, G., Méndez, J. C., Merchant, H., & Haegens,
S. (2023). Distinct beta frequencies reflect categorical decisions.
Nature Communications, 14(1), 2923. https://doi.org/10.1038/s41467-
023-38675-3
Rerko, L., & Oberauer, K. (2013). Focused, unfocused, and defocused
information in working memory. *Journal of Experimental Psychology:*
Learning, Memory, and Cognition, 39(4), 1075–1096.
Rerko, L., Souza, A. S., & Oberauer, K. (2014). Retro-cue benefits in working
memory without sustained focal attention. *Memory & Cognition, 42*(5),
712–728.
Riddle, J., McPherson, T., Sheikh, A., Shin, H., Hadar, E., & Frohlich, F.
(2024). Internal representations are prioritized by frontoparietal theta
connectivity and suppressed by alpha oscillation dynamics: Evidence
from concurrent transcranial magnetic stimulation EEG and invasive
EEG. *Journal of Neuroscience, 44*(15).
https://www.jneurosci.org/content/44/15/e1381232024.abstract
Riddle, J., Scimeca, J. M., Cellier, D., Dhanani, S., & D'Esposito, M. (2020).
Causal evidence for a role of theta and alpha oscillations in the control
of working memory. *Current Biology, 30*(9), 1748–1754.
Roux, F., & Uhlhaas, P. J. (2014). Working memory and neural oscillations:
Alpha–gamma versus theta–gamma codes for distinct WM
information? *Trends in Cognitive Sciences, 18*(1), 16–25.
Salazar, R. F., Dotson, N. M., Bressler, S. L., & Gray, C. M. (2012). Content-
Specific Fronto-Parietal Synchronization During Visual Working
Memory. *Science, 338*(6110), 1097–1100.
Schmidt, R., Ruiz, M. H., Kilavik, B. E., Lundqvist, M., Starr, P. A., & Aron, A.
R. (2019). Beta Oscillations in Working Memory, Executive Control of
Movement and Thought, and Sensorimotor Function. *Journal of*
Neuroscience, 39(42), 8231–8238.
https://doi.org/10.1523/JNEUROSCI.1163-19.2019
Schneider, D., Barth, A., & Wascher, E. (2017). On the contribution of motor
planning to the retroactive cuing benefit in working memory: Evidence
by mu and beta oscillatory activity in the EEG. *NeuroImage, 162*, 73–
85.
Shepard, R. N., & Metzler, J. (1971). Mental Rotation of Three-Dimensional
Objects. *Science, 171*(3972), 701–703.
https://doi.org/10.1126/science.171.3972.701
Souza, A. S., & Oberauer, K. (2016). In search of the focus of attention in
working memory: 13 years of the retro-cue effect. *Attention,*

Perception, & Psychophysics, 1–22.

Spitzer, B., & Haegens, S. (2017). Beyond the status quo: A role for beta
oscillations in endogenous content (re) activation. *Eneuro, 4*(4).
https://www.eneuro.org/content/4/4/ENEURO.0170-17.2017.short

Sprague, T. C., Ester, E. F., & Serences, J. T. (2016). Restoring Latent Visual
Working Memory Representations in Human Cortex. *Neuron, 91*(3),
694–707.

van Ede, F., Niklaus, M., & Nobre, A. C. (2016). Temporal expectations guide
dynamic prioritization in visual working memory through attenuated
alpha oscillations. *The Journal of Neuroscience*, 2272–16.

Van Ede, F., & Nobre, A. C. (2023). Turning Attention Inside Out: How
Working Memory Serves Behavior. *Annual Review of Psychology*,
74(1), 137–165. https://doi.org/10.1146/annurev-psych-021422-041757

van Loon, A. M., Olmos-Solis, K., Fahrenfort, J. J., & Olivers, C. N. (2018).
Current and future goals are represented in opposite patterns in object-
selective cortex. *eLife, 7*, 1427.

Wallis, G., Stokes, M., Cousijn, H., Woolrich, M., & Nobre, A. C. (2015).
Frontoparietal and Cingulo-opercular Networks Play Dissociable Roles
in Control of Working Memory. *Journal of Cognitive Neuroscience, 15*,
1–16.

Wessel, J. R., & Aron, A. R. (2013). Unexpected events induce motor slowing
via a brain mechanism for action-stopping with global suppressive
effects. *Journal of Neuroscience, 33*(47), 18481–18491.

Wessel, J. R., & Aron, A. R. (2017). On the globality of motor suppression:
Unexpected events and their influence on behavior and cognition.
Neuron, 93(2), 259–280.

Wolff, M. J., Jochim, J., Akyürek, E. G., Buschman, T. J., & Stokes, M. G.
(2020). Drifting codes within a stable coding scheme for working
memory. *PLoS Biology, 18*(3), e3000625.

Wolff, M. J., Jochim, J., Akyürek, E. G., & Stokes, M. G. (2017). Dynamic
hidden states underlying working-memory-guided behavior. *Nature*
Neuroscience, 20(6), 864–871.

Wolff, M. J., Kandemir, G., Stokes, M. G., & Akyürek, E. G. (2020). Unimodal
and Bimodal Access to Sensory Working Memories by Auditory and
Visual Impulses. *Journal of Neuroscience, 40*(3), 671–681.

Yu, Q., Teng, C., & Postle, B. R. (2020). Different states of priority recruit
different neural representations in visual working memory. *PLoS*
Biology, 18(6), e3000769.

1108

1109 **Supplementary Results to**
1110 **Human Beta Oscillations Reflect Magnitude and Fidelity of Priority Shifts**
1111 **in Working Memory**
1112

1113
Figure S1. *Independent Effects of Swap Errors and Item Distance on Switch*
trials. **A.** *Psychometric curves showing proportion of trials judged clockwise,*
sorted by probe orientation relative to cued WM item. **B.** *Responses sorted*
relative to the uncued item, indicating swap errors. Dots show data (whiskers
are standard error of the mean), lines show predicted response from fit of a
mixture model (shaded area is standard error the mean across participants).
C.-E. *Mixture model parameters (inverse precision, guess/lapse rate, and*
swap rate) are shown in the right panels. **F.** *Effect of item distance (left panel)*
and congruency (right panel) on accuracy. **G.** *Effect on reaction time.*
1123
1124 Inverse precision (σ), p_{guess}, and p_{swap} were fit using maximum likelihood
1125 estimation separately for Switch and Repeat trials.
1126
1127 Swap errors were significantly higher on Switch trials (Switch: 10.1±1.3%)
1128 than on first Repeat trials (Repeat1: 6.2±0.9%, t_{42}=3.99, p=2.566*10^{-4},
1129 Cohen's d=0.61) or subsequent Repeat trials (Repeat2+: 6.6±1.0%, t_{42}=3.16,
1130 p=2.930*10^{-3}, d=0.48). Swap errors did not differ between Repeat1 and
1131 Repeat2+ trials (t_{42}=-0.73, p=0.469, d=-0.11).
1132
1133 Precision still did not differ significantly between Switch (22.5±1.2º) and
1134 Repeat1 (21.3±0.9º) trials (t_{42}=1.84, p=0.073, d=0.28), although there was a
1135 trend. Precision on Repeat2+ trials (21.6±0.8º) was no different from Switch

1136 trials (t_{42}=1.38, p=0.174, d=0.21) or Repeat1 trials (t_{42}=-0.64, p=0.526, d=-
1137 0.10).

1138
1139 Similarly, the guess/lapse rate was still significantly higher on Switch
1140 (5.0±0.9%) than on Repeat1 (2.5±0.5%) trials (t_{42}=3.66, p=6.987*10^{-4}, d=0.56)
1141 and Repeat2+ trials (1.7±0.4%, t_{42}=3.61, p=8.15*10^{-4}, d=0.55). Guess rates
1142 did not differ between Repeat1 and Repeat2+ trials (t_{42}=1.42, p=0.164,
1143 d=0.22).

1144
While the mixture modeling suggests that swap errors contributed to switch
costs, it does not rule out that item distance contributed as well. We
quantified both effects in tandem using multiple regression. We reasoned
that if swap errors contributed to behaviour, they would reduce accuracy
when the correct response relative to the uncued WM item was different from
that to the cued item (i.e., incongruent trials). For example, if the cued item
was 30º and the probe orientation was 45º, the correct response would be
'counterclockwise'. If the uncued item was 60º, a swap error might produce a
'clockwise' judgment, resulting in an incorrect response. By contrast, swap
errors should not reduce accuracy on congruent trials. We used trialwise
logistic regression to predict accuracy from item distance and congruency.
We also included a number of nuisance regressors (the absolute angle
between the probe and the cued item, between the probe and the uncued
item, the block number, and the trial number within the block) which we had
found to influence behaviour in our previous publication (Muhle-Karbe et al.,
2021). We then compared regression coefficients against 0 and across
Switch vs. Repeat trials. We repeated the same analysis (using linear
regression) on reaction time (on correct trials only).

1163
Even after taking potential swap errors into account (as measured by
congruency), item distance had a significant negative effect on accuracy on
Switch trials (t_{42}=-4.36, p=8.2*10^{-5}, Cohen's d=-0.66) and on the first Repeat
trial (t_{42}=-3.29, p=2.04*10^{-3}, d=-0.50), but not subsequent trials (t_{42}=-1.182,
p=0.244, d=-0.18). The effect on Switch trials was marginally stronger
compared to the first Repeat trial (t_{42}=2.03, p=0.0486, d=-0.31) and
compared to subsequent Repeat trials (t_{42}=3.35, p=0.00171, d=-0.51). The
effect was not different on the first vs. subsequent Repeat trials (t_{42}=-1.82,
p=0.0756, d=-0.28), although there was a trend.

1173
At the same time, congruency also affected accuracy on all three trial types
(Switch: t_{42}=3.65, p=7.25*10^{-4}, d=0.56; Repeat1: t_{42}=2.6427, p=0.0115,
d=0.40; Repeat2+: t_{42}=4.67, p=3.05*10^{-5}, d=0.71). The congruency effect was
marginally stronger on Switch trials than the first Repeat trial (t_{42}=-2.001,
p=0.0519, d=0.31), consistent with the psychometric modeling results above,
but not on Switch vs. subsequent Repeat trials (t_{42}=-0.345, p=0.732, d=0.05).
The effect was also marginally but non-significantly smaller on the first
Repeat trial compared to subsequent Repeats (t_{42}=-1.837, p=0.0733, d=-
0.28).

1183

1184   In sum, item distance and congruency (as a measure of swap errors) made
1185   independent contributions to performance, and both appeared to be
1186   modulated on Switch trials.
1187
1188   For completeness, we also examined the effect of item distance and
1189   congruency on reaction time (RT). In brief, item distance had a significant
1190   effect on RT that was strongest on Switch trials, while congruency had no
1191   effect on RT.
1192
Item Distance had a significant slowing effect on RT on all trial types (Switch:
$t_{42}=9.12$, $p=1.62*10^{-11}$, $d=1.39$; Repeat1: $t_{42}=5.57$, $p=1.65*10^{-6}$, $d=0.85$;
Repeat2+: $t_{42}=4.27$, $p=1.11*10^{-4}$, $d=0.65$). The effect was stronger on Switch
trials than on the first ($t_{42}=-5.89$, $p=5.71*10^{-7}$, $d=0.90$) or subsequent Repeats
($t_{42}=-8.35$, $p=1.83*10^{-10}$, $d=1.27$). The effect was also stronger on the first
Repeat compared to subsequent Repeat trials ($t_{42}=2.84$, $p=7.00*10^{-3}$, $d=0.43$).
1199
Congruency had no significant effect on RT on any trial type (Switch: $t_{42}=-$
1.21, $p=0.232$; Repeat1: $t_{42}=0.85$, $p=0.399$; Repeat2+: $t_{42}=-1.14$, $p=0.261$).
There were no differences between conditions (Switch vs. Repeat1: $t_{42}=1.59$,
$p=0.118$, $d=-0.24$; Switch vs Repeat2+: $t_{42}=0.55$, $p=0.582$, $d=-0.08$; Repeat1
vs Repeat2+: $t_{42}=1.51$, $p=0.137$, $d=0.23$).

Figure S2. A. *Posterior Alpha Power desynchronised on Switch trials.*
Desynchronisation was most pronounced in the low alpha band (<10Hz) on
Switch trials (left panel). Desynchronisation was strongest at posterior-parietal
sensors (second panel). The desynchronisation appears limited to the theta-
low alpha range. We also observed a desynchronisation between 30-40Hz on
Switch trials (third panel). The timecourse of desynchronisation is limited to a
short period in the second half of the cue-target interval (right panel). Plotting
conventions as in Fig. 2. **B.** *Posterior Alpha Power does not scale with item*
distance on Switch trials. There is, however, a trend towards a positive
relationship between alpha power and item distance on Repeat trials. Plotting
conventions as in Fig. 3. **C.** *No relationship between alpha-band*
desynchronisation and decoding of the cued item, **D.** *uncued item, or* **E.**
cross-decoding of the uncued item (trained on the cued item). **F.** *There is also*
no effect when alpha-band power is used without baseline-correction.
Plotting conventions as in Fig. 4.
1221
Supplementary Results of the Analysis of Item Distance Effects on Alpha
Power
Alpha power in the late cue period (400-800ms, 8-14Hz, posterior channels)
was lower on Switch trials ($F_{1,42}=5.80, p=0.021,$ partial $\eta^2=0.12$) but did not
scale with item distance ($F_{6,252}=1.41, p=0.21,$ partial $\eta^2=0.006,$ left panel).
There was also no interaction between Switch and item distance
($F_{6,252}=1.162, p=0.33,$ partial $\eta^2=0.004$). A linear regression of item distance on
alpha power did not show any effect on Switch trials ($t_{42}=-1.25, p=0.22, d=-$
0.19) or Repeat trials ($t_{42}=0.88, p=0.38, d=0.13$). Nevertheless, when
examining the difference in regression slopes between Switch and Repeat
trials separately for each timepoint and frequency, we found a small but
significant cluster (p=0.0234) late in the cue-target interval that extended into
the beta band (approximately 500-800ms, 7-24Hz, second panel). This effect
was restricted to posterior sensors (third panel) and appeared to be driven by
a small positive effect on Repeat trials late in the delay (Repeat>0, 562-
782ms, p=0.0353, Repeat>Switch, p=0.0352, upper right panel). The effect
was not restricted to the alpha band, ranging from 10-21Hz (p=0.016). It thus
may have depended to some extent on the beta-band effect (Fig. 3). An
additional effect appeared in the low gamma band (31-40Hz, p=0.019).
1241

1242

Figure S3. *Beta- and Theta-power differences between Switch trials, first Repeats (Repeat1), and subsequent Repeats (Repeat2). Switch trials differed from Repeat1 trials (rows 1-2, from Fig. 2, repeated here for comparison) and from Repeat 2 trials (rows 3-4). While Repeat1 and Repeat2 trials were qualitatively similar, Repeat1 trials showed lower central alpha/beta power. There were no theta-band differences. Plotting conventions as in Figure 2.*

Figure S4. A. *Omitting baseline correction has no effect on the relationship between beta-power and decoding on Switch trials.* **B.** *However, it substantially affects the relationship between theta-power and decoding, where now a positive correlation emerges for decoding late in the trial.* **C.** *Alternatively, regressing baseline power out of the signal also has no effect on the relationship between beta-power and decoding on Switch trials.* **D.** *Again it affects the relationship between theta-power and decoding, where a positive*

1259 *correlation emerges for decoding late in the trial. +, p<0.10. Other plotting*
1260 *conventions as in Figure 4.*
1261

1262
Figure S5. *Decoding emerges around the time of probe onset (0.8s), but is*
much weaker in the cue-probe interval (left panels). Cued item decoding is
indistinguishable on Switch (blue) and Repeat (grey) trials. Beta power on
Switch trials (0.4-0.8s) only predicted decoding after probe onset (right panels).
When EEG features used for decoding were not demeaned over time (bottom
row), the results remained virtually unchanged.
1269