1 **Title:** Strigolactone effects on *Sorghum bicolor* ecophysiology and symbioses
2

3 **Authors:** Chloee M. McLaughlin[1,2,3*], Margarita Takou[2,4], Joel Masanga[2], Erica H. Lawrence-
4 Paul[2], Evelyn J. Abraham[1], Melanie Perryman[5], Anna Calabritto[6], Amandeep Cheema[5,7], Yuxing
5 Xu[2], Baloua Nebie[8], Steven Runo[9], Joshua J. Kellogg[1,10], Roberta Croce[6], Daniel P.
6 Schachtman[11], Huirong Gao[12], Ruairidh J. H. Sawers[5], Jesse R. Lasky[2,13]
7
[1] Intercollege Graduate Degree Program in Plant Biology, Pennsylvania State University,
University Park, PA, USA
[2] Department of Biology, Pennsylvania State University, University Park, PA, USA
[3] Genome Sequencing Center, HudsonAlpha Institute for Biotechnology, Huntsville, AL, USA
[4] Thünen Institute of Forest Genetics, Großhansdorf, Germany
[5] Department of Plant Science, Pennsylvania State University, University Park, PA, USA
[6] Department of Physics and Astronomy and Institute for Lasers, Life and Biophotonics, Faculty
of Sciences, VU University Amsterdam, 1081 HV Amsterdam, Netherlands
[7] Department of Biology and Program in Cell and Molecular Biology, Colorado State University,
Fort Collins, CO USA
[8] West and Central Africa sorghum improvement Program, International Maize and Wheat
Improvement Center (CIMMYT), Dakar, Senegal
[9] Department of Biochemistry, Microbiology and Biotechnology, Kenyatta University, Nairobi,
Kenya
[10] Department of Veterinary & Biomedical Sciences, Pennsylvania State University, University
Park, PA, USA
[11] Department of Agronomy and Horticulture, and Center for Plant Science Innovation,
University of Nebraska-Lincoln, Lincoln, NE, USA
[12] Corteva Agriscience, Johnston, IA, USA
[13] PAC Herbarium, Pennsylvania State University, University Park, Pennsylvania, PA, USA
28
29 *Correspondence CMM (cmclaughlin@hudsonalpha.org)

34
35 **Running head:** Strigolactone effects on *Sorghum bicolor*
36
37
38

Abstract: Strigolactones are ecologically, developmentally, and physiologically important
hormones, but much remains unknown about their evolution and role in non-model species.
Sorghum is a globally important C_4 cereal and exhibits natural variation in root-exuded
strigolactones. Differences in sorghum strigolactone stereochemistry are associated with
resistance to parasitic plants, but with evidence for potential trade-offs. We studied sorghum
mutants of loci in the strigolactone biosynthetic pathway, *CAROTENOID CLEAVAGE*
DIOXYGENASE 8 (Sb*CCD8b*) and *LOW GERMINATION STIMULANT 1* (*LGS1*), previously
shown to be *Striga* resistant by stimulating little germination of the parasite. Sb*CCD8b* CRISPR-
Cas9 deletions changed the accumulation of low abundance metabolites, reduced net carbon
assimilation rate, altered root architecture and anatomy, and diminished the establishment and
benefit of mycorrhizal symbionts. For *Striga*-resistant *LGS1* CRISPR-Cas9 deletions,
differentially expressed genes were enriched with promoter motifs for stress response and growth
pathways, net carbon assimilation rate was reduced, and the colonization of mycorrhizal
symbionts was delayed. We additionally restored functional *LGS1* into the RTx430 genetic
background, which normally has the *lgs1*-2 natural deletion allele. While root exudates from
LGS1 insertion mutants rescued *Striga* susceptibility, we did not see consistent rescue of other
traits impacted in *LGS1* loss-of-function mutants. We hypothesize that epistasis with a
neighboring strigolactone synthesis gene, which is rarely lost without concomitant loss of *LGS1,*
may alter the phenotypic effects of *LGS1* variation. Our study gives context to potential trade-
offs associated with host resistance to parasitic plants and, more broadly, builds on the
contribution of strigolactones in shaping sorghum physiological processes, growth, and
development.

Introduction

Strigolactones, originally named for their role as germination stimulants to the parasitic plant *Striga* spp. (Cook et al., 1966) are well established as an important class of hormones for plant biology and ecological interactions (Saeed et al., 2017; Wu et al., 2022). Strigolactones play key roles in various aspects of plant growth, physiology, and symbiotic interactions, including the regulation of shoot architecture, root development, light harvesting complexes, and associations with mycorrhizal fungi (Akiyama et al., 2005; Gomez-Roldan et al., 2008; Mayzlish-Gati et al., 2010; Ruyter-Spira et al., 2011). As signaling molecules, strigolactones enable plants to regulate their growth and development in response to the external environment (Barbier et al., 2023). To date, more than 30 strigolactones have been characterized which differ in their structure (Yoneyama and Brewer, 2021), though much of the functional variation associated with this structural diversity remains to be elucidated.

Strigolactones are synthesized from carotenoids which undergo catalytic reactions by several enzymes. Their biosynthesis begins with the isomerization of β-carotene from a trans to a cis configuration by β-carotene isomerase (*DWARF27*, *D27*). 9-cis-β-carotene is then cleaved by two carotenoid cleavage dioxygenases, *CAROTENOID CLEAVAGE DIOXYGENASE 7* (*CCD7*) and *CAROTENOID CLEAVAGE DIOXYGENASE 8* (*CCD8*), resulting in the formation of carlactone (Alder et al., 2012; Seto et al., 2014). Mutants of *CCD7* and *CCD8* have undetectable or extremely reduced exudation of strigolactones (tomato, Sl*CCD8* (Kohlen et al., 2012); sorghum, Sb*CCD8b* (Hao et al., 2023); rice, Os*CCD7* (Butt et al., 2018)), demonstrating that *CCD*s function as critical enzymes for the biosynthesis of strigolactones.

Mutants of *CCD7* and *CCD8* homologs are not only deficient in strigolactone exudation, but also have altered shoot tillering and branching (*Arabidopsis* (Sorefan et al., 2003); rice (Arite et al., 2007); tobacco (Gao et al., 2018); tomato (Vogel et al., 2010); maize (Guan et al., 2012); sorghum (Hao et al., 2023)) and altered root system architecture (rice (Arite et al., 2012); tomato (Kohlen et al., 2012); *Arabidopsis* and pea (Rasmussen et al., 2012); maize (Guan et al., 2012); sorghum (Hao et al., 2023)). More recently, strigolactone signaling has been linked to variation in root anatomy. For example, (Zhao et al., 2025) showed that strigolactones negatively regulate cambium-derived xylem vessel formation in *Arabidopsis*, while (Shrestha et al., 2026) demonstrated strigolactone-deficient Os*PSY2* mutants in rice to have impaired aerenchyma formation that can be rescued by exogenous strigolactone application. Beyond modulating plant architecture and morphology, strigolactones also play a role in shaping physiological traits and have been linked to differences in stomatal development and behavior (Van Ha et al., 2014), chlorophyll content and the efficiency of light capture through transcriptional regulation (Mayzlish-Gati et al., 2010). Transcriptome analysis of strigolactone deficient tomato mutants (Sl*ORT1*) revealed decreased expression of genes involved in light harvesting which can be rescued with exogenous treatment of the synthetic strigolactone analog GR24 (Mayzlish-Gati et al., 2010), suggesting that strigolactones are positive regulators of light harvesting processes. Altogether, the number of plant phenotypes regulated by or associated with differences in strigolactone profiles establishes their vital role in plant response to the environment and downstream impacts on numerous complex traits.

Along with other plant hormones, strigolactones play a key role in signaling pathways
that integrate nutrient and environmental cues to form beneficial symbioses. For example, plants
have evolved the phosphate starvation response (PSR) which reprograms plant processes to
acquire more phosphorus from the environment and to remobilize phosphorus under nutrient-
deprived conditions (Zhang et al., 2014; Chien et al., 2018). PSR results in an increase of
synthesis and exudation of strigolactones (Wang et al., 2021; Haider et al., 2023). This increase
in strigolactones provides two crucial functions: first to serve as endogenous hormones that
control plant development and stress response, and secondly, to serve as exogenous signaling
molecules and promote symbiosis with beneficial arbuscular mycorrhizal fungi (AMF). AMF are
obligate heterotrophs that interact with the roots of more than 80 percent of land plants, including
crop plants, and provide difficult to access nutrients in return for photosynthetically fixed carbon
and lipids (Brundrett and Tedersoo, 2018). AMF colonizes the host root cortex before
differentiating into highly branched structures called arbuscules, the main site of nutrient
exchange between the fungal and plant symbiotic partners. Strigolactones play a key role in the
initiation of the symbiosis by inducing AMF hyphal branching, enhancing the possibility of
contact between AMF and a host root (Giovannetti et al., 1993; Giovannetti et al., 1994; Kodama
et al., 2022).
123
Host plant exudation of strigolactones to recruit AMF symbionts has been hijacked by
parasitic plant members of the Orobanchaceae family. Obligate root parasites in this family
germinate after sensing host root exuded strigolactones (Yoneyama et al., 2010) and parasites in
this family generally use host strigolactones as chemoattractants (Ogawa et al., 2022). Of
particular interest is the obligate root parasite *Striga hermonthica,* which accounts for billions of
dollars' worth of annual cereal crop damage in tropical Africa and is considered one of the
greatest constraints to African food security (Spallek et al., 2013). One of the best characterized
host resistance mechanisms is low germination of *S. hermonthica* by the host's root exudates
(Gurney et al., 2003; Ruyter-Spira et al., 2011; Mallu et al., 2021). In sorghum, naturally
occurring deletion mutants of the *LOW GERMINATION STIMULANT 1* (*LGS1*) quantitative trait
locus (QTL) confer shifts in strigolactone stereochemistry, where orobanchol-type rather than 5-
deoxystrigol (5DS) becomes the more abundant strigolactone in root exudates (Gobena et al.,
2017). The gene causing the low germination stimulation phenotype (*Sobic.005213600, LGS1*)
encodes a sulfotransferase that functions downstream of CYP711A (Sb*MAX1a*) converting 18-
hydroxy-carlactonoic acid (18-hydroxy-CLA) into 5DS and 4-deoxyorobanchol (4DO)
simultaneously (Wu and Li, 2021; Yoda et al., 2021). In addition to altering the strigolactone
composition, loss-of-function at the *LGS1* gene may reduce the quantity of total strigolactone
exudation (Bellis et al., 2020). Directly upstream of the *LGS1* gene is *Sobic.005G213500*
(*Sb3500*), a 2-oxoglutarate-dependent dioxygenase involved in stereoselective strigolactone
production and increases the quantity of 5DS relative to 4DO when expressed with *LGS1* (Yoda
et al., 2023). Previously described naturally occurring deletion variants at the QTL either lack
both *LGS1* and *Sb3500* (e.g. *lgs1*-1, *lgs1*-2), or retain *Sb3500* with absence of *LGS1* (e.g. *lgs1*-3)
(Gobena et al., 2017). Thus, the genomic region where *LGS1* resides may function as a multi-
gene module where epistasis influences strigolactone profiles, though much of the work to date
on *LGS1*-based resistance in sorghum has focused on the gene encoding the sulfotransferase
(*LGS1, Sobic.005213600*).
150

Despite the potential for loss-of-function in the *LGS1* gene to increase sorghum's fitness
in landscapes with parasite pressure, the alterations to strigolactone biosynthesis may impair
endogenous processes in sorghum. Supporting this hypothesis, Bellis et al., (2020) found
evidence of potential trade-offs: natural loss-of-function *lgs1* alleles are maintained at a modest
frequency in sorghum landraces within *S. hermonthica*-prone regions (up to ~30% in northern
Nigeria) and are at very low frequency in *S. hermonthica*-free landscapes. Also, CRISPR-Cas9
deletions of *LGS1* altered expression of >2,000 genes compared to the wild type, and there was
a significant downregulation of light harvesting genes in shoots (Bellis et al., 2020). Given the
broad roles strigolactones play in plant processes, we hypothesized that sorghum with loss-of-
function at *LGS1* is subject to trade-offs through pleiotropic impacts.
161
In the present study, we explore the impact of mutations at two strigolactone biosynthesis
genes, Sb*CCD8b* and *LGS1,* on sorghum physiology, growth, and biotic interactions with
parasitic plants and arbuscular mycorrhizal fungi to ask the following questions: (Q1) How does
variation at strigolactone biosynthesis genes in sorghum impact biotic interactions with *Striga*
hermonthica and arbuscular mycorrhizae fungi? (Q2) What regulatory pathways may be altered
in mutants of strigolactone biosynthesis genes and are phenotypic changes consistent with altered
pathways? (Q3) How does variation at strigolactone biosynthesis genes in sorghum affect
sorghum performance and physiology? (Q4) Are strigolactone impacts consistent across diverse
genetic backgrounds?
171
172　**Materials & Methods**
173
174　*Generation of mutants*
175
We studied loss-of-function in the genes Sb*CCD8* and *LGS1* using gene-edited *S. bicolor*
of two different genetic backgrounds (Table 1). CRISPR-Cas9 edited lines of Sb*CCD8b* in the
'RTx430' background were included in our study to represent a strigolactone deficient line
(RTx430 *ccd8b*, see greater detail in Hao et al., 2023). Native sorghum ecotypes have natural
variation at the *LGS1* locus*,* impacting its functionality, and we included genetic two
backgrounds, Macia and RTx430, to determine if *LGS1* effects are consistent. Functional *LGS1*
was knocked out in the normally *LGS1*-functional background Macia (PI 565121) using
CRISPR-Cas9 to produce independent genome-edited lines (Macia *lgs1*-d, see greater detail for
generation of S_2 plants in Bellis et al. 2020). Unless otherwise indicated, we used a more
advanced version of the material described in Bellis et al. (2020) which was backcrossed to the
recurrent background Macia and selfed for an additional generation (BC_0F_2) in this study (Figure
S1, Supplementary Methods).
188
We additionally sought to test the effect of inserting functional *LGS1* alone back into a
normally non-functional background. RTx430 (PI 655996) is an inbred background, developed
at Texas A&M using a partially converted Ethiopian sorghum variety (Miller, 1984). The
RTx430 background carries the *lgs1-2* allele, which contains a large deletion that spans three
predicted genes, including the sulfotransferase *LGS1* and the 2-oxoglutarate-dependent
dioxygenase *Sb3500* (Gobena et al., 2017). *LGS1* was inserted via transformation with
Agrobacterium into the RTx430 genetic background, to produce two independent mutants with
restored functionality of the locus (RTx430 *LGS1*) (Supplementary Methods). For both *LGS1*

insertion (RTx430) and deletion events (Macia), null event mutants were also produced and
included in our analysis as "wild-type like" (WT-L) comparisons. See Table 1 for descriptions of
the plant material used in this study.
200
201
202 **Table 1 | Description of plant material used in this study. All gene models are reported from the BTx623v5.1**
203 **reference genome.**
204

Genetic Background	Genotype	Description
RTx430	RTx430 *ccd8b*	CRISPR/Cas9-edited deletion mutant of the Sb*CCD8b* (*Sobic.003G293600*) gene described in Hao et al., (2023).
	RTx430 WT (*lgs1*-2)	Wild type with the *lgs1*-2 natural loss-of-function allele. The *lgs1*-2 deletion allele spans the three gene models *Sobic.005G213400*, *Sb3500* (*Sobic.005G213500*), and *LGS1* (S*obic.005G213600*).
	RTx430 WT-L (*lgs1*-2)	Insertion transformation event null segregant.
	RTx430 *LGS1*	Mutants with functional *LGS1* (*Sobic.005G213600*) introduced via *Agrobacterium* random transformation, described in this study.
Macia	Macia WT	Wild type with intact *LGS1* (S*obic.005G213600*).
	Macia WT-L	Deletion transformation event null segregant.
	Macia *lgs1*-d	CRISPR/Cas9-edited deletion mutant of the gene *Sobic.005G213600* (*LGS1*). Material grown in this study includes the S_2 generation (described in Bellis et al., (2020)) and a more advanced generation backcrossed to the Macia wild type to generate BC_0F_2 seed (Figure S1).

205
206
207
208 *Evaluation of allelic diversity in natural sorghum populations*
209
Because Yoda et al. (2023) found the effect of presence-absence of *LGS1*
(*Sobic.005213600*) and *Sb3500* (*Sobic.005213500*) had interactive effects on strigolactones
when transiently expressed in *Nicotiana benthamiana*, we evaluated patterns of presence-
absence variation and linkage of these genes in sorghum landraces (Table 2)(Morris et al., 2026).
Gene presence was determined using a local exon-level *de novo* reassembly pipeline, in which
sequencing reads were assembled without mapping to a primary reference. Exon contigs
including short flanking regions were stitched according to the BTx623v5.1 (Morris et al., 2026)
exon order to reconstruct pseudo-mRNA sequences. Genes with reassembled proteins showing
≥90% identity and ≥90% coverage relative to BTx623v5.1 were classified as present.
Strigolactone profiles reported in Table 2 were not directly measured, but inferred based on
published functional annotations of *LGS1* and *Sb3500* and their roles in strigolactone
biosynthesis (Gobena et al., 2017; Yoda et al., 2021; Yoda et al., 2023).
222

Striga hermonthica *germination assay*

Root exudates of the mutants with functional *LGS1* inserted, "wild type-like" (WT-L)
null segregants, and the WT in the RTx430 background were collected to test their *S.*
hermonthica germination activity. Plants were grown in a greenhouse in University Park, PA
(16-hour photoperiod; day temperature 28°C; night temperature 22°C). Sorghum seeds were
surface sterilized in 1.2% sodium hypochlorite for 10 minutes, rinsed with sterile deionized
water (diH$_2$O) and germinated in autoclaved vermiculite. One-week old seedlings were
transferred to a hydroponics system comprising 50 mL falcon tubes filled with half-strength
modified Hoagland solution. Each seedling was suspended in the solution by a bottomless 1.5
mL centrifuge tube held at the center of the 50 mL falcon tube cap (Wang et al., 2022). Plants
were grown in this medium for seven days and tubes were manually agitated every other day to
improve airflow. The medium was discarded, then tubes and seedling roots rinsed with diH$_2$O.
Tubes were refilled with half-strength Hoagland's solution without phosphate, and seedlings
grown for an additional 7 days. Root exudates were collected in 50 mL diH$_2$O over a 48-hr
period under darkness. Each seedling's root volume was recorded by water displacement in a
graduated cylinder, and exudate concentrations standardized to the lowest recorded root volume
prior to use for induction of *S. hermonthica* germination.

Germination assays used *S. hermonthica* seed collected in 2016 from a sorghum field in
Kibos, Kenya (0°40′S; 34°49′E) and in 2018 from a sorghum field in Siby, Mali (12°23′N;
8°20′W). Since *S. hermonthica* requires conditioning to be responsive to germination stimulants
(Matusova et al., 2004), parasite seeds were surface sterilized in 1.2% sodium hypochlorite with
a drop of Tween 20 for 20 mins, rinsed thoroughly with sterile diH$_2$O, and incubated at 30°C in
the dark for 10 days. After conditioning, the *S. hermonthica* seed-diH$_2$O mixture was
homogenized before pipetting 100 μL (approximately 100 seeds) into a single well of a 12-well
microtiter plate (CytoOne). Excess water was aspirated out of the wells before adding 3 mL of
sorghum root exudates. 15 biological replicates of each sorghum genotype, and 2 technical reps
(wells per biological rep) were tested for germinability of each *S. hermonthica* population. A
similar volume of 0.1ppm GR24 (synthetic strigolactone standard) and diH$_2$O were used as
positive and negative controls, respectively. *S. hermonthica* germination, characterized by
emergence of a radicle, was checked after incubating the plates for 48 hours at 30°C in the dark,
and quantified by calculating the percentage of germinated seeds in each well. Germination
percentage of *S. hermonthica* in Figure 1 is plotted with a broken axis using functions described
in R/ggbreak (Xu et al., 2021).

Identifying potential transcription factor binding sites related to differential gene expression in
lgs1-*d plants*

Using previously generated transcriptome data from Bellis et al. 2020, we identified
impacted regulatory pathways in Macia *lgs1*-d plants (S$_2$ generation). We used samtools v1.10
(Li et al., 2009) to extract the flanking genomic sequence of differentially expressed genes in
roots and shoots separately. Specifically, we extracted the sequence 500 bp upstream of the
gene's start codon and the sequence 500 bp downstream of the gene's stop codon, to capture
differences in upstream and downstream regulatory coding regions. We then used STREME
(Bailey, 2021) to determine significantly enriched motifs within the regulatory coding regions of

differentially expressed genes compared to non-differentially expressed genes within each
specific tissue (roots and shoots). We searched for overlap between enriched motifs and the
JASPAR 2020 CORE (Anon) plant transcription factor binding sites database (Sandelin et al.,
2004) with MEME suite's online tool TOMTOM (Gupta et al., 2007).

Growth conditions for mycorrhizal assay
We tested for differences in *LGS1* and Sb*CCD8b* functional and non-functional
genotypes in the two genetic backgrounds (Maica, RTx430) under variable phosphate levels and
grown with and without mycorrhizae. Plants were grown in a greenhouse in University Park, PA
(16-hour photoperiod; day temperature minimum 21°C and maximum 28°C; night temperature
minimum 21°C and maximum 26°C; relative humidity 20%) and subjected to three treatments:
high phosphate without mycorrhizae (P+ AMF-), low phosphate without mycorrhizae (P- AMF-
), and low phosphate with mycorrhizae (P- AMF+). The plants were grown in 10 inch "deepots"
(Stuewe & Sons Inc., https://stuewe.com/product/2-5-x-10-heavyweight-deepot-cell/) in a mix of
90% sterile silica sand (Quickrete) and 10% kaolin clay (CB Minerals,
https://cbminerals.com/kaolin/). After sowing, pots were watered with tap water every day until
seven days after plant emergence. One-week post-emergence, plants were watered every other
day with 200 mL water and treated with a supplemental fertigation of 1/3 Hoagland with
different phosphate concentrations every other watering (every 4 days) until time of harvest. The
phosphate level of the nutrient solution used at each fertigation was 333 μM for the high
phosphate treatment (P+) or 33 μM for the low phosphate treatments (P-). Pots with a
mycorrhizal treatment (AMF+) were inoculated with 50 mL of a sandy mix containing ~20
spores/mL of *Rhizophagus intraradices* prior to planting. The spores used for inoculation were
bulked in University Park, PA two months before the greenhouse experiment. Plants were
harvested and destructive measurements were collected at two timepoints, 21 and 42 days after
plant emergence (*n* = 7 per genotype, treatment, and time point; total *n* = 462).
The non-destructive measurements of light-saturated net carbon assimilation rate (steady
state, measured via LI-6400), chlorophyll fluorescence and stomatal conductance (measured via
LI-600 fluorometer/porometer), and shoot traits (plant height, vegetative phase change markers)
were measured on greenhouse-grown plants prior to harvest dates. At the time of harvest (21 and
42 days) several destructive measurements were taken, and root tissue was collected for
quantification of mycorrhizal colonization. Above ground tissue was collected and dried in an
oven at 70°C for 72 hours for collection of dry biomass. As we could not measure root dry
weight for plants grown in mycorrhizal treatments, fresh weight of the entire root system was
measured. For the low phosphate non-mycorrhizal and mycorrhizal treatments, root systems
were divided into biopsy cassettes (Fisher Scientific) for staining to quantify mycorrhizal
colonization and confirmation of uncontaminated negative controls.

Mycorrhizae counting assay
To test for differences in colonization of the fungal species *R. intraradices*, root samples
of the mycorrhizal treatments stored in biopsy cassettes from both harvest time points (21 and 42
days) were stained (https://invam.ku.edu/staining-roots) and measured for relative mycorrhizal
colonization (https://www2.dijon.inrae.fr/mychintec/Protocole/protoframe.html) (Phillips and
Hayman, 1970; Koske and Gemma, 1989; McGONIGLE et al., 1990). At the time of harvest,
cassettes were packed with ~0.1 grams of root system and placed in 75% EtoH until time of
staining. Roots were cleared using 10% KOH that was brought to boiling and then removed from

the heat prior to adding cassettes. Cassettes containing root samples were left in the hot KOH
solution for 10 minutes. Samples were then washed five times using diH_2O and acidified by
being immersed in a 2% HCl solution for 20 minutes at room temperature. After acidification,
roots were stained from incubation in a mixture of 1:1:1 water, glycerin, and lactic acid
containing 0.05% direct blue at room temperature for at least 12 hours. Stained samples were
again rinsed with diH_2O five times and stored suspended in diH_2O in a fridge at 15°C for one
week to allow for excess leaching of the stain prior to quantification of mycorrhizae.
Relative mycorrhizal colonization was quantified using a modified version of the grid
line intersect method (Newman, 1966; Tennant, 1975; Giovannetti and Mosse, 1980). Stained
roots were randomly dispersed in a nine-cm diameter petri plate with an acetate sheet containing
10x10 grid lines under the petri plate. Using an Amscope SM-2, intersections where a root
crossed a grid line were designated as either non-mycorrhizal or colonized. The presence of any
fungal structure (intracellular hyphae, vesicle, arbuscule, spore) resulted in the intersection being
scored as "colonized". To assess the relative abundance of mycorrhizae in each sample, 100
intersections were scored, and relative colonization was calculated as a percentage.
330
331 *Collection of root phenotypes*
332
Parameters of root system architecture and anatomy were collected for plants grown in
the high phosphate no mycorrhizae treatment at the 42 days after plant emergence time point. To
characterize architecture, entire root systems were scanned on an Epson scanner and processed in
RhizoVision Explorer v2.0.3 (Seethepalli and York, 2020; Seethepalli et al., 2021) using
algorithms described by (Seethepalli et al., 2021). For root anatomy, we collected two
representative axial roots from node three and excised two four-cm root samples five to nine cm
from the most basal portion of the whole root sample. Root samples were stored in 75% ethanol
until being sectioned by laser ablation tomography (LAT; (Strock et al., 2019)). In LAT, a
sample is moved via an automatic stage towards a 355-nm Avia 7000 pulsed laser and ablated in
the focal plane of a camera. A Canon T3i camera with a 53micro lens (MP-E 65 mm) was used
to capture images of the root cross-section. Two representative images for each root sample
sectioned one to three cm apart and were saved for later image analysis with Rootscan
(https://plantscience.psu.edu/research/labs/roots/methods/computer/rootscan) (Burton et al.,
2012). For each plant, anatomical phenotypes were averaged across images collected for each
nodal root and reported for the two sampled roots.
348
349 *Photosynthesis, stomatal conductance, and density of stomata*
350
All steady state photosynthesis measurements were measured from plants grown without
nutrient limitation using a Li☐6400 portable photosynthesis machine (Li☐Cor Environmental).
Prior to collecting gas exchange measurements, leaves were acclimated to the starting chamber
conditions and the first data point was logged after CO_2 and transpiration rates reached steady
state. For all measurements, a constant leaf temperature of 25°C, light level of 1800 µmol m^{-2} s
$^{-1}$, and a flow rate of 400 µmol s^{-1} were maintained. Light saturated photosynthetic rate (A_{sat})
was measured at a reference CO_2 level of 400 ppm. A_{sat} was recorded every 20 seconds for two
minutes and reported as the average across all data points logged.
359

360   The Li-600 Porometer/Fluorometer machine (Li□Cor Environmental) was also used to
361   measure stomatal conductance and parameters of dark (leaves were kept in the dark for a
362   minimum of 30 minutes) and light-adapted chlorophyll fluorescence. For all samples, light and
363   dark-adapted measurements were collected on the same leaves with the fixed flow rate set to
364   high (150 µmol s^{-1}) and a flash intensity of 6000 µmol m^{-2} s^{-1}.
365

We further collected CO_2 response curves, steady state photosynthesis, and ran
untargeted metabolomics for the previous S_2 generation of Macia *lgs1*-d CRISPR deletion
mutants using plant material as described in (Bellis et al., 2020) in a separate greenhouse
experiment. Here, we also collected steady state photosynthesis for RTx430 *ccd8b*. Plants were
grown in a media mix that consisted of 80% commercial sand (Quickrete) and 20% turface
(LESCO turface, https://www.siteone.com/en/088782-lesco-turface-all-sport-soil-conditioner-
infields-50-lb/p/23795) for 16-hour days. After germination, plants were watered with 100 mL of
water every other day and supplementally fertigated with 100 mL half strength Hoagland once a
week starting at 7 days after plant emergence. CO_2 response curves were built by measuring net
photosynthetic rate (A_{net}) at the reference CO_2 levels of 50, 100, 200, 300, 400, 600, 800, 1000
ppm. Across all CO_2 levels, a constant leaf temperature of 25°C, light level of 1800 µmol m^{-2} s
$^{-1}$, and a flow rate of 400 µmol s^{-1} were maintained. Leaves were given a chance to acclimate to
the chamber CO_2 conditions for at least two minutes before data logging at each CO_2 level. We
used methods described in (Zhou et al., 2019) to fit A_{net} □ C_i curves and to determine apparent
V_{cmax} and apparent J_{max} using procedures of (Sharkey et al., 2007), accounting for carbonic
anhydrase activity.
382

Leaves used in gas exchange measurements were also analysed to determine stomatal
density. Here, we collected three non□overlapping fields of view from comparable regions on
the abaxial and adaxial surface. Images were captured using an AmScope MU1400 digital
camera mounted on a compound microscope equipped with a 20×/0.25 finite objective (F-
LD□20/0.25□160/2) and the camera was connected through a 0.5× C□mount adapter, resulting
in an effective magnification of approximately 10×. Number and area of stomata were quantified
using ImageJ (Schneider et al., 2012). All area measurements were converted from pixel units to
micrometers using the calibrated pixel size ($\approx$□0.14□µm□px^{-1}).
391

Metabolomics

To quantify differences in abundance of small molecules for sorghum mutants of
Sb*CCD8b* and *LGS1*, root and shoot tissue samples were taken for $n = 5$ plants of the genotypes
RTx430 WT, RTx430 *ccd8b*, Macia WT, and the S_2 generation of Macia *lgs1*-d deletion lines
(Bellis et al. 2020) and for evaluation with untargeted metabolomics. Root and shoot samples
were immediately flash-frozen separately in liquid nitrogen and transferred to a 1.5 mL
Eppendorf tube for further processing. Samples were then lyophilized at -80°C overnight before
being ground to a fine powder by placing a metal ball in each Eppendorf tube and shaking in a
TissueLyser II (Qiagen) for 5 minutes at 25 Hz.

All solvents were chromatography grade and acquired from VWR (Radnor, PA, U.S.A.)
or Sigma Aldrich (St. Louis, MO, U.S.A.). Metabolites were extracted using 1 mL 80% MeOH +
0.1% formic acid for every 100 mg of powdered sample. Samples were vortexed for 10 seconds

and sonicated for five minutes. Then, each sample was centrifuged at 10,000 rpm for 10 minutes
and the supernatant was transferred to a 1 mL glass scintillation vial. The extraction process was
repeated with the same tissue and the supernatants were combined before being air-dried to
completion in a fume hood. All samples were reconstituted at a concentration of 1 mg/mL in LC-
MS grade methanol with 1 uM chlorpropamide as an internal standard. A Vanquish Duo UHPLC
system connected to a Thermo Fusion Lumos Orbitrap Mass Spectrometer (Thermo Fisher
Scientific, Waltham, MA) was used for metabolomics analysis with a Waters Acquity UPLC
BEH C18 (21.x150 mm x 1.7 um) column. The flow rate was 0.1 mL/min at 55°C. Solvent A
was 0.1% formic acid (v/v) in LC-MS water and solvent B was 0.1% formic acid (v/v) in LC-MS
acetonitrile. The mobile phase gradient of solvent B was: 3% for 0.01 minutes, 45% for 10
minutes, 75% for 2 minutes, 100% for 4.5 minutes, and 3% for 0.2 minutes. A 2 ul injection was
used for all samples.

Mass spectrometry was conducted using an electrospray ionization source with a positive
ion spray voltage of 3500 V, sheath gas pressure of 25 Arb, auxiliary gas pressure of 5 Arb, ion
transfer temperature of 275°C, and vaporizer temperature of 75°C. MS1 data was acquired with
an Orbitrap resolution of 120,000, scan range of 100-1000 Da, and RF lens of 50% in the profile
mode. MS2 data was collected in a data-dependent manner using intensity (2.5E4) and dynamic
exclusion (ions excluded after 1 detection for 30 seconds).
423
All .raw mass spectral files were converted to .mzML using MSconvert (Proteowizard)
and then loaded into MzMine3.1 (Schmid et al., 2023). Root and shoot files were analyzed
separately. Supplemental Table 5 outlines the MZMine3 processing parameters. Following
processing in MzMine3.1, all features with a peak area less than 5-fold the intensity of the
metabolite concentration in the blanks were removed. The peak area was Hellinger transformed
and auto-scaled. To make volcano plots, the compounds' P-values and $\log_2$fold changes were
calculated between each mutant and its corresponding wild type. Compounds with a $\log_2$fold
change greater than 0.6 and a P-value less than 0.05 were labeled as "up-regulated" and those
with a $\log_2$fold change less than 0.6 and a P-value less than 0.05 were labeled as "down-
regulated." Key compounds were annotated by comparing their MS2 profiles to potential
matches found via molecular networking performed using the Global Natural Products Social
Molecular Networking (GNPS) library search feature (Wang et al., 2016). To compare statistical
differences between mutants and their respective wild type lines, a Permutational Multivariate
Analysis of Variance (PerMANOVA) was performed using the R/vegan (Oksanen et al., 2022).

Quantification of chlorophyll content and light harvesting proteins
440
We assessed if chlorophyll content (Chl*a*/Chl*b*), and concentration of photosystem
proteins (ratio of PSAA/CP43 and LHCB1/CP43) were altered in sorghum genotypes
segregating for functional Sb*CCD8b* and *LGS1*. Plants used in this quantification were watered
every day for the first week and then every other day until time of harvest. At every other
watering, plants were supplementally fertigated with 1/3 Hoagland solution. At the time of
harvest, five-week-old plants were flash frozen in liquid nitrogen. Chlorophylls were extracted
from leaf tissue and thylakoid samples in acetone 80% and the absorption spectra were analyzed
using the Cary and fitted as previously described (Croce et al., 2002).

Total proteins were extracted by grinding the frozen leaf tissue in a mortar with the
BufferA/4 from (Schägger, 2006) (3% SDS (wt/vol), 1.5% mercaptoethanol (vol/vol), 7.5%
glycerol (wt/vol), 0.012% CoomassieblueG-250 (Serva), 37.5mMTris/HCl (pH7.0))
supplemented with protease inhibitor (cOmplete™ Protease Inhibitor Cocktail (Roche)).
Samples were then centrifugated and the supernatant was stored at -20°C. Immunoblotting was
performed as described in (Schägger, 2006). All primary antibodies, PsaA(AS06172),
PsbC(AS111787), and LHCB1(AS01004), were purchased from Agrisera, Sweden.
Chemilluminescence was collected by a LAS 4000 Image Analyzer and analyzed with the
software ImageJ.
458
Statistical analysis

For statistical analysis of the impact of exudate on *Striga* germination, we built negative
binomial generalized linear models using R/MASS::glm.nb (Venables and Ripley, 2002). Nested
models accounted for transformation event and *S. hermonthica* ecotype as effects, with one
model additionally including an allele term. The two models were compared using a likelihood
ratio test and significance of the *LGS1* allele effect was assessed using a chi-square distribution.
465
For all other collected phenotypes, we tested for the significance of *LGS1* deletion (*lgs1-*
d) and *LGS1* insertion alleles by fitting linear mixed effects models using R/lme4::lmer (Doran et
al., 2007). Full models included the fixed effect of allele and the random effect of transformation
event and a reduced model excluded the fixed effect allele term. To determine if the inclusion of
the allele term improved model fit, the full and reduced models were assessed with a likelihood
ratio test implemented with the R 'anova' function and significance was determined based on the
resulting chi-square statistic and associated *P*-value. Separate models were built for plants of the
Macia and RTx430 background. For Sb*CCD8b* deletions, we used a student's *t*-test to assess
significance of the deletion allele. R statistical software (R Core Team, 2022) was used
throughout for data processing and statistical analysis.
476
477 **Results**
478
479 *Presence-absence variation of* LGS1 *and* Sb3500 *in natural populations and predicted*
480 *strigolactone profiles*
481
Previous studies demonstrate substantial structural variation across the *LGS1* genomic
region in sorghum (Gobena et al., 2017; Bellis et al., 2020; Maina et al., 2025). To contextualize
our experimental genotypes, we assessed presence-absence variation of two genes that impact
strigolactone profiles, *Sb3500* (*Sobic.005213500*) and *LGS1* (*Sobic.005213600*), in both our
experimental lines and natural populations, and used published functional data to infer expected
strigolactone profile (Table 2).

494 **Table 2 | For material of the *LGS1* gene used in this study, summary of gene presence-absence at *Sb3500* and**
495 ***LGS1*, observations of the allele combinations in landraces, predicted strigolactone profiles, and observed**
496 **effects of synthetic mutants on symbiotic interactions.**
497

	Genetic Background			
	Macia		RTx430	
	Wild Type (WT / WT-L)	Deletion Mutant	Wild Type (WT / WT-L)	Insertion Mutant
Gene presence-absence				
Sb3500 (*Sobic.005213500*)	Present	Present	Absent	Absent
LGS1 (*Sobic.005213600*)	Present	Absent. Deleted with CRISPR-Cas9 to generate *lgs1*-d mutant.	Absent	Present. Reintroduced via agrobacterium random transformation to generate *LGS1* insertion mutant.
Observations of allele combinations (presence-absence at *Sb3500* and *LGS1*) in *Sorghum biocolor* landraces	*N* = 668 Most common combination of alleles observed in Sorghum landraces. Predominantly georeferenced to localities less suitable for *Striga hermonthica* (Bellis et al. 2020).	*N* = 15 Combination of alleles observed in the natural deletion variant *lgs1*-3 (Gobena et al. 2017).	*N* = 20 Combination of alleles observed in the natural deletion variant *lgs1*-2 (Gobena et al. 2017).	*N* = 1
Predicted strigolactone profile	High 5DS, low orobanchol (Gobena et al. 2017). Higher production of 5DS (relative to 4DO) when *LGS1* and *Sb3500* are co-expressed (Yoda et al. 2023).	High orobanchol, low 5DS (Gobena et al. 2017). Hypothesized lower overall strigolactone exuded (Bellis et al. 2020).	Orobanchol is the predominant strigolactone in RTx430 (Yoda et al. 2021). Hypothesized lower overall strigolactone exuded (this study).	4DO and 5DS in approximately equal amounts when *LGS1* is expressed without *Sb3500* (Yoda et al. 2023).
Observed impact to symbioses *Striga hermonthica* germination		Reduced germination of *Striga* when treated with root exudates from Macia *LGS1* deletion (*lgs1*-d) mutants (Bellis et al 2020).		Germination of *Striga* rescued when treated with root exudates from RTx430 *LGS1* insertion mutants (this study).
Arbuscular mycorrhizal fungi (AMF) colonization		Reduced colonization of the AMF species *R. intraradices* at earlier timepoint for Macia *LGS1* deletion (*lgs1*-d) mutants, difference not significant at later timepoints (this study).		No difference in colonization of the AMF species *R. intraradices* at either time point for RTx430 *LGS1* insertion mutants (this study).

498
Genotypes with putatively functional copies of both genes (e.g. Macia WT) were the
most common in sorghum landraces (*N* = 668) and are expected to produce predominantly 5DS
type strigolactone (Gobena et al., 2017; Yoda et al., 2023). In contrast, genotypes with *LGS1*
deletion alleles and presence of *Sb3500* (e.g. Macia *lgs1*-d, *lgs1*-3) are expected to primarily

exude orobanchol (Gobena et al., 2017; Bellis et al., 2020) and occur in sorghum landraces at
low frequency (N = 15). The RTx430 WT carries the natural *lgs1*-2 deletion allele which spans
both *Sb3500* and *LGS1*. This absence at both these genes also occurs at a low frequency in
sorghum landraces (N = 20), and orobanchol is the expected predominant strigolactone in
exudate (Gobena et al., 2017; Yoda et al., 2021). Importantly, the RTx430 *LGS1* insertion mutant
represents a distinct and extremely rare haplotype (N = 1) in which *LGS1* is present in the
absence of *Sb3500*. Transient expression of *LGS1* in *E. coli* incubated with a strigolactone
precursor produced 5DS and 4DO in approximately equal amounts, while co-expression with
Sb3500 resulted in the stereoselective biosynthesis of 5DS in *Nicotiana benthamiana* (Yoda et
al., 2023). As such, we hypothesize that the RTx430 *LGS1* insertion genotype, absent of
functional *Sb3500*, produces an intermediate strigolactone profile of 5DS and 4DO and is distinct
from the strigolactone profile of RTx430 WT (orobanchol), and the vast majority of genotypes
with the *LGS1* gene present (high 5DS) (Mohemed et al., 2018).
516
517 *Inserting functional* LGS1 *into a naturally loss-of-function background restores* Striga
518 hermonthica *germination*
519
We sought to confirm that the new mutants with functional *LGS1* inserted into the
RTx430 background changed strigolactone profiles in a manner consistent with the deletion
effects (Q1). We tested the germinability of two *S. hermonthica* ecotypes known to be
differentially sensitive to (+) 5DS and (±) orobanchol (Bellis et al., 2020)(Figure 1C). Root
exudate collected from the RTx430 *LGS1* insertion mutants induced significantly more
germination of both *S. hermonthica* ecotypes as compared to RTx430 WT and WT-L null
segregants with the *lgs1*-2 natural deletion allele (χ^2 = 53.95, P < 0.001, likelihood ratio test for
fixed effect of *LGS1* deletion; Figure 1A-B). The magnitude of RTx430 *LGS1* effect on *S.*
hermonthica germinability differed between the two ecotypes, supporting that there is genetic
variation in parasite sensitivity to host-derived strigolactone profiles of root exudates. For the *S.*
hermonthica ecotype from Siby, Mali, average germination was <0.02% when treated with
exudate from RTx430 WT and WT-L (*lgs1*-2) plants and 6.5% when treated with exudate from
the RTx430 *LGS1* insertion mutant plants (Figure 1A). The *S. hermonthica* ecotype from Kibos,
Kenya had slightly higher germinability, and an average germination rate of <1% when treated
with RTx430 WT and WT-L (*lgs1*-2) exudate and 10.56% when treated with exudate from
RTx430 *LGS1* (Figure 1B). Variation in parasite sensitivity to strigolactone profiles for the same
two ecotypes was also reported by Bellis et al. (2020), in which Macia *lgs1*-d mutants had lower
germinability of the parasite compared to Macia WT, with this trend being weaker for the
Kenyan ecotype. Notably, this Kenyan *S. hermonthica* ecotype has a similar sensitivity to (+)
5DS and (±) orobanchol (Figure 1C) (Bellis et al., 2020). Here, we observe little to no
germination for both *S. hermonthica* ecotypes when exposed to exudates from RTx430 WT and
WT-L plants, suggesting that the *lgs1*-2 natural deletion allele may not only shift strigolactone
profiles in matter that does not strongly induce germination (i.e. orobanchol), but also reduce the
overall strigolactone in root exudate, as exemplified by low germination in the Kenyan
population. Together, insertion of functional *LGS1* into a natural loss-of-function background,
RTx430 (*lgs1*-2), changes strigolactone profiles in a manner that increased the germination of *S.*
hermonthica likely due to an increase of 5DS and a larger quantity of strigolactone in root
exudate.
548

Figure 1 / Restoring functional *LGS1* in the RTx430 background rescues *Striga hermonthica* germination.
Percent *S. hermonthica* germination of ecotypes sourced from A) Siby, Mali and B) Kibos, Kenya when exposed to
root exudate from RTx430 WT (*lgs1*-2), "wild type-like" null segregants (RTx430 L1,2 WT-L), insertion mutants of
the *LGS1* gene (RTx430 L1,2 *LGS1*), a synthetic strigolactone analog (GR24), and water. L1,2 indicate mutant and
null segregant pairs that originate from the same transformation event. Boxplots colored by *LGS1* allele of root
exudate used as germination stimulants or controls. Asterisks denote statistically significant differences from the
RTx430 WT by Student's *t*-test. C) Source location (large white dots) of the two *S. hermonthica* ecotypes used in
germination experiments. Denoted is each ecotype's response to synthetic strigolactones, (+) 5-deoxystrigol (5DS)
or (±) orobanchol, as reported in Bellis et al. 2020. **$P<0.01$, ****$P<0.0001$ (*n*=15).

Pathways involved in gene regulatory effects of LGS1 deletion

   Bellis et al. (2020) reported that deleting functional *LGS1* from the otherwise functional
Macia background resulted in substantial changes in gene expression. In *LGS1* knockout plants
(Macia *lgs1*-d), 505 differentially expressed genes (DEGs) were identified in roots (244 down-
regulated, 261 up-regulated) and 2,167 DEGs were identified in shoots (917 down-regulated,

1,250 up-regulated). To identify regulatory pathways that relate to differences in gene expression
(Q2), we searched for sequence motifs in putative promoter regions of *lgs1*-d DEGs using
STREME (Bailey, 2021). Within putative promoter regions of Macia *lgs1*-d root DEGs, we
discovered three enriched sequences that overlapped with 16 known motifs, and within shoot
DEGs we found six enriched sequences that overlapped with 92 known motifs. We next sought
to match enriched motifs identified in putative promoter regions of DEGs to known transcription
factor binding site (TFBS) motifs. In roots, the most commonly discovered motif
(*GACCGTCCATC*) was not significantly enriched in DEGs ($P = 0.48$), indicating that it is
broadly prevalent across expressed root genes, rather than specific to differential expression.
This motif overlapped with members of six known TFBS families (Table S1), including
ethylene-responsive element-binding factors (ERFs) and Squamosa Promoter Binding Protein-
Like (SPL) genes (overlap of discovered motif with TFBS sequences, $P < 0.004$), which are
involved in abiotic stress response and lateral root development and nodulation, respectively
(Fujimoto et al., 2000; Yu et al., 2015; Nasrollahi et al., 2022). Together, the identified root
TFBS motifs reflect regulatory patterns among expressed genes in roots; however, these motifs
were not significantly associated with the regulatory variation underlying differential expression
in Maica *lgs1*-d roots. For shoots, the most common discovered motif (*CCGGCAGCG*) was
significantly enriched in 41 DEGs ($P = 0.002$, Table S2). Of known TFBS significantly
overlapping motifs enriched in promoters of shoot DEGs, the majority belonged to the
APETALA 2/ethylene-responsive element binding proteins (AP2/EREBP) family (overlap of
discovered motif with TFBS sequences, $P < 0.02$), which are key regulators of leaf and floral
development and participate in response to biotic and environmental stress (Riechmann and
Meyerowitz, 1998). AP2/ERFs regulate numerous target genes, suggesting that deleting *LGS1* in
a normally functional background may lead to broad pleiotropic effects, and stress and growth
responses may be particularly disrupted in Macia *lgs1*-d.
592
593   *Reduced CO$_2$ assimilation rate in strigolactone biosynthesis deletion mutants*
594
595   Strigolactones have been demonstrated to be positive regulators of light-harvesting genes
in tomato (Mayzlish-Gati et al., 2010) and Bellis et al., (2020) reported significant
downregulation of photosystem I, photosystem II, and light-harvesting genes in Macia *lgs1*-d
knockout lines. To assess whether this transcriptional signal translates into downstream
phenotypic effects, we evaluated mutant plants for differences in assimilation rate, chlorophyll
content, stomatal conductance, stomatal density, and the relative abundance of core light-
harvesting proteins (Q3).
602
603   In the strigolactone-deficient RTx430 *ccd8b* line, light-saturated net CO$_2$ assimilation
rate (A_{sat}) was significantly reduced compared to RTx430 WT (t(10) = 2.36, $P = 0.040$,
difference of genotypes by Student's *t*-test; Figure 2D). This reduction may reflect changes in
stomatal regulation and/or photosynthetic capacity. RTx430 *ccd8b* mutants had reduced stomatal
density on both abaxial and adaxial leaf surfaces (t(34) = -2.1684, $P = 0.037$, difference of
genotypes by Student's *t*-test; Figure S2A-B). Under non-limiting nutrient conditions, stomatal
conductance did not significantly differ between RTx430 *ccd8b* and WT plants, although it was
elevated in the mutant earlier in development (Figure S2C-D). RTx430 *ccd8b* also had lower
values for chlorophyll a/b ratio (Chl*a*/Chl*b*), the ratio of antenna complexes to PSII core proteins
(LHCB1/CP43), and the ratio of PSI to PSII core proteins (PSAA/CP43) potentiating physical

differences in photosynthetic machinery relative to the RTx430 WT; however, these differences
were not statistically significant (Figure S3). Taken together, these results suggest the
contribution of stomatal traits for reduced carbon assimilation in RTx430 *ccd8b*, although the
roles of stomatal versus biochemical limitations remain unresolved.
617
618
619

620
621
Figure 2 | Light saturated assimilation rate is reduced when strigolactone biosynthesis genes are knocked out
of normally functional backgrounds.
A) Net carbon assimilation rate of *LGS1* deletion (*lgs1*-d) mutants and null segregants (WT-L) of the same
transformation event (L2) generated in the Macia background. B) Assimilation rate of the Macia WT and *LGS1*
deletion (*lgs1*-d) mutants at varying levels of C_i. Data is collected with mutants of the S_2 generation, see Figure S4
for net carbon assimilation rate of S_2 generation Macia *lgs1*-d. C) Net carbon assimilation rate of mutants with
functional *LGS1* inserted and "wild type-like" null segregants (WT-L *lgs1*-2) of the same transformation event (L2)
generated in the RTx430 background. D) Net carbon assimilation rate of *ccd8b* mutants and WT in the RTx430
background. Asterisks denote a significant *P*-value comparing each mutant and wild type (WT or WT-L)
comparison by Student's *t*-test. *$P<0.05$, ***$P<0.001$ (*n*=6).
632

For mutants of the *LGS1* gene, the rate of light-saturated photosynthesis (A_{sat}) was on average 25% lower in the deletion mutant (Macia *lgs1*-d; $\chi^2(1) = 6.8157$, $P = 0.009$, likelihood ratio test for fixed effect of *LGS1* deletion; Figure 2A, Figure S4A), but was not significantly different in the insertion mutant (RTx430 *LGS1*; Figure 2C) relative to each WT comparison. Despite a reduced stomatal density on both abaxial and adaxial leaf surfaces for Macia *lgs1*-d (Figure S4D-E), stomatal conductance was not significantly different from the WT (Figure S4F), suggesting differences in Macia *lgs1*-d A_{sat} is not attributable to stomatal constraints. We next assessed if reductions in A_{sat} for Macia *lgs1*-d reflected changes in photosynthetic biochemistry, and additionally examined the maximum Rubisco carboxylation rate (V_{cmax}) and the maximum electron transport rate for ribulose-1,5-bisphosphate regeneration (J_{max}), estimated from A_{net}–C_i curves, along with fluorescence-derived traits (Fv/Fm, maximum quantum efficiency of photosystem II (PSII); ΦPSII, quantum yield of PSII; non-photochemical quenching (NPQ); electron transport rate (ETR)), and the relative abundance of core light-harvesting proteins.

In Macia *lgs1*-d lines, V_{cmax} was significantly reduced ($\chi^2(1) = 5.00$, $P = 0.025$, likelihood ratio test for fixed effect of *LGS1* deletion; Figure S4B) and apparent J_{max} showed a similar but non-significant trend toward reduction ($\chi^2(1) = 3.25$, $P = 0.07$, likelihood ratio test for fixed effect of *LGS1* deletion; Figure S4C). In contrast, fluorescence traits did not differ significantly between genotypes. Reductions in V_{cmax} and J_{max} are consistent with altered photosynthetic capacity, and may relate to differences in nitrogen allocation or photosystem investment (Walker et al., 2014). Despite reduced assimilation rates of Macia *lgs1*-d, the overall quantification of chlorophyll content and relative abundance of antenna and core photosystem proteins revealed no detectable differences between *lgs1*-d mutants and WT or WT-L lines in the Macia background (Figure S3). Together, the reductions in net carbon assimilation rate may be attributable to photosynthetic efficiency, such as Rubisco activity or activation state (as indicated by reduced V_{cmax} in Macia *lgs1*-d), rather than structural changes to the photosynthetic machinery. Notably, differences in assimilation rate between Macia and RTx430 functional and non-functional *LGS1* suggest *LGS1* effects as dependent on genetic background (Q4), and underscores the importance of evaluating QTL in multiple genetic backgrounds, when possible.

Mutation of strigolactone genes impacts the metabolome

Previous research has reported differences in the metabolic profiles of unrelated, non-isogenic sorghum genotypes that segregate for *LGS1* (Kawa et al., 2021) and some metabolites, especially flavonols, are observed to have strigolactone-dependent accumulation (*A. thaliana* (Walton et al., 2016)). We sought to identify any differences in the root and shoot metabolome of Sb*CCD8b* and *LGS1* knockout lines through untargeted metabolomics using ultra-high performance liquid chromatography-MS/MS (UHPLC-MS/MS). For both deletion mutants, we observed differences in metabolite intensity compared to wild type (Figure S5). Most of these shifts were in low-abundance compounds that we were unable to identify via library matching and molecular networking (Table S3, S4) and did not dramatically change the overall profile of the metabolome (PerMANOVA of root and shoot untargeted metabolomic profiles, both genes, $P > 0.5$; Figure S6). An exception was a major shift in the production of a disaccharide that we have tentatively identified as maltose, which was increased in the roots and shoots of RTx430 *ccd8b* and had variable shifts in the roots and shoots of Macia *lgs1*-d (Figure S5).

For RTx430 *ccd8b*, the overall base peak chromatogram (BPC) intensity of both roots
and shoots was higher than the RTx430 WT, indicating elevated total metabolite production
when Sb*CCD8b* is deleted (Figure S5A-B). The majority of RTx430 *ccd8b* differentially
accumulated metabolites were up-regulated in the roots (Figure S7A). Of the RTx430 *ccd8b* up-
regulated root metabolites, we putatively identified dihydroorotate which has been previously
reported as up-regulated in drought-sensitive sorghum (Li et al., 2022) and is consistent with
observations of strigolactone biosynthesis mutants' increased sensitivity to drought (Van Ha et
al., 2014). Further, dihydroorotate dehydrogenase (*DHODH*) is an enzyme involved in
pyrimidine biosynthesis and its overexpression in rice both increased *DHODH* activity and
enhanced plant tolerance to salt and drought stresses (Os*DHODH1*; (Liu et al., 2009)).
688
Though we did not detect significant differences between the overall metabolome of
Macia *lgs1*-d as compared to Macia WT Figure S6A-B, differentially accumulated metabolites
were more down-regulation in the roots (Figure S7C,E,G) and up-regulated in the shoots (Figure
S7D,F,H). A metabolite with a putative annotation of 2,6-Dimethylpyrazine, a natural product
belonging to the class of pyrazine-class volatile organic compounds (VOCs), was significantly
down-regulated in roots of Macia *lgs1*-d (Table S4). Though direct studies on 2,6-
dimethylpyrazine are limited, it is assigned to a pyrazine-class of VOCs that can activate defense
pathways (Patel et al., 2021), and act as signal molecules perceived by plants (Khashaveh et al.,
2025). Its down-regulation in the roots of Macia *lgs1*-d once again suggests alteration to stress
response when *LGS1* is knocked out of a normally functional background. For both Sb*CCD8b*
and *LGS1* loss-of-function mutants, differences in low-abundance metabolites suggesting some
metabolic pathways were affected, though the most abundant metabolites remained largely
unchanged.
702
703   *Strigolactone synthesis mutations impact root architecture and anatomy*
704
Strigolactones are well established to play critical roles in modulating root architecture
and more recently have been linked to root anatomical differences to optimize vessel element
formation (Zhao et al., 2025) and in aerenchyma formation (Shrestha et al., 2026). Consistent
with previous reports by Hoa et al., (2023), which demonstrated RTx430 *ccd8b* to have fewer
root tips and overall smaller root systems, we also report RTx430 *ccd8b* plants had significantly
smaller root systems (t(14) = 4.54, *P* < 0.001, difference of genotypes by Student's t-test; Figure
3C). Additionally, we found that RTx430 *ccd8b* root systems had significantly more nodal roots
(t(14) = -3.17, *P* = 0.007, difference of genotypes by Student's t-test; Figure 3D), with roots that
were smaller in cross section area (t(23) = -3.43, *P* = 0.002, difference of genotypes by Student's
t-test; Figure 3E), and had altered root anatomy (Figure 3A-B). Most notably, RTx430 *ccd8b*
aerenchyma area (air space in the root cortex) was significantly reduced (t(17) = -2.60, *P* =
0.019, difference of genotypes by Student's t-test; Figure 3H) and despite maintaining the same
number of metaxylem vessels, trended towards reduction in total metaxylem vessel area (t(9) = -
1.71, *P* = 0.123, difference of genotypes by Student's t-test; Figure 3F-G).

724
725
Figure 3 | Root Architecture and anatomy is altered in mutants with Sb*CCD8b* deleted. Two representative
root cross sections captured by laser ablation tomography (LAT) for the RTx430 background
A) WT and B) *ccd8b* mutants. The scale bar in each photo is 0.5 mm. Boxplots compare the RTx430 WT (black)
and *ccd8b* mutants (gray) for the root architecture phenotypes of C) total root system network area D) number of
nodal roots and the root anatomical traits of E) root cross-section area F) total metaxylem vessel area G) number of
metaxylem vessels H) total aerenchyma area. Asterisks denote a significant *P*-value comparing WT and mutant by
Student's *t*-test. *$P<0.05$, **$P<0.01$, ***$P<0.001$.
733
Macia *lgs1*-d maintained a greater number of metaxylem vessel elements ($\chi^2(1) = 5.67$,
$P = 0.017$, likelihood ratio test for fixed effect of *LGS1* deletion; Figure S8E) with individual
elements that were smaller in size ($\chi^2(1) = 4.20$, $P = 0.04$, likelihood ratio test for fixed effect of

LGS1 deletion; Figure S8F). For the aerenchyma area trait, the Macia *lgs1*-d mutant had a
reduced formation of airspace in the cortex as compared to the Macia WT and WT-L null
segregants, which showed greater plasticity in the formation of nodal root aerenchyma (χ^2(1)
=1.26, *P* = 0.26, likelihood ratio test for fixed effect of *LGS1* deletion; Figure S8G). Differences
in root architecture anatomy for the RTx430 mutants with functional *LGS1* inserted were not as
pronounced and not consistent with observations in Macia *lgs1*-d mutants. RTx430 *LGS1* mutant
had a greater number of nodal roots as compared to the RTx430 WT (t(43)=-2.715, *P* = 0.001,
likelihood ratio test for fixed effect of *LGS1* deletion; Figure S9B). All other root traits were not
significantly different. Interestingly, both genotypes with functional strigolactone genes knocked
out (RTx430 *ccd8b*, Macia *lgs1*-d) had reduced aerenchyma area and is consistent with recent
reports by (Shrestha et al., 2026) that strigolactones play a role in aerenchyma formation.
748
749   *Strigolactone synthesis mutants grown in nutrient and mycorrhizal treatments*
750
We hypothesized the alteration of sorghum strigolactone profiles from loss-of-function at
LGS1 that reduce parasite stimulation (Figure 1, Bellis et al. 2020), would also be coupled with
trade-offs in plant growth and interactions with arbuscular mycorrhizal fungi (AMF). As
strigolactones mediate rhizosphere signaling for symbiosis, we predicted that loss-of-function at
Sb*CCD8b* and *LGS1* would impair mycorrhizal establishment (Q1). To test this, all genotypes
(Table 1) were evaluated under three conditions: low phosphate without mycorrhizae, low
phosphate with mycorrhizae, and high phosphate without mycorrhizae (Figure 4).
758
We observed strong treatment effects, and plants were the tallest and largest in the high
phosphate treatment (P+ AMF-, Figure 5A). The application of phosphate (P+) and mycorrhizal
inoculation (AMF+) each independently increased plant height, shoot weight, root weight, a
SPAD-based measurement of chlorophyll concentration, and stomatal conductance, as compared
to the low phosphate non-mycorrhizal treatment (P- AMF-)(Figure 5A-E). For all genotypes,
chlorophyll concentration and stomatal conductance were the lowest in the low phosphate no
mycorrhizae treatment (P- AMF-), as compared to the treatments with high phosphate (P+,
AMF-) and with mycorrhizae (P- AMF+) (Figure 5D-E), indicating that the low phosphate no
mycorrhizae treatment (P- AMF-) was indeed stressful.
768

Treatment

■ High Phosphate, No AMF (P+ AMF -)   ■ Low Phosphate, AMF (P- AMF +)   ■ Low Phosphate, No AMF (P- AMF -)

Figure 4 | Representative images of greenhouse grown plants taken at 28 days after plant emergence.
For a genotype (denoted above image), three replicates grown in the three treatments are included in a photo. The
color bar corresponds to the treatment a plant was grown in. Treatments are high phosphate and no arbuscular
mycorrhizal fungi (P+ AMF- ; blue), low phosphate and arbuscular mycorrhizal fungi (P- AMF+ ; orange), and low
phosphate and no arbuscular mycorrhizal fungi (P- AMF- ; red). A,B) Comparison of "wild type -like" null
segregants (WT-L) and deletion mutants of the *LGS1* gene (*lgs1*-d) in the Macia background. C,D) Comparison of
"wild type -like" null segregants (WT-L *lgs1*-2) and insertion mutants of the *LGS1* gene RTx430 background. E,F)
Comparison of WT and deletion mutants of the Sb*CCD8b* gene (*ccd8b*) in the RTx430 background. Description of
genotypes are reported in Table 1.

      In all three treatments, the RTX430 *ccd8b* mutant was significantly shorter (Figure 4F,
Figure 5A), had lower shoot weight (Figure 5B), and lower root weight (Figure 5C) as compared
to the RTx430 WT. RTx430 *ccd8b* plants grown in the treatment without nutrient limitation (P+
AMF-) displayed phenotypic differences classically observed in strigolactone biosynthetic
mutants. RTx430 *ccd8b* had increased shoot branching and tillering later in growth (data not
shown), though this phenotype was not observed in RTx430 *ccd8b* plants grown in either low
phosphate treatments with (P- AMF+) or without (P- AMF-) mycorrhizae, which may be
attributable to a reduced growth rate under stressful conditions. These observations in the
RTx430 *ccd8b* mutant support that Sb*CCD8b* loss-of-function results in pleiotropic effects,
independent of nutrient availability. Nevertheless, under nutrient limitation, phenotypic effects
were especially pronounced and growth was severely stunted compared to the RTx430 wild type
(Figure 4E,F).

792
793

794
Figure 5 | Phenotypes collected in greenhouse experiment.
A) Plant height B) shoot dry weight C) root fresh weight of plants collected at Timepoint 2 (42 days). D) Single-
photon avalanche diode (SPAD)-based measurement of leaf chlorophyll and E) Stomatal conductance collected at
28 days after plant emergence. Boxplots colored by treatment: high phosphate and no arbuscular mycorrhizal fungi
(P+ AMF- ; blue), low phosphate and arbuscular mycorrhizal fungi (P- AMF+ ; orange), low phosphate and no
arbuscular mycorrhizal fungi (P- AMF- ; red). Description of genotypes are reported in Table 1.
801
802
803

There were no significant differences in root or shoot weight for Macia *lgs1*-d grown in
the high phosphate (P+ AMF-) treatment earlier in growth (21 days); however, later in growth
(42 days) dry root biomass was significantly lower ($\chi^2(1) = 4.18$, $P = 0.03$, likelihood ratio test
for fixed effect of *LGS1* deletion; Figure S10) as compared to the Macia WT and WT-L null
segregants. As such, for plants of the Macia background differing in functionality of *LGS1*
grown without nutrient limitation, differences in growth may be subtle, only detectable at later
points in development, and *lgs1*-d plants may benefit less from the application of phosphate. For
plants differing for the functionality of *LGS1* in the RTx430 background grown in the high
phosphate treatment (P+ AMF-), we did not find evidence for any of the traits we measured (root
weight, shoot weight, SPAD, or plant height) to be significantly different, again supporting that
LGS1 effects may depend on variation in other genes (Q4).
815
816 *Strigolactone biosynthesis deletion mutants have impaired colonization of mycorrhizae*
817
At the first time point (21 days), RTx430 *ccd8b* plants had significantly delayed and
overall reduced mycorrhizal colonization ($t(9) = -6.65$, $P < 0.001$, difference of genotypes by
Student's t-test; Figure 6B, Figure 7C), implicating Sb*CCD8b* to be involved in the early
establishment of mycorrhizal partners. Furthermore, across both timepoints (21 and 42 days)
RTx430 *ccd8b* root weight, shoot weight, and plant height remained significantly reduced as
compared to RTx430 WT plants grown in the same treatment (Figure 5, Figure 6D, Figure 7C-
D), indicating a relatively low benefit from the addition of mycorrhizae for plants with loss of
function at Sb*CCD8b*. For RTx430 genotypes grown with (P- AMF+) and without (P- AMF-)
mycorrhizae at 42 days, the relative increase in shoot weight was 20% lower for RTx430 *ccd8b*
($t(8.2) = -4.53$, $P = 0.002$, difference of genotypes by Student's t-test; Figure 6D), suggesting
that the impaired establishment of AMF, in turn, influenced the benefit mycorrhizal symbionts
provided to RTx430 *ccd8b*.
830
For plants of the Macia background at the earlier time point (21 days), mycorrhizal
colonization was on average 48% lower in Macia *lgs1*-d as compared to the colonization of WT
and WT-L null segregants ($\chi^2(1) = 11.38$, $P < 0.001$, likelihood ratio test for fixed effect of
LGS1 deletion; Figure 6A, Figure 7A). This disruption in Macia *lgs1*-d early mycorrhizal
colonization was also apparent from differences in shoot traits. At the earlier harvest time point
(21 days) of the low phosphate mycorrhizal treatment (P- AMF+), Macia with functional *LGS1*
had greater shoot weight than the Macia *lgs1*-d mutants ($t(20) = -1.7327$, $P = 0.1$, difference of
genotypes by Student's t-test; Figure 6C, Figure 4A-B). Additionally, plants of the Macia
background with functional *LGS1* gained a greater benefit from the addition of mycorrhizae
($t(38) = -2.69$, $P = 0.01$, Macia *LGS1* difference between low phosphate mycorrhizal and non-
mycorrhizal treatment by Student's t-test), where Macia *lgs1*-d did not show any difference in
average shoot weight between mycorrhizal (P- AMF+) and non-mycorrhizal (P- AMF-)
treatments (Figure 6C).

844
Figure 6| Mycorrhizal colonization is disrupted when strigolactone biosynthesis genes are knocked out of
normally functional backgrounds.
A) Percent of roots colonized by the mycorrhizal species *R. intraradices* at Timepoint 1 (21 days) and Timepoint 2
(42 days) for WT, "wild type-like" null segregants (WT-L), and *LGS1* deletion (*lgs1*-d) lines of the Macia
background. B) Percent of roots colonized by mycorrhizal species *R. intraradices* at Timepoint 1 (21 days) and
Timepoint 2 (42 days) for WT, *ccd8b* mutants, "wild type-like" null segregants (WT-L), and *LGS1* insertion mutants
(*LGS1*) of the RTx430 background. WT-L and mutants of the same transformation event (L1,2,3) indicate paired
genotypes. Significance of the allele effect (*LGS1* or Sb*CCD8b*, respectively) from linear models is reported. Dry
shoot weight at Timepoint 1 (21 days) and Timepoint 2 (42 days) for C) Macia genotypes and D) RTx430 genotypes
grown under low phosphate conditions, with (AMF+ P-) and without mycorrhizae (AMF- P-). The mean dry shoot
weight for genotypes with the same allele grown in a treatment is plotted as a large dot with whiskers extending to
the 95% confidence intervals.

At the later time point (42 days), while mycorrhizal colonization remained 15% lower for
Macia *lgs1*-d as compared to colonization of WT and WT-L null segregants, the difference in
colonization was no longer significant ($\chi^2(1) = 2.48$, $P = 0.115$, likelihood ratio test for fixed
effect of *LGS1* deletion; Figure 6A,C, Figure 7B). Shoot dry weight between Macia *lgs1*-d and
Macia lines with functional *LGS1* (WT, WT-L null segregants) at the later time point (42 days)
increased in a similar manner for plants grown with (P- AMF+) and without (P- AMF-)
mycorrhizae. This indicates that all Macia genotypes, regardless of *LGS1* functionality, benefit
from the addition of mycorrhizae given sufficient time to associate with mycorrhizal symbionts.
For plants of the RTx430 background, we did not find evidence for the functionality of *LGS1* to

be associated with differences in mycorrhizal colonization. Across timepoints, mycorrhizal
colonization was not significantly different between the RTx430 WT or WT-L plants
maintaining the *lgs1*-2 deletion allele and the mutants with functional *LGS1* inserted (Figure
6B,D, Figure 7C,D).
872

873
874 **Figure 7 | Percent roots colonized by mycorrhizae plot against root fresh weight.**
875 Genotypes of the Macia background at A) Timepoint 1 (21 days) and B) Timepoint 2 (42 days). Genotypes of the
876 RTx430 background at C) Timepoint 1 (21 days) and D) Timepoint 2 (42 days). Points and regression lines are
877 colored by allele.
878
879
880

Discussion

Strigolactones are hormones that regulate plant development and physiology, while also acting as rhizosphere signals that mediate interactions with symbiotic microbes and parasitic plants. Our study explored the impact of strigolactones in sorghum, a crop vital to semiarid regions with often nutrient-poor soils, by examining mutants of the strigolactone biosynthesis genes Sb*CCD8b* and *LGS1*.

Artificial CRISPR-Cas9 knockouts of strigolactone biosynthesis genes affected many traits. Loss-of-function in Sb*CCD8b* led to changes in growth (Figure 5E,F), altered physiology (Figure 2D), disrupted mycorrhizal establishment (Figure 6B,D), significant changes to low-abundance root metabolites (Figure S7A), and changes in root architecture and anatomy (Figure 3). Similarly, deletion of functional *LGS1* in the Macia background (Macia *lgs1*-d) resulted in transcriptional changes of shoots enriched for promoter motifs for stress response and growth pathways (Table S2), a down-regulation of root metabolites and up-regulation of shoot metabolites (Figure S7C-H), reduced carbon assimilation (Figure 2A), impaired early mycorrhizal colonization (Figure 6A,C), and reductions in root growth under non-limiting conditions (Figure S10). Together, our study demonstrates that deletion of the tested strigolactone biosynthesis genes impacts both endogenous developmental processes and exogenous signaling functions, leading to measurable trade-offs in plant performance (Q3). At the same time, we hypothesize epistasis with a neighboring strigolactone biosynthesis gene may modify the effects of loss-of-function at *LGS1*.

Genetic background dependent trade-offs and evidence for epistasis

A key finding of our study is that the magnitude and direction of *LGS1*-associated trade-offs depend strongly on genetic background (Q4). While deletion of *LGS1* in the Macia background resulted in clear impeded photosynthesis and AMF colonization, reintroduction of functional *LGS1* into the RTx430 background restored parasite susceptibility (Figure 1) without enhancing photosynthesis or AMF colonization. Aside from increased nodal root number under high phosphate conditions (Figure S9B), RTx430 mutants with functional *LGS1* inserted showed minimal evidence for differences compared to RTx430 WT and WT-L null segregants (*lgs1*-2). This apparent epistasis may be due to another gene near *LGS1* involved in strigolactone biosynthesis. In addition to *LGS1* (*Sobic.005G213600*), the *lgs1*-2 deletion allele carried by RTx430 also includes deletion of *Sobic.005G213400* and *Sobic.005G213500* (*Sb3500*) (Table 2), genes we did not insert into RTx430 along with *LGS1*. *Sb3500* is a 2-oxoglutarate-dependent dioxygenase affecting the conversion of carlactone to 5DS (Yoda et al., 2023) and is ~3 kb upstream of the *LGS1* gene. Yoda et al. 2023 previously showed epistasis between *Sb3500* and *LGS1* for strigolactone synthesis when expressed with other strigolactone biosynthetic genes in *Nicotiana benthamiana* (expected strigolactones shown in Table 2), suggesting the insertion mutant RTx430 *LGS1* did not have the typical strigolactone profile of Macia WT. It may be that restoration of functional *Sb3500* and a primarily 5DS strigolactone profile would be required to enhance photosynthesis and AMF colonization of RTx430. Though we found only minimal phenotypic differences between RTx430 WT and the RTx430 *LGS1* insertion mutant, the combination of alles in the *LGS1* insertion mutant, absent *Sb3500* and present *LGS1*, is extremely rare in natural sorghum genotypes (only 1 of 704 landraces, Table 2) and may indicate fitness costs associated with this allelic combination. RTx430 is an extensively used breeding

line (Gurel et al., 2009) for both research and hybrid production, and other natural mutations in
RTx430 may influence *LGS1* deletion effects. Additionally, recent studies suggest some
endogenous strigolactone signalling is driven by non-canonical strigolactones and not canonical
(e.g. 5DS, orobanchol) strigolactones (Ito et al., 2022), and so the effects of mutations in
canonical strigolactone biosynthesis genes like *LGS1* and resulting feedbacks require more
dissection. Future work would benefit from disentangling the allelic effects of *LGS1* and *Sb3500*
from differences in genetic background, and functionally validating the impact of *Sb3500* on
sorghum strigolactone exudate profiles.

Linking strigolactones, stress responses, and differences in photosynthesis

The pleiotropic effects observed in the strigolactone biosynthesis mutants used in
experiments are consistent with known crosstalk between strigolactones and stress-response
pathways (Q2). Strigolactones interact with abscisic acid (ABA) signaling and contribute to plant
responses to drought and other abiotic stresses (Korek and Marzec, 2023). In our study,
Sb*CCD8b* mutants in the RTx430 background trended towards elevated stomatal conductance
earlier in growth (Figure S3C) and RTx430 *ccd8b* roots had an increased accumulation of
dihydroorotate (Table S3), a metabolite associated with drought-sensitive genotypes (Li et al.,
2022). Additionally, both Sb*CCD8b* and *LGS1* knockout mutants exhibited reduced net carbon
assimilation (Figure 2A,C; Figure S4A). In Macia *lgs1*-d, this reduction was related to decreases
in maximum rate of carboxylation (V_{cmax}) (Figure S4D), suggesting limitations to Rubisco
activity when functional *LGS1* is deleted. Relatedly, enrichment of AP2/EREBP transcription
factor binding sites in the promoter regions of Macia *LGS1* knockouts (Macia *lgs1*-d)
differentially expressed genes supports a regulatory basis for these effects. This family of
transcription factors is involved in both stress responses (Xu et al., 2011) and photosynthetic
regulation (Hong et al., 2009). As such, AP2/EREBPs have been suggested as ideal candidates
for crop improvement and their overexpression enhances tolerances to multiple abiotic and biotic
stressors (Xu et al., 2011). For example, overexpression of the At*DREB1A* (AP2/EREBP
subfamily transcription factor) in Chrysanthemum resulted in significantly higher photosynthetic
capacity, and elevated activity of Rubisco (Hong et al., 2009). Together, the decreased carbon
assimilation rate in Macia *lgs1*-d may be attributable to the AP2/EREP or other associated
regulatory networks, although additional work is needed to resolve the underlying mechanisms.
957
958 *Developmental regulation and the miR156–SPL pathway*
959
Many of the phenotypes altered in the strigolactone biosynthesis knockout mutants,
including - reduced stomatal density, decreased mycorrhizal colonization, altered photosynthetic
rates, and increased nodal root production - are known to be regulated by the miR156-SPL
pathway (Poethig, 2013; Aung et al., 2015; Lawrence et al., 2021) which regulates
developmental timing and the transition of vegetative characteristics from juvenile to adult (Q2).
The majority of phenotypes we collected in greenhouse experiments were collected from plants
transitioning from the juvenile to the adult stage (average last juvenile leaf = node 3, average
first adult leaf = node 7, all leaves between these positions are considered transition leaves) and
traits observed in the RTx430 *ccd8b* and Macia *lgs1*-d mutants are consistent with a more
"juvenile-like" developmental state, associated with elevated miR156 and reduced SPL
expression. Previous studies have linked the miR156-SPL pathway with strigolactone
biosynthesis and signaling. SPL proteins mediate strigolactone signaling through direct

interaction with the signaling molecule DWARF53 (D53), which binds SPLs in the absence of
strigolactones suppressing their transcriptional activity (Barbier et al., 2023). Further, SPLs
promote expression of *CCD* and *NCED* genes, key enzymes in strigolactone biosynthesis
(Arango et al., 2016). Supporting this, the miR156-SPL pathway regulates miR172 expression,
which targets AP2 transcription factors (Ó'Maoiléidigh et al., 2021), the largest group of
transcription factors enriched in our analysis of regulatory pathways related to patterns of
differential expression in Macia *lgs1*-d mutants (Table S2). Consequently, the range of
phenotypes altered when functional strigolactone biosynthesis genes are knocked out (RTx430
ccd8b and Macia *lgs1*-d) may be associated with, or potentially mediated by, changes in
developmental timing through the miR156-SPL regulatory network.

Impacts on nutrient signaling and mycorrhizal symbiosis

Strigolactones play a central role in nutrient signaling, particularly under phosphate
limitation, where they promote root architectural changes and facilitate recruitment of
mycorrhizal symbionts (Mayzlish-Gati et al., 2010; Ruyter-Spira et al., 2011; Alder et al., 2012;
Sun et al., 2014). Previous observations by Hao et al. (2023) demonstrated significant shifts in
the root-associated microbiome in Sb*CCD8b* knockouts. In agreement with these findings, we
found the same RTx430 *ccd8b* knockout lines exhibited significantly reduced colonization by
arbuscular mycorrhizal fungi and diminished benefits from this symbiotic partner (Q1, Figure
6B,D). We additionally demonstrate the disruption of early mycorrhizal colonization for Macia
lgs1-d (Figure 6A, Figure 7A), though given sufficient time, the relative colonization and benefit
of mycorrhizae appears similar for plants of the Macia background regardless of *LGS1*
functionality (Q1, Figure 7B). The effects of impaired symbiosis with AMF may arise from
multiple mechanisms, including altered strigolactone stereochemistry, reduced quantify
strigolactone in root exudates, or both. Results from *S. hermonthica* germination assays reported
in this study and previously (Bellis et al. 2020) are consistent with the hypothesis that absence of
functional *LGS1* reduces total strigolactone exudation. Additionally, both orobanchol and 5DS
are known to stimulate mycorrhizal branching (Akiyama et al., 2005; Akiyama et al., 2010).
Reduced early colonization for Macia *lgs1*-d may therefore reflect insufficient signaling through
reduced strigolactone exudation (Figure 7A), while the recovery at the later time point may
indicate cumulative and sufficient signaling over time. In contrast, RTx430 *LGS1* insertion
mutants showed no difference in mycorrhizal colonization at either time point (Figure 6B,D),
despite a clear difference in strigolactone profiles from RTx430 WT and WT-L plants (*lgs1*-2),
as evidenced by elevated germinability of parasite populations (Q1, Figure 1A-B). The *LGS1*
insertion in the RTx430 background is expected to shift strigolactone composition from
orobanchol toward 5DS, though likely to a lesser extent than in Macia WT due to the absence of
Sb3500 (Table 2). As plant responses to mycorrhizal colonization can vary across fungal species
and strains (Angelard et al., 2010) and the same sorghum genotype can have substantial variation
in colonization rates between fungal species (Gobena et al., 2017), our conclusions are limited to
this *R. intraradices* experiment.

Implications for breeding and deployment of LGS1 alleles

From an applied perspective, *LGS1* loss-of-function alleles are of considerable interest
for improving sorghum's resistance to *S. hermonthica* (Hess et al., 1992; Ejeta, 2007). Our
results demonstrate that such alleles, specifically deleting functional *LGS1* from a normally

functional background (Macia), reduce parasite germination (Bellis et al. 2020) but may also be coupled with trade-offs in growth, photosynthesis, and early establishment of mycorrhizal symbiosis. The absence of strong trade-offs in comparing the RTx430 WT (*lgs1*-2) to mutants with functional *LGS1* inserted suggests that these costs are not inevitable and may be mitigated by genetic background; however, the rarity of the combination of alleles present in the RTx430 *LGS1* insertion mutant may suggest other fitness trade-offs we did not observe. Epistasis, in particular genetic backgrounds or between multiple genes that impact sorghum strigolactone profiles (e.g. *Sb3500* and *LGS1*), may therefore represent vital variation for optimizing plant performance while maintaining resistance to *S. hermonthica* in farmer-preferred varieties. At the same time, we caution that the experiments described here likely do not fully capture the dynamics of field-grown sorghum, and any trade-offs incurred from deleting *LGS1* from a normally functional background could be preferable to crop loss caused by *S. hermonthica* pressure. Our study advances understanding of strigolactone biology in sorghum and highlights the complexity of co-evolutionary dynamics, where functional variation at a single locus can confer parasite resistance but with evidence for costs to host fitness.

Acknowledgements & Funding We thank Scott Diloreto and Sam Gruneberg for sharing their greenhouse and plant care expertise. We are grateful to Amanda Penn, Shiran Ben-Zeev, Diana Gamba, and Eric Nascari for their assistance with greenhouse data collection. We also thank the Corteva Sorghum transformation team for generation of RTx430 transgenic events, control environment team for taking care of and backcrossing plants, and the analysis team for molecular characterization of the material. RJHS is funded by the USDA National Institute of Food and Agriculture and Hatch Appropriations under Project No. PEN05039 and Accession No. 7008935, as well as by the USDA project "The Genetic Basis of Maize Response to Arbuscular Mycorrhizal Fungi" (Grant No. 2022-67013-38264). DPS acknowledges support from the USDA National Institute of Food and Agriculture (Grant No. 2020-67019-31796). JJK is supported by USDA NIFA and Hatch Appropriations under Project No. PEN04956 and Accession No. 7006496. JRL, JM, and CMM acknowledge support from the Bill & Melinda Gates Foundation.

Author Contributions CMM, JRL, RHJS conceived and planned the research. CMM, MT, JM, EJA, AC, AC, YX, EHL-P, and MP performed the experiments and analyzed data. Plant genetic material was developed by DPS and HG. *Striga hermonthica* seed was collected by SR and BN. CMM, JRL, and EHL-P wrote the paper with input from all authors.

References

Akiyama K, Matsuzaki K-I, Hayashi H (2005) Plant sesquiterpenes induce hyphal branching in arbuscular mycorrhizal fungi. Nature 435: 824–827

Akiyama K, Ogasawara S, Ito S, Hayashi H (2010) Structural requirements of strigolactones for hyphal branching in AM fungi. Plant Cell Physiol 51: 1104–1117

Alder A, Jamil M, Marzorati M, Bruno M, Vermathen M, Bigler P, Ghisla S, Bouwmeester H, Beyer P, Al-Babili S (2012) The path from β-carotene to carlactone, a strigolactone-like plant hormone. Science 335: 1348–1351

Angelard C, Colard A, Niculita-Hirzel H, Croll D, Sanders IR (2010) Segregation in a mycorrhizal fungus alters rice growth and symbiosis-specific gene transcription. Curr Biol 20: 1216–1221

Arango J, Beltrán J, Nuñez J, Chavarriaga P (2016) Evidence of epigenetic mechanisms affecting carotenoids. Subcell Biochem 79: 295–307

Arite T, Iwata H, Ohshima K, Maekawa M, Nakajima M, Kojima M, Sakakibara H, Kyozuka J (2007) DWARF10, an RMS1/MAX4/DAD1 ortholog, controls lateral bud outgrowth in rice. Plant J 51: 1019–1029

Arite T, Kameoka H, Kyozuka J (2012) Strigolactone positively controls crown root elongation in rice. J Plant Growth Regul 31: 165–172

Aung B, Gruber MY, Hannoufa A (2015) The MicroRNA156 system: A tool in plant biotechnology. Biocatal Agric Biotechnol 4: 432–442

Bailey TL (2021) STREME: accurate and versatile sequence motif discovery. Bioinformatics 37: 2834–2840

Barbier F, Fichtner F, Beveridge C (2023) The strigolactone pathway plays a crucial role in integrating metabolic and nutritional signals in plants. Nat Plants 9: 1191–1200

Bellis ES, Kelly EA, Lorts CM, Gao H, DeLeo VL, Rouhan G, Budden A, Bhaskara GB, Hu Z, Muscarella R, et al (2020) Genomics of sorghum local adaptation to a parasitic plant. Proc Natl Acad Sci U S A 117: 4243–4251

Brundrett MC, Tedersoo L (2018) Evolutionary history of mycorrhizal symbioses and global host plant diversity. New Phytol 220: 1108–1115

Burton AL, Williams M, Lynch JP, Brown KM (2012) RootScan: Software for high-throughput analysis of root anatomical traits. Plant Soil 357: 189–203

Butt H, Jamil M, Wang JY, Al-Babili S, Mahfouz M (2018) Engineering plant architecture via CRISPR/Cas9-mediated alteration of strigolactone biosynthesis. BMC Plant Biol 18: 174

Chien P-S, Chiang C-P, Leong SJ, Chiou T-J (2018) Sensing and signaling of phosphate starvation: From local to long distance. Plant Cell Physiol 59: 1714–1722

Cook CE, Whichard LP, Turner B, Wall ME, Egley GH (1966) Germination of witchweed (Striga lutea Lour.): Isolation and properties of a potent stimulant. Science 154: 1189–1190

Croce R, Canino G, Ros F, Bassi R (2002) Chromophore organization in the higher-plant photosystem II
antenna protein CP26. Biochemistry 41: 7334–7343

Doran H, Bates D, Bliese P, Dowling M (2007) Estimating the multilevel Rasch model: With
thelme4Package. J Stat Softw. doi: 10.18637/jss.v020.i02

Ejeta G (2007) Breeding for *Striga* resistance in sorghum: Exploitation of an intricate host–parasite
biology. Crop Sci 47: S–216–S–227

Fujimoto SY, Ohta M, Usui A, Shinshi H, Ohme-Takagi M (2000) Arabidopsis ethylene-responsive
element binding factors act as transcriptional activators or repressors of GCC box-mediated gene
expression. Plant Cell 12: 393–404

Gao J, Zhang T, Xu B, Jia L, Xiao B, Liu H, Liu L, Yan H, Xia Q (2018) CRISPR/Cas9-mediated
Mutagenesis of carotenoid cleavage dioxygenase 8 (CCD8) in tobacco affects shoot and root
architecture. Int J Mol Sci. doi: 10.3390/ijms19041062

Giovannetti M, Mosse B (1980) An evaluation of techniques for measuring vesicular arbuscular
mycorrhizal infection in roots. New Phytol 84: 489–500

Giovannetti M, Sbrana C, Avio L, Citernesi AS, Logi C (1993) Differential hyphal morphogenesis in
arbuscular mycorrhizal fungi during pre-infection stages. New Phytol 125: 587–593

Giovannetti M, Sbrana C, Logi C (1994) Early processes involved in host recognition by arbuscular
mycorrhizal fungi. New Phytol 127: 703–709

Gobena D, Shimels M, Rich PJ, Ruyter-Spira C, Bouwmeester H, Kanuganti S, Mengiste T, Ejeta G
(2017) Mutation in sorghum LOW GERMINATION STIMULANT 1 alters strigolactones and
causes Striga resistance. Proc Natl Acad Sci U S A 114: 4471–4476

Gomez-Roldan V, Fermas S, Brewer PB, Puech-Pagès V, Dun EA, Pillot J-P, Letisse F, Matusova R,
Danoun S, Portais J-C, et al (2008) Strigolactone inhibition of shoot branching. Nature 455: 189–194

Guan JC, Koch KE, Suzuki M, Wu S, Latshaw S, Petruff T, Goulet C, Klee HJ, McCarty DR (2012)
Diverse roles of strigolactone signaling in maize architecture and the uncoupling of a branching-
specific subnetwork. Plant Physiol 160: 1303–1317

Gupta S, Stamatoyannopoulos JA, Bailey TL, Noble WS (2007) Quantifying similarity between motifs.
Genome Biol 8: R24

Gurel S, Gurel E, Kaur R, Wong J, Meng L, Tan H-Q, Lemaux PG (2009) Efficient, reproducible
Agrobacterium-mediated transformation of sorghum using heat treatment of immature embryos.
Plant Cell Rep 28: 429–444

Gurney AL, Grimanelli D, Kanampiu F, Hoisington D, Scholes JD, Press MC (2003) Novel sources of
resistance to Striga hermonthica in Tripsacum dactyloides, a wild relative of maize. New Phytol 160:
557–568

Haider I, Yunmeng Z, White F, Li C, Incitti R, Alam I, Gojobori T, Ruyter-Spira C, Al-Babili S,
Bouwmeester HJ (2023) Transcriptome analysis of the phosphate starvation response sheds light on
strigolactone biosynthesis in rice. Plant J 114: 355–370

Hao J, Yang Y, Futrell S, Kelly EA, Lorts CM, Nebie B, Runo S, Yang J, Alvarez S, Lasky JR, et al

(2023) CRISPR/Cas9-mediated Mutagenesis of *carotenoid cleavage dioxygenase* (*CCD*) genes in sorghum alters strigolactone biosynthesis and plant biotic interactions. Phytobiomes J. doi: 10.1094/pbiomes-08-22-0053-r

Hess DE, Ejeta G, Butler LG (1992) Selecting sorghum genotypes expressing a quantitative biosynthetic trait that confers resistance to Striga. Phytochemistry 31: 493–497

Hong B, Ma C, Yang Y, Wang T, Yamaguchi-Shinozaki K, Gao J (2009) Over-expression of AtDREB1A in chrysanthemum enhances tolerance to heat stress. Plant Mol Biol 70: 231–240

Ito S, Braguy J, Wang JY, Yoda A, Fiorilli V, Takahashi I, Jamil M, Felemban A, Miyazaki S, Mazzarella T, et al (2022) Canonical strigolactones are not the major determinant of tillering but important rhizospheric signals in rice. Sci Adv 8: eadd1278

Kawa D, Taylor T, Thiombiano B, Musa Z, Vahldick HE, Walmsley A, Bucksch A, Bouwmeester H, Brady SM (2021) Characterization of growth and development of sorghum genotypes with differential susceptibility to Striga hermonthica. J Exp Bot 72: 7970–7983

Khashaveh A, Yi C, Tang H, Song X, Zhang G, Xie J, Zhang Y (2025) Biological roles of pyrazines in insect chemical communication. Chem Biol Technol Agric. doi: 10.1186/s40538-025-00830-x

Kodama K, Rich MK, Yoda A, Shimazaki S, Xie X, Akiyama K, Mizuno Y, Komatsu A, Luo Y, Suzuki H, et al (2022) An ancestral function of strigolactones as symbiotic rhizosphere signals. Nat Commun 13: 3974

Kohlen W, Charnikhova T, Lammers M, Pollina T, Tóth P, Haider I, Pozo MJ, de Maagd RA, Ruyter-Spira C, Bouwmeester HJ, et al (2012) The tomato CAROTENOID CLEAVAGE DIOXYGENASE8 (SlCCD8) regulates rhizosphere signaling, plant architecture and affects reproductive development through strigolactone biosynthesis. New Phytol 196: 535–547

Korek M, Marzec M (2023) Strigolactones and abscisic acid interactions affect plant development and response to abiotic stresses. BMC Plant Biol 23: 314

Koske RE, Gemma JN (1989) A modified procedure for staining roots to detect VA mycorrhizas. Mycol Res 94: 486–488

Lawrence EH, Springer CJ, Helliker BR, Poethig RS (2021) MicroRNA156-mediated changes in leaf composition lead to altered photosynthetic traits during vegetative phase change. New Phytol 231: 1008–1022

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R, 1000 Genome Project Data Processing Subgroup (2009) The Sequence Alignment/Map format and SAMtools. Bioinformatics 25: 2078–2079

Liu W-Y, Wang M-M, Huang J, Tang H-J, Lan H-X, Zhang H-S (2009) The OsDHODH1 gene is involved in salt and drought tolerance in rice. J Integr Plant Biol 51: 825–833

Li Y, Tan B, Wang D, Mu Y, Li G, Zhang Z, Pan Y, Zhu L (2022) Proteomic analysis revealed different molecular mechanisms of response to PEG stress in drought-sensitive and drought-resistant sorghums. Int J Mol Sci 23: 13297

Maina F, Faye JM, Kena AW, Diatta C, Tankari MI, Ardaly OA, Diakite OS, Ibrahim AM, Moussa OA, Harou A, et al (2025) Delivering trait-enhanced varieties to African smallholders through a

1173   pangenomic breeding network. bioRxiv. doi: 10.1101/2025.08.07.667917

1174 Mallu TS, Mutinda S, Githiri SM, Achieng Odeny D, Runo S (2021) New pre-attachment Striga resistant
1175   sorghum adapted to African agro-ecologies. Pest Manag Sci 77: 2894–2902

1176 Matusova R, van Mourik T, Bouwmeester HJ (2004) Changes in the sensitivity of parasitic weed seeds to
1177   germination stimulants. Seed Sci Res 14: 335–344

1178 Mayzlish-Gati E, LekKala SP, Resnick N, Wininger S, Bhattacharya C, Lemcoff JH, Kapulnik Y, Koltai
1179   H (2010) Strigolactones are positive regulators of light-harvesting genes in tomato. J Exp Bot 61:
1180   3129–3136

1181 McGONIGLE TP, Miller MH, Evans DG, Fairchild GL, Swan JA (1990) A new method which gives an
1182   objective measure of colonization of roots by vesicular-arbuscular mycorrhizal fungi. New Phytol
1183   115: 495–501

1184 Miller FR (1984) Registration of RTx430 sorghum parental line. Crop Sci 24: 1224–1224

1185 Mohemed N, Charnikhova T, Fradin EF, Rienstra J, Babiker AGT, Bouwmeester HJ (2018) Genetic
1186   variation in Sorghum bicolor strigolactones and their role in resistance against Striga hermonthica. J
1187   Exp Bot 69: 2415–2430

1188 Morris GP, Harder AM, Healey AL, McLaughlin CM, Rifkin JL, Cruet-Burgos C, Jenkins JW, Shu S,
1189   Spiekerman JJ, VanGessel CJ, et al (2026) A sorghum pangenome reference improves global crop
1190   trait discovery. Nature. doi: 10.1038/s41586-026-10229-9

1191 Nasrollahi V, Yuan Z-C, Lu QSM, McDowell T, Kohalmi SE, Hannoufa A (2022) Deciphering the role
1192   of SPL12 and AGL6 from a genetic module that functions in nodulation and root regeneration in
1193   Medicago sativa. Plant Mol Biol 110: 511–529

1194 Newman EI (1966) A method of estimating the total length of root in a sample. J Appl Ecol 3: 139

1195 Ogawa S, Cui S, White ARF, Nelson DC, Yoshida S, Shirasu K (2022) Strigolactones are
1196   chemoattractants for host tropism in Orobanchaceae parasitic plants. Nat Commun 13: 4653

1197 Oksanen J, Simpson G, Blanchet FG, Kindt R, Legendre P, Minchin P, O'hara RB, Solymos P, Stevens
1198   H, Szöcs E, et al (2022) vegan community ecology package version 2.6-2 April 2022. The
1199   Comprehensive R Archive Network. Available online: http://cran. r-project. org (accessed on 15
1200   August 2022)

1201 Ó'Maoiléidigh DS, van Driel AD, Singh A, Sang Q, Le Bec N, Vincent C, de Olalla EBG, Vayssières A,
1202   Romera Branchat M, Severing E, et al (2021) Systematic analyses of the MIR172 family members of
1203   Arabidopsis define their distinct roles in regulation of APETALA2 during floral transition. PLoS
1204   Biol 19: e3001043

1205 Patel A, Kumar A, Sheoran N, Kumar M, Sahu KP, Ganeshan P, Ashajyothi M, Gopalakrishnan S, Gogoi
1206   R (2021) Antifungal and defense elicitor activities of pyrazines identified in endophytic
1207   Pseudomonas putida BP25 against fungal blast incited by Magnaporthe oryzae in rice. J Plant Dis
1208   Prot (2006) 128: 261–272

1209 Phillips JM, Hayman DS (1970) Improved procedures for clearing roots and staining parasitic and
1210   vesicular-arbuscular mycorrhizal fungi for rapid assessment of infection. Trans Br Mycol Soc 55:
1211   158–IN18

Poethig RS (2013) Vegetative phase change and shoot maturation in plants. Curr Top Dev Biol 105: 125–
152

Rasmussen A, Mason MG, De Cuyper C, Brewer PB, Herold S, Agusti J, Geelen D, Greb T,
Goormachtig S, Beeckman T, et al (2012) Strigolactones suppress adventitious rooting in
Arabidopsis and pea. Plant Physiol 158: 1976–1987

R Core Team (2022) R: A Language and Environment for Statistical Computing. R Foundation for
Statistical Computing, Vienna.

Riechmann JL, Meyerowitz EM (1998) The AP2/EREBP family of plant transcription factors. Biol Chem
379: 633–646

Ruyter-Spira C, Kohlen W, Charnikhova T, van Zeijl A, van Bezouwen L, de Ruijter N, Cardoso C,
Lopez-Raez JA, Matusova R, Bours R, et al (2011) Physiological effects of the synthetic
strigolactone analog GR24 on root system architecture in Arabidopsis: another belowground role for
strigolactones? Plant Physiol 155: 721–734

Saeed W, Naseem S, Ali Z (2017) Strigolactones biosynthesis and their role in abiotic stress resilience in
plants: A critical review. Front Plant Sci 8: 1487

Sandelin A, Alkema W, Engström P, Wasserman WW, Lenhard B (2004) JASPAR: an open-access
database for eukaryotic transcription factor binding profiles. Nucleic Acids Res 32: D91–4

Schägger H (2006) Tricine-SDS-PAGE. Nat Protoc 1: 16–22

Schmid R, Heuckeroth S, Korf A, Smirnov A, Myers O, Dyrlund TS, Bushuiev R, Murray KJ, Hoffmann
N, Lu M, et al (2023) Integrative analysis of multimodal mass spectrometry data in MZmine 3. Nat
Biotechnol 41: 447–449

Schneider CA, Rasband WS, Eliceiri KW (2012) NIH Image to ImageJ: 25 years of image analysis. Nat
Methods 9: 671–675

Seethepalli A, Dhakal K, Griffiths M, Guo H, Freschet GT, York LM (2021) RhizoVision Explorer:
open-source software for root image analysis and measurement standardization. AoB Plants 13:
lab056

Seethepalli A, York LM (2020) RhizoVision Explorer - Interactive software for generalized root image
analysis designed for everyone. doi: 10.5281/ZENODO.4095629

Seto Y, Sado A, Asami K, Hanada A, Umehara M, Akiyama K, Yamaguchi S (2014) Carlactone is an
endogenous biosynthetic precursor for strigolactones. Proc Natl Acad Sci U S A 111: 1640–1645

Sharkey TD, Bernacchi CJ, Farquhar GD, Singsaas EL (2007) Fitting photosynthetic carbon dioxide
response curves for C(3) leaves. Plant Cell Environ 30: 1035–1040

Shrestha JK, Lin C-Y, Wang JY, Tang I-C, Hu C-H, Tsednee M, Zhang Y, Jamil M, Berqdar L, Blilou I,
et al (2026) Carotenoid biosynthesis drives root plasticity through aerenchyma and iron plaque
formation in rice. Nat Plants 12: 179–190

Sorefan K, Booker J, Haurogné K, Goussot M, Bainbridge K, Foo E, Chatfield S, Ward S, Beveridge C,
Rameau C, et al (2003) MAX4 and RMS1 are orthologous dioxygenase-like genes that regulate
shoot branching in Arabidopsis and pea. Genes Dev 17: 1469–1474

Spallek T, Mutuku M, Shirasu K (2013) The genus Striga: a witch profile. Mol Plant Pathol 14: 861–869

Strock CF, Schneider HM, Galindo-Castañeda T, Hall BT, Van Gansbeke B, Mather DE, Roth MG,
Chilvers MI, Guo X, Brown K, et al (2019) Laser ablation tomography for visualization of root
colonization by edaphic organisms. J Exp Bot 70: 5327–5342

Sun H, Tao J, Liu S, Huang S, Chen S, Xie X, Yoneyama K, Zhang Y, Xu G (2014) Strigolactones are
involved in phosphate- and nitrate-deficiency-induced root development and auxin transport in rice.
J Exp Bot 65: 6735–6746

Tennant D (1975) A test of a modified line intersect method of estimating root length. J Ecol 63: 995

Van Ha C, Leyva-González MA, Osakabe Y, Tran UT, Nishiyama R, Watanabe Y, Tanaka M, Seki M,
Yamaguchi S, Van Dong N, et al (2014) Positive regulatory role of strigolactone in plant responses
to drought and salt stress. Proc Natl Acad Sci U S A 111: 851–856

Venables WN, Ripley BD (2002) Modern applied statistics with S. Springer, New York, NY

Vogel JT, Walter MH, Giavalisco P, Lytovchenko A, Kohlen W, Charnikhova T, Simkin AJ, Goulet C,
Strack D, Bouwmeester HJ, et al (2010) SlCCD7 controls strigolactone biosynthesis, shoot
branching and mycorrhiza-induced apocarotenoid formation in tomato. Plant J 61: 300–311

Walker AP, Beckerman AP, Gu L, Kattge J, Cernusak LA, Domingues TF, Scales JC, Wohlfahrt G,
Wullschleger SD, Woodward FI (2014) The relationship of leaf photosynthetic traits - V cmax and J
max - to leaf nitrogen, leaf phosphorus, and specific leaf area: a meta-analysis and modeling study.
Ecol Evol 4: 3218–3235

Walton A, Stes E, Goeminne G, Braem L, Vuylsteke M, Matthys C, De Cuyper C, Staes A,
Vandenbussche J, Boyer F-D, et al (2016) The response of the root proteome to the synthetic
strigolactone GR24 in Arabidopsis. Mol Cell Proteomics 15: 2744–2755

Wang JY, Chen G-TE, Jamil M, Braguy J, Sioud S, Liew KX, Balakrishna A, Al-Babili S (2022)
Protocol for characterizing strigolactones released by plant roots. STAR Protoc 3: 101352

Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J, Kapono CA,
Luzzatto-Knaan T, et al (2016) Sharing and community curation of mass spectrometry data with
Global Natural Products Social Molecular Networking. Nat Biotechnol 34: 828–837

Wang Y, Duran HGS, van Haarst JC, Schijlen EGWM, Ruyter-Spira C, Medema MH, Dong L,
Bouwmeester HJ (2021) The role of strigolactones in P deficiency induced transcriptional changes in
tomato roots. BMC Plant Biol 21: 349

Wu F, Gao Y, Yang W, Sui N, Zhu J (2022) Biological functions of strigolactones and their crosstalk
with other phytohormones. Front Plant Sci 13: 821563

Wu S, Li Y (2021) A unique sulfotransferase-involving strigolactone biosynthetic route in sorghum. Front
Plant Sci 12: 793459

Xu S, Chen M, Feng T, Zhan L, Zhou L, Yu G (2021) Use ggbreak to effectively utilize plotting space to
deal with large datasets and outliers. Front Genet 12: 774846

Xu Z-S, Chen M, Li L-C, Ma Y-Z (2011) Functions and application of the AP2/ERF transcription factor
family in crop improvement. J Integr Plant Biol 53: 570–585

Yoda A, Mori N, Akiyama K, Kikuchi M, Xie X, Miura K, Yoneyama K, Sato-Izawa K, Yamaguchi S,
Yoneyama K, et al (2021) Strigolactone biosynthesis catalyzed by cytochrome P450 and
sulfotransferase in sorghum. New Phytol 232: 1999–2010

Yoda A, Xie X, Yoneyama K, Miura K, McErlean CSP, Nomura T (2023) A stereoselective strigolactone
biosynthesis catalyzed by a 2-oxoglutarate-dependent dioxygenase in sorghum. Plant Cell Physiol
64: 1034–1045

Yoneyama K, Awad AA, Xie X, Yoneyama K, Takeuchi Y (2010) Strigolactones as germination
stimulants for root parasitic plants. Plant Cell Physiol 51: 1095–1103

Yoneyama K, Brewer PB (2021) Strigolactones, how are they synthesized to regulate plant growth and
development? Curr Opin Plant Biol 63: 102072

Yu N, Niu Q-W, Ng K-H, Chua N-H (2015) The role of miR156/SPLs modules in Arabidopsis lateral
root development. Plant J 83: 673–685

Zhang Z, Liao H, Lucas WJ (2014) Molecular mechanisms underlying phosphate sensing, signaling, and
adaptation in plants. J Integr Plant Biol 56: 192–220

Zhao J, Shi D, Kaeufer K, Song C, Both D, Thier AL, Cao H, Lassen L, Xu X, Hamamura Y, et al (2025)
Strigolactones optimise plant water usage by modulating vessel formation. Nat Commun 16: 3854

Zhou H, Akçay E, Helliker BR (2019) Estimating C4 photosynthesis parameters by fitting intensive A/Ci
curves. Photosynth Res 141: 181–194