Macrophage Immune-Competent Colon Assembloids for Functional Interrogation of Neuroinflammation-Induced Colonic Dysmotility

Claudia A. Collier[1], Karla I. Ortega Sandoval[1], Aelita Salikhova[1], Shrinarayanee Rengarajan[1], Anvitha Tharakesh[1], Alana Aristimuno Millan[1],Shanthi Srinivasan[2], and Shreya A. Raghavan*[1]

[1]Department of Biomedical Engineering, Texas A&M University
[2]Department of Medicine, Division of Digestive Diseases, Emory University

*Corresponding author:
Department of Biomedical Engineering at Texas A&M University
3120 TAMU
College Station, TX, 77843, United States
E-mail: sraghavan@tamu.edu

ETHICAL CONSIDERATIONS

This work was conducted in a laboratory approved by the Institutional Biosafety Committee at Texas A&M University. No animal use or institutional review board approval was required, since there is no human or vertebrate animal research.

FUNDING

This work was supported by the Department of Defense Gulf War Illness Research Program W81XWH-21-0477 (S.A.R.), Department of Defense Toxic Exposures Research Program HT9425-24-1-0998 (S.A.R.), Texas A&M University Avilés-Johnson Graduate Fellowship Program (C.A.C.), the National Defense Science and Engineering Graduate (NDSEG) Fellowship Program (C.A.C.) through the Army Research Office, and NIH R01 DK080684 (S.S.).

DATA TRANSPARENCY STATEMENT

Data supporting the findings of this study will be available from the corresponding author upon written request. RNA-Seq datasets are deposited on the NCBI Gene Expression Omnibus database, under ID: GSE316347.

REPORTING GUIDELINES: The data reported in this manuscript is not related to randomized trials, case reports or animal studies. Therefore, reporting guidelines are not applicable.

CONFLICTS OF INTEREST: The authors report no conflicting conflicts for interests.

AUTHOR CONTRIBUTIONS

Conceptualization: S.A.R.;SS Data curation: C.A.C., K.O.S; Formal analysis: C.A.C.,K.O.S, S.A.R.; Funding acquisition: S.A.R.; S.S; Investigation: C.A.C.,K.O.S, A.S.; Methodology: C.A.C., K.O.S, S.A.R.; Project administration: S.A.R.; Resources: S.A.R., S.S.; Software: C.A.C.; Supervision: S.A.R.; Validation: C.A.C., S.A.R.;

46 Visualization: C.A.C., K.O.S, A.S., A.V.,A.A.M, S.A.R.; Writing – original draft: C.A.C.,
47 S.A.R. ; Writing – review & editing: S.A.R, S.S.
48
49 **PREPRINT**: https://doi.org/10.1101/2025.08.07.669171
50
LIST OF ABBREVIATIONS
α-SMA – alpha smooth muscle actin
β-NGF – beta nerve growth factor
ACh - acetylcholine
BMP-2 – bone morphogenetic protein-2
CCL2, CCL11 – C-C-motif chemokine ligand 2/11
ChAT – Choline acetyltransferase
DAPI- 4,6- diamidino-2-phenylindole
ENS – enteric nervous system
FGID – functional gastrointestinal disorders
GWI – Gulf War Illness
iBMDM – immortalized murine bone marrow derived macrophages
IBD – inflammatory bowel disease
iCSMC – immortalized colonic smooth muscle cells
IM-FEN – immorto mouse fetal enteric neuron
IL-1β,6,10 - interleukin
NGFR – nerve growth factor receptor
nNOS/NOS1 – neuronal nitric oxide synthase
PB – pyridostigmine bromide
TIMP-1- tissue inhinitor of metalloproteinase
TNF-α - tumor necrosis factor alpha
TTX - tetrodotoxin

ABSTRACT

Functional gastrointestinal disorders (FGIDs) affect ~40% of the global population and are frequently characterized by colonic dysmotility. Symptomatic manifestations of colonic dysmotility significantly reduce quality of life in inflammatory bowel disease (IBD), diabetes, and Gulf War Illness (GWI). Current in vitro models lack the integration of functional physiology with immune and neuronal complexity required to establish causal links between neuroinflammation and dysmotility. Here, an immune-competent bioengineered colon assembloid is introduced that integrates multiple cell types of the external colonic wall, along with functional readouts of motility. Within bioengineered colon assembloids, various inflammatory insults resulted in enteric neuroinflammation, cascading to changes in colonic motility. Key mechanisms of dysmotility following inflammatory insult within the bioengineered colon assembloids included impaired neuronal regeneration, and aberrant smooth muscle remodeling. The bioengineered colon assembloid model mimicked diverse aspects of enteric neuroinflammation. Ultimately, the platform offers a physiologically relevant avenue to interrogate neuroimmune crosstalk and dissect mechanisms of colonic dysmotility, paving the way to new therapeutic strategies to improve colonic motility.

1. INTRODUCTION

Functional gastrointestinal disorders (FGIDs) are a significant global health burden[1], affecting an estimated 40% of people globally[2]. FGIDs occur in individuals with Inflammatory Bowel Disease (IBD)[3], Gulf War Illness (GWI)[4, 5], and even secondary to diabetes[6]. At the heart of FGIDs is altered colonic motility (colonic dysmotility). Yet, the mechanistic underpinnings of FGID-related colonic dysmotility are still poorly understood, and consequently, our ability to design targeted therapeutics to curtail it lag. This gap is partly due to the lack of nuanced in vitro model systems that can functionally integrate multi-cellular communication with motility-focused read outs.

The smooth muscle layers of the colonic muscularis externa integrate input from the enteric nervous system (ENS), tissue-resident immune cells like macrophages, and other interstitial cells to produce colonic motility[7]. Coordinated colonic motility occurs via reciprocal interactions between many of these individual cell types, with a beautifully complex regulatory network[7]. Chief among these are neuro-immune interactions, well studied in infection, inflammation, and mucosal barrier function[8,9]. From whole animal models, we deeply appreciate the evolving role of neuro-immune interactions in colonic motility, dysmotility and FGIDs[10,11], especially those between macrophages and enteric nerves[12].

Subtle alterations in the connectivity and biochemistry of the ENS impact colonic motility patterns[13]. Tolerogenic macrophages support the regeneration of damaged enteric nerves from a tissue-resident enteric neuronal progenitor pool, by clearing neuronal debris[14]. Activated pro-inflammatory macrophages release cytokines that impair neuronal regeneration and smooth muscle contraction[15, 16], thereby affecting colonic motility. Collectively, the presence of altered neuroplasticity, increased pro-inflammatory cytokines and altered colonic motility supports a model in which macrophages, smooth muscle cells of the colon, and enteric neurons interact to contribute to the symptoms of colonic dysmotility[12, 17, 18].

These foundational insights have been obtained from experimental mouse models, linking functional physiology to histological observations. A hallmark study demonstrated that macrophage-ENS interactions directed colonic motility via bone morphogenetic protein-2 (BMP-2) secretion[12]. Another experimental mouse model correlated the role of macrophages in myenteric plexus repair to intestinal dysmotility following surgical trauma[17]. Our own research with mouse models of Gulf War Illness demonstrated that toxic exposures reprogrammed macrophages and induced enteric neuronal progenitor cell apoptosis, leading to chronically altered colonic motility[19, 20]. Similar disruptions in neuroplasticity and altered colonic motility are routinely reported in mouse models of inflammatory bowel disease and FGIDs, often accompanied by increased pro-inflammatory cytokines[11, 18, 21]. However, establishing signal specificity and interrogating molecular mechanisms directing colonic motility independent of the microbiome and other systemic factors is tedious and challenging in experimental animal models. Herein, bioengineering and in vitro modeling offer a fantastic opportunity for tight variable isolation for mechanistic studies[22, 23].

Ongoing advances in organoid technologies[24], discovery and use of postnatal and adult human enteric neural progenitor cells[25], and gene editing have paved the way towards in vitro bioengineered systems capable of nuanced mechanistic dissection in enteric neuro-inflammation. Murine colonic organoids co-cultured with sensory neurons demonstrated a shared calcium influx pathway in macrophage-nerve interactions that may lead to pain in FGIDs[26]. Colonic organoids incorporating epithelial and immune components have been previously used to investigate immune and cytokine signaling in IBD[27]. Organoids, however, are largely epithelial-dominant systems with limited utility in studying neuro-immune crosstalk and motility. For example, co-culture of human intestinal organoids with induced pluripotent stem cell-derived neural progenitor cells showed neuroglial structures like the myenteric plexus. While this system demonstrated transient calcium waves, similar to neuronal activity that regulates colonic motility, it falls short of being able to quantify motility and contractility[28]. This is because organoids lack the active smooth muscle component of the outer layers of the colon that generate motility forces[29].

Assembloid systems on the other hand offer a construction strategy that enables multi-cellular structures within directed architectures[30, 31]. Colon assembloid systems have been successfully used to study stromal cell sustenance of epithelial crypts[32], and macrophage involvement in mucosal immunity[33]. Silk scaffold directed assembly of intestinal epithelium, myofibroblasts and differentiated enteric nerves enabled investigation of mucosal barrier function[34]. More recently, ENS-rich assembloids were used to study the dynamics and mechanisms of ENS regeneration and neural elongation during inflammation[35]. Despite these successes, current assembloid models also do not integrate functional readouts of colonic motility including spontaneous or evoked contractions in response to exogenous stimuli, thereby limiting their ability to capture how neuroinflammation drives colonic motility (and dysmotility).

The objective of the current study, therefore, was to create a bioengineered colon assembloid that integrated macrophage-ENS crosstalk with smooth muscle contractility. Our model guides the self-assembly of key cell types of the colonic wall that regulate colonic motility. These cell types include smooth muscle cells, macrophages, and an enteric neural circuit derived from enteric neuronal progenitor cells. This platform structurally mimics the outer colon layer and enables real-time assessment of tissue contractility. In this study, we show that bioengineered colon assembloids respond to inflammatory insults by altering macrophage activation, neural structure and composition, ultimately impacting colonic motility. Beyond its relevance to specific disease models, this system serves as a foundational platform for deeper mechanistic inquiry into neuroimmune processes driving colonic motility and dysmotility.

2. METHODS
2.1 Materials and reagents
All tissue culture media, and supplements were purchased from ThermoFisher Scientific (Waltham, MA, USA), unless otherwise specified. Growth factors and proteins were purchased from Peprotech (Cranbury, NJ, USA). All conjugated antibodies were purchased from Santa Cruz (Dallas, TX, USA), Cell Signaling Technology (Danvers, MA,

182 USA), or Abcam (Cambridge, MA, USA), unless specified otherwise. Pyridostigmine
183 bromide, Acetycholine (ACh), Tetrodotoxin (TTX) were purchased from Sigma Aldrich (St.
184 Louis, MO, USA).
185
2.2 Cell lines and culture
2.2.1 Isolation And Culture of Immortalized Colonic Smooth Muscle Cells (icSMCs)
Primary murine colonic smooth muscle cells were isolated from adult mouse colons aged
20-28 weeks. During dissection, the sigmoid colon was isolated via sharp dissection[19, 36,]
[37]. Dissected sigmoid colons were cleaned and prepped for enzymatic digestion. Briefly,
dissected colonic tissue was washed in sterile phosphate buffered saline, minced with
sharp scissors, and digested in type II collagenase. Digested cells were filtered serially
through a 70 and 40 µm mesh to remove debris, and plated in complete Dulbecco's
Modified Eagle Medium (DMEM) supplemented with 10% fetal bovine serum + 1%
penicillin and streptomycin. After initial expansion, cells were transduced with EF1α-SV40
large T antigen (SV40T) lentiviral particles (GeneCopoeia, puromycin-selectable;
Rockville, MD, USA) to establish a stable immortalized smooth muscle cell line (icSMC).
Successfully transduced cells were selected using puromycin and maintained at 37°C
with media changes every 48 hours.
200
201 *2.2.2 Culture Of Immortalized Bone Marrow Derived Macrophages*
202 Immortalized bone marrow derived macrophages[38] (iBMDMs) were a gift from Dr. Philip
203 West's laboratory at Jackson Labs, Bar Harbor ME. Cells were maintained in complete
204 DMEM supplemented with 10% fetal bovine serum + 1% antibiotic-antimycotic at 37°C
205 with 5% CO_2.
206
2.2.3 Culture Of Immortalized Fetal Enteric Neural Cells (IM-FENs):
The immortalized murine fetal enteric neuron (IM-FEN) cell line used in this study was
generated and characterized by the Srinivasan Laboratory at Emory University[39, 40].
Briefly, IM-FENs were derived from E13 H-2Kb-tsA58 immortomouse embryos. Neural
precursors were isolated using magnetic bead immunoselection targeting the low-affinity
NGF receptor (Ngfr-p75). IM-FENs were maintained in a modified N2 medium
supplemented with 10% fetal bovine serum, 10 ng/mL glial cell line-derived neurotrophic
factor (GDNF), 20 U/mL recombinant mouse interferon-γ, and 1% penicillin/streptomycin.
Cells were cultured at 33°C in a humidified incubator with 5% CO_2 to preserve their
progenitor state. To induce neuronal differentiation, cells were transitioned to a 1:1 ratio
of conditioned media collected from icSMCs and neurobasal-A medium supplemented
with B-27 (serum-free), 2 mmol/L L-glutamine, 1% heat-inactivated fetal bovine serum,
1% penicillin/streptomycin, and 10 ng/mL GDNF. Cultures were maintained at 39°C and
5% CO_2 for up to 7 days to promote differentiation into mature enteric neurons.
221
2.3 Bioengineered colon assembloid formation and characterization
2.3.1 Assembly
Bioengineered colon assembloids were fabricated using a two-layer collagen hydrogel
deposition technique to enable directed self-assembly within cell-laden hydrogels. Briefly,
35mm dishes were prepared by coating with a thin layer of non-adhesive polydimethyl
siloxane (PDMS; Sylgard® 184 silicone elastomer kit, Denver, CO, USA). The coating

also contained a central silicone post (3mm diameter) to define the hollow lumen.
Following curing over 48hrs, 35mm culture dishes had a non-adherent surface with a
defined central post. Around this central post, two sequential layers of collagen hydrogels
(1 mg/ml type I collagen; rat tail; ibidi; Madison, WI, USA) were patterned one after the
other.
The following cell types were uniformly suspended within the first base collagen gel layer:
(i) 400,000 cells/mL IM-FENs; (ii) 1,500 cells/mL iBMDMs; and (iii) 200,000 cells/mL
icSMCs. 300µl of cell-laden gel mixture was cast around the central post and allowed to
gel at 37°C for 15 minutes.
A second cell-laden collagen gel suspension was prepared using 600,000 cells/mL
iCSMCs only. Once gelation of the first base layer with all three cell types was complete,
300µL of cell-laden second layer was patterned around the first layer and the post. The
assembled structures were allowed to gel for an additional 30 minutes at 37°C. Following
gelation, 2 mL of iCSMC media was added to each dish, and constructs were incubated
at 37°C to promote self-assembly, alignment and compaction. After 24 hours, media was
supplemented with neuronal differentiation media for a final 1:1 ratio of iCSMC media:
neural differentiation media to support differentiation of enteric neuronal progenitor cells.
Formation of bioengineered colon assembloids was monitored via daily photography,
ensuring compaction and tight assembly around luminal post. Compaction of the gel area
was measured and monitored over 7 days.
248
2.3.2 Visualization of multi-cellular compartments within bioengineered colon
assembloids using whole mount immunofluorescence
Bioengineered colon assembloids were fixed in 4% paraformaldehyde for 30 minutes at
room temperature, followed by three washes in phosphate-buffered saline (PBS).
Assembloids were washed in 0.1M glycine for 20 minutes at room temperature followed
by permeabilizing and blocking for 2 hours in 0.2% Triton-X and 10% goat serum.
Fluorophore-conjugated antibodies targeting lineage-specific markers (**Supplementary**
Table 1) were diluted in blocking buffer and incubated overnight at 4°C. Following
incubation with antibodies, samples were then washed thoroughly, mounted between
coverslips and imaged using Olympus Fluoview FV3000 Confocal Laser Scanning
Microscope (RRID:SCR_021637). Each experimental condition was replicated across
multiple biological samples (n ≥ 5). Maximum intensity projections of the XY plane were
visualized to demonstrate overall structural organization of cell types within the
bioengineered colon assembloids.
263
2.4 Inflammatory Stimulation of Bioengineered Colon Assembloids
Bioengineered colon assembloids were established and allowed to fully form between
Days 1-3. Once compaction was achieved, inflammatory insult was initiated by adding
100µM Pyridostigmine Bromide (PB) or 100ng/mL Tumor Necrosis Factor-alpha (TNF-α)
into the culture media of assembloids for 24hrs. PB is a toxic exposure linked to
neuroinflammation and FGIDs in Gulf War Illness[19, 20], while TNF-α is a cytokine observed
in many FGIDs[41, 42]. Inflammatory insult was initiated at Day 3 and removed on Day 4.
271
2.4.1 Defining "acute" and "chronic" lingering effects of inflammation following one-time
inflammatory insult

An "acute" immediate effect was assessed by collecting assembloids at Day 4 for
downstream processing. Assembloids were also allowed to recover for 3 days following
one-time inflammatory insult (Days 4-7), based on previously documented studies
establishing that in vitro macrophage activation dynamics changed over 72hrs[43, 44]. Any
lingering "chronic" effects of one-time inflammatory insult were therefore assessed
following recovery, at Day 7. Time-matched untreated assembloids were maintained as
controls.
281
2.5 Characterization of the bioengineered colon assembloids following
inflammatory insult for acute and chronic effects
Following assembly (1-3 days), and inflammatory insult, colon assembloids were
processed for downstream analysis to investigate acute (Day 4) and chronic effects (Day
7) of inflammation. Analyses focused on targeted gene expression analysis, unbiased
RNA-Sequencing analysis, cytokine profiling and quantitative whole mount
immunofluorescence.
289
2.5.1 RNA extraction and gene expression analysis
RNA was extracted from cultured cells or assembloids using a RNeasy extraction kit
(Qiagen, Hilden, Germany). Bioengineered colon assembloids were first flash frozen in
liquid nitrogen, minced in cold TRIzol™ LS Reagent (Invitrogen, Carlsbad, CA, USA) and
extracted using the same kit following manufacture's protocol. RNA concentrations were
measured using a Nanodrop 2000 (ThermoFisher Scientific, Carlsbad, CA, USA)
spectrophotometer, transcribed to cDNA using the high-fidelity cDNA transcription kit (Life
Technologies, Carlsbad, CA, USA), and q-PCR was carried out in the 96-well format
using a QuantStudio 3(Applied Biosystems, Foster City, CA, USA). Gene expression
differences were quantified using the 2ΔΔCT method, using *Gapdh* as the housekeeping
control, and reported as fold changes compared to a control sample. qPCR experiments
were run in triplicates with 5 independent samples. Primer sets used are detailed in
Supplementary Table 2. Gene expression was normalized to time-matched control colon
assembloids that received no inflammatory insult. Fold changes in gene expression were
reported at acute (Day 4) or chronic (Day 7) time points due to PB or TNF-α treatment,
compared to time-matched untreated controls.
306
2.5.2 RNA Sequencing and analysis
RNA was isolated from PB or TNF-α treatment bioengineered colon assembloids, or
control untreated assembloids at Day 4 (acute; n=4) and Day 7 (chronic; n=3). Library
preparation with poly(A) selection and 150-bp paired-end sequencing on an Illumina
HiSeq 2500 were performed by Azenta (South Plainfield, NJ, USA). Raw sequencing data
in FASTQ format were generated by Azenta Life Sciences using an Illumina platform.
Azenta performed initial quality assessment, adapter trimming, alignment, and gene count
quantification. Specifically, adapter sequences and low-quality bases were trimmed using
Trimmomatic v0.36, and trimmed reads were aligned to the *Mus musculus* GRCm38
reference genome (Ensembl) using STAR aligner v2.5.2b. Gene-level quantification was
performed using featureCounts from the Subread package v1.5.2, generating a raw gene
count matrix for each sample.
319

320 All subsequent computational and statistical analyses were conducted in-house using
321 RStudio (2025.05.1+513) and R packages listed below. Differential gene expression
322 analysis (DGE) was performed using the DESeq2 package. Genes with p-value < 0.05
323 and absolute log2 fold change > 1.25 were defined as differentially expressed genes
324 (DEGs).
325
Volcano plots were generated using ggplot2 and dplyr, and 3D principal component
analysis (PCA) was performed using plotly and dplyr to visualize sample clustering based
on transcriptomic profiles. Venn diagrams were used to compare the number of
upregulated and downregulated DEGs across conditions relative to controls. Gene
ontology (GO) enrichment analysis was conducted using the EnRichGO, clusterProfiler,
org.Mm.eg.db, AnnotationDbi, and tidyverse packages. Biological process (BP) terms
were assessed for DEGs with log2FC > 1.25 and p-value < 0.05.
333
334
2.5.3 Cytokine profiling within bioengineered colon assembloids
Bioengineered colon assembloids were maintained as untreated controls, or subject to
inflammatory insult (PB or TNF-α) at Day 3 with removal at Day 4. "Acute" effects were
evaluated at Day 4, while "chronic" lingering effects were evaluated following 3 days of
recovery, at Day 7. Assembloids were flash frozen with liquid nitrogen and collected in a
RIPA+HALT solution. Supernatants were clarified by centrifugation at 500× g for 5
minutes, sterile filtered, and stored at −80°C until analysis. Protein concentration was
measured using a Pierce BCA assay (Thermo-Fisher) following manufacturer's protocols.
Cytokine levels were evaluated using bead-based multiplex immunoassays and done
according to the manufacturer's protocol (R&D Systems, Minneapolis, MN). Equal
amounts of protein/sample were used per assay, across acute and chronic time points
for each inflammatory insult, compared to untreated time-matched controls. Magnetic
bead washes and intermediary steps were performed using a Bio-Rad Bio-Plex Pro II
plate washer and final luminescence was read on a Bio-Rad Bio-Plex analyzer. Each
experimental condition was replicated across multiple biological samples (n ≥ 3). The
following cytokines were assayed: CCL2 (C-C motif chemokine ligand 2), IL-1β
(Interleukin-1 beta), IL-10 (Interleukin-10), TIMP-1 (Tissue inhibitor of metalloproteinases-
1), CCL11 (C-C motif chemokine ligand 11), β-NGF (beta-nerve growth factor), IL-6
(Interleukin-6).
354
355 *2.5.4 Evaluation of architectural changes within bioengineered colon assembloids with*
356 *inflammatory insult via whole mount immunofluorescence and quantification*
357
Assembloids were formed, exposed to inflammatory insult, and fixed and stained at
"acute" Day 4 or "chronic" Day 7 time points. Time-matched untreated controls were also
fixed and stained using methods outlined in Section *2.3.2*. Whole mount
immunofluorescence was performed to visualize macrophages and their activation via co-
staining for F480 (pan-macrophage marker), and either Cd40 (pro-inflammatory
activation marker) or Ym1 (tolerogenic macrophage marker). Nerves with stained with
pan-neuronal β-III Tubulin, with subtype specific staining using ChAT (choline
acetyltransferase) and neuronal nitric oxide synthase (Nos1). Enteric neuronal progenitor

cells were visualized with Ngfr-p75 staining, and smooth muscle structures with α-smooth muscle action (α-SMA). XY maximum intensity projections were obtained using the Olympus FluoView software, following consistent acquisition settings across all experimental conditions (control, PB, TNF-α). Quantitative fluorometry was performing using the corrected total fluorescence (CTF) method in FIJI/Image J using previously established protocols[19, 20, 45]. For each image, integrated density, area, and mean background fluorescence were measured for each ROI. CTF was calculated using the following formula:

CTF = Integrated Density − (Area of ROI × Mean Background Fluorescence)

To account for variations in cell number, CTF was divided by the number of nuclei (DAPI-positive cells) in each image, yielding an average CTF per cell. These CTF values were then normalized to the mean value from the control group to allow for relative comparison across conditions. Each experimental condition was replicated across multiple biological samples (n ≥ 5).

To assess macrophage activation, iBMDMs were co-stained for F480 (pan-macrophage marker) and Cd40 (pro-inflammatory activation marker) and imaged using standardized acquisition settings across all experimental groups. Quantification was performed using the same CTF method in ImageJ. For each image, Cd40 CTF was divided by the corresponding F480 CTF to generate a normalized Cd40/F480 fluorescence ratio, representing the degree of pro-inflammatory polarization relative to total macrophage content. Similar quantification was performed for Ym1/F480, to identify tolerogenic macrophage populations. Neuronal density changes with inflammatory insult were quantified via CTF of β-III Tubulin, ChAT, Nos1 and p75, while smooth muscle architectural changes were quantified via CTF of α-SMA. Each experimental condition was replicated across multiple biological samples (n ≥ 5).

2.6 Evaluation of contractility in bioengineered colon assembloids
Physiological contractility of bioengineered colon assembloids were assessed using a custom-built organ bath system[19, 20]. Each assembloid was mounted by securing one luminal end to a fixed reference post and the other to the arm of a force transducer (FT20, Harvard Apparatus). The organ bath was filled with DMEM buffered with 1 M HEPES and maintained at 37°C. A 30% passive stretch was applied using a micromanipulator and maintained throughout the experiment to record isometric force. Assembloids were equilibrated under these conditions for 40–60 minutes. Baseline force post-equilibration was arbitrarily set to zero to allow for direct comparison of pharmacologically induced changes in contractility.

To assess neural responsiveness and muscle function, assembloids were treated with the following agents: Acetylcholine (ACh, 1 µM/mL) to induce cholinergic contraction; Tetrodotoxin (TTX, 1 µM/mL) to reversibly inhibit neuronal sodium channels and block neuronal activity.

411   Pharmacologic agents were added directly to the bath, and contractile responses were
412   recorded. Force traces were analyzed for peak amplitude and area under the curve (AUC)
413   to quantify physiological responses. Each experimental condition was replicated across
414   multiple biological samples (n ≥ 5).
415
2.7 Evaluation of macrophage mono-culture exposure to inflammatory stimuli
To initiate inflammatory insult on macrophage mono-cultures, iBMDMs were seeded and
allowed to adhere overnight in densities ranging from 3,000 – 8,000 cells/mL. Following
adhesion, on Day 1, macrophages were exposed to 100µM PB or 100ng/mL TNF-α for
24 hours. iBMDMs were assessed at different time points: 'Acute' was collected
immediately after 24 hours of exposure (Day 2). To assess recovery following one-time,
24hr exposure to PB or TNF-α, removal of insult was followed by recovery in freshly
supplemented growth media. Recovery was assessed longitudinally at two time points: 3
days post removal, or 5 days post removal of PB or TNF-α. Both these recovery time
points were given the nomenclature of "Day 3 Chronic" to mean 3 days of recovery to
assess lingering effects of inflammation, or "Day 5 Chronic" to mean 5 days of recovery
to assess lingering effects of inflammation.
Macrophage activation in response to PB or TNF-α treatment was evaluated in multiple
ways: (1) Gene expression analysis following methods outlined in Section 2.5.1,
comparing time-matched untreated macrophages to PB or TNF-α treated macrophages.
(2) Immunofluorescent co-staining of macrophages with pan-macrophage marker F480
and inflammatory marker Cd40, with CTF based quantitative fluorometry; (3) Cytokine
profiling of cell culture supernatants following methods outlined in Section 2.5.3.
434
435   Media was also collected from iBMDMs following acute exposure to PB (Day 2), sterile
436   filtered and termed as 'PB-conditioned media (PB-CM)'. This media was used in future
437   experiments, to mimic an inflammatory insult to cultured enteric neuronal progenitor cells.
438
439   **2.8 Effect of inflammatory insult and prolonged macrophage-derived factors on**
440   **enteric neuronal progenitor cell (IM-FEN) proliferation and identity**
441
IM-FENs (8000 cells/mL) were allowed to adhere onto fibronectin-coated dishes overnight
at 33°C, prior to initiation of differentiation. Once cells were adhered, progenitor IM-FENs
were maintained at 33°C with no exposure (control) or constant inflammatory
macrophage-derived conditioned media (PB-CM) for 7 days. Cells were fixed, and
immunostained with the proliferation marker, Ki67 following immunofluorescence
protocols.
448
The primary antibody, Ki67 (Alexa Fluor 488) was used to mark active cell proliferation.
Ki67+ cells were identified by the presence of green fluorescence within DAPI (blue
fluorescence) counterstained nuclei and were quantified manually using previously
demonstrated protocols[46]. Analysis was performed to determine the number of actively
proliferating cells (Ki67 antigen in the nuclei) versus the total cell number (number of
nuclei) to determine the percentage of proliferating cells. Exposed cells were compared
to controls. Each experimental condition was replicated across multiple biological
samples (n ≥ 5).

IM-FENs were characterized using flow cytometry for Ngfr-p75 expression[47], by adapting
previously used flow analysis protocols[19, 20]. Briefly, IM-FENs were collected and
resuspended in a flow cytometry amenable buffer (PBS + 2% FCS) at a concentration of
200,000 cells/ml. Cells were incubated with a p75-AF647 antibody, with a matched
isotype control. Cells were then processed through a flow cytometer (Attune NxT,
Waltham, MA, USA). Flow cytometry analysis was completed with FlowJo (Ashland, OR).
Polygon gates were drawn on isotype flow cytometry graphs to establish a background
cut off gate at 0.5% (gating strategy is presented in **Supplementary Figure 6**). The same
gate was applied to antibody-stained cells to establish a percent positive population.
Each experimental condition was replicated across multiple biological samples (n ≥ 3).
468

2.9 Effect of inflammatory insult and prolonged macrophage-driven factors on IM-
FEN differentiation
IM-FENs (8000 cells/mL) were allowed to adhere onto fibronectin-coated dishes overnight
at 33°C, prior to initiation of differentiation. Once cells were adhered, enteric neuronal
progenitor cells were either maintained as untreated controls, or exposed to PB-CM in its
progenitor state prior to differentiation for 24hrs, initiating the inflammatory insult in the
progenitor cell state. Differentiation was initiated by increasing the temperature to 39°C
on Day 2, supplemented with neural differentiation media. Cells were differentiated as
untreated controls, or under the constant prolonged exposure of PB-CM over 5 additional
days up to Day 7. Ability or changes in differentiation with prolonged PB-CM exposure
was characterized using immunostaining for differentiated neuronal markers including β-
III Tubulin, ChAT, and Nos1 and CTF-based quantification. Comparisons were drawn
between untreated controls and PB-CM treated IM-FENs. Similar quantitative analyses
were also carried out using gene expression analysis, to assess changes in PB-CM
treated IM-FENs.
484

485   **3. RESULTS AND DISCUSSION**
486

487   **3.1 Formation of a bioengineered 3D colon assembloid incorporating neural,**
488   **smooth muscle, and immune lineages**
489

Colonic motility arises from the muscularis externa of the colon, the outer layers that
comprise smooth muscle cells, an integrated neuronal network and tissue-resident
macrophages. To study functional consequences of enteric neuro-inflammation on
colonic motility, we bioengineered the colonic muscularis externa. The bioengineered
assembloid combined three key cell types within two layers of collagen hydrogel: (i)
immortalized colonic smooth muscle cells (icSMCs) as the smooth muscle component
that generates the forces relevant to colonic motility; (ii) immortalized enteric neuronal
progenitor cells (IM-FENs) that differentiate within the bioengineered construct to form
mature neuronal networks; and (iii) immortalized macrophages (iBMDMs) to add immune
competency. The base inner layer contained all three cell types, and was cast loosely
around a luminal post to define the hollow lumen. A second outer layer contained smooth
muscle cells only, and enveloped the base layer and allowed to gel over 30 minutes.
Figure 1A outlines the sequential cell seeding process into the collagen hydrogel, and

subsequent self-assembly into concentric layered structures. Bioengineered colon assembloids matured over 1-7 days, enabling integrated neuro-immune structures and concentrically aligned smooth muscle architecture (**Figure 1B**). Compaction of the collagen hydrogel also occurred during the self-assembly process (>75% compaction over 3 days; **Figure 1C**).

Spatial and cellular maturation and organization within the bioengineered colon assembloids was confirmed via whole mount immunofluorescence (**Figure 1D**). By day 5, α-SMA[+] smooth muscle cells were concentrically aligned, similar to the structural organization of smooth muscle cells in the colon[48]. Differentiated neurons derived from neuronal progenitor cells robustly expressed βIII-tubulin[+], organized in reticular network-like structures similar to their organization within the myenteric plexus of the colon[49, 50]. Differentiated neuronal subtypes were further identified via stains specific to the synthesis of neurotransmitters, namely choline acetyltransferase (ChAT) and nitric oxide synthase (Nos1)[51]. Interestingly, our stains also demonstrated that bioengineered colon assembloids retained a population of neuronal progenitor cells in their progenitor state, evidenced by Ngfr-p75[+] staining, similar to tissue resident adult enteric neuronal stem cells present in the human and mouse colons[52, 53]. Macrophage presence was additionally verified by F4/80[+] expression interspersed within the bioengineered structure.

3.2 Bioengineered colon assembloids have distinct gene expression changes to stimulated neuro-inflammation

We next assessed the transcriptional consequences of inflammatory exposure in the bioengineered colon assembloids. We stimulated enteric neuro-inflammation using two different methods commonly associated with functional gastrointestinal disorders. The first stimulant was Pyridostigmine bromide (PB), a toxic exposure associated with Gulf War Illness[19, 20], resulting in severe colonic motility disruption. The second stimulant we used in this study was Tumor Necrosis Factor-alpha (TNF-α), central to enteric neuro-inflammation in inflammatory bowel diseases like Crohn's Disease[21], where colonic motility is similarly disrupted. **Figure 2A** displays a timeline for when inflammatory stimuli are added to bioengineered colon assembloids on Day 3 once their compaction and self-assembly is complete. PB or TNF-α stimulation occurred for 24hrs (Day 3-4), following which an immediate response to inflammation was assessed (acute; Day 4). Once inflammatory stimulants were removed, bioengineered colon assembloids were allowed to recover for 3 additional days, and any chronic lingering effects of inflammation was assessed (chronic; Day 7). The recovery time window of 3 days was chosen based on previous studies assessing the dynamics of macrophage activation[43, 44]. All comparisons were made to untreated control bioengineered colon assembloids that were time-matched in culture.

3.2.1 PB-induced gene expression changes in bioengineered colon assembloids mimic key aspects of those observed in Gulf War Illness mouse colons

Acute PB exposure significantly increased the gene expression of *Il6* (**3.68-fold**), with a subsequent drop to 2.62-fold chronically, indicating mild but incomplete recovery (**Figure 2B**). The elevated low-grade inflammation in bioengineered colon assembloids was in line with observations from mouse models of persistent enteric neuro-inflammation in Gulf

549   War Illness[20]. Markers of reparative macrophages, *Mrc1* and *Ch3l3*, were acutely
550   downregulated, with a partial restoration following recovery.
551
Acute PB exposure also resulted in a reduction of neuronal gene expression compared
to untreated bioengineered colon assembloids (**Figure 2C**). Specifically, *Tubb3* (**0.85-
fold; ****p<0.0001**), *Chat* (**0.68-fold; ****p<0.0001**), and *Nos1* (**0.73-fold; ****p<0.0001**)
were all significantly decreased, indicating impaired neuronal programming. Interestingly,
upon recovery, *Chat* expression trended towards significant upregulation (**2.97-fold;
****p<0.0001**). These findings in bioengineered colon assembloids also mimicked studies
in Gulf War Illness mouse models, where one-time PB exposure chronically destroyed
enteric neuronal integrity and altered neuronal subtype balance[20]. Neuronal recovery,
despite subtype imbalance, was supported by increased gene expression of *Ngfr-p75* and
Sox2 (**Figure 2D**), in line with adult enteric neo-neurogenesis following inflammatory
insult[54, 55], captured within bioengineered colon assembloids. Interestingly, PB exposure
did not alter the gene expression of smooth muscle markers (**Figure 2E)**, also in line with
our previous observations in mouse models[20].
565
3.2.2 TNF-α–induced gene expression changes are distinctly different compared to
PB-induced changes in bioengineered colon assembloids
Bioengineered colon assembloids exposed acutely to TNF-α exhibited strong pro-
inflammatory activation (**Figure 2F**), with significant upregulation of *Il6* (1.94-fold;
p<0.01). Reparative macrophage markers, *Mrc1* and *Ch3l3,* were reduced (0.32-** and
0.44-fold, ***p<0.001) compared to control, with no evident recovery in gene expression
levels. This is unsurprising, because TNF-α is a potent inflammatory mediator in
macrophages, including in inflammatory bowel disease[21].
574
Neuronal damage and mild recovery were observed in bioengineered rings with TNF-α
treatment, with acute downregulation of *Tubb3* and *Nos1,* but not *Chat* (**Figure 2G**).
Markers of neuronal stemness were more broadly suppressed with TNF-α treatment,
compared to PB, both acutely and chronically despite recovery. *Sox2* expression partially
recovered to 0.89-fold (******p<0.0001**) but remained significantly lower than control
(***p<0.05;   Figure   2H**). *Ngfr-p75* slightly   increased   to   0.78-fold   (****p<0.01**),
while *Nes* declined   further   to   0.52-fold   (******p<0.0001**)   compared   to   acute   rings,
demonstrating complex and marker-specific neural stem cell regulation over chronic TNF-
α induced inflammation.
584
Acute TNF-α exposure induced pronounced upregulation of smooth muscle contractile
genes *Acta2* (**2.27-fold;   ****p<0.0001**), *Cald1* (**1.75-fold; *p<0.05**),   and *Tagln* (**1.19-
fold; ns),** compared to control rings (**Figure 2I)**. In line with observations from
experimental mouse models of inflammatory bowel disease, TNF-α exposure significantly
alters expression of thin filament contractile elements within smooth muscle like
Caldesmon (*Cald1*) and Smoothelin (*Tagln),* that ultimately result in impaired colonic
contractility and motility[56, 57]. Bioengineered colon assembloids capture this important
nuance, where TNF-α treatment resulted in changes in the smooth muscle compartment
within the bioengineered colon, while PB did not.
594

3.3 Unbiased transcriptomics captures distinctions in PB and TNF-α induced enteric neuro-inflammation within bioengineered colon assembloids

In order to capture the differences in assembloid response to both acute and chronic PB or TNF-α stimulation, we performed three-dimensional principal component analysis (3D PCA) on bulk RNA-seq data from treated and untreated bioengineered colon assembloids. The acute 3D PCA plot (**Figure 2J**) illustrates distinct clustering of the three groups along principal components 1, 2, and 3, which collectively explain 68.9% of the total variance (PC1 = 52.7%, PC2 = 9.8%, PC3 = 6.4%). Samples from PB- and TNF-α treated groups separate clearly from controls, demonstrating transcriptomic alterations in response to inflammatory challenge within the bioengineered colon assembloids. These transcriptional distinctions persisted into the chronic phase, as shown in the chronic PCA plot (**Figure 2K**), where the three groups again clustered separately, with total variance explained reaching 70.4% (PC1 = 41.8%, PC2 = 16.8%, PC3 = 11.8%). This continued separation suggests lasting and distinct transcriptional remodeling despite removal of the initial stimulus.

The upregulated gene sets (**Figure 2L**) revealed both overlapping and distinct genes: a core subset of 91 genes was commonly upregulated by both PB and TNF-α, likely indicative of shared inflammatory pathways. Additionally, TNF-α uniquely upregulated 344 genes, while PB uniquely upregulated 133 genes, reflecting the operation of stimulus-specific transcriptional programs. Similarly, the downregulated gene sets (**Figure 2M**) revealed an overlap of 8 genes, with TNF-α driving broader suppression of 169 unique genes compared to PB's 9 unique genes. After recovery, chronic lingering effects of inflammation led to even more divergent transcriptional responses. A total of 408 genes were uniquely upregulated by PB, and 680 by TNF-α, with 66 genes commonly upregulated between the two treatments (**Figure 2N**). Similarly, 366 genes were uniquely downregulated by PB, 227 by TNF-α, and 53 genes were shared (**Figure 2O**).

3.4 Gene Ontology and biological process analyses reveal shared and distinct mechanisms of PB and TNF-α induced inflammation and impaired neurogenesis

Noting that PB and TNF-α exposure generated some distinct inflammatory signatures within bioengineered colon assembloids (differentially regulated genes also visualized as volcano plots in **Supplementary Figure 1A, B**), we first explored unbiased RNA-Seq analysis to further determine distinctions. **Supplementary Figure 1** also demonstrates distinct genes that drive unique but strong immunologic signatures upon PB and TNF-α exposure, with distinctions between the two exposures broken down further in **Supplementary Figure 2**. The involvement of shared and distinct gene signatures was further highlighted via unbiased gene ontology and biological process (GO; BP) enrichment analyses (**Figure 3A-D**). Both stimuli (PB and TNF-α) resulted in positive regulation and activation of innate immune response within the bioengineered colon assembloids. Within the colonic muscularis externa, macrophages are key regulators of immune response to inflammation and stress, resulting in changes in colonic motility and neuronal health[16]. Transcriptomic changes confirmed that macrophages included within

the assembloids were able to respond to inflammatory stimuli like PB or TNF-α, the functional consequences of which we explore later in the study.

Despite the seeming similarities of inflammation responses initiated by PB and TNF-α, a deeper investigation of enriched gene sets highlighted subtle differences in molecular mechanisms underlying the exposures (**Figure 3A-D, Supplementary Figure 3**). For each identified pathway, a z-score was calculated based on the proportion of significant genes within the pathway.

The toxic exposure-based stimulant PB resulted in pronounced interferon-gamma based inflammatory responses in both acute and chronic settings (**Figure 3A-B; Supplementary Figure 3A,B)**, in line with many observations in Gulf War veterans that have higher serum levels of interferon-gamma[58, 59]. In fact, interferon-gamma based neuro-inflammation is documented to drive many neurological disorders central to Gulf War Illness, priming the innate immune system to sustain neuro-inflammation chronically[60].

In addition to inflammation via interferon-gamma based response, acute TNF-α exposure also drove inflammation via NF-κB signaling, with stronger enrichment of cellular response to interleukin-1 (**Figure 3C-D**; **Supplementary Figure 2D, Supplementary Figure 3C**). TNF-α and interleukin-1 are elevated in functional gastrointestinal disorders and inflammatory bowel disease[61,62]. Interestingly, chronic TNF-α exposure strongly enriched chemotaxis (**Supplementary Figure 3D**), where persistent immune cell recruitment contributes to tissue remodeling and fibrosis[63]. These findings were also supported by changes in regulatory transcription factor-gene interaction maps (**Supplementary Figure 4**) between the two inflammatory stimuli.

Overall, unbiased transcriptomic analysis revealed that bioengineered colon assembloids were capable of responding to inflammatory stimuli, initiated either by PB exposure or by TNF-α exposure. Importantly, even after removal of the initial insult (chronic conditions), each exposure sustained slightly divergent signatures during recovery: PB-driven responses remained centered on interferon-gamma mediated neuroinflammation and impaired neurogenesis, while TNF-α shifted toward extracellular matrix remodeling and fibrotic pathways, mimicking key aspects of published work in GWI and IBD.

3.5 Cytokine profiling reveals distinct protein levels changes in bioengineered colon assembloids under neuro-inflammatory stimuli

Secreted cytokines are key orchestrators of tissue damage downstream of inflammatory initiation. Given the prevalent, yet subtly distinct, inflammatory gene signatures in bioengineered colon assembloids with PB and TNF-α exposure, we next investigated changes in a panel of secreted cytokines that amplify inflammation[64-68], mediate neural repair[69, 70] or are associated with tissue damage within the colonic muscularis externa[71]. PB and TNF-α treatment demonstrated partially distinct profiles (**Figure 4**).

3.5.1 Single PB exposure results in chronic low-grade neuro-inflammation and pathological tissue remodeling

Acute exposure to PB triggered a robust pro-inflammatory response in bioengineered colon assembloids, characterized by elevated CCL2, IL-1β, and IL-6 (***p<0.001; **Figure 4A; 'Acute'**) and decreased IL-10 (****p<0.0001). Cytokine production was also in-line with identified GO processes identified within bioengineered colon assembloids enriched following PB-exposure (**Supplementary Figure 3A**). This mirrors neuro-inflammatory profiles reported in the muscularis externa of GWI-affected colons[19, 20].

Following removal of PB and subsequent recovery, the lingering inflammatory environment in bioengineered colons assembloids evolved (noted as 'Chronic' in **Figure 4**): TIMP-1 and CCL11 levels increased (****p<0.0001), reflecting pathological tissue remodeling commonly seen within the inflamed colon[72, 73]. Low-grade, unresolved inflammation persisted through elevated IL-6 and IL-1β, along with low IL-10 levels, similar to GWI[74]. Additionally, persistent elevation of β-NGF (**p<0.01) was indicative of neuroinflammation driven altered neuronal plasticity[75]. Together, these findings showed that PB exposure induced a chronic low-grade inflammatory state in the macrophage-immune competent bioengineered colon assembloids.

3.5.2 TNF-α induced secretome partially reflects inflammation resolution with time

Acute TNF-α exposure prompted pro-inflammatory cytokine secretion (IL-1β, IL-6) in bioengineered colon assembloids (**Figure 4B; 'Acute'**). Interestingly, TNF-α (but not PB) immediately increased TIMP-1, a sign of acute pathological tissue remodeling. Following the removal of TNF-α, bioengineered colon assembloids progressed partially towards recovery (**Figure 4B; 'Chronic'**), with plummeting levels of IL-1β, IL-6 and TIMP-1. Mild signatures of neuro-inflammation were still persistent, including a potential recovery, with elevated CCL11 and β-NGF.

Overall, bioengineered colon assembloids continued to capture key aspects of inflammatory cytokine landscapes and remodeling processes similar to that observed in the colon, that contribute to motility dysfunction within the muscularis externa[76, 77].

3.6 Wholemount immunostaining reveals structural changes in bioengineered colon assembloids with neuro-inflammatory stimulation

Next, we used whole mount immunostaining to visualize structural changes that resulted from secreted factors and gene expression associated with neuro-inflammation, tissue remodeling, and neural plasticity. As the likely responder and orchestrater of inflammation, we first assayed macrophage numbers and phenotypes within the bioengineered colon assembloids. Overall neuronal network integrity was assessed using βIII-tubulin, with cholinergic subtypes visualized via ChAT (choline acetyltransferase), and inhibitory nitrergic subtypes using Nos1 (neuronal nitric oxide synthase). The continued presence of neuronal progenitor cells was also assayed, since these progenitor cells contribute to neuronal repair. Lastly, the effectors of motility, the smooth muscle cells were identified for their arrangement using α-smooth muscle actin to additionally note

tissue structure integrity changes (**Figure 5**). Changes were compared between control untreated and PB or TNF-α stimulated bioengineered colon assembloids.

3.6.1 PB exposure drives sustained inflammation with impaired neural regeneration

Inflammatory Cd40[+] macrophages significantly increased with PB-treatment, rising 15% acutely and staying elevated by 37% in chronic conditions compared to time-matched untreated controls (***p<0.05; Figure 5A-C**). No changes were observed in reparative Ym1[+] macrophages, consistent with its associated gene expression of *Ch3l3* (**Figure 2B**) and relatively stable levels of IL-10 (**Figure 4A**).

Acute and sustained inflammation lead to neuronal degeneration[19, 78, 79]. Structural damage of the neuronal network was evident with visual changes in βIII-tubulin organization (84% reduction with PB, compared to untreated controls; **Figure 5A-C**). Cholinergic and nitrergic neuronal subtypes were also reduced (79% and 73% respectively; ***p<0.05; Figure 5B, C**). Acute cholinergic nerve loss is unsurprising following PB treatment, since PB reversibly inhibits acetylcholine esterase, thereby directly targeting cholinergic nerves[80]. Sustained loss of neuronal integrity despite recovery ('Chronic' in **Figure 5B-C**) reflects a compromised neuronal repair system[81]. Consistent with this notion, PB treatment reduced neuronal progenitor cells by 87% (Ngfr-75; ***p<0.05; Figure 5A-C**) with no recovery, contributing to the disarrayed neuronal network.

Smooth muscle alignment and structural integrity largely remained unchanged, with only minor fluctuations (9% increase acutely, 3% decrease chronically; **Figure 5A-C**), in line with gene expression data (**Figure 2E**). This stability suggested that smooth muscle structural integrity was preserved despite neuroimmune disruption.

3.6.2 TNF-α exposure drives distinct neuronal changes along with reactive smooth muscle hypertrophy

Similar to PB-exposure, TNF-α also increased inflammatory Cd40[+] macrophages in bioengineered colon assembloids (23% acutely and 37% chronically compared to untreated controls; ***p<0.05; Figure 5D-F**). Unchanging reparative Ym1 dynamics continued to suggest an impaired resolution of inflammation in the timeframes the bioengineered colon assembloids were examined.

TNF-α exposure also contributed to acute disintegration of neural network integrity (69% loss in βIII-tubulin fluorescence, ***p<0.05, Figure 5D-F**). While cholinergic ChAT[+] neurons were not affected with TNF-α like PB, a severe loss of nitrergic Nos1[+] neurons occurred. The loss in nitrergic nerves is in line with therapeutic evidence that anti-TNF monoclonal antibodies aid in robust nitrergic neuronal recovery in experimental IBD models[82]. Interestingly, following recovery from TNF-α exposure, neuronal network integrity and ChAT[+] neurons were fully restored (**'Chronic' TNF-α, Figure 5E**). This mild neurotrophic effect of TNF-α on enteric neurons, especially in co-culture systems where smooth muscle cells are present has been observed previously[83], and replicated within bioengineered colon assembloids.

777 As expected, α-SMA expression increased acutely by 67% (**p<0.01**), consistent with
778 massive increases in secreted TIMP1, and reactive smooth muscle hypertrophy observed
779 in IBD[84, 85]. This contrasts with the stable α-SMA levels seen following PB exposure.
780 During recovery, α-SMA expression returned to baseline, indicating a resolution of the
781 hypertrophic response over time.
782
783 **3.7 Inflammatory exposures impair colonic motility in a 3D bioengineered colon**
784 **model**
785
Altered colonic motility is a hallmark of both GWI[86] and IBD[87, 88], contributing significantly
to functional gastrointestinal disorders and reduced quality of life[89-91]. Coordinated
contractile function is a complex interplay of homeostatic macrophage immune activation,
structural integrity of smooth muscle cells, intact neuronal architecture and balanced
neuronal subtypes. In response to inflammatory stimuli in bioengineered colon
assembloids, we observed pervasive changes in inflammatory macrophage activation
(Cd40 enrichment in Figure 5; cytokine secretome in Figure 4; genotype in Figures 2-3),
the structure of smooth muscle arrangement (like those observed in Figure 5E), and
neuronal damage and imbalance in neurochemical coding (seen as imbalanced
ChAT/Nos neurons in Figure 5). We hypothesized that these changes will alter colonic
motility in PB or TNF-α exposed colon assembloids compared to control, untreated colon
assembloids. To this end, we assessed contractile behavior in bioengineered colon
assembloids using a force transducer-based organ bath system[19, 20, 92]. Bioengineered
colon assembloids were stabilized in organ baths, and contractions were recorded
following exogenous stimuli designed to interrogate both neuronal and smooth muscle
contributions. Representative contractility traces from control and inflamed assembloids
(PB- and TNF-α exposed) are shown in **Figure 6**, with time on the X-axis and force
amplitude relative to baseline on the Y-axis.
804
3.7.1 Inflammation causes persistent impairment of colonic contractile function
mediated by cholinergic neuronal activity
Contractile function was assessed by exogenous addition of acetylcholine (ACh; timing
indicated by arrows in **Figure 6**). To isolate neuronal functionality in addition to the direct
effect of ACh on smooth muscle cells, bioengineered colon assembloids were pre-treated
with a nerve blocker, tetrodotoxin (TTX; green dotted traces in **Figure 6**). Control
untreated bioengineered colon assembloids displayed robust ACh-induced contractions
with an average peak force of 132.0 ± 0.8 µN (black trace, **Figure 6A, C**). Pretreatment
with TTX significantly reduced contractile amplitude by 67% (green dotted trace,
******p<0.0001; Figure 6B**), confirming the presence of a functional neuronal component
capable of mediating contractions within bioengineered colon assembloids[92].
816
Both PB and TNF-α treatment reduced ACh-induced contractile magnitude (53% lower in
PB; 44% lower in TNF-α; ******p<0.0001, ***p<0.001; Figure 6B, C**). Interestingly, acute
structural loss of cholinergic nerves with PB treatment (but not TNF-α) manifested
functionally as muted sensitivity to TTX pre-treatment (only 23% sensitive in PB-exposed
assembloids, compared to 67% sensitivity in untreated controls and 64% sensitivity in
TNF-α treated assembloids). Following removal of PB and recovery, contractile

dysfunction persisted in chronic PB-treated assembloids (**Figure 6D-F**), similar to loss of
contractility in GWI colons[19]. Only partial recovery of contractility was observed in chronic
TNF-α treated assembloids (***p<0.001; **p<0.01**; **Figure 6E, F**), despite a recovery of
structural neuronal integrity. The broader colonic dysmotility is further supported by
immunostaining and gene expression data, where an imbalance in excitatory and
inhibitory neuronal populations is evident, which can impair coordinated motility[20, 93, 94].
Especially in the TNF-α treated colons, a further contributing factor to diminished
contractility is the reactive changes in smooth muscle that can alter muscle
responsiveness, limiting contractile efficiency[95, 96]. Future testing will expand to different
pharmacological agents capable of testing neuronal connectivity and muscarinic receptor
sensitivity, for deeper characterization of the impact of neuro-inflammation on colonic
contractility.
835
836 **3.8 Inflammatory secretome from PB-exposed macrophages disrupts neural stem**
837 **cell maintenance and impairs neuronal differentiation**
838
While the 3D bioengineered colon assembloid model captured key aspects of the
complex interactions between neuro-inflammation and motility, the presence of multiple
interacting cell types and extracellular matrix components makes it difficult for finer
mechanistic dissection. To disentangle the direct effects of inflammatory stimuli, such as
PB and TNF-α, we first employed a simplified macrophage monoculture system.
Macrophages treated with PB for 24hrs in cell culture released cytokines, creating a
soluble inflammatory milieu supporting low-grade inflammation, notably with elevated IL-
6 (**Supplementary Figure 6**). This inflammatory milieu was captured in the form of
conditioned media derived from macrophages that were treated with PB for 24hrs (PB-
CM). This resolution observed in macrophage monocultures contrasts with the persistent
inflammation seen within 3D bioengineered colon assembloids, where complex
multicellular interactions and extracellular matrix components likely sustain pro-
inflammatory signaling and impede full recovery. Thus, while macrophages have an
intrinsic capacity to resolve inflammation after PB exposure in vitro, the 3D
microenvironment introduces additional regulatory factors that complicate resolution,
highlighting the importance of studying inflammation in physiologically relevant contexts.
855
Next, we employed a monoculture model of immortalized p75[+] enteric neuronal progenitor
cells (IM-FENs)[39], which possess the ability to differentiate into motor neuronal
subtypes[92]. First, we directly exposed IM-FENs to PB for 24hrs to assess their intrinsic
sensitivity to inflammatory insult (**Supplementary Figure 6**). While PB treatment induced
some changes in proliferation and viability, the overall effects of PB on differentiation
capability were modest. One-time exposure to PB in IM-FENs did not significantly alter
the differentiation of IM-FENs into cholinergic or nitrergic neurons (**Supplementary**
Figure 6F-H). These results implied that even though PB treatment induced a loss in the
progenitor pool, differentiation capacity was only modestly impacted in monocultures.
865
866 However, our previous observations in mouse models of PB exposure[20] and
867 bioengineered colon assembloids (**Figure 5,6**) demonstrated significant damage in
868 neuronal structural integrity resulting in functional dysmotility, implying that multi-cellular

interactions may drive neuronal progenitor damage. Indeed, from PB mouse models, we observed inflammatory Cd40 macrophages in close proximity to p75+ enteric neuronal progenitor cells within the colonic myenteric plexus[20]. Therefore, we hypothesized that impairment in enteric neuronal progenitor differentiation might be macrophage mediated. To test this hypothesis, we next applied conditioned media collected from PB-treated macrophages (PB-CM; iBMDMs) to IM-FEN cultures (**Figure 7A**). Macrophages were only treated with PB for 24hrs, and media was collected acutely right away following exposure. IM-FENs were chronically exposed to PB-CM conditioned medium, to simulate ongoing low-grade enteric neuro-inflammation consistently.

3.8.1 PB-CM reduces enteric neuronal progenitor proliferation and depletes progenitor identity

Ki67 immunostaining revealed a marked reduction in proliferative IM-FENs following continuous exposure to PB-CM (**Figure 7B**), compared to untreated controls. Quantitative analysis showed a 31% decrease in Ki67$^+$ cells at day 5, and 28% at day 7 in PB-CM treated cultures relative to untreated controls (****p<0.01; Figure 7C**), suggesting a potential impairment in self-renewal ability. This reduced proliferative capacity was accompanied by an 82% decline in Ngfr-p75$^+$ cells (a marker of stem-like identity) (******p<0.0001; Figure 7D**), indicating that prolonged exposure to inflammatory CM severely compromises neural progenitor maintenance. The gating strategy for all flow analysis is provided in **Supplementary Figure 7.** Gene expression analysis supported these findings, showing significant downregulation of neural stem cell markers *Ngfr-p75* (**0.36-fold**)*, Nestin (***0.94-fold**), and *Sox2 (***0.38-fold***) in PB-CM treated IM-FENs (******p<0.0001; Figure 7E**), consistent with loss of stem-like phenotype.

3.8.2 Inflammatory PB-CM impairs neuronal differentiation of IM-FENs

Neural progenitor cells are essential for maintaining productive neuroplasticity in response to inflammation. To understand how inflammation affects this regenerative potential in neuronal progenitor cells, we evaluated neuronal differentiation in the presence of PB-CM. When stimulated to differentiate with exogenously added factors, neuronal progenitor cells demonstrated an impaired ability to differentiate with reduction in βIII-tubulin, ChAT, and Nos1 in PB-CM treated cultures compared to controls (**Figure 7F, G**). This loss of neuronal markers was supported by gene expression analysis, which revealed 49% and 52% reductions in *Chat* and *Nos1* expression (*****p<0.001; Figure 7H**). These results reveal a marked impairment in neurogenesis under PB-mediated inflammation, characterized by loss of neuronal identity and differentiation capacity.

These findings support our hypothesis that PB-induced, macrophage-mediated inflammation impairs enteric neural repair and contributes to dysmotility. In PB-exposed bioengineered colon assembloids, we observed sustained inflammation, and activation of cell death pathways, as evidenced by transcriptomic profiling (**Figures 3**) and elevated levels of inflammatory cytokines (**Figure 4**), alongside major neuronal losses and inflammatory macrophage accumulation (**Figure 5**). Collectively, this suggests that PB exposure initiates a strong macrophage-driven inflammatory response that persists beyond the initial insult, leading to long-term disruption of neurogenesis and neural circuit

integrity. This chronic neuroinflammation (despite PB removal) likely explains the loss of
functional motility observed in the PB-exposed bioengineered colon assembloid.

CONCLUSIONS
Through our work, we have demonstrated the fabrication of a multi-cellular bioengineered
colon assembloid, that structurally and functionally integrates key cell types of the colonic
muscularis externa: smooth muscle cells, macrophages and an enteric neural circuit. The
platform enables real-time quantitative assessment of colonic contractility, directly linking
neuro-immune interactions to functional contractility outputs. Our results demonstrate the
bioengineered colon assembloid is sensitive to inflammatory perturbations, inducing
measurable changes in macrophage activation, enteric neuronal plasticity, and ultimately,
colonic contractility. The platform also provides a tightly controlled environment for
isolating cellular variables to interrogate molecular mechanisms in the future to
investigate how neuro-inflammation contributes to colonic dysmotility in FGIDs.

FIGURE LEGENDS
Figure 1. *Multicellular 3D bioengineered colon assembloid integrating smooth muscle,*
neural, and macrophage immune cells. A) Schematic of the sequential collagen casting
and cell seeding procedure. Layer 1 contained immortalized murine bone marrow-derived
macrophages (iBMDM), immortalized murine fetal enteric neural progenitor cells (IM-
FEN), and immortalized murine colonic smooth muscle cells (icSMC). These cells were
embedded in an inner collagen gel, cast around a silicone post placed to define a hollow
lumen. Once the first gel layer formed (~15minutes), a second collagen layer was cast,
laden with icSMCs alone (outer collagen gel). Constructs were cultured for 7 days at 37°C
to self-assemble into a ring-like structure with a defined hollow lumen. B) Representative
photographs of bioengineered colon assembloids during directed layer-by-layer
assembly, and every day from days 1-7. A central post defining the hollow luminal space
is still visualized in these photographs, with the bioengineered circular colon assembloid
around it. Scale bar = 15 mm (C) Quantification of bioengineered colon assembloid area
over time with progressive compaction. ****p<0.0001, ordinary one-way ANOVA, n>14.
(D) Wholemount immunofluorescence staining at days 5 and 7 demonstrating cellular
markers: α-SMA (α-smooth muscle actin), βIII-tubulin (neuronal tubulin), ChAT (choline
acetyltransferase), and Nos1 (neuronal nitric oxide synthase), Ngfr-p75 (receptor for the
neural growth factor p75 labeling neural progenitor cells), F4/80 (pan-macrophage
marker). Scale bar = 20 μm.

Figure 2. *Transcriptomic changes in bioengineered colon assembloids induced by*
exposure to Pyridostigmine bromide (PB) or Tumor necrosis factor (TNF-α).
A) Experimental timeline demonstrating timeline of inflammatory insult initiation at Day 3,
and removal at Day 4; Acute effects were evaluated on Day 4; Chronic lingering effects
of inflammation were evaluated after 3 days of recovery, on Day 7. (B–E) Gene
expression analysis in PB-exposed bioengineered colon assembloids for macrophage
(*Il6, Ch3l3, Mrc1*) (B), enteric neuron (*Tubb3, Chat, Nos1*) (C), neural stem cells (*Ngfr-*
p75, Nes, Sox2) (D), and smooth muscle (*Acta2, Tagln, Cald1*) (E) markers, comparing
acute and chronic conditions to untreated control. (F–I) Corresponding gene expression
changes in TNF-α exposed rings for macrophage (F), enteric neuron (G), neural stem cell
(H), and smooth muscle (I) markers. The dotted line at 1 represents the normalized level

of gene expression for time-matched control colon assembloids. Data are presented as
mean ± SEM. ****p<0.0001, ***p<0.001, **p<0.01, *p<0.05, 'ns' = not significant, one
sample t-test, n>3. (J, K) Three-dimensional principal component analysis (PCA) of bulk
RNA-seq showing clustering of control (brown), PB-treated (green), and TNF-α treated
(blue) bioengineered colon assembloids under acute (n=4; J) and chronic (n=3; K)
conditions. Each dot represents an individual biological replicate. % values indicate the
proportion of total variance within each principal component. (L, M) Venn diagram
comparing overlapping of significantly upregulated (L) and downregulated (M) genes
between acute PB and TNF-α exposure. (N, O) Venn diagram comparing overlapping of
significantly upregulated (N) and downregulated (O) genes in lingering chronic PB and
TNF-α induced inflammation.

Figure 3. *Transcriptomic profiling and biological process enrichment following exposure*
to PB and TNF-α in bioengineered colon assembloids. (A, B) Top unbiased Gene
Ontology (GO) Biological Processes (BP) significantly enriched in PB-treated
assembloids acutely (A) and chronic effects of one-time PB treatment (B). (C, D) Top
unbiased GO BP significantly enriched in assembloids treated with TNF-α and assessed
acutely (C) and for chronic effects (D). Bars represent the –log10 adjusted p-value for
each GO term.

Figure 4. *Cytokine profiling of bioengineered colon assembloids following exposure to*
PB and TNF-α. (A, B) Multiplex Luminex quantification of inflammatory and regulatory
cytokines from bioengineered colon assembloids under untreated control, or PB or TNF-
α treatment for 24hrs. Treatment was assessed either acutely (immediately upon 24hr
exposure), or lingering chronic effects were assessed following a 3-day recovery after
initial inflammatory insult. Analyzed cytokines include: CCL2 (monocyte chemoattractant
protein-1), IL-1β (interleukin-1 beta), IL-10 (interleukin-10), TIMP-1 (tissue inhibitor of
metalloproteinases-1), CCL11 (eotaxin), β-NGF (beta-nerve growth factor), and IL-6
(interleukin-6). A) Cytokine levels following PB treatment in bioengineered colon
assembloids. B) Cytokine levels following TNF-α treatment in bioengineered colon
asembloids. Data are presented as mean ± SEM. ****p<0.0001, ***p<0.001, **p<0.01,
*p<0.05, 'ns' = not significant; ordinary one-way ANOVA; n>3.

Figure 5. *Wholemount immunostaining in bioengineered colon assembloids as untreated*
controls or following PB or TNF-α exposure. (A–C) Representative immunofluorescence
images of control (A) and PB-exposed (B) bioengineered colon assembloids at acute
(24hrs following stimulation) and chronic (3 days of recovery after initial inflammatory
stimuli) timepoints. Scale bar = 20 μm. (C) Quantification of macrophage phenotypes
(F4/80, yellow; Cd40, red; Ym1, red), neuronal subtypes (βIII-tubulin, ChAT, Nos1- all
green), neural progenitors (Ngfr-p75, green), and smooth muscle (α-SMA, red) in control
and PB-treated bioengineered colons under acute (24-hr exposure) and chronic (3-day
recovery) conditions. (D–F) Representative immunofluorescence images of control
untreated (D) and TNF-α (E) treated bioengineered colon assembloids at acute and
chronic timepoints. Scale bar = 20 μm. (F) Quantification of control and TNF-α treated
rings at acute and chronic time-points. Data are presented as mean ± SEM. ****p<0.0001,
***p<0.001, **p<0.01, *p<0.05, 'ns' = not significant; ordinary one-way ANOVA; n>5.

Figure 6. *Inflammatory exposures impair neuronally mediated contractility in 3D bioengineered colon assembloids.* Bioengineered colon assembloids were mounted in a force transducer-fitted organ bath, and contractile force was measured in response to exogenous addition of acetylcholine (ACh). ACh addition is indicated by the blue arrow, after the establishment of a stable baseline of force (arbitrarily set to zero). (A) Representative traces in response to 1 μM ACh (black traces) stimulation in control untreated or acute exposure to PB-, or TNF-α. Neuronal contributions were isolated by pre-treatment with 1 μM tetrodotoxin (TTX, blue dotted lines. (B) Quantification of contractile output as area under the curve for various treatment conditions under acute exposure. Data are presented as mean ± SEM. ****$p<0.0001$, ***$p<0.001$, **$p<0.01$, 'ns' = not significant; two-way ANOVA; $n>5$. (C) Quantification of contractile output as peak magnitude of force generated following ACh stimulation in acute conditions. Data are presented as mean ± SEM. ****$p<0.0001$, **$p<0.01$, *$p<0.05$, 'ns' = not significant; two-way ANOVA; $n>5$. (D) Representative traces in response to ACh stimulation in chronic bioengineered colons, maintained as untreated controls or 3 days recovered from PB or TNF-α treatment. (E) Quantification of contractile output as area under the curve for various treatment conditions under chronic exposure. Data are presented as mean ± SEM. ***$p<0.001$, **$p<0.01$, 'ns' = not significant; two-way ANOVA; $n>3$. (F) Quantification of contractile output as peak magnitude of force generated following ACh stimulation in chronic conditions. Data are presented as mean ± SEM. **$p<0.01$, *$p<0.05$, 'ns' = not significant; two-way ANOVA; $n>3$.

Figure 7. *Inflammatory secretome from PB-exposed macrophages disrupts IM-FEN proliferation and neurogenic potential.* (A) Schematic of experimental timeline for treating IM-FENs with conditioned media collected from PB-exposed macrophages (PB-CM). Macrophages were exposed to PB for 24hrs, and conditioned media was collected immediately following exposure and stored (PB-CM). Enteric neuronal progenitor cells were adhered to culture surfaces, and treated with PB-CM prior to differentiation on Day 1. Cultures of IM-FENs were maintained in undifferentiated progenitor states to evaluate proliferation and stem cell identity. Cultures of IM-FENs were differentiated beginning at Day 2, for 5 days leading up to Day 7 in the presence or absence of PB-CM, to analyze their differentiation capability. (B) Representative immunofluorescence images showing Ki67 (green), a marker of proliferating cells, in untreated and PB-CM treated IM-FEN cultures at the indicated timepoints. Nuclei are counterstained with DAPI (blue). Scale bars = 15 μm. (C) Quantification of Ki67$^+$ proliferating cells over 7 days. Data are presented as mean ± SEM. ****$p<0.0001$, **$p<0.01$, two-way ANOVA; $n>5$. (D) Flow cytometry analysis of neural progenitor marker Ngfr-p75 expression in control and PB-CM treated IM-FENs. Data are presented as mean ± SEM. ****$p<0.0001$, unpaired t-test; $n>9$. (E) Relative gene expression of neural stem markers *(Ngfr-p75, Nestin, Sox2)* in PB-CM treated IM-FENs, normalized to untreated controls (dotted line = 1). Data are presented as mean ± SEM. ****$p<0.0001$, 'ns' = not significant; two-way ANOVA; $n>4$. (F, G) Untreated or PB-CM treated IM-FENs were differentiated into several neuronal subtypes. (F) Representative immunofluorescence images showing expression of neuronal markers: βIII-tubulin (green), ChAT (red), and Nos1 (magenta) in control and PB-CM treated differentiated IM-FENs. Nuclei are counterstained with DAPI (blue). Scale bars = 20 μm. (G) Quantification of differentiated neuronal populations expressing βIII-

tubulin, ChAT, and Nos1. Data are presented as mean ± SEM. ****p<0.0001, ***p<0.001; unpaired t-test; n>5. (H) Heatmap displaying relative gene expression fold changes of *Tubb3*, *Chat*, and *Nos1* in untreated and PB-CM treated differentiated IM-FENs, normalized to undifferentiated controls.

REFERENCES

1. Sperber AD, Bangdiwala SI, Drossman DA, et al. Worldwide Prevalence and Burden of Functional Gastrointestinal Disorders, Results of Rome Foundation Global Study. Gastroenterology 2021;160:99-114.e3.
2. Fikree A, Byrne P. Management of functional gastrointestinal disorders. Clin Med (Lond) 2021;21:44-52.
3. Walker C, Boland A, Carroll A, et al. Concurrent functional gastrointestinal disorders in patients with inflammatory bowel disease. Frontiers in Gastroenterology 2022;Volume 1 - 2022.
4. Collier CA, Salikhova A, Sabir S, et al. Crisis in the gut: navigating gastrointestinal challenges in Gulf War Illness with bioengineering. Military Medical Research 2024;11:45.
5. Johnson AG, Hu Z, Cost AA. Incidence and recent trends in functional gastrointestinal disorders, active component, US Armed Forces, 2005-2014. MSMR 2016;23:10-15.
6. Fagherazzi G, Gusto G, Balkau B, et al. Functional gastrointestinal disorders and incidence of type 2 diabetes: Evidence from the E3N–EPIC cohort study. Diabetes & Metabolism 2016;42:178-183.
7. Hou X, Artis D. Neuro-immune cell interactions in the regulation of intestinal immune homeostasis. Curr Opin Gastroenterol 2025;41:38-45.
8. Wallrapp A, Chiu IM. Neuroimmune Interactions in the Intestine. Annu Rev Immunol 2024;42:489-519.
9. Kulkarni S, Kurapati S, Bogunovic M. Neuro-innate immune interactions in gut mucosal immunity. Curr Opin Immunol 2021;68:64-71.
10. Lakhan SE, Kirchgessner A. Neuroinflammation in inflammatory bowel disease. Journal of Neuroinflammation 2010;7:37.
11. Akiho H, Ihara E, Motomura Y, et al. Cytokine-induced alterations of gastrointestinal motility in gastrointestinal disorders. World J Gastrointest Pathophysiol 2011;2:72-81.
12. Muller PA, Koscsó B, Rajani GM, et al. Crosstalk between muscularis macrophages and enteric neurons regulates gastrointestinal motility. Cell 2014;158:300-313.
13. Niesler B, Kuerten S, Demir IE, et al. Disorders of the enteric nervous system - a holistic view. Nat Rev Gastroenterol Hepatol 2021;18:393-410.
14. Kulkarni S, Micci MA, Leser J, et al. Adult enteric nervous system in health is maintained by a dynamic balance between neuronal apoptosis and neurogenesis. Proc Natl Acad Sci U S A 2017;114:E3709-E3718.
15. Wang H, Foong JPP, Harris NL, et al. Enteric neuroimmune interactions coordinate intestinal responses in health and disease. Mucosal Immunology 2022;15:27-39.
16. De Schepper S, Stakenborg N, Matteoli G, et al. Muscularis macrophages: Key players in intestinal homeostasis and disease. Cellular Immunology 2018;330:142-150.
17. Breßer M, Siemens KD, Schneider L, et al. Macrophage-induced enteric neurodegeneration leads to motility impairment during gut inflammation. EMBO Mol Med 2025;17:301-335.
18. Akiho H, Ihara E, Nakamura K. Low-grade inflammation plays a pivotal role in gastrointestinal dysfunction in irritable bowel syndrome. World J Gastrointest Pathophysiol 2010;1:97-105.
19. Collier CA, Foncerrada S, Clevenger AJ, et al. Acute Exposure to Pyridostigmine Bromide Disrupts Cholinergic Myenteric Neuroimmune Function in Mice. Advanced Biology 2023;7:2200254.
20. Collier CA, Salikhova A, Sabir S, et al. Persistent enteric neuroinflammation chronically impairs colonic motility in a pyridostigmine bromide-induced mouse model of Gulf War Illness. Biology Open 2025.
21. Spear ET, Mawe GM. Enteric neuroplasticity and dysmotility in inflammatory disease: key players and possible therapeutic targets. Am J Physiol Gastrointest Liver Physiol 2019;317:G853-g861.
22. Llorente C. The Imperative for Innovative Enteric Nervous System-Intestinal Organoid Co-Culture Models: Transforming GI Disease Modeling and Treatment. Cells 2024;13.
23. Collier CA, Mendiondo C, Raghavan S. Tissue engineering of the gastrointestinal tract: the historic path to translation. Journal of Biological Engineering 2022;16.

24. Sarvestani SK, Signs S, Hu B, et al. Induced organoids derived from patients with ulcerative colitis
recapitulate colitic reactivity. Nat Commun 2021;12:262.
25. Schmitteckert S, Mederer T, Roth R, et al. Postnatal human enteric neurospheres show a
remarkable molecular complexity. Neurogastroenterol Motil 2019;31:e13674.
26. Zhang Z, Engel MA, Koch E, et al. Menthacarin induces calcium ion influx in sensory neurons,
macrophages and colonic organoids of mice. Life Sci 2021;264:118682.
27. Hibiya S, Tsuchiya K, Hayashi R, et al. Long-term Inflammation Transforms Intestinal Epithelial
Cells of Colonic Organoids. J Crohns Colitis 2017;11:621-630.
28. Workman MJ, Mahe MM, Trisno S, et al. Engineered human pluripotent-stem-cell-derived intestinal
tissues with a functional enteric nervous system. Nat Med 2017;23:49-59.
29. Flood P, Hanrahan N, Nally K, et al. Human intestinal organoids: Modeling gastrointestinal
physiology and immunopathology — current applications and limitations. European Journal of
Immunology 2024;54:2250248.
30. Luo J, Meng J, Gu Z, et al. Topography-Induced Cell Self-Organization from Simple to Complex
Aggregates. Small 2019;15:e1900030.
31. Zhu T, Hu Y, Cui H, et al. 3D Multispheroid Assembly Strategies towards Tissue Engineering and
Disease Modeling. Adv Healthc Mater 2024;13:e2400957.
32. Lin M, Hartl K, Heuberger J, et al. Establishment of gastrointestinal assembloids to study the
interplay between epithelial crypts and their mesenchymal niche. Nat Commun 2023;14:3025.
33. Noel G, Baetz NW, Staab JF, et al. A primary human macrophage-enteroid co-culture model to
investigate mucosal gut physiology and host-pathogen interactions. Sci Rep 2017;7:45270.
34. Manousiouthakis E, Chen Y, Cairns DM, et al. Bioengineered in vitro enteric nervous system. J
Tissue Eng Regen Med 2019;13:1712-1723.
35. Sari I, Uzun B, Oguz MO, et al. Self-Organizing Assembloids Reveal Enteric Nervous System
Dynamics in Gut Homeostasis and Regeneration. bioRxiv 2025:2025.01.14.632538.
36. Hashish M, Raghavan S, Somara S, et al. Surgical implantation of a bioengineered internal anal
sphincter. J Pediatr Surg 2010;45:52-8.
37. Raghavan S, Miyasaka EA, Hashish M, et al. Successful implantation of physiologically functional
bioengineered mouse internal anal sphincter. Am J Physiol Gastrointest Liver Physiol
2010;299:G430-9.
38. Spera I, Sánchez-Rodríguez R, Favia M, et al. The J2-Immortalized Murine Macrophage Cell Line
Displays Phenotypical and Metabolic Features of Primary BMDMs in Their M1 and M2 Polarization
State. Cancers (Basel) 2021;13.
39. Anitha M, Joseph I, Ding X, et al. Characterization of fetal and postnatal enteric neuronal cell lines
with improvement in intestinal neural function. Gastroenterology 2008;134:1424-1435.
40. Balasubramaniam A, Li G, Ramanathan A, et al. SIRT3 activation promotes enteric neurons
survival and differentiation. Scientific reports 2022;12:22076.
41. Liebregts T, Adam B, Bredack C, et al. Immune activation in patients with irritable bowel syndrome.
Gastroenterology 2007;132:913-920.
42. Billiet T, Paul R, Marc F, et al. Targeting TNF-α for the treatment of inflammatory bowel disease.
Expert Opinion on Biological Therapy 2014;14:75-101.
43. Chau SM, Herting SM, Noltensmeyer DA, et al. Macrophage activation in response to shape
memory polymer foam-coated aneurysm occlusion devices. J Biomed Mater Res B Appl Biomater
2022;110:1535-1544.
44. Donehoo DA, Collier CA, VandenHeuvel SN, et al. Degrees of macrophage-facilitated healing in
aneurysm occlusion devices. J Biomed Mater Res B Appl Biomater 2024;112:e35385.
45. Bankhead P. Analyzing fluorescence microscopy images with ImageJ, 2014.
46. Clevenger AJ, McFarlin MK, Collier CA, et al. Peristalsis-Associated Mechanotransduction Drives
Malignant Progression of Colorectal Cancer. Cellular and Molecular Bioengineering 2023;16:261-
281.
47. Metzger M, Bareiss PM, Danker T, et al. Expansion and differentiation of neural progenitors derived
from the human adult enteric nervous system. Gastroenterology 2009;137:2063-2073. e4.
48. Huycke TR, Miller BM, Gill HK, et al. Genetic and Mechanical Regulation of Intestinal Smooth
Muscle Development. Cell 2019;179:90-105.e21.
49. Sharkey KA, Mawe GM. The enteric nervous system. Physiological Reviews 2023;103:1487-1564.

50. Kigata T, Koyama T, Lu X, et al. Quantitative analysis of the distribution of the myenteric plexus in
the rabbit large intestine. Anatomia, Histologia, Embryologia 2023;52:1029-1033.
51. Furness JB. Types of neurons in the enteric nervous system. Journal of the Autonomic Nervous
System 2000;81:87-96.
52. Heanue TA, Pachnis V. Prospective Identification and Isolation of Enteric Nervous System
Progenitors Using Sox2. Stem Cells 2011;29:128-140.
53. Metzger M, Bareiss PM, Danker T, et al. Expansion and Differentiation of Neural Progenitors
Derived From the Human Adult Enteric Nervous System. Gastroenterology 2009;137:2063-
2073.e4.
54. Belkind-Gerson J, Hotta R, Nagy N, et al. Colitis Induces Enteric Neurogenesis Through a 5-HT4–
dependent Mechanism. Inflammatory Bowel Diseases 2015;21:870-878.
55. Belkind-Gerson J, Graham HK, Reynolds J, et al. Colitis promotes neuronal differentiation of Sox2+
and PLP1+ enteric cells. Scientific reports 2017;7:2525.
56. Nair DG, Miller KG, Lourenssen SR, et al. Inflammatory cytokines promote growth of intestinal
smooth muscle cells by induced expression of PDGF-Rβ. Journal of Cellular and Molecular
Medicine 2014;18:444-454.
57. Alkahtani R, Mahavadi S, Al-Shboul O, et al. Changes in the expression of smooth muscle
contractile proteins in TNBS- and DSS-induced colitis in mice. Inflammation 2013;36:1304-15.
58. Khaiboullina SF, DeMeirleir KL, Rawat S, et al. Cytokine expression provides clues to the
pathophysiology of Gulf War illness and myalgic encephalomyelitis. Cytokine 2015;72:1-8.
59. Kalahasty K, Lee Y, Locatelli E, et al. Ocular and inflammatory markers associated with Gulf War
illness symptoms. Scientific Reports 2023;13:3512.
60. Trageser KJ, Sebastian-Valverde M, Naughton SX, et al. The Innate Immune System and
Inflammatory Priming: Potential Mechanistic Factors in Mood Disorders and Gulf War Illness.
Frontiers in Psychiatry 2020;Volume 11 - 2020.
61. Souza RF, Caetano MAF, Magalhães HIR, et al. Study of tumor necrosis factor receptor in the
inflammatory bowel disease. World J Gastroenterol 2023;29:2733-2746.
62. Vulliemoz M, Brand S, Juillerat P, et al. TNF-Alpha Blockers in Inflammatory Bowel Diseases:
Practical Recommendations and a User's Guide: An Update. Digestion 2020;101:16-26.
63. Speca S, Giusti I, Rieder F, et al. Cellular and molecular mechanisms of intestinal fibrosis. World J
Gastroenterol 2012;18:3635-61.
64. Popivanova BK, Kostadinova FI, Furuichi K, et al. Blockade of a Chemokine, CCL2, Reduces
Chronic Colitis-Associated Carcinogenesis in Mice. Cancer Research 2009;69:7884-7892.
65. Leal RF, Planell N, Kajekar R, et al. Identification of inflammatory mediators in patients with Crohn's
disease unresponsive to anti-TNFα therapy. Gut 2015;64:233-242.
66. Tagore A, Gonsalkorale W, Pravica V, et al. Interleukin-10 (IL-10) genotypes in inflammatory bowel
disease. Tissue antigens 1999;54:386-390.
67. Polosukhina D, Singh K, Asim M, et al. CCL11 exacerbates colitis and inflammation-associated
colon tumorigenesis. Oncogene 2021;40:6540-6546.
68. Mudter J, Neurath MF. Il-6 signaling in inflammatory bowel disease: pathophysiological role and
clinical relevance. Inflammatory bowel diseases 2007;13:1016-1023.
69. Woolf CJ, Safieh-Garabedian B, Ma QP, et al. Nerve growth factor contributes to the generation of
inflammatory sensory hypersensitivity. Neuroscience 1994;62:327-31.
70. Rai MF, Cai L, Chinzei N, et al. Distinct patterns of cytokines, chemokines, and growth factors in
synovial fluid after ACL injury in comparison to osteoarthritis. Journal of Orthopaedic Research
2024;42:1448-1462.
71. Kapsoritakis AN, Kapsoritaki AI, Davidi IP, et al. Imbalance of tissue inhibitors of metalloproteinases
(TIMP) – 1 and – 4 serum levels, in patients with inflammatory bowel disease. BMC
Gastroenterology 2008;8:55.
72. Mimura T, Bateman AC, Lee RL, et al. Up-regulation of collagen and tissue inhibitors of matrix
metalloproteinase in colonic diverticular disease. Dis Colon Rectum 2004;47:371-8; discussion
378-9.
73. Polosukhina D, Singh K, Asim M, et al. CCL11 exacerbates colitis and inflammation-associated
colon tumorigenesis. Oncogene 2021;40:6540-6546.
74. Whistler T, Fletcher MA, Lonergan W, et al. Impaired immune function in Gulf War Illness. BMC
Medical Genomics 2009;2:12.

75. Longange B. Defining the Neurochemical Phenotype of Myenteric Neurons that Display Nerve
Growth Factor Promotor Activity in Adult Transgenic Mice - Implications for Colonic Inflammation.
Canada -- Ontario, CA: Queen's University (Canada), 2022:66.
76. Shi X-Z, Winston JH, Sarna SK. Differential immune and genetic responses in rat models of Crohn's
colitis and ulcerative colitis. American Journal of Physiology-Gastrointestinal and Liver Physiology
2011;300:G41-G51.
77. Vrees MD, Pricolo VE, Potenti FM, et al. Abnormal Motility in Patients With Ulcerative Colitis: The
Role of Inflammatory Cytokines. Archives of Surgery 2002;137:439-446.
78. Winston JH, Li Q, Sarna SK. Paradoxical regulation of ChAT and nNOS expression in animal
models of Crohn's colitis and ulcerative colitis. American Journal of Physiology-Gastrointestinal and
Liver Physiology 2013;305:G295-G302.
79. Becker L, Nguyen L, Gill J, et al. Age-dependent shift in macrophage polarisation causes
inflammation-mediated degeneration of enteric nervous system. Gut 2018;67:827-836.
80. Candiotti K. A primer on nerve agents: what the emergency responder, anesthesiologist, and
intensivist needs to know. Can J Anaesth 2017;64:1059-1070.
81. Grundmann D, Loris E, Maas-Omlor S, et al. Enteric Neurogenesis During Life Span Under
Physiological and Pathophysiological Conditions. Anat Rec (Hoboken) 2019;302:1345-1353.
82. Souza RF, Machado FA, Caetano MAF, et al. Effect of Anti-TNF Monoclonal Antibody on Enteric
Neurons and Enteric Glial Cells in Experimental Colitis. Dig Dis Sci 2025;70:1375-1394.
83. Gougeon PY, Lourenssen S, Han TY, et al. The pro-inflammatory cytokines IL-1beta and TNFalpha
are neurotrophic for enteric neurons. J Neurosci 2013;33:3339-51.
84. Veisman I, Massey WJ, Goren I, et al. Muscular hyperplasia in Crohn's disease strictures: through
thick and thin. American Journal of Physiology-Cell Physiology 2024;327:C671-C683.
85. Chen W, Lu C, Hirota C, et al. Smooth muscle hyperplasia/hypertrophy is the most prominent
histological change in Crohn's fibrostenosing bowel strictures: a semiquantitative analysis by using
a novel histological grading scheme. Journal of Crohn's and Colitis 2017;11:92-104.
86. Wang WF, Guo XX, Yang YS. Gastrointestinal problems in modern wars: clinical features and
possible mechanisms. Mil Med Res 2015;2:15.
87. Bassotti G, Antonelli E, Villanacci V, et al. Colonic motility in ulcerative colitis. United European
Gastroenterol J 2014;2:457-62.
88. da Silva Watanabe P, Cavichioli AM, de Lima Mendes JDA, et al. Colonic motility adjustments in
acute and chronic DSS-induced colitis. Life Sciences 2023;321:121642.
89. Ansari R, Attari F, Razjouyan H, et al. Ulcerative colitis and irritable bowel syndrome: relationships
with quality of life. European journal of gastroenterology & hepatology 2008;20:46-50.
90. Piche T, Ducrotté P, Sabate J, et al. Impact of functional bowel symptoms on quality of life and
fatigue in quiescent Crohn disease and irritable bowel syndrome. Neurogastroenterology & Motility
2010;22:626-e174.
91. Trivedi KH, Schlett CD, Tribble DR, et al. The impact of post-infectious functional gastrointestinal
disorders and symptoms on the health-related quality of life of US military personnel returning from
deployment to the Middle East. Digestive diseases and sciences 2011;56:3602-3609.
92. Raghavan S, Gilmont RR, Miyasaka EA, et al. Successful implantation of bioengineered,
intrinsically innervated, human internal anal sphincter. Gastroenterology 2011;141:310-9.
93. Li S, Fei G, Fang X, et al. Changes in Enteric Neurons of Small Intestine in a Rat Model of Irritable
Bowel Syndrome with Diarrhea. J Neurogastroenterol Motil 2016;22:310-20.
94. Takahashi T. Pathophysiological significance of neuronal nitric oxide synthase in the
gastrointestinal tract. J Gastroenterol 2003;38:421-30.
95. Wells R, Blennerhassett M. Persistent and selective effects of inflammation on smooth muscle cell
contractility in rat colitis. Pflügers Archiv 2004;448:515-524.
96. MacDonald J. Smooth muscle phenotypic plasticity in mechanical obstruction of the small intestine.
Neurogastroenterology & Motility 2008;20:737-740.
1268

ACUTE

- response to virus
- defense response to virus
- regulation of innate immune response
- response to interferon-beta
- positive regulation of innate immune response
- cellular response to interferon-beta
- negative regulation of response to biotic stimulus
- negative regulation of innate immune response
- cellular response to type II interferon
- defense response to bacterium
- antigen processing and presentation of exogenous antigen
- antigen processing and presentation
- defense response to protozoan
- antigen processing and presentation of exogenous peptide antigen
- negative regulation of immune response
- negative regulation of viral process
- antigen processing and presentation of peptide antigen
- response to protozoan

CHRONIC

- response to type II interferon
- regulation of innate immune response
- cellular response to type II interferon
- response to interferon-beta
- cellular response to interferon-beta
- defense response to virus
- antigen processing and presentation
- response to virus
- response to protozoan
- positive regulation of response to biotic stimulus
- cell killing
- defense response to bacterium
- antigen processing and presentation of exogenous antigen
- positive regulation of innate immune response
- antigen processing and presentation of peptide antigen
- negative regulation of response to biotic stimulus
- positive regulation of leukocyte activation
- antigen processing and presentation of exogenous peptide antigen
- positive regulation of lymphocyte activation
- positive regulation of cell activation

ACUTE

- regulation of innate immune response
- positive regulation of response to biotic stimulus
- positive regulation of innate immune response
- response to type II interferon
- cellular response to molecule of bacterial origin
- response to molecule of bacterial origin
- cellular response to lipopolysaccharide
- cellular response to biotic stimulus
- response to lipopolysaccharide
- response to virus
- defense response to bacterium
- cellular response to type II interferon
- cytokine-mediated signaling pathway
- immune response-activating signaling pathway
- immune response-regulating signaling pathway
- defense response to virus
- activation of innate immune response
- regulation of inflammatory response
- canonical NF-kappaB signal transduction
- regulation of tumor necrosis factor superfamily cytokine production

CHRONIC

- extracellular structure organization
- extracellular matrix organization
- external encapsulating structure organization
- regulation of peptidyl-tyrosine phosphorylation
- ossification
- positive regulation of peptidyl-tyrosine phosphorylation
- peptidyl-tyrosine phosphorylation
- peptidyl-tyrosine modification
- leukocyte migration
- bone development
- tissue remodeling
- blood
- chemotaxis
- myeloid leukocyte migration
- cell chemotaxis
- gland morphogenesis
- morphogenesis of a branching structure
- reproductive structure development
- morphogenesis of a branching epithelium
- reproductive system development