Noc1 Downregulation Induces Nucleolar Stress and Upregulates p53 Isoforms, with a Robust Increase of the Truncated p53E Isoform in Drosophila Wing Discs.

Andrea Vutera Cuda[1*], Shivani Bajaj[1*], Valeria Manara[1], and Paola Bellosta[1,2,**]

[1] Department of Cellular, Computational and Integrative Biology (CIBIO),
University of Trento, Trento, TN 38123, Italy.

[2] Department of Medicine, NYU Langone Medical Center, New York, NY 10016, United States.

* co-first authorship

** correspondence: paola.bellosta@unitn.it

Running title: p53 Isoform Upregulation in Noc1 Nucleolar Stress.

Key words: p53 isoforms, nucleolar stress, Noc1, apoptosis, γ-H2AV, primer specificity, isoform primer design, mutant screen, *Drosophila* imaginal discs.

Abstract

Disruption of ribosome biogenesis triggers nucleolar stress, a conserved cellular response that activates p53. We previously demonstrated that depletion of Nucleolar Complex Protein 1 (Noc1) in Drosophila wing imaginal discs impairs rRNA maturation and ribosome assembly, resulting in elevated p53 levels and apoptosis, hallmarks of nucleolar stress. The Drosophila p53 gene produces four mRNA isoforms, yet their individual contributions to nucleolar stress responses remain poorly understood. Using newly designed isoform-specific qPCR primers, we found that although all p53 isoforms exhibit moderate transcriptional changes following Noc1 reduction, the truncated isoform p53E is robustly and preferentially upregulated. Notably, p53E lacks the N-terminal transactivation domain and has been reported to negatively regulate p53-induced apoptosis in specific tissues. Furthermore, our analyses indicate that γ-H2AV accumulation arises from caspase-dependent apoptosis rather than primary genomic lesions, suggesting the activation of a p53-dependent stress pathway distinct from canonical genotoxic pathways. Together, these findings suggest that p53E may be part of a novel mechanism activated during nucleolar stress, providing insight into how cells adapt to defects in ribosome biogenesis.

Introduction

Nucleolar stress is a cellular response to unbalanced ribosomal biogenesis (Boukoura and Larsen 2024). The nucleolus acts as a key stress sensor, responding to perturbations in ribosomal RNA (rRNA) processing and ribosome biogenesis by inducing the upregulation of p53 (Lindstrom et al. 2022).

Drosophila p53 exhibits significant structural and functional similarities with human p53 (Brodsky et al. 2000; Ingaramo et al. 2018; Ollmann et al. 2000; Bourdon et al. 2005). The *p53* gene (FBgn0039044) maps to chromosome 3R and encodes four mRNA isoforms (*p53A, p53B, p53C,* and *p53E*). According to the current deep RNA sequencing-based genome annotation, these isoforms encode three distinct proteins (Boley et al. 2014). Notably, *p53A,* also known as ∆Np53, and *p53C* differ in their 5'

untranslated regions (5'UTRs), and while the protein product of p53C remains

unconfirmed, its open reading frame (ORF) is predicted to be identical to that of p53A,

and to encode a 44 KDa protein (Zhang et al. 2015; Chakravarti et al. 2022). This

suggests that differential promoter regulation underlies their distinct functions in different

cellular contexts. *p53B* is the longer isoform, carrying a unique N-terminal domain and

encoding for a 56 KDa protein considered the closest homolog to human p53 (TP53)

(Ingaramo et al. 2018). *p53E* is the shorter isoform that encodes a predicted 38 kDa

protein lacking part of the N-terminal transactivation domain, and is considered to act as

a dominant negative in specific tissues and stress conditions (Wylie et al. 2022).

p53A, and p53B are the most characterized protein isoforms. While their expression is

comparable in the germline, they begin to exhibit distinct expression and regulation

patterns across various developmental stages (Chakravarti et al. 2022). In somatic

tissues, p53A mRNA and protein were found to be more abundant, likely reflecting

differential roles during tissue development (Chakravarti et al. 2022). This may reflect

their role in controlling apoptosis.

Initial studies demonstrated that overexpression of each p53 protein isoform leads to

embryonic lethality. In addition, it was demonstrated that overexpression of p53A or

p53B and of p53E induced apoptosis in the developing eye when driven by the GMR

promoter, while in the ovary, overexpression of p53A and p53B proteins markedly

enhanced apoptosis following irradiation, whereas p53E overexpression reduced the

number of apoptotic cells, suggesting a dominant-negative role for this isoform in the

ovary under irradiation-induced stress conditions (Zhang et al. 2015). Numerous studies

have demonstrated that both p53 protein A and B isoforms play distinct roles in

regulating stress responses and inducing apoptosis; indeed, p53A was shown to be

both necessary and sufficient to activate the proapoptotic genes *rpr* and *hid* in response

to DNA damage in somatic and germline tissues (Zhang et al. 2015; Chakravarti et al.

2022).

Although p53B can induce apoptosis when experimentally overexpressed, its behavior

in normal tissues appears to be different. In several somatic tissues, the mRNA levels of

p53B are only slightly lower than those of p53A, yet the amount of p53B protein

detected is considerably lower. This discrepancy suggests that p53B may be translated

less efficiently or undergo more rapid protein degradation (Zhang et al. 2015; Zhang et

al. 2014). These differences in protein abundance and activity among tissues may stem

from competition between p53 isoforms that have opposing regulatory functions.

Specifically, p53B primarily acts as a transactivator, promoting the expression of target

genes, whereas p53A, which is more broadly expressed across tissues, can function as

both a transactivator and a transrepressor ,(Wylie et al. 2022). Thus, the relative levels

and functional properties of these isoforms likely contribute to the tissue-specific

outcomes of p53 signaling. In support of this hypothesis, studies in photoreceptor

neurons have shown that the balance between autophagy-mediated survival and

apoptosis-driven cell death is governed by antagonistic regulatory inputs from p53A and

p53B, whose mRNA levels shift differentially in response to oxidative stress (Robin et al.

2019).

Although p53 is best known for its conserved role in apoptosis triggered by DNA

damage, it also exerts broad control over tissue physiology and metabolic homeostasis.

Under nutrient-limiting conditions, *p53*-null mutant flies fail to sustain normal survival,

revealing that p53 is essential for coordinating metabolic adaptation and efficient

nutrient consumption during stress (Barrio et al. 2014). p53 also regulates growth and

proliferation in a coordinated, non-autonomous manner and modulates apoptosis in

response to stress, such as rRNA depletion (Mesquita et al. 2010). Furthermore, p53 is

necessary to maintain imaginal disc plasticity and regenerative potential (Wells and

Johnston 2012) and to act as a general sensor of competitive confrontation during

MYC-induced cell competition (de la Cova et al. 2014). Notably, p53 isoform-specific

protein functions have not been assessed in these experiments despite growing

evidence that their function can differ significantly depending on the cellular context in

which they are expressed. Thus, it may be important to consider that distinct signals

and tissue-specific expression programs could drive isoform-specific roles in stress

response and gene regulation.

Our previous study demonstrates that reducing the nucleolar gene *Noc1* (FBgn0036124)

impairs rRNA processing and ribosome biogenesis, leading to apoptosis and the

upregulation of p53 (Rambaldelli et al. 2025; Destefanis et al. 2022), both hallmarks of

nucleolar stress. Using our newly designed p53 isoform-specific qPCR primers, we
found that *p53E* is the most upregulated isoform following *Noc1* reduction. This
suggests that nucleolar stress may induce a novel regulatory mechanism for p53E,
whose strong upregulation upon nucleolar disruption points to a potential autoregulatory
interaction in which one p53 isoform may influence the activity or stability of other
isoforms. In addition, γ-H2AV phosphorylation accumulates because of caspase-
dependent cell death rather than direct genomic lesions, consistent with the activation of
a p53 stress pathway distinct from the canonical genotoxic pathway. Together, these
findings suggest that p53E may be part of a novel mechanism activated during
nucleolar stress, providing insight into how cells adapt to defects in ribosome
biogenesis.

Materials and Methods
137
138
139 Drosophila husbandry and lines

140
Animals were raised at low density in vials containing standard fly food, composed of 9
g/l agar, 75 g/l corn flour, 50 g/l fresh yeast, 30 g/l yeast extract, 50 g/l white sugar, and
30 ml/l molasses, along with nipagin (in ethanol) and propionic acid. The crosses and
flies used for the experiments are kept at 25°C, unless otherwise stated.
Line used: w^{1118} from VDRC #6000, *Rotund*-Gal4 BDSC#76179, *UAS-Noc1-RNAi*
BDSC #25992, *UAS-p35* (Destefanis et al. 2022), $p53^{5-1-4}$ BDSC #6815, $p53^{A2.3}$ and
$p53^{B41.5}$ from (Chakravarti et al. 2022).

Alkaline Comet Assay

Using the *UAS-Gal4* system (Brand and Perrimon 1993), *Noc1* was downregulated in
the entire imaginal disc using a temporally controlled expression system under the
Actin-Gal4 promoter in combination with the temperature-sensitive *Gal80-ts* allele that
inhibits Gal4 (Germani et al. 2018). Animals were left to develop for five days after egg
laying (AEL) at the permissive temperature of 18°C, then shifted to 30°C for 60 hours,

during which Gal4 was active and induced the ubiquitous expression of *Noc1-RNAi*. Wing imaginal discs were dissected and incubated in Ringer's solution, then enzymatically digested with Collagenase in TrypLE at 37–38 °C for 30 minutes following the protocol in (Pederzolli et al. 2025). After electrophoresis and Hoechst staining, photos of nuclei were imaged using a Zeiss Axio Imager fluorescent microscope with a 40X objective, and DNA damage was analyzed with CometScore 2.0.

RNA Extraction and RT-qPCR

RNA extraction was performed with the RNeasy Mini Kit (Qiagen). Five larvae or 30 wing discs were collected and placed in 250µl of lysis buffer, then processed following the manufacturer's instructions. The isolated RNA was quantified using a Nanodrop (Thermo Scientific) and stored at -80 °C for extended periods. RNA (1µg) was retro-transcribed into cDNA using the SuperScript IV VILO Master Mix (Invitrogen). DNase was added to the samples. To perform retro-transcription, the mix was incubated at 25°C for 10 minutes, 50°C for 10 minutes, and 85°C for 5 minutes. The obtained cDNA was used to perform quantitative PCR (qPCR) with the SYBR Green PCR Kit (Qiagen). The final mix for each sample to be tested was composed by 8µl of cDNA, 2µl of primers (10µM), and 10µl of SYBR Green reaction mix. Reactions were carried out using a BioRad CFX96 machine with the following protocol: 50°C for 2 minutes, 95°C for 2 minutes, 95°C for 15 seconds, and 60°C for 30 seconds (repeated for 45 cycles), followed by 72°C for 1 minute. Analyses were done using Bio-Rad CFX Manager software. The relative abundance level for each transcript was calculated by subtracting the value of actin from the value of p53 and using the $2^{-\Delta\Delta Ct}$ method. For qPCR in wing discs analysis, the RNA was extracted from at least 20 wing imaginal discs, or from at least five larvae for each genotype. Data are expressed as arbitrary units (A.U.). The number of independent biological replicates is cited in the figure legend for each data set. The sequences of the primers used are in Table 1.

Dissections and immunofluorescence

Larvae were collected at the third-instar stage, dissected in 1× phosphate-buffered
saline (PBS), and fixed for 30 min in 4% paraformaldehyde (PFA) at room temperature
(RT). After 15 minutes of tissue permeabilization with 0.3% Triton X-100, samples were
washed in PBS with 0.04% Tween 20 (PBS-T) and blocked in 1% bovine serum
albumin (BSA) for 1 hour at RT. Samples were incubated overnight at 4°C with primary
antibodies in 1% BSA and, after washing, with Alexa-Fluor-488- or 555-conjugated
secondary antibodies in BSA. During washing in PBS-T, nuclei were stained with
Hoechst 33342 (Thermo Fisher Scientific). Imaginal discs were dissected from the
carcasses and mounted on slides with Vectashield (LsBio-Vector Laboratories). Images
were acquired using a Leica SP8 confocal microscope or Zeiss Axio Imager fluorescent
microscope and analyzed with the Fiji program to measure fluorescence. Images were
assembled using Adobe Photoshop 2025. Primary antibodies used were rabbit anti-
cleaved Caspase3, 1:400 (Cell Signaling 9661); and anti p2HAV, 1:1000
(DSHB#UNC93-5.2.1)(Lake et al. 2013).
201
202 Image processing
203
204 Raw confocal images were processed with Fiji to obtain a z-stack projection (using the
205 Average Intensity or Max Intensity options). Then, all images from the same experiment
206 were processed, enhancing the brightness of all channels equally across all images.
207
208 **Results**
209
210 **1.1 Reduction of Noc1 triggers DNA Double-Strand Breaks, p53 upregulation, and**
211 **apoptotic signaling that regulates γH2AV phosphorylation.**
212
We previously demonstrated that downregulation of the nucleolar factor Noc1 in
Drosophila tissues leads to the accumulation of rRNA (Destefanis et al. 2022;
Rambaldelli et al. 2025). Furthermore, *Noc1-RNAi* expression driven by the *rotund*
promoter in cells of the wing imaginal discs induces DNA damage as evidenced by the
comet-like morphology of the nuclei in the affected cells (Figure 1A-B, and (Pederzolli et

al. 2025). These nuclei show a statistically significant increase in the comet tail length
(or tail moment, calculated as tail length multiplied by the percentage of DNA in the
comet head), indicating the presence of double-strand breaks absent in the nuclei of
wild-type w^{1118} cells (Figure 1C). Notably, reduction of Noc1 is accompanied by a
significant increase in apoptosis in cells of the wing pouch (Figure 1D-E and F), together
with the upregulation of p53 at both transcriptional and protein levels (Figure 1G-H).

224
To assess the impact of Noc1 downregulation on DNA damage signaling, we monitored
γH2Av phosphorylation, a well-established marker of double-strand breaks.
Interestingly, blocking apoptosis prevented γH2AV accumulation (Figure 1I), indicating
that in this context, γH2Av phosphorylation is downstream of apoptotic signaling rather
than a direct consequence of Noc1 loss. This suggests that apoptosis itself contributes
to the observed DNA damage response in these tissues. Together, these findings are
consistent with a cellular response to nucleolar stress resulting from impaired rRNA
processing upon Noc1 reduction. Based on these data, we designed isoform-specific
qPCR primers to determine which individual p53 isoforms respond to Noc1 reduction,
as they may respond specifically to nucleolar stress.

235
1.2 p53 locus and novel primers for detecting distinct p53 isoforms

The *Drosophila* genome contains a single *p53* locus (Figure 2A, FBgn0039044 and
NCBI gene ID: 2768677) on chromosome 3R, which produces four distinct mRNA
transcripts through alternative promoter usage and splicing, resulting in four protein
isoforms (Figure 2B). The canonical *p53A* (also referred to as ΔNp53) lacks part of the
N-terminal Transactivation Domain (TAD), producing a protein of approximately 44 kDa.
The *p53B* transcript retains the full TAD, encoding for a protein of about 56 kDa, while
the rest of the sequence is identical to that of *p53A* mRNA. p53C protein shares the
same amino acid sequence as p53A, but their mRNAs originate from an alternative
transcription start site and are predicted to produce a longer transcript. Finally, *p53E* is
the shortest isoform, predicted to encode a 38 kDa protein that lacks the N-terminal
TADs of p53A and p53B, but retains the DNA-binding domain. This isoform has been
shown to inhibit the apoptotic response to ionizing radiation, suggesting that it may act

as a dominant-negative regulator of the other p53 isoforms (Wylie et al. 2022; Zhang et
al. 2015).

Although primer sets have previously been reported for distinguishing among p53
mRNA isoforms (Supplementary Figure 1), none have been fully optimized to reliably
differentiate between *p53A* and *p53B*. Additionally, primers capable of specifically
discriminating between *p53B* and *p53C,* or uniquely amplifying *p53C*, remain
unavailable. To overcome these limitations, we designed and optimized a novel set of
primers to enable precise and isoform-specific analysis of p53 transcripts (Table 1).

To ensure isoform specificity and accurate validation, primers were designed using the
sequences of the available *p53^{5A-1-4}* null lines and the isoform-specific *p53A* and *p53B*
mRNA mutant lines. Briefly, the *p53A* null line carries a 23bp deletion and 7bp insertion
at the end of the first exon which expands also in the first intron (Robin et al. 2019); the
mutant *p53B* null line has a 14 bp deletion and 1 bp insertion in the second exon,
resulting in multiple early STOP codons (Chakravarti et al. 2022); and the *p53^{5A-1-4}* null
line carries a large deletion of approximately 3.3 Kb that removes four exons, preventing
the transcription of a functioning mRNA, resulting in a complete loss of p53 protein
expression (Figure 2C) (Rong et al. 2002).

The new primers for the *p53A* isoform span the exon-exon junction between exons 1
and 2 (Figure 2D-i, shown as light green arrows in the panel, and Table1), providing
enhanced specificity compared to earlier designs (shown as arrows in pink, and Table
1). Like the previously available primers, this set can be used to assess *p53A*
expression in the p53A null line as the forward primer anneals upstream of the deleted
region, while the reverse primer anneals right after the deletion, allowing the
amplification of a specific region of the mutant (Figure 2D-ii). A limitation of the new
primers, as with previously published ones, is their inability to distinguish between *p53A*
and *p53C* transcripts, due to the high sequence similarity between them. To design
primers capable of distinguishing *p53B* from *p53C* isoform, we took advantage of the
difference in the length of their 5′ untranslated region (5'UTR) (Figure 2E). While the
reverse primer anneals in a sequence common between *p53B* and *p53C*, the forward

primer for *p53B* anneals at the end of the second exon and extends into the beginning

of exon 3 (green arrows in Figure 2E-i and 2E-iii). This design maximizes specificity,

ensuring amplification of only *p53B*. Moreover, this set of primers is suitable for

screening *p53B* in *p53B* null lines, as they do not anneal within the deleted region

(Figure 2E-ii).

Through detailed analysis of the 5′UTR sequence of *p53C*, we identified a region unique

to this isoform (highlighted in orange in Figure 2Fi). Thus, we designed primers that

specifically target this region, enabling the selective detection of *p53C* mRNA and

providing a distinct primer set for this isoform. Furthermore, the availability of p53C-

specific primers allows for a more accurate interpretation of results obtained with the

p53A primer set, which cannot distinguish between the *p53A* and *p53C* transcripts.

In addition to p53A, p53B, and p53C, we developed isoform-specific primers for the

truncated p53E isoform. These primers target the unique 5' sequence generated by the

p53E transcription start site (Figure 2G), providing selective amplification and

preventing cross-detection with other isoforms.

Primer specificity was tested by performing qPCRs on RNA extracted from imaginal

discs of w^{1118} third-instar larvae. All primers amplified a single product, as evidenced by

a single band observed in the gel (Figure 2H) and confirmed by melting curve analysis,

which showed a distinct peak for each primer (Supplementary Figure 2).

1.3 Validation of isoform-specific primers in wild-type and p53 mutant

background.

To further assess the specificity and to complete the characterization of the newly

designed primer sets, we conducted qPCR analysis on RNA from wild-type control w^{1118}

and *p53* mutant lines using primers targeting the single *p53* mRNA isoforms (Table 1).

As shown in Figure 3A, the p53 pan primers, which target a region common to all the

p53 isoforms (Figure 2B-i), detect p53 transcript levels in the *p53A* null line that are

comparable to those of control w^{1118}. In contrast, the *p53B* null line exhibits a small but

not significant increase in *p53* mRNA expression, suggesting a potential compensatory
upregulation in response to the loss of *p53B.*

Using the p53A/C primers, we observed that the *p53A/C* mRNA levels were significantly
reduced by approximately 50% in the *p53A* null line (Figure 3B). In contrast, the *p53B*
null line showed a significant twofold increase in *p53A/C* mRNA expression. Consistent
with the data in Figure 3A, these findings may suggest the presence of a compensatory
response to the other isoforms when p53B is eliminated in null animals, whereas loss of
p53A has minimal effect on overall transcript abundance.

In the *p53B* null line (Figure 3C), we observed an unexpected significant upregulation of
p53B mRNA. However, since the mutation does not prevent transcription and the
primers amplify regions downstream of the deletion, it is likely that the mutant transcript
can still be detected by qPCR. Overall, this finding suggests that compensatory or
autoregulatory feedback mechanisms may act to increase *p53B* transcription or stabilize
the mutant mRNA in response to reduced levels of p53B protein.
Finally, analysis of the *p53C* isoform, using primers that specifically detect the
transcript, reveals a significant reduction in its expression in both *p53A* and *p53B* null
lines (Figure 3D), suggesting that both *p53A* and *p53B* isoforms, directly or indirectly,
participate in the regulation of *p53C* expression.
Importantly, the complete absence of *p53* transcripts in the $p53^{5A-1-4}$ null line confirms
the specificity and reliability of the isoform-targeted primer set used in this study.

1.4 p53E is the predominantly upregulated isoform upon Noc1 downregulation.

Analysis of p53 expression in cells of the wing imaginal disc in which Noc1 was reduced
in the pouch revealed a significant upregulation of *p53* mRNA levels when assessed
using pan-isoform primers (Figure 4A and Figure 1G). Because these primers recognize
a conserved region shared by all four p53 isoforms, as illustrated in the magnified view
(Figure 2B-i), this initial analysis did not allow discrimination between individual isoform
contributions. Given that Noc1 reduction leads to rRNA accumulation and increased p53

levels (Figure 1B-C), and considering that p53 isoforms can be expressed at different
levels under various stress conditions, we therefore designed isoform-specific primers
to determine which p53 isoform(s) are preferentially upregulated in response to *Noc1-*
RNAi-induced nucleolar stress.

Among the isoforms, p53E exhibited the strongest induction ($p < 0.001$), whereas all the
other isoforms showed a moderately significant increase ($p < 0.05$). Notably, p53E has
been reported to act as a dominant negative in a tissue-specific manner. Thus, our
analysis suggests that p53E may contribute to the repression of specific target genes
during the nucleolar stress response.

Collectively, these findings highlight an isoform-specific regulation of p53 in response to
nucleolar stress, with p53E emerging as a potential novel key mediator of transcriptional
repression. The differential upregulation of p53 isoforms suggests a fine-tuned cellular
strategy, where transrepressor isoforms may selectively modulate gene expression to
orchestrate an appropriate stress response, while other isoforms potentially maintain
canonical p53 functions. This underscores the importance of considering isoform-
specific roles in understanding p53-mediated pathways under nucleolar stress
conditions.

Discussion

In this study, we present an optimized and validated set of isoform-specific primers
designed for the accurate detection and quantification of *Drosophila* p53 transcript
variants. Here, we also designed primers to specifically detect the unique p53C isoform,
enabling precise isoform discrimination. We first utilized p53 mutant lines to validate the
specificity of these primers. These analyses show that the *p53A* null line retains some of
p53A and *p53C* transcript levels (Figure 3B and D). Moreover, despite the introduction
of premature stop codons within the coding sequence of the *p53B* mutation, our qPCR
analysis revealed an upregulation of *p53B*-mRNA in *p53A/C* and *B* mutant lines (Figure
3D and C), which may suggest the presence of a compensatory mechanism, to buffer

the loss of p53B function, or to compensate for the reduction of p53A/C protein, highlighting isoform-specific plasticity of the p53 network, where alternative isoforms may adjust expression to preserve critical stress response functions.

Our analysis of isoform-specific p53 expression under nucleolar stress induced by Noc1 reduction (Milkereit et al. 2001; Destefanis et al. 2022) (Figure 1) reveals that all *p53* isoforms were upregulated ($p < 0.05$), with *p53E* showing the most robust increase ($p < 0.01$), marking its first correlation with nucleolar stress and its role in stress-induced apoptosis. This truncated variant, characterized by a unique N-terminal domain, has been shown to counteract p53-induced apoptosis in eye imaginal discs (Zhang et al. 2015).

Indeed, the transcriptional repression activity of p53 isoforms is highly context-dependent. While p53A was shown to act as both an activator and a repressor in salivary glands, p53E functions only as a transrepressor and is dispensable for transactivation in irradiated embryos, where p53A was essential (Wylie et al. 2022). However, our study specifically addresses nucleolar stress, which is mechanistically distinct from stress induced by DNA damage. In wing imaginal discs subjected to Noc1 depletion, p53E may be selectively induced to engage stress-specific or compensatory mechanisms that modulate the nucleolar stress response in a manner that differs across tissues and developmental stages. p53E could influence the relative abundance and composition of p53 oligomeric complexes, thereby fine-tuning p53 signaling output.

In this context, p53E may alter the balance of p53A-p53B-containing complexes. Depending on the cellular milieu, p53A can affect the formation of homo- or hetero-oligomers, ultimately shaping transcriptional responses and cell fate decisions. Consistent with this model, shorter p53 isoforms have been shown to repress the activity of longer isoforms through heterotetramer formation, a conserved mechanism of p53 isoform regulation (Anbarasan and Bourdon 2019). Supporting isoform-specific regulation in vivo, Chakravarti et al. (Chakravarti et al. 2022) used isoform-specific mutants with the *hid-GFP* reporter in the germline, demonstrating differential transcription outputs among p53 isoforms. Together, these studies highlight the

importance of heterotetramer formation among p53 isoforms (Zhang et al. 2014) and

suggest a potential role for post-translational modifications in modulating isoform-

specific functions. Further investigation will be required to determine how these

regulatory mechanisms integrate under distinct cellular stress conditions, including

nucleolar stress.

Post-translational modifications (PTMs) also contribute to isoform-specific regulation of

p53. Indeed, ubiquitination, phosphorylation, sumoylation, and acetylation may differ

among p53 isoforms due to variations in their amino acid sequences. Lok/Chk2-

dependent phosphorylation at serine 4 activates DNA repair and apoptotic responses

(Brodsky et al. 2004; Peters et al. 2002). The E3 ligase dSyx5 acts as a negative

regulator of p53 protein levels, while the role of the deacetylase dSir2 remains unclear

(Bauer and Helfand 2009). Sumoylation at lysines 26 and 302 is essential for its full

transcriptional and apoptotic activity (Mauri et al. 2008).

Moreover, our findings also highlight key mechanistic differences in p53 regulation

between flies and humans. In vertebrates, nucleolar stress primarily activates p53 post-

translationally through protein stabilization, mediated by ribosomal proteins that inhibit

the E3 ubiquitin ligase MDM2 (Lindstrom et al. 2022). In contrast, *Drosophila* lacks a

clear MDM2 homolog with E3 ligase activity, suggesting an alternative regulatory

mechanism. The gene *companion of reaper* (*Corp*) has been proposed as a structural

and partial functional analog of mammalian MDM2. Corp binds to p53 and negatively

regulates its levels through a feedback loop, similar to MDM2; however, it lacks an E3

ligase domain and thus cannot mediate canonical p53 degradation (Chakraborty et al.

2015). These distinctions point to a divergence in the evolution of p53 regulatory

circuits, where transcriptional control may play a more prominent role in flies. This

difference emphasizes the evolutionary divergence in p53 regulation and raises

important questions about whether isoform-specific transcriptional control might also

serve as a stress adaptation mechanism in vertebrates, including humans.

Further supporting the evolutionary conservation of nucleolar stress responses, our

previous data show that downregulation of the human homolog of Noc1, called *CEBPZ*

(Rambaldelli et al. 2025), also activates p53 both transcriptionally and at the protein

level, consistent with our findings in *Drosophila* (Destefanis et al. 2022). Noc1 functions as part of a conserved nucleolar heterodimeric complex with Noc2 and Noc3, which is essential for pre-rRNA processing and 60S ribosomal subunit assembly, as demonstrated in both yeast and *Drosophila* (Milkereit et al. 2001; Destefanis et al. 2022). Our finding that Noc1 depletion triggers p53 upregulation in both flies and humans highlights the evolutionary conservation of nucleolar stress responses. This conserved activation highlights the broader relevance of nucleolar stress-induced p53 signaling and underscores the value of isoform-specific tools in elucidating the underlying regulatory mechanisms.

In conclusion, our analysis, enabled by isoform-specific primers and p53 mutant lines, reveals previously unrecognized residual activity and compensatory regulation, underscoring the intricate and multifaceted control of p53 in *Drosophila*. Our results indicate that p53E may participate in the apoptotic response triggered by nucleolar stress, suggesting it may have an isoform-specific role in this process. Furthermore, the observation that γ-H2AV phosphorylation reflects caspase-dependent apoptosis rather than direct genomic lesions indicates activation of a p53 stress response distinct from canonical genotoxic pathways. Collectively, these findings suggest that p53E is a component of a novel nucleolar stress–responsive mechanism that influences cellular adaptation to defects in ribosome biogenesis.

While our findings are consistent with potential transcriptional regulation of stress responses, further studies will be required to determine the relative contributions of transcriptional and post-translational mechanisms in maintaining cellular homeostasis.

Data Availability Statement

All data are available in the published manuscript and its online Supplementary Files: Supplementary Figure 1, which reports the sequences of all the primers published for p53 isoforms, along with their corresponding recognition region in the *p53* locus; Supplementary Figure 2, which includes the melting curve of the primer sets used; and

Supplementary Figure 3. Sequences showing the mutation present in the $p53^{A2.3}$,
$p53^{B41.5}$ mutant lines.

Acknowledgments

We thank the Bloomington Stock Center (NIH P40OD018537). Department CIBIO Core
Facilities is supported by the European Regional Development Fund (FESR) 2021–
2027. The input from members of the COST Action CA21154 TRANSLACORE,
supported by COST (European Cooperation in Science and Technology and by the
Dipartimento di Eccellenza 2023-2027, Legge 232/2016 project n 40613, funded by the
MUR. This document includes language and clarity improvements made using
ChatGPT (OpenAI).

Author contributions

AVC performed the experiments, conceived the study, and contributed to the writing of
the manuscript. SB contributed to the final version of the manuscript. VM performed the
experiments and analyzed the data. PB participated in the design, coordinated and
supervised the team, and contributed to the writing and editing of the final manuscript.
All authors have read and agreed to the published version of the manuscript.

Conflict of Interest

No competing interests are declared.

Funder Information

No specific funds for this manuscript are declared.

Table 1: Sequences of the primers used in this work.

Target	Sequence	Reference
p53A/C	F: TATATCACAGCCAATGTCGTG	This work
	R: CCTCCGTGGAGTCATCC	

p53B	F: CCACAAGGGCACTGATTC	This work
	R: ATTCCGATCCCGATACCTC	
p53C	F: GTGAGCAAATTCAAAACACGC	This work
	R: CCATTCGCCACGCAG	
p53E	F: GAGCGCAAACAATCCGT	This work
	R: TGTGATTCTCTAGCTTGGGC	
p53 pan isoform	F: CCAAGCTAGAGAATCACAACATCG	(Zhang et al. 2014)
	R: TCGAGTACATCCAAAGAGACTTGG	

Bibliography

Anbarasan, T., and J.C. Bourdon, 2019 The Emerging Landscape of p53 Isoforms in
Physiology, Cancer and Degenerative Diseases. *Int J Mol Sci* 20 (24).

Barrio, L., A. Dekanty, and M. Milan, 2014 MicroRNA-mediated regulation of Dp53 in
the Drosophila fat body contributes to metabolic adaptation to nutrient
deprivation. *Cell Rep* 8 (2):528–541.

Bauer, J.H., and S.L. Helfand, 2009 Sir2 and longevity: the p53 connection. *Cell Cycle* 8
(12):1821.

Boley, N., M.H. Stoiber, B.W. Booth, K.H. Wan, R.A. Hoskins *et al.*, 2014 Genome-
guided transcript assembly by integrative analysis of RNA sequence data. *Nat*
Biotechnol 32 (4):341–346.

Boukoura, S., and D.H. Larsen, 2024 Nucleolar organization and ribosomal DNA
stability in response to DNA damage. *Curr Opin Cell Biol* 89:102380.

Bourdon, J.C., K. Fernandes, F. Murray-Zmijewski, G. Liu, A. Diot *et al.*, 2005 p53
isoforms can regulate p53 transcriptional activity. *Genes Dev* 19 (18):2122–2137.

Brand, A.H., and N. Perrimon, 1993 Targeted gene expression as a means of altering
cell fates and generating dominant phenotypes. *Development* 118 (2):401–415.

Brodsky, M.H., W. Nordstrom, G. Tsang, E. Kwan, G.M. Rubin *et al.*, 2000 Drosophila
p53 binds a damage response element at the reaper locus. *Cell* 101 (1):103–
113.

Brodsky, M.H., B.T. Weinert, G. Tsang, Y.S. Rong, N.M. McGinnis *et al.*, 2004
    Drosophila melanogaster MNK/Chk2 and p53 regulate multiple DNA repair and
    apoptotic pathways following DNA damage. *Mol Cell Biol* 24 (3):1219–1231.

Chakraborty, R., Y. Li, L. Zhou, and K.G. Golic, 2015 Corp Regulates P53 in Drosophila
    melanogaster via a Negative Feedback Loop. *PLoS Genet* 11 (7):e1005400.

Chakravarti, A., H.N. Thirimanne, S. Brown, and B.R. Calvi, 2022 Drosophila p53
    isoforms have overlapping and distinct functions in germline genome integrity
    and oocyte quality control. *Elife* 11.

de la Cova, C., N. Senoo-Matsuda, M. Ziosi, D.C. Wu, P. Bellosta *et al.*, 2014
    Supercompetitor Status of Drosophila Myc Cells Requires p53 as a Fitness
    Sensor to Reprogram Metabolism and Promote Viability. *Cell Metab*.

Destefanis, F., V. Manara, S. Santarelli, S. Zola, M. Brambilla *et al.*, 2022 Reduction of
    nucleolar NOC1 leads to the accumulation of pre-rRNAs and induces Xrp1,
    affecting growth and resulting in cell competition. *J Cell Sci* 135 (23).

Germani, F., C. Bergantinos, and L.A. Johnston, 2018 Mosaic Analysis in Drosophila.
    Genetics 208 (2):473–490.

Ingaramo, M.C., J.A. Sanchez, and A. Dekanty, 2018 Regulation and function of p53: A
    perspective from Drosophila studies. *Mech Dev* 154:82–90.

Lake, C.M., J.K. Holsclaw, S.P. Bellendir, J. Sekelsky, and R.S. Hawley, 2013 The
    development of a monoclonal antibody recognizing the Drosophila melanogaster
    phosphorylated histone H2A variant (gamma-H2AV). *G3 (Bethesda)* 3 (9):1539–
    1543.

Lindstrom, M.S., J. Bartek, and A. Maya-Mendoza, 2022 p53 at the crossroad of DNA
    replication and ribosome biogenesis stress pathways. *Cell Death Differ* 29
    (5):972–982.

Mauri, F., L.M. McNamee, A. Lunardi, F. Chiacchiera, G. Del Sal *et al.*, 2008
    Modification of Drosophila p53 by SUMO modulates its transactivation and pro-
    apoptotic functions. *J Biol Chem* 283 (30):20848–20856.

Mesquita, D., A. Dekanty, and M. Milan, 2010 A dp53-dependent mechanism involved
    in coordinating tissue growth in Drosophila. *PLoS Biol* 8 (12):e1000566.

Milkereit, P., O. Gadal, A. Podtelejnikov, S. Trumtel, N. Gas *et al.*, 2001 Maturation and
    intranuclear transport of pre-ribosomes requires Noc proteins. *Cell* 105 (4):499–
    509.

Ollmann, M., L.M. Young, C.J. Di Como, F. Karim, M. Belvin *et al.*, 2000 Drosophila p53
    is a structural and functional homolog of the tumor suppressor p53. *Cell* 101
    (1):91–101.

Pederzolli, M., E. Barion, A. Valerio, A.V. Cuda, V. Manara *et al.*, 2025 Optimized
protocol for single-cell isolation and alkaline comet assay to detect DNA damage
in cells of Drosophila wing imaginal discs. *STAR Protoc* 6 (1):103590.

Peters, M., C. DeLuca, A. Hirao, V. Stambolic, J. Potter *et al.*, 2002 Chk2 regulates
irradiation-induced, p53-mediated apoptosis in Drosophila. *Proc Natl Acad Sci U*
S A 99 (17):11305–11310.

Rambaldelli, G., V. Manara, A. Vutera Cuda, G. Bertalot, M. Penzo *et al.*, 2025 From
Flies to Humans: Conserved Roles of CEBPZ, NOC2L, and NOC3L in rRNA
Processing and Tumorigenesis. *bioRxiv*.

Robin, M., A.R. Issa, C.C. Santos, F. Napoletano, C. Petitgas *et al.*, 2019 Drosophila
p53 integrates the antagonism between autophagy and apoptosis in response to
stress. *Autophagy* 15 (5):771–784.

Rong, Y.S., S.W. Titen, H.B. Xie, M.M. Golic, M. Bastiani *et al.*, 2002 Targeted
mutagenesis by homologous recombination in D. melanogaster. *Genes Dev* 16
(12):1568–1581.

Wells, B.S., and L.A. Johnston, 2012 Maintenance of imaginal disc plasticity and
regenerative potential in Drosophila by p53. *Dev Biol* 361 (2):263–276.

Wylie, A., A.E. Jones, S. Das, W.J. Lu, and J.M. Abrams, 2022 Distinct p53 isoforms
code for opposing transcriptional outcomes. *Dev Cell* 57 (15):1833–1846 e1836.

Zhang, B., S. Mehrotra, W.L. Ng, and B.R. Calvi, 2014 Low levels of p53 protein and
chromatin silencing of p53 target genes repress apoptosis in Drosophila
endocycling cells. *PLoS Genet* 10 (9):e1004581.

Zhang, B., M. Rotelli, M. Dixon, and B.R. Calvi, 2015 The function of Drosophila p53
isoforms in apoptosis. *Cell Death Differ* 22 (12):2058–2067.

Figure Legends

Figure 1. Noc1 Downregulation induces DNA double-strand breaks (DSBs),
leading to p53 upregulation and apoptosis, which contributes to γH2AV
phosphorylation.

(A-C) Comet assay showing fluorescent images of nuclei stained with Hoechst (white)

from cells of the wing imaginal discs of w^{1118} third instar larvae (A), and from animals in

which *Noc1* was transiently reduced for 60 hours under the temporal control of the *Act-*

Gal80^{ts} system. (B), scale bar is 50μm. (C) Quantification of single-cell tail moment from

the comet assay performed on nuclei from the genotypes shown in A and B. These data

are obtained from pooling at least 20 wing imaginal discs of each genotype, and one

representative experiment is shown. Statistical analysis was performed using the non-

parametric Kruskal-Wallis as described in (Pederzolli et al. 2025). Experiments in (D-I)

are from wing imaginal discs from third instar control larvae ($rn>w^{1118}$) or from larvae in

which expression of the transgenes was reduced using the indicated Rnai lines in the

wing pouch under the control of the *rotund* (*rn*) promoter. (D-E) Confocal images of wing

imaginal discs from third instar larvae immunostained for the expression of p53 protein

(green) and for Caspase3, as a marker of apoptosis (red). Nuclei were stained with

Hoechst (blue). Scale bar is 100μm. (F) Quantification of Caspase3 by

immunofluorescence in cells from the pouch of wing discs from third instar larvae of the

indicated genotype. (G) qPCR analysis from RNA extracted from at least 20 wing

imaginal discs from larvae of the indicated genotype, *p53* mRNA expression was

analyzed using p53 pan isoform primers (Table 1). Data are expressed as arbitrary units

(A.U.) (H). Quantification of p53 by immunofluorescence was analyzed in the pouch of

wing imaginal discs from animals of the indicated genotype. (I) Quantification of γH2AV

phosphorylation by immunofluorescence was analyzed in the wing pouch form larvae of

the indicated genotype. In C, the asterisks represent the *p-values* from Student's *t*-test

**** = $p < 0.0001$, in one out of three independent experiments performed, where the

dots represent the number of nuclei analyzed. In F, G, H, and I, the dots represent the

number of independent biological replicates, with at least five imaginal discs analyzed

for each genotype. The asterisks represent the *p-values* from Student's *t*-test * = $p <$

0.05 and ** = $p < 0.01$; the error bars indicate the standard deviations.

Figure 2. Schematic representation of the p53 locus and primer design used to

detect specific isoforms. (A-B) Schematic representation of the *p53* locus (gene ID

2768677) and of the p53 isoforms *A, B, C,* and *E*. (B-i), magnification of a common exon

with the representation of the region where the primers that capture all *p53* isoforms

anneal (Zhang et al. 2015). These primers detect mRNA transcripts for all isoforms and

were used in Figure 1F; they will be referred to as p53 pan. (C) Schematic

representation of the $p53^{A2.3}$, $p53^{B41.5}$, and $p53^{5A-1-4}$ null mutant alleles. The $p53^{A2.3}$ null

line was generated with a 23bp deletion and a 7bp insertion spanning from the end of the first exon until the first base of the first intron (in red), generating multiple early STOP codons (Supplementary Figure 3A). This disrupts protein production from the p53A and p53C isoforms, but mRNA may still be detectable if primers anneal outside the deleted region. $p53^{B41.5}$ null line carries a 14bp deletion and a 1bp insertion in exon 2, resulting in a frameshift and premature stop codons leading to a truncated, non-functional p53B protein (Supplementary Figure 3B)(Chakravarti et al. 2022). $p53^{5A-1-4}$ null line, contains a 3.3 kb deletion that removes the common 3' coding region (in red), eliminating protein expression from all isoforms. (D) *p53A* isoform. (D-i and ii). Magnification of exons 1 and 2 of *p53A* and $p53^{A2.3}$ null isoform, respectively, showing where the primers designed in this study anneal (green) compared to previous primers (in pink). As shown in D-ii, the new primers can also be used to detect *p53A* mRNA in the $p53^{A2.3}$ null mutant, because the forward primer anneals outside the deleted region. In contrast, the previous reverse primer (pink) has 1 bp mismatch and may fail to detect the transcript in the *p53A* null line. The new primers also detect *p53C* mRNA and will therefore be referred to as p53A/C. (E) *p53B* and *p53C* isoforms. (E-i, ii, and iii). Magnification of exons 2 and 3, shared by both isoforms, showing where the primers designed in this study anneal (in green). The new primers specifically detect *p53B* mRNA, whereas the old primers (pink) also detect *p53C* mRNA. Moreover, the new pair is also suitable for analyzing the *p53B* null line because the forward primer does not anneal in the deleted region marked in red (E-ii). This pair of primers will be referred to as p53B. (F) *p53C* isoform. (F-i). Magnification of the region in exon 2 that includes the sequence of 78 bp, from 440 to 518 nucleotides from the beginning of the *p53C* transcript (in orange), unique to this isoform. Optimized primers (green) detect *p53C* mRNA and are referred to as p53C. (G) *p53E* isoform. (G-i). Magnification of the region between exons, highlighting the amplified region specific to this isoform. (H) Agarose gel (2%) showing qPCR-amplified products from RNA extracted from wing imaginal discs of w^{1118} third-instar larvae, using the indicated primers and *actin* as a control. A single band is observed for each primer pair, demonstrating the specificity for mRNA detection.

Figure 3. Expression analysis of *p53 mRNA* isoforms by RT-qPCR in p53 mutant lines using isoform-specific primers. (A-D) Relative expression of *p53 mRNA* isoforms compared to *actin5C,* used as a control. The relative p53 isoforms analyzed with the specific primers are indicated at the top of each graph. The genotype of the lines is shown below. Relative expression of *p53 mRNA* is shown as arbitrary units (A.U.). *p53* pan detects all p53 isoforms; p53A/C specifically detects *p53A* and the predicted *p53C* isoforms; p53B, p53D, and p53E detect only the respective isoform; w^{1118} animals are used as control. The asterisks indicate the *p*-values obtained from Student's t-test analysis from two independent biological experiments. RNA was extracted from five whole larvae of the respective genotype for each experiment. * = $p < 0.05$, *** = $p < 0.001$, and **** = $p < 0.0001$, and the error bars indicate the standard deviations.

Figure 4: Upregulation of p53 isoforms in Noc1-downregulated cells. (A-E) qPCRs from RNA extracted from wing imaginal discs of w^{1118} animals and expressing *Noc1-RNAi* in the wing pouch using the *rotund* promoter. Data are expressed as relative expression of *p53 mRNA* isoforms as arbitrary units (A.U.) compared to *actin5C,* used as a control. The asterisks represent the *p*-values obtained from Student's t-test analysis: * = $p < 0.05$, and *** = $p < 0.001$. The error bars indicate the standard deviations. The dots represent the number of independent biological experiments. In each biological experiment, 30-40 wing imaginal discs were dissected per genotype and pooled for RNA extraction.