Surgical Removal of Visceral Adipose Tissue has Therapeutic Benefit in Male APP$^{NL-F}$ Mice

Samuel A. McFadden[1†], Yimin Fang[1†], Kathleen Quinn[1], Mackenzie R. Peck[1], Jenelle E.

Chapman[1], Tiarra Hill[1], Andrzej Bartke[2,3], Erin R. Hascup[1,4], Kevin N. Hascup[1,3,4*]

[1]Dale and Deborah Smith Center for Alzheimer's Research and Treatment, [2]Internal Medicine,

[3]Medical Microbiology, Immunology and Cell Biology, and [4]Pharmacology, at Southern Illinois

University School of Medicine, Springfield, IL, USA

[†]These authors contributed equally to this work and share first authorship.

*Corresponding Author: Kevin N. Hascup, Dale and Deborah Smith Center for Alzheimer's

Research and Treatment, Southern Illinois University School of Medicine, P.O. Box 19628,

Springfield, IL 62794-9628, USA Tel: 217-545-6994, Email: khascup49@siumed.edu

Keywords: Alzheimer's disease, lipectomy, white adipose tissue, cell senescence, glucose and

lipid metabolism, inflammation, visfatin

Manuscript Length: 5,557 Words

Figures: 7

Tables: 2

**Abstract**

Purpose:  Visceral white adipose tissue (vWAT) accumulation causes systemic inflammation,

insulin resistance, metabolic syndrome, and senescent cell accumulation that are risk factors for

Alzheimer's disease (AD). Visceral fat removal (VFR) improves metabolism and reduces pro-

inflammatory cytokines.  We hypothesized that VFR removal in AD mice would improve

metabolism and cognition.

Methods:  Male and female APP$^{NL-F}$ mice underwent sham or vWAT surgical resection

(periovarian or epididymal and perirenal) at 4 (pre-symptomatic) and 16 (symptomatic) months

of age to understand interventional and therapeutic effects, respectively.  At 18 months of age,

glucose metabolism and novel object recognition (NOR) memory were assayed followed by

assessment of body composition and tissue-specific markers of metabolism, cell senescence,

inflammation, or amyloid accumulation.

Results:  Male and female APP$^{NL-F}$ mice showed distinct VFR responses. In pre-symptomatic

males, increased vWAT lipolysis and hepatic lipogenesis led to ectopic liver lipid accumulation,

with reduced adiponectin and leptin, elevated visfatin, and impaired glucose metabolism.

Symptomatic males showed reduced vWAT lipogenesis, enhanced hepatic lipolysis, glycolysis,

and glycogenesis, lowering liver lipids and improving insulin sensitivity. Only symptomatic males

improved NOR, linked to elevated hippocampal learning and memory markers. Female vWAT

reaccumulation was due to increased lipogenesis and lower lipolysis.  Pre-symptomatic females

had lower hepatic lipogenesis, while glycolysis and glycogenesis declined with disease

progression. Hippocampal senescence and inflammation were elevated early in the disease that

persisted symptomatically.

Conclusions:  Sex-specific differences in glucose and lipid metabolism and lipid accumulation

underlie the divergent responses to VFR in APP$^{NL-F}$ mice, with symptomatic males showing the

only beneficial outcomes in metabolism and cognition.

## Graphical Abstract

Male and female APP$^{NL-F}$ mice exhibited distinct VFR responses to AD progression. In males, early disease stages were marked by vWAT lipolysis and hepatic lipid accumulation with metabolic dysfunction, while symptomatic stages showed a metabolic shift that improved insulin sensitivity and NOR performance. In contrast, females displayed progressive vWAT reaccumulation, reduced hepatic metabolism, and persistent hippocampal senescence and inflammation from early stages onward.

## Introduction

AD is a progressive, neurodegenerative disorder, and the most prevalent form of dementia. AD is historically characterized by senile plaques composed of aggregated amyloid beta (Aβ) and neurofibrillary tangles formed from hyperphosphorylated tau protein. However, AD is a multifactorial disorder characterized not only by aberrant protein aggregation, but also by mitochondrial dysfunction, oxidative stress, cellular senescence, and chronic inflammation - all of which contribute to disease progression (Breijyeh and Karaman, 2020). The deleterious cascade underlying AD progression is hypothesized to begin in midlife and is not isolated to the CNS.  Peripheral organs, such as vWAT, regulate brain function through hormone secretion as well as glucose and lipid metabolism (Gustafson, 2012, Dove et al., 2021). Accumulation of vWAT in midlife is a predictor for future development of AD (Huang et al., 2022).

Adipose tissue regulates many aspects of whole-body physiology, including food intake, maintenance of energy levels, insulin sensitivity, body temperature, and immune responses (Sakers et al., 2022). Adipose tissue is located under the skin (subcutaneous; SubQ), between and around organs (vWAT), interscapular brown adipose tissue (BAT), in bone marrow, and in mammary glands (Wong Zhang et al., 2023). Beyond total adipose tissue mass, the distribution of fat depots is a stronger predictor of obesity-related disease risk and associated complications (Chusyd et al., 2016). Excess vWAT accumulation causes systemic inflammation, insulin resistance, reduced cerebral blood flow, and metabolic syndrome, which contributes to brain volume shrinkage and cognitive decline (Richard et al., 2000, Wong Zhang et al., 2023, Baglietto-Vargas et al., 2016, Biessels and Despa, 2018, Dove et al., 2021, Nguyen et al., 2014).

The link between systemic inflammation with vWAT and metabolic dysfunction has been reported in diabetic fatty rodents and in aged rats (Borst et al., 2005, Gabriely et al., 2002, Pitombo et al., 2006). In humans, vWAT is a stronger predictor of age-related cognitive

impairment than body mass index (Whitmer et al., 2008, Elias et al., 2005, Kamogawa et al.,

2010). A cross-sectional study also showed that vWAT dysfunction and inflammatory status in

obesity, is associated with increased cerebral Aβ burden in midlife (Kim et al., 2022).

Transplantation of vWAT from obese mice led to impaired cognition in lean wild-type recipient

mice mediated by NLRP3/IL-1β signaling (Guo et al., 2020).  The effects of VFR are broadly

similar to observations in calorie restricted rodents (Masternak et al, 2012).  VFR prevented

obesity-induced insulin resistance, hyperinsulinemia, and decreased plasma pro-inflammatory

cytokine levels (Borst et al., 2005, Gabriely et al., 2002, Pitombo et al., 2006), which improved

glucose metabolism (Barzilai et al., 1999, Foster et al., 2011, Franczyk et al., 2021). Aging also

increases vWAT senescent cell burden leading to a chronic pro-inflammatory state (Tchkonia et

al., 2010). Reduction of senescent cells in vWAT improves metabolic and cognitive performance

in normal aging (Fang et al., 2023) and AD mice (Fang et al, 2025).

Considering the contribution of vWAT accumulation on cognitive decline and AD progression,

we hypothesized that surgical resection of this adipose depot would improve the metabolic

profile and reduce cognitive impairments in the APP$^{NL-F}$ AD mouse model.  APP$^{NL-F}$ mice contain

the human Aβ isoform of the Swedish (KM670/671NL) and Iberian (I716F) mutations knocked

into the *App* locus leading to upregulation of Aβ$_{42}$ without overexpression of APP (Saito et al.,

2014, Benitez et al., 2021). Plaque accumulation begins at ~6 months of age while cognitive

impairments occur between 12-18 months of age.  Our prior research indicates that both sexes

of APP$^{NL-F}$ mice have increased markers of senescent cell (p16$^{Ink4α}$ and p21$^{Cip1}$) and

senescence-associated secretory phenotype (SASP; IL-6, TNFα, MIP1α) in vWAT by 4 months

of age (Fang et al., 2025), which may account for their reduced glucose tolerance (Abi-Ghanem

et al., 2025). Besides metabolic dysfunction, senescent adipocytes also contribute to chronic

systemic inflammation which diminishes cognitive reserve, promotes the accumulation of

misfolded proteins, and accelerates AD progression (Huang et al, 2022).

To test our hypothesis, we surgically resected vWAT (periovarian or epididymal and perirenal,

hereafter perigonadal) from male and female APP$^{NL-F}$ mice at either 4 or 16 months of age.

These time points allowed us to examine the effects of VFR for pre-symptomatic intervention (4

months) or symptomatic therapy (16 months).  All metabolic and cognitive assays were

conducted at 18 months of age.  Our results indicate that VFR effects were disease stage- and

sex-dependent with metabolic and cognitive benefits observed in symptomatic male APP$^{NL-F}$

mice.

**Material and Methods**

<u>Chemicals</u>

All chemicals were stored according to manufacturer recommendations unless stated otherwise.

The Phosphovanilin reagent for lipid determination was prepared by using 6 milligrams of

vanillin (Fisher Scientific Cat # AC140821000) dissolved in 100 mL of 37°C $H_2O$ and further

diluted to 500 mL with 85% phosphoric acid.

<u>Animals</u>

Founder APP$^{NL-F}$ mice on a C57BL/6 background (RRID: IMSR_RBRC06343) were obtained

from Riken (Japan). C57BL/6 (RRID:IMSR_JAX:000664) mice were originally obtained from The

Jackson Laboratory (Bar Harbor, ME) and used to maintain the genetic background of the

breeding colonies.  At weaning, pups were randomly assigned to mixed-litter cages and group-

housed with mice of the same sex and genotype to control for litter size, maternal care, and

other potential litter-specific variables.  Protocols for animal use were approved by the

Institutional Animal Care and Use Committee at Southern Illinois University School of Medicine

(Protocol number: 2025-055), which is accredited by the Association for Assessment and

Accreditation of Laboratory Animal Care. All studies were conducted in accordance with the

United States Public Health Service's Policy on Humane Care and Use of Laboratory Animals.

Mice were group housed on a 12:12 h light-dark cycle, with food (Chow 5001 with 23.4%

protein, 5% fat, and 5.8% crude fiber; LabDiet PMI Feeds) and water available *ad libitum*.

Genotypes were confirmed by collecting a 5 mm tail tip for DNA analysis by TransnetYX, Inc

(Cordova, TN).  The number of mice in each cohort used for the study are shown in Table 1.

<u>VFR Surgery</u>

4 and 16-month-old male and female APP$^{NL-F}$ mice were randomized into two groups – VFR

(perigonadal) and sham-operated.  Mice were isoflurane anesthetized, abdominal fur shaved,

and betadine scrubbed over the incision site.  An ~1.5 cm vertical midline incision along the

lower abdomen in males or ventral incisions on either lateral abdominal side in females were

made to access epidydimal adipose depots. The perinephric adipose depots were removed by

flank incisions. Adipose depots were removed by blunt dissection.  Mice that underwent the

sham operation had their vWAT mobilized, but not removed.  Absorbable sutures were used to

close the body cavity while wound clips were used on the skin. Tylenol (1 mg/mL) was

administered in the drinking water from one day pre- to 3 days post-surgery.  Table 1 shows the

number of mice that were euthanized two weeks or less post-surgery for each group.

<u>Glucose tolerance test (GTT) and insulin tolerance test (ITT)</u>

Glucose metabolism was assessed via GTT and ITT when mice reached 18 months of age (14

or 2 months post VFR surgery) as previously described (Fang et al., 2023). Fifteen-hour-fasted

mice underwent GTT by intraperitoneal (ip) injection with 2.0 g glucose per kg of body weight

(BW). Blood glucose levels were measured at 0, 15, 30, 45, 60, and 120 min with a PRESTO

glucometer (AgaMatrix). For ITT, mice were injected ip with 1 IU insulin glargine (Sanofi-

Aventis) per kg of BW. Blood glucose levels were measured at 0, 15, 30, 45, 60, and 120 min.

<u>NOR task</u>

Object recognition memory was assessed when mice reached ~19 months of age (15 or 3

months post VFR surgery). The NOR was used to evaluate memory retention based on the

premise that mice spend more time exploring a novel object rather than a familiar object if the

animal remembers the familiar object.  Mice were habituated in the open field chamber for 30

min on the first day. 24 hours after, the mouse was returned to the chamber and presented with

two similar objects for 5 min. A 24 hr inter-session-interval was used between introduction and

retention phases to promote long-term memory retrieval.  During the retention phase one of the

familiar objects was replaced with a novel object and the mouse was given 5 minutes of

exploration. The ANY-maze video tracking system (Stoelting Co., Wood Dale, IL) was used to

record mouse navigation during the familiarization and retention phases and used to calculate

the retention index and novelty preference.

<u>Body composition and tissue procurement</u>

At 19 months of age, BW was recorded and unfasted mice were anesthetized using isoflurane.

A cardiac puncture was used to collect the plasma. The blood was mixed with EDTA, followed

by centrifugation at 10,000 g for 15 min at 4°C for plasma collection. Immediately following

cardiac puncture, mice were decapitated and the peripheral tissues (liver, pancreas,

perigonadal and SubQ, WAT, and BAT were weighed to calculate their contribution to total BW.

Tissue was immediately flash frozen and stored at -80°C until processing as previously

described (Fang et al., 2023).

<u>vWAT Histomorphology and Quantification</u>

Frozen vWAT was transferred to ice cold 4% paraformaldehyde in phosphate buffer (pH 7.2)

and fixed at 4°C for 72 hrs. Tissue was then cryoprotected in 30% sucrose in phosphate buffer

(pH 7.2) and stored at 4°C for a minimum of 24 hrs. Following fixation, samples were embedded

in optimal cutting temperature compound (Fisher Scientific Cat #23-730-571) using 15 mm

cryomolds (Fisher Scientific Cat #22-363-553) on dry ice.  Sections (10 µm) were cut using a

Leica CM1950 cryostat (chamber -30°C; specimen holder -40°C) and mounted on gelatin-

coated, charged glass slides (Fisher Scientific, Cat #12-550-15).  Slides were air-dried at room

temperature (~25°C) for at least 15 minutes and stored at 4°C until staining. Slide-mounted

vWAT sections were stained according to the manufacturer's instructions using a commercial

hematoxylin and eosin (H&E) staining kit (Abcam Cat # ab245880).  Following staining and

dehydration, sections were cleared with xylene substitute (Fisher Scientific Cat #9990505) and

coverslipped with Permount mounting medium (Fisher Scientific Cat # SP15-100). Brightfield

images were acquired at 40x magnification using a BZ-X810 All-in-One Flourescence

microscope (Keyence Corp). For each sample, four fields of view from four separate section

were imaged (16 images per sample). Adipocyte number and interior area ($\mu m^2$) were quantified

for each field using the Hybrid Cell Count software (Keyence Corp.). Measurements were

averaged across fields and sections to generate a single mean adipocyte count and sample

area per mouse.

<u>Total Liver Lipid Quantification</u>

A modified micro-scale colorimetric sulfo-phospho-vanillin method was used to measure total

liver lipids (Tran et al., 2019). Approximately 50 mg of liver tissue was homogenized in 1000 $\mu L$

of chloroform/methanol (2:1) followed by addition of 500 $\mu L$ of saline. The mixture was vortexed

then centrifuged at 10,000 rpm for 5 minutes. The chloroform layer was separated and

evaporated at 90°C  for ~20 min, then incubated at 90°C with 200 $\mu L$ of 98% sulfuric acid for 20

min. After adding 750 $\mu L$ of vanillin–phosphoric acid reagent, samples were incubated at 37°C

for 15 min, cooled to 25°C, and 200 $\mu L$ was transferred to a 96-well microplate. Absorbance was

measured at 530 nm, and lipid content was normalized to tissue weight.

<u>Assessment of blood chemistry</u>

Per the manufacturer's protocol, adiponectin or leptin was measured with respective ELISA kits

from Crystal Chem (Elk Grove Village, IL; Cat# 80569, 90030).  Visfatin was measured by an

ELISA kit from Mybiosource (San Diego, CA; Cat# MBS760980).

<u>RT–PCR</u>

mRNA expression was analyzed by quantitative RT–PCR as previously described (Fang et al.,

2020) using Bio-Rad cDNA synthesis kits (Cat # 1708897 and 1708891) and SYBR green mix

(Cat # 1725121) and the QuantStudio 3 Real-Time PCR (Thermo Fisher Scientific). RNA was

extracted using a RNeasy mini kit or RNeasy Lipid Tissue Mini Kit (Qiagen) following the

manufacturer's instructions. Relative expression was calculated as previously described

(Masternak et al., 2012) and primers were purchased from Integrated DNA Technologies.  A list

of forward and reverse primers used in this study is shown in Table 2.

Soluble Aβ$_{42}$ Quantification

The hippocampus from one hemisphere was dissected and stored at -80°C until tissue

processing.  Soluble Aβ$_{42}$ concentrations were determined using the Human / Rat β amyloid

ELISA kits (WAKO Chemicals; Cat: 290-62601) according to the manufacturer recommended

protocols.

Statistical analysis

Statistical analyses were conducted to determine differences between age- and sex-matched

sham and VFR groups.  Statistical tests and post hoc analyses are indicated in each Figure

legend. A Grubb's test was used to identify outliers. Significance was defined as $p < 0.05$ and

investigatory as $p < 0.10$. Data are presented as mean ± SEM. All statistical analyses and

graphs were completed using Prism 10 (GraphPad Inc, La Jolla, CA, USA).

**Results**

<u>VFR at the symptomatic time point improved object recognition memory in male APP$^{NL-F}$ mice.</u>

To explore whether VFR could improve object recognition memory in APP$^{NL-F}$ mice, surgery was

performed at a pre-symptomatic stage (4 months of age, prior to detectable plaque

accumulation and cognitive decline) or a symptomatic stage (16 months of age, during

advanced pathology and functional decline). Both male and female APP$^{NL-F}$ cohorts were

included to assess sex-specific effects of VFR. Object recognition memory was assayed at 18

months of age via the NOR task.  Only male APP$^{NL-F}$ mice that underwent VFR at the

symptomatic time point had improved recognition memory as indicated by increased retention

index (Fig. 1A) and novelty preference (Fig. 1B) compared to the age- and sex-matched sham

surgery controls.  VFR surgery performed at the pre-symptomatic time point in male, or either

time point of female APP$^{NL-F}$, had no effect on object recognition memory (Fig 1A-B).

<u>VFR resulted in changes to BW and composition in an age- and sex-dependent manner.</u>

To assess physiological changes caused by VFR, we examined body composition at ~19

months of age upon euthanasia. VFR decreased BW at the symptomatic stage in male APP$^{NL-F}$

mice compared to the sham group (Fig 1C).  No differences were observed in the other VFR

groups.  The body composition analyses indicated that after VFR, vWAT accumulated in all

ages and sexes but at different percentages. After VFR, APP$^{NL-F}$ mice had vWAT

reaccumulation occurring in 92% and 100% of mice in the pre-symptomatic and symptomatic

cohorts, respectively (Fig. 1D).  In contrast, ~50% of the male mice undergoing VFR surgery at

either age observed vWAT reaccumulation (Fig. 1D). This difference in vWAT was the major

contributor to the observed changes of total percent adipose mass when including SubQ,

interscapular BAT, and vWAT (Fig. 1E). In summary, the BW of symptomatic male APP$^{NL-F}$ mice

after VFR was reduced mainly due to less vWAT reaccumulation.

<u>Reaccumulated vWAT in pre-symptomatic male APP<sup>NL-F</sup> mice had increased adipocyte number</u>

<u>and smaller size.</u>

H&E staining was used to examine if reaccumulated vWAT had morphological differences.

Representative 40x images of pre-symptomatic male sham and VFR APP<sup>NL-F</sup> cohorts are shown

in Fig. 1F.  Reaccumulated vWAT following VFR in this cohort exhibited adipocyte hyperplasia

characterized by a reduced mean internal diameter (Fig. 1G), and may account for the observed

decrease in vWAT mass (Fig. 1E).  No differences were observed in the other VFR treated

groups compared to sham littermates.

<u>VFR during the pre-symptomatic time point in male APP<sup>NL-F</sup> mice caused ectopic liver lipid</u>

<u>deposition via increased de novo lipogenesis.</u>

Although vWAT reaccumulated similarly after VFR regardless of disease stage in male APP<sup>NL-F</sup>

mice, BW at the time of euthanasia was comparable between pre-symptomatic males after VFR

and their sham controls. Body composition changes without effects on BW are known to happen

with aging (St-Onge and Gallagher, 2010).  We sought to identify tissue composition changes

that caused the increased BW despite reduced adiposity in the pre-symptomatic male mice after

VFR.  Lipodystrophy leads to the excessive deposition of lipids in other organs, such as the liver

or muscle, contributing to insulin resistance (Petersen and Shulman, 2018). To test whether the

liver had more ectopic lipid deposition, the percentage of total liver lipids was measured.  Pre-

symptomatic male APP<sup>NL-F</sup> mice had a higher percentage of liver lipids after VFR compared with

sham controls (Fig 2A).  In symptomatic APP<sup>NL-F</sup> male mice, VFR reduced total liver lipids

compared with sham surgery controls while no differences were observed in females at either

disease stage.

Liver gene expression showed that key *de novo* lipogenesis genes including acetyl-coenzyme A

carboxylase (ACC), fatty acid synthase (FAS), stearoyl-CoA desaturase 1 (SCD1), elongation of

long-chain fatty acids family member 6 (Elovl6), sterol regulatory element-binding protein-1

(SREBP1) were increased after VFR in pre-symptomatic male APP[NL-F] mice compared with

sham controls (Fig 2B).  In pre-symptomatic female APP[NL-F] mice, expression of these same

genes (except for ACC) were decreased after VFR.  Interestingly, liver gene expression of

monoglyceride lipase (MGL), the rate-limiting enzyme in lipolysis, was only upregulated after

VFR in symptomatic male APP[NL-F] mice (Fig. 2B). This is one explanation for the decreased BW

and vWAT observed in this group.  These data indicated that VFR in pre-symptomatic male

APP[NL-F] mice cause more ectopic liver lipids likely due to higher *de novo* lipogenesis, while the

contrary was observed in males at the symptomatic disease stage due to higher lipolysis.

<u>VFR in female APP[NL-F] mice led to increased gene expression of de novo lipogenesis and</u>

<u>decreased expression of lipolysis genes in reaccumulated vWAT.</u>

Nearly all the APP[NL-F] female mice reaccumulated vWAT after VFR, with similar morphology to

levels observed in sham surgery controls matched for age and sex (Fig. 1D, G).  To understand

the reason for this reaccumulation, we examined gene expression of lipogenesis and lipolysis in

vWAT (Fig 2C).  Expression levels of two key *de novo* lipogenesis genes, ACC and FAS, were

increased in APP[NL-F] female mice at either disease stage after VFR. The expression of ACC and

FAS were decreased following VFR in male APP[NL-F] mice at the symptomatic, but not at the pre-

symptomatic time point.  Lipolysis gene expression, including protein kinase A (PKA), adipose

triglyceride lipase (ATGL), and hormone sensitive lipase (HSL), were decreased at both disease

stages after VFR in APP[NL-F] female mice compared with sham controls. In male mice, VFR

increased the expression of PKA, ATGL, and HSL at the pre-symptomatic, but unchanged, at

the symptomatic time point. The data suggests that the accumulation of vWAT after VFR in

female APP[NL-F] mice at either stage of disease progression is a result of increased lipogenesis

and decreased lipolysis gene expression.  In male APP[NL-F] mice, the reduced accumulation of

vWAT after VFR was disease stage specific with pre-symptomatic time points increasing

expression of lipolysis genes, while surgery at the symptomatic time point decreased

lipogenesis gene expression.  Interestingly, SREBP1 vWAT expression was upregulated after

VFR only in pre-symptomatic APP[NL-F] male mice. SREBP1 is a transcription factor activated by

high blood glucose (Bertolio et al., 2019, Zhao et al., 2024), suggesting altered glucose

metabolism in this cohort.

<u>VFR surgery impacted either glucose tolerance or insulin sensitivity in male APP[NL-F] mice.</u>

Lipid metabolism plays an important role in regulating glucose utilization, and the two processes

are tightly interconnected (Parhofer, 2015).  Accordingly, the observed changes in lipid

metabolizing gene expression could affect glucose metabolism. To test the effects of VFR on

blood glucose metabolism, we monitored fasting and fed glucose levels as well as glucose

tolerance with a GTT, and insulin sensitivity with an ITT, at 18 months of age.  VFR during the

pre-symptomatic disease stage in APP[NL-F] male mice reduced their glucose tolerance, but had

no effects in symptomatic males (Fig 3A).  VFR improved insulin sensitivity in symptomatic

APP[NL-F] male mice, but effects were not observed during the pre-symptomatic stage in males

(Fig 3B).  Consistent with the GTT, VFR increased fasting glucose in pre-symptomatic APP[NL-F]

male mice compared with sham controls (Fig. 3C).  VFR did not affect fasting blood glucose in

symptomatic APP[NL-F] male mice, nor were differences in fed glucose observed in either disease

stage of the male cohorts (Fig. 3D). Blood glucose levels were unchanged after VFR in either

disease stage of APP[NL-F] female mice.

The liver plays a major role in the control of whole body glucose homeostasis (Han et al., 2016,

Guerra and Gastaldelli, 2020, Scoditti et al., 2024). To understand the observed blood glucose

differences, we examined liver gene expression (Fig 4A) associated with gluconeogenesis

(synthesis of glucose from non-carbohydrate sources), glycolysis (breakdown of glucose for

energy), and glycogenesis (conversion of glucose to glycogen). VFR increased expression of

gluconeogenesis genes, glucose-6-phosphatase (G6PC) and phosphoenolpyruvate

carboxykinase (PEPCK), in pre-symptomatic male mice, consistent with their increased fasting

blood glucose levels (Fig 3C).  In contrast, glucose transporter 2 (Glut2) was increased in

symptomatic males, while GCK (Glucokinase) and glycogen synthase 2 (GYS2) key genes in

glycolysis and glycogenesis, respectively, were decreased in female APP$^{NL-F}$ mice after VFR.

Taken together, the blood glucose and liver gene expression measures indicate VFR affected

glucose metabolism in a sex- and disease stage-dependent manner. Pre-symptomatic APP$^{NL-F}$

male mice had increased fasting blood glucose and made them less glucose tolerant after VFR,

likely mediated via increasing gluconeogenesis. At symptomatic time points, VFR improved

glucose metabolism by increasing insulin sensitivity via enhancing glycolytic and glycogenesis

gene expression.  Although this gene expression was reduced after VFR in symptomatic female

mice, no effects on insulin sensitivity or blood glucose levels were observed.

<u>VFR alters plasma visfatin, adiponectin, and leptin in pre-symptomatic male APP$^{NL-F}$ mice.</u>

Adipokines secreted from adipose tissue are important for whole body glucose metabolism and

insulin sensitivity (Rabe et al., 2008, Kojta et al., 2020).  Given the altered lipid and glucose

metabolism after VFR in the APP$^{NL-F}$ mice, we examined plasma adipokines.  Visfatin is

primarily produced in vWAT, with plasma levels increasing in response to high-fat diet feeding

and obesity (Adeghate, 2008, Sethi and Vidal-Puig, 2005), and is associated with glucose

intolerance (Abdalla, 2022).  VFR increased plasma visfatin in pre-symptomatic male APP$^{NL-F}$

mice compared with age-matched sham controls (Fig 4B), consistent with their higher ectopic

liver accumulation, fasting blood glucose, and glucose intolerance. No plasma visfatin

differences were observed in the other mouse cohorts.

Adiponectin enhances glucose and lipid metabolism (Yanai and Yoshida, 2019), decreases

lipogenesis (Richard et al., 2000), and has neuroprotective effects (Khoramipour et al., 2021).

VFR reduced plasma adiponectin in pre-symptomatic VFR APP$^{NL-F}$ male mice (Fig 4C).  This

would also contribute to the reduced glucose metabolism and increased liver lipids observed in

this cohort. Plasma adiponectin changes were not observed after VFR in the other cohorts.

Leptin regulates appetite and energy expenditure, and its deficiency can disrupt glucose and

lipid metabolism, leading to neuronal damage that impairs neuroplasticity and memory

consolidation (Fernandes et al., 2024). Higher leptin levels are also associated with a lower

dementia risk (Lieb et al., 2009). Plasma leptin was decreased in pre-symptomatic APP$^{NL-F}$

male mice after VFR compared with age-matched sham controls (Fig. 4D). This likely

contributed to the imbalance of glucose and lipid metabolism in this male cohort. Differences in

plasma leptin levels were not observed in the other cohorts of mice.

<u>VFR increases WAT senescent cell and SASP markers in pre-symptomatic male and</u>

<u>symptomatic female APP$^{NL-F}$ mice.</u>

Adipose tissue, particularly vWAT, has the highest burden of senescent cells (Palmer et al.,

2021) and we previously showed senescent cell markers in vWAT were increased in male and

female APP$^{NL-F}$ mice (Fang et al., 2025). VFR can decrease systemic and neuroinflammation in

aged mice (Shin et al., 2015). We examined gene expression of senescent cell and SASP

markers in vWAT (Fig 5A). Gene expression of p16$^{Ink4\alpha}$, p21$^{Cip1}$, and IL-10 were increased in

vWAT after VFR in pre-symptomatic APP$^{NL-F}$ male mice. Expression of vWAT p16$^{Ink4\alpha}$ and

TNFα were also increased in symptomatic APP$^{NL-F}$ female mice (Fig 5A) while no differences in

IL-6 and MCP1 were observed in any cohort (data not shown) after VFR. SASP markers were

also examined in SubQ WAT where VFR increased TNFα, IL-10, IL-6, and MCP1 in pre-

symptomatic APP$^{NL-F}$ male mice (Fig 5B). SubQ IL-6 was also increased in symptomatic

APP$^{NL-F}$ female mice after VFR. Overall, VFR increased gene expression of senescent cell and

SASP markers in WAT of pre-symptomatic APP$^{NL-F}$ male mice, with a modest increase

observed in symptomatic females.

<u>VFR modulates gene expression of hippocampal senescence, SASP, and cognitive markers.</u>

Excess vWAT accumulation is linked to accelerated brain aging, neuroinflammation, and

dementia risk (Moser and Pike, 2016, Oh et al., 2023, Budamagunta et al., 2023).  To

understand how VFR affected hippocampal function, the brain region strongly associated with

learning and memory, we examined gene expression of cell senescence and cytokines (Fig 6A).

VFR had no effect on p16$^{Ink4\alpha}$, p21$^{Cip1}$, or TNFα expression in APP$^{NL-F}$ male mice at either stage

of disease progression.  Alternatively, p16$^{Ink4\alpha}$ gene expression was increased after VFR in pre-

symptomatic APP$^{NL-F}$ female mice.  VFR caused a corresponding TNFα gene expression

increase in this cohort as well as the symptomatic females. VFR had deleterious effects on

hippocampal senescence and inflammation in female, but not male APP$^{NL-F}$ mice.

Since we observed improvements in object recognition memory after VFR in symptomatic male

APP$^{NL-F}$ mice, we examined markers associated with learning and memory in the hippocampus

(Fig 6A).  Both adiponectin and leptin receptors (AdipoR, LepR) are found in the hippocampus

and play a role in synaptic plasticity and cognition. Gene expression of both receptors were

upregulated after VFR in symptomatic APP$^{NL-F}$ male mice (Fig 6A).  In females, VFR increased

LepR in the pre-symptomatic disease stage.  Cyclic AMP response element-binding protein 1

(Creb1), plays a role in long-term memory formation.  VFR decreased Creb1 gene expression in

pre-symptomatic APP$^{NL-F}$ male mice, but elevated expression in the symptomatic male cohort.

Progranulin (PRGN) is important for neuronal survival and growth in the hippocampus.  PRGN

gene expression was decreased after VFR in pre-symptomatic APP$^{NL-F}$ female mice, but elicited

an increase in symptomatic males. Overall, VFR altered the hippocampal gene expression

profile of symptomatic APP$^{NL-F}$ male mice in a manner that would improve their object

recognition memory.

<u>VFR increases soluble Aβ$_{42}$ in pre-symptomatic male and female mice.</u>

Soluble $A\beta_{42}$ is the neurotoxic species associated with synaptic dysfunction and cognitive

decline in AD (Findley et al., 2019).  An ELISA indicated hippocampal soluble $A\beta_{42}$ (Fig 6B)

production was only increased in pre-symptomatic stage of $APP^{NL-F}$ male and female mice after

VFR.  VFR did not affect soluble $A\beta_{42}$ during the symptomatic stage in either sex.

**Discussion**

Adipose tissue accumulation and distribution exhibit sex-specific differences, yet these patterns are evolutionarily conserved across humans and mice (Chusyd et al., 2016).  Females typically have a higher overall fat percentage with a large proportion stored subcutaneously for energy supply during pregnancy and lactation during food scarcity, while males accumulate fat in visceral regions. Numerous variables influence adipose development including sex chromosomes, gonadal hormone levels, and hormone receptors. Sex chromosome complement impacts adipose accumulation whereby the presence of two "X" chromosomes leads to higher accumulation while the presence of the "Y" chromosome has no influence. The gonadal hormones, testosterone and estrogen, drive vWAT and SubQ accumulation in males and females, respectively. Age-related hormonal changes increase vWAT accumulation regardless of sex and is a key factor in the development of metabolic syndrome that increases dementia risk.

Besides its role as an energy reservoir, adipose tissue is an endocrine organ that secretes peptides such as TNFα, IL-6, and adiponectin, which can influence either tissue-specific or whole body metabolism. Prior studies examining VFR effects in rodents were conducted to elucidate mechanisms contributing to metabolic improvements. VFR improves insulin sensitivity and glucose tolerance in young, aged, and obese rodents with some of the metabolic changes occurring through decreased free fatty acids, leptin, and TNFα expression (Kim et al., 1999, Gabriely et al., 2002, Shi et al., 2007a). VFR in aged C57BL/6 mice led to reduced expression of pro-inflammatory cytokines in the plasma, WAT, and cortex (Shin et al., 2015).  Although adiponectin plays a key role in glucose and lipid metabolism and suppresses inflammatory cytokines, serum levels are found to decrease following VFR in male and female Sprague Dawley rats (Borst et al., 2005, Ling et al., 2014), which may be due to the absence of these

WAT depots. Considering the metabolic benefits of VFR and the known link between metabolic

dysfunction and AD risk, we aimed to investigate the impact of VFR in an AD model.

Surprisingly, the VFR response was disease stage- and sex-dependent. The removal of vWAT

in symptomatic APP[NL-F] male mice improved object recognition memory potentially due to

increased hippocampal AdipoR, LepR, Creb1, and PRGN expression.  Lipogenesis vWAT gene

expression was down-regulated, while hepatic genes associated with lipolysis and glucose

transport were up-regulated.  These effects were associated with a decreased percentage of

total liver lipids. All of these alterations were reflected by a reduced percentage of total fat,

specifically reaccumulated vWAT, BW, and improved blood glucose maintenance.  The opposite

was true for VFR during the pre-symptomatic stage in males.  After VFR, an imbalance of lipid

metabolism was prevalent, reflected by an elevated expression of vWAT lipolysis and hepatic

lipogenesis genes.  This tissue metabolic imbalance may be a result of adipocyte hyperplasia

with reduced cell size as well as the redistribution of lipids into other organs, such as the liver,

where higher ectopic fat accumulation was observed.  Plasma adiponectin and leptin were

decreased along with increased visfatin resulting in impaired glucose metabolism and peripheral

inflammation. Typically, adiponectin and leptin are inversely related, but plasma levels of both

were decreased in pre-symptomatic male APP[NL-F] mice. The decrease in leptin might be due to

the loss of the adipose depot while the anticipated compensatory increase in adiponectin may

not have been observed because the adiponectin source was also depleted. Further studies are

needed to determine why this effect was not observed in the other cohorts after VFR.  In the

brain, reduced hippocampal Creb1 expression and increased soluble $A\beta_{42}$, combined with the

aforementioned adverse physiological changes, failed to improve object recognition memory.

In contrast to males, vWAT reaccumulated in nearly all female mice and was similar in number

and size to sham littermates regardless of when VFR occurred. This could be attributed to

higher lipogenesis and lower lipolysis gene expression.  Although ectopic lipid deposition did not

accumulate in the liver, VFR in pre-symptomatic female APP$^{NL-F}$ mice led to decreased hepatic

lipogenesis gene expression, while expression of genes involved in glycolysis and glycogenesis

were decreased in symptomatic females. These results were in opposition to age-matched

males. Hippocampal soluble A$\beta_{42}$, senescent cell, and inflammatory markers were elevated in

pre-symptomatic APP$^{NL-F}$ female mice while the later was only observed at the symptomatic

time point. No changes to blood glucose levels or object recognition memory were observed in

females after VFR.

Lipectomy in mice, rats, and other small mammals does not result in long-term BW loss with

regain occurring 12-14 weeks post-surgery (Murillo et al., 2019). In the majority of the studies

this was attributed to regeneration of the lipectomized depot, redistribution of fat to other

adipose depots, or ectopic fat accumulation in other organs. In our present study, female

APP$^{NL-F}$ mice regained BW due to reaccumulating vWAT. Although vWAT accumulation was

lower in pre-symptomatic males, they regained BW due to ectopic fat distribution in the liver and

possibly other organs.  Only symptomatic APP$^{NL-F}$ male mice had lower BW attributed to less

visceral adiposity.  Redistribution of fat to other adipose depots was not observed in any cohort.

Our present study was designed to examine disease stage-specific interventional or therapeutic

AD strategies rather than examining the contribution of aging on adiposity and AD progression.

Regardless, effects of aging on adipose accumulation likely contributed to the observed

differences in lipid distribution and the comparable end-of-study BW between sham and VFR

cohorts matched for age and sex.  The gradual decline in estrous cycling and the associated

drop in estrogen levels, which promotes visceral rather than SubQ accumulation, may partially

explain post-resection visceral accumulation in female mice, whereas aging males exhibit

ectopic liver lipid deposition (Donato et al., 2014).

The exact mechanisms behind this adipose rebound are not well known, but suggest biological

feedback systems that act to preserve energy balance.  For example, the adipostat hypothesis

postulates that alterations to energy reserves triggers compensatory mechanisms in metabolism

and/or thermogenesis to restore the reservoir.  Since gonadal hormones and sex chromosome

complement mediate adipose storage, distribution, and utilization, male and female mice use

different behavioral strategies to restore fat after lipectomy. Males increase caloric consumption

while females reduce heat production and energy expenditure (Shi et al., 2007b).  Our findings

are in support of this whereby pre-symptomatic APP$^{NL-F}$ male mice had increased

gluconeogenesis gene expression and fasting blood glucose, and worse glucose tolerance after

VFR.  While VFR in Female APP$^{NL-F}$ mice reduced glycolysis and lipolysis, but increased

lipogenesis gene expression.  It is important to note, however, that our study was not intended

to serve as a preclinical model for liposuction, as that procedure primarily targets SubQ rather

than vWAT depots.  However, lipectomy in humans causes a transient decrease in BW and

adipose depots that gradually return to preoperative levels within months (Hernandez et al.,

2011, Seretis et al., 2015), suggesting the preservation of this energy balance is conserved

across species.

The accumulation of WAT and subsequent increases in BW begins in midlife and continues into

early aging in both humans (40-65 years) and mice (12-18 months). The reason for this

increase is attributed to adipogenesis and adipocyte hypertrophy that are prominent during

midlife (Wang et al., 2025). Remodeling of vWAT, like its distribution, varies between sexes,

with estrogen levels playing a key role in regulating these processes and shifting notably with

age. Female mice have increased adipose progenitor cells leading to hyperplasia while males

experience hypertrophy (Wu et al., 2017).  Both increase adipose accumulation, but hyperplasia

is associated with better metabolic health (Steiner and Berry, 2022), which may explain why

blood glucose levels were unaffected in female mice after VFR.  Hypertrophic adipocytes

secrete reduced adiponectin and increased pro-inflammatory cytokines, the latter promoting

ectopic lipid distribution into peripheral organs.  These pathological features were evident

following VFR in pre-symptomatic APP$^{NL-F}$ male mice, despite the reaccumulated vWAT

exhibiting a less hypertrophic phenotype. Accordingly, adipose cell size rather than number

correlates with increased risk for metabolic syndrome (Vishvanath and Gupta, 2019).

Senescent cell accumulation in both the periphery and brain contribute to aging and age-related

neurodegenerative disorders such as AD. The mechanisms behind this are multifaceted and

dependent upon the type of senescent cell (Zhu et al., 2024). Senescent brain cells alter

amyloid precursor protein processing by shifting it from the non-amyloidogenic to the

amyloidogenic pathway, promoting the formation of Aβ (Sun et al., 2018). The pro-inflammatory

SASP response coupled with decreased microglial phagocytosis of misfolded proteins can lead

to the subsequent Aβ accumulation and deposition in the brain. We observed elevated

hippocampal soluble Aβ$_{42}$ after VFR in pre-symptomatic APP$^{NL-F}$ mice compared with sex-

matched sham controls, which was contradictory to our hypothesis. VFR increased WAT

senescent cell burden and SASP in pre-symptomatic males, while these were increased in the

hippocampus of pre-symptomatic females. Considering senescent cell and SASP increases

were predominantly observed in the pre-symptomatic groups, this may indicate a long-term

effect of VFR. In AD mouse models, reducing peripheral and hippocampal senescent cell

burden with senolytics has been shown to reduce amyloid pathology and improve cognitive

performance - albeit with benefits favoring female mice (Currais et al., 2014, Zhang et al., 2019,

Fang et al., 2025, Ng et al., 2024).

Given that AD pathology is thought to initiate in midlife prior to cognitive decline, the concurrent

increase in adipose tissue accumulation during this period may play a causal role in disease

progression.  Our present study indicates accumulation or redistribution of adipose tissue during

the pre-symptomatic AD stage can cause metabolic disturbances resulting in sex-specific

peripheral and cerebral senescent cell accumulation that exacerbates pathological AD

progression.  However, during symptomatic AD stages, elimination of vWAT has beneficial

metabolic and cognitive effects in male, but not female APP$^{NL-F}$ mice. These sex-specific

differences are attributed to increasing energy storage in female symptomatic mice.

Several study limitations should be noted when interpreting the results of the present study.

Although the study design was focused on identifying within-model sex-specific and disease-

stage dependent alterations relevant for ameliorating AD progression; inclusion of age-matched

genetic background control C57BL/6 mice could enhance the interpretability and translational

relevance of the findings. Second, we did not measure circulating gonadal hormones since our

study design did not include repeated sampling or estrous cycle staging. A single-point

hormone measurement would not have yielded biologically meaningful data due to the large

physiological variability inherent in their sampling. Third, we only measured ectopic lipid

deposition in the liver, so therefore it is possible for lipids to accumulate in other organs such as

muscle. Finally, we focused on specific comparisons of sham and VFR surgery animals to their

age- and sex-matched controls, consistent with the hypothesis, by means of a t-test. This was

done to avoid exploratory-type findings for possible age and sex effects since our primary goal

was to assess whether VFR improved cognitive performance at distinct disease stages within

each sex.

**Disclosures**

The authors state that they have no financial or non-financial competing interests to disclose.

SAM, YF, MRP, KQ, TH, and JEC conducted the experiments and analyzed the data. YF, AB,

KNH, and ERH conceived the study, designed the experiments, interpreted the data, and wrote

the manuscript. All authors approved the final version of the manuscript.

**Funding Sources**

This work was supported by the National Institutes of Health NIA R01-AG057767 and NIA R01-

AG061937, Kenneth Stark Endowment, and Dale and Deborah Smith Center for Alzheimer's

Research and Treatment (YF, MRP, KQ, JEC, SM, TH, KNH, ERH). Geriatrics Initiative (AB).

## References

**References**

ABDALLA, M. M. I. 2022. Role of visfatin in obesity-induced insulin resistance. *World J Clin Cases,* 10**,**
10840-10851.
ABI-GHANEM, C., KELLY, R. D., GROOM, E. A., VALERIAN, C. G., PAUL, A. S., THRASHER, C. A., SALINERO,
A. E., BATCHELDER, M. R., LAFRICAN, J. J., WANG, M., SMITH, R. M., TEMPLE, S., ZULOAGA, D. G.
& ZULOAGA, K. L. 2025. Interactions between menopause and high-fat diet on cognition and
pathology in a mouse model of Alzheimer's disease. *Alzheimers Dement,* 21**,** e70026.
ADEGHATE, E. 2008. Visfatin: structure, function and relation to diabetes mellitus and other
dysfunctions. *Curr Med Chem,* 15**,** 1851-62.
BAGLIETTO-VARGAS, D., SHI, J., YAEGER, D. M., AGER, R. & LAFERLA, F. M. 2016. Diabetes and
Alzheimer's disease crosstalk. *Neurosci Biobehav Rev,* 64**,** 272-87.
BARZILAI, N., SHE, L., LIU, L., WANG, J., HU, M., VUGUIN, P. & ROSSETTI, L. 1999. Decreased visceral
adiposity accounts for leptin effect on hepatic but not peripheral insulin action. *Am J Physiol,*
277**,** E291-8.
BENITEZ, D. P., JIANG, S., WOOD, J., WANG, R., HALL, C. M., PEERBOOM, C., WONG, N., STRINGER, K. M.,
VITANOVA, K. S., SMITH, V. C., JOSHI, D., SAITO, T., SAIDO, T. C., HARDY, J., HANRIEDER, J., DE
STROOPER, B., SALIH, D. A., TRIPATHI, T., EDWARDS, F. A. & CUMMINGS, D. M. 2021. Knock-in
models related to Alzheimer's disease: synaptic transmission, plaques and the role of microglia.
*Mol Neurodegener,* 16**,** 47.
BERTOLIO, R., NAPOLETANO, F., MANO, M., MAURER-STROH, S., FANTUZ, M., ZANNINI, A., BICCIATO, S.,
SORRENTINO, G. & DEL SAL, G. 2019. Sterol regulatory element binding protein 1 couples
mechanical cues and lipid metabolism. *Nat Commun,* 10**,** 1326.
BIESSELS, G. J. & DESPA, F. 2018. Cognitive decline and dementia in diabetes mellitus: mechanisms and
clinical implications. *Nat Rev Endocrinol,* 14**,** 591-604.
BORST, S. E., CONOVER, C. F. & BAGBY, G. J. 2005. Association of resistin with visceral fat and muscle
insulin resistance. *Cytokine,* 32**,** 39-44.
BREIJYEH, Z. & KARAMAN, R. 2020. Comprehensive Review on Alzheimer's Disease: Causes and
Treatment. *Molecules,* 25.
BUDAMAGUNTA, V., KUMAR, A., RANI, A., BEAN, L., MANOHAR-SINDHU, S., YANG, Y., ZHOU, D. &
FOSTER, T. C. 2023. Effect of peripheral cellular senescence on brain aging and cognitive decline.
*Aging Cell,* 22**,** e13817.
CHUSYD, D. E., WANG, D., HUFFMAN, D. M. & NAGY, T. R. 2016. Relationships between Rodent White
Adipose Fat Pads and Human White Adipose Fat Depots. *Front Nutr,* 3**,** 10.
CURRAIS, A., PRIOR, M., DARGUSCH, R., ARMANDO, A., EHREN, J., SCHUBERT, D., QUEHENBERGER, O. &
MAHER, P. 2014. Modulation of p25 and inflammatory pathways by fisetin maintains cognitive
function in Alzheimer's disease transgenic mice. *Aging Cell,* 13**,** 379-90.
DONATO, A. J., HENSON, G. D., HART, C. R., LAYEC, G., TRINITY, J. D., BRAMWELL, R. C., ENZ, R. A.,
MORGAN, R. G., REIHL, K. D., HAZRA, S., WALKER, A. E., RICHARDSON, R. S. & LESNIEWSKI, L. A.
2014. The impact of ageing on adipose structure, function and vasculature in the B6D2F1
mouse: evidence of significant multisystem dysfunction. *J Physiol,* 592**,** 4083-96.
DOVE, A., SHANG, Y., XU, W., GRANDE, G., LAUKKA, E. J., FRATIGLIONI, L. & MARSEGLIA, A. 2021. The
impact of diabetes on cognitive impairment and its progression to dementia. *Alzheimers*
*Dement,* 17**,** 1769-1778.
ELIAS, M. F., ELIAS, P. K., SULLIVAN, L. M., WOLF, P. A. & D'AGOSTINO, R. B. 2005. Obesity, diabetes and
cognitive deficit: The Framingham Heart Study. *Neurobiol Aging,* 26 Suppl 1**,** 11-6.

FANG, Y., MCFADDEN, S., DARCY, J., HASCUP, E. R., HASCUP, K. N. & BARTKE, A. 2020. Lifespan of long-lived growth hormone receptor knockout mice was not normalized by housing at 30 degrees C since weaning. *Aging Cell,* 19**,** e13123.

FANG, Y., MEDINA, D., STOCKWELL, R., MCFADDEN, S., QUINN, K., PECK, M. R., BARTKE, A., HASCUP, K. N. & HASCUP, E. R. 2023. Sexual dimorphic metabolic and cognitive responses of C57BL/6 mice to Fisetin or Dasatinib and quercetin cocktail oral treatment. *Geroscience,* 45**,** 2835-2850.

FANG, Y., PECK, M. R., QUINN, K., CHAPMAN, J. E., MEDINA, D., MCFADDEN, S. A., BARTKE, A., HASCUP, E. R. & HASCUP, K. N. 2025. Senolytic intervention improves cognition, metabolism, and adiposity in female APP(NL)(-F/NL-F) mice. *Geroscience,* 47**,** 1123-1138.

FERNANDES, C., FORNY-GERMANO, L., ANDRADE, M. M., LYRA, E. S. N. M., RAMOS-LOBO, A. M., MEIRELES, F., TOVAR-MOLL, F., HOUZEL, J. C., DONATO, J., JR. & DE FELICE, F. G. 2024. Leptin receptor reactivation restores brain function in early-life Lepr-deficient mice. *Brain,* 147**,** 2706-2717.

FINDLEY, C. A., BARTKE, A., HASCUP, K. N. & HASCUP, E. R. 2019. Amyloid Beta-Related Alterations to Glutamate Signaling Dynamics During Alzheimer's Disease Progression. *ASN Neuro,* 11**,** 1759091419855541.

FOSTER, M. T., SHI, H., SEELEY, R. J. & WOODS, S. C. 2011. Removal of intra-abdominal visceral adipose tissue improves glucose tolerance in rats: role of hepatic triglyceride storage. *Physiol Behav,* 104**,** 845-54.

FRANCZYK, M. P., HE, M. & YOSHINO, J. 2021. Removal of Epididymal Visceral Adipose Tissue Prevents Obesity-Induced Multi-organ Insulin Resistance in Male Mice. *J Endocr Soc,* 5**,** bvab024.

GABRIELY, I., MA, X. H., YANG, X. M., ATZMON, G., RAJALA, M. W., BERG, A. H., SCHERER, P., ROSSETTI, L. & BARZILAI, N. 2002. Removal of visceral fat prevents insulin resistance and glucose intolerance of aging: an adipokine-mediated process? *Diabetes,* 51**,** 2951-8.

GUERRA, S. & GASTALDELLI, A. 2020. The role of the liver in the modulation of glucose and insulin in non alcoholic fatty liver disease and type 2 diabetes. *Curr Opin Pharmacol,* 55**,** 165-174.

GUO, D. H., YAMAMOTO, M., HERNANDEZ, C. M., KHODADADI, H., BABAN, B. & STRANAHAN, A. M. 2020. Visceral adipose NLRP3 impairs cognition in obesity via IL-1R1 on CX3CR1+ cells. *J Clin Invest,* 130**,** 1961-1976.

GUSTAFSON, D. R. 2012. Adiposity and cognitive decline: underlying mechanisms. *J Alzheimers Dis,* 30 Suppl 2**,** S97-112.

HAN, H. S., KANG, G., KIM, J. S., CHOI, B. H. & KOO, S. H. 2016. Regulation of glucose metabolism from a liver-centric perspective. *Exp Mol Med,* 48**,** e218.

HERNANDEZ, T. L., KITTELSON, J. M., LAW, C. K., KETCH, L. L., STOB, N. R., LINDSTROM, R. C., SCHERZINGER, A., STAMM, E. R. & ECKEL, R. H. 2011. Fat redistribution following suction lipectomy: defense of body fat and patterns of restoration. *Obesity (Silver Spring),* 19**,** 1388-95.

HUANG, X., WANG, Y. J. & XIANG, Y. 2022. Bidirectional communication between brain and visceral white adipose tissue: Its potential impact on Alzheimer's disease. *EBioMedicine,* 84**,** 104263.

KAMOGAWA, K., KOHARA, K., TABARA, Y., UETANI, E., NAGAI, T., YAMAMOTO, M., IGASE, M. & MIKI, T. 2010. Abdominal fat, adipose-derived hormones and mild cognitive impairment: the J-SHIPP study. *Dement Geriatr Cogn Disord,* 30**,** 432-9.

KHORAMIPOUR, K., CHAMARI, K., HEKMATIKAR, A. A., ZIYAIYAN, A., TAHERKHANI, S., ELGUINDY, N. M. & BRAGAZZI, N. L. 2021. Adiponectin: Structure, Physiological Functions, Role in Diseases, and Effects of Nutrition. *Nutrients,* 13.

KIM, S., YI, H. A., WON, K. S., LEE, J. S. & KIM, H. W. 2022. Association between Visceral Adipose Tissue Metabolism and Alzheimer's Disease Pathology. *Metabolites,* 12.

KIM, Y. W., KIM, J. Y. & LEE, S. K. 1999. Surgical removal of visceral fat decreases plasma free fatty acid
and increases insulin sensitivity on liver and peripheral tissue in monosodium glutamate (MSG)-
obese rats. *J Korean Med Sci,* 14**,** 539-45.
KOJTA, I., CHACINSKA, M. & BLACHNIO-ZABIELSKA, A. 2020. Obesity, Bioactive Lipids, and Adipose Tissue
Inflammation in Insulin Resistance. *Nutrients,* 12.
LIEB, W., BEISER, A. S., VASAN, R. S., TAN, Z. S., AU, R., HARRIS, T. B., ROUBENOFF, R., AUERBACH, S.,
DECARLI, C., WOLF, P. A. & SESHADRI, S. 2009. Association of plasma leptin levels with incident
Alzheimer disease and MRI measures of brain aging. *JAMA,* 302**,** 2565-72.
LING, B. L., CHIU, C. T., LU, H. C., LIN, J. J., KUO, C. Y. & CHOU, F. P. 2014. Short and long-term impact of
lipectomy on expression profile of hepatic anabolic genes in rats: a high fat and high cholesterol
diet-induced obese model. *PLoS One,* 9**,** e108717.
MASTERNAK, M. M., BARTKE, A., WANG, F., SPONG, A., GESING, A., FANG, Y., SALMON, A. B., HUGHES, L.
F., LIBERATI, T., BOPARAI, R., KOPCHICK, J. J. & WESTBROOK, R. 2012. Metabolic effects of intra-
abdominal fat in GHRKO mice. *Aging Cell,* 11**,** 73-81.
MOSER, V. A. & PIKE, C. J. 2016. Obesity and sex interact in the regulation of Alzheimer's disease.
*Neurosci Biobehav Rev,* 67**,** 102-18.
MURILLO, A. L., KAISER, K. A., SMITH, D. L., JR., PETERSON, C. M., AFFUSO, O., TIWARI, H. K. & ALLISON,
D. B. 2019. A Systematic Scoping Review of Surgically Manipulated Adipose Tissue and the
Regulation of Energetics and Body Fat in Animals. *Obesity (Silver Spring),* 27**,** 1404-1417.
NG, P. Y., ZHANG, C., LI, H. & BAKER, D. J. 2024. Senescence Targeting Methods Impact Alzheimer's
Disease Features in 3xTg Mice. *J Alzheimers Dis,* 97**,** 1751-1763.
NGUYEN, J. C., KILLCROSS, A. S. & JENKINS, T. A. 2014. Obesity and cognitive decline: role of
inflammation and vascular changes. *Front Neurosci,* 8**,** 375.
OH, H. S., RUTLEDGE, J., NACHUN, D., PALOVICS, R., ABIOSE, O., MORAN-LOSADA, P., CHANNAPPA, D.,
UREY, D. Y., KIM, K., SUNG, Y. J., WANG, L., TIMSINA, J., WESTERN, D., LIU, M., KOHLFELD, P.,
BUDDE, J., WILSON, E. N., GUEN, Y., MAURER, T. M., HANEY, M., YANG, A. C., HE, Z., GREICIUS,
M. D., ANDREASSON, K. I., SATHYAN, S., WEISS, E. F., MILMAN, S., BARZILAI, N., CRUCHAGA, C.,
WAGNER, A. D., MORMINO, E., LEHALLIER, B., HENDERSON, V. W., LONGO, F. M.,
MONTGOMERY, S. B. & WYSS-CORAY, T. 2023. Organ aging signatures in the plasma proteome
track health and disease. *Nature,* 624**,** 164-172.
PALMER, A. K., TCHKONIA, T. & KIRKLAND, J. L. 2021. Senolytics: Potential for Alleviating Diabetes and Its
Complications. *Endocrinology,* 162.
PARHOFER, K. G. 2015. Interaction between Glucose and Lipid Metabolism: More than Diabetic
Dyslipidemia. *Diabetes Metab J,* 39**,** 353-62.
PETERSEN, M. C. & SHULMAN, G. I. 2018. Mechanisms of Insulin Action and Insulin Resistance. *Physiol*
*Rev,* 98**,** 2133-2223.
PITOMBO, C., ARAUJO, E. P., DE SOUZA, C. T., PAREJA, J. C., GELONEZE, B. & VELLOSO, L. A. 2006.
Amelioration of diet-induced diabetes mellitus by removal of visceral fat. *J Endocrinol,* 191**,** 699-
706.
RABE, K., LEHRKE, M., PARHOFER, K. G. & BROEDL, U. C. 2008. Adipokines and insulin resistance. *Mol*
*Med,* 14**,** 741-51.
RICHARD, A. J., WHITE, U., ELKS, C. M. & STEPHENS, J. M. 2000. Adipose Tissue: Physiology to Metabolic
Dysfunction. *In:* FEINGOLD, K. R., AHMED, S. F., ANAWALT, B., BLACKMAN, M. R., BOYCE, A.,
CHROUSOS, G., CORPAS, E., DE HERDER, W. W., DHATARIYA, K., DUNGAN, K., HOFLAND, J.,
KALRA, S., KALTSAS, G., KAPOOR, N., KOCH, C., KOPP, P., KORBONITS, M., KOVACS, C. S.,
KUOHUNG, W., LAFERRERE, B., LEVY, M., MCGEE, E. A., MCLACHLAN, R., MUZUMDAR, R.,
PURNELL, J., REY, R., SAHAY, R., SHAH, A. S., SINGER, F., SPERLING, M. A., STRATAKIS, C. A.,
TRENCE, D. L. & WILSON, D. P. (eds.) *Endotext.* South Dartmouth (MA).

SAITO, T., MATSUBA, Y., MIHIRA, N., TAKANO, J., NILSSON, P., ITOHARA, S., IWATA, N. & SAIDO, T. C.
2014. Single App knock-in mouse models of Alzheimer's disease. *Nat Neurosci,* 17**,** 661-3.
SAKERS, A., DE SIQUEIRA, M. K., SEALE, P. & VILLANUEVA, C. J. 2022. Adipose-tissue plasticity in health
and disease. *Cell,* 185**,** 419-446.
SCODITTI, E., SABATINI, S., CARLI, F. & GASTALDELLI, A. 2024. Hepatic glucose metabolism in the
steatotic liver. *Nat Rev Gastroenterol Hepatol,* 21**,** 319-334.
SERETIS, K., GOULIS, D. G., KOLIAKOS, G. & DEMIRI, E. 2015. Short- and Long-Term Effects of Abdominal
Lipectomy on Weight and Fat Mass in Females: a Systematic Review. *Obes Surg,* 25**,** 1950-8.
SETHI, J. K. & VIDAL-PUIG, A. 2005. Visfatin: the missing link between intra-abdominal obesity and
diabetes? *Trends Mol Med,* 11**,** 344-7.
SHI, H., STRADER, A. D., WOODS, S. C. & SEELEY, R. J. 2007a. The effect of fat removal on glucose
tolerance is depot specific in male and female mice. *Am J Physiol Endocrinol Metab,* 293**,** E1012-
20.
SHI, H., STRADER, A. D., WOODS, S. C. & SEELEY, R. J. 2007b. Sexually dimorphic responses to fat loss
after caloric restriction or surgical lipectomy. *Am J Physiol Endocrinol Metab,* 293**,** E316-26.
SHIN, J. A., JEONG, S. I., KIM, M., YOON, J. C., KIM, H. S. & PARK, E. M. 2015. Visceral adipose tissue
inflammation is associated with age-related brain changes and ischemic brain damage in aged
mice. *Brain Behav Immun,* 50**,** 221-231.
ST-ONGE, M. P. & GALLAGHER, D. 2010. Body composition changes with aging: the cause or the result of
alterations in metabolic rate and macronutrient oxidation? *Nutrition,* 26**,** 152-5.
STEINER, B. M. & BERRY, D. C. 2022. The Regulation of Adipose Tissue Health by Estrogens. *Front
Endocrinol (Lausanne),* 13**,** 889923.
SUN, R., HE, T., PAN, Y. & KATUSIC, Z. S. 2018. Effects of senescence and angiotensin II on expression and
processing of amyloid precursor protein in human cerebral microvascular endothelial cells.
*Aging (Albany NY),* 10**,** 100-114.
TCHKONIA, T., MORBECK, D. E., VON ZGLINICKI, T., VAN DEURSEN, J., LUSTGARTEN, J., SCRABLE, H.,
KHOSLA, S., JENSEN, M. D. & KIRKLAND, J. L. 2010. Fat tissue, aging, and cellular senescence.
*Aging Cell,* 9**,** 667-84.
TRAN, T. M., DAI, T. X. T., TRAN, D. B., NGUYEN, Q. C. T. & NGUYEN, D. H. Y. 2019. A simple
spectrophotometric method for quantifying total lipids in plants and animals. *CTU Journal of
Innovation and Sustainable Development,* 11**,** 106-110.
VISHVANATH, L. & GUPTA, R. K. 2019. Contribution of adipogenesis to healthy adipose tissue expansion
in obesity. *J Clin Invest,* 129**,** 4022-4031.
WANG, G., LI, G., SONG, A., ZHAO, Y., YU, J., WANG, Y., DAI, W., SALAS, M., QIN, H., MEDRANO, L., DOW,
J., LI, A., ARMSTRONG, B., FUEGER, P. T., YU, H., ZHU, Y., SHAO, M., WU, X., JIANG, L., CAMPISI,
J., YANG, X. & WANG, Q. A. 2025. Distinct adipose progenitor cells emerging with age drive
active adipogenesis. *Science,* 388**,** eadj0430.
WHITMER, R. A., GUSTAFSON, D. R., BARRETT-CONNOR, E., HAAN, M. N., GUNDERSON, E. P. & YAFFE, K.
2008. Central obesity and increased risk of dementia more than three decades later. *Neurology,*
71**,** 1057-64.
WONG ZHANG, D. E., TRAN, V., VINH, A., DINH, Q. N., DRUMMOND, G. R., SOBEY, C. G., JELINIC, M. & DE
SILVA, T. M. 2023. Pathophysiological Links Between Obesity and Dementia. *Neuromolecular
Med,* 25**,** 451-456.
WU, Y., LEE, M. J., IDO, Y. & FRIED, S. K. 2017. High-fat diet-induced obesity regulates MMP3 to
modulate depot- and sex-dependent adipose expansion in C57BL/6J mice. *Am J Physiol
Endocrinol Metab,* 312**,** E58-E71.
YANAI, H. & YOSHIDA, H. 2019. Beneficial Effects of Adiponectin on Glucose and Lipid Metabolism and
Atherosclerotic Progression: Mechanisms and Perspectives. *Int J Mol Sci,* 20.

ZHANG, P., KISHIMOTO, Y., GRAMMATIKAKIS, I., GOTTIMUKKALA, K., CUTLER, R. G., ZHANG, S.,
ABDELMOHSEN, K., BOHR, V. A., MISRA SEN, J., GOROSPE, M. & MATTSON, M. P. 2019. Senolytic
therapy alleviates Abeta-associated oligodendrocyte progenitor cell senescence and cognitive
deficits in an Alzheimer's disease model. *Nat Neurosci,* 22**,** 719-728.
ZHAO, Z., XIANG, X., CHEN, Q., DU, J., ZHU, S., XU, X., SHEN, Y., WEN, S., LI, Y., XU, W., MAI, K. & AI, Q.
2024. Sterol Regulatory Element Binding Protein 1: A Mediator for High-Fat Diet-Induced
Hepatic Gluconeogenesis and Glucose Intolerance in Fish. *J Nutr,* 154**,** 1505-1516.
ZHU, J., WU, C. & YANG, L. 2024. Cellular senescence in Alzheimer's disease: from physiology to
pathology. *Transl Neurodegener,* 13**,** 55.

**Figure Legends**

*Figure 1 - NOR and body composition analysis after VFR.* The retention index (A) and novelty

preference (B) for the NOR task were calculated at 18 months of age for all groups. BW (C) and

the percentage of vWAT (D) and total fat mass for BAT, SubQ, and vWAT (E) in relation to BW

were determined upon euthanization at 19 months of age. The percentage of VFR mice that

regenerated vWAT (reg. ratio) are shown in (D). Representative 40x images of H&E stained

vWAT in pre-symptomatic male sham (top) and VFR (bottom) cohorts (F). The average number

of vWAT adipocytes and their internal diameter were determined for all groups (G). Data are

represented as means ± SEM. *p<0.05, **p<0.01 based on a two-tailed Welch's *t* test (A-C, E),

a two-way ANOVA with Sidak's post hoc (D), and an unpaired two-tailed t test (G); n=4-20.

*Figure 2 – Lipid metabolism in hepatic and vWAT.* The sulfo-phospho-vanillin method was used

to measure the percentage of total liver lipids (A). qRT-PCR was used to determine relative

gene expression profile of lipolysis and lipogenesis genes in the liver (B) and vWAT (C). Data

are represented as means ± SEM. *p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 based on a two-

tailed Welch's t test (n=8-15). Abbreviations: ACC – acetyl CoA-carboxylase, FAS – fatty acid

synthase, SCD1 - stearoyl-CoA Desaturase, Elovl6 – fatty acid elongase enzyme 6, SREBP1 –

sterol regulatory element-binding protein 1, MGL – monoglyceride lipase, PKA – protein kinase

A, ATGL – adipose triglyceride lipase, HSL – hormone sensitive lipase

*Figure 3 – Blood glucose levels in pre-symptomatic and symptomatic APP$^{NL-F}$ mice.* Mice were

fasted for 15 hours before they received an ip injection of glucose and blood glucose levels

were monitored with a glucometer for two hours to determine glucose tolerance (A). Unfasted

mice received an ip injection of insulin and blood glucose levels were monitored with a

glucometer for 2 hours to determine insulin sensitivity (B). The area under the curve (AUC) for

each assay are inset in pre-symptomatic (left) and symptomatic (right) cohorts. Fasting (C) and

fed (D) blood glucose levels were determined from t=0 during the GTT and ITT, respectively.

Data are represented as means ± SEM. *p<0.05 based on a two-way ANOVA with Sidak's post

hoc (A-D) and a two-tailed Welch's t test (n=9-16).

*Figure 4 – Liver glucose metabolism and plasma adipokines*. qRT-PCR was used to determine

hepatic relative gene expression profiles (A) of gluconeogenesis (G6PC, PEPCK), glycolysis

(GLUT2, GCK), and glycogenesis (GYS2). Circulating visfatin (B), adiponectin (C), and leptin

(D) determined by ELISA from plasma collected at the time of euthanasia. Data are represented

as means ± SEM. *p<0.05 based on a two-tailed Welch's t test (n=4-12). Abbreviations: G6PC -

glucose-6-phosphatase, PEPCK - phosphoenolpyruvate carboxykinase, Glut2 - glucose

transporter 2, GCK - glucokinase, GYS2 - glycogen synthase 2 (GYS2)

*Figure 5 – Senescent cell and inflammation markers in vWAT and SubQ.* qRT-PCR was used

to determine vWAT (A) and SubQ (B) relative gene expression profiles of senescence ($p16^{Ink4\alpha}$,

$p21^{Cip1}$) and inflammation (TNFα, IL-10, IL-6, MCP1). Data are represented as means ± SEM.

*p<0.05, **p<0.01, ***p<0.001, ****p<0.0001 based on a two-tailed Welch's t test (n=4-12).

Abbreviations: TNFα – tumor necrosis factor alpha, IL – interleukin, MCP1 – monocyte

chemoattractant protein 1.

Figure 6 - *Senescent cell and inflammation markers in the hippocampus.* qRT-PCR was used

to determine hippocampal relative gene expression profiles (A) of senescence ($p16^{Ink4\alpha}$, $p21^{Cip1}$),

inflammation (TNFα), and memory (AdipoR, LepR, Creb1, PRGN), while soluble $A\beta_{42}$ (B) was

determined by ELISA. Data are represented as means ± SEM. *p<0.05, **p<0.01 based on a

two-tailed Welch's t test (n=4-12). Abbreviations: TNFα – tumor necrosis factor alpha, AdipoR

– adiponectin receptor, LepR – leptin receptor, Creb1 – cyclic AMP Response Element-Binding

Protein 1, PRGN – progranulin.

Table 1: The number of mice in each cohort as well as the number of mice that experienced post-surgical mortality for that cohort.

| | Pre-Symptomatic APP$^{NL-F}$ | | | | Symptomatic APP$^{NL-F}$ | | | |
|---|---|---|---|---|---|---|---|---|
| Cohort | Male Sham | Male VFR | Female Sham | Female VFR | Male Sham | Male VFR | Female Sham | Female VFR |
| Mice | 10 | 20 | 15 | 17 | 25 | 9 | 14 | 15 |
| Mortality | 0 | 0 | 0 | 2 | 0 | 4 | 2 | 2 |

Table 2: A list of forward and reverse primers.

| Gene Name (Abbreviation) | Forward Primer | Reverse Primer |
|---|---|---|
| Acetyl-CoA Carboxylase (ACC) | 5'-ATGGGCGGAATGGTCTCTTTC-3' | 5'-TGGGGACCTTGTCTTCATCAT-3' |
| Adiponectin Receptor 1 (AdipoR1) | 5'-TCTTCGGGATGTTCTTCCTGG-3' | 5'-TTTGGAAAAAGTCCGAGAGACC-3' |
| Adipose Triglyceride Lipase (ATGL) | 5'-GGATGGCGGCATTTCAGACA-3' | 5'-CAAAGGGTTGGGTTGGTTCAG-3' |
| Beta-2 Microglobulin (BDM) | 5'-AAGTATACTCACGCCACCCA-3' | 5'-AGGACCAGTCCTTGCTGAAG-3' |
| Cyclic AMP Response Element-Binding Protein 1 (CREB1) | 5'-CCCAAAAACGAAGGGAAATCCT-3' | 5'-CCTGGTGCATCAGAAGATAAGTC-3' |
| Fatty Acid Elongase Enzyme 6 (Elovl6) | 5'-GAAAAGCAGTTCAACGAGAACG-3' | 5'-AGATGCCGACCACCAAAGATA-3' |
| Fatty Acid Synthase (FAS) | 5'-GGAGGTGGTGATAGCCGGTAT-3' | 5'-TGGGTAATCCATAGAGCCCAG-3' |
| Glucokinase (GLK) | 5'-TGAGCCGGATGCAGAAGGA-3' | 5'-GCAACATCTTTACACTGGCCT-3' |
| Glucose-6-Phosphatase (G6PC) | 5'-CGACTCGCTATCTCCAAGTGA-3' | 5'-GTTGAACCAGTCTCCGACCA-3' |
| Glucose Transporter 2 (GLUT2) | 5'-TCAGAAGACAAGATCACCGGA-3' | 5'-GCTGGTGTGACTGTAAGTGGG-3' |
| Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH) | 5'-AGGTCGGTGTGAACGGATTTG-3' | 5'-TGTAGACCATGTAGTTGAGGTCA-3' |
| Glycogen Synthase (GYS2) | 5'-ACCAAGGCCAAAACGACAG-3' | 5'-GGGCTCACATTGTTCTACTTGA-3' |
| Hormone Sensitive Lipase (HSL) | 5'-CCAGCCTGAGGGCTTACTG-3' | 5'-CTCCATTGACTGTGACATCTCG-3' |
| Interleukin-6 (IL-6) | 5'-CTGCAAGAGACTTCCATCCAG-3' | 5'-AGTGGTATAGACAGGTCTGTTGG-3' |
| Interleukin-10 (IL-10) | 5'-GCTCTTACTGACTGGCATGAG-3' | 5'-CGCAGCTCTAGGAGCATGTG-3' |
| Leptin Receptor (LEPR) | 5'-TGGTCCCAGCAGCTATGGT-3' | 5'-ACCCAGAGAAGTTAGCACTGT-3' |
| Monocyte Chemoattractant Protein-1 (MCP1) | 5'-CCACTCACCTGCTGCTACTCAT-3' | 5'-TGGTGATCCTCTTGTAGCTCTCC-3' |
| Monoglyceride Lipase (MGL) | 5'-CGGACTTCCAAGTTTTTGTCAGA-3' | 5'-GCAGCCACTAGGATGGAGATG-3' |
| p16$^{Ink4\alpha}$ | 5'-CGGTCGTACCCCGATTCAG-3' | 5'-GCACCGTAGTTGAGCAGAAGA-3' |
| p21$^{Cip1}$ | 5'-CCTGGTGATGTCCGACCTG-3' | 5'-CCATGAGCGCATCGCAATC-3' |
| Phosphoenolpyruvate Carboxykinase (PEPCK) | 5'-CTGCATAACGGTCTGGACTTC-3' | 5'-CAGCAACTGCCCGTACTCC-3' |
| Progranulin (PGRN) | 5'-ATGTGGGTCCTGATGAGCTG-3' | 5'-GCTCGTTATTCTAGGCCATGTG-3' |
| Protein Kinase A (PKA) | 5'-AGATCGTCCTGACCTTTGAGT-3' | 5'-GGCAAAACCGAAGTCTGTCAC-3' |
| Sterol Regulatory Element-Binding Protein 1 (SREBP1) | 5'-GATGTGCGAACTGGACACAG-3' | 5'-CATAGGGGGCGTCAAACAG-3' |
| Stearoyl-CoA Desaturase (SCD1) | 5'-TTCTTGCGATACACTCTGGTGC-3' | 5'-CGGGATTGAATGTTCTTGTCGT-3' |
| Tumor Necrosis Factor α (TNFα) | 5'-CAGGCGGTGCCTATGTCTC-3' | 5'-CGATCACCCCGAAGTTCAGTAG-3' |
| Ubiquitin Conjugating Enzyme E2 D2 (UBE2D2) | 5'-TGCCTGAGATTGCTCGGATCT-3' | 5'-TCGCATACTTCTGAGTCCATTCC-3' |

# Figure 1

Pre-symptomatic Sham    Symptomatic Sham
Pre-symptomatic VFR    Symptomatic VFR

A — Retention Index
B — Novelty Preference (% time)
C — Body Weight (BW) (g)

D — Visceral Fat Mass %

reg.ratio: 55%    92%    50%    100%

male sham    female sham    male sham    female sham
male VFR    female VFR    male VFR    female VFR

Pre-Symptomatic    Symptomatic

E — Total Fat Mass %

Sham    BAT    SubQ    vWAT
VFR

male    female    male    female
Pre-symptomatic    Symptomatic

F

Pre-symptomatic Male Sham

Pre-symptomatic Male VFR

G — Adipocyte Count    Adipocyte Size (µm²)

male    female    male    female

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6