Surgical Removal of Visceral Adipose Tissue has Therapeutic Benefit in Male APP[NL-F] Mice

Samuel A. McFadden[1†], Yimin Fang[1†], Kathleen Quinn[1], Mackenzie R. Peck[1], Jenelle E.

Chapman[1], Tiarra Hill[1], Andrzej Bartke[2,3], Erin R. Hascup[1,4], Kevin N. Hascup[1,3,4*]

[1]Dale and Deborah Smith Center for Alzheimer's Research and Treatment, [2]Internal Medicine,

[3]Medical Microbiology, Immunology and Cell Biology, and [4]Pharmacology, at Southern Illinois

University School of Medicine, Springfield, IL, USA

[†]These authors contributed equally to this work and share first authorship.

*Corresponding Author: Kevin N. Hascup, Dale and Deborah Smith Center for Alzheimer's

Research and Treatment, Southern Illinois University School of Medicine, P.O. Box 19628,

Springfield, IL 62794-9628, USA Tel: 217-545-6994, Email: khascup49@siumed.edu

Keywords: Alzheimer's disease, lipectomy, white adipose tissue, cell senescence, glucose and

lipid metabolism, inflammation, visfatin

Manuscript Length: 5,557 Words

Figures: 7

Tables: 1

Abstract

<u>Purpose:</u> Visceral white adipose tissue (vWAT) accumulation causes systemic inflammation,

insulin resistance, metabolic syndrome, and senescent cell accumulation that are risk factors for

Alzheimer's disease (AD). Visceral fat removal (VFR) improves metabolism and reduces

proinflammatory cytokines. We hypothesized that VFR removal in AD mice would improve

metabolism and cognition.

<u>Methods:</u> Male and female APP^{NL-F} mice underwent sham or vWAT surgical resection

(epididymal and perirenal) at 4 (pre-symptomatic) and 16 (symptomatic) months of age to

understand interventional and therapeutic effects, respectively. At 18 months of age, glucose

metabolism and novel object recognition (NOR) memory were assayed followed by assessment

of body composition and tissue-specific markers of metabolism, cell senescence, inflammation,

or amyloid accumulation.

<u>Results:</u> Male and female APP^{NL-F} mice showed distinct VFR responses. In pre-symptomatic

males, increased vWAT lipolysis and hepatic lipogenesis led to ectopic liver lipid accumulation,

with reduced adiponectin and leptin, elevated visfatin, and impaired glucose metabolism.

Symptomatic males showed reduced vWAT lipogenesis, enhanced hepatic lipolysis, glycolysis,

and glycogenesis, lowering liver lipids and improving insulin sensitivity. Only symptomatic males

improved NOR, linked to elevated hippocampal learning and memory markers. Female vWAT

reaccumulation was due to increased lipogenesis and lower lipolysis. Pre-symptomatic females

had lower hepatic lipogenesis, while glycolysis and glycogenesis declined with disease

progression. Hippocampal senescence and inflammation were elevated early in the disease that

persisted symptomatically.

<u>Conclusions:</u> Sex-specific differences in glucose and lipid metabolism and lipid accumulation

underlie the divergent responses to VFR in APP^{NL-F} mice, with symptomatic males showing the

only beneficial outcomes in metabolism and cognition.

Graphical Abstract

Male and female APP^{NL-F} mice exhibited distinct visceral fat responses to AD progression. In males, early disease stages were marked by vWAT lipolysis and hepatic lipid accumulation with metabolic dysfunction, while symptomatic stages showed a metabolic shift that improved insulin sensitivity and NOR performance. In contrast, females displayed progressive vWAT reaccumulation, reduced hepatic metabolism, and persistent hippocampal senescence and inflammation from early stages onward.

Introduction

Alzheimer's disease (AD) is a progressive, neurodegenerative disorder, and the most prevalent form of dementia. AD is historically characterized by senile plaques composed of aggregated amyloid beta (Aβ) and neurofibrillary tangles formed from hyperphosphorylated tau protein. However, AD is a multifactorial disorder characterized not only by aberrant protein aggregation, but also by mitochondrial dysfunction, oxidative stress, cellular senescence, and chronic inflammation - all of which contribute to disease progression (Breijyeh and Karaman, 2020). The deleterious cascade underlying AD progression is hypothesized to begin in midlife and is not isolated to the CNS. Rather peripheral organs, such as visceral white adipose tissue (vWAT), regulate brain function through hormone secretion as well as glucose and lipid metabolism (Gustafson, 2012, Dove et al., 2021). Accumulation of vWAT in midlife is a predictor for future development of AD (Huang et al., 2022).

Adipose tissue regulates many aspects of whole-body physiology, including food intake, maintenance of energy levels, insulin sensitivity, body temperature, and immune responses (Sakers et al., 2022). Adipose tissue is located under the skin (subcutaneous; SubQ), between and around organs (vWAT), interscapular brown adipose tissue (BAT), in bone marrow, and in mammary glands (Wong Zhang et al., 2023). Beyond total adipose tissue mass, the distribution of fat depots is a stronger predictor of obesity-related disease risk and associated complications (Chusyd et al., 2016). Excess vWAT accumulation causes systemic inflammation, insulin resistance, reduced cerebral blood flow, and metabolic syndrome, which contributes to brain volume shrinkage and cognitive decline (Richard et al., 2000, Wong Zhang et al., 2023, Baglietto-Vargas et al., 2016, Biessels and Despa, 2018, Dove et al., 2021, Nguyen et al., 2014).

The link between systemic inflammation with vWAT and metabolic dysfunction has been reported in diabetic fatty rodents and in aged rats (Borst et al., 2005, Gabriely et al., 2002,

Pitombo et al., 2006). In humans, vWAT is a stronger predictor of age-related cognitive

impairment than body mass index (Whitmer et al., 2008, Elias et al., 2005, Kamogawa et al.,

2010). A cross-sectional study also showed that vWAT dysfunction and inflammatory status in

obesity, is associated with increased cerebral Aβ burden in midlife (Kim et al., 2022).

Transplantation of vWAT from obese mice led to impaired cognition in lean wild-type recipient

mice mediated by NLRP3/IL-1β signaling (Guo et al., 2020). Removal of vWAT prevented

obesity-induced insulin resistance, hyperinsulinemia, and decreased plasma proinflammatory

cytokine levels (Borst et al., 2005, Gabriely et al., 2002, Pitombo et al., 2006), which improved

glucose metabolism (Barzilai et al., 1999, Foster et al., 2011, Franczyk et al., 2021). Aging also

increases vWAT senescent cell burden leading to a chronic pro-inflammatory state (Tchkonia et

al., 2010). Reduction of senescent cells in vWAT improves metabolic and cognitive performance

in normal aging (Fang et al., 2023) and AD mice (Fang et al, 2025).

Considering the contribution of vWAT accumulation on cognitive decline and AD progression,

we hypothesized that surgical resection of this adipose depot would improve the metabolic

profile and reduce cognitive impairments in the APP^{NL-F} AD mouse model. APP^{NL-F} mice contain

the human Aβ isoform of the Swedish (KM670/671NL) and Iberian (I716F) mutations knocked

into the *App* locus leading to upregulation of Aβ$_{42}$ without overexpression of APP (Saito et al.,

2014, Benitez et al., 2021). Plaque accumulation begins at ~6 months of age while cognitive

impairments occur between 12-18 months of age. Our prior research indicates that both sexes

of APP^{NL-F} mice have increased markers of senescent cell (p16^{Ink4α} and p21^{Cip1}) and

senescence-associated secretory phenotype (SASP; IL-6, TNFα, MIP1α) in vWAT by four

months of age (Fang et al., 2025). To test our hypothesis, we surgically resected vWAT

(epididymal and perirenal) from male and female APP^{NL-F} mice at either 4 or 16 months of age.

These time points allowed us to examine the effects of visceral fat removal (VFR) for pre-

symptomatic intervention (4 months) or symptomatic therapy (16 months). All metabolic and

cognitive assays were conducted at 18 months of age. Our results indicate that VFR effects

were disease stage- and sex-dependent with metabolic and cognitive benefits observed in

symptomatic male APP^{NL-F} mice.

Material and Methods

Chemicals

All chemicals were stored according to manufacturer recommendations unless stated otherwise. The Phosphovanilin (PV) reagent for lipid determination was prepared by using six milligrams of vanillin (Fisher Scientific Cat # AC140821000) dissolved in 100 mL of 37°C water and further diluted to 500 mL with 85% phosphoric acid.

Animals

Founder APP^{NL-F} mice on a C57BL/6 background (RRID: IMSR_RBRC06343) were obtained from Riken (Japan). C57BL/6 (RRID:IMSR_JAX:000664) mice were originally obtained from The Jackson Laboratory (Bar Harbor, ME) and used to maintain the genetic background of the breeding colonies. At weaning, pups were randomly assigned to mixed-litter cages and group-housed with mice of the same sex and genotype to control for litter size, maternal care, and other potential litter-specific variables. Protocols for animal use were approved by the Institutional Animal Care and Use Committee at Southern Illinois University School of Medicine (Protocol number: 2025-055), which is accredited by the Association for Assessment and Accreditation of Laboratory Animal Care. All studies were conducted in accordance with the United States Public Health Service's Policy on Humane Care and Use of Laboratory Animals. Mice were group housed on a 12:12 h light-dark cycle, with food (Chow 5001 with 23.4% protein, 5% fat, and 5.8% crude fiber; LabDiet PMI Feeds) and water available ad libitum. Genotypes were confirmed by collecting a 5 mm tail tip for DNA analysis by TransnetYX, Inc (Cordova, TN).

Visceral Fat Removal (VFR) Surgery

Four and sixteen-month-old male and female APP^{NL-F} mice were randomized into two groups – VFR (epidydimal and perirenal) and sham-operated. Mice were isoflurane anesthetized,

abdominal fur shaved, and betadine scrubbed over the incision site. An ~1.5 cm vertical midline

incision along the lower abdomen in males or ventral incisions on either lateral abdominal side

in females were made to access epidydimal adipose depots. The perinephric adipose depots

were removed by flank incisions. Adipose depots were removed by blunt dissection. Mice that

underwent the sham operation had their vWAT mobilized, but not removed. Absorbable sutures

were used to close the body cavity while wound clips were used on the skin. Tylenol (1 mg/mL)

was administered in the drinking water from one day pre- to 3 days post-surgery.

<u>Glucose tolerance test (GTT) and insulin tolerance test (ITT)</u>

Glucose metabolism was assessed via GTT and ITT when mice reached 18 months of age (14

or 2 months post VFR surgery) as previously described (Fang et al., 2023). Fifteen-hour-fasted

mice underwent GTT by intraperitoneal (i.p.) injection with 2.0 g glucose per kg of body weight

(BW). Blood glucose levels were measured at 0, 15, 30, 45, 60, and 120 min with a PRESTO

glucometer (AgaMatrix). For ITT, mice were injected i.p. with 1 IU porcine insulin from sigma

(St. Louis, MO) (Cat# I5523) per kg of BW. Blood glucose levels were measured at 0, 15, 30,

45, 60, and 120 min.

<u>Novel object recognition (NOR) task</u>

The NOR was used to evaluate memory retention based on the premise that mice spend more

time exploring a novel object rather than a familiar object if the animal remembers the familiar

object. Mice were habituated in the open field chamber for 30 min on the first day. Twenty-four

hours after, the mouse was returned to the chamber and presented with two similar objects for 5

min. A 24 hr inter-session-interval was used between introduction and retention phases to

promote long-term memory retrieval. During the retention phase one of the familiar objects was

replaced with a novel object and the mouse was given 5 minutes of exploration. The ANY-maze

video tracking system (Stoelting Co., Wood Dale, IL) was used to record mouse navigation

during the familiarization and retention phases and used to calculate the retention index and

novelty preference.

<u>Body composition and tissue procurement</u>

At 19 months of age, body weight was recorded and mice were anesthetized using isoflurane.

A cardiac puncture was used to collect the plasma. The blood was mixed with EDTA, followed

by centrifugation at 10,000 g for 15 min at 4°C for plasma collection. Immediately following

cardiac puncture, mice were decapitated and the peripheral tissues (liver, pancreas,

perigonadal and subcutaneous (SubQ) WAT, and BAT) were weighed to calculate their

contribution to total body weight. Tissue was immediately flash frozen and stored at -80˚C until

processing as previously described (Fang et al., 2023).

<u>Total Liver Lipid Quantification</u>

A modified micro-scale colorimetric sulfo-phospho-vanillin (SPV) method was used to measure

total liver lipids (Tran et al., 2019). Approximately 50 mg of liver tissue was homogenized in

1000 µL of chloroform/methanol (2:1) followed by addition of 500 µL of saline. The mixture was

vortexed then centrifuged at 10,000 rpm for 5 minutes. The chloroform layer was separated and

evaporated at 90°C for ~20 min, then incubated at 90°C with 200 µL of 98% sulfuric acid for 20

min. After adding 750 µL of vanillin–phosphoric acid reagent, samples were incubated at 37˚C

for 15 min, cooled to 25˚C, and 200 µL was transferred to a 96-well microplate. Absorbance was

measured at 530 nm, and lipid content was normalized to tissue weight.

<u>Assessment of blood chemistry</u>

Per the manufacturer's protocol, adiponectin or leptin was measured with respective ELISA kits

from Crystal Chem (Elk Grove Village, IL; Cat# 80569, 90030). Visfatin was measured by an

ELISA kit from Mybiosource (San Diego, CA; Cat# MBS760980).

RT–PCR

mRNA expression was analyzed by quantitative RT–PCR as previously described (Fang et al.,

2020) using Bio-Rad cDNA synthesis kits (Cat # 1708897 and 1708891) and SYBR green mix

(Cat # 1725121) and the QuantStudio 3 Real-Time PCR (Thermo Fisher Scientific). RNA was

extracted using a RNeasy mini kit or RNeasy Lipid Tissue Mini Kit (Qiagen) following the

manufacturer's instructions. Relative expression was calculated as previously described

(Masternak et al., 2012) and primers were purchased from Integrated DNA Technologies. A list

of forward and reverse primers used in this study is shown in Table 1.

Soluble $A\beta_{42}$ Quantification

The hippocampus from one hemisphere was dissected and stored at -80°C until tissue

processing. Soluble $A\beta_{42}$ concentrations were determined using the Human / Rat β amyloid

ELISA kits (WAKO Chemicals; Cat: 290-62601) according to the manufacturer recommended

protocols.

Statistical analysis

Statistical analyses were conducted to determine differences between age- and sex-matched

sham and VFR groups using unpaired two-tailed Student's t tests with significance defined as p

< 0.05. Data are presented as mean ± SEM. All statistical analyses and graphs were completed

using Prism 10 (GraphPad Inc, La Jolla, CA, USA).

Results

<u>VFR at the symptomatic time point improved object recognition memory in male APP^{NL-F} mice.</u>

To explore whether VFR could improve object recognition memory in APP^{NL-F} mice, surgery was performed at a pre-symptomatic stage (4 months of age, prior to detectable plaque accumulation and cognitive decline) or a symptomatic stage (16 months of age, during advanced pathology and functional decline). Both male and female APP^{NL-F} cohorts were included to assess sex-specific effects of VFR. Object recognition memory was assayed at 18 months of age via the NOR task. Only male APP^{NL-F} mice with VFR surgery at the symptomatic time point had improved recognition memory as indicated by increased retention index (Fig. 1A) and novelty preference (Fig. 1B) compared to the age- and sex-matched sham surgery controls. VFR surgery performed at the pre-symptomatic time point in male, or either time point of female APP^{NL-F}, had no effect on object recognition memory (Fig 1A-B).

<u>VFR resulted in changes to body weight and composition in an age- and sex-dependent manner.</u>

To assess physiological changes caused by VFR, we examined body composition at ~19 months of age upon euthanization. Symptomatic male APP^{NL-F} mice with VFR had decreased BW compared to the sham group (Fig 1C). No differences were observed in the other VFR groups. The body composition analyses indicated that after VFR, vWAT accumulated in all ages and sexes but at different percentages. Female VFR APP^{NL-F} mice had vWAT reaccumulation occurring in 92% and 100% of mice in the pre-symptomatic and symptomatic cohorts, respectively (Fig. 1D). In contrast, ~50% of the male mice with VFR surgery at either age observed vWAT reaccumulation (Fig. 1D). This difference in vWAT was the major contributor to the observed changes of total percent adipose mass when including SubQ,

interscapular BAT, and vWAT (Fig. 1E). In summary, the body weight of symptomatic VFR male

APP[NL-F] mice was reduced mainly due to less vWAT reaccumulation.

<u>VFR during the pre-symptomatic time point in male APP[NL-F] mice caused ectopic liver lipid</u>

<u>deposition via increased de novo lipogenesis.</u>

Although reduced reaccumulation of vWAT was similar between pre-symptomatic and

symptomatic VFR male APP[NL-F] mice, surgery at the pre-symptomatic time point did not affect

BW. Body composition changes without effects on BW are known to happen with aging (St-

Onge and Gallagher, 2010). We sought to identify tissue composition changes that caused the

increased body weight despite reduced adiposity in the pre-symptomatic VFR male mice.

Lipodystrophy leads to the excessive deposition of lipids in other organs, such as the liver,

contributing to insulin resistance (Petersen and Shulman, 2018). To test whether the liver had

more ectopic fat deposition, the percentage of total liver lipids was measured. Pre-symptomatic

VFR male APP[NL-F] mice had a higher percentage of liver lipids compared with sham controls

(Fig 2A). In symptomatic APP[NL-F] VFR male mice, total liver lipids were reduced compared with

sham surgery controls while no differences were observed in females at either disease stage.

Liver gene expression showed that key *de novo* lipogenesis genes including acetyl-coenzyme A

carboxylase (ACC), fatty acid synthase (FAS), stearoyl-CoA desaturase 1 (SCD1), elongation of

long-chain fatty acids family member 6 (Elovl6), sterol regulatory element-binding protein-1

(SREBP1) were increased in pre-symptomatic VFR male APP[NL-F] mice compared with sham

controls (Fig 2B). In pre-symptomatic VFR female APP[NL-F] mice, expression of these same

genes (except for ACC) were decreased. Interestingly, liver gene expression of monoglyceride

lipase (MGL), the rate-limiting enzyme in lipolysis, was upregulated only in symptomatic VFR

male APP[NL-F] mice (Fig. 2B). This is one explanation for the decreased BW and vWAT observed

in this group. These data indicated that pre-symptomatic VFR male APP[NL-F] mice accumulated

more ectopic liver lipids likely due to higher *de novo* lipogenesis, while the contrary was

observed in symptomatic males due to higher lipolysis.

<u>VFR in female APP^NL-F mice led to increased gene expression of de novo lipogenesis and</u>

<u>decreased expression of lipolysis genes in reaccumulated vWAT.</u>

Nearly all the VFR APP^{NL-F} female mice reaccumulated vWAT to levels observed in sham

surgery controls matched for age and sex (Fig. 1D). To understand the reason for this

reaccumulation, we examined gene expression of lipogenesis and lipolysis in vWAT (Fig 2C).

Expression levels of two key *de novo* lipogenesis genes, ACC and FAS, were increased in VFR

APP^{NL-F} female mice at either disease stage. The expression of ACC and FAS were decreased

in VFR male APP^{NL-F} mice at the symptomatic, but not at the pre-symptomatic time point.

Lipolysis gene expression, including protein kinase A (PKA), adipocyte triglyceride lipase

(ATGL), and hormone sensitive lipase (HSL), were decreased at both disease stages in VFR

APP^{NL-F} female mice compared with sham controls. In male mice, expression of PKA, ATGL,

and HSL were increased at the pre-symptomatic, but unchanged, at the symptomatic time point.

The data suggests that the accumulation of vWAT after VFR in female APP^{NL-F} mice at either

stage of disease progression is a result of increased lipogenesis and decreased lipolysis gene

expression. In male APP^{NL-F} mice, the reduced accumulation of vWAT after VFR was disease

stage specific with pre-symptomatic time points increasing expression of lipolysis genes, while

surgery at the symptomatic time point decreased lipogenesis gene expression. Interestingly,

SREBP1 vWAT expression was upregulated only in VFR APP^{NL-F} male mice during the pre-

symptomatic time point. SREBP1 is a transcription factor activated by high blood glucose

(Bertolio et al., 2019, Zhao et al., 2024), suggesting altered glucose metabolism in this cohort.

<u>VFR surgery impacted either glucose tolerance or insulin sensitivity in male APP^{NL-F} mice.</u>

Lipid metabolism plays an important role in regulating glucose utilization, and the two processes

are tightly interconnected (Parhofer, 2015). Accordingly, the observed changes in lipid

metabolizing gene expression could affect glucose metabolism. To test the effects of VFR on

blood glucose metabolism, we monitored fasting and fed glucose levels as well as glucose

tolerance with a GTT, and insulin sensitivity with an ITT, at 18 months of age. Pre-symptomatic

VFR APP^{NL-F} male mice were less glucose tolerant, but no effects were observed in

symptomatic males (Fig 3A). Insulin sensitivity was improved in symptomatic VFR APP^{NL-F} male

mice, but differences were not observed in pre-symptomatic males (Fig 3B). Consistent with the

GTT, fasting glucose was higher in pre-symptomatic VFR APP^{NL-F} male mice compared with

sham controls (Fig. 3C). No fasting glucose changes were observed in symptomatic APP^{NL-F}

male mice, nor were differences in fed glucose observed (Fig. 3D) in either male cohort. No

blood glucose differences were observed at either disease stage in VFR APP^{NL-F} female mice.

The liver plays a major role in the control of whole body glucose homeostasis (Han et al., 2016,

Guerra and Gastaldelli, 2020, Scoditti et al., 2024). To understand the observed blood glucose

differences, we examined liver gene expression (Fig 4A) associated with gluconeogenesis

(synthesis of glucose from non-carbohydrate sources), glycolysis (breakdown of glucose for

energy), and glycogenesis (conversion of glucose to glycogen). Expression of gluconeogenesis

genes, glucose-6-phosphatase (G6PC) and phosphoenolpyruvate carboxykinase (PEPCK),

were increased in pre-symptomatic VFR APP^{NL-F} male mice, which was consistent with their

increased fasting blood glucose levels (Fig 3C). In contrast, glucose transporter 2 (Glut2), GCK

(Glucokinase), and glycogen synthase 2 (GYS2), key genes in glycolysis and glycogenesis,

respectively, were increased in male, but decreased in female APP^{NL-F} mice after symptomatic

VFR.

Taken together, the blood glucose and liver gene expression measures indicate VFR affected

glucose metabolism in a sex- and disease stage-dependent manner. Pre-symptomatic VFR in

APP[NL-F] male mice increased their fasting blood glucose and made them less glucose tolerant,

likely mediated via increasing gluconeogenesis. At symptomatic time points, VFR improved

glucose metabolism by increasing insulin sensitivity via enhancing glycolytic and glycogenesis

gene expression. Although this gene expression was reduced in symptomatic VFR female

mice, no effects on insulin sensitivity or blood glucose levels were observed.

<u>VFR alters plasma visfatin, adiponectin, and leptin in pre-symptomatic male APP[NL-F] mice.</u>

Adipokines secreted from adipose tissue are important for whole-body glucose metabolism and

insulin sensitivity (Rabe et al., 2008, Kojta et al., 2020). Given the altered lipid and glucose

metabolism after VFR in the APP[NL-F] mice, we examined plasma adipokines. Visfatin is

primarily produced in vWAT, with plasma levels increasing in response to high-fat diet feeding

and obesity (Adeghate, 2008, Sethi and Vidal-Puig, 2005), and is associated with glucose

intolerance (Abdalla, 2022). Plasma visfatin was increased in pre-symptomatic VFR male

APP[NL-F] mice compared with age-matched sham controls (Fig 4B), consistent with their higher

ectopic liver accumulation, fasting blood glucose, and glucose intolerance. No plasma visfatin

differences were observed in the other mouse cohorts.

Adiponectin enhances glucose and lipid metabolism (Yanai and Yoshida, 2019), decreases

lipogenesis (Richard et al., 2000), and has neuroprotective effects (Khoramipour et al., 2021).

Plasma adiponectin was reduced in pre-symptomatic VFR APP[NL-F] male mice (Fig 4C). This

would also contribute to the reduced glucose metabolism and increased liver lipid observed in

this cohort. Plasma adiponectin changes were not observed after VFR in the other cohorts.

Leptin regulates appetite and energy expenditure, and its deficiency can disrupt glucose and

lipid metabolism, leading to neuronal damage that impairs neuroplasticity and memory

consolidation (Fernandes et al., 2024). Higher leptin levels are also associated with a lower

dementia risk (Lieb et al., 2009). Plasma leptin was decreased in pre-symptomatic VFR APP[NL-F]

male mice compared with age-matched sham controls (Fig. 4D), which contributed to the

imbalance of glucose and lipid metabolism in this male cohort. Differences in plasma leptin

levels were not observed in the other cohorts of mice.

<u>VFR increases WAT senescent cell and SASP markers in pre-symptomatic male and</u>

<u>symptomatic female APP^{NL-F} mice.</u>

Adipose tissue, particularly vWAT, has the highest burden of senescent cells (Palmer et al.,

2021) and we previously showed senescent cell markers in vWAT were increased in male and

female APP^{NL-F} mice (Fang et al., 2025). VFR can decrease systemic and neuroinflammation in

aged mice (Shin et al., 2015). We examined gene expression of senescent cell and SASP

markers in vWAT (Fig 5A). In pre-symptomatic VFR APP^{NL-F} male mice, gene expression of

p16$^{Ink4\alpha}$, p21^{Cip1}, and IL-10 were increased in vWAT. Expression of vWAT p16$^{Ink4\alpha}$ and TNFα

were also increased in symptomatic VFR APP^{NL-F} female mice (Fig 5A) while no differences in

IL-6 and MCP1 were observed in any cohort (data not shown). SASP markers were also

examined in SubQ WAT where TNFα, IL-10, IL-6, and MCP1 were all increased in pre-

symptomatic VFR APP^{NL-F} male mice (Fig 5B). SubQ IL-6 was also increased in symptomatic

VFR APP^{NL-F} female mice. Overall, VFR increased gene expression of senescent cell and

SASP markers in WAT of pre-symptomatic APP^{NL-F} male mice, with a modest increase

observed in symptomatic females.

<u>VFR modulates gene expression of hippocampal senescence, SASP, and cognitive markers.</u>

Excess vWAT accumulation is linked to accelerated brain aging, neuroinflammation, and

dementia risk (Moser and Pike, 2016, Oh et al., 2023, Budamagunta et al., 2023). To

understand how VFR affected hippocampal function, the brain region strongly associated with

learning and memory, we examined gene expression of cell senescence and cytokines (Fig 6A).

VFR had no effect on p16$^{Ink4\alpha}$, p21^{Cip1}, or TNFα expression in APP^{NL-F} male mice at either stage

of disease progression. Alternatively, $p16^{Ink4\alpha}$ gene expression was increased in symptomatic

VFR APP^{NL-F} female mice. There was a corresponding increase in TNFα expression in this

cohort as well as the symptomatic females. VFR had deleterious effects on hippocampal

senescence and inflammation in female, but not male APP^{NL-F} mice.

Since we observed improvements in object recognition memory after VFR in symptomatic male

APP^{NL-F} mice, we examined markers associated with learning and memory in the hippocampus

(Fig 6A). Both adiponectin and leptin receptors (AdipoR, LepR) are found in the hippocampus

and play a role in synaptic plasticity and cognition. Gene expression of both receptors were

upregulated in symptomatic VFR APP^{NL-F} male mice (Fig 6A). In females, VFR increased LepR

in the pre-symptomatic disease stage. Cyclic AMP response element-binding protein 1 (Creb1),

plays a role in long-term memory formation. Creb1 gene expression was decreased in pre-

symptomatic VFR APP^{NL-F} male mice, but elevated in the symptomatic male cohort. Progranulin

(PRGN) is important for neuronal survival and growth in the hippocampus. PRGN gene

expression was decreased in pre-symptomatic VFR APP^{NL-F} female mice, but increased in

symptomatic males. Overall, VFR altered the hippocampal gene expression profile of

symptomatic APP^{NL-F} male mice in a manner that would improve their object recognition

memory.

<u>Soluble Aβ$_{42}$ is increased in pre-symptomatic male and female mice.</u>

Soluble Aβ$_{42}$ is the neurotoxic species associated with synaptic dysfunction and cognitive

decline in AD (Findley et al., 2019). An ELISA indicated hippocampal soluble Aβ$_{42}$ (Fig 6B)

production was increased in VFR APP^{NL-F} male and female mice, but only during the pre-

symptomatic disease stage. VFR during the symptomatic stage had no effect on soluble Aβ$_{42}$.

Discussion

Adipose accumulation and distribution differ between sexes and are evolutionarily conserved. Females typically have a higher overall fat percentage with a large proportion stored subcutaneously for energy supply during pregnancy and lactation during food scarcity, while males accumulate fat in visceral regions. Numerous variables influence adipose development including sex chromosomes, gonadal hormone levels, and hormone receptors. Sex chromosome complement impacts adipose accumulation whereby the presence of two "X" chromosomes leads to higher accumulation while the presence of the "Y" chromosome has no influence. The gonadal hormones, testosterone and estrogen, drive visceral and subcutaneous adipose accumulation in males and females, respectively. Age-related hormonal changes increase vWAT accumulation regardless of sex and is a key factor in the development of metabolic syndrome that increases dementia risk.

Besides its role as an energy reservoir, adipose tissue is an endocrine organ that secretes peptides such as TNFα, IL-6, and adiponectin, which can influence either tissue-specific or whole body metabolism. Prior studies examining VFR effects in rodents were conducted to elucidate mechanisms contributing to metabolic improvements. VFR improves insulin sensitivity and glucose tolerance in young, aged, and obese rodents with some of the metabolic changes occurring through decreased free fatty acids, leptin, and TNFα expression (Kim et al., 1999, Gabriely et al., 2002, Shi et al., 2007a). VFR in aged C57BL/6 mice led to reduced expression of proinflammatory cytokines in the plasma, WAT, and cortex (Shin et al., 2015). Although adiponectin plays a key role in glucose and lipid metabolism and suppresses inflammatory cytokines, serum levels are found to decrease following VFR in male and female Sprague Dawley rats (Borst et al., 2005, Ling et al., 2014), which may be due to the absence of these WAT depots after VFR. Considering the metabolic benefits of VFR and the known link between metabolic dysfunction and AD risk, we aimed to investigate the impact of VFR in an AD model.

Prior research has shown APP^{NL-F} mice, particularly females, have worse glucose tolerance

(Abi-Ghanem et al., 2025), potentially due to increased senescent cell and SASP markers in

vWAT (Fang et al., 2025). Senescent adipocytes contribute to chronic systemic inflammation

and metabolic dysfunction, which diminishes cognitive reserve, promotes the accumulation of

misfolded proteins, and accelerates AD progression (Huang et al., 2022). We hypothesized that

surgical resection of vWAT would improve metabolic function and recognition memory in APP^{NL-}

^{F} mice. We performed VFR at two different stages of disease progression to explore

interventional and therapeutic strategies.

Surprisingly, the VFR response was disease stage- and sex-dependent. The removal of vWAT

in symptomatic APP^{NL-F} male mice improved object recognition memory potentially due to

increased hippocampal AdipoR, LepR, Creb1, and PRGN expression. Lipogenesis vWAT gene

expression was down-regulated, while hepatic genes associated with lipolysis, glycolysis, and

glycogenesis were up-regulated. These effects were associated with a decreased percentage

of total liver lipids. All of these alterations were reflected by a reduced percentage of total fat,

specifically reaccumulated vWAT, BW, and improved blood glucose maintenance. The opposite

was true for VFR during the pre-symptomatic stage in males. After VFR, an imbalance of lipid

metabolism was prevalent, reflected by an elevated expression of vWAT lipolysis and hepatic

lipogenesis genes. This tissue metabolic imbalance accounts for the redistribution of lipids into

other organs, such as the liver, where higher ectopic fat accumulation was observed. Plasma

adiponectin and leptin were decreased along with increased visfatin resulting in impaired

glucose metabolism and peripheral inflammation. Reduced hippocampal Creb1 expression and

increased soluble Aβ_{42}, combined with the aforementioned adverse physiological changes,

failed to improve object recognition memory.

In contrast to males, vWAT reaccumulated in nearly all female mice regardless of when VFR

occurred, attributed to higher lipogenesis and lower lipolysis gene expression. Although ectopic

lipid deposition did not accumulate in the liver, VFR in pre-symptomatic female APP^{NL-F} mice led

to decreased hepatic lipogenesis gene expression, while expression of genes involved in

glycolysis and glycogenesis were decreased in symptomatic females. These results were in

opposition to age-matched males. Hippocampal soluble Aβ_{42}, senescent cell, and inflammatory

markers were elevated in pre-symptomatic APP^{NL-F} female mice while the later was only

observed at the symptomatic time point. No changes to blood glucose levels or object

recognition memory were observed in females after VFR.

Lipectomy in mice, rats, and other small mammals does not result in long-term BW loss with

regain occurring 12-14 weeks post-surgery (Murillo et al., 2019). In the majority of the studies

this was attributed to regeneration of the lipectomized depot, redistribution of fat to other

adipose depots, or ectopic fat accumulation in other organs. In our present study, female APP^{NL-}

^{F} mice regained BW due to accumulating vWAT. Although vWAT accumulation was lower in

pre-symptomatic males, they regained BW due to ectopic fat distribution in the liver. Only

symptomatic APP^{NL-F} male mice had lower BW attributed to less visceral adiposity.

Redistribution of fat to other adipose depots was not observed in any cohort. Our present study

was designed to examine disease stage-specific interventional or therapeutic AD strategies

rather than examining the contribution of aging on adiposity and AD progression. Regardless,

effects of aging on adipose accumulation likely contributed to the observed differences in lipid

distribution and the comparable end-of-study BW between sham and VFR cohorts matched for

age and sex. Estropause and the gradual decrease in estrogen levels that favors vWAT rather

than SubQ accumulation may partially account for visceral accumulation after resection in

female mice, while in males, ectopic liver lipid distribution has been observed with aging

(Donato et al., 2014).

The exact mechanisms behind this adipose rebound are not well known, but suggest biological

feedback systems that act to preserve energy balance. For example, the adipostat hypothesis

postulates that alterations to energy reserves triggers compensatory mechanisms in metabolism

and/or thermogenesis to restore the reservoir. Since gonadal hormones and sex chromosome

complement mediate adipose storage, distribution, and utilization, male and female mice use

different behavioral strategies to restore fat after lipectomy. Males increase caloric consumption

while females reduce heat production and energy expenditure (Shi et al., 2007b). Our findings

are in support of this whereby pre-symptomatic VFR APP^{NL-F} male mice had increased

gluconeogenesis gene expression and fasting blood glucose, and worse glucose tolerance.

Female VFR APP^{NL-F} mice had reduced glycolysis and lipolysis, but increased lipogenesis gene

expression. It is important to note, however, that our study was not intended to serve as a

preclinical model for liposuction, as that procedure primarily targets SubQ rather than vWAT

depots. However, lipectomy in humans causes a transient decrease in body fat and weight that

gradually return to preoperative levels within months (Hernandez et al., 2011, Seretis et al.,

2015), suggesting the preservation of this energy balance is conserved across species.

The accumulation of WAT and subsequent increases in BW begins in midlife and continues into

early aging in both humans (40-65 years) and mice (12-18 months). The reason for this

increase is attributed to adipogenesis and adipocyte hypertrophy that are prominent during

midlife (Wang et al., 2025). Remodeling of vWAT, like its distribution, varies between sexes,

with estrogen levels playing a key role in regulating these processes and shifting notably with

age. Female mice have increased adipose progenitor cells leading to hyperplasia while males

experience hypertrophy (Wu et al., 2017). Both increase adipose accumulation, but hyperplasia

is associated with better metabolic health (Steiner and Berry, 2022), which may explain why

blood glucose levels were unaffected in female mice after VFR. Hypertrophic adipocytes

secrete less adiponectin and release more proinflammatory cytokines with the latter causing

ectopic lipid distribution into other organs, which were all prevalent in pre-symptomatic VFR

APP[NL-F] male mice. Accordingly, adipose cell size rather than number correlates with increased

risk for metabolic syndrome (Vishvanath and Gupta, 2019).

Senescent cell accumulation in both the periphery and brain contribute to aging and age-related

neurodegenerative disorders such as AD. The mechanisms behind this are multifaceted and

dependent upon the type of senescent cell (Zhu et al., 2024). Senescent brain cells alter

amyloid precursor protein (APP) processing by shifting it from the non-amyloidogenic to the

amyloidogenic pathway, promoting the formation of Aβ (Sun et al., 2018). The proinflammatory

SASP response coupled with decreased microglial phagocytosis of misfolded proteins can lead

to the subsequent Aβ accumulation and deposition in the brain. We observed elevated

hippocampal soluble $Aβ_{42}$ in pre-symptomatic VFR APP[NL-F] mice compared with sex-matched

sham controls. Senescent cell and SASP accumulation were prevalent in the WAT of pre-

symptomatic males, while these were increased in the hippocampus of pre-symptomatic

females. In AD mouse models, reducing peripheral and hippocampal senescent cell burden with

senolytics has been shown to reduce amyloid pathology and improve cognitive performance -

albeit with benefits favoring female mice (Currais et al., 2014, Zhang et al., 2019, Fang et al.,

2025, Ng et al., 2024).

Given that AD pathology is thought to initiate in midlife prior to cognitive decline, the concurrent

increase in adipose tissue accumulation during this period may play a causal role in disease

progression. Our present study indicates accumulation or redistribution of adipose tissue during

the pre-symptomatic AD stage can cause metabolic disturbances resulting in sex-specific

peripheral and cerebral senescent cell accumulation that exacerbates pathological AD

progression. However, during symptomatic AD stages, elimination of vWAT has beneficial

metabolic and cognitive effects in male, but not female APP[NL-F] mice. These sex-specific

differences are attributed to increasing energy storage in female symptomatic mice.

Disclosures

The authors state that they have no financial or non-financial competing interests to disclose.

SAM, YF, MRP, KQ, TH, and JEC conducted the experiments and analyzed the data. YF, AB,

KNH, and ERH conceived the study, designed the experiments, interpreted the data, and wrote

the manuscript. All authors approved the final version of the manuscript.

Funding Sources

This work was supported by the National Institutes of Health NIA R01-AG057767 and NIA R01-

AG061937, Kenneth Stark Endowment, and Dale and Deborah Smith Center for Alzheimer's

Research and Treatment (YF, MRP, KQ, JEC, SM, TH, KNH, ERH). Geriatrics Initiative (AB).

References

ABDALLA, M. M. I. 2022. Role of visfatin in obesity-induced insulin resistance. *World J Clin Cases,* 10**,**
10840-10851.
ABI-GHANEM, C., KELLY, R. D., GROOM, E. A., VALERIAN, C. G., PAUL, A. S., THRASHER, C. A., SALINERO,
A. E., BATCHELDER, M. R., LAFRICAN, J. J., WANG, M., SMITH, R. M., TEMPLE, S., ZULOAGA, D. G.
& ZULOAGA, K. L. 2025. Interactions between menopause and high-fat diet on cognition and
pathology in a mouse model of Alzheimer's disease. *Alzheimers Dement,* 21**,** e70026.
ADEGHATE, E. 2008. Visfatin: structure, function and relation to diabetes mellitus and other
dysfunctions. *Curr Med Chem,* 15**,** 1851-62.
BAGLIETTO-VARGAS, D., SHI, J., YAEGER, D. M., AGER, R. & LAFERLA, F. M. 2016. Diabetes and
Alzheimer's disease crosstalk. *Neurosci Biobehav Rev,* 64**,** 272-87.
BARZILAI, N., SHE, L., LIU, L., WANG, J., HU, M., VUGUIN, P. & ROSSETTI, L. 1999. Decreased visceral
adiposity accounts for leptin effect on hepatic but not peripheral insulin action. *Am J Physiol,*
277**,** E291-8.
BENITEZ, D. P., JIANG, S., WOOD, J., WANG, R., HALL, C. M., PEERBOOM, C., WONG, N., STRINGER, K. M.,
VITANOVA, K. S., SMITH, V. C., JOSHI, D., SAITO, T., SAIDO, T. C., HARDY, J., HANRIEDER, J., DE
STROOPER, B., SALIH, D. A., TRIPATHI, T., EDWARDS, F. A. & CUMMINGS, D. M. 2021. Knock-in
models related to Alzheimer's disease: synaptic transmission, plaques and the role of microglia.
Mol Neurodegener, 16**,** 47.
BERTOLIO, R., NAPOLETANO, F., MANO, M., MAURER-STROH, S., FANTUZ, M., ZANNINI, A., BICCIATO, S.,
SORRENTINO, G. & DEL SAL, G. 2019. Sterol regulatory element binding protein 1 couples
mechanical cues and lipid metabolism. *Nat Commun,* 10**,** 1326.
BIESSELS, G. J. & DESPA, F. 2018. Cognitive decline and dementia in diabetes mellitus: mechanisms and
clinical implications. *Nat Rev Endocrinol,* 14**,** 591-604.
BORST, S. E., CONOVER, C. F. & BAGBY, G. J. 2005. Association of resistin with visceral fat and muscle
insulin resistance. *Cytokine,* 32**,** 39-44.
BREIJYEH, Z. & KARAMAN, R. 2020. Comprehensive Review on Alzheimer's Disease: Causes and
Treatment. *Molecules,* 25.
BUDAMAGUNTA, V., KUMAR, A., RANI, A., BEAN, L., MANOHAR-SINDHU, S., YANG, Y., ZHOU, D. &
FOSTER, T. C. 2023. Effect of peripheral cellular senescence on brain aging and cognitive decline.
Aging Cell, 22**,** e13817.
CHUSYD, D. E., WANG, D., HUFFMAN, D. M. & NAGY, T. R. 2016. Relationships between Rodent White
Adipose Fat Pads and Human White Adipose Fat Depots. *Front Nutr,* 3**,** 10.
CURRAIS, A., PRIOR, M., DARGUSCH, R., ARMANDO, A., EHREN, J., SCHUBERT, D., QUEHENBERGER, O. &
MAHER, P. 2014. Modulation of p25 and inflammatory pathways by fisetin maintains cognitive
function in Alzheimer's disease transgenic mice. *Aging Cell,* 13**,** 379-90.
DONATO, A. J., HENSON, G. D., HART, C. R., LAYEC, G., TRINITY, J. D., BRAMWELL, R. C., ENZ, R. A.,
MORGAN, R. G., REIHL, K. D., HAZRA, S., WALKER, A. E., RICHARDSON, R. S. & LESNIEWSKI, L. A.
2014. The impact of ageing on adipose structure, function and vasculature in the B6D2F1
mouse: evidence of significant multisystem dysfunction. *J Physiol,* 592**,** 4083-96.
DOVE, A., SHANG, Y., XU, W., GRANDE, G., LAUKKA, E. J., FRATIGLIONI, L. & MARSEGLIA, A. 2021. The
impact of diabetes on cognitive impairment and its progression to dementia. *Alzheimers*
Dement, 17**,** 1769-1778.
ELIAS, M. F., ELIAS, P. K., SULLIVAN, L. M., WOLF, P. A. & D'AGOSTINO, R. B. 2005. Obesity, diabetes and
cognitive deficit: The Framingham Heart Study. *Neurobiol Aging,* 26 Suppl 1**,** 11-6.

FANG, Y., MCFADDEN, S., DARCY, J., HASCUP, E. R., HASCUP, K. N. & BARTKE, A. 2020. Lifespan of long-
lived growth hormone receptor knockout mice was not normalized by housing at 30 degrees C
since weaning. *Aging Cell,* 19**,** e13123.
FANG, Y., MEDINA, D., STOCKWELL, R., MCFADDEN, S., QUINN, K., PECK, M. R., BARTKE, A., HASCUP, K.
N. & HASCUP, E. R. 2023. Sexual dimorphic metabolic and cognitive responses of C57BL/6 mice
to Fisetin or Dasatinib and quercetin cocktail oral treatment. *Geroscience,* 45**,** 2835-2850.
FANG, Y., PECK, M. R., QUINN, K., CHAPMAN, J. E., MEDINA, D., MCFADDEN, S. A., BARTKE, A., HASCUP,
E. R. & HASCUP, K. N. 2025. Senolytic intervention improves cognition, metabolism, and
adiposity in female APP(NL)(-F/NL-F) mice. *Geroscience,* 47**,** 1123-1138.
FERNANDES, C., FORNY-GERMANO, L., ANDRADE, M. M., LYRA, E. S. N. M., RAMOS-LOBO, A. M.,
MEIRELES, F., TOVAR-MOLL, F., HOUZEL, J. C., DONATO, J., JR. & DE FELICE, F. G. 2024. Leptin
receptor reactivation restores brain function in early-life Lepr-deficient mice. *Brain,* 147**,** 2706-
2717.
FINDLEY, C. A., BARTKE, A., HASCUP, K. N. & HASCUP, E. R. 2019. Amyloid Beta-Related Alterations to
Glutamate Signaling Dynamics During Alzheimer's Disease Progression. *ASN Neuro,* 11**,**
1759091419855541.
FOSTER, M. T., SHI, H., SEELEY, R. J. & WOODS, S. C. 2011. Removal of intra-abdominal visceral adipose
tissue improves glucose tolerance in rats: role of hepatic triglyceride storage. *Physiol Behav,*
104**,** 845-54.
FRANCZYK, M. P., HE, M. & YOSHINO, J. 2021. Removal of Epididymal Visceral Adipose Tissue Prevents
Obesity-Induced Multi-organ Insulin Resistance in Male Mice. *J Endocr Soc,* 5**,** bvab024.
GABRIELY, I., MA, X. H., YANG, X. M., ATZMON, G., RAJALA, M. W., BERG, A. H., SCHERER, P., ROSSETTI, L.
& BARZILAI, N. 2002. Removal of visceral fat prevents insulin resistance and glucose intolerance
of aging: an adipokine-mediated process? *Diabetes,* 51**,** 2951-8.
GUERRA, S. & GASTALDELLI, A. 2020. The role of the liver in the modulation of glucose and insulin in non
alcoholic fatty liver disease and type 2 diabetes. *Curr Opin Pharmacol,* 55**,** 165-174.
GUO, D. H., YAMAMOTO, M., HERNANDEZ, C. M., KHODADADI, H., BABAN, B. & STRANAHAN, A. M.
2020. Visceral adipose NLRP3 impairs cognition in obesity via IL-1R1 on CX3CR1+ cells. *J Clin
Invest,* 130**,** 1961-1976.
GUSTAFSON, D. R. 2012. Adiposity and cognitive decline: underlying mechanisms. *J Alzheimers Dis,* 30
Suppl 2**,** S97-112.
HAN, H. S., KANG, G., KIM, J. S., CHOI, B. H. & KOO, S. H. 2016. Regulation of glucose metabolism from a
liver-centric perspective. *Exp Mol Med,* 48**,** e218.
HERNANDEZ, T. L., KITTELSON, J. M., LAW, C. K., KETCH, L. L., STOB, N. R., LINDSTROM, R. C.,
SCHERZINGER, A., STAMM, E. R. & ECKEL, R. H. 2011. Fat redistribution following suction
lipectomy: defense of body fat and patterns of restoration. *Obesity (Silver Spring),* 19**,** 1388-95.
HUANG, X., WANG, Y. J. & XIANG, Y. 2022. Bidirectional communication between brain and visceral
white adipose tissue: Its potential impact on Alzheimer's disease. *EBioMedicine,* 84**,** 104263.
KAMOGAWA, K., KOHARA, K., TABARA, Y., UETANI, E., NAGAI, T., YAMAMOTO, M., IGASE, M. & MIKI, T.
2010. Abdominal fat, adipose-derived hormones and mild cognitive impairment: the J-SHIPP
study. *Dement Geriatr Cogn Disord,* 30**,** 432-9.
KHORAMIPOUR, K., CHAMARI, K., HEKMATIKAR, A. A., ZIYAIYAN, A., TAHERKHANI, S., ELGUINDY, N. M. &
BRAGAZZI, N. L. 2021. Adiponectin: Structure, Physiological Functions, Role in Diseases, and
Effects of Nutrition. *Nutrients,* 13.
KIM, S., YI, H. A., WON, K. S., LEE, J. S. & KIM, H. W. 2022. Association between Visceral Adipose Tissue
Metabolism and Alzheimer's Disease Pathology. *Metabolites,* 12.

583 KIM, Y. W., KIM, J. Y. & LEE, S. K. 1999. Surgical removal of visceral fat decreases plasma free fatty acid
584   and increases insulin sensitivity on liver and peripheral tissue in monosodium glutamate (MSG)-
585   obese rats. *J Korean Med Sci,* 14**,** 539-45.
586 KOJTA, I., CHACINSKA, M. & BLACHNIO-ZABIELSKA, A. 2020. Obesity, Bioactive Lipids, and Adipose Tissue
587   Inflammation in Insulin Resistance. *Nutrients,* 12.
588 LIEB, W., BEISER, A. S., VASAN, R. S., TAN, Z. S., AU, R., HARRIS, T. B., ROUBENOFF, R., AUERBACH, S.,
589   DECARLI, C., WOLF, P. A. & SESHADRI, S. 2009. Association of plasma leptin levels with incident
590   Alzheimer disease and MRI measures of brain aging. *JAMA,* 302**,** 2565-72.
591 LING, B. L., CHIU, C. T., LU, H. C., LIN, J. J., KUO, C. Y. & CHOU, F. P. 2014. Short and long-term impact of
592   lipectomy on expression profile of hepatic anabolic genes in rats: a high fat and high cholesterol
593   diet-induced obese model. *PLoS One,* 9**,** e108717.
594 MASTERNAK, M. M., BARTKE, A., WANG, F., SPONG, A., GESING, A., FANG, Y., SALMON, A. B., HUGHES, L.
595   F., LIBERATI, T., BOPARAI, R., KOPCHICK, J. J. & WESTBROOK, R. 2012. Metabolic effects of intra-
596   abdominal fat in GHRKO mice. *Aging Cell,* 11**,** 73-81.
597 MOSER, V. A. & PIKE, C. J. 2016. Obesity and sex interact in the regulation of Alzheimer's disease.
598   *Neurosci Biobehav Rev,* 67**,** 102-18.
599 MURILLO, A. L., KAISER, K. A., SMITH, D. L., JR., PETERSON, C. M., AFFUSO, O., TIWARI, H. K. & ALLISON,
600   D. B. 2019. A Systematic Scoping Review of Surgically Manipulated Adipose Tissue and the
601   Regulation of Energetics and Body Fat in Animals. *Obesity (Silver Spring),* 27**,** 1404-1417.
602 NG, P. Y., ZHANG, C., LI, H. & BAKER, D. J. 2024. Senescence Targeting Methods Impact Alzheimer's
603   Disease Features in 3xTg Mice. *J Alzheimers Dis,* 97**,** 1751-1763.
604 NGUYEN, J. C., KILLCROSS, A. S. & JENKINS, T. A. 2014. Obesity and cognitive decline: role of
605   inflammation and vascular changes. *Front Neurosci,* 8**,** 375.
606 OH, H. S., RUTLEDGE, J., NACHUN, D., PALOVICS, R., ABIOSE, O., MORAN-LOSADA, P., CHANNAPPA, D.,
607   UREY, D. Y., KIM, K., SUNG, Y. J., WANG, L., TIMSINA, J., WESTERN, D., LIU, M., KOHLFELD, P.,
608   BUDDE, J., WILSON, E. N., GUEN, Y., MAURER, T. M., HANEY, M., YANG, A. C., HE, Z., GREICIUS,
609   M. D., ANDREASSON, K. I., SATHYAN, S., WEISS, E. F., MILMAN, S., BARZILAI, N., CRUCHAGA, C.,
610   WAGNER, A. D., MORMINO, E., LEHALLIER, B., HENDERSON, V. W., LONGO, F. M.,
611   MONTGOMERY, S. B. & WYSS-CORAY, T. 2023. Organ aging signatures in the plasma proteome
612   track health and disease. *Nature,* 624**,** 164-172.
613 PALMER, A. K., TCHKONIA, T. & KIRKLAND, J. L. 2021. Senolytics: Potential for Alleviating Diabetes and Its
614   Complications. *Endocrinology,* 162.
615 PARHOFER, K. G. 2015. Interaction between Glucose and Lipid Metabolism: More than Diabetic
616   Dyslipidemia. *Diabetes Metab J,* 39**,** 353-62.
617 PETERSEN, M. C. & SHULMAN, G. I. 2018. Mechanisms of Insulin Action and Insulin Resistance. *Physiol*
618   *Rev,* 98**,** 2133-2223.
619 PITOMBO, C., ARAUJO, E. P., DE SOUZA, C. T., PAREJA, J. C., GELONEZE, B. & VELLOSO, L. A. 2006.
620   Amelioration of diet-induced diabetes mellitus by removal of visceral fat. *J Endocrinol,* 191**,** 699-
621   706.
622 RABE, K., LEHRKE, M., PARHOFER, K. G. & BROEDL, U. C. 2008. Adipokines and insulin resistance. *Mol*
623   *Med,* 14**,** 741-51.
624 RICHARD, A. J., WHITE, U., ELKS, C. M. & STEPHENS, J. M. 2000. Adipose Tissue: Physiology to Metabolic
625   Dysfunction. *In:* FEINGOLD, K. R., AHMED, S. F., ANAWALT, B., BLACKMAN, M. R., BOYCE, A.,
626   CHROUSOS, G., CORPAS, E., DE HERDER, W. W., DHATARIYA, K., DUNGAN, K., HOFLAND, J.,
627   KALRA, S., KALTSAS, G., KAPOOR, N., KOCH, C., KOPP, P., KORBONITS, M., KOVACS, C. S.,
628   KUOHUNG, W., LAFERRERE, B., LEVY, M., MCGEE, E. A., MCLACHLAN, R., MUZUMDAR, R.,
629   PURNELL, J., REY, R., SAHAY, R., SHAH, A. S., SINGER, F., SPERLING, M. A., STRATAKIS, C. A.,
630   TRENCE, D. L. & WILSON, D. P. (eds.) *Endotext.* South Dartmouth (MA).

SAITO, T., MATSUBA, Y., MIHIRA, N., TAKANO, J., NILSSON, P., ITOHARA, S., IWATA, N. & SAIDO, T. C.
2014. Single App knock-in mouse models of Alzheimer's disease. *Nat Neurosci,* 17**,** 661-3.
SAKERS, A., DE SIQUEIRA, M. K., SEALE, P. & VILLANUEVA, C. J. 2022. Adipose-tissue plasticity in health
and disease. *Cell,* 185**,** 419-446.
SCODITTI, E., SABATINI, S., CARLI, F. & GASTALDELLI, A. 2024. Hepatic glucose metabolism in the
steatotic liver. *Nat Rev Gastroenterol Hepatol,* 21**,** 319-334.
SERETIS, K., GOULIS, D. G., KOLIAKOS, G. & DEMIRI, E. 2015. Short- and Long-Term Effects of Abdominal
Lipectomy on Weight and Fat Mass in Females: a Systematic Review. *Obes Surg,* 25**,** 1950-8.
SETHI, J. K. & VIDAL-PUIG, A. 2005. Visfatin: the missing link between intra-abdominal obesity and
diabetes? *Trends Mol Med,* 11**,** 344-7.
SHI, H., STRADER, A. D., WOODS, S. C. & SEELEY, R. J. 2007a. The effect of fat removal on glucose
tolerance is depot specific in male and female mice. *Am J Physiol Endocrinol Metab,* 293**,** E1012-
20.
SHI, H., STRADER, A. D., WOODS, S. C. & SEELEY, R. J. 2007b. Sexually dimorphic responses to fat loss
after caloric restriction or surgical lipectomy. *Am J Physiol Endocrinol Metab,* 293**,** E316-26.
SHIN, J. A., JEONG, S. I., KIM, M., YOON, J. C., KIM, H. S. & PARK, E. M. 2015. Visceral adipose tissue
inflammation is associated with age-related brain changes and ischemic brain damage in aged
mice. *Brain Behav Immun,* 50**,** 221-231.
ST-ONGE, M. P. & GALLAGHER, D. 2010. Body composition changes with aging: the cause or the result of
alterations in metabolic rate and macronutrient oxidation? *Nutrition,* 26**,** 152-5.
STEINER, B. M. & BERRY, D. C. 2022. The Regulation of Adipose Tissue Health by Estrogens. *Front
Endocrinol (Lausanne),* 13**,** 889923.
SUN, R., HE, T., PAN, Y. & KATUSIC, Z. S. 2018. Effects of senescence and angiotensin II on expression and
processing of amyloid precursor protein in human cerebral microvascular endothelial cells.
Aging (Albany NY), 10**,** 100-114.
TCHKONIA, T., MORBECK, D. E., VON ZGLINICKI, T., VAN DEURSEN, J., LUSTGARTEN, J., SCRABLE, H.,
KHOSLA, S., JENSEN, M. D. & KIRKLAND, J. L. 2010. Fat tissue, aging, and cellular senescence.
Aging Cell, 9**,** 667-84.
TRAN, T. M., DAI, T. X. T., TRAN, D. B., NGUYEN, Q. C. T. & NGUYEN, D. H. Y. 2019. A simple
spectrophotometric method for quantifying total lipids in plants and animals. *CTU Journal of
Innovation and Sustainable Development,* 11**,** 106-110.
VISHVANATH, L. & GUPTA, R. K. 2019. Contribution of adipogenesis to healthy adipose tissue expansion
in obesity. *J Clin Invest,* 129**,** 4022-4031.
WANG, G., LI, G., SONG, A., ZHAO, Y., YU, J., WANG, Y., DAI, W., SALAS, M., QIN, H., MEDRANO, L., DOW,
J., LI, A., ARMSTRONG, B., FUEGER, P. T., YU, H., ZHU, Y., SHAO, M., WU, X., JIANG, L., CAMPISI,
J., YANG, X. & WANG, Q. A. 2025. Distinct adipose progenitor cells emerging with age drive
active adipogenesis. *Science,* 388**,** eadj0430.
WHITMER, R. A., GUSTAFSON, D. R., BARRETT-CONNOR, E., HAAN, M. N., GUNDERSON, E. P. & YAFFE, K.
2008. Central obesity and increased risk of dementia more than three decades later. *Neurology,*
71**,** 1057-64.
WONG ZHANG, D. E., TRAN, V., VINH, A., DINH, Q. N., DRUMMOND, G. R., SOBEY, C. G., JELINIC, M. & DE
SILVA, T. M. 2023. Pathophysiological Links Between Obesity and Dementia. *Neuromolecular
Med,* 25**,** 451-456.
WU, Y., LEE, M. J., IDO, Y. & FRIED, S. K. 2017. High-fat diet-induced obesity regulates MMP3 to
modulate depot- and sex-dependent adipose expansion in C57BL/6J mice. *Am J Physiol
Endocrinol Metab,* 312**,** E58-E71.
YANAI, H. & YOSHIDA, H. 2019. Beneficial Effects of Adiponectin on Glucose and Lipid Metabolism and
Atherosclerotic Progression: Mechanisms and Perspectives. *Int J Mol Sci,* 20.

ZHANG, P., KISHIMOTO, Y., GRAMMATIKAKIS, I., GOTTIMUKKALA, K., CUTLER, R. G., ZHANG, S.,
ABDELMOHSEN, K., BOHR, V. A., MISRA SEN, J., GOROSPE, M. & MATTSON, M. P. 2019. Senolytic
therapy alleviates Abeta-associated oligodendrocyte progenitor cell senescence and cognitive
deficits in an Alzheimer's disease model. *Nat Neurosci,* 22**,** 719-728.
ZHAO, Z., XIANG, X., CHEN, Q., DU, J., ZHU, S., XU, X., SHEN, Y., WEN, S., LI, Y., XU, W., MAI, K. & AI, Q.
2024. Sterol Regulatory Element Binding Protein 1: A Mediator for High-Fat Diet-Induced
Hepatic Gluconeogenesis and Glucose Intolerance in Fish. *J Nutr,* 154**,** 1505-1516.
ZHU, J., WU, C. & YANG, L. 2024. Cellular senescence in Alzheimer's disease: from physiology to
pathology. *Transl Neurodegener,* 13**,** 55.

 Figure Legends

Figure 1 - Novel object recognition and body composition analysis after VFR. Retention index

(A) and novelty preference (B) for the NOR task. Body weight (C) and the percentage of vWAT

(D) and total fat mass for three adipose depots (brown, subcutaneous, and visceral; E) in

relation to body weight. The percentage of VFR mice that regenerated vWAT (reg. ratio) are

shown in (D). Data are represented as means ± SEM. *p<0.05, ***p<0.001 based on a two-

tailed Student's *t* test (n=9-20).

Figure 2 – Lipid metabolism in hepatic and visceral white adipose tissue. Percentage of total

liver lipids (A). Relative gene expression profile of lipolysis and lipogenesis genes in the liver (B)

and vWAT (C). Data are represented as means ± SEM. *p<0.05, **p<0.01, ***p<0.001,

****p<0.0001 based on a two-tailed Student's *t* test (n=8-15). Abbreviations: ACC – acetyl CoA-

carboxylase, FAS – fatty acid synthase, SCD1 - stearoyl-CoA Desaturase, Elovl6 – fatty acid

elongase enzyme 6, SREBP1 – sterol regulator element-binding protein 1, MGL –

monoglyceride lipase, PKA – protein kinase A, ATGL – adipose triglyceride lipase, HSL –

hormone sensitive lipase

Figure 3 – Blood glucose levels in pre-symptomatic and symptomatic APP^{NL-F} mice. Glucose

tolerance test (GTT; A) and insulin tolerance test (ITT; B) and their respective area under the

curve (AUC; inset) in pre-symptomatic (left) and symptomatic (right) cohorts. Fasting (C) and

fed (D) blood glucose levels were determined from t=0 during the GTT and ITT, respectively.

Data are represented as means ± SEM. *p<0.05 based on a two-tailed Student's t test (n=9-16).

Figure 4 – Liver glucose metabolism and plasma adipokines. Hepatic relative gene expression

profiles (A) of gluconeogenesis (G6PC, PEPCK), glycolysis (GLUT2, GCK), and glycogenesis

(GYS2). Circulating visfatin (B), adiponectin (C), and Leptin (D) determined by ELISA from

plasma collected at the time of euthanization. Data are represented as means ± SEM. *p<0.05

based on a two-tailed Student's t test (n=4-12). Abbreviations: G6PC - glucose-6-phosphatase,

PEPCK - phosphoenolpyruvate carboxykinase, Glut2 - glucose transporter 2, GCK -

glucokinase, GYS2 - glycogen synthase 2 (GYS2)

Figure 5 – Senescent cell and inflammation markers in visceral and subcutaneous white

adipose tissue. Visceral (A) and subcutaneous white adipose tissue (SubQ; B) relative gene

expression profiles of senescence ($p16^{Ink4\alpha}$, $p21^{Cip1}$) and inflammation (TNFα, IL-10, IL-6,

MCP1). Data are represented as means ± SEM. *p<0.05, **p<0.01, ***p<0.001 based on a two-

tailed Student's t test (n=4-12). Abbreviations: TNFα – tumor necrosis factor alpha, IL –

interleukin, MCP1 – monocyte chemoattractant protein 1.

Figure 6 - *Senescent cell and inflammation markers in the hippocampus.* Hippocampal relative

gene expression profiles (A) of senescence ($p16^{Ink4\alpha}$, $p21^{Cip1}$), inflammation (TNFα), and

memory (AdipoR, LepR, Creb1, PRGN) and soluble $A\beta_{42}$ (B) as determined by ELISA. Data are

represented as means ± SEM. *p<0.05, **p<0.01, ***p<0.001 based on a two-tailed Student's t

test (n=4-12).   Abbreviations: TNFα – tumor necrosis factor alpha, AdipoR – adiponectin

receptor, LepR – leptin receptor, Creb1 – cyclic AMP Response Element-Binding Protein 1.

Table 1: A list of forward and reverse primers. The last gene listed for each tissue type was used as that tissue's housekeeping gene.

Gene Name (Abbreviation)	Forward Primer	Reverse Primer
Liver		
Acetyl-CoA Carboxylase (ACC)	5'-ATGGGCGGAATGGTCTCTTTC-3'	5'-TGGGGACCTTGTCTTCATCAT-3'
Fatty Acid Synthase (FAS)	5'-GGAGGTGGTGATAGCCGGTAT-3'	5'-TGGGTAATCCATAGAGCCCAG-3'
Stearoyl-CoA Desaturase (SCD1)	5'-TTCTTGCGATACACTCTGGTGC-3'	5'-CGGGATTGAATGTTCTTGTCGT-3'
Fatty Acid Elongase Enzyme 6 (Elovl6)	5'-GAAAAGCAGTTCAACGAGAACG-3'	5'-AGATGCCGACCACCAAAGATA-3'
Sterol Regulator Element-Binding Protein 1 (SREBP1)	5'-GATGTGCGAACTGGACACAG-3'	5'-CATAGGGGGCGTCAAACAG-3'
Monoglyceride Lipase (MGL)	5'-CGGACTTCCAAGTTTTTGTCAGA-	5'-GCAGCCACTAGGATGGAGATG-3'
Glucose-6-Phosphatase (G6PC)	5'-CGACTCGCTATCTCCAAGTGA-3'	5'-GTTGAACCAGTCTCCGACCA-3'
Phosphoenolpyruvate Carboxykinase (PEPCK)	5'-CTGCATAACGGTCTGGACTTC-3'	5'-CAGCAACTGCCCGTACTCC-3'
Glucose Transporter 2 (GLUT2)	5'-TCAGAAGACAAGATCACCGGA-3'	5'-GCTGGTGTGACTGTAAGTGGG-3'
Glucokinase (GLK)	5'-TGAGCCGGATGCAGAAGGA-3'	5'-GCAACATCTTTACACTGGCCT-3'
Glycogen Synthase (GYS2)	5'-ACCAAGGCCAAAACGACAG-3'	5'-GGGCTCACATTGTTCTACTTGA-3'
Beta-2 Microglobulin (BDM)	5'-AAGTATACTCACGCCACCCA-3'	5'-AGGACCAGTCCTTGCTGAAG-3'
Visceral White Adipose Tissue		
Acetyl-CoA Carboxylase (ACC)	5'-ATGGGCGGAATGGTCTCTTTC-3'	5'-TGGGGACCTTGTCTTCATCAT-3'
Fatty Acid Synthase (FAS)	5'-GGAGGTGGTGATAGCCGGTAT-3'	5'-TGGGTAATCCATAGAGCCCAG-3'
Protein Kinase A (PKA)	5'-AGATCGTCCTGACCTTTGAGT-3'	5'-GGCAAAACCGAAGTCTGTCAC-3'
Adipose Triglyceride Lipase (ATGL)	5'-GGATGGCGGCATTTCAGACA-3'	5'-CAAAGGGTTGGGTTGGTTCAG-3'
Hormone Sensitive Lipase (HSL)	5'-CCAGCCTGAGGGCTTACTG-3'	5'-CTCCATTGACTGTGACATCTCG-3'
Sterol Regulator Element-Binding Protein 1 (SREBP1)	5'-GATGTGCGAACTGGACACAG-3'	5'-CATAGGGGGCGTCAAACAG-3'
p16$^{Ink4\alpha}$	5'-CGGTCGTACCCCGATTCAG-3'	5'-GCACCGTAGTTGAGCAGAAGA-3'
p21^{Cip1}	5'-CCTGGTGATGTCCGACCTG-3'	5'-CCATGAGCGCATCGCAATC-3'
Tumor Necrosis Factor α (TNFα)	5'-CAGGCGGTGCCTATGTCTC-3'	5'-CGATCACCCCGAAGTTCAGTAG-3'
Interleukin-10 (IL-10)	5'-GCTCTTACTGACTGGCATGAG-3'	5'-CGCAGCTCTAGGAGCATGTG-3'
Glyceraldehyde-3-Phosphate Dehydrogenase (GAPDH)	5'-AGGTCGGTGTGAACGGATTTG-3'	5'-TGTAGACCATGTAGTTGAGGTCA-3'
Subcutaneous White Adipose Tissue		
Tumor Necrosis Factor α (TNFα)	5'-CAGGCGGTGCCTATGTCTC-3'	5'-CGATCACCCCGAAGTTCAGTAG-3'

Interleukin-10 (IL-10)	5'-GCTCTTACTGACTGGCATGAG-3'	5'-CGCAGCTCTAGGAGCATGTG-3'
Interleukin-6 (IL-6)	5'-CTGCAAGAGACTTCCATCCAG-3'	5'-AGTGGTATAGACAGGTCTGTTGG-3'
Monocyte Chemoattractant Protein-1 (MCP1)	5'-CCACTCACCTGCTGCTACTCAT-3'	5'-TGGTGATCCTCTTGTAGCTCTCC-3'
Beta-2 Microglobulin (BDM)	5'-AAGTATACTCACGCCACCCA-3'	5'-AGGACCAGTCCTTGCTGAAG-3'
Hippocampus		
p16$^{Ink4\alpha}$	5'-CGGTCGTACCCCGATTCAG-3'	5'-GCACCGTAGTTGAGCAGAAGA-3'
p21^{Cip1}	5'-CCTGGTGATGTCCGACCTG-3'	5'-CCATGAGCGCATCGCAATC-3'
Tumor Necrosis Factor α (TNFα)	5'-CAGGCGGTGCCTATGTCTC-3'	5'-CGATCACCCCGAAGTTCAGTAG-3'
Adiponectin Receptor 1 (AdipoR1)	5'-TCTTCGGGATGTTCTTCCTGG-3'	5'-TTTGGAAAAAGTCCGAGAGACC-3'
cAMP Response Element-Binding Protein 1 (CREB1)	5'-CCCAAAAACGAAGGGAAATCCT-3'	5'-CCTGGTGCATCAGAAGATAAGTC-3'
Leptin Receptor (LEPR)	5'-TGGTCCCAGCAGCTATGGT-3'	5'-ACCCAGAGAAGTTAGCACTGT-3'
Progranulin (PGRN)	5'-ATGTGGGTCCTGATGAGCTG-3'	5'-GCTCGTTATTCTAGGCCATGTG-3'
Ubiquitin Conjugating Enzyme E2 D2 (UBE2D2	5'-TGCCTGAGATTGCTCGGATCT-3'	5'-TCGCATACTTCTGAGTCCATTCC-3'

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6