Inflammation increases the penetrance of behavioral impairment in *Shank3*

haploinsufficiency mice – can it explain the behavioral regression in Autism?

Authors: Sheng-Nan Qiao, PhD[1], Sung Eun Wang, PhD[1, 6], Kun-Yong Kim, PhD[1],

Sungsin Jo, PhD[7], Yong-Hui Jiang, MD, PhD [1-5]

Affiliation: [1]Department of Genetics, [2]Department of Neuroscience, [3]Department of

Pediatrics, and [4]Department of Obstetrics, Gynecology & Reproductive Sciences, Yale

University School of Medicine, 333 Cedar Street, New Haven, CT 06510 USA

[5]Wu-Tsai Institute, Yale University, 100 College Street, New Haven CT 06520 USA

[6]Department of Systems Biology, Yonsei University, Yonsei-ro 50, Seodaemun-gu, Seoul,

South Korea

[7]Department of Biology, Soonchunhyang University, Soonchunhyang-ro 22, Sinchang-

myeon, Asan, South Korea

Correspondence:

Yong-Hui Jiang, MD, PhD.

Department of Genetics, Neuroscience, Pediatrics, and OB/Gyn & Reproductive

Sciences

TAC S317/S320

Yale University School of Medicine

300 Cedar St, New Haven, CT 06519

Tel: 203-785-2429

Email:yong-hui.jiang@yale.edu

Running title: Neuroinflammation Susceptibility in Autism

Abstract

Behavioral regression occurs in ~40% of *SHANK3*-associated autism spectrum disorder (ASD). We previously reported that significant behavioral regression in a small cohort with *SHANK3* haploinsufficiency patients, triggered by subclinical infections, responded to immunomodulator treatments. We hypothesize that behavioral regression results from the interplay between *SHANK3* deficiency and neuroinflammation. Using *Shank3* exon 4-22 deletion heterozygous mutant (*Sh3*[+/-]) mouse, which shows no significant behavior impairments, we established a preclinical model – *Shank3* haploinsufficiency mouse undergoing systemic inflammation challenge via intraperitoneal injection of lipopolysaccharides (LPS). We found that, two weeks after LPS challenge, wild-type mice (WT) recovered but *Sh3*[+/-] mice exhibited motor impairment, anxiety-like behaviors, and excessive grooming, similar to *Shank3* exon 4-22 deletion homozygous mutants. Anti-inflammatory treatment partially reversed LPS-induced behavioral changes. Transcriptomic analysis revealed upregulation of neuroinflammation-related genes and downregulation of synaptic function-related genes in *Sh3*[+/-] mice in response to LPS. Especially, pro-inflammatory genes and microglia markers were overly activated that may result from the increased toll-like receptor 4 (TLR4) in *Sh3*[+/-] mice. Microglia overactivation elevated synapse engulfment and disrupted synaptic protein may underlie LPS-triggered worsen behavior phenotypes in *Sh3*[+/-] mice. Our findings indicate that neuroinflammation increases the penetrance of behavioral impairment in *Shank3* haploinsufficiency mice and support a potential mechanism for the behavioral regression in human *SHANK3* disorders for future investigations.

Introduction

Autism spectrum disorder (ASD) is a group of neurodevelopmental conditions with typical onset before the age of three [1]. A significant proportion of children with ASD shows developmental arrest at an early developmental age and behavioral regression throughout the lifespan. Behavioral regression is characterized by loss of previously acquired skills, including loss of purposeful hand use, loss of acquired speech, and gait abnormalities [2-4]. Catatonic features are frequently associated with behavioral regression observed in ASD [5,6]. However, the underlying mechanism for the early developmental arrest and behavioral regression remains poorly understood.

The *SHANK3* mutation is one of the most common monogenic defects associated with ASD in humans [7-10]. Genetic mutations in *SHANK3* gene can manifest as Phelan-McDermid Syndrome (PMS), characterized by a chromosomal deletion of 22q13.3 or single nucleotide variants, indels, exonic deletions within the *SHANK3* gene in individuals with primary ASD [11]. Patients with *SHANK3* haploinsufficiency also exhibit severe intellectual disability, motor deficits, and speech delays [12-14]. Developmental regression with catatonic features has been reported in ~40% of cases with *SHANK3* haploinsufficiency [15-20]. We previously described four patients with pathogenic *SHANK3* mutations who experienced acute and severe behavioral regression following possible subclinical viral infections [21]. Notably, treatment with intravenous immunoglobulins (IVIG) combined with anti-inflammatory drugs and immunomodulators significantly reversed the behavioral regression in these cases [21]. This clinical report strongly suggests a potential link between inflammation and behavioral regression associated with *SHANK3*

haploinsufficiency. However, the direct causality could not be investigated and determined

from human studies.

SHANK3 protein is primarily characterized as a scaffold protein located at the

postsynaptic density of neurons, interacting with other synaptic proteins and affecting

synaptic function [22]. However, the role of SHANK3 in function of non-neuronal cells such

as microglia has been poorly studied [23-26]. Microglia activation can sculpt synapses and

brain circuits and change animal behavioral phenotypes during normal development as

well as in response to neuroinflammation [27,28]. Aberrant microglia activation has been

associated with ASD[29-32]. Our previous studies showed that inflammation induces acute

behavioral changes [33] and aggravates hypothermia in *Shank3* mutant mouse model [34].

These initial findings suggest a potential mechanistic link between *SHANK3* deficiency

and inflammatory response.

Our lab previously generated a *Shank3* mutant mouse line with a deletion of exon

4 to 22 (*Shank3$^{Δe4-22}$*) that disrupts most of the *Shank3* coding region[35]. *Shank3$^{Δe4-22}$*

homozygous but not heterozygous mice showed robust face validity for autism-like

behaviors[35]. Therefore, *Shank3$^{Δe4-22}$* heterozygous mice offer a unique opportunity to

examine the interaction between genes and environment, specifically whether

neuroinflammation may modify the penetrance of behavioral phenotypes associated with

SHANK3 deficiency in humans. In this study, we utilized the *Shank3$^{Δe4-22}$* heterozygous

mice and investigated the behavioral outcome, transcriptomic profiles, microglia

activation, and synapse marker after systemic inflammatory challenge induced by

lipopolysaccharides (LPS). These findings provide a general insight on the role of

neuroinflammation in ASD susceptibility.

Results

Systemic inflammation via LPS modified the penetrance of motor dysfunction, anxiety-like behavior, and repetitive behavior in *Shank3*[Δe4-22] heterozygous (*Sh3*[+/-]) mice.

To explore whether inflammation plays a role in regression of ASD-associated behaviors, we challenged *Sh3*[+/-] mice by intraperitoneal injection of LPS (*Sh3*[+/-]+ LPS). Control mice were given the same volume of normal saline (PBS) (*Sh3*[+/-] + PBS)*.* Both LPS-treated *Sh3*[+/-] and wild-type (WT) mice experienced weight loss at 24 hours compared to PBS-treated mice (**Fig. 1A**). This indicated systemic inflammation induced by LPS, consistent with previous reports [36]. LPS-treated mice regained body weight at 72 hours post-injection, suggesting recovery from the LPS challenge (**Fig. 1A**). Acute behavioral changes, including decreased locomotor activity [37], social avoidance [38], and depressive-like behavior [39], have been reported in LPS-treated mice around 24 hours post-injection. However, interpretating these behavioral changes in the context of ASD model may be confounded by immediate sickness of test mice [33,40,41]. Therefore, we performed behavioral analysis two weeks after LPS injection to distinguish "LPS and SHANK3 haploinsufficiency interaction effect" vs. "acute LPS effect" (**Fig. 1B**). The *Shank3*[Δe4-22] homozygous (*Sh3*[-/-]) group was included for a comparison [35].

To assess the motor coordination, latency to fall from the rotarod was measured in both steady speed (16 rpm) and accelerated speed (4-40 rpm) conditions. As expected, *Sh3*[-/-] mice showed significantly reduced latency to fall compared to WT+PBS group. No significant difference was observed between WT+PBS and WT+LPS groups, but *Sh3*[+/-] mice treated with LPS showed significantly shorter latency to fall than the PBS group (**Fig.**

1C-D). The duration of mouse spent in center of open field test (OFT) showed no significant difference between WT+PBS and WT+LPS groups, but $Sh3^{+/-}$+LPS spent significantly less time in the center compared to $Sh3^{+/-}$+PBS (**Fig. 1E**). In the light/dark box test, distance of mouse moved in light chamber showed no significant difference between WT+PBS and WT+LPS groups while $Sh3^{+/-}$+LPS significantly moved less in light chamber compared to $Sh3^{+/-}$+PBS (**Fig. 1F**). Repetitive behavior was assayed through scoring grooming behavior. $Sh3^{+/-}$+LPS showed significantly longer grooming duration than $Sh3^{+/-}$+PBS while WT+PBS and WT+LPS groups were comparable (**Fig. 1G**). We also conducted a three-chamber social preference test. As expected, $Sh3^{-/-}$ mice showed no preference for social targets compared to WT. However, both LPS- and PBS-treated $Sh3^{+/-}$ and WT mice spent significantly more time in the chamber with the social target (S) or in the proximity region of the social target (SP) (**Fig. 1H-I**).

To explore potential sex effect, we compared motor function and sociability in both PBS- and LPS-treated female and male $Sh3^{+/-}$ mice. LPS-treated male $Sh3^{+/-}$ mice exhibited significant reduced latency to fall in rotarod, although no significant difference found between LPS and PBS group in female $Sh3^{+/-}$ mice (**Supplementary Fig. 1A-B**). Both LPS- and PBS-treated male $Sh3^{+/-}$ mice spent significantly more time in S or SP, but no significant difference found in female $Sh3^{+/-}$ mice (**Supplementary Fig. 1C-D**). Taken together, our results indicated that, LPS treatment in $Sh3^{+/-}$ mice significantly impaired the motor performance on rotarod, changed exploratory or anxiety-like behaviors in OFT and light/dark box test, and increased grooming duration, but did not affect sociability in three-chamber test. By comparison, LPS treatment in WT mice did not have any significant effect on those behaviors. These findings suggest the interplay of inflammation and

Shank3 haploinsufficiency in select domains of behaviors.

To assess whether anti-inflammation treatment can reverse the behavioral

changes induced by LPS in *Sh3+/-* mice, we applied mefenamic acid (MFA), a nonsteroidal

anti-inflammatory drug (NSAID). One of the MFA targets is the intracellular

cyclooxygenase (COX) signaling, which can be activated by LPS [42-44]. Previous study

also reported that MFA has neuroprotective effect [45]. Given that both WT+LPS and *Sh3+/-*

+LPS group exhibited impaired motor function at 1 week after the LPS injection

(**Supplementary Fig. 2A-B**) but such behavior deficits of WT+LPS group recovered to

baseline at 2-week post-injection (**Fig. 1C-D**), we chose day 7 post-injection as the

endpoint of MFA treatment (**Fig. 2A**). We administered MFA via intraperitoneal injection

once a day for seven days following LPS injection and then performed rotarod test 24

hours after the final dose of MFA to evaluate the treatment efficacy (**Fig. 2A**). In the steady

speed (16 rpm) rotarod test, both WT+LPS+MFA and *Sh3+/-*+LPS+MFA groups showed

significantly increased latency to fall compared to vehicle-treated groups (**Fig. 2B**),

suggesting that MFA effectively mitigated the motor function deficits induced by LPS.

Sh3+/-+LPS+MFA showed trend of shorter latency to fall than WT+LPS+MFA (**Fig. 2B**, *p*

= 0.091), suggesting potential less pronounced anti-inflammatory efficacy and the

presence of exacerbated inflammatory responses in *Sh3+/-* mice. In contrast, in the

accelerated speed (4-40 rpm) rotarod test, MFA did not restore motor function in either

WT or *Sh3+/-* mice (**Fig. 2C**). Moreover, when comparing MFA treatment in female and

male *Sh3+/-* mice, MFA significantly increased latency to fall in female *Sh3+/-* mice but not

in male *Sh3+/-* mice (**Fig. 2D-E**). Overall, our results showed that LPS-induced motor

function impairment in *Sh3+/-* mice can be partially reversed by anti-inflammatory

treatment. Future studies will include additional behavioral assays, as well as time-course

and dose-dependent responses to MFA treatment.

LPS-induced differentially expressed genes (DEGs) were associated with

neuroinflammation and neurotransmission in $Sh3^{+/-}$ mice.

163   To understand how LPS-induced inflammation modifies the penetrance of motor

impairment in $Sh3^{+/-}$ mice, we analyzed the transcriptomic profiles in the forebrains by

bulk RNA-sequencing (RNA-seq) (**Supplementary Fig. 3A-B, Supplementary Tables 1**

and 2). LPS treatment induced hundreds of differentially expressed genes (DEGs) in both

WT and $Sh3^{+/-}$ mice compared to PBS-treated groups. The number of upregulated genes

(488 DEGs in $Sh3^{+/-}$+LPS vs. $Sh3^{+/-}$+PBS and 433 DEGs in WT+LPS vs. WT+PBS) was

greater than that of downregulated genes (75 DEGs in $Sh3^{+/-}$+LPS vs. $Sh3^{+/-}$+PBS and

108 DEGs in WT+LPS vs. WT+PBS) (**Fig. 3A-B**). We identified 88 upregulated and 37

downregulated DEGs between $Sh3^{+/-}$+LPS vs. WT+LPS (**Fig. 3C**). Although we found 67

upregulated and 33 downregulated genes when comparing transcriptomic profiles

between $Sh3^{+/-}$+PBS and WT+PBS (**Fig. 3D**), those genes barely overlapped with any

LPS-induced DEGs observed in $Sh3^{+/-}$ mice. Next, we compared the overlapping genes

among four comparisons (**Fig. 3E-F, Supplementary Tables 3 and 4**). A total of 313

shared upregulated and 12 shared downregulated DEGs were identified between

WT+LPS vs. WT+PBS and $Sh3^{+/-}$+LPS vs. $Sh3^{+/-}$+PBS. Those shared DEGs probably

represent the LPS-induced neuroinflammatory responses in mouse brains. Eight

upregulated genes shared among $Sh3^{+/-}$+LPS vs. WT+LPS, $Sh3^{+/-}$+LPS vs. $Sh3^{+/-}$+PBS

and WT+LPS vs. WT+PBS were well-known for their roles in neuroinflammation (**Fig. 3E**

highlighted with white underline, **Table 1**). Interestingly, 10 upregulated genes and 5

downregulated genes were found only in $Sh3^{+/-}$ mice not in WT mice in response to LPS

(**Fig. 3E-F** highlighted with green underline, **Table 2-3**), suggesting that LPS-induced

neuroinflammation in $Sh3^{+/-}$ mice is mediated by distinct signaling pathways. Additionally,

when we compared the overlapping genes between $Sh3^{+/-}$+PBS vs. WT+PBS and $Sh3^{+/-}$

+LPS vs. WT+LPS, only 4 genes in total (two are protein-coding genes) were identified

(**Fig. 3E-F** highlighted with red underline). Particularly, *Acr* is a *Shank3* subordinate

adjacent gene, whose expression increased when *Shank3* is deleted [46]. To further

characterize the function of DEGs from RNA-seq, we performed the pathway analysis

using the QIAGEN Ingenuity Pathway Analysis (IPA) [47]. We identified more activated

inflammation-associated pathways in response to LPS challenges in both $Sh3^{+/-}$ and WT

(**Supplementary Fig. 4 and 5**). In contrast, inhibited neurotransmission-related pathways

including NMDA receptor and calcium signaling pathways were only observed in LPS-

treated $Sh3^{+/-}$ but not in LPS-treated WT (**Supplementary Fig. 4 and 5**). To understand

the distinct molecular pathways in LPS-treated $Sh3^{+/-}$ mice, we delved into the pathway

analysis from $Sh3^{+/-}$+LPS vs. WT+LPS (**Supplementary Fig. 6**). The top five activated

pathways were all related to inflammatory processes. The top two inhibited pathways,

cardiac conduction and CREB signaling in neurons, represent inhibited neuron signal

transmission and neuronal functions. The other three inhibited pathways, HIF1α signaling,

eNOS signaling and phagosome formation pathways, represent increased neuronal

vulnerability to stress and inflammatory stimuli. In addition to the top five pathways with

highest z score value, we noticed that most of the upregulated signaling pathways in

response to LPS were all inflammation-related, including cytokine storm pathway and

neuroinflammation pathway (**Fig. 3G**). In contrast, the downregulated pathways in LPS-

treated *Sh3⁺ᐟ⁻* mice were neuronal and synaptic function-related, including potassium

channel pathway, glutamatergic signaling pathway, acetylcholine signaling pathway, and

gap junction signaling pathway (**Fig. 3G**). Overall, our RNA-seq results suggested that a

combination of exacerbated neuroinflammatory responses and impaired neuronal

functions is attributed to the LPS-induced behavioral regression in *Sh3⁺ᐟ⁻* mice.

Neuroinflammation was increased in *Sh3⁺ᐟ⁻* mice

To investigate how LPS leads to aggravated neuroinflammation in *Sh3⁺ᐟ⁻* brains,

we first measured the RNA expression of the pro-inflammatory cytokines *Il1b* and

chemokines *Cxcl10* using RT-qPCR. *Il1b* and *Cxcl10* have been identified as common

upregulated genes across all LPS-treated groups in RNA-seq (**Table 1**). RT-qPCR

confirmed that LPS significantly increased *Il1b* and *Cxcl10* expression in both WT and

Sh3⁺ᐟ⁻ mice compared to PBS groups. Although no significant difference in *Il1b* and *Cxcl10*

expression was found between *Sh3⁺ᐟ⁻*+PBS vs. WT+PBS, *Sh3⁺ᐟ⁻*+LPS showed

significantly increased expression compared to WT+LPS group (**Fig. 4A-B**). A previous

study showed that LPS treatment upregulates microglia marker gene *Cx3cr1* while

downregulates microglia homeostasis gene *P2ry12* expression [48]. Therefore, we

examined the expression of microglia markers *Cx3cr1* and *P2ry12* using RT-qPCR. We

found that, as expected, LPS significantly increased *Cx3cr1* expression and decreased

P2ry12 expression in both WT and *Sh3⁺ᐟ⁻* mice compared to PBS groups (**Fig. 4C-D**).

However, *Cx3cr1* expression was comparable between *Sh3⁺ᐟ⁻*+PBS and WT+PBS but

significantly increased in *Sh3⁺ᐟ⁻*+LPS compared to WT+LPS group (**Fig. 4C**). In contrast,

P2ry12 expression was significantly higher in *Sh3⁺ᐟ⁻*+PBS than in WT+PBS and higher in

Sh3[+/-]+LPS than in WT+LPS (**Fig. 4D**).

We further examined microglia activation by examining the expression of IBA-1, a

marker widely used for the assessment of microglia activation state [49] (**Fig. 4E-F**).

Compared to PBS treatment, LPS increased IBA-1 staining intensity significantly at 3

hours and 24 hours after LPS treatment in both WT and *Sh3*[+/-] mice. However, *Sh3*[+/-]

+LPS showed significantly higher microglia activation than WT+LPS group at 24 hours

but not at 3 hours post-injection. These results indicated that the prolonged activation of

microglia contributed to increase of neuroinflammatory responses in *Sh3*[+/-] mice.

Next, to understand the mechanism underlying the LPS-induced microglia

overactivation in *Sh3*[+/-] mice, we analyzed the LPS binding receptor toll-like receptor-4

(TLR-4) expression in WT+LPS and *Sh3*[+/-]+LPS groups at 24 hours post-injection [50]. We

found that TLR4 expression was significantly higher in *Sh3*[+/-] than WT with either PBS or

LPS treatment (**Fig. 4G-H**). Additionally, we measured the SHANK3 protein expression

and found that either PBS- or LPS-treated *Sh3*[+/-] groups showed a reduction in SHANK3

protein expression compared to WT groups as expected, but no changes in SHANK3

expression were observed after LPS treatment (**Fig. 4G, I**). TLR4 is known to associate

with LPS-induced microglia activation[51-54]. To investigate whether observed TLR4

augmentation directly facilitates microglia overactivation in LPS treated *Sh3*[+/-] mice, we

analyzed colocalization between TLR4 and IBA1 (**Fig. 4J-P**). We found that TLR4 and

IBA1 expression were well-colocalized in WT, but a small fraction of TLR4 signals in *Sh3*[+/-]

mice were not colocalized with IBA1 (**Fig 4J-K**). Although colocalization Pearson's

coefficient was similar between WT+LPS and *Sh3*[+/-]+LPS group, colocalization coefficient

normalized by TLR4 was lower in *Sh3*[+/-] mice and the colocalization coefficient normalized

by IBA1 was similar (**Fig. 4L-N**). Interestingly, cellular immunofluorescence intensity of

TLR4 in *Sh3*[+/-] mice is slightly lower than WT (**Fig. 4P**). In addition, cellular

immunofluorescence intensity of IBA1 was significantly higher in *Sh3*[+/-] brains (**Fig. 4O**).

Given that TLR4 was expressed in many different cell types including astrocytes,

neutrophils, and brain endothelial cells [55,56], these findings suggested that elevated TLR4

expression in other cell types might contribute to microglia activation in *Sh3*[+/-] mice.

Microglia activation affected synaptic structure in *Sh3*[+/-] mice

To uncover how microglia overactivation leads to behavioral changes in *Sh3*[+/-] mice,

we examined microglia engulfment and synaptic structures. We compared lysosome

engulfment (lysosome marker CD68 occupancy) and synapse engulfment (presynaptic

marker vGluT1 occupancy) between WT+LPS and *Sh3*[+/-]+LPS brains at 24 hours post-

injection. While CD68 occupancy was comparable between the two groups, vGluT1

occupancy — both within CD68-positive lysosomal puncta in microglia (CD68[+]IBA1[+]) and

within the microglial cytoplasm (IBA1[+]) — was significantly higher in *Sh3*[+/−] mice

compared to WT mice (**Fig. 5A-D**). These findings suggest that microglial activation is

associated with increased phagocytosis of synaptic materials in the *Sh3*[+/−]+LPS group.

To determine whether increased synaptic engulfment leads to synapse dismantling

in the *Sh3*[+/−] brain, we examined the colocalization of the presynaptic marker vGluT1 with

the postsynaptic density marker PSD95 at 24 hours following LPS injection. In WT mice,

vGluT1 and PSD95 puncta were well colocalized (**Fig. 5E**, arrowhead). In contrast, *Sh3*[+/−]

mice displayed discrete PSD95 clusters that did not colocalize with vGluT1 puncta (**Fig.**

5E, arrow). Quantitative analysis revealed that the *Sh3*[+/−]+LPS group showed a

significantly lower Pearson's correlation coefficient for colocalization, a reduced

colocalization coefficient normalized by PSD95, and decreased overall synaptic puncta

density compared to the WT+LPS group (**Fig. 5F-H**).

Next, we compared the colocalization of vGluT1 and PSD95 among four groups

— WT+PBS, WT+LPS, *Sh3*$^{+/-}$+PBS, and *Sh3*$^{+/-}$+LPS — at two weeks after injection. In

both WT+PBS and WT+LPS groups, vGluT1 and PSD95 puncta remained closely aligned

(**Fig. 5I**). The *Sh3*$^{+/-}$+PBS group also exhibited substantial colocalization of vGluT1 and

PSD95; however, some displaced PSD95 fragments were observed, which became more

pronounced in the *Sh3*$^{+/-}$+LPS group (**Fig. 5J,** arrow). Consistent with the findings at 24

hours post-injection, the *Sh3*$^{+/-}$+LPS group showed a significantly lower Pearson's

correlation coefficient for colocalization, a reduced colocalization coefficient normalized

by PSD95, and decreased synapse density compared to the other groups (**Fig. 5K-M**).

These results support the conclusion that microglial overactivation enhances synapse

engulfment and contributes to synaptic structure disorganization in *Sh3*$^{+/-}$ mice in

response to LPS-induced inflammation.

To investigate whether disrupted synaptic structure caused PSD protein reduction

and synaptic dysfunction in *Sh3*$^{+/-}$ mice, we explored excitatory postsynaptic scaffold

proteins PSD95 and HOMER 1b/c expression (**Fig. 5N**). No significant change of PSD95

was observed between *Sh3*$^{+/-}$+PBS vs. WT+PBS. However, PSD95 was significantly

decreased in the *Sh3*$^{+/-}$+LPS compared to both the WT+LPS group (~28% reduction) and

the *Sh3*$^{+/-}$+PBS group (~17% reduction), suggesting that LPS treatment reduced the

PSD95 protein level in *Sh3*$^{+/-}$ mice not in WT mice (**Fig. 5O**). In contrast, HOMER 1b/c

expression was significantly decreased in *Sh3*$^{+/-}$+PBS vs. WT+PBS (~41% reduction) and

in *Sh3*$^{+/-}$+LPS vs. WT+LPS (~23% reduction) but showed no significant change between

Sh3[+/-]+LPS vs. *Sh3*[+/-]+PBS. It indicated that LPS treatment did not further reduce

HOMER expression (**Fig. 5P**). PSD95 is critical for excitatory synapse function through

its interactions with NMDA and AMPA receptors[57]. We examined whether the change of

PSD95 may lead to the change of the expression of major subunits of NMDA receptor

subunit NR2A and the AMPA receptor subunit GluA1 associated with LPS treatment (**Fig.**

5Q). We found that NR2A expression was reduced in the *Sh3*[+/-]+LPS group compared to

both the WT+LPS group and the *Sh3*[+/-]+PBS group (**Fig. 5R**). The GluA1 expression was

significantly reduced in *Sh3*[+/-]+LPS mice compared to both the WT+LPS group and

Sh3[+/-]+PBS group **(Fig. 5S)**. Taken together, these results suggested a mechanistic

convergence between LPS-induced neuroimmune responses, and the synaptic

disruptions known to arise from Shank3 deficiency.

Table 1. Eight Upregulated DEGs shared among *Sh3*[+/-]+LPS vs. WT+LPS, *Sh3*[+/-]+LPS vs. *Sh3*[+/-]+PBS and WT+LPS vs. WT+PBS.

Gene name	Protein function
Bcl3	BCL3 is an autonomous determinant of memory/terminal effector cell balance during CD8[+] T cell differentiation in response to acute viral infection [58].
Ccl3	CCL3 as a cytokine contributes to progressive tissue damage and functional impairment during secondary injury after spinal cord injury [59].
Cxcl10	CXCL10 as a lymphocyte chemoattractant contributes to the intraparenchymal trafficking of lymphocytes during acute CNS inflammation [60].
Gbp5	GBP5 as a GTPase participates in the regulation of immunity and neuroinflammation [61].
Ifitm6	IFITM6 encodes cell surface proteins that modulate cell-cell adhesion and cell differentiation, is induced in various macrophages, and increases expression 24 hours after LPS [62].
Il1b	IL-1β can be acutely induced in the brain following peripheral bacterial infections, and microglia facilitate exaggerated IL-1β production upon exposure to acute systemic inflammation induced by bacterial LPS [63].
Steap4	STEAP4 as a key effector molecule contributes to the pathogenesis of neuroinflammation in experimental autoimmune encephalomyelitis [64].
Tlr2	Toll-like receptors (TLRs) are fundamental sensor molecules of the host innate immune system, which detect conserved molecular signatures of a wide range of microbial pathogens and initiate innate immune responses via distinct signaling pathways [65].

Table 2. Ten upregulated DEGs only shared between *Sh3*^{+/-}+LPS vs. WT+LPS and *Sh3*^{+/-}+LPS vs. *Sh3*^{+/-}+PBS, not include in WT+LPS vs. WT+PBS.

Gene name	Protein function
Alms1-ps1	ALMS1, centrosome and basal body associated, pseudogene 1.
Atf3	ATF3 as a transcription factor responds to and exert dual effects on inflammatory responses and in involved in neuroinflammatory response [66].
Ccl12	CCL12 as a chemokine expressed by microglia is involved in inflammation [67].
Fas	Fas/FasL pathway plays a key role in immune homeostasis and leads to excessive neuroinflammation during excessive neuroinflammation [68].
Gm11681	long non-coding RNA.
Hspa1a *Hspa1b* *Hspb1*	HSPA1A and HSPA1B are two subtypes of HSPA1 proteins. HSPA1 is a member of the stress-inducible 70-kDa heat shock protein (HSP70) family, which plays a critical role in the cellular stress response [69,70].
Oas1g	OAS1G is an important murine antiviral response factor, function through activating RNase L to degrade viral RNA in turn inhibiting viral replication and propagation [71].
Prok2	PROK2 is a secreted protein involved in the pathogenesis of several acute and chronic neurological diseases. It shows that PROK2 mediates neuronal cell deaths in traumatic brain injury via ferroptosis [72].

Table 3. Five downregulated DEGs only shared between *Sh3*[+/-]+LPS vs. WT+LPS and *Sh3*[+/-]+LPS vs. *Sh3*[+/-]+PBS, not include in WT+LPS vs. WT+PBS.

Gene name	Protein function
4930415F15Rik	SPMIP7 protein is mainly involved in spermatogenesis but is also expressed in the brain and its function is unclear.
AC173482.5	Predicted gene, Gm47797
Gm14393	Predicted gene
Gm5111	Predicted gene
Tgif2	TGIF2 is a transcriptional repressor. Downregulation of TGIF2 is correlated with neuronal apoptosis and development of autism in mice [73].

Discussion

In this study, we successfully established a preclinical model and experimental protocol to assess the role of neuroinflammation induced by LPS in *Sh3*[+/-] mice (**Fig. 6**). We demonstrated that LPS challenge modified the penetrance of motor function deficits, anxiety-like behavior, and repetitive behavior but did not alter social preference in *Sh3*[+/-] mice. More interestingly, the motor and emotion specific behavioral changes are reminiscent of catatonia features that are often observed in humans with *SHANK3* deficiency and idiopathic ASD [74-76]. Additionally, our extensive comparative transcriptome analysis among the different groups validated the experimental protocol of LPS induction and revealed distinct underlying molecular mechanisms. The significantly upregulated genes ubiquitously associated with inflammatory pathways in LPS treated groups indicated the effectiveness of LPS induction. The downregulated genes in multiple pathways associated with neuronal and synaptic functions strongly support a potential causal link between LPS treatment and increased penetrance of abnormal behaviors in *Sh3*[+/-] mice. Moreover, increased microglia activation and reduced synaptic function from LPS treatment were further validated by molecular experiments. Finally, the partial reversal of motor impairment in LPS treated *Sh3*[+/-] mice through the anti-inflammation treatment with MFA further supports this causality. Notably, MFA treatment did not restore LPS-treated WT and *Sh3*[+/-] mice performance in the accelerating speed rotarod. These findings suggest that the mechanisms underlying inflammation-induced motor impairment are different between steady speed and accelerating speed rotarod tests [77]. Additionally, one-week MFA treatment was more effective in female *Sh3*[+/-] mice than male mice (**Fig. 2D-E**) and LPS-treated male *Sh3*[+/-] mice performance was worse than female ones

(**Supplementary Fig. 1A-B**), indicating potential sex difference in motor function[78] and

neuroinflammation process[79].

The behavior impairments induced by LPS in *Sh3*[+/-] mice could attribute to the

dysfunction of neuronal cells resulted from neuroinflammation. In the previous study that

our lab contributed, LPS challenge increased the excitability in D1 but not D2 medium

spiny neurons using whole-cell patch clamp recordings and decreased neuron calcium

transients in nucleus accumbens in response to social stimuli using *in vivo* Ca^{2+} imaging[33].

Our data showed that increased neuroinflammation downregulated genes associated

with neuronal and synaptic functions in *Sh3*[+/-] mice compared to WT mice and reduced

synaptic protein PSD95 in *Sh3*[+/-] mice.

LPS-induced neuroinflammation could also alter the function of non-neuronal cells

in *Sh3*[+/-] mice. Our findings of augmented TLR4 expression and microglia activation

associated with LPS treatment suggested that microglia are primarily involved in more

significant neuroinflammatory responses in *Sh3*[+/-] mice. Consistent with the findings in

many reports in literature supporting that TLR4 mediates microglial activation in brains [51-

54], our findings also support that the upregulation of TLR4 contributes to the observed

microglial activation following LPS administration. The direct causality between

upregulation of TLR4 and microglia activation due to LPS induction has been

demonstrated in a recent study[80]. Our observations are consistent with this finding and

support the augmented TLR4 as the cause of microglia activation in SHANK3 deficiency

mice. Further analysis of TLR4 and IBA1 cellular colocalization indicated that additional

cell types might also be involved in LPS-TLR4 (ligand-receptor) mediated

neuroinflammation. Cellular fluorescence intensity of TLR4 in fixed brain cortex regions

decreased while total protein level of TLR4 in freshly prepared forebrain tissue increased

in $Sh3^{+/-}$ mice than WT. This inconsistence implied that changes of TLR4 in $Sh3^{+/-}$ might

be brain region specific and cell type specific, which warranted a future study on the TLR4

transduction signaling pathways in $Shank3$ haploinsufficiency mouse model.

Moreover, we uncovered a mechanistic convergence between LPS-induced

neuroimmune responses, and the synaptic disruptions known to arise from Shank3

deficiency. We discovered increased microglia engulfment and disrupted synaptic

structure in $Sh3^{+/-}$ mice. Such dysfunction sustained until 2 weeks later when we identified

LPS-triggered behavioral changes only in $Sh3^{+/-}$ mice instead of WT mice. Moreover, in a

previous study, we reported that the scaffold protein PSD95 expression did not change

significantly while HOMER1b/c was significantly reduced in $Sh3^{+/-}$ mice[35]. In the current

study, we replicated the finding that HOMER1b/c expression is significantly decreased in

$Sh3^{+/-}$ mice. However, LPS treatment did not further reduce HOMER levels. We also

replicated the finding that PSD95 is not changed in $Sh3^{+/-}$ mice. However, LPS treatment

significantly reduced PSD95 expression in $Sh3^{+/-}$ mice not WT mice. This pattern of

PSD95 changes in response to LPS, influenced by Shank3 haploinsufficiency, differs from

that observed for HOMER1b/c. Although we previously reported no change of NR2A or

GluA1 in $Sh3^{+/-}$ mice[35], we found that LPS treatment reduced NR2A and GluA1

expression in $Sh3^{+/-}$ mice instead of WT mice.

Our findings represent a first step toward understanding how Shank3

haploinsufficiency is linked to neuroimmune signaling. Our previous study showed that

SHANK3 deficiency in periphery nerve system contributes to the exaggerated

inflammatory responses[81]. Our recently published collaborative study has shown that

conditional knock-out of SHANK3 major isoform in brain endothelial cells recapitulated

social behavior impairment and other synaptic phenotypes [82]. Our findings are aligned

with previous studies showing that *Shank3* deficiency exacerbates the inflammatory

responses in peripheral neurons either induced by LPS [34,81] or by maternal immune

activation during the pregnancy [83]. Overall, our findings showed that LPS exacerbated

neuroinflammation in *Sh3*[+/-] mouse brains through unique molecular mechanisms as

summarized in **Fig. 6** for further investigations in future.

In   humans,   haploinsufficiency   of   *SHANK3*   is   highly   penetrant   for

neurodevelopmental impairments [84]. Approximately ~70% of PMS and *SHANK3*

deficiency patients met the diagnostic criteria for ASD. The comorbidity of intellectual

disability, anxiety, and epilepsy are highly penetrant or common [75]. There is an increasing

recognition of behavioral regression and catatonia among young teens with *SHANK3*

deficiency [74]. However, the triggers and mechanisms underlying the behavioral

regression are largely unknown. Although few clinical case reports implicated

neuroinflammation as a trigger [21,85], our findings may offer a mechanistic explanation for

the behavioral regressions observed in *SHANK3* patients after experiencing a subclinical

infection responded to the immune modulator treatment. Our study may support a link

between neuroinflammation and autism susceptibility in general.

Our results establish an important proof-of-concept study to dissect the link

between neuroinflammation and autism susceptibility. The apparent limitations of our

studies include that we have not provided definitive causality for the cell types and specific

molecular mechanism directly implicated in the increased neuroinflammatory responses

indued by LPS in SHANK3 haploinsufficiency mice. Future studies are warranted to

examine dosage- or treatment duration-dependent effects of LPS on behavioral outcomes

at different neurodevelopmental stages and in different sexes of *Shank3* deficiency mice.

Despite these limitations, our results suggest that *Shank3* haploinsufficiency increases

susceptibility to neuroinflammation-induced behavior impairment through distinct

molecular pathways. These findings provide new insights into the role of

neuroinflammation in ASD and for developing anti-inflammatory treatments as both

preventive and therapeutic strategies for behavior regression in ASD that are warranted

for further investigations.

Materials and methods

Animals

Animals were housed under standard 12 h/12 h light/dark cycles (7 AM-7 PM) with food, water ad libitum, controlled temperature and humidity in the Yale University animal facility. Experiments procedures conducted were following the Institutional Animal Care and Use Committee (IACUC) protocol at Yale University (protocol No. 2025-20271). $Shank3^{\Delta e4-22}$ mouse line (Jax Strain #:039524) was generated in-house and had been crossed with C57BL/6J mouse line (Jax Strain #:000664) over 20 generations. Wild-type (WT), $Shank3^{\Delta e4-22}$ homozygous ($Sh3^{-/-}$) and $Shank3^{\Delta e4-22}$ heterozygous mice ($Sh3^{+/-}$) were obtained through heterozygotes breeding as previously described [35]. All experiment procedures were performed within a time frame of 9 AM-5 PM. Animals of the same sex were group housed (2-5 mice per cage). Both male and female mice were used. One cage of mice with same genotype and same sex were assigned to PBS or LPS group randomly. Experimental order of different group was randomized. Mice from multiple groups for same experiments were carried out at same time. Before experiments, each mouse was cataloged with identification number in an excel sheet which included detailed information such as genotype and drug treatment. During experiments, experimenter handled mice based on its identification number and attempted to be semi-blinded. Post-processing, data handling and statistical analysis were semi-automated performed using software stated below. Mice with health problems such as runt, fight wounds, malocclusion, and eye problems, were excluded from experiments. For RNA and protein extraction, mice were deeply anesthetized with 5% isoflurane, then mouse brains were quickly collected, flash frozen in liquid nitrogen and stored in -80 °C freezer until further

use. The number of animals used in each experiment was determined based on our lab

prior experience and a prior power analysis was calculated using G*power statistics

software[86] (**Supplementary table 5**). Total number of mice used for each experiment was

as follows: 86 mice for MFA treatment rotarod test, 52-95 mice for open field, rotarod and

three-chamber behavior test, 49-66 mice for light/dark box and grooming behavior test,

15 mice for RNA sequencing, 40 mice for quantitative PCR, 24 mice for wide-field IBA1

staining, 6 mice for TLR4/IBA1 co-staining, 6 mice for microglia engulfment and

vGluT1/PSD95 staining at 24 hours post-injection, 12 mice for vGluT1/PSD95 staining at

24 hours post-injection, 18 mice for whole-cell protein analysis and 24 mice for synaptic

protein analysis. All experiments were replicated at least once, except that RNA

sequencing was done one time.

Drug

Three to five-week-old female and male mice were intraperitoneally injected with

LPS (Sigma, L6511) in saline at 1 mg/kg or same volume of saline (PBS). To test the

mefenamic acid (MFA) anti-inflammatory effect on behavior improvement, MFA

(Thermofisher, AAJ6270522) dissolved in DMSO at 5 mg/kg or the same volume of

DMSO (vehicle) was injected intraperitoneally to the mice with LPS injection once a day

for 7 days. Before injection, freshly made stock solution of MFA in DMSO (MFA treatment)

or DMSO (vehicle treatment) was diluted with 1:100 ratio in PBS to minimize DMSO

toxicity.

Behavior assay

For behavioral experiments, mice were handled for 3 days and habituated in the

procedure room for 1 hour before behavioral assays. All behavioral tests were conducted

once for each mouse.

Anxiety-like behaviors were tested by open field test (OFT) and light/dark box test

followed previous studies[35,87]. In OFT, mouse spontaneous activity in Noldus Open Field

(4 chambers, 40 (L) x 40 (W) x 30 (H) cm per chamber) over 10 minutes was recorded

with EthoVision XT video tracking software (Noldus). The Noldus two-chambered

light/dark box is equipped with the lid of dark chamber made by infrared translucent

material. The dimensions of the compartment are one third for the dark chamber (20 (L)

x 20 (W) x 20 (H) cm) and two thirds for the light chamber (20 (L) x 40 (W) x 20 (H) cm).

Mice were firstly placed into the open-light chamber (~700 lux) and given 5 min to freely

explore both light and dark (0 lux) chambers. The video was recorded with EthoVision XT

video tracking software. Ethovision software was used to automatically score the travel

distance and duration in arena.

Repetitive behavior was tested by evaluating mouse grooming behaviors. Mouse

was placed in a Noldus phenotyper (30 (L) x 30 (W) x 45 (H) cm) and recorded with

EthoVision XT video tracking software. Mouse freely explored for 10 minutes and last 5

minutes of grooming behavior was hand scored.

Motor coordination was tested by rotarod (Med Associate) performance as

described [35,88]. Mice were tested in two 5 min trials with an inter-trial interval of 30 min. In

accelerating session, revolutions per minute (rpm) of the turning barrel is accelerating

from 4 to 40. In steady speed session, the rpm of the turning barrel is fixed at 16. Latency

to fall, or to rotate off the top of the turning barrel, was measured by the timer.

Social preference was tested by a three-chamber assay as described[88]. An

unfamiliar age- and sex-matched mouse was used as a social stimuli mouse in the three-

chamber test and habituated under a wired enclosure for 3 days before testing. A

customized chamber composed of a rectangular Plexiglas arena (42 (L) x 56.5(W) x 35.8

(H) cm) divided into three chambers (each chamber 18.5 x 42 cm), with 2 doors

connecting chambers and two enclosures (wired pencil cups with metal vertical bars)

located in the two outer chambers. Briefly, the test mouse was habituated in the center

chamber for 5 min. Then the social stimuli mouse was placed under one of the pencil

cups and 2 doors were lifted allowing the experimental mouse to freely explore the whole

arena. The location of the empty and mouse enclosures, and the social stimuli mouse

were changed and counterbalanced for each trial. The EthoVision XT video tracking

software was used to score the duration of mouse spending in the empty chamber, the

mouse chamber, proximity regions (5 cm in distance) close to the empty enclosure, and

proximity regions close to the mouse enclosure.

Bulk RNA sequencing

Total RNA was isolated from the forebrain hemisphere using Trizol (Invitrogen,

15596018) followed manufacturer protocol and then treated with DNase I (Invitrogen,

AM1907) to remove contaminating DNA. RNA samples were sent to Yale Center for

Genome Analysis (YCGA) for bulk RNA sequencing (RNA-seq). See details about

sequencing process in Supplementary Information. After obtaining the data, low quality

reads were trimmed, and adaptor contamination were removed using Trim Galore (v0.5.0).

Trimmed reads were mapped to the mouse reference genome (mm10) using HISAT2

(v2.1.0) [89]. Gene expression levels were quantified using StringTie (v1.3.3b) [90].

Differentially expressed genes were identified using DESeq2 (v 1.22.1) [91]. Unpaired *t*-test

p value was used for filtering DEGs with significant differences between the two groups.

Log$_2$fold cutoff threshold at 1 and *p* < 0.05 was used to filter DEGs from two groups

comparison (WT+LPS vs. WT+PBS, *Sh3*$^{+/-}$+LPS vs. *Sh3*$^{+/-}$+PBS, *Sh3*$^{+/-}$+LPS vs.

WT+LPS and *Sh3*$^{+/-}$+PBS vs. WT+PBS) for genes comparison and analysis. *Pathway*

analysis: QIAGEN Ingenuity Pathway Analysis (IPA) library of canonical pathways was

used to identify the pathways most significant to the data set. Molecules from the dataset

meeting the criteria in which log$_2$fold cutoff threshold at 0.25 and *p* < 0.05 were associated

with a canonical pathway in the QIAGEN Knowledge Base were considered for the

analysis. Because using log$_2$fold cutoff threshold at 1 only revealed three activated

pathways in *Sh3*$^{+/-}$+LPS vs. WT+LPS comparison which were all inflammation-related, so

we expanded the log$_2$fold cutoff threshold to 0.25. The significance of the association

between the data set and the canonical pathway was measured based on: 1) a ratio of

the number of molecules from the data set that map to the pathway divided by the total

number of molecules that map to the canonical pathway was displayed, 2) a right-tailed

Fisher's Exact Test was used to calculate a p-value determining the probability that the

association between the genes in the dataset and the canonical pathway was explained

by chance alone, 3) A z-score was calculated to indicate the likelihood of activation

or inhibition of that pathway. In this paper, pathways with right-tailed Fisher's Exact Test

p < 0.05, z-score > 0.1 was considered as activated pathways and z-score < -0.1 was

considered as inhibited pathways. The volcano plot was created using VolcaNoseR [92].

Veen diagram was plotted using InteractiVenn [93].

Real-Time Quantitative Reverse Transcription PCR (RT-qPCR)

523   Total RNA was isolated using same protocol as RNA-seq. RNA was transcribed

524 into cDNA using iScript cDNA Synthesis Kit (Bio-Rad, 1708890) and qPCR was performed

525 in triplicate using iQ SYBR Green Supermix (Bio-Rad, 1708880). Gene expressions were

526 normalized to the expression of GAPDH. Primers used to amplify cDNAs are: *Il1b,*

527 forward, TGGACCTTCCAGGATGAGGACA, reverse, GTTCATCTCGGAGCCTGTAGTG;

528 *Cxcl10,* forward, ATCATCCCTGCGAGCCTATCCT, reverse, GACCTTTTTTGGCTAAA

529 CGCTTTC; *Cx3cr1,* forward, GAGCATCACTGACATCTACCTCC, reverse, AGAAGGCA-

530 GTCGTGAGCTTGCA; *P2ry12* forward, CATTGACCGCTACCTGAAGACC, reverse,

531 GCCTCCTGTTGGTGAGAATCATG; *Gapdh* forward, CAAAATGGTGAAGGTCGGTG,

532 reverse, AATGAAGGGGTCGTTGATGG.

533 **Immunohistochemistry**

534   Transcardiac perfusion was conducted at both 3 hours and 24 hours after LPS or

535 PBS injection with 20 mL PBS then 20 mL 4% paraformaldehyde (PFA, Santa Cruz, sc-

536 281692). Mouse brains were collected, fixed in 4% PFA for 24 hours, then changed to 30%

537 (w/v) sucrose before sectioning. 50 μM brain sections between Bregma 1 mm to -1 mm

538 were obtained using vibratome (Leica, VT1200S) or using cryostat (Leica, CM1860).

539 Primary antibody including anti-rabbit-IBA1 (1:200, FUJIFILM Wako Chemicals, 019-

540 19741), anti-mouse-TLR4 (1:100, Abcam, ab22048), anti-rat-CD68 (1:100, Abcam,

541 ab53444), anti-mouse-vGluT1 (1:100, Synaptic Systems, 135511), anti-guineapig-

542 vGluT1 (1:200, Synaptic Systems, 135318) and anti-mouse-PSD95 (1:100, Invitrogen,

543 MA1-045), and secondary antibody including goat anti-mouse Alexa Fluor 488 (Abcam,

544 ab150113), donkey anti-mouse Alexa Fluor 488 (Abcam, ab150105), donkey anti-rabbit

545 Alexa Fluor 594 (Invitrogen, A-21207), goat anti-rat Alexa Fluor 594 (Invitrogen, A-11007),

goat anti-guineapig Alexa Fluor 647 (Abcam, ab150187), goat anti-rabbit Alexa Fluor 647

(Jackson ImmunoResearch, 111-605-144) were used at 1:500 dilution. Zeiss LSM 980

confocal microscopy with a 20X objective lens was used for wide-field IBA1 imaging, a

63X oil objective lens with Z stack (0.2 μm thickness/slice, total 30-40 slices/image) was

used for engulfment imaging, and Airyscan 2 with a 63X oil objective lens was used for

super-resolution imaging of vGluT1 and PSD95 colocalization. Zeiss LSM 900 confocal

microscopy with a 63X oil objective lens was used for TLR4/IBA1 co-staining imaging. In

total, 49 images at 3 hours and 52 images at 24 hours post-injection were analyzed for

IBA1 fluorescence intensity, 20 images were analyzed for IBA1/TLR4 colocalization, 47

images were analyzed for microglia engulfment, and 26 images at 24 hours and 61

images at 2 weeks post-injection were analyzed for vGluT1/PSD95 colocalization. Images

were randomly selected from cortex regions of brain slices. Immunostaining and imaging

were performed under same conditions and same settings at same time across different

groups within same cohort. Cohort 1 with 8 groups was for IBA1 staining. Cohort 2 with 2

groups for TLR4/IBA1 co-staining. Cohort 3 with 2 groups was for IBA1/CD68/vGluT1 co-

staining. Cohort 4 with 2 groups was for 24 hours post-injection vGluT1/PSD95 co-

staining. Cohort 5 with 4 groups was for 2 weeks post-injection vGluT1/PSD95 co-staining.

Biological triplicates of brain slices from one mouse brain were employed. Results from

at least three mouse brains per group were analyzed and compared. ImageJ software

was used for IBA1 fluorescence intensity quantification. Imaris software (Oxford

Instrument) was used for microglia engulfment 3D rendering analysis. ZEN Blue software

was used for colocalization analysis. See details on 3D rendering and colocalization

analysis protocols in the Supplementary Information.

Western blot

Western blot was performed as previously described [94]. Whole cell lysates from mouse forebrain hemisphere were prepared using 1x lysis buffer (Cell Signaling Technology, 9803S) containing 1x protease/phosphatase inhibitor (Cell Signaling Technology, 5872S). Crude postsynaptic density (PSD) fraction from mouse forebrain hemisphere were prepared following previously described protocol with some modifications[35,95]. See details in Supplementary Information. Samples were boiled in 4x Laemlli buffer (Bio-Rad, 161-0747) at 98 °C for 5 min before gel electrophoresis. Home-made anti-rabbit SHANK3 C-terminus monoclonal antibody (1:1000), anti-rabbit TLR4 monoclonal antibody (1:1000, Santa Cruz, sc-293072), anti-mouse PSD95 monoclonal antibody (1:1000, Invitrogen, MA1-046), HOMER 1b/c monoclonal HRP-conjugated antibody (1:1000, Santa Cruz, sc-17842-HRP), anti-rabbit GluA1 (AMPA receptor) polyclonal antibody (1:1000, Abcam, ab31232), anti-NR2A (NMDA receptor) polyclonal antibody (1:1000, Millipore, ab31232), and β-ACTIN monoclonal HRP-conjugated antibody (1:1000, Invitrogen, MA5-15739-HRP) were used.

Statistical analysis

Graphpad Prism 10.0 (GraphPad Software) was used for the statistical analysis and data plot. Differences were analyzed by student's *t*-test if comparing two variables or using ANOVA if comparing more than two variables followed by multiple comparisons either using statistical hypothesis testing with correction or planned comparisons without correction. Significance was determined by two-tailed *t*-test *p* value or multiplicity adjusted *p* value. For data, which was not sampled from Gaussian distribution, a nonparametric test was performed. Since unequal variance observed in LPS-evoked cytokines and

chemokines, Brown-Forsythe and Welch ANOVA tests were performed (**Fig. 4A-B**). See

detailed statistical analysis for each figure in **Supplementary Table 5**. n represents the

total number of data points per group, usually meaning the mice number with an exception

that n represents number of images in immunohistochemistry results. Both sexes of mice

were used and balanced across genotype and treatment, with an exception that WT+PBS

and WT+LPS groups were all male mice in RNA-seq experiments. Behavioral difference

between male and female mice were explored in **Supplementary Fig. 1**. In box plot,

individual data points were plotted as: whiskers go down to the smallest value and up to

the largest value, box extends from the 25th to 75th percentiles and line in the middle of

the box is plotted at the median. In bar graph, individual data points were plotted and

presented as average ± SEM. *p < 0.05* was considered statistically significant. BioRender

was used to create schematic diagram.

Authorship contribution statement

S-N.Q. and Y-H.J. designed the experiments. S-N.Q. collected and analyzed the data. S.W. manually scored grooming behavior. S.W., K-Y.K. and S.J. performed western blot. S-N.Q. and Y-H.J. wrote and revised the manuscripts.

Acknowledgement

This work was supported by the National Institutes of Health (NIH) grant HD088007 MH104316, MH098114, MH117289, HD087795 (Y-H.J), the Autism Science Foundation (ASF) fellowship (S-N.Q), and the National Research Foundation of Korea (NRF) grant (RS-2024-00401971 (S.J), Brain Pool program RS-2024-00409006 (S.E.W)). This work was supported by the rodent behavior analysis facility and imaging core facility in Yale University funded by the Kavli Institute of Neuroscience. The bulk RNA sequencing was supported by the National Institute of General Medical Sciences of the National Institutes of Health under Award Number 1S10OD030363-01A1.

Conflict of Interest

There are no competing financial interests in relation to the work described.

Reference

1   Regier, D. A., Kuhl, E. A. & Kupfer, D. J. The DSM-5: Classification and criteria changes.

World Psychiatry **12**, 92-98 (2013). https://doi.org/10.1002/wps.20050

2   Stefanatos, G. A. Regression in autistic spectrum disorders. *Neuropsychol Rev* **18**,

305-319 (2008). https://doi.org/10.1007/s11065-008-9073-y

3   Manelis, L. *et al*. Language regression is associated with faster early motor

development in children with autism spectrum disorder. *Autism Res* **13**, 145-156

(2020). https://doi.org/10.1002/aur.2197

4   Boterberg, S., Van Coster, R. & Roeyers, H. Characteristics, Early Development and

Outcome of Parent-Reported Regression in Autism Spectrum Disorder. *J Autism Dev*

Disord **49**, 4603-4625 (2019). https://doi.org/10.1007/s10803-019-04183-x

5   Kohlenberg, T. M. *et al*. Psychiatric illness and regression in individuals with Phelan-

McDermid syndrome. *J Neurodev Disord* **12**, 7 (2020).

https://doi.org/10.1186/s11689-020-9309-6

6   Ghaziuddin, M. Catatonia: A Common Cause of Late Regression in Autism. *Front*

Psychiatry **12**, 674009 (2021). https://doi.org/10.3389/fpsyt.2021.674009

7   Betancur, C. & Buxbaum, J. D. SHANK3 haploinsufficiency: a "common" but

underdiagnosed highly penetrant monogenic cause of autism spectrum disorders.

Mol Autism **4**, 17 (2013). https://doi.org/10.1186/2040-2392-4-17

8   Satterstrom, F. K. *et al*. Large-Scale Exome Sequencing Study Implicates Both

Developmental and Functional Changes in the Neurobiology of Autism. *Cell* **180**,

568-584 e523 (2020). https://doi.org/10.1016/j.cell.2019.12.036

9   Zhou, X. *et al*. Integrating de novo and inherited variants in 42,607 autism cases

identifies   mutations   in   new   moderate-risk   genes.   *Nat   Genet*   (2022).

https://doi.org/10.1038/s41588-022-01148-2

10   Fu, J. M. *et al*. Rare coding variation provides insight into the genetic architecture and

phenotypic context of autism. *Nat Genet* (2022). https://doi.org/10.1038/s41588-

022-01104-0

11   Nevado, J. *et al*. Variability in Phelan-McDermid Syndrome in a Cohort of 210

Individuals.   *Front   Genet*   **13**,   652454   (2022).

https://doi.org/10.3389/fgene.2022.652454

12   Soorya, L. *et al*. Prospective investigation of autism and genotype-phenotype

correlations in 22q13 deletion syndrome and SHANK3 deficiency. *Mol Autism* **4**, 18

(2013). https://doi.org/10.1186/2040-2392-4-18

13   Leblond, C. S. *et al*. Meta-analysis of SHANK Mutations in Autism Spectrum Disorders:

a gradient of severity in cognitive impairments. *PLoS Genet* **10**, e1004580 (2014).

https://doi.org/10.1371/journal.pgen.1004580

14   Serrada-Tejeda, S., Martínez-Piédrola, R. M., Máximo-Bocanegra, N., Sánchez-

Herrera-Baeza, P. & Pérez-de-Heredia-Torres, M. Descriptive Analysis of Adaptive

Behavior in Phelan-McDermid Syndrome and Autism Spectrum Disorder. *Frontiers in*

neuroscience **16**, 893003 (2022). https://doi.org/10.3389/fnins.2022.893003

15   Reierson, G. *et al*. Characterizing regression in Phelan McDermid Syndrome (22q13

deletion   syndrome).   *J   Psychiatr   Res*   **91**,   139-144   (2017).

https://doi.org/10.1016/j.jpsychires.2017.03.010

16   Denayer, A. *et al*. Neuropsychopathology in 7 Patients with the 22q13 Deletion

Syndrome: Presence of Bipolar Disorder and Progressive Loss of Skills. *Molecular*

Syndromology **3**, 14-20 (2012). https://doi.org/10.1159/000339119

17   Zwanenburg, R. J., Ruiter, S. A. J., van den Heuvel, E. R., Flapper, B. C. T. & Van

Ravenswaaij-Arts, C. M. A. Developmental phenotype in Phelan-McDermid (22q13.3

deletion) syndrome: a systematic and prospective study in 34 children. *Journal of*

Neurodevelopmental Disorders **8**, 16 (2016). https://doi.org/10.1186/s11689-016-

9150-0

18   Serret, S. *et al*. Lithium as a rescue therapy for regression and catatonia features in

two SHANK3 patients with autism spectrum disorder: case reports. *BMC Psychiatry*

15, 107 (2015). https://doi.org/10.1186/s12888-015-0490-1

19   Phelan, K. & McDermid, H. E. The 22q13.3 Deletion Syndrome (Phelan-McDermid

Syndrome). *Mol Syndromol* **2**, 186-201 (2012). https://doi.org/10.1159/000334260

20   Phelan, K. *et al*. Phelan-McDermid syndrome: a classification system after 30 years

of experience. *Orphanet J Rare Dis* **17**, 27 (2022). https://doi.org/10.1186/s13023-

022-02180-5

21   Bey, A. L. *et al*. Subacute Neuropsychiatric Syndrome in Girls With SHANK3 Mutations

Responds to Immunomodulation. *Pediatrics* **145**, e20191490 (2020).

https://doi.org/10.1542/peds.2019-1490

22   Durand, C. M. *et al*. Mutations in the gene encoding the synaptic scaffolding protein

SHANK3 are associated with autism spectrum disorders. *Nat Genet* **39**, 25-27 (2007).

https://doi.org/10.1038/ng1933

23   He, G. *et al*. Dexmedetomidine attenuates neuroinflammation and microglia

activation in LPS-stimulated BV2 microglia cells through targeting circ-Shank3/miR-

140-3p/TLR4 axis. *Eur J Histochem* **67** (2023). https://doi.org/10.4081/ejh.2023.3766

24   Fischer, I. *et al*. Hyperbaric Oxygen Therapy Alleviates Social Behavior Dysfunction

and Neuroinflammation in a Mouse Model for Autism Spectrum Disorders. *Int J Mol*

Sci **23** (2022). https://doi.org/10.3390/ijms231911077

25   Cope, E. C. *et al*. Immature Neurons and Radial Glia, But Not Astrocytes or Microglia,

Are Altered in Adult Cntnap2 and Shank3 Mice, Models of Autism. *eNeuro* **3** (2016).

https://doi.org/10.1523/eneuro.0196-16.2016

26   Urrutia-Ruiz, C. *et al*. Deletion of the Autism-Associated Protein SHANK3 Abolishes

Structural Synaptic Plasticity after Brain Trauma. *Int J Mol Sci* **23** (2022).

https://doi.org/10.3390/ijms23116081

27   Eyo, U. & Molofsky, A. V. Defining microglial-synapse interactions. *Science* **381**, 1155-

1156 (2023). https://doi.org/doi:10.1126/science.adh7906

28   Pereira-Iglesias, M. *et al*. Microglia as hunters or gatherers of brain synapses. *Nature*

Neuroscience **28**, 15-23 (2025). https://doi.org/10.1038/s41593-024-01818-w

29   Suzuki, K. *et al*. Microglial activation in young adults with autism spectrum disorder.

JAMA Psychiatry **70**, 49-58 (2013). https://doi.org/10.1001/jamapsychiatry.2013.272

30   Kumar, A., Williams, M. T. & Chugani, H. T. Evaluation of basal ganglia and thalamic

inflammation in children with pediatric autoimmune neuropsychiatric disorders

associated with streptococcal infection and tourette syndrome: a positron emission

tomographic (PET) study using 11C-[R]-PK11195. *J Child Neurol* **30**, 749-756 (2015).

https://doi.org/10.1177/0883073814543303

31   Han, V. X., Patel, S., Jones, H. F. & Dale, R. C. Maternal immune activation and

neuroinflammation in human neurodevelopmental disorders. *Nature Reviews*

Neurology **17**, 564-579 (2021). https://doi.org/10.1038/s41582-021-00530-8

32   Michels, S., Mali, A., Jäntti, H., Rezaie, M. & Malm, T. Microglial involvement in autism

spectrum disorder: insights from human data and iPSC models. *Brain, Behavior, and*

Immunity **130**, 106071 (2025).

https://doi.org/https://doi.org/10.1016/j.bbi.2025.106071

33   Tzanoulinou, S. *et al.* Inhibition of Trpv4 rescues circuit and social deficits unmasked

by acute inflammatory response in a Shank3 mouse model of Autism. *Molecular*

Psychiatry **27**, 2080-2094 (2022). https://doi.org/10.1038/s41380-021-01427-0

34   Zhang, L. *et al.* SHANK3 in vagal sensory neurons regulates body temperature,

systemic inflammation, and sepsis. *Front Immunol* **14**, 1124356 (2023).

https://doi.org/10.3389/fimmu.2023.1124356

35   Wang, X. *et al.* Altered mGluR5-Homer scaffolds and corticostriatal connectivity in a

Shank3 complete knockout model of autism. *Nat Commun* **7**, 11459 (2016).

https://doi.org/10.1038/ncomms11459

36   Piirsalu, M. *et al.* Treatment With Lipopolysaccharide Induces Distinct Changes in

Metabolite Profile and Body Weight in 129Sv and Bl6 Mouse Strains. *Frontiers in*

Pharmacology **11** (2020). https://doi.org/10.3389/fphar.2020.00371

37 Jung, H. *et al*. LPS induces microglial activation and GABAergic synaptic deficits in

the hippocampus accompanied by prolonged cognitive impairment. *Scientific*

Reports **13**, 6547 (2023). https://doi.org/10.1038/s41598-023-32798-9

38 Decker Ramirez, E. B. *et al*. The effects of lipopolysaccharide exposure on social

interaction, cytokine expression, and alcohol consumption in male and female mice.

Physiology & Behavior **265**, 114159 (2023).

https://doi.org/https://doi.org/10.1016/j.physbeh.2023.114159

39 O'Connor, J. C. *et al*. Lipopolysaccharide-induced depressive-like behavior is

mediated by indoleamine 2,3-dioxygenase activation in mice. *Mol Psychiatry* **14**, 511-

522 (2009). https://doi.org/10.1038/sj.mp.4002148

40 Parrott, J. M., Oster, T. & Lee, H. Y. Altered inflammatory response in FMRP-deficient

microglia. *iScience* **24**, 103293 (2021). https://doi.org/10.1016/j.isci.2021.103293

41 Hodges, S. L. *et al*. Lipopolysaccharide-induced inflammation leads to acute

elevations in pro-inflammatory cytokine expression in a mouse model of Fragile X

syndrome. *Physiol Behav* **215**, 112776 (2020).

https://doi.org/10.1016/j.physbeh.2019.112776

42 Eliopoulos, A. G., Dumitru, C. D., Wang, C. C., Cho, J. & Tsichlis, P. N. Induction of

COX-2 by LPS in macrophages is regulated by Tpl2-dependent CREB activation

signals. *Embo j* **21**, 4831-4840 (2002). https://doi.org/10.1093/emboj/cdf478

43 Blais, V., Turrin, N. P. & Rivest, S. Cyclooxygenase 2 (COX-2) inhibition increases the

inflammatory response in the brain during systemic immune stimuli. *J Neurochem* **95**,

1563-1574 (2005). https://doi.org/10.1111/j.1471-4159.2005.03480.x

44   Cimolai, N. The potential and promise of mefenamic acid. *Expert Review of Clinical*

Pharmacology **6**, 289-305 (2013). https://doi.org/10.1586/ecp.13.15

45   Khansari, P. S. & Halliwell, R. F. Evidence for neuroprotection by the fenamate NSAID,

mefenamic   acid.   *Neurochemistry   International*   **55**,   683-688   (2009).

https://doi.org/https://doi.org/10.1016/j.neuint.2009.06.014

46   Lu, X. *et al.* Transcriptional determinism and stochasticity contribute to the

complexity of autism-associated SHANK family genes. *Cell Rep* **43**, 114376 (2024).

https://doi.org/10.1016/j.celrep.2024.114376

47   Krämer, A., Green, J., Pollard, J., Jr. & Tugendreich, S. Causal analysis approaches in

Ingenuity   Pathway   Analysis.   *Bioinformatics*   **30**,   523-530   (2014).

https://doi.org/10.1093/bioinformatics/btt703

48   Sousa, C. *et al.* Single-cell transcriptomics reveals distinct inflammation-induced

microglia   signatures.   *EMBO   Rep*   **19**   (2018).

https://doi.org/10.15252/embr.201846171

49   Norden, D. M., Trojanowski, P. J., Villanueva, E., Navarro, E. & Godbout, J. P.

Sequential activation of microglia and astrocyte cytokine expression precedes

increased Iba-1 or GFAP immunoreactivity following systemic immune challenge.

Glia **64**, 300-316 (2016). https://doi.org/10.1002/glia.22930

50   Tiwari, P. C. & Pal, R. The potential role of neuroinflammation and transcription

factors in Parkinson disease. *Dialogues Clin Neurosci* **19**, 71-80 (2017).

https://doi.org/10.31887/DCNS.2017.19.1/rpal

51   Cartarozzi, L. P., de Melo Lima, B. H., Tomiyama, A. L. R. M., Santos, F. R. & de Oliveira,

A. L. R. Toll-like receptors 2 and 4 influence neuronal survival and glial reactions

following ventral root crush injury in mice. *Journal of Neuroimmunology* **406**, 578655

(2025). https://doi.org/https://doi.org/10.1016/j.jneuroim.2025.578655

52   Yao, L. *et al*. Toll-like receptor 4 mediates microglial activation and production of

inflammatory mediators in neonatal rat brain following hypoxia: role of TLR4 in

hypoxic   microglia.   *Journal   of   Neuroinflammation*   **10**,   785   (2013).

https://doi.org/10.1186/1742-2094-10-23

53   Jiang, Y. *et al*. Evobrutinib mitigates neuroinflammation after ischemic stroke by

targeting M1 microglial polarization via the TLR4/Myd88/NF-κB pathway. *Mol Med* **31**,

148 (2025). https://doi.org/10.1186/s10020-025-01203-8

54   Zhu, J. *et al*. Toll-like receptor 4 deficiency in Purkinje neurons drives cerebellar ataxia

by impairing the BK channel-mediated after-hyperpolarization and cytosolic calcium

homeostasis. *Cell Death Dis* **15**, 594 (2024). https://doi.org/10.1038/s41419-024-

06988-w

55   *TLR4   expression   in   the   Human   Protein   Atlas*,

<https://www.proteinatlas.org/ENSG00000136869-TLR4/single+cell> (

56   Sjöstedt, E. *et al*. An atlas of the protein-coding genes in the human, pig, and mouse

brain. *Science* **367**, eaay5947 (2020). https://doi.org/10.1126/science.aay5947

57   Chen, X. *et al*. PSD-95 family MAGUKs are essential for anchoring AMPA and NMDA

receptor complexes at the postsynaptic density. *Proc Natl Acad Sci U S A* **112**, E6983-

6992 (2015). https://doi.org/10.1073/pnas.1517045112

58   Jaiswal, H. *et al*. The NF-κB regulator Bcl-3 restricts terminal differentiation and

promotes memory cell formation of CD8+ T cells during viral infection. *PLoS Pathog*

17, e1009249 (2021). https://doi.org/10.1371/journal.ppat.1009249

59   Pelisch, N., Rosas Almanza, J., Stehlik, K. E., Aperi, B. V. & Kroner, A. CCL3 contributes

to secondary damage after spinal cord injury. *J Neuroinflammation* **17**, 362 (2020).

https://doi.org/10.1186/s12974-020-02037-3

60   Klein, R. S. Regulation of neuroinflammation: the role of CXCL10 in lymphocyte

infiltration during autoimmune encephalomyelitis. *J Cell Biochem* **92**, 213-222 (2004).

https://doi.org/10.1002/jcb.20052

61   Zhong, K. *et al*. A Synthetic Steroid 5α-Androst-3β, 5, 6β-triol Alleviates Radiation-

Induced Brain Injury in Mice via Inhibiting GBP5/NF-κB/NLRP3 Signal Axis. *Mol*

Neurobiol **61**, 4074-4089 (2024). https://doi.org/10.1007/s12035-023-03831-9

62   Han, J. H. *et al*. IFITM6 expression is increased in macrophages of tumor-bearing mice.

Oncol Rep **25**, 531-536 (2011). https://doi.org/10.3892/or.2010.1092

63   Cunningham, C., Wilcockson, D. C., Campion, S., Lunnon, K. & Perry, V. H. Central

and systemic endotoxin challenges exacerbate the local inflammatory response and

increase neuronal death during chronic neurodegeneration. *J Neurosci* **25**, 9275-

9284 (2005). https://doi.org/10.1523/jneurosci.2614-05.2005

64   Zhao, J. *et al*. STEAP4 expression in CNS resident cells promotes Th17 cell-induced

autoimmune encephalomyelitis. *J Neuroinflammation* **18**, 98 (2021).

https://doi.org/10.1186/s12974-021-02146-7

65   Lester, S. N. & Li, K. Toll-like receptors in antiviral innate immunity. *J Mol Biol* **426**,

1246-1264 (2014). https://doi.org/10.1016/j.jmb.2013.11.024

66   Li, Y. *et al*. The multifaceted roles of activating transcription factor 3 (ATF3) in

inflammatory responses - Potential target to regulate neuroinflammation in acute

brain   injury.   *J   Cereb   Blood   Flow   Metab*   **43**,   8-17   (2023).

https://doi.org/10.1177/0271678x231171999

67   Ross, J. L. *et al*. Microglia and monocyte-derived macrophages drive progression of

pediatric high-grade gliomas and are transcriptionally shaped by histone mutations.

Immunity   **57**,   2669-2687.e2666   (2024).

https://doi.org/10.1016/j.immuni.2024.09.007

68   Krzyzowska, M., Kowalczyk, A., Skulska, K., Thörn, K. & Eriksson, K. Fas/FasL

Contributes to HSV-1 Brain Infection and Neuroinflammation. *Front Immunol* **12**,

714821 (2021). https://doi.org/10.3389/fimmu.2021.714821

69   Wu, S. *et al*. HSPA1A Protects Cells from Thermal Stress by Impeding ESCRT-0–

Mediated Autophagic Flux in Epidermal Thermoresistance. *Journal of Investigative*

Dermatology   **141**,   48-58.e43   (2021).

https://doi.org/https://doi.org/10.1016/j.jid.2020.05.105

70   Kampinga, H. H. *et al*. Guidelines for the nomenclature of the human heat shock

proteins.   *Cell   Stress   and   Chaperones*   **14**,   105-111   (2009).

https://doi.org/https://doi.org/10.1007/s12192-008-0068-7

71   Frankiw, L., Mann, M., Li, G., Joglekar, A. & Baltimore, D. Alternative splicing coupled

with transcript degradation modulates OAS1g antiviral activity. *Rna* **26**, 126-136

(2020). https://doi.org/10.1261/rna.073825.119

72   Bao, Z. *et al*. Prokineticin-2 prevents neuronal cell deaths in a model of traumatic

brain injury. *Nat Commun* **12**, 4220 (2021). https://doi.org/10.1038/s41467-021-

24469-y

73   Lei, J., Deng, Y. & Ma, S. Downregulation of TGIF2 is possibly correlated with neuronal

apoptosis and autism-like symptoms in mice. *Brain Behav* **12**, e2610 (2022).

https://doi.org/10.1002/brb3.2610

74   Dhossche, D. *et al*. Early-onset catatonia associated with SHANK3 mutations:

looking at the autism spectrum through the prism of psychomotor phenomena. *Front*

Psychiatry **14**, 1186555 (2023). https://doi.org/10.3389/fpsyt.2023.1186555

75   Srivastava, S. *et al*. Updated consensus guidelines on the management of Phelan-

McDermid syndrome. *Am J Med Genet A* **191**, 2015-2044 (2023).

https://doi.org/10.1002/ajmg.a.63312

76   Yurumez, E., Temelturk, R. D., Kaymak, B., Karabıyıkoğlu, Y. & Zengin, N. D. Catatonia

in autism spectrum disorders: Prevalence, functional impact, and diagnostic insights.

Psychiatry Res **342**, 116250 (2024). https://doi.org/10.1016/j.psychres.2024.116250

77   Monville, C., Torres, E. M. & Dunnett, S. B. Comparison of incremental and

accelerating protocols of the rotarod test for the assessment of motor deficits in the

6-OHDA model. *J Neurosci Methods* **158**, 219-223 (2006).

https://doi.org/10.1016/j.jneumeth.2006.06.001

78   Hernandez, A. R., Truckenbrod, L. M., Campos, K. T., Williams, S. A. & Burke, S. N. Sex

differences in age-related impairments vary across cognitive and physical

assessments in rats. *Behav Neurosci* **134**, 69-81 (2020).

https://doi.org/10.1037/bne0000352

79   Han, J., Fan, Y., Zhou, K., Blomgren, K. & Harris, R. A. Uncovering sex differences of

rodent microglia. *Journal of Neuroinflammation* **18**, 74 (2021).

https://doi.org/10.1186/s12974-021-02124-z

80   Hines, D. J., Choi, H. B., Hines, R. M., Phillips, A. G. & MacVicar, B. A. Prevention of

LPS-induced microglia activation, cytokine production and sickness behavior with

TLR4 receptor interfering peptides. *PLoS One* **8**, e60388 (2013).

https://doi.org/10.1371/journal.pone.0060388

81   Han, Q. *et al*. SHANK3 Deficiency Impairs Heat Hyperalgesia and TRPV1 Signaling in

Primary Sensory Neurons. *Neuron* **92**, 1279-1293 (2016).

https://doi.org/10.1016/j.neuron.2016.11.007

82   Kim, Y. E. *et al*. Endothelial SHANK3 regulates tight junctions in the neonatal mouse

blood-brain barrier through β-Catenin signaling. *Nat Commun* **16**, 1407 (2025).

https://doi.org/10.1038/s41467-025-56720-1

83   Atanasova, E. *et al*. Immune activation during pregnancy exacerbates ASD-related

alterations in Shank3-deficient mice. *Mol Autism* **14**, 1 (2023).

https://doi.org/10.1186/s13229-022-00532-3

84 Leblond, C. S. *et al*. Meta-analysis of SHANK Mutations in Autism Spectrum Disorders:

A Gradient of Severity in Cognitive Impairments. *PLOS Genetics* **10**, e1004580 (2014).

https://doi.org/10.1371/journal.pgen.1004580

85 Jesse, S. *et al*. Phelan McDermid Syndrome: Multiple Sclerosis as a Rare but Treatable

Cause for Regression-A Case Report. *Int J Mol Sci* **22** (2021).

https://doi.org/10.3390/ijms22052311

86 Faul, F., Erdfelder, E., Lang, A. G. & Buchner, A. G*Power 3: a flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behav Res*

Methods **39**, 175-191 (2007). https://doi.org/10.3758/bf03193146

87 Bourin, M. & Hascoët, M. The mouse light/dark box test. *European Journal of*

Pharmacology **463**, 55-65 (2003). https://doi.org/https://doi.org/10.1016/S0014-

2999(03)01274-3

88 Moy, S. S. *et al*. Mouse behavioral tasks relevant to autism: phenotypes of 10 inbred

strains. *Behav Brain Res* **176**, 4-20 (2007). https://doi.org/10.1016/j.bbr.2006.07.030

89 Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**,

907-915 (2019). https://doi.org/10.1038/s41587-019-0201-4

90 Pertea, M. *et al*. StringTie enables improved reconstruction of a transcriptome from

RNA-seq reads. *Nature Biotechnology* **33**, 290-295 (2015).

https://doi.org/10.1038/nbt.3122

91   Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and

dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

https://doi.org/10.1186/s13059-014-0550-8

92   Goedhart, J. & Luijsterburg, M. S. VolcaNoseR is a web app for creating, exploring,

labeling and sharing volcano plots. *Scientific Reports* **10**, 20560 (2020).

https://doi.org/10.1038/s41598-020-76603-3

93   Heberle, H., Meirelles, G. V., da Silva, F. R., Telles, G. P. & Minghim, R. InteractiVenn: a

web-based tool for the analysis of sets through Venn diagrams. *BMC Bioinformatics*

16, 169 (2015). https://doi.org/10.1186/s12859-015-0611-3

94   Wang, S. E. *et al*. TRPV1 Regulates Stress Responses through HDAC2. *Cell Rep* **19**,

401-412 (2017). https://doi.org/10.1016/j.celrep.2017.03.050

95   Carlin, R. K., Grab, D. J., Cohen, R. S. & Siekevitz, P. Isolation and characterization of

postsynaptic densities from various brain regions: enrichment of different types of

postsynaptic densities. *J Cell Biol* **86**, 831-845 (1980).

https://doi.org/10.1083/jcb.86.3.831

Figure legend:

Figure 1. LPS challenge induced selective behavioral impairments in $Sh3^{+/-}$ mice.

A. The body weight loss after LPS intraperitoneal injection. Mouse was weighed

before injection, and at 3, 6, 24, 48, and 72 hours after LPS (1 mg/kg) or the same volume

of PBS injection. Each mouse's body weight was normalized to the weight before injection.

Two-way repeated measures ANOVA was performed (genotype/LPS effect $F_{(3, 50)}$ = 34.32,

$p < 0.0001$; Hour effect $F_{(4, 200)}$ = 84.15, $p < 0.0001$; genotype/LPS X Hour interaction F

$_{(12, 200)}$ = 22.64, $p < 0.0001$). Post-hoc Tuckey's test showed no significant difference

between WT+PBS (n = 8) and $Sh3^{+/-}$+PBS (n = 17) or between WT+LPS (n = 8) and

$Sh3^{+/-}$+LPS (n = 21) at 3 hours, 6 hours, 24 hours, 48 hours and 72 hours after injection.

Mouse body weight started to reduce significantly after LPS injection compared to PBS

injection at 24 hours, 48 hours and 72 hours. Mouse body weight from LPS groups

significantly increased at 72 hours compared to 48 hours after injection in both WT+LPS

and $Sh3^{+/-}$+LPS groups. See multiple comparison statistics in Supplementary Table 5.

B. Schematic drawing for the experimental design.

C-D. LPS challenge impaired motor function in $Sh3^{+/-}$ mice. Steady speed rotarod (16

rpm, **C**) and accelerated speed rotarod (4-40 rpm, **D**) tests were performed and latency

to fall was measured. No significant difference was found between WT+PBS and $Sh3^{+/-}$

+PBS (16 rpm, p =0.5487, 4-40 rpm, p =0.3283) or between WT+PBS and WT+LPS (16

rpm, p =0.9797, 4-40 rpm, p =0.6711). $Sh3^{+/-}$+LPS significantly reduced latency to fall

compared to $Sh3^{+/-}$+PBS (16 rpm, p =0.0126, 4-40 rpm, p =0.0061) or compared to

WT+LPS (16 rpm, p =0.002, 4-40 rpm, p =0.001). $Sh3^{-/-}$ mice significantly reduced latency

to fall compared to WT+PBS (16 rpm, p < 0.0001, 4-40 rpm, p < 0.0001).

E-F. LPS challenge increased anxiety-like behavior in $Sh3^{+/-}$ mice. Mouse duration

spent in the center of the open field test and travel distance in light chamber of light/dark

box test were measured. No significant difference found between WT+PBS and WT+LPS

(OFT, p = 0.5956, light/dark box p = 0.6004). $Sh3^{+/-}$+LPS showed less center duration

than $Sh3^{+/-}$+PBS (p = 0.0033) despite that $Sh3^{+/-}$+PBS showed increased center duration

than WT+PBS (p = 0.0005), and $Sh3^{+/-}$+LPS showed less travel distance in light chamber

than $Sh3^{+/-}$+PBS (p = 0.0158) or WT+LPS (p = 0.0107).

G. LPS challenge increased repetitive behavior in $Sh3^{+/-}$ mice. Grooming behavior

was hand scored and compared. $Sh3^{+/-}$+LPS showed longer grooming duration than

$Sh3^{+/-}$+PBS (p = 0.0348) or WT+LPS (p = 0.0135).

H-I. LPS challenge had no effect on sociability $Sh3^{+/-}$ mice. Three-chamber tests were

conducted and duration spent in mouse chamber (S), empty chamber (E), proximity

regions in mouse chamber (SP) and empty chamber (EP) were compared (Two-way

repeated measures ANOVA. **H.** genotype effect F $_{(4, 68)}$ = 1.590, p = 0.1869, S/E effect F

$_{(1, 68)}$ = 25.46, p < 0.0001, genotype X S/E interaction F $_{(4, 68)}$ = 0.8135, p = 0.5209; **I.**

genotype effect F $_{(4, 68)}$ = 1.398, p = 0.2441, SP/EP effect F $_{(1, 68)}$ = 34, p < 0.0001, genotype

X SP/EP interaction F $_{(4, 68)}$ = 0.4236, p = 0.7911). No significant difference of duration

was found between S vs. E (p = 0.5325) or SP vs. EP (p = 0.2529) in $Sh3^{-/-}$ mice. All other

groups of mice showed significantly increased duration in S vs. E or SP vs. EP. * p < 0.05,

** p < 0.005, *** p < 0.0005, **** p < 0.0001

Figure 2. Mefenamic acid (MFA) partially improved the motor behavior deterioration in WT and *Sh3*$^{+/-}$ mice.

A. Schematic drawing for experiment design. Anti-inflammatory treatment with intraperitoneal MFA injection was given to mice once every day for 7 days after LPS injection. Rotarod tests were performed 24 hours after the last does of MFA injection.

B-C. MFA partially improved the rotarod test performance. After 1-week treatment of MFA, Steady speed rotarod (16 rpm, **B**) and accelerated speed rotarod (4-40 rpm, **C**) tests were performed and latency to fall was compared. (Two-way ANOVA, **B. 16 rpm,** LPS/MFA effect $F_{(1, 37)}$ = 23.66, $p < 0.0001$, genotype effect $F_{(1,46)}$ = 1.859, $p = 0.1794$, LPS/MFA X genotype interaction $F_{(1, 37)}$ = 0.8697, $p = 0.3571$. **C. 4-40 rpm,** LPS/MFA effect $F_{(1, 85)}$ = 1.721, $p = 0.1932$, genotype effect $F_{(1,85)}$ = 4.822, $p = 0.0308$, LPS/MFA X genotype interaction $F_{(1, 85)}$ = 0.8220, $p = 0.3672$). Compared to vehicle group, MFA-treated *Sh3*$^{+/-}$ and WT mice showed increased latency to fall in 16 rpm rotarod test (*Sh3*$^{+/-}$, $p = 0.0091$; WT, $p = 0.0002$).

D-E. MFA treatment was more effective in female *Sh3*$^{+/-}$ mice than male ones. Two-way ANOVA was performed to compare sex difference in MFA treatment (**D. 16 rpm,** LPS/MFA effect $F_{(1, 43)}$ = 4.626, $p = 0.0372$, sex effect $F_{(1,43)}$ = 9.227, $p = 0.004$, LPS/MFA X sex interaction $F_{(1, 43)}$ = 1.858, $p = 0.1799$. **E. 4-40 rpm,** LPS/MFA effect $F_{(1, 16)}$ = 2.065, $p = 0.17$, sex effect $F_{(1,29)}$ = 2.369, $p = 0.1346$, LPS/MFA X sex interaction $F_{(1, 16)}$ = 8.046, $p = 0.0119$). MFA-treated female *Sh3*$^{+/-}$ showed significantly higher latency to fall compared to vehicle-treated female *Sh3*$^{+/-}$ (16 rpm, $p = 0.0229$; 40rpm, $p = 0.0106$) and MFA-treated male *Sh3*$^{+/-}$ (16 rpm, $p = 0.0009$; 40rpm, $p = 0.0011$). * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$, **** $p < 0.0001$

Figure 3. Bulk RNA sequencing revealed distinct molecular mechanism underlying

LPS effect on *Sh3*$^{+/-}$ mice.

A-D. The volcano plot shows differentially expressed genes (DEGs) distribution.

DEGs were identified in four comparisons: *Sh3*$^{+/-}$+LPS vs *Sh3*$^{+/-}$+PBS (**A**), WT+LPS vs.

WT+PBS (**B**), *Sh3*$^{+/-}$+LPS vs. WT+LPS (**C**), *Sh3*$^{+/-}$+PBS vs. WT+PBS (**D**). Upregulated

genes are indicated in red color, downregulated genes are indicated in blue color and

DEGs below threshold are indicated in grey color.

E-F. Venn diagrams show shared upregulated and downregulated genes among

four comparisons. Shared upregulated (**E**) and downregulated genes (**F**) numbers are

shown within the diagrams. Comparison of *Sh3*$^{+/-}$+LPS vs *Sh3*$^{+/-}$+PBS is red diagram,

WT+LPS vs. WT+PBS is blue diagram, *Sh3*$^{+/-}$+LPS vs. WT+LPS is green diagram, and

Sh3$^{+/-}$+PBS vs. WT+PBS is yellow diagram.

G. Bubble chart shows functions of DEGs of *Sh3*$^{+/-}$+LPS vs. WT+LPS comparison.

Volcano plot of the negative log of p-value (Y-axis) vs. the z-score (X-axis). *p*-value was

calculated using the right-tailed Fisher's Exact Test. A negative log of p-value cutoff of

1.3 (*p* < 0.05) was used to identify significantly changed functional pathways. A z-score

range cutoff of ±0.1 was used to identify the activation or inhibition state of a canonical

pathway. Some names of pathways are annotated in the chart. Activated pathways are

orange bubbles and inhibited pathways are blue bubbles. The size of the bubbles reflects

the number of genes overlapped with the pathway database.

Figure 4. Neuroinflammation increased in *Sh3*$^{+/-}$ mice compared to WT mice in response to LPS.

A-D. Differential expression of neuroinflammatory marker genes *Il1b, Cxcl10, Cx3cr1* and *P2ry12*. At 24 hours after LPS injection, forebrain RNA was extracted, and RT-qPCR was performed. Target gene expression was normalized to the expression of the housekeeping gene GAPDH. One-way ANOVA was performed to compare gene expression in four groups (*Il1b*: F= 14.93, *p* = 0.0004, *Cxcl10*: F= 16.73, *p* = 0.0002, *Cx3cr1*: F= 14.8, *p* < 0.0001, *P2ry12*: F= 8.442, *p* = 0.0002). As expected, compared to PBS treatment, LPS significantly increased expression of *Il1b, Cxcl10*, and *Cx3cr1*, and decreased expression of *P2ry12* in both WT and *Sh3*$^{+/-}$ mice (*Il1b*: WT+PBS (n=10) vs. WT+LPS (n=9), *p* = 0.0005, *Cxcl10*: WT+PBS (n=10) vs. WT+LPS (n=6), *p* = 0.0105, *Cx3cr1*: WT+PBS (n=10) vs. WT+LPS (n=9), *p* = 0.0388, *P2ry12*: WT+PBS (n=10) vs. WT+LPS (n=8), *p* = 0.0241; *Il1b*: *Sh3*$^{+/-}$ +PBS (n=12) vs. *Sh3*$^{+/-}$ +LPS (n=11), *p* = 0.002, *Cxcl10*: *Sh3*$^{+/-}$ +PBS (n=12) vs. *Sh3*$^{+/-}$ +LPS (n=11), *p* = 0.0015, *Cx3cr1*: *Sh3*$^{+/-}$ +PBS (n=12) vs. *Sh3*$^{+/-}$ +LPS (n=12), *p* < 0.0001, *P2ry12*: *Sh3*$^{+/-}$ +PBS (n=11) vs. *Sh3*$^{+/-}$ +LPS (n=11), *p* = 0.0046). Significant increase of *Il1b, Cxcl10, Cx3cr1* was observed in *Sh3*$^{+/-}$ +LPS vs. WT+LPS (*Il1b*: *p* = 0.0166, *Cxcl10*: *p* = 0.0076, *Cx3cr1*: *p* = 0.0004). In contrast, significantly increased expression of *P2ry12* was observed in *Sh3*$^{+/-}$+PBS vs. WT+PBS (*p* = 0.0102) and *Sh3*$^{+/-}$+LPS vs. WT+LPS (*p* = 0.036).

E-F. Increased immunohistochemistry fluorescence intensity of IBA1 in *Sh3*$^{+/-}$ +LPS group. Mouse brain slices were collected at 3 hours and 24 hours after injection, respectively. Microglia activation was labeled with IBA1 antibody (**E1-E4, F1-F4**). Quantitative fluorescence intensity of IBA1 at cortex regions from WT+LPS, *Sh3*$^{+/-}$+PBS,

and $Sh3^{+/-}$+LPS groups were normalized to WT+PBS group and then compared (Kruskal-

Wallis test, **E5.** p = 0.0002; **F5.** p < 0.0001). LPS increased microglia activation in both

WT and $Sh3^{+/-}$ mice at both 3 hours and 24 hours after injection compared to PBS groups

(3 hours: WT+PBS (n=10) vs. WT+LPS (n=13), p = 0.0067, $Sh3^{+/-}$ +PBS (n=12) vs. $Sh3^{+/-}$

+LPS (n=11), p = 0.0004. 24 hours: WT+PBS (n=9) vs. WT+LPS (n=16), p = 0.0053,

$Sh3^{+/-}$ +PBS (n=15) vs. $Sh3^{+/-}$ +LPS (n=12), p < 0.0001). Moreover, LPS induced

significant overactivation of microglia in $Sh3^{+/-}$ mice at 24 hours after injection compared

to WT mice, not 3 hours (WT+LPS vs. $Sh3^{+/-}$ +LPS, 3 hours, p = 0.6204, 24 hours, p =

0.0017).

G-I. Increased TLR4 and decreased SHANK3 protein expression in S$h3^{+/-}$ mice. LPS

receptor TLR4 and SHANK3 expression were examined in whole-cell lysates using

western blot at 24 hours after injection. Target protein expression was normalized to β-

ACTIN expression (**G**). Quantitative protein expression from WT+LPS, $Sh3^{+/-}$+PBS, and

$Sh3^{+/-}$+LPS were normalized to WT+PBS group and one-way ANOVA was performed

(TLR4, F = 14.38, p = 0.0001, SHANK3, F = 7.681, p = 0.0033). Significantly elevated

TLR4 expression was observed in $Sh3^{+/-}$ mice compared to WT mice (**H**, WT+PBS vs.

$Sh3^{+/-}$ +PBS, p < 0.0001; WT+LPS vs. $Sh3^{+/-}$ +LPS, p = 0.0081). As expected, significantly

reduced SHANK3 expression was detected in $Sh3^{+/-}$ mice (**I**, WT+PBS vs. $Sh3^{+/-}$ +PBS,

p = 0.0172; WT+LPS vs. $Sh3^{+/-}$ +LPS, p = 0.002).

J-P. Colocalization analysis of TLR4/IBA1 between WT+LPS vs. $Sh3^{+/-}$ +LPS group.

Mouse brain slices were collected at 24 hours after LPS injection in WT and $Sh3^{+/-}$ mice.

IBA1 and TLR4 antibody were used for immunocytochemistry. Colocalization of IBA1 and

TLR4 at cortex regions were analyzed and compared (**J-K**). Unpaired t-test was

conducted. No difference in colocalization Pearson's coefficient between WT+LPS vs.

$Sh3^{+/-}$ +LPS was observed (**L**, two-tailed p = 0.2309). $Sh3^{+/-}$ showed smaller normalized

colocalization by TLR4 (**M**, two-tailed p = 0.0093) but similar normalized colocalization by

IBA1 (**N**, two-tailed p = 0.9141) compared to WT. $Sh3^{+/-}$ showed decreased TLR4 cellular

fluorescence intensity (**O**, two-tailed p = 0.0183) but increased IBA1 cellular fluorescence

intensity (**P**, two-tailed p = 0.0001) than WT. * p < 0.05, ** p < 0.005, *** p < 0.0005, **** p <

0.0001.

Figure 5. Microglia activation disrupted synaptic structure in *Sh3*+/- mice compared to WT mice in response to LPS.

A-D. Increased vGluT1 engulfment in *Sh3*+/- mice at 24 hours after LPS injection. A. Representative images of maximum projection of z-stack imaging of IBA1/CD68/vGluT1 co-staining (left), 3D rendering masked by IBA1 and CD68 (center), and enlarged images of 3D rendering (right). Total volumes of masked vGluT1 and CD68 were calculated and unpaired t-test was conducted. No significant difference of IBA1 masked CD68 occupancy was detected (**B.** two-tailed p = 0.7149). IBA1 and CD68 double masked vGluT1 occupancy and IBA1 masked vGluT1 occupancy was higher in *Sh3*+/- mice than WT mice (**C.** two-tailed p < 0.0001, **D.** two-tailed p = 0.0003).

E-H. Reduced synapses in *Sh3*+/- mice at 24 hours after LPS injection. E. Representative images of super-resolution imaging of vGluT1/PSD95 co-staining at 24 hours post-injection. Colocalized puncta of vGluT1and PSD95 were indicated by arrowheads and were counted as synapse puncta. Non-colocalized puncta of PSD95 were indicated by arrows. Colocalization of vGluT1and PSD95 was analyzed, and number of colocalized puncta were counted as synapse density. Unpaired t-test was conducted. *Sh3*+/- mice showed a lower colocalization Pearson's coefficient (**F**, two-tailed p = 0.0002), a reduced normalized colocalization coefficient by PSD95 (**G**, two-tailed p = 0.0056) and decreased synapse density (**H**, two-tailed p = 0.0107) compared to WT mice.

I-M. Reduced synapses in *Sh3*+/- mice at 2 weeks after LPS injection. I-J. Representative images of super-resolution imaging of vGluT1/PSD95 co-staining at 2 weeks post-injection among four groups, WT+PBS, WT+LPS, *Sh3*+/-+PBS, and *Sh3*+/-+LPS. Mask was generated based on the colocalized puncta of vGluT1and PSD95. Non-

colocalized puncta of PSD95 were indicated by arrows in **J**. One-way ANOVA was

conducted for comparing colocalization Pearson's coefficient (**K.** F = 4.254, *p* = 0.0088),

colocalization coefficient by PSD95 (**L.** F = 26.51, *p* < 0.0001), and synapse density (**M.**

F = 10.94, *p* < 0.0001). *Sh3*$^{+/-}$+LPS group showed a lower colocalization Pearson's

coefficient than WT+LPS (*p* = 0.0067) or *Sh3*$^{+/-}$+PBS groups (*p* = 0.0022), a reduced

normalized colocalization coefficient by PSD95 than WT+LPS (*p* < 0.0001) or *Sh3*$^{+/-}$+PBS

groups (*p* < 0.0001) , and decreased synapse density than WT+LPS (*p* < 0.0001) or *Sh3*$^{+/-}$

+PBS groups (*p* < 0.0001).

N-S. Reduced PSD95, HOMER 1b/c, NR2A and GluA1 expression in *Sh3*$^{+/-}$ mice at 2

weeks after LPS injection. Postsynaptic scaffold proteins PSD95 and HOMER 1b/c (**N**),

and excitatory receptors NMDA subunit NR2A and AMPA subunit GluA1 expression (**Q**)

were examined in forebrain PSD fractions using western blot at 2 weeks after injection.

Target protein expression was normalized to β-ACTIN expression. Quantitative protein

expression from WT+LPS, *Sh3*$^{+/-}$+PBS, and *Sh3*$^{+/-}$+LPS were normalized to WT+PBS

group and one-way ANOVA was conducted (PSD95, F = 4.763, *p* = 0.006, HOMER 1b/c,

F = 12.28, *p* < 0.0001, NR2A, F = 5.06, *p* = 0.0091, GluA1, F = 4.951, *p* = 0.0313). **O.**

Significantly decreased PSD95 expression was detected in *Sh3*$^{+/-}$ mice not WT mice after

LPS treatment (*Sh3*$^{+/-}$+LPS (n=11) vs. WT+LPS (n=12), *p* = 0.0009; *Sh3*$^{+/-}$+LPS (n=11)

vs. *Sh3*$^{+/-}$+PBS (n=11), *p* = 0.0354). No significant change in PSD95 was detected

between WT+PBS (n=12) vs. WT+LPS (*p* = 0.5153) or WT+PBS vs. *Sh3*$^{+/-}$+PBS (*p* =

0.4845). **P.** Significantly reduced HOMER 1b/c expression was observed in *Sh3*$^{+/-}$ mice

compared to WT mice (WT+PBS (n=6) vs. *Sh3*$^{+/-}$ +PBS (n=6), *p* < 0.0001; WT+LPS (n=6)

vs. *Sh3*$^{+/-}$+LPS (n=6), *p* = 0.0088). No significant change of HOMER 1b/c was detected

after LPS treatment (WT+PBS vs. WT+LPS (p = 0.1781) or $Sh3^{+/-}$+PBS vs. $Sh3^{+/-}$+LPS

(p = 0.318)). **R.** $Sh3^{+/-}$+LPS group (n=6) showed significant decreased NR2A expression

compared to both the $Sh3^{+/-}$+PBS group (n=6) (p = 0.0053) and the WT+LPS group (n=6)

(p = 0.002). No significant change of NR2A was observed between WT+PBS (n=6) vs.

WT+LPS (p = 0.1404) or WT+PBS vs. $Sh3^{+/-}$+PBS (p = 0.2827). **S.** $Sh3^{+/-}$+LPS group

(n=3) showed significantly reduced GluA1 expression compared to both the $Sh3^{+/-}$ +PBS

group (n=3) (p = 0.0378) and the WT+LPS group (n=3) (p = 0.0169). No significant

change in GluA1 was found between WT+PBS (n=3) vs. WT+LPS (p = 0.5898) or

WT+PBS vs. $Sh3^{+/-}$+PBS (p = 0.3102). * p < 0.05, ** p < 0.005, *** p < 0.0005, **** p < 0.0001.

1105

Figure 6. Summary of increased neuroinflammation susceptibility in *Sh3*$^{+/-}$ mice.

The schematic drawing proposes a model of increased neuroinflammation and altered

behavioral phenotypes in *Shank3* haploinsufficiency mouse after LPS treatment.

Generally speaking, LPS evokes neuroinflammation through three distinct pathways, 1)

LPS binds to TLR4 expressed on neutrophils, promoting neutrophil infiltration through

blood-brain barrier and activation of microglia cells; 2) LPS binds to TLR4 expressed in

cerebral endothelial cells located in blood-brain barrier, compromising the integrity of

blood-brain barrier and increasing permeability of additional immune components

entering the brain; 3) LPS that penetrates blood-brain barrier binds to TLR4 expressed in

microglia, activating microglia and triggering proinflammatory cytokines release. In *Sh3*$^{+/-}$

brain vs. WT brain, we observed excessive proinflammatory gene expression,

proinflammatory cytokines, overactivation of microglia and increased expression of TLR4

at 24 hours after LPS injection. It suggests that LPS might exacerbate neuroinflammation

responses through above routes in *Sh3*$^{+/-}$ mice.

Microglia activation can sculpt synapses and brain circuits and change animal behavioral

phenotypes long-term. In *Sh3*$^{+/-}$ brain vs. WT brain, we observed elevated microglia

engulfment, reduced synapse, and decreased synaptic transmission gene expression at

24 hours after LPS injection, suggesting increased synapse elimination in *Sh3*$^{+/-}$ mice. At

2 weeks after LPS injection, we observed reduced synapse density, decreased PSD95

expression and lower expression of excitatory receptor expression. It indicates that

neuroinflammation-induced synaptic dysfunction was associated with behavioral changes

found in *Sh3*$^{+/-}$ mice.

Figure 1

A — Normalized body weight across time after injection (3 hour, 6 hour, 24 hour, 48 hour, 72 hour) for WT+PBS, WT+LPS, $Sh3^{+/-}$+PBS, $Sh3^{+/-}$+LPS.

B — Experimental timeline: LPS injection Day 0; Rotarod Day 15; Open field Day 16; Sociability (Social target, Empty cage) Day 17; Light/dark box; Grooming.

Rotarod

C — 16 rpm Latency to fall (s): WT+PBS (n=15), WT+LPS (n=21), $Sh3^{+/-}$+PBS (n=23), $Sh3^{+/-}$+LPS (n=23), $Sh3^{-/-}$ (n=12). **** ** *

D — 4-40 rpm Latency to fall (s): WT+PBS (n=15), WT+LPS (n=21), $Sh3^{+/-}$+PBS (n=23), $Sh3^{+/-}$+LPS (n=23), $Sh3^{-/-}$ (n=12). **** ** **

Open field

E — Center duration (s): WT+PBS (n=11), WT+LPS (n=14), $Sh3^{+/-}$+PBS (n=15), $Sh3^{+/-}$+LPS (n=15). *** **

Light/dark box

F — Moving distance in light chamber (cm): WT+PBS (n=14), WT+LPS (n=16), $Sh3^{+/-}$+PBS (n=16), $Sh3^{+/-}$+LPS (n=15). * *

Grooming

G — Duration (s): WT+PBS (n=12), WT+LPS (n=16), $Sh3^{+/-}$+PBS (n=17), $Sh3^{+/-}$+LPS (n=19). * *

Social preference

H — Duration (s); S vs E: WT+PBS *, WT+LPS **, $Sh3^{+/-}$+PBS *, $Sh3^{+/-}$+LPS **, $Sh3^{-/-}$ ns.

I — Duration (s); SP vs EP: *, **, ****, ***, ns.

WT+PBS (n = 11) — WT+LPS (n = 11) — $Sh3^{+/-}$+PBS (n = 20) — $Sh3^{+/-}$+LPS (n = 23) — $Sh3^{-/-}$ (n = 8)

Figure 2

Figure 3

A. *Sh3*[+/-]+LPS vs. *Sh3*[+/-]+PBS

75 downregulated genes
488 upregulated genes

B. WT+LPS vs. WT+PBS

108 downregulated genes
433 upregulated genes

C. *Sh3*[+/-]+LPS vs. WT+LPS

37 downregulated genes
88 upregulated genes

D. *Sh3*[+/-]+PBS vs. WT+PBS

33 downregulated genes
67 upregulated genes

E. **Shared Upregulated Genes**

Sh3[+/-]+LPS vs. WT+LPS (88)
WT+LPS vs. WT+PBS (433)
Sh3[+/-]+PBS vs. WT+PBS (67)
Sh3[+/-]+LPS vs. *Sh3*[+/-]+PBS (488)

F. **Shared Downregulated Genes**

Sh3[+/-]+LPS vs. WT+LPS (37)
WT+LPS vs. WT+PBS (108)
Sh3[+/-]+PBS vs. WT+PBS (33)
Sh3[+/-]+LPS vs. *Sh3*[+/-]+PBS (75)

G. *Sh3*[+/-]+LPS vs. WT+LPS

Activation (positive z-score)
Inhbition (negative z-score)

Number of genes that overlap the pathways

Interferon gamma signaling
Interferon alpha/beta signaling
Pathogen Induced Cytokine Storm Signaling Pathway
Neuroinflammation Signaling Pathway
Acetylcholine Receptor Signaling Pathway
CREB Signaling in Neurons
Gap Junction Signaling Pathway
Multiple Sclerosis Signaling Pathway
Potassium Channel Signaling Pathway
Glutamatergic Receptor Signaling Pathway
Cachexia Signaling Pathway

Figure 4

Figure 5

24 hours after LPS/PBS injection

A IBA1/CD68/vGluT1 3D rendering — IBA1, CD68, vGluT1

WT+LPS

Sh3+/-+LPS

5 µM   1 µM

B CD68 occupancy /IBA1+

C vGluT1 occupancy /CD68+ IBA1+   ****

D vGluT1 occupancy /IBA1+   ***

WT+LPS (n=19)   Sh3+/-+LPS (n=28)

E vGluT1   PSD95   Overlay

WT+LPS

Sh3+/-+LPS

1 µM

F Colocalization Pearson's coefficient   ***

G Colocalization /PSD95   **

H Synapse density (n/um²)   *

WT+LPS (n=12)   Sh3+/-+LPS (n=14)

2 weeks after LPS/PBS injection

I vGluT1   PSD95   Mask   Overlay

WT+PBS

WT+LPS

J vGluT1   PSD95   Mask   Overlay

Sh3+/-+PBS

Sh3+/-+LPS

1 µM

K Colocalization Pearson's coefficient   **   **

L Colocalization / PSD95   ****   ****

M Synapse density (n/um²)   ****   ****

WT+PBS (n=19)   WT+LPS (n=12)   Sh3+/-+PBS (n=18)   Sh3+/-+LPS (n=12)

N   WT+PBS   WT+LPS   Sh3+/-+PBS   Sh3+/-+LPS

PSD95

HOMER 1b/c

β-ACTIN

Q   WT+PBS   WT+LPS   Sh3+/-+PBS   Sh3+/-+LPS

NR2A

GluA1

β-ACTIN

O Normalized PSD95 expression   ***   *

P Normalized HOMER1 expression   ****   **

R Normalized NR2A expression   **   **

S Normalized GluA1 expression   *   *

WT+PBS   WT+LPS   Sh3+/-+PBS   Sh3+/-+LPS

Figure 6

$Shank3^{+/+}$ $Shank3^{+/-}$

Blood brain barrier

PSD

Capillary   Brain

Endothelial cells   LPS   Proinflammatory cytokines and chemokines

Neutrophil   TLR-4   SHANK3   HOMER   PSD95

Microglia   Activated microglia   PSD: Postsynaptic density region