Structure-Activity Relationship Studies Towards Analogues of Pleconaril as Novel Enterovirus-D68 Capsid-Targeting Antivirals

David L. Cousins,[*,†,⊥] Ed J. Griffen,[†,⊥] Jessica Stacey,[†,⊥] Alpha A. Lee,[*,‡,⊥] Yuliia Filimonova,[¶] Anton Hlavin,[¶] Yuliia Holota,[¶] Roman Khmil,[¶] Mykyta Kordubailo,[¶] Oleksii Kostinov,[¶] Dmytro Lesyk,[¶] Ivan Logvinenko,[¶] Mariia Lototska,[¶] Viacheslav Lysenko,[¶] Anna Pashchenko,[¶] Mariia Pavlichenko,[¶] Anzhela Rodnichenko,[¶] Anton Tkachenko,[¶] Brett L. Hurst,[§] Justin G. Julander,[§] Hong Wang,[§] Rebecca Pearl,[∥,⊥] Jared Benjamin,[∥,⊥] Randy Diaz-Tapia,[∥,⊥] Mary E. Gordon,[∥,⊥] Randy A. Albrecht,[∥,⊥] and Kris White[∥,⊥]

†*MedChemica Consultancy Ltd, Macclesfield, Cheshire, SK11 6DU, UK.*
https://medchemica.com/
‡*PostEra, 1 Broadway, 14th floor, Cambridge, MA 02142, USA. https://postera.ai/*
¶*Enamine Ltd., Winston Churchill Street 78, Kyiv, 02094 Ukraine. https://enamine.net/*
§*Institute for Antiviral Research, Utah State University, Logan, UT 84322, USA.*
https://caas.usu.edu/iar/
∥*Department of Microbiology, Global Health and Emerging Pathogens Institute, Icahn School of Medicine at Mount Sinai, New York, NY, USA.*
https://icahn.mssm.edu/research/global-health

⊥*The ASAP Discovery Consortium, https://asapdiscovery.org/*

E-mail: david.cousins@medchemica.com; alpha.lee@postera.ai

Abstract

Non-polio enteroviruses (NPEV) such as enterovirus D68 (EV-D68) that are highly infectious and associated with polio-like neurological complications have caused outbreaks, globally, in recent years. While some clinical and preclinical compounds have shown efficacy against NPEV *in-vitro*, liabilities that caused historical compounds such as pleconaril to fall short of FDA approval still remain. We present herein SAR and SPR studies of analogues of clinical compounds such as pleconaril and vapendavir against EV-D68 as a representative NPEV. Numerous structurally differentiated analogues with EV-D68 antiviral activity and useful ADME properties were discovered, which could serve as starting points for future EV drug discovery campaigns. Screening against a panel of enteroviruses revealed moderately broad-spectrum anti-EV activity of compound 26.

Keywords

Pandemic Preparedness, Antiviral, Drug Discovery, Hit To Lead, Enterovirus, EV-D68, Pleconaril, Vapendavir, Capsid

Introduction

Poliovirus has almost been eradicated through immunisation and other approaches.[1–3] There are, however, a number of emerging non-polio enteroviruses (NPEV) belonging to the same viral family (*Picornaviridae*) whose outbreaks have been confirmed globally in recent years (Figure 1) and represent a pandemic risk.[4–13] These include Enterovirus-D68 (EV-D68),

Figure 1: Phylogenetic tree depicting the evolutionary history of EV-D68 for the period 1995-2023, The location of the US/MO/47 strain discovered during the 2014 outbreak of EV-D68 in the US is highlighted. Interactive view generated by NextStrain (https://nextstrain.org/enterovirus/d68/genome) [21,22]

Enterovirus-A71 (EV-A71), Coxsackievirus-A6 (CV-A6), Coxsackievirus-B3 (CV-B3), variants thereof and more.

EV-D68 was associated with an outbreak of severe respiratory illness in the United States in 2014. [14,15] Clinical presentations during the outbreak were typically mild to severe respiratory disease, however, a minority of patients (usually those with naïve or compromised immune systems) developed serious neurological complications, such as acute flaccid paralysis and cranial nerve dysfunction. [16] There is a growing body of evidence linking EV-D68 infection with these polio-like neurological symptoms. [17–19] Furthermore, a 2018 study, [20] compared the infectivity of historical strains of EV-D68 with contemporary strains and found that only the more recently discovered US/MO/47 strain (Figure 1) was able to replicate in the SH-SY5Y neuronal cell line, suggesting a changing risk of neurological sequelae after EV-D68 infection. Enterovirus infections have also been linked to the onset of islet cell autoimmunity (Type 1 Diabetes), clinical trials involving anti-EV compounds are ongoing to investigate this link. [23–25]

Though EV-D68 is a non-enveloped virus, it is cold-adapted and pH-sensitive, [31,32] putting

Figure 2: Known capsid binding antiviral compounds. Published EC_{50} values against EV-D68 in-vitro for (A) pleconaril,[26] (B) CP-11526092[27] and (C) vapendavir.[28,29] Inset structures visualised using PyMOL from left to right: pleconaril (green) bound to the EV-D68 capsid (pdb ID: 4WM7), 11526092 (cyan) bound to the EV-D68 capsid (pdb ID: 7TAF), vapendavir (magenta) bound to the hRV-A2 capsid (pdb ID: 3VDD) AlogP98 = calculated logP.[30]

the primary site of viral replication and transmission in the upper respiratory tract. This heightens the risk of rapid airborne transmission and wider spread of the virus. Despite the risk posed by pandemic EV-D68, there are currently no approved drugs, vaccines or monoclonal antibodies available. The closest point to an approved drug capable of treating EV-D68 was pleconaril (**1**, Figure 2A), an orally bioavailable small molecule that shows broad spectrum picornavirus antiviral activity *in-vitro*.[33–35] Pleconaril was developed as a treatment for the common cold (primarily human rhinoviruses - hRVs), but FDA approval was rejected in 2002 due to a combination of: lack of efficacy, safety concerns and other factors.[36]

The mode of action (MoA) of pleconaril (Figure 3) is mimicry of the natural lipid pocket factor that sits in the hydrophobic cavity of Viral Protein 1 (VP1, one of 4 protomers that make up the icosahedral viral capsid: VP1-4).[26,37,38] This pocket factor regulates capsid stability and is released when in the acidic environment of the endosome after cell entry,

Figure 3: (A) Representation of the EV-D68 replication cycle. (B) Cartoon depicting the topology of the hydrophobic lipid pocket of VP1 with regard to the bound pose of pleconaril, 1 and other capsid-binding antivirals.

leading to viral uncoating and injection of viral RNA into the host cell. Pleconaril replaces the pocket factor and prevents one or both of: (1) virus-cell attachment and (2) viral uncoating. The cavity of VP1 lies at the bottom of a cleft in the viral capsid, sometimes referred to as the "canyon" (Figure 3B),[39–41] which is thought to selectively recognise cellular receptors such as the Intercellular Adhesion Molecule (ICAM-5 and ICAM-1),[42–45] the human Scavenger Receptor class B member 2 (hSCARB2)[46] and sialic acid receptors[47–49] while resisting immune system surveillance. Other molecular targets investigated for anti-EV-D68 drug discovery include key viral proteases, such as 2A, 2C, 3C proteases and the RNA-dependant RNA polymerase (RdRp). All these targets and approaches to anti EV-D68 drug discovery have been reviewed extensively in recent years.[50–53]

Other capsid binders analogous to pleconaril have been investigated recently, such as vapendavir (3, Figure 2), which was also developed as a hRV treatment but was similarly halted due to efficacy concerns.[54,55] The *in-vitro* antiviral activity of vapendavir against

EV-D68 is ambiguous, with one source showing activity[29] and another showing no activity against four strains of EV-D68.[28] Since the discovery of pleconaril, there have been numerous structure-activity relationship (SAR) studies against NPEV of concern.[27,56–65]

For repurposing pleconaril towards the treatment of emerging NPEV, mutations in the VP1 pocket identified as potentially causing pleconaril resistance should be addressed.[66,67] Makarov and coworkers have carried out systematic work in recent years,[27,58,61–63,68] testing close analogues of pleconaril against both pleconaril-sensitive and pleconaril-resistant NPEV strains. Intriguing insights into the SAR of this series have been revealed, including a new ligand-bound cryo-EM structure of the EV-D68 capsid,[27] resulting in the discovery of the next generation antiviral compound: CP-11526092 (**2**, Figure 2). While representing a step-improvement in broad spectrum NPEV activity from pleconaril, **2** is still highly lipophilic and poorly soluble. The physicochemical property profiles of **1** and **2** are not conducive to an orally bioavailable candidate drug without the need for lipid formulations and special drug delivery systems. This introduces another known risk with the pleconaril series: that of cytochrome P450 (CYP) enzyme induction after repeated dosing. This can lead to drug-drug interactions that are unacceptable for a community use antiviral drug. CYP induction is a well-known side effect of **1**,[69] and is still present in **2** according to *in-vitro* data.[27] Neither **1** nor **2** are agonists of the human pregnane-X receptor (PXR),[27] which is a common cause of this phenomenon. Further work is required to determine the root cause of CYP induction and what *in-vitro* data are clinically relevant.

The mechanisms of CYP induction are complex and hard to predict due to overlap of nuclear hormone receptor signalling pathways and other factors.[70,71] However, we speculate that the shared property elements of **1** and **2**, including high lipophilicity (logP), high numbers of rotational bonds (nRotB), high plasma protein binding (PPB) and relatively slow metabolism combine to give a unique CYP induction risk profile. Data describing the CYP induction status of **3** are not in the public domain, to our knowledge.

Thus, our approach to the continued development of the pleconaril series of compounds

as EV-D68 antivirals was to find the minimal pharmacophoric elements of antiviral activity and introduce structural novelty to diverge significantly from the established property space (Figure 4). This will be presented here in two parts, firstly, we will present our attempts at rationally editing the present series including combining structural features of literature compounds where structurally and synthetically plausible. Secondly, we will share our results in wider library design and antiviral screening.

Figure 4: Summary of the medicinal chemistry strategy adopted in this work

Synthetic Chemistry

The majority of analogues presented here were made according to Scheme 1, whereby the alcohol **5** was prepared from **4**, and reacted with selected (hetero)phenols under Mitsunobu conditions.[72,73] The conditions found proved to be generally applicable and were sufficient to rapidly produce analogues **6** in both parallel and singleton fashion.

Scheme 1: Reagents and conditions: (i) dimethylamine, HATU, DIPEA, MeCN, RT, 12h (ii) phenol, PPh₃ or PBu₃, DEAD or DIAD or ADDP, THF, RT 16h

In contrast, the preparation of ether analogues **10** (Scheme 2) of 2-methyl-4-[5-(trifluoromethyl)-1,2,4-oxadiazol-3-yl]phenol proved to be less generally applicable, primarily due to the chem-

ical sensitivity of the CF$_3$-oxadiazole ring towards basic workup procedures. It is for this reason that cyclised analogues **14** (Scheme 3) were prepared according to Scheme 3, where the CF$_3$-oxadiazole was constructed in the very last step from the corresponding amidoxime and TFAA.

Scheme 2: Reagents and conditions: (i) hydroxylamine hydrochloride, K$_2$CO$_3$, EtOH (absolute), reflux, 16h (ii) TFAA, pyridine, 80 °C, 2h (iii) 3-(hetero)arylpropyl alcohol, PPh$_3$, DEAD, THF, RT, 12h

Scheme 3: : Reagents and conditions: (i) methyl 5-(3-bromopropyl)-1,2-oxazole-3-carboxylate, NaH, DMF, 0 to 50 °C, 16 h (ii) LiOH, THF/H$_2$O, RT, 16h (iii) dimethylamine, HATU, DIPEA, RT, 16h (iv) hydroxylamine hydrochloride, NaHCO$_3$, EtOH, 70 °C, 16h (v) TFAA, THF, RT, 16h

Regarding the preparation of vapendavir-pleconaril hybrid compounds **19** (Scheme 4), a general route that proved successful can be seen in Scheme 4. 2,4-difluoro benzoic acids **15** proved to be very useful double electrophiles in both the synthesis of benzisoxazoles **17** and the final phenolate ester compounds **19**. Other schemes to obtain similar products were

attempted, such as in Scheme 5, where 3-chloro-6-fluoro-1,2-benzoxazole **20** was reacted sequentially with the appropriate alcohol under $S_N Ar$ conditions. Interestingly, this scheme resulted in regioisomeric products **21a** and **21b** (isolated in 1:1 ratio), upon testing, it was found that an example of the originally undesired isomer **21b** (see compound **51**, Table 5) was also active against EV-D68.

Scheme 4: Reagents and conditions: (i) hydroxylamine hydrochloride, oxalyl chloride (ii) K_2CO_3, EtOAc/H_2O, 0 °C to RT, 16 h (iii) KOH, ^{n}BuOH, 120 °C, 4 h (iv) ethyl iodide, Ag_2O, $CHCl_3$, reflux, 16 h (v) 5-(3-hydroxypropyl)-N,N-dimethyl-1,2-oxazole-3-carboxamide, NaH, DMF, 100 °C, 16h

Scheme 5: Reagents and conditions: (i) aliphatic primary alcohol, NaH, DMF, 0 °C to RT, 16 h (ii) 5-(3-hydroxypropyl)-N,N-dimethyl-1,2-oxazole-3-carboxamide, NaH, DMF, 0 °C to RT, 16h

Results and discussion

In-Vitro Antiviral and ADME Characterisation of Known EV Capsid-Binding Antivirals

Table 1: Measured EV-D68 antiviral activity (RD cells) and *in-vitro* ADME profile of relevant known capsid binders from the literature. EC_{50} and CC_{50} are reported as the geometric mean. P_{app} = Apparent Permeability (MDCKII-MDR1 cell system), Cl_{int} = Intrinsic Clearance, HLM = Human Liver Microsomes, MLM = Mouse Liver Microsomes. See the Supporting Information for further details.

Structure			
Identifier	Pleconaril (**1**)	CP-11526092 (**2**)	Vapendavir (**3**)
EC_{50} EV-D68 $[CC_{50}](\mu M)$	<0.029[>50]	0.043[>50]	0.170[>50]
AlogP98 [30]	4.7	3.8	4.0
logD (pH 7.4)	4.2	$\geq$4.5	3.0
Solubility, Kinetic (KSOL, μM)	18	9	16
P_{app}A$\rightarrow$B (10^{-6} cm s^{-1})	1.42	2.26	2.41
Cl_{int} HLM , MLM (μL min^{-1} mg^{-1})	12 , 22	11 , 24	42 , 95

Our first round of testing was on the key clinical and preclinical capsid-binding anti-EV compounds found in the literature, on which we would base our studies. We initially screened these compounds in our EV-D68 antiviral assay and a suite of ADME tests, including determination of: logD, kinetic aqueous solubility (KSOL), human and mouse microsomal stability (HLM and MLM, respectively) and MDCK-MDR1 A->B permeability. Our results for **1**, **2** and **3** can be seen in Table 1 (for a full table of data including additional literature capsid binders, see **Table S1** in the supporting information). Obtaining antiviral EC_{50} values against EV-D68 confirmed the sub-μM potency of compounds **1**-**3**, with **1** having the highest potency at <29 nM. Close analogue of pleconaril: **2** and the clinical antiviral **3** were similarly potent in our hands, with EC_{50} values of 43 nM and 170 nM, respectively.

The ADME properties of the published compounds were measured (Table 1), these compounds generally exhibit: high lipophilicity, high permeability and low aqueous solubility with varying levels of intrinsic microsomal clearance. **1** is known to have useful PK properties,[35,74] although the oral bioavailability (%F) is heavily dependent on whether the patient is in a fed or fasted state. The high %F of **1** is in spite of its high lipophilicity (measured logD = 4.20), due to a number of factors including: high tissue penetration[75] and a "global protective" effect from P450 metabolism provided by the CF_3 group attached to the oxadiazole ring.[76]

Structural Editing of Known EV Capsid-Binding Antivirals

Table 2: Measured EV-D68 antiviral activity (RD cells) and and *in-vitro* ADME profile of linker-variant compounds from Compound **2**. See the Supporting Information for further details.

Identifier	Structure	$EC_{50}[CC_{50}]$ (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 $P_{app}A{\to}B$ (10^{-6} cm s^{-1})	Cl_{int} HLM , MLM (μL min^{-1} mg^{-1})
22		0.115[>10]	3.5	68	15.2	37 , 79
23		0.216[>45]	4.0	48	4.0	48 , 87
24		>30.4[>50]	ND	ND	ND	ND , ND

Building on the extensive SAR studies by Makarov and coworkers cited above, we began our investigations by modifying compound **2**. Moving the oxygen atom of the ether linker along the chain by one (**22**, Table 2) and two (**23**) atoms resulted in a drop in logD and a jump in the measured kinetic aqueous solubility. Both **22** and **23** displayed slightly reduced anti-EV-D68 potency compared to **2**, with EC_{50} values of 115 nM and 216 nM respectively. Moving to an ethanolamine linker (**24**) resulted in much diminished antiviral activity. Al-

though it wasn't experimentally determined, the cell permeability was expected to be lower than **2** due to the introduction of the aniline NH into the chain.[77] To probe the effects of conformational restriction by scaffold-hopping, (aza)indole-linked compounds **25-30** (Table 3) were prepared according to Scheme 3. The intermediate nitrile **25** was tested and found to have an EC_{50} of 2.87 μM, significantly less potent against EV-D68 than **2**, but appreciably active. The first indole oxadiazole derivative **26** was found to be very potent against EV-D68, with an EC_{50} value of 17 nM. While the high antiviral potency of **26** against EV-D68 was a promising result, the measured ADME properties suggested limited improvement from **2** in terms of lipophilicity (logD = $\geq$4.5) and aqueous solubility (KSOL = 8 μM).

Thus, nitrogen substituted analogues **27-30** were also prepared and profiled similarly. All of these aza-substituted analogues dropped the measured logD and increased KSOL relative to **2**, as expected, but this was found to be at the cost of EV-D68 antiviral potency. Benzene and pyridine linker replacements for the central propyl chain were investigated in **31** and **32**, respectively. Neither of these provided potency gains, nor were expected to offer any benefit in terms of ADME properties so were deprioritised from further testing. Finally, pyrollidine-linked enantiomers **33** and **34** were prepared. Both enantiomers were found to have sub-micromolar EC_{50} values, but **34** was 10-fold more potent than **33**. Intrigued by the departure from closer analogues of **2** described above: represented by these rigid pyrollidine-linked compounds, we further profiled **33** and **34** in ADME assays. Compared to **2**, they maintained a similar profile, with high lipophilicity (logD = 4.2), low kinetic solubility (<10 μM), high permeability and moderate to low microsomal clearance. While the properties of **34** as a whole were not sufficient for progression, 2 rotatable bonds were eliminated from **2** through this change to a pyrollidine linker, making the compound significantly more conformationally rigid. This novel chemotype is a highly attractive core for further derivatisation and property optimisation.

We then turned our attention to the VP1 pore-binding (Figure 3B) region, by preparing analogues of **2** according to Scheme 2, using readily available (heteroaryl)propyl alcohol

Table 3: Measured EV-D68 antiviral activity (RD cells) and and *in-vitro* ADME profile of scaffold-hopped designs. See the Supporting Information for further details.

Identifier	Structure	EC$_{50}$[CC$_{50}$] (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 P$_{app}$A→B (10^{-6} cm s^{-1})	Cl$_{int}$ HLM , MLM (μL min^{-1} mg^{-1})
25		2.87[>50]	ND	ND	ND	ND , ND
26		0.017[>50]	≥4.5	8	1.0	29 , 26
27		>1.78[>50]	3.6	52	10.9	15 , 111
28		1.03[>50]	2.7	205	20.7	51 , 44
29		1.89[>100]	3.8	28	15.5	30 , 133
30		8.43[>100]	3.0	30	3.5	20 , 29
31		4.87[>50]	ND	ND	ND	ND , ND
32		4.35[>50]	ND	ND	ND	ND , ND
33		0.437[>50]	4.2	9	4.0	24 , 38
34		0.058[>50]	4.2	7	2.5	22 , 41

building blocks (Table 4). 1,3,4-oxadiazoles **35-38** were all found to be active against EV-D68. Unsubstituted azole **35** was potent, with an EC$_{50}$ of 41 nM while also offering a lower logD and higher solubility than **2**. Methyl substituted analogue **36** lost some EV-D68 potency

(EC_{50} = 932 nM), but further improved the solubility (KSOL = 78 μM). **36**, however, was found to have moderate microsomal clearance for both human and mouse (MLM/HLM Cl_{int} = 75/17 μL min^{-1} mg^{-1} protein). Interestingly, **39** (the triazole matched pair of **36**) had much lower microsomal clearance values (MLM/HLM Cl_{int} = 15/<10 μL min^{-1} mg^{-1} protein) and was significantly more potent against EV-D68, but it did display a lower solubility (KSOL = 18 μM) compared to **36**. Fluoroalkyl substitution of the oxadiazole was explored, first with difluoromethyl derivative, **37**, which showed high antiviral potency but very low solubility (KSOL = 6 μM). Trifluoromethyl derivative **38** was found to be much less potent (EV-D68 EC_{50} = 39.9 μM) and was not characterised further.

Table 4: Measured EV-D68 antiviral activity (RD cells) and and *in-vitro* ADME profile of compounds with alternative capsid VP1 pore-binding groups. See the Supporting Information for further details.

Identifier	Structure	$EC_{50}[CC_{50}]$ (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 P_{app} A→B (10^{-6} cm s^{-1})	Cl_{int} HLM , MLM (μL min^{-1} mg^{-1})
35		0.067[>43]	4.0	21	12.1	<10 , 13
36		0.932[>50]	4.4	78	8.9	17 , 75
37		<0.065[>48]	4.4	6	1.9	10 , 19
38		39.9[>50]	ND	ND	ND	ND , ND
39		0.117[>50]	4.1	18	13.8	<10 , 15

Identifier	Structure	EC$_{50}$[CC$_{50}$] (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 P$_{app}$ A$\rightarrow$B (10^{-6} cm s^{-1})	Cl$_{int}$ HLM , MLM (μL min^{-1} mg^{-1})
40		0.445[>50]	4.0	5	0.5	<10 , 18
41		0.287[>50]	4.2	4	0.7	18 , 73
42		0.179[>50]	≥4.5	10	0.6	11 , 31
43		1.31[>50]	≥4.5	7	0.6	15 , 46
44		1.17[>50]	≥4.5	4	0.5	11 , 24
45		0.409[>50]	≥4.5	9	2.0	<10 , 34

Pyrazole (**40**, **41**) and pyridine (**42-44**) isomers were found to have useful, but variable levels of EV-D68 potency, these could be considered as useful isosteres for the isoxazole pore-binding groups found in **1** and **2**, for later SAR/SPR combinations. Finally, the phenyl tetrazole compound **45** was found to have useful anti-EV-D68 potency, with an EC$_{50}$ of 55 nM. This tetrazole functionality bears some resemblance to the EV-D68 antiviral R856932 recently discovered by Wang *et al.*[60] whose capsid-binding MoA was investigated extensively

and is expected to bind the pore region of VP1 as we also expect of the tetrazole of **45**.

To explore SAR and SPR mergers between the pleconaril (**1**) and vapendavir (**2**) compound series, novel hybrid compounds **46**-**50** (Table 5) were prepared according to Schemes 4 and 5. The simplest benzisoxazole combination, **46**, was found to have modest potency against EV-D68 (EC$_{50}$ = 1.04 μM), much lower lipophilicity than **2** (logD = 3.5) and moderate human and mouse microsomal clearance HLM/MLM Cl$_{int}$ = 31/140 μL min^{-1} mg^{-1} protein).

Table 5: Measured EV-D68 antiviral activity (RD cells) and and *in-vitro* ADME profile of compounds representing structural mergers between compounds **2** and **3**. See the Supporting Information for further details.

Identifier	Structure	EC$_{50}$[CC$_{50}$] (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 P$_{app}$ A→B (10^{-6} cm s^{-1})	Cl$_{int}$ HLM , MLM (μL min^{-1} mg^{-1})
46		1.04[>50]	3.5	18	14.8	31 , 140
47		2.87[>50]	4.0	7	4.1	46 , 166
48		0.285[>50]	4.0	25	15.8	97 , 418
49		0.903[>50]	3.7	7	5.1	36 , 132
50		1.33[>50]	4.4	11	2.7	41 , 100

Identifier	Structure	EC$_{50}$[CC$_{50}$] (μM)	LogD (pH 7.4)	KSOL (μM)	MDCKII-MDR1 P$_{app}$ A$\rightarrow$B (10^{-6} cm s^{-1})	Cl$_{int}$ HLM , MLM (μL min^{-1} mg^{-1})
51		0.412[>50]	4.5	15	7.1	95 , 69

Methyl substitution in the 5-position of the benzisoxazole, giving compound **47**, was detrimental to EV-D68 potency and tested ADME properties. Interestingly, 7-methyl analogue **48** was found to be much more potent (EC$_{50}$ = 152 nM) than **47** while also having slightly higher solubility (KSOL = 25 μM). The comparison of potency between isolipophilic compounds **47** and **48** is reminiscent of a "magic methyl" effect,[78,79] where a significant and non-obvious change in activity is yielded from methyl substitution. With the context of methylation SAR from the work of Makarov et al.[58,63] and alignment of published capsid structures of **2** (pdb ID = 7TAF) and **3** (pdb ID = 3VDD) (alignment not shown, see **Figures S1 and S2** in the supporting information), we can infer that the optimal location to place the single methyl group on the benzisoxazole would be the 7-position, as confirmed by the activity of **48**. Regrettably, **48** displayed metabolic instability, with HLM and MLM Cl$_{int}$ values of 97 and 418 mL min^{-1} mg^{-1} protein, respectively. Despite having a lower logD than **2**, benzisoxazole compounds **46**, **47** and **48** displayed much lower microsomal stability. This could suggest a structural metabolic vulnerability of the alkoxy benzisoxazole itself, or it could signify a return to a more conventional relationship between lipophilicity and metabolic stability from the anomalous profile of **1** and **2**.

Figure 5: Plot of EV-D68 antiviral potency (pEC50 = -log10(EC50)) vs calculated lipophilicity (AlogP98) for compounds designed via rational editing of known antivirals. Points are coloured based on their measured kinetic solubility values (KSOL / μM). Lipophilic Cellular Efficiency (LipCellE) displayed as normalisation factors of cellular potency against the lipophilicity of the compound (LipCellE = cellular pEC50 - AlogP98)

Alternative alkyl chain termini were investigated in compounds **49** (cyclopropyl methyl) and **50** (isobutyl), these compounds were similar in potency to the ethyl parent **46** and didn't offer any improvement in ADME properties. As outlined in Scheme 5, sequential addition of alcohols to the bifunctional reagent **20** resulted in isomeric structures **21a** and **21b**.

One such example of isomer **21b**, compound **51** (Table 5), was tested and found to have an EV-D68 EC_{50} of just 412 nM. This intriguing result encouraged us to profile **51** further, revealing broadly similar ADME properties to its isomer **50**, lower MLM clearance (MLM $Cl_{int} = 69$ mL min^{-1} mg^{-1} protein) but interestingly, slightly higher HLM clearance (HLM $Cl_{int} = 95$ mL min^{-1} mg^{-1} protein).

As high lipophilicity is one of the key risks with most of these compounds, a key metric for ranking was the Lipophilic Ligand Efficiency (LLE),[80,81] or in our case of cellular potency measurements: Lipophilic Cellular Efficiency (LipCellE). LipCellE is defined as the compound's measured pEC_{50} value minus its calculated lipophilicity (AlogP98). Figure 5 shows

an overview of EV-D68 cell potency (pEC_{50}) for the compounds discussed in the previous section, against their AlogP98 values and coloured by KSOL.

Figure 5, <u>A</u> shows the position of **2**, highlighting that although it is highly lipophilic, its high potency places the LipCellE at >3 indicating that the lipophilic groups on **2** add potency more efficiently than many others in this set (I.e. compounds with LipCellE<3). Moving to Figure 5, <u>B</u>, shows us compound **28**, which displays excellent aqueous KSOL and a lower logP than **2**, but the loss in potency is enough to lower the LipCellE value to below 3. Almost equipotent to **28** is vapendavir-hybrid compound **46** (Figure 5, <u>C</u>), with a lower lipophilicity, reinstating the higher LipCellE. 5'-methylation of the benzisoxazole of **46** gives us **47** (Figure 5, <u>D</u>), this was a move that increased the lipophilicity and lowered the potency which is the most direct way to lower LipCellE. In contrast, 7'-methylation in compound **48** (Figure 5, <u>E</u>) provided a 19-fold increase in potency from **47** for the same lipophilicity, providing a drastic increase in LipCellE. Finally, moving to the highly constrained pyrollidine **34** (Figure 5, <u>F</u>) resulted in yet higher potency along with only a slight increase in lipophilicity, further increasing the LipCellE, just above that of **2**.

Design and Synthesis of Pleconaril Analogue Libraries

During the course of our synthetic work, we discovered an efficient set of Mitsunobu coupling conditions for the production of phenol ethers (Scheme 1). These conditions were robust enough to be employed in parallel synthesis, so we designed a library of compounds based on available phenol building blocks (BBs) aimed at exploring the VP1 pocket more thoroughly.

We began this process by characterising a set of ˜4000 phenols (Figure 6) with calculated properties, including: 3D shape descriptors (Normalised Principal Moment of Inertia, NPMI1, NPMI2)[82] and pharmacophoric descriptors (identity and topological bond count from reacting group of pharmacophoric features). A summary of this full characterisation can be seen in Figure 6B, with the NPMI plot showing the 3D shape distribution of the total BB set lying largely on the "Rod-Disk" axis, as expected for a phenol library. The

Figure 6: Method for selection of building block (BB) phenols for application in parallel and singleton Mitsunobu reactions, enabling further exploration of the capsid VP1 lipophilic pocket. **A**: Methods through which phenol BB libraries were characterised, **B**(left): Pharmacophore feature heatmap for starting phenol BB library, **B**(right): NPMI[82] plot for starting phenol BB library, **C**(left): Pharmacophore feature heatmap for selected phenol BBs, **C**(right): NPMI plot for selected phenol BBs, **D**: Scheme for Mitsunobu reactions suitable for parallel synthesis, used to make products. Base = basic group, Acid = Acidic group, HBA = hydrogen bond acceptor, HBD = hydrogen bond donor, Lipo = lipophilic group, Aro = aromatic group, Aliph = aliphatic group, Halo = halogen, phco4 = pharmacophore, NPMI = Normalised principal moment of Inertia

pharmacophoric feature map displayed good coverage of features among the total set, with the exception of acidic groups ("Acid"). This is due to carboxylic acids being filtered to avoid alkylation under Mitsunobu conditions.

Our phenol picking criteria were centred on whether or not it added diversity in either 3D-shape or pharmacophoric feature coverage (within the confines of chemical space provided by the input phenol BBs), alongside a final ranking based on heavy atom count (HAC) to ensure only the most efficient cluster representatives were taken forward. Out of 123 systematically picked phenols, 80 enumerated ether products were proposed for synthesis,

Figure 7: Plot of EV-D68 antiviral potency (pEC50 = -log10(EC50)) vs calculated lipophilicity (AlogP98) for compounds designed via diversity-based library approach. Points are coloured based on their measured kinetic solubility values (KSOL / μM). Lipophilic Cellular Efficiency (LipCellE) displayed as normalisation factors of cellular potency against the lipophilicity of the compound (LipCellE = cellular pEC50 - AlogP98)

of these, 75 were made and tested. The characteristics of the building blocks associated with these 75 compounds can be seen in Figure 6C. Some gaps in pharmacophoric feature coverage relative to the starting set (Fig. 6A) can be seen, such as "Base 5" and "Aro 6". This suggests that our selection lacked coverage of basic groups 5 atoms away from the phenol and aromatic groups 6 atoms away. Nevertheless, numerous compounds active against EV-D68 were discovered, a summary of their antiviral pEC_{50} values vs their AlogP98 can be seen in Figure 7.

The compounds that initially caught our eye as having maintained some EV-D68 antiviral activity, were the fluoroaryls **52**, **53** and **54**. Simple fluorobenzene derivatives **52** and **54** had modest EV-D68 antiviral activity (EC_{50} values of 9.88 μM and 9.49 μM, respectively), but showed reduced lipophilicity and increased KSOL values, indicating that a compromise between potency and lipophilicity could possibly be reached with further development. The more heavily substituted **53** was more potent, but at the expense of higher lipophilicity and

lower solubility (logD = 4.2, KSOL = 7 μM). Propargyl derivative **55** was found to have low micromolar potency (EC$_{50}$ = 4.88 μM) and a modest logD value of 3.1, alongside a useful KSOL of 198 μM. Interestingly, the extended homologue of **55**: **56**, was found to have the highest LipCellE out of the entire library and it maintained a lower logD and higher KSOL.

Assessment of Broad-Spectrum Antiviral Activity Through EV Panel Screening

To test the most promising compounds for broad spectrum anti-enterovirus activity, we profiled **26**, **48** and **56** against a panel of 8 enteroviruses (Table 6).

Compound **26** gave the most promising profile, showing broad spectrum low-micromolar to nanomolar EC$_{50}$ values against 7 out of 8 viral strains tested with CC$_{50}$ values of >32 μM across all cell lines. Sub-micromolar EC$_{50}$ values were seen for EV-D68 (US/ KY/14-18953 strain), Echovirus-11 (Echo-11), Echovirus-30 (Echo-30) and Human Rhinovirus-14 (HRV-14). Slightly lower potency was seen for EV-A71 (3.9 μM), Poliovirus-1 (POV-1, 4.1 μM) and Coxsackievirus-B3 (Cox-B3, 1.8 μM). The notable exception to the broad-spectrum activity of **26** was Coxsackievirus-A6 (Cox-A6), for which no antiviral activity was observed.

Moving to the vapendavir hybrid compound **48**, robust activity against EV-D68 (US/KY/14-18953 strain) was observed plus some other low micromolar potencies, however, cytotoxicity was also observed across the cell lines used which complicated the assessment of antiviral activity.

Finally, Mitsunobu library compound **56** displayed low micromolar EC$_{50}$ values against EV-D68 (4.5 μM) and EV-A71 (3.5 μM) and slightly less cytotoxicity throughout the panel than **48**. However, lower antiviral potency for **56** was also observed across the board. This is somewhat unsurprising, given that **56** was discovered as part of the screening campaign against EV-D68 (US/MO/14-18949 strain) (Figure 7) and did not contain as many highly optimised structural features such as those from CP-11526092 and vapendavir in compounds **26** and **48** (respectively).

Table 6: Summary of results from the screening of **26**, **48** and **56** against a panel of 8 enteroviruses. EV = Enterovirus, POV = Poliovirus, Echo = Echovirus, HRV = Human Rhinovirus, Cox = Coxsackievirus. See the supporting information for further details.

Compound	Virus	Strain	Cell Line	EC_{50} (μM)	CC_{50} (μM)
	EV-D68	US/KY/14-18953	RD	<0.01	>32
	EV-A71	Tainan/4643/98	Vero-76	3.9	>32
	POV-1	Mahoney	Vero 76	4.1	>32
26	Echo-11	Gregory	RD	0.18	>32
	Echo-30	Bastianni	RD	0.72	>32
	HRV-14	1059	HeLa Ohio	0.67	>32
	Cox-B3	HA 201933	Vero-76	1.8	>32
	Cox-A6	Gdula	RD	>32	>32
	EV-D68	US/KY/14-18953	RD	0.57	>32
	EV-A71	Tainan/4643/98	Vero-76	3	9
	POV-1	Mahoney	Vero 76	2.4	9.4
48	Echo-11	Gregory	RD	1.7	9.5
	Echo-30	Bastianni	RD	1.4	9.3
	HRV-14	1059	HeLa Ohio	1.9	>32
	Cox-B3	HA 201933	Vero-76	>5.6	5.6
	Cox-A6	Gdula	RD	>11	11
	EV-D68	US/KY/14-18953	RD	4.5	>32
	EV-A71	Tainan/4643/98	Vero-76	3.5	26
	POV-1	Mahoney	Vero 76	18	23
56	Echo-11	Gregory	RD	18	18
	Echo-30	Bastianni	RD	>14	14
	HRV-14	1059	HeLa Ohio	5.7	>32
	Cox-B3	HA 201933	Vero-76	>19	19
	Cox-A6	Gdula	RD	>32	>32

Summary and Conclusions

In summary, we have surveyed numerous approaches in the literature towards repurposing analogues of **1** from its original rhinovirus indication. **1** failed to gain FDA approval for its targeted indication, but remains an interesting clinical tool for treating NPEV, which are of potentially greater public health concern. Among known pleconaril analogues, compound **2** was found to have increased potency towards NPEV than pleconaril, however, it resides in a similar property space and still holds the risk of CYP induction, based on *in-vitro* data. The clinical compound **3** occupies a much more favourable property space, but does not offer the NPEV potency of **1** and **2**, according to the literature.

We prepared analogues of the pleconaril and vapendavir series of compounds and tested them against EV-D68 *in-vitro*, as a representative of the NPEV class. We also tested interesting analogues in our tier 1 ADME assays to probe clinical usefulness in the development of orally bioavailable community-use anti-EV-D68 drugs. Compounds were designed by both rationally editing known antivirals in a matched molecular pair approach and by building block-based library generation. Numerous compounds that are significantly differentiated from published examples, in both structure and properties, displayed potent antiviral activity against EV-D68.

We screened 3 compounds against a panel of 8 enteroviruses and identified **26** as a broad-spectrum enterovirus inhibitor. Other compounds **48** and **56** displayed some antiviral activity against NPEV of concern, but also displayed cytotoxicity in a similar concentration window. No antiviral activity was observed against Coxsackievirus-A6 from any of the compounds tested.

In conclusion, the SAR studies undertaken here provide potentially useful data and offer new starting points for those looking to develop potent and orally bioavailable drugs to treat NPEV.

Acknowledgement

The authors thank Karla Kirkegaard (ORCID: 0000-0001-7628-3770) for engaging discussions.

Funding

This work was supported by the NIH and NIAID Antiviral Drug Discovery (AViDD) grant (U19AI171399). This work was also supported in part by services provided by the NIAID Preclinical Services Program under Contract No. 75N93019D00021, Task Order No. B11.

The ISMMS BSL-3 Biocontainment Shared Research Resource is supported by subsidies from the ISMMS Dean's Office and by investigator support through a cost recovery mechanism. Research reported in this publication was supported by NIAID of the NIH under Award Number G20AI174733 (R.A. Albrecht).

Disclaimers

The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Disclosures

DLC, EJG and JS are employees of MedChemica Consultancy Ltd. AAL is an employee of PostEra Inc., which has commercial interests in the discovery, development, and commercialisation of therapeutics, including antivirals and results disclosed in this manuscript. YF, AH, YH, RK, MK, OK, DL, IL, ML, VL, AP, MP, AR and AT are employees of Enamine Ltd. BLH, JGJ and HW are employees of Utah State University. RP, JB, RD-T, MEG,

RAA and KW are employees of the Icahn School of Medicine at Mount Sinai.

Supporting Information Available

Experimental procedures and characterisation data for all new compounds, protocols and references for antiviral and ADME assays.

The deposition of all data related to this work into ChEMBL is in progress. In the meantime, data (including raw data and data not presented here) can be made available on reasonable request.

References

(1) Bhutta, Z. A. The Last Mile in Global Poliomyelitis Eradication. *The Lancet* **2011**, *378*, 549–552.

(2) Bigouette, J. P.; Wilkinson, A. L.; Tallis, G.; Burns, C. C.; Wassilak, S. G. F.; Vertefeuille, J. F. Progress Toward Polio Eradication — Worldwide, January 2019–June 2021. *MMWR. Morbidity and Mortality Weekly Report* **2021**, *70*, 1129–1135.

(3) Global Wild AFP Cases and Environmental Samples 2018-2025. https://polioeradication.org/wild-poliovirus-count/.

(4) Ang, P. Y.; Chong, C. W. H.; Alonso, S. Viral Determinants That Drive Enterovirus-A71 Fitness and Virulence. *Emerging Microbes & Infections* **2021**, *10*, 713–724.

(5) Benschop, K. S. et al. Re-Emergence of Enterovirus D68 in Europe after Easing the COVID-19 Lockdown, September 2021. *Eurosurveillance* **2021**, *26*.

(6) Gao, F.; Bian, L.; Hao, X.; Hu, Y.; Yao, X.; Sun, S.; Chen, P.; Yang, C.; Du, R.; Li, J.; Zhu, F.; Mao, Q.; Liang, Z. Seroepidemiology of Coxsackievirus B5 in Infants and

Children in Jiangsu Province, China. *Human Vaccines & Immunotherapeutics* **2018**, *14*, 74–80.

(7) Park, S. W.; Messacar, K.; Douek, D. C.; Spaulding, A. B.; Metcalf, C. J. E.; Grenfell, B. T. Predicting the Impact of COVID-19 Non-Pharmaceutical Intervention on Short- and Medium-Term Dynamics of Enterovirus D68 in the US. *Epidemics* **2024**, *46*, 100736.

(8) Pons-Salort, M.; Parker, E. P.; Grassly, N. C. The Epidemiology of Non-Polio Enteroviruses: Recent Advances and Outstanding Questions. *Current Opinion in Infectious Diseases* **2015**, *28*, 479–487.

(9) Sooksawasdi Na Ayudhya, S.; Laksono, B. M.; Van Riel, D. The Pathogenesis and Virulence of Enterovirus-D68 Infection. *Virulence* **2021**, *12*, 2060–2072.

(10) Sun, S.; Gao, F.; Hu, Y.; Bian, L.; Wu, X.; Su, Y.; Du, R.; Fu, Y.; Zhu, F.; Mao, Q.; Liang, Z. A Cross-Sectional Seroepidemiology Study of EV-D68 in China. *Emerging Microbes & Infections* **2018**, *7*, 1–7.

(11) Sun, Z.; Wang, J.; Chi, X.; Niu, P.; Zhang, R.; Gao, Q.; Zhang, Y.; Ma, X. Epidemiology and Genetic Characteristics of Coxsackievirus A16 Associated with Hand-Foot-and-Mouth Disease in Yantai City, China in 2018–2021. *Biosafety and Health* **2023**, *5*, 181–186.

(12) Tee, H. K.; Zainol, M. I.; Sam, I.-C.; Chan, Y. F. Recent Advances in the Understanding of Enterovirus A71 Infection: A Focus on Neuropathogenesis. *Expert Review of Anti-infective Therapy* **2021**, *19*, 733–747.

(13) Xiang, Z.; Li, L.; Ren, L.; Guo, L.; Xie, Z.; Liu, C.; Li, T.; Luo, M.; Paranhos-Baccalà, G.; Xu, W.; Wang, J. Seroepidemiology of Enterovirus D68 Infection in China. *Emerging Microbes & Infections* **2017**, *6*, 1–5.

(14) Midgley, C. M.; Jackson, M. A.; Selvarangan, R.; Turabelidze, G.; Obringer, E.; Johnson, D.; Louise Giles, B.; Patel, A.; Echols, F.; Steven Oberste, M.; Allan Nix, W.; Watson, J. T.; Gerber, S. I. Severe Respiratory Illness Associated with Enterovirus D68—Missouri and Illinois, 2014. *American Journal of Transplantation* **2014**, *14*, 2662–2663.

(15) Oermann, C. M.; Schuster, J. E.; Conners, G. P.; Newland, J. G.; Selvarangan, R.; Jackson, M. A. Enterovirus D68. A Focused Review and Clinical Highlights from the 2014 U.S. Outbreak. *Annals of the American Thoracic Society* **2015**, *12*, 775–781.

(16) Messacar, K.; Schreiner, T. L.; Maloney, J. A.; Wallace, A.; Ludke, J.; Oberste, M. S.; Nix, W. A.; Robinson, C. C.; Glodé, M. P.; Abzug, M. J.; Dominguez, S. R. A Cluster of Acute Flaccid Paralysis and Cranial Nerve Dysfunction Temporally Associated with an Outbreak of Enterovirus D68 in Children in Colorado, USA. *The Lancet* **2015**, *385*, 1662–1671.

(17) Messacar, K.; Asturias, E. J.; Hixon, A. M.; Van Leer-Buter, C.; Niesters, H. G. M.; Tyler, K. L.; Abzug, M. J.; Dominguez, S. R. Enterovirus D68 and Acute Flaccid Myelitis—Evaluating the Evidence for Causality. *The Lancet Infectious Diseases* **2018**, *18*, e239–e247.

(18) Park, S. W.; Pons-Salort, M.; Messacar, K.; Cook, C.; Meyers, L.; Farrar, J.; Grenfell, B. T. Epidemiological Dynamics of Enterovirus D68 in the United States and Implications for Acute Flaccid Myelitis. *Science Translational Medicine* **2021**, *13*, eabd2400.

(19) Zhang, Y.; Cao, J.; Zhang, S.; Lee, A. J.; Sun, G.; Larsen, C. N.; Zhao, H.; Gu, Z.; He, S.; Klem, E. B.; Scheuermann, R. H. Genetic Changes Found in a Distinct Clade of Enterovirus D68 Associated with Paralysis during the 2014 Outbreak. *Virus Evolution* **2016**, *2*, vew015.

(20) Brown, D. M.; Hixon, A. M.; Oldfield, L. M.; Zhang, Y.; Novotny, M.; Wang, W.;

Das, S. R.; Shabman, R. S.; Tyler, K. L.; Scheuermann, R. H. Contemporary Circulating Enterovirus D68 Strains Have Acquired the Capacity for Viral Entry and Replication in Human Neuronal Cells. *mBio* **2018**, *9*, e01954–18.

(21) Hadfield, J.; Megill, C.; Bell, S. M.; Huddleston, J.; Potter, B.; Callender, C.; Sagulenko, P.; Bedford, T.; Neher, R. A. Nextstrain: Real-Time Tracking of Pathogen Evolution. *Bioinformatics* **2018**, *34*, 4121–4123.

(22) Sagulenko, P.; Puller, V.; Neher, R. A. TreeTime: Maximum-likelihood Phylodynamic Analysis. *Virus Evolution* **2018**, *4*.

(23) Faulkner, C. L.; Luo, Y. X.; Isaacs, S.; Rawlinson, W. D.; Craig, M. E.; Kim, K. W. The Virome in Early Life and Childhood and Development of Islet Autoimmunity and Type 1 Diabetes: A Systematic Review and Meta-analysis of Observational Studies. *Reviews in Medical Virology* **2021**, *31*, 1–14.

(24) Isaacs, S. R.; Roy, A.; Dance, B.; Ward, E. J.; Foskett, D. B.; Maxwell, A. J.; Rawlinson, W. D.; Kim, K. W.; Craig, M. E. Enteroviruses and Risk of Islet Autoimmunity or Type 1 Diabetes: Systematic Review and Meta-Analysis of Controlled Observational Studies Detecting Viral Nucleic Acids and Proteins. *The Lancet Diabetes & Endocrinology* **2023**, *11*, 578–592.

(25) Krogvold, L.; Mynarek, I. M.; Ponzi, E.; Mørk, F. B.; Hessel, T. W.; Roald, T.; Lindblom, N.; Westman, J.; Barker, P.; Hyöty, H.; Ludvigsson, J.; Hanssen, K. F.; Johannesen, J.; Dahl-Jørgensen, K. Pleconaril and Ribavirin in New-Onset Type 1 Diabetes: A Phase 2 Randomized Trial. *Nature Medicine* **2023**, *29*, 2902–2908.

(26) Liu, Y.; Sheng, J.; Fokine, A.; Meng, G.; Shin, W.-H.; Long, F.; Kuhn, R. J.; Kihara, D.; Rossmann, M. G. Structure and Inhibition of EV-D68, a Virus That Causes Respiratory Illness in Children. *Science* **2015**, *347*, 71–74.

(27) Lane, T. R. et al. Efficacy of an Isoxazole-3-Carboxamide Analog of Pleconaril in Mouse Models of Enterovirus-D68 and Coxsackie B5. *Antiviral Research* **2023**, *216*, 105654.

(28) Rhoden, E.; Zhang, M.; Nix, W. A.; Oberste, M. S. *In Vitro* Efficacy of Antiviral Compounds against Enterovirus D68. *Antimicrobial Agents and Chemotherapy* **2015**, *59*, 7779–7781.

(29) Sun, L.; Meijer, A.; Froeyen, M.; Zhang, L.; Thibaut, H. J.; Baggen, J.; George, S.; Vernachio, J.; Van Kuppeveld, F. J. M.; Leyssen, P.; Hilgenfeld, R.; Neyts, J.; Delang, L. Antiviral Activity of Broad-Spectrum and Enterovirus-Specific Inhibitors against Clinical Isolates of Enterovirus D68. *Antimicrobial Agents and Chemotherapy* **2015**, *59*, 7782–7785.

(30) Wildman, S. A.; Crippen, G. M. Prediction of Physicochemical Parameters by Atomic Contributions. *Journal of Chemical Information and Computer Sciences* **1999**, *39*, 868–873.

(31) Liu, Y.; Sheng, J.; Van Vliet, A. L. W.; Buda, G.; Van Kuppeveld, F. J. M.; Rossmann, M. G. Molecular Basis for the Acid-Initiated Uncoating of Human Enterovirus D68. *Proceedings of the National Academy of Sciences* **2018**, *115*.

(32) Oberste, M. S.; Maher, K.; Schnurr, D.; Flemister, M. R.; Lovchik, J. C.; Peters, H.; Sessions, W.; Kirk, C.; Chatterjee, N.; Fuller, S.; Hanauer, J. M.; Pallansch, M. A. Enterovirus 68 Is Associated with Respiratory Illness and Shares Biological Features with Both the Enteroviruses and the Rhinoviruses. *Journal of General Virology* **2004**, *85*, 2577–2584.

(33) Pevear, D. C.; Tull, T. M.; Seipel, M. E.; Groarke, J. M. Activity of Pleconaril against Enteroviruses. *Antimicrobial Agents and Chemotherapy* **1999**, *43*, 2109–2115.

(34) Rogers, J. M.; Diana, G. D.; McKinlay, M. A. In *Antiviral Chemotherapy 5*; Mills, J., Volberding, P. A., Corey, L., Eds.; Springer US: Boston, MA, 1999; Vol. 458; pp 69–76.

(35) Romero, J. R. Pleconaril: A Novel Antipicornaviral Drug. *Expert Opinion on Investigational Drugs* **2001**, *10*, 369–379.

(36) Senior, K. FDA Panel Rejects Common Cold Treatment. *The Lancet Infectious Diseases* **2002**, *2*, 264.

(37) Anasir, M. I.; Zarif, F.; Poh, C. L. Antivirals Blocking Entry of Enteroviruses and Therapeutic Potential. *Journal of Biomedical Science* **2021**, *28*, 10.

(38) Egorova, A.; Ekins, S.; Schmidtke, M.; Makarov, V. Back to the Future: Advances in Development of Broad-Spectrum Capsid-Binding Inhibitors of Enteroviruses. *European Journal of Medicinal Chemistry* **2019**, *178*, 606–622.

(39) Colonno, R. J.; Condra, J. H.; Mizutani, S.; Callahan, P. L.; Davies, M. E.; Murcko, M. A. Evidence for the Direct Involvement of the Rhinovirus Canyon in Receptor Binding. *Proceedings of the National Academy of Sciences* **1988**, *85*, 5449–5453.

(40) Rossmann, M. G.; Arnold, E.; Erickson, J. W.; Frankenberger, E. A.; Griffith, J. P.; Hecht, H.-J.; Johnson, J. E.; Kamer, G.; Luo, M.; Mosser, A. G.; Rueckert, R. R.; Sherry, B.; Vriend, G. Structure of a Human Common Cold Virus and Functional Relationship to Other Picornaviruses. *Nature* **1985**, *317*, 145–153.

(41) Rossmann, M. G. The Canyon Hypothesis. *Journal of Biological Chemistry* **1989**, *264*, 14587–14590.

(42) Greve, J. M.; Davis, G.; Meyer, A. M.; Forte, C. P.; Yost, S. C.; Marlor, C. W.; Kamarck, M. E.; McClelland, A. The Major Human Rhinovirus Receptor Is ICAM-1. *Cell* **1989**, *56*, 839–847.

(43) Staunton, D. E.; Merluzzi, V. J.; Rothlein, R.; Barton, R.; Marlin, S. D.; Springer, T. A. A Cell Adhesion Molecule, ICAM-1, Is the Major Surface Receptor for Rhinoviruses. *Cell* **1989**, *56*, 849–853.

(44) Tomassini, J. E.; Graham, D.; DeWitt, C. M.; Lineberger, D. W.; Rodkey, J. A.; Colonno, R. J. cDNA Cloning Reveals That the Major Group Rhinovirus Receptor on HeLa Cells Is Intercellular Adhesion Molecule 1. *Proceedings of the National Academy of Sciences* **1989**, *86*, 4907–4911.

(45) Wei, W.; Guo, H.; Chang, J.; Yu, Y.; Liu, G.; Zhang, N.; Willard, S. H.; Zheng, S.; Yu, X.-F. ICAM-5/Telencephalin Is a Functional Entry Receptor for Enterovirus D68. *Cell Host & Microbe* **2016**, *20*, 631–641.

(46) Yamayoshi, S.; Yamashita, Y.; Li, J.; Hanagata, N.; Minowa, T.; Takemura, T.; Koike, S. Scavenger Receptor B2 Is a Cellular Receptor for Enterovirus 71. *Nature Medicine* **2009**, *15*, 798–801.

(47) Baggen, J. et al. Enterovirus D68 Receptor Requirements Unveiled by Haploid Genetics. *Proceedings of the National Academy of Sciences* **2016**, *113*, 1399–1404.

(48) Imamura, T.; Okamoto, M.; Nakakita, S.-i.; Suzuki, A.; Saito, M.; Tamaki, R.; Lupisan, S.; Roy, C. N.; Hiramatsu, H.; Sugawara, K.-e.; Mizuta, K.; Matsuzaki, Y.; Suzuki, Y.; Oshitani, H. Antigenic and Receptor Binding Properties of Enterovirus 68. *Journal of Virology* **2014**, *88*, 2374–2384.

(49) Liu, Y.; Sheng, J.; Baggen, J.; Meng, G.; Xiao, C.; Thibaut, H. J.; Van Kuppeveld, F. J. M.; Rossmann, M. G. Sialic Acid-Dependent Cell Entry of Human Enterovirus D68. *Nature Communications* **2015**, *6*, 8865.

(50) Baggen, J.; Thibaut, H. J.; Strating, J. R. P. M.; Van Kuppeveld, F. J. M. The Life Cycle of Non-Polio Enteroviruses and How to Target It. *Nature Reviews Microbiology* **2018**, *16*, 368–381.

(51) Grizer, C. S.; Messacar, K.; Mattapallil, J. J. Enterovirus-D68 – a Reemerging Non-Polio Enterovirus That Causes Severe Respiratory and Neurological Disease in Children. *Frontiers in Virology* **2024**, *4*, 1328457.

(52) Hu, Y.; Musharrafieh, R.; Zheng, M.; Wang, J. Enterovirus D68 Antivirals: Past, Present, and Future. *ACS Infectious Diseases* **2020**, *6*, 1572–1586.

(53) Roux, H. F. G.; Touret, F.; Rathelot, P.; Sciò, P.; Coluccia, A.; Vanelle, P.; Roche, M. Non-Polio Enterovirus Inhibitors: Scaffolds, Targets, and Potency-What's New? *ACS Infectious Diseases* **2025**, *11*, 21–46.

(54) Aviragen Antiviral Drug Vapendavir Fails Phase IIb Study in Asthma Patients. https://www.genengnews.com/news/aviragen-antiviral-drug-vapendavir-fails-phase-iib-study-in-asthma-patients/.

(55) Feil, S. C.; Hamilton, S.; Krippner, G. Y.; Lin, B.; Luttick, A.; McConnell, D. B.; Nearn, R.; Parker, M. W.; Ryan, J.; Stanislawski, P. C.; Tucker, S. P.; Watson, K. G.; Morton, C. J. An Orally Available 3-Ethoxybenzisoxazole Capsid Binder with Clinical Activity against Human Rhinovirus. *ACS Medicinal Chemistry Letters* **2012**, *3*, 303–307.

(56) Da Costa, L.; Scheers, E.; Coluccia, A.; Casulli, A.; Roche, M.; Di Giorgio, C.; Neyts, J.; Terme, T.; Cirilli, R.; La Regina, G.; Silvestri, R.; Mirabelli, C.; Vanelle, P. Structure-Based Drug Design of Potent Pyrazole Derivatives against Rhinovirus Replication. *Journal of Medicinal Chemistry* **2018**, *61*, 8402–8416.

(57) De Colibus, L. et al. More-Powerful Virus Inhibitors from Structure-Based Analysis of HEV71 Capsid-Binding Molecules. *Nature Structural & Molecular Biology* **2014**, *21*, 282–288.

(58) Egorova, A.; Kazakova, E.; Jahn, B.; Ekins, S.; Makarov, V.; Schmidtke, M. Novel Pleconaril Derivatives: Influence of Substituents in the Isoxazole and Phenyl Rings on the Antiviral Activity against Enteroviruses. *European Journal of Medicinal Chemistry* **2020**, *188*, 112007.

(59) Ke, Y.-Y.; Lin, T.-H. Modeling the Ligand-Receptor Interaction for a Series of Inhibitors of the Capsid Protein of Enterovirus 71 Using Several Three-Dimensional Quantitative Structure-Activity Relationship Techniques. *Journal of Medicinal Chemistry* **2006**, *49*, 4517–4525.

(60) Ma, C.; Hu, Y.; Zhang, J.; Musharrafieh, R.; Wang, J. A Novel Capsid Binding Inhibitor Displays Potent Antiviral Activity against Enterovirus D68. *ACS Infectious Diseases* **2019**, *5*, 1952–1962.

(61) Makarov, V. A.; Riabova, O. B.; Granik, V. G.; Wutzler, P.; Schmidtke, M. Novel [(Biphenyloxy)Propyl]Isoxazole Derivatives for Inhibition of Human Rhinovirus 2 and Coxsackievirus B3 Replication. *Journal of Antimicrobial Chemotherapy* **2005**, *55*, 483–488.

(62) Muratov, E. N.; Varlamova, E. V.; Artemenko, A. G.; Khristova, T.; Kuz'min, V. E.; Makarov, V. A.; Riabova, O. B.; Wutzler, P.; Schmidtke, M. Qsar Analysis of [(Biphenyloxy)Propyl]Isoxazoles: Agents Against Coxsackievirus B3. *Future Medicinal Chemistry* **2011**, *3*, 15–27.

(63) Schmidtke, M.; Wutzler, P.; Zieger, R.; Riabova, O. B.; Makarov, V. A. New Pleconaril and [(Biphenyloxy)Propyl]Isoxazole Derivatives with Substitutions in the Central Ring Exhibit Antiviral Activity against Pleconaril-Resistant Coxsackievirus B3. *Antiviral Research* **2009**, *81*, 56–63.

(64) Shia, K.-S.; Li, W.-T.; Chang, C.-M.; Hsu, M.-C.; Chern, J.-H.; Leong, M. K.; Tseng, S.-N.; Lee, C.-C.; Lee, Y.-C.; Chen, S.-J.; Peng, K.-C.; Tseng, H.-Y.; Chang, Y.-L.; Tai, C.-L.; Shih, S.-R. Design, Synthesis, and Structure-Activity Relationship of Pyridyl Imidazolidinones: A Novel Class of Potent and Selective Human Enterovirus 71 Inhibitors. *Journal of Medicinal Chemistry* **2002**, *45*, 1644–1655.

(65) Zhang, M.; Wang, Y.; He, W.; Sun, Y.; Guo, Y.; Zhong, W.; Gao, Q.; Liao, M.;

Wang, X.; Cai, Y.; Guo, Y.; Rao, Z. Design, Synthesis, and Evaluation of Novel Enterovirus 71 Inhibitors as Therapeutic Drug Leads for the Treatment of Human Hand, Foot, and Mouth Disease. *Journal of Medicinal Chemistry* **2020**, *63*, 1233–1244.

(66) Braun, H.; Makarov, V. A.; Riabova, O. B.; Wutzler, P.; Schmidtke, M. Amino Acid Substitutions At Residue 207 of Viral Capsid Protein 1 (VP1) Confer Pleconaril Resistance in Coxsackievirus B3 (CVB3). *Antiviral Research* **2011**, *90*, A54–A55.

(67) Ledford, R. M.; Collett, M. S.; Pevear, D. C. Insights into the Genetic Basis for Natural Phenotypic Resistance of Human Rhinoviruses to Pleconaril. *Antiviral Research* **2005**, *68*, 135–138.

(68) Volobueva, A.; Egorova, A.; Galochkina, A.; Ekins, S.; Zarubaev, V.; Makarov, V. The Evolution of Pleconaril: Modified O-Alkyl Linker Analogs Have Biological Activity towards Coxsackievirus B3 Nancy. *Molecules* **2020**, *25*, 1345.

(69) Hayden, F. G.; Herrington, D. T.; Coats, T. L.; Kim, K.; Cooper, E. C.; Villano, S. A.; Liu, S.; Hudson, S.; Pevear, D. C.; Collett, M.; McKinlay, M.; Pleconaril Respiratory Infection Study Group Efficacy and Safety of Oral Pleconaril for Treatment of Colds Due to Picornaviruses in Adults: Results of 2 Double-Blind, Randomized, Placebo-Controlled Trials. *Clinical Infectious Diseases* **2003**, *36*, 1523–1532.

(70) Dickins, M. Induction of Cytochromes P450. *Current Topics in Medicinal Chemistry* **2004**, *4*, 1745–1766.

(71) Tompkins, L. M.; Wallace, A. D. Mechanisms of Cytochrome P450 Induction. *Journal of Biochemical and Molecular Toxicology* **2007**, *21*, 176–181.

(72) Fletcher, S. The Mitsunobu Reaction in the 21st Century. *Organic Chemistry Frontiers* **2015**, *2*, 739–752.

(73) Mitsunobu, O.; Yamada, M. Preparation of Esters of Carboxylic and Phosphoric Acid *via* Quaternary Phosphonium Salts. *Bulletin of the Chemical Society of Japan* **1967**, *40*, 2380–2382.

(74) Abdel-Rahman, S. M.; Kearns, G. L. Single-Dose Pharmacokinetics of a Pleconaril (VP63843) Oral Solution and Effect of Food. *Antimicrobial Agents and Chemotherapy* **1998**, *42*, 2706–2709.

(75) Abdel-Rahman, S. M.; Kearns, G. L. Single Oral Dose Escalation Pharmacokinetics of Pleconaril (VP 63843) Capsules in Adults. *The Journal of Clinical Pharmacology* **1999**, *39*, 613–618.

(76) Diana, G. D. et al. Picornavirus Inhibitors: Trifluoromethyl Substitution Provides a Global Protective Effect against Hepatic Metabolism. *Journal of Medicinal Chemistry* **1995**, *38*, 1355–1371.

(77) Kenny, P. W. Hydrogen-Bond Donors in Drug Design. *Journal of Medicinal Chemistry* **2022**, *65*, 14261–14275.

(78) Lowe, D. More Magic Methyls, Please. https://www.science.org/content/blog-post/more-magic-methyls-please.

(79) Schönherr, H.; Cernak, T. Profound Methyl Effects in Drug Discovery and a Call for New C-H Methylation Reactions. *Angewandte Chemie International Edition* **2013**, *52*, 12256–12267.

(80) Hopkins, A. L.; Keserü, G. M.; Leeson, P. D.; Rees, D. C.; Reynolds, C. H. The Role of Ligand Efficiency Metrics in Drug Discovery. *Nature Reviews Drug Discovery* **2014**, *13*, 105–121.

(81) Johnson, T. W.; Gallego, R. A.; Edwards, M. P. Lipophilic Efficiency as an Important Metric in Drug Design. *Journal of Medicinal Chemistry* **2018**, *61*, 6401–6420.

(82) Sauer, W. H. B.; Schwarz, M. K. Molecular Shape Diversity of Combinatorial Libraries: A Prerequisite for Broad Bioactivity. *Journal of Chemical Information and Computer Sciences* **2003**, *43*, 987–1003.