Heartbeat-evoked responses in M/EEG: A systematic review of methods with suggestions for analysis and reporting

Paul Steinfath[1,2*#], Maria Azanova[1,3#], Nikolai Kapralov[1,2#], Thomas Loesche[1], Lioba Enk[1,3], Vadim Nikulin[1##], Arno Villringer[1,3,4,5,6##]

[1]Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany

[2]International Max Planck Research School NeuroCom, Leipzig, Germany

[3]Max Planck School of Cognition, Leipzig, Germany

[4]LIFE – Leipzig Research Center for Civilization Diseases, University of Leipzig, Leipzig, Germany

[5]Department of Cognitive Neurology, University Hospital Leipzig, Leipzig, Germany

[6]MindBrainBody Institute, Berlin School of Mind and Brain, Humboldt University Berlin, Berlin, Germany

[#,##]These authors contributed equally to this work

*Corresponding author: steinfath@cbs.mpg.de,

Department of Neurology, Max Planck Institute for Human Cognitive and Brain Sciences, Leipzig, Germany, Stephanstr. 1a, 04103, Leipzig, Germany

Abstract

Heartbeat-evoked responses (HER), as measured by electroencephalography (EEG) or magnetoencephalography (MEG), represent neural activity time-locked to heartbeats and are widely used as a marker of cardiac interoception in the study of brain-body interactions. However, HER studies report largely variable findings, at least partially due to methodological variability. To achieve consensus on HER processing and improve the reproducibility of findings, the field urgently requires a structured summary of the methods employed so far. To this end, we conducted a systematic review of 132 HER studies using non-invasive M/EEG recordings in humans. Our results reveal substantial heterogeneity across most steps of HER analysis, ranging from data acquisition and preprocessing to HER estimation and statistical approaches. The large diversity in the processing choices is accompanied by considerable proportions of unreported methodological information across reviewed studies, reaching up to 80% for key processing steps. In addition, less than 33% of studies had enough statistical power to reliably detect meta-level HER effects, while their reported spatiotemporal locations varied substantially. We provide a comprehensive reporting and quality control checklist to aid in the development of more standardized procedures, highlighting critical steps for robust HER investigations. Additionally, we share the full extracted dataset, including an interactive version, to support other researchers in answering additional specific questions they may have. We hope that these resources will improve the robustness, reproducibility, and transparency of research in the growing HER field.

Keywords

Heartbeat-evoked response, heartbeat-evoked potential, methodological variability, reporting guidelines, MEG, EEG, ECG

1. Introduction

The dynamic interplay between the central nervous system and other bodily organs is increasingly recognized as a crucial factor in understanding cognition, behavior, and overall well-being (Azzalini et al., 2019; Criscuolo et al., 2022; Villringer et al., 2025). A growing body of research has demonstrated how the interaction between the brain and body shapes many functional aspects, ranging from the direct modulation of neuronal activity by blood pressure (Jammal Salameh et al., 2024) up to higher cognition (Umeda et al., 2016) and complex emotional states (Feldman et al., 2024; Garfinkel et al., 2014).

Among the various organs involved, the heart plays a particularly important role in brain-body interaction research. Beyond its essential function of delivering oxygenated blood and vital nutrients to the brain and the rest of the body (Hall & Hall, 2021), its rhythmic activity itself appears to serve as a pacemaker, providing an important reference frame for the brain (Tallon-Baudry et al., 2018). Accumulating evidence suggests that distinct phases of the cardiac cycle are advantageous for either perception or action; therefore, monitoring and adjustment of cardiac activity by the brain might be necessary to achieve optimal performance (Galvez-Pol et al., 2022; Skora et al., 2022).

One key methodology for studying heart-brain interactions with electroencephalography (EEG) or magnetoencephalography (MEG) involves analyzing neural activity time-locked to the heartbeat. The resulting heartbeat-evoked responses[1] (HERs) reflect the processing of heartbeat-related information in the central nervous system (Dirlich et al., 1997; Schandry et al., 1986). HERs are commonly used as a marker for cardiac interoception (Pollatos & Schandry, 2004) and have been linked to a broad range of functions, such as perception (Al

---

[1] In the EEG field, these responses are commonly referred to as heartbeat-evoked potentials (HEPs) due to the nature of the electric potentials recorded with EEG. MEG studies, however, refer to them as heartbeat-evoked fields (HEFs) or heartbeat-evoked responses (HERs). Since we include both EEG and MEG studies in this review, we use the more general term 'response' (HER) throughout the text.

et al., 2020; Park et al., 2014), attention (Banellis & Cruse, 2020), motor activity (Al et al.,

2023), or emotions (Di Bernardi Luft & Bhattacharya, 2015).

Despite the growing interest and number of studies, the field of HER research is facing a

challenge of methodological heterogeneity and variability in reported findings (Coll et al.,

2021). There appears to be no single, unified manifestation of HER. Instead, heartbeat-locked

neural activity has been associated with a wide range of cognitive and emotional functions

and observed across diverse topographic and temporal regions (Engelen, Solcà, et al., 2023;

Park & Blanke, 2019). While some of this variability can be attributed to participant

characteristics (e.g., patients or controls) or the experimental context in which HERs are

investigated, variations in data acquisition, preprocessing techniques, and statistical analyses

likely also play a role.

Beyond the general methodological decisions that apply to all studies of evoked responses

(such as how to filter the data and whether to use baseline correction), analysis of HERs

requires special treatment, most notably due to the cardiac field artifact (CFA). CFA is caused

by the instantaneous spread of magnetic and electric fields from the heart's activity due to

volume conduction, and is directly captured by M/EEG sensors (Dirlich et al., 1997). In

classical task-based studies, the CFA is widely believed to play a minor role, as cardiac-related

noise is not time-locked to stimuli and thus suppressed by trial averaging. However, due to

the inherent coupling between heartbeat and CFA, it is not reduced by averaging across trials

when obtaining HERs and poses a serious challenge for HER research. To investigate the

brain activity related to the heartbeat, without the CFA, researchers typically either try to avoid

time ranges corresponding to high CFA, remove the CFA from the data, or include the heart's

activity as a confound in the statistical analysis. Various approaches are employed for this

purpose, and the outcome may depend on the selected approach and even on decisions such

as the choice of settings for electrocardiogram (ECG) recording (Gray et al., 2007). To note,

other confounds, such as pulse-related movement of tissue and sensors, can also introduce

heartbeat-locked artifacts (Kern et al., 2013), but their influence on measured HERs remains
unclear.

Apart from cardiac artifacts, the overlap of heartbeat- and task-evoked activity additionally
complicates the analysis of HERs. When HERs are studied in task paradigms, the potential
overlap with task-evoked activity must be accounted for (Azzalini et al., 2019; Steinfath et al.,
2025). Furthermore, due to the cyclic nature of the heartbeat, neural responses to successive
heartbeats can superimpose, especially when inter-beat intervals (IBIs) are short. The
necessity to address these problems further increases the number of analysis decisions
researchers have to make.

Overall, it remains unclear to what extent methodological choices influence the HER effects
reported across different studies, and thorough examination of the employed methods has
been repeatedly advocated (Coll et al., 2021; Park & Blanke, 2019). This situation mirrors the
reproducibility challenge in the broader field of neuroscience (Botvinik-Nezer & Wager, 2023),
which is currently emphasized and addressed by newly appearing analytic approaches such
as large-scale collaborative data analyses (EEGmanypipelines, Trübutschek et al., 2024),
multiverse analyses (Clayson et al., 2021), or meta-methods reviews (Elson, 2019).

In light of these challenges, it is an essential first step to systematically collect and assess the
methodological approaches used in HER research. To achieve improved consistency and
reproducibility in HER analysis, a clear understanding of current practices is necessary. This
review, therefore, focuses specifically on the methods used in HER research. To this end, we
extracted and analyzed the methodological decisions, from data acquisition through
preprocessing to statistical analysis, from published HER studies.

In addition, we highlight aspects that may be especially important for the HER analysis, which
we hope will stimulate further HER research, discussion, increased transparency, and
potential consensus. To facilitate this, we publicly share the full resulting dataset, its interactive
version, the accompanying codebook, and the HER analysis checklist with this article.

2. Methods

2.1. Literature search

We conducted a systematic literature search using the PubMed database on three separate dates to identify studies investigating HERs using noninvasive neuroimaging. Table 1 specifies the search terms used, with corresponding dates and the number of studies retrieved, as well as the total number of studies after duplicate removal.

Table 1. Search strategy

Date	Search query	Studies retrieved
15.01.2024	(eeg OR electroencephalograph*) AND (heartbeat evoked potential OR heartbeat evoked response) NOT (Review[Publication Type] OR Systematic Review[Publication Type]) AND English[Language]	647
24.07.2024 (rerun: 05.08.2024)	(eeg OR electroencephalograph* OR meg OR magnetoencephalograph*) AND (heartbeat evoked potential OR heartbeat evoked response) NOT (Review[Publication Type] OR Systematic Review[Publication Type]) AND English[Language]	682 (rerun: 683)
05.08.2024	("heartbeat evoked" OR "heart evoked") AND Journal Article[Publication Type] AND English[Language]	166
Total amount of unique studies after removing duplicate entries		**749**

After the removal of duplicates, all retrieved records (N = 749) were screened following the guidelines of the Preferred Reporting Items for Systematic Reviews and Meta-Analyses

(PRISMA; Moher et al., 2009). We included studies that: (1) examined HERs, (2) applied noninvasive EEG or MEG recordings, (3) were conducted in human participants, (4) reported original data, (5) were in English language, and (6) had full text available. Exclusion criteria were: (1) intracranial recordings (due to the different data analysis approaches), (2) non-original research (reviews, commentary papers, and toolboxes), (3) studies not analyzing evoked responses (e.g., time-frequency analysis only), and (4) lack of a peer review. Because this review focuses on HER analysis methods, we also included studies analyzing data in source space and those without statistical testing. After screening titles and abstracts, we evaluated the full texts of potentially relevant articles to determine their eligibility. After applying the above-mentioned criteria, 127 articles were included. In addition, we identified seven references through the bibliographies of the reviewed publications, of which five were included, resulting in a final sample of 132 articles used in the systematic review. The screening process is detailed in the PRISMA flow diagram (Fig. 1A).

2.2. Extraction of methodological choices

In the following, we use the term *processing steps* when referring to the various data transformations or parameters that researchers choose (for example, the cutoff frequency of a high-pass filter). A particular value (e.g., 1 Hz) that is then selected for a processing step is referred to as a *(methodological) choice*, and a combination of methodological choices for all processing steps is considered as a *pipeline* in this review. If a study uses different methodological choices for any processing step (e.g., performs the analysis with and without baseline correction), we say that several pipelines correspond to this same study.

For each article included in the review, we extracted methodological choices for a wide range of processing steps, from the acquisition of physiological data (EEG/MEG/ECG) to the calculation and statistical analysis of HERs (Table 2). Three reviewers, independently, extracted the data and then additionally checked for the consistency of the extracted information for each processing step. We developed a standardized codebook to unify and

ensure consistency of labels referring to the same methodological categories (e.g., "linked

mastoids" and "averaged mastoids"). The codebook is publicly available and can be found

together with the final dataset at https://osf.io/znrbm.

Figure 1. Overview of the studies included in the review. (A) Flowchart describing the literature

selection process according to the PRISMA guidelines. (B) The majority of the included studies

were published within the last 10 years. (C) Most studies use EEG data to analyze heartbeat-

evoked responses (HERs). Diamond indicates that most MEG studies come from one

research group. (D) HERs are typically analyzed during task performance rather than in the

resting state.

Table 2. Overview of the extracted processing steps from pipelines used for studying HERs.

Abbreviations: HERs – heartbeat-evoked responses, M/EEG – magneto- or

electroencephalography, ECG – electrocardiogram, ICA – independent component analysis,

ICs – independent components, CFA – cardiac field artifact.

Category	Processing step
Study	Research topic, whether new data or an existing dataset were analyzed
Experiment	Experimental setting (task or resting state), sample size, whether patients of any disorder are present among participants, age group(s), number of participant groups, number of experimental conditions, length of the recording used to analyze HERs, number of epochs used to obtain HERs (per participant and condition)
M/EEG acquisition	Imaging modality (M/EEG), number and locations (layout) of M/EEG sensors, number of gradiometers and magnetometers (MEG only), reference electrode used during the recording (online reference; EEG only), sampling frequency
ECG acquisition	Number and locations of ECG electrodes, ECG lead, ECG ground electrode, sampling frequency
ECG preprocessing	Cutoff frequencies for the low- and high-pass filters, sampling frequency (in case of downsampling), method, and, if used, toolbox for detecting reference events (R/T)
M/EEG preprocessing	Reference electrode used in the data analysis (offline reference; EEG only), sampling frequency (in case of downsampling), cutoff frequencies for the low- and high-pass filters, whether ICA was applied, type of ICs that were rejected, other approaches for cleaning the data
Removal of the CFA	Whether ICA was applied on epoched or continuous data, the number of rejected CFA-related ICs, the approach and criteria

	used for rejecting CFA-related ICs, other approaches for removal of CFA apart from ICA
HER computation	Approach to determine locations and times of interest (primary/secondary), epoch onset (R-/T-peak), start and end latencies of the analyzed time window, start and end latencies of the baseline window, selected M/EEG sensors, extracted value, whether channels or time points were averaged
Statistical analysis	Statistical test, type of association tested (difference or correlation), whether cluster-based permutation test was applied, number of permutations, control variables, whether the result of the test was significant
Clusters (in case of a significant test)	Epoch onset (R-/T-peak), channels, and time-points that belong to the cluster

2.3. Data analysis

Data analysis and visualization were performed using R (version 4.4.3; R Core Team, 2025). For most processing steps, we assessed the overall scope of methodological choices and the proportion of studies or pipelines that utilized each choice. Since the information relevant for this review is not reported for some studies and pipelines, we specify the exact number of data points above each plot. Some studies perform several analyses with different selections of methods — if this is the case for a particular plot, we refer to the data points as pipelines; otherwise, we refer to them as studies. Importantly, the number of studies/pipelines using a specific methodological choice can increase not only because of community consensus, but also when the same research group repeats acquisition or analysis settings across multiple

studies. To address it in histogram plots, we add a diamond symbol above choices, for which

the same first or last author contributes at least 50% of the occurrences.

Entropy. To quantify and compare the variability of methodological choices for different

processing steps, we used the concept of Shannon entropy (Shannon, 1948). Let *k* denote

the number of possible choices, while $p_1, p_2, \ldots, p_k$ stand for their occurrence proportions across

studies or pipelines. Then, Shannon entropy is computed as:

$$H = -\sum_{i=1}^{k} p_i \cdot log_2(p_i) \tag{1}$$

185 Larger entropy values indicate a greater heterogeneity in the choices, while entropy is zero

186 when all studies use the same method. As the maximum possible entropy depends on the

187 number of available options, we normalized it to allow for comparisons across processing

188 steps with different numbers of choices:

$$H_{norm} = H / log_2(k) \tag{2}$$

Additionally, we binned continuous parameters (e.g., the number of trials) that contained more

than 10 unique values to make them comparable to categorical variables before computing

entropy.

Aggregation of EEG locations. To analyze the spatial distribution of observed HER effects, we

collected information about the EEG channels included in the HER analysis and, if applicable,

about those belonging to at least one of the clusters reported for significant tests. To combine

studies that used different EEG layouts during data recording, we calculated the number of

occurrences for each channel and layout separately. The occurrence counts were interpolated

to form topographies independent of actual sensor locations. The topographies were then

summed across the considered layouts, that is, variations of the 10-20 system with 19, 32,

and 64 channels, as well as the 128-channel BioSemi layout.

HER peak timing comparison. To statistically compare the timing of effects observed with different HER analysis strategies (e.g., clustering vs. averaging), we implemented a permutation test. For each strategy, the distribution of analysis time window start and end times was aggregated as a cumulative sum, and the peak was identified. Next, the peak time difference between the two strategies was calculated. To test if the differences between strategies were statistically significant, all analysis time windows from both strategies were pooled and randomly assigned to two groups of the original sizes. The peak timing differences were recalculated, and the permutation procedure was repeated 1000 times. Lastly, p-values were obtained by comparing the original peak timing difference to the permutation distribution.

Definition of cluster-based tests. Note that throughout the manuscript, by "cluster-based permutation tests" (also "permutation clustering", "cluster-based analysis", and similar) we specifically refer to cluster-sum tests, which establish significance based on the summed statistic (e.g., t-values) across spatiotemporal points within a cluster (Maris & Oostenveld, 2007). While this approach is currently popular in the M/EEG field, it comes with the limitation that the precise timing and spatial extent of clusters should be interpreted with caution (Sassenhagen & Draschkow, 2019). Other cluster-based approaches exist (Frossard & Renaud, 2022; Rousselet, 2025), but, due to their novelty and lack of implementation in neuroscientific toolboxes, their prevalence in HER research was not assessed in this review.

Minimal detectable effect sizes. To compare minimal effect sizes that could be robustly detected across pipelines, we estimated the smallest effect that a study can reliably identify given a specified level of confidence. Such estimates provide a practical way to assess whether a study is sufficiently powered for the effects it aims to detect. Accordingly, we (1) computed minimal detectable effect size for one- and two-sample t-tests, ANOVAs, and correlations using pwr R package (version 1.3-0; Champely et al., 2020) based on the information about sample sizes, numbers of groups, conditions, and required power of 80% with significance level set at 5%, (2) converted these effect sizes to Cohen's d, when required, and finally (3) converted Cohen's d into Hedges' g using the exact correction approach

(Hedges & Olkin, 1985). Hedges' g was used instead of Cohen's d because it provides a less

biased effect size estimate for small sample sizes, which are typical for HER studies. The

resulting Hedges' g estimates of minimal detectable effect size were compared to previously

reported effect sizes in HER literature (Coll et al., 2021). Details on the estimation of Hedges'

g are provided in Section A of the Supplementary Material. Finally, we examined how

detectable effect sizes could change if within-participant variability in HER values across

epochs was taken into account (see Section B of the Supplementary Material for more details

on these estimations).

Estimation of the number of epochs. As M/EEG signals are prone to high amounts of

environmental and physiological noise, obtaining a clear evoked response often requires

averaging over a substantial number of epochs. For each pipeline, we extracted the minimum

(preferred) or mean number of heartbeat-locked epochs per participant and condition, where

possible. Those were not readily available in some studies. For such cases, we estimated

approximate numbers using strategies such as (1) dividing the overall reported number of

epochs by the number of analyzed participants, (2) multiplying the reported length of the

recording in minutes within which HERs were analyzed by the average heart rate, when

reported, or otherwise by 60 beats per minute. If reported, those were adjusted by the number

of epochs removed during cleaning.

3. Results

The results of the review are structured as follows. We start with an overview of included

studies (Section 3.1), followed by an overview of processing steps that were extracted and

analyzed (Section 3.2). Sections 3.3-3.9 provide a detailed description of methodological

choices that occur in the literature for various processing steps from data acquisition to

statistical analysis.

3.1. Overview of the included studies

In total, 132 studies published between 1986 and 2024 were included in the review (Fig. 1B). The vast majority of studies used EEG recordings (123/132, 93%), while only nine studies (6.8%) employed MEG to measure HERs (Fig. 1C). As shown in Fig. 1D, most studies (93/132; 70%) focused on the analysis of HERs during performance of various tasks involving perception of own heartbeats (see Coll et al., 2021 for a review) or external stimuli (e.g., Al et al., 2020; Marshall et al., 2020; Park et al., 2014). A smaller proportion of studies (32/132; 24%) performed resting-state analyses, testing the potential relationship between the amplitude of HER and diverse psycho-physiological states (e.g., Flasbeck et al., 2021; Giusti et al., 2024; Yoris et al., 2024) as well as diverse clinical conditions (e.g., Birba et al., 2022; Kumral et al., 2022; Schmitz et al., 2021; Solcà et al., 2020). Seven studies (5.3%) analyzed HERs both during task and resting state.

The included studies were published in a wide range of journals (Table S2) and varied in the patient or age groups studied, research topics, imaging modalities, and specific HER research groups. All these factors likely contribute to the variability in methods and findings that we document in this review and may affect some processing choices more than others. We invite readers to use the provided interactive resource (https://paulsteinfath.shinyapps.io/her-systematic-review/) to investigate the importance of these factors, which are outside the scope of the current review.

3.2. Overview of methodological choices

For each processing step of Table 2, we extracted methodological choices made by researchers (e.g., which value of a parameter to use or whether to apply a certain method). In total, we extracted 410 pipelines (note that some studies included multiple pipelines due to the use of different parameters or strategies). In the subsequent sections, we only consider distinct pipelines with respect to each processing step we discuss. For example, when discussing preprocessing steps, multiple pipelines that differ only in terms of statistical

analysis are treated as one. This should ensure that our analyses reflect methodological

variability within each processing step without being inflated by differences in unrelated steps.

The total number of pipelines (or studies, when only one pipeline per study was reported) is

specified above each figure panel or in parentheses after each reported value. Note that

several resulting pipelines may come from one laboratory (max. 12 studies from the same last

author) or from re-analyses of the same dataset (max. 2 studies).

Based on the information we extracted for data processing steps, from acquisition to statistics,

we observe a substantial lack of consensus in most methodological choices. We quantify this

heterogeneity for each processing step using normalized Shannon entropy (Eq. 2), where

values close to 1 indicate high variability in choices, and values closer to 0 indicate a single

dominant choice. Most steps showed large entropy (Fig. 2A). For example, frequency cutoffs

of high- and low-pass filters used in preprocessing show great variability (Fig. 3E). Similarly,

researchers select widely varying time windows of interest for their HER analysis (see Section

3.7 for more details). Few steps show consistency across the field: most studies use EEG

data, select the R-peak as the reference event for time-locking HERs, and analyze HER

amplitude.

During acquisition and preprocessing, most studies apply exactly one choice per step, e.g.,

they either perform ICA for artifact correction or do not, but rarely both (Fig. 2B). In contrast,

using several approaches for HER computation and statistical analysis within the same study

is quite common: 56% of studies (74/132) use multiple statistical tests, and 45% (59/132) vary

the start or the end latency of the time window of interest. Many such cases involve primary

and secondary data analysis, or several related hypotheses within the same study, and

therefore, typically do not require additional statistical correction.

However, almost no study systematically compares several methodological choices per

processing step to assess the effects of researcher-dependent decisions on HER results.

Among the reviewed studies, Candia-Rivera, Catrambone, et al. (2021) is a single such study

that varies a preprocessing step (offline reference). Future studies should address the

outstanding question of how processing decisions affect HER findings. Additionally, HER

research articles could show if findings are robust to decisions such as baseline correction or

different CFA cleaning methods.

Adding to that, a substantial proportion of methodological decisions is unreported (Fig. 2C). In

71% of studies reporting CFA-related independent component rejection (49/69), information

on the number of rejected components is not provided. In 80% of studies (105/132), the ECG

ground location is not explicitly stated. As natural contamination with CFA makes HERs

distinct from other event-related responses, it is critical to provide as much information as

possible about heartbeat-related processing and acquisition decisions. In addition, we observe

a high percentage of unreported values for the following critical methodological choices: the

number of epochs averaged to obtain HERs (73/132 studies, 55%), recording length (78/132,

59%), cutoff frequencies of high-pass (76/132, 58%) and low-pass (80/132, 61%) filters

applied to ECG, algorithm for detection of R-/T-peaks in the ECG (69/132, 52%), and the start

or end of the baseline window for HER (59/132, 45%). In 28% of studies (37/132), ECG

locations are not reported. Overall, these results highlight that even for parameters critical to

HER computation and analysis, there is a lack of transparency in the field, which contributes

to reduced reproducibility of results. To remedy this issue, we provide a comprehensive

reporting and quality control checklist with this review (Box 1).

Figure 2. Overview of the extracted processing steps, heterogeneity in methodological choices, and unreported information. (A) Entropy was calculated to illustrate the diversity in processing choices across studies. Within categories, the processing steps are sorted by entropy in descending order. (B) Some studies use several analysis approaches, specifically to estimate heartbeat-evoked response (HER) values and perform the statistics. (C) The highest percentage of unreported information was observed for ECG ground location and the number of rejected cardiac independent components (ICs), a relevant parameter for artifact

correction in the HER field. Abbreviations: ICA – independent component analysis, CFA – cardiac field artifact, ECG – electrocardiography, EEG – electroencephalography, MEG – magnetoencephalography.

3.3. M/EEG acquisition and preprocessing of evoked responses

Since HERs typically occur in the 0.5–2 µV range (Pérez et al., 2005; Pollatos & Schandry, 2004; Schandry & Weitkunat, 1990), M/EEG recordings of these signals often have a relatively low signal-to-noise ratio (SNR), requiring careful data preprocessing and artifact removal strategies. At the same time, the preprocessing itself can affect the shape and amplitude of ERP components (Clayson et al., 2021). In this section, we focus on the processing steps that are common to all ERPs, while the specific methods for suppressing CFA are discussed in more detail in Section 3.5.

In EEG, the choice of the reference electrode is important as it affects the shape and distribution of the ERP (Luck, 2014), making direct comparisons across studies with different references difficult. As shown in Fig. 3A, we observe some variability with respect to the online reference used during data acquisition, with linked mastoids (23%, 28/123 EEG studies) and Cz (17%, 21/123) as the most common options. This variability is reduced during data processing, where the common average reference (CAR, Fig. 3B) emerges as the most dominant choice (45%, 57/127 pipelines). This observation is complemented by Candia-Rivera, Catrambone, et al. (2021), who compared the influence of different re-referencing schemes on HERs and concluded that CAR leads to the most consistent results across a range of brain-heart interaction measures. However, they also note that the optimal re-referencing choice depends on the experimental settings and can vary across studies.

When analyzing relatively slow evoked responses, such as the HER, optimal filter settings are an important consideration to avoid removing or inducing effects (Luck, 2014; Tanner et al., 2015). We observe that, across studies, filtering is generally done between 0.1 and 40 Hz, with the most dominant high-pass filter cutoffs at 0.1, 0.5, and 1 Hz (Fig. 3E). In standard ERP

analyses, frequencies above 40 Hz are typically excluded because they often are

contaminated with different artifacts like muscle or line noise (Luck, 2014). Optionally, the data

can be downsampled, but the final sampling frequency should be at least twice as high as the

maximal frequency present in the recordings to avoid aliasing (e.g., at least 80 Hz for a 40 Hz

low-pass filter). As shown in Fig. S3, all reviewed studies used sufficiently high sampling

frequencies (at least 100 Hz) during recording and in offline analysis, compared to the popular

choice of a 40 Hz low-pass filter.

Meanwhile, the high-pass filter cutoff must balance the removal of low-frequency noise while

preserving the slowly evolving components of the heartbeat-locked signal. Given that some

neurons track changes in inter-beat intervals (De Falco et al., 2024) and assuming the resting

heart rate in humans is approximately 1 Hz, a high-pass filter of 1 Hz might inadvertently

remove genuine HER activity. Ultimately, the filter selection represents a trade-off between

the specific aims of the HER analysis and the noise characteristics of the dataset. Future

studies are needed to shed light on the frequency components of the HER across contexts,

their relationship to heart rate, and the impact of different filter cutoffs on the observed

associations between HERs and behavior or cardiac artifacts. Approaches such as multiverse

analysis could be used for the systematic exploration of such effects (Clayson et al., 2021). In

the meantime, we recommend reporting the filter settings used to obtain HER (Box 1).

Due to the low amplitude of HERs, additional artifact-reduction strategies are often used to

clean the data and obtain robust HERs that can be compared across groups or conditions.

Our review shows that 55% of the studies (72/132) removed noisy epochs, while 11% (15/132)

explicitly mention that bad channels were excluded. The majority of pipelines (73%, 97/133;

Fig. 3C) applied independent component analysis (ICA), a commonly used decomposition

method designed to separate the recorded M/EEG signal into its underlying brain and non-

brain sources.

The primary artifact sources that are addressed by ICA include eye movement (89%, 85 out

of 96 pipelines that reported the types of rejected components), cardiac (72%, 69/96), muscle

(23%, 22/96), or other nuisance sources (Fig. 3D). While the removal of cardiac artifacts is

discussed in Section 3.5, it is important to recognize that other physiological activity such as

eye movements (Ohl et al., 2016), muscle activity (Niizeki et al., 1993), or respiration (Yasuma

& Hayano, 2004) also co-vary with the heartbeat. The extent to which these factors influence

the measured HERs, and whether ICA can mitigate their effects, remains an open question

that should be addressed in future studies.

Figure 3. Heterogeneity in M/EEG acquisition settings and preprocessing methods. (A) Linked

mastoids and Cz are most commonly used during the recording as reference. Only studies

that used EEG are shown. The category Other denotes references that appeared only in a

single study. Diamond symbols indicate choices driven by one research group (at least 50%

of occurrences). (B) Common average is the most common reference used in the data

analysis. Only studies that used EEG are shown. (C) Most studies use ICA for artifact removal.

(D) ICA components related to eye movements (ocular) and CFA are rejected most frequently.

(E) Frequency ranges analyzed by the reviewed studies. Popular high-pass filter choices

include 0.1, 0.5, or 1 Hz. Abbreviations: LinkM – linked mastoids, CMS – common mode

sense, CAR – common average reference, LinkE – linked earlobes, LM – left mastoid, LAP –

laplacian or current source density analysis, N/M – not mentioned, REST – reference electrode

standardization technique, ICA – independent component analysis, CFA – cardiac field

artifact.

403 3.4. ECG acquisition and preprocessing

ECG recordings are primarily used to extract cardiac markers for time-locking the HER

(typically, either R-peak or T-peak). At the same time, they are also often (by 59/132 studies,

45%) employed in control analyses to ensure that observed HER effects are not accompanied

and thus cannot be explained by differences in cardiac activity. Placement of the ECG

electrodes can play an important role in such analyses, as the presence of differences in ECG

between experimental conditions has been shown to depend on the ECG derivation of choice

(Gray et al., 2007). Moreover, using just one ECG lead to control for the spread of CFA might

not be sufficient to capture the three-dimensional CFA in full (Pérez et al., 2005). With these

motivations in mind, we evaluated the consistency of ECG electrode setups used in the

literature.

Most studies (67%, 89/132) dedicate two electrodes to record ECG (Fig. 4A), but the

placement of the electrodes on the body varies substantially. Although lead II is used most

frequently (by 36% of studies, 48/132), some studies also use lead I (11%, 15/132) and lead III (2.3%, 3/132), or locations that do not correspond to any of the standard leads (27%, 35/132; marked as 'N/C' in Fig. 4B). Even with standard lead configurations, electrode locations vary (e.g., torso vs. limb placement; Fig. 4C) though the significance of this variability remains unclear and may differ between clinical and non-clinical populations (Sheppard et al., 2011).

In addition, we extracted information about ECG preprocessing settings, including filtering cutoffs and methods for detection of reference events (R-/T-peak). Across the studies, high- and low-pass filter cutoffs largely mirror those used in M/EEG preprocessing, with the most common high-pass cutoffs being 0.1, 0.5, and 1 Hz (Fig. 4D). Especially if researchers plan to use the ECG in statistical control analyses, it may be advisable to match the filter settings across M/EEG and ECG.

Peak detection is typically performed automatically and often followed by visual inspection of the results. As shown in Fig. 4E and 4F, the choice of the method and toolbox for peak detection is highly dependent on the research group. Most common algorithms can be grouped into three major families based on their core principle and target features (Fig. 4E): thresholding of the ECG derivative (Pan & Tompkins, 1985; 15/63, 24%; Niazy et al., 2005; 9/63, 14%), analysis of changes in the derivative's sign (13/63, 21%), and convolution with a template cardiac cycle (13/63, 21%). As shown in Fig. 4F, commonly used toolboxes include EEGLAB plug-ins HEPLAB (Perakakis, 2019; 11/60, 18%) and FMRIB (Niazy et al., 2005; 9/60, 15%), Peakfinder function (Yoder, 2026; 11/60, 18%), or standalone software, e.g., Kubios (Tarvainen et al., 2014). Notably, peak detection tools might automatically perform data processing steps (e.g., filtering) that are often not explicitly reported, potentially hampering reproducibility.

ECG acquisition parameters

A Number of ECG electrodes

n = 132 studies

B ECG lead

n = 132 studies

C Locations of ECG electrodes

n = 51 studies

Number of studies
— 5
— 10
— 15

ECG lead
— Lead I (n = 15)
— Lead II (n = 34)
— Lead III (n = 3)

ECG preprocessing parameters

D Filter cutoffs

n = 49 studies

Number of studies

0 10 20 30 40

E Algorithm for ECG peak detection

n = 63 studies

F Software for ECG peak detection

n = 60 studies

Figure 4. Various ECG leads, electrode locations, and preprocessing settings are used in the literature. Diamond symbols indicate choices driven by one research group (at least 50% of occurrences). (A) Most studies use two electrodes to record ECG. (B) While lead II is the most commonly used derivation, many studies also use electrode placements that do not correspond to any of the standard leads (N/C) or do not specify the lead used (N/M). None denotes studies which reconstructed ECG from EEG. (C) Locations of ECG electrodes that are placed to measure leads I, II, and III. (D) The filtering cutoffs for ECG largely mirror the

ones used for M/EEG during preprocessing. (E) Peak detection algorithms based on a

template cardiac cycle or ECG derivative are used most often. Category 'Other' includes

methods that appear only in 1 study. (F) Diverse toolboxes are used for ECG peak detection.

Category 'Other' includes toolboxes that appear only in 2 studies or less. Abbreviations: RC –

right clavicle, LC – left clavicle, RS – right shoulder, LCh – left side of the chest, LAb – lower

left part of the abdomen, RW – right wrist, LW – left wrist, LL – left leg/ankle, N/M – not

mentioned, N/C – not classified.

455 3.5. Dealing with cardiac artifacts

Since HERs are time-locked to the cardiac cycle, it is essential to consider the confounding

influence of cardio-vascular artifacts on HER effects. The two major factors that contribute to

these artifacts are the electric field generated by the heart (Dirlich et al., 1997) and pulse-

related movement of tissue or EEG electrodes (Cataldi et al., 2025; Kern et al., 2013). As

mentioned above, researchers predominantly employ one of three overarching strategies to

mitigate those effects: avoiding HER space-time points where the influence of artifacts is

potentially present, cleaning the data from artifacts, or including the ECG in the statistical

analyses. This pattern also held in the reviewed sample of studies, with 89% (117/132)

applying at least one of those strategies. However, the applied strategies and their effect on

HER might differ substantially.

In terms of artifact removal, ICA is used by 52% of all studies (69/132) to remove cardiac

components. This trend is corroborated by research that demonstrates the effectiveness of

ICA in attenuating the influence of cardiac artifacts on the measured HERs (Buot et al., 2021).

However, there is no standard way of performing ICA, and approaches vary across studies.

Most frequently, ICA is applied to continuous data (76%, 74/97 studies that applied ICA). In

contrast, around 24% of studies (23/97) choose to perform it after data epoching, likely to

specifically target artifacts occurring around the R-peak and T-peak (Dirlich et al., 1997).

While cardiac-related activity often displays relatively consistent spatial patterns across participants (Dirlich et al., 1997), ICA may not always accurately differentiate those from genuine brain signals. Variability of the ICA decomposition across participants can introduce additional variance in the data (Pontifex et al., 2017), potentially influencing HER findings. Given that 86% of the studies that remove cardiac ICs (59/69) also report the approach used for identifying these components, the field is moving towards transparent reporting, facilitating the reproducible removal of cardiac artifacts.

Among the reports, we find that manual identification of CFA components (37/69, 54%) is used more often than automatic (13/69, 19%) or semi-automatic (9/69, 13%) detection approaches (Fig. 5A). The CFA-related components are commonly identified based on the time course (32/69, 46%), topography (31/69, 45%), and power spectrum (13/69, 19%; Fig. 5B). The fully automatic approaches often rely on algorithms such as CORRMAP (Campos Viola et al., 2009), ICLabel (Pion-Tonachini et al., 2019), and SASICA (Chaumon et al., 2015). Semi-automatic cardiac artifact removal is often guided by measures such as phase consistency (7/69, 10%) or correlation (2/69, 2.9%) between ICs and ECG.

Only 29% of studies (20/69) that removed CFA-related ICs reported the number of removed components, which ranged between 1 and 3 with a median of 1.7 (Fig. 5C). Since many factors, such as the SNR or the M/EEG sensor configuration, influence the number of cardiac ICs obtained by the ICA, it could also be informative to report the combined explained variance of removed cardiac ICs. The explained variance might be a more generalizable measure across studies, allowing for better comparison of CFA removal efficiency. Thus, we suggest considering it for reporting in HER studies (Box 1). Note that differences in cardiac parameters and cleaning may induce spurious HER differences between groups or conditions; it is important to compare the cleaning (number of removed components, explained variance), the number of remaining epochs, and the results without the CFA removal.

Besides ICA, the reviewed studies employ several other techniques to suppress CFA (Fig. 5D). These include current source density analysis (CSD, 11/132, 8.3%), subtracting or regressing out the ECG artifact from the M/EEG data (9/132, 6.8%), principal component analysis (PCA, 2/132, 1.5%), and subtracting the average resting state HERs from task HERs (1/132, 0.8%). However, it remains an open question as to how these different methods compare with one another.

In addition, 21% of studies (28/132) aim to avoid the influence of cardiac artifacts by restricting the time range of interest used in their analyses. Some studies pursue this by excluding early time periods within the cardiac cycle (0-200 ms after the R-peak) from their analyses (Callara et al., 2023; Cambi et al., 2024; Engelen, Buot, et al., 2023; Giusti et al., 2024; Immanuel et al., 2014; Melo et al., 2022). Another common strategy is to choose a time window of approximately 455-595 ms post-R-peak (Gray et al., 2007; Lutz et al., 2019; Müller et al., 2015; Rapp et al., 2023; Schmitz et al., 2020, 2021; Schulz et al., 2013), following Dirlich et al. (1997), who demonstrated the minimal influence of the heart's electric field during this period. However, given the large variability across studies, there does not appear to be a consensus on a time range free from cardiac artifacts (Fig. 5F).

If HER trials are close to the surrounding heartbeats, it is reasonable to remove such trials to avoid contamination with surrounding HERs or heartbeat-related artifacts. Overall, 13% of studies (17/132) reported the removal of HER trials based on a minimal RR interval, the shortest time allowed between two consecutive R-peaks. We observe that the minimal allowed RR interval varies from 350 to 750 ms across these studies (Fig. 5E). Such large variability could be attributed to diverse participant populations and experimental designs, warranting study-specific definitions of the cutoff. Note that a high minimal RR interval threshold may impose unintended imbalance across the numbers of epochs among conditions or groups if there are underlying differences in heart rate. On top of that, such differences may signify distinct arousal states and distort the interpretation of HER findings. Therefore, it is advisable to report the number of rejected epochs per group and condition (Box 1).

Figure 5. Approaches to address cardiac field artifacts. (A) CFA-related independent components (ICs) are rejected manually, based on automatic approaches or a combination of those (semi-automatic). (B) Decision criteria that are used to define CFA-related ICs. The inset plot displays the number of studies using different algorithms for CFA IC removal. (C) The number of rejected CFA-related ICs per participant ranges from one to three across studies. (D) Additional approaches to address cardiac artifacts. Across the reviewed studies, overlap with artifacts is reduced by (E) excluding trials with short RR intervals, or (F) by using specific time periods of interest for analysis. Simulated ECG traces are shown in gray for reference in panels E and F. T-peak-based time windows (in orange) are shifted by 300 ms relative to the

536 R-peak for illustrative purposes. Abbreviations: ICA – independent component analysis, ICs –

537 independent components, CFA – cardiac field artifact, CSD – current source density, PCA –

538 principal component analysis, rsHER – resting state HER, N/M – not mentioned.

3.6. Epoching and baseline correction

540 Critical decisions in HER research include the choice of the cardiac reference event for

541 epoching (e.g. R-peak vs. T-peak) and the spatiotemporal window used for statistical analysis.

542 These choices can directly influence the detection of HER effects and may define susceptibility

543 to the influence of artifacts. In this review, we find that the vast majority of reviewed studies

544 use the R-peak as reference (123/132, 93.2%). In comparison, only 4.5% (6/132) used the T-

545 peak, and 2.3% (3/132) performed analyses using both peaks within a single study (Fig. 6A).

546 After epoching, baseline correction is typically performed to remove vertical amplitude offsets

547 and slow drifts by subtracting the average pre-stimulus amplitude from the whole epoch.

548 Although it is a standard procedure in conventional ERP analyses (Luck, 2014), it is

549 controversial in HER research, because the heartbeat has a short inter-beat interval (~1s),

550 potentially allowing activity from one cycle to affect the pre-stimulus time window of the next

551 cardiac cycle (Azzalini et al., 2019).

552 Overall, only 61% of studies (81/132) clearly state whether baseline correction was performed.

553 Out of these, 91% (74/81) include baseline correction and 8.6% (7/81) state that they do not

554 use it (Fig. 6B). The remaining 39% (51/132) do not mention it in their manuscript, leaving it

555 unclear if they performed baseline correction or just omitted details on this step. Of note, none

556 of the reviewed studies have directly compared baseline-corrected to uncorrected values.

557 Such a comparison could be of value to ensure the observed effects are not due to baseline

558 differences (Box 1). In particular, additional tests may be performed on the baseline window

559 itself or, when performing cluster-based permutation tests, the analysis should be run on the

560 whole epoch, including the baseline.

While 58% of the studies using the R-peak as a reference event (73/126) clearly state the use

of baseline correction, only two out of the nine studies using the T-peak (22%) did so. In both

studies, the baseline interval was set to -100 and 0 ms relative to the T-peak. A possible

explanation for the lower usage of baseline correction, here, could be that the time period

preceding the T-peak is close to the cardiac contraction, and hence more prone to

contamination by cardiac artifacts.

In R-peak locked analyses, baseline windows show considerable variability, ranging from -300

(min) to 50 (max) ms relative to the R-peak. Notably, 23% of these studies (17/73) define the

end of the baseline period at least 25 ms before the R-peak, potentially to avoid including

cardiac field artifacts (Coll et al., 2021). One study used a regression-based baseline

correction approach to avoid introducing baseline differences in the HER time window of

interest (Zaccaro et al., 2024). In future studies, it would be valuable to investigate the

effectiveness of baseline-correction methods and contrast them with the caveats associated

with not performing any baseline correction.

3.7. Predefined spatiotemporal HER regions and resulting findings

When it comes to quantifying the HER signal, amplitude is by far the most prevalent outcome

measure (128/132, 97% of studies). Alternative measures are rare, with only three studies

using HER latency (2.3%) and two studies examining global field power (1.5%). Note that

studies investigating heartbeat-locked neural activity using time-frequency analysis (e.g., Kim

& Jeong, 2019) were not considered in this review.

When it comes to choosing regions and time windows of interest, we observe that, overall,

analyses locked to T-peak cover a smaller time range (from -10 to 400 ms after T-peak). In

comparison, R-peak locked analyses start as early as -250 ms before the R-peak and range

until 1000 ms thereafter. Since studies often use averaging over a space-time window of

interest or cluster-based permutation tests to identify HER effects (Fig. 6C, studies using only

averaging: 47/115, 41%; only clustering: 30/115, 26%; both approaches: 38/115, 33%), we

analyzed the data separately for each of these two approaches.

The regions (analyzed for EEG only) and time windows of interest (pooled across EEG and

MEG) differ substantially between cluster- and averaging-based analysis approaches. As

expected, we see that all EEG channels are typically included in the cluster-based analysis

(Fig. 6F). However, averaging is performed more often only over frontocentral EEG channels

(Fig. 6E), often motivated by the results of previous studies (Billeci et al., 2021; Elkommos et

al., 2023; Schulz et al., 2020). Similarly, clustering approaches use wider time windows more

often than averaging approaches do. Nonetheless, there is a clear trend across both

categories, averaging as well as clustering, to limit analysis to a time range between 200-600

ms relative to the R-peak (Fig. 6E-F). In MEG studies, it is also common to restrict sensor-

space analysis to magnetometers only (Table S3).

Additionally, we also extracted information about channels and time points that show

statistically significant test results and pooled them across all reviewed studies. Overall,

significant HER effects were found between -10 and 1000 ms for R-peak locked analyses and

between 50 and 389 ms for T-peak locked analyses. Interestingly, there is a difference

between averaging- and cluster-based analysis in terms of where in time effects are most

often detected, as illustrated in the cumulative sum plots in Fig. 6E-F. Specifically for the R-

peak locked analyses, comparison of cumulative sum scores indicated that cluster-based

approaches identified the most common latencies significantly earlier than averaging (mode =

250 ms vs. 490 ms; permutation test on the time difference between peaks, p = 0.021; 1000

permutations). Although the reason for this discrepancy is unclear, one possibility is that

cluster-based permutation tests capture effects that are outside of narrow averaging windows

by scanning a wider time range. Another possibility is that shorter effects are diluted when

averaged across wider time ranges. In contrast, T-peak locked analyses showed no significant

difference between clustering (mode = 307 ms) and averaging (mode = 234 ms; permutation

test on time difference between peaks, p = 0.744; 1000 permutations), likely related to the

overall smaller time range of interest in T-peak locked analyses. However, it is important to

note that permutation clustering does not allow for any precise inference about the

spatiotemporal localization of effects (Sassenhagen & Draschkow, 2019), which means that

the comparison to averaging-based analysis should be interpreted with caution. Also note that

between-participant variability in R-T intervals may affect the presence or distribution of HER

effects in R- or T-peak locked analyses, so it may be worthwhile to compare such analyses.

However, few reviewed studies did so (2.3%, 3/132).

When performing analysis on high-dimensional data, one way to alleviate the multiple

comparisons problem is to define spatiotemporal regions of interest in advance (Luck, 2014).

Our review indicates that most studies follow this procedure and have predefined analysis

windows (88/132, 67%, referred to as primary in Fig. 6D). However, we also observe that in

33% of the studies (44/132), researchers perform a two-step procedure, where the selected

space-time window is based on the analyzed data within the same study (referred to as

secondary in Fig. 6D). While this approach can be useful to identify the electrode sites and

time windows where an effect appears in the studied population (Luck & Gaspelin, 2017), it is

not always trivial to prevent the analyses from becoming circular (Kriegeskorte et al., 2010;

Sassenhagen & Draschkow, 2019).

Figure 6. Predefined spatiotemporal regions and detected HER effects. Proportion of studies

using: (A) R- vs. T-peak as reference event, (B) baseline correction (applied vs. not applied,

unreported cases omitted), (C) averaging approaches vs. cluster-based permutation tests (or

both, if performed alongside each other), (D) only primary or both primary and secondary

analyses. For (E) averaging approaches or (F) cluster-based permutation tests, baseline

windows (top row), time ranges and regions of interest (second row), as well as spatiotemporal

distributions of significant effects (third row) are summarized. Panels E and F use cumulative

sum plots to visualize the different HER window types, pooled across EEG and MEG studies. Results are split by heartbeat reference R-peak (gray) or T-peak (orange) and aligned to the respective time range of the simulated ECG trace in the bottom row (gray / orange dashed line). T-peak based time windows are shifted by 300 ms relative to the R-peak for illustrative purposes. Topographies include only EEG channels, and colorbars below them indicate the number of studies. Diamond symbol indicates choices driven by one research group (at least 50% of occurrences).

3.8. Statistical analysis and power considerations

Neuroscientific studies are often underpowered, resulting in low research reliability and a potential waste of resources (Button et al., 2013). To assess power in the growing field of HER research, we estimate the minimal effect sizes that the reviewed pipelines can detect with a power of 80% at a 5% significance level, based on their sample size, number of groups and conditions, as well as the type of chosen statistical test.

Across studies, the number of included participants lies between 1 and 552, with a median of 32 (Fig. 7A), the number of studied groups lies between 1 and 6 (Fig. 7B), and the number of conditions lies between 1 and 20 (Fig. 7C). Many pipelines compare groups or conditions (301/410, 73%), or perform a correlational/regression analysis to study a linear association (109/410, 27%). As mentioned earlier, and where possible (Section 2.3), we approximate the number of heartbeat-related epochs used to obtain HERs for the many studies (73/132, 55%) that did not report this information. The estimated and reported numbers of heartbeat-locked epochs range from 22 to 5517 with a median of 300 averaged epochs per participant and condition (Fig. 7D). As shown in Fig. 7E, the t-test family (one-sample, two-sample) is the predominant choice in statistical analysis (74/222 pipelines, 33%), closely followed by variations of correlation (62/222, 28%) and ANOVA (52/222, 23%).

The resulting minimal detectable effect sizes range from 0.18 to 2.37, with a median of 0.86 Hedges' g. Fig. 7F shows their distribution across pipelines that used averaged HER values

in one- or two-sample t-tests, correlations, or ANOVAs (estimates including clustering and

regression pipelines, which require more assumptions, are shown in Fig. S1). In general, the

smaller the minimal detectable effect size, the broader the range of effect sizes that the

pipeline can robustly detect. To contextualize these minimal detectable effect sizes, we

compared them to meta-level effect sizes from a HER review by Coll et al. (2021). This review

investigated the frequently proposed role of HERs as markers of interoceptive functions

(Pollatos & Schandry, 2004). The meta-analyses were grouped into categories by topic and

assessed whether HER amplitudes are influenced by manipulations of attention (Attention

category) or arousal (Arousal category), whether HERs are related to measures of cardiac

interoception (Performance category), or whether they differ in clinical populations compared

to controls (Clinical category). When comparing the estimated detectable effect sizes to these

four meta-level effect sizes, we observe that only 0.7% (1/134) of pipelines had enough power

to detect Hedges' g of 0.35 (Performance meta-estimate), 1.5% (2/134) of pipelines could

reliably detect Hedges' g of 0.37 (Attention meta-estimate), 8.2% (11/134) of pipelines could

reliably detect Hedges' g of 0.49 (Clinical meta-estimate), and 33% (44/134) of pipelines could

reliably detect Hedges' g of 0.72 (Arousal meta-estimate). These observations suggest that

67% (90/134) to 99% (133/134) of pipelines may have yielded underpowered and potentially

unreliable findings. Such evidence draws attention to potential reproducibility and reliability

problems of current findings within the HER field. Studies with larger sample sizes could test

the robustness of previous findings, and future studies may consider adjusting their sampling

strategy to detect smaller effect sizes more reliably.

It is essential to note three key points. First, the review by Coll et al. (2021) focuses on a

subset of 45 HER studies that report interoception-related HER effects. We did not assess the

actual effect sizes among the studies we reviewed, as they varied in topics and experimental

designs. Second, the Hedges' g estimates presented here are based on approximations and

therefore should be interpreted with caution. Third, as mentioned previously, HERs are

susceptible to noise contamination due to their relatively low amplitudes of 0.5-2 μV (Pollatos

& Schandry, 2004; Schandry & Weitkunat, 1990). It is unclear how many trials, or heartbeat-

locked epochs, are required to obtain a stable evoked response per participant. Previous ERP

studies note that there is no straightforward answer, but it is clear that smaller effects (as

typically observed in HER amplitudes) generally require larger samples and more epochs to

be reliably detected (Boudewyn et al., 2018; Gibney et al., 2020). However, Hedges' g

estimates discussed in the previous paragraphs do not account for the levels of within-

participant noise across HER epochs, i.e., for trial averaging. To examine its effect on minimal

detectable effect sizes, we varied the hypothetical within- and between-participant variabilities

in HER values and used the numbers of averaged HER epochs from the reviewed studies to

obtain adjusted minimal detectable effect sizes (Supplementary Section B and Fig. S2). By

construction, they are increasing after adjusting for within-participant variations in HER across

epochs. Therefore, in Fig. 7F, we demonstrate optimistic estimates, unadjusted for possible

measurement noise. Future methodological studies should examine the levels of HER

variability across trials and participants and recommend recording settings. In practice, pilot

data can be used to assess HER variability and adjust the number of epochs (see

Supplementary Section C on some suggestions). At the very least, we recommend reporting

the number of averaged HER epochs per participant and condition (Box 1).

A lack of transparency related to sample size calculation is characteristic of evoked response

studies in neuroscience (Larson & Carbine, 2017). To confront this, we recommend

preregistering the planned HER analysis alongside power calculations (Box 1). Preregistration

is a declaration of planned analysis in a public registry, and it reduces the risk of bias (adjusting

methods after knowing outcomes), as well as increases transparency by allowing other

researchers to judge the reliability of outcomes (Hardwicke & Wagenmakers, 2023). Amongst

the reviewed studies, only 3.8% (5/132) mentioned having published a preregistration. Further,

there were no registered reports, advanced forms of preregistration in which the methods are

reviewed before data collection and a publication is guaranteed regardless of the outcome

(Chambers & Tzavella, 2022). It is possible that the high variability in methods (Sections 3.2-

3.6) and findings (Section 3.7) prevents HER researchers from preregistering their

hypotheses. This problem can be best remedied if, firstly, most research groups working in

the HER field preregister their studies by default, and secondly, more research is devoted to

the advancement of robust HER analysis methods.

Another potential benefit of preregistrations is overcoming publication bias, the tendency for

publications with significant findings to be published more frequently than those with null

findings (Thornton & Lee, 2000). There is evidence for publication bias in the HER field: we

observe that only 6.8% of studies (9/132) were published with no significant HER results. We

suggest that a combination of preregistrations with formal null testing could reduce publication

bias and make null findings more visible and reliable (Box 1). Such results would contribute to

the structured accumulation of knowledge in the field, as future studies could use this

information to refine their approach. Statistical assessment of "absent" effects is possible using

Bayesian statistics (Wagenmakers et al., 2018) and frequentist equivalence testing (Lakens

et al., 2018). Few studies in the reviewed sample have used Bayesian statistics to assert the

absence of effects, and we did not observe any use of equivalence testing or, importantly, any

definitions of the smallest effect sizes of interest. Determining the smallest effect size of

interest (Lakens et al., 2018) or the region of practical equivalence (Kruschke, 2021) is not

only useful for establishing the absence of effects, but also for making sure that a study does

not focus on significant effects too small to be reliably replicated (Box 1).

Figure 7. Statistical analysis and minimal detectable effect sizes. The settings used in experimental design and statistical analysis across studies are as follows: (A) sample size (x-axis is log-scaled); (B) number of groups; (C) number of conditions; and (D) estimated number of epochs averaged to obtain HERs per participant and condition (x-axis is log-scaled). (E) The statistical tests that were used in the reviewed studies. (F) Minimal detectable effect sizes with 80% power at 5% significance level, and four HER effect sizes from the meta-analysis by Coll et al. (2021), visualized by colored triangles. Estimated minimal effect sizes are based on the type of statistical test, sample size, number of participants, groups, and conditions, and are only shown for pipelines that use averaged HER values for t-tests, correlations, and ANOVAs. In Panels B and E, diamond symbols indicate choices driven by one research group (at least 50% of occurrences).

3.9. Control analysis

While small effect sizes could pose serious problems for the reproducibility of HER findings, even the presence of strongly significant, "large" effects does not guarantee that the observed HER effect reflects genuine heartbeat-evoked neural activity (Park & Blanke, 2019; Steinfath et al., 2025). Although many strategies exist to avoid or remove cardiac artifacts (discussed in Section 3.5), it is unclear how effective these are. To ensure that HER effects do not simply reflect cardiac artifacts, 65% of the studies (86/132) include ECG and other heartbeat-related controls in their statistical analysis (Fig. 8A). Specifically, it is common to repeat the analysis on ECG data alone or include it as a regressor of no interest in analyses (combined: 59/132, 45% of studies; Fig. 8B) to demonstrate that effects found at the level of M/EEG sensors do not only reflect cardiac field artifacts. Furthermore, many studies control for the length of the RR interval (45/132, 34%), for instance, by including it as a regressor of no interest in their analyses or by comparing the RR intervals of the studied groups. In addition, 26% of studies (34/132) control for heart rate variability, 6.1% (8/132) for respiration, and 3% (4/132) for blood pressure (Fig. 8A).

Apart from confounding factors that are directly related to cardiac activity, various heartbeat-independent factors can also affect HER. For instance, if HERs were recorded on-task, the concurrent activity evoked by external stimuli can overlap with HERs, leading to spurious results. Similarly, group-related differences in SNR, for example, due to an imbalanced number of trials, systematic differences in artifacts, comparisons between patients and controls, or differences across age groups, can lead to contaminated HER results. These problems can often be addressed by surrogate heartbeat control analysis or pseudotrial correction (Park et al., 2014; Steinfath et al., 2025). Pseudotrial correction works by subtracting heartbeat-independent confounding activity from the HERs. In contrast, surrogate heartbeat control analyses verify that observed effects are truly locked to the heartbeat by shuffling the onset timings of the cardiac markers within each condition, and repeating the same analyses multiple times (e.g., >100 iterations). The outcome statistic of the original

analysis is then compared to the distribution of statistics obtained from these permutations.

Hence, it can only be concluded that the original effect is truly related to the heartbeat if it

exceeds most of the permuted effects. In our review, we find that 19% of the studies (25/132)

perform surrogate heartbeat control analyses or other variations of data shuffling. Note that

both methods require a temporal jitter between task and heartbeat onsets. In addition, the way

that heartbeats are shuffled within an iteration affects the quality of control analysis. We

recommend staying as close to the actual data as possible: same number of epochs, similar

IBI and distance to task events, etc., and to clearly describe the motivation for these choices.

Currently, robust control methods for HER research are still being developed (most are listed

in the quality control checklist; Box 1). For a full overview of control variables that appeared in

the reviewed studies, see Fig. S4.

Figure 8. Variables and approaches commonly used for control analysis sorted by the

frequency of occurrence in descending order. (A) Control variables grouped by category. (B)

Detailed breakdown of ECG- and heartbeat-related control variables. Blue color indicates that

items shown in Panel B compose the category ECG and Heartbeat Related Controls from

Panel A.

Box 1. Quality-control checklist for HER analysis and reporting. We suggest key reporting

points and recommendations for HER studies based on the extracted information from the

reviewed studies and prior experience analyzing HERs.

Step 1 – Designing an experiment

A. <u>Preregistration</u>. Supports the reproducibility and reliability of findings, as well as the publication of null results. May incorporate several control analysis pipelines (e.g., with and without baseline correction).

B. <u>Relationship between heartbeats and experimental design</u>. HERs can overlap with task-related activity, and they might co-vary with differences in background activity across groups. Avoid overlaps, if possible, or address them in analysis (see Step 6 – Controls). Exact time locking between heartbeats and task triggers can be difficult to deal with, as correction methods require some jitter.

C. <u>ECG electrode leads, ground, and location</u>. Affect ECG-based control analyses. Report lead configuration used for analysis. Consider recording several leads (e.g. Lead I, II, III, and neck electrodes) to account for the three-dimensional field distribution.

D. <u>Recording of physiological activity.</u> Breathing, eye movements, cardiac and other parameters can confound HER measures.

Step 2 – Data preprocessing

A. <u>Filtering</u>. High- and low-pass filter settings can affect the HER shape. High-pass filtering could attenuate genuine HER activity. Hence, for slow HER components, consider a high-pass of 0.5 Hz, which can also diminish respiration-related artifacts.

B. Reference electrode. The choice of online and offline references can affect HER results and artifact removal. Common average reference is widely used and facilitates comparability across studies.

C. R-peak or T-peak. Or other points of reference. Can be compared to each other as they may result in different findings.

D. Minimal IBI. Prevents contamination by overlapping heartbeats. Longer thresholds may induce imbalance in the number of epochs if the heart rate is different between conditions or groups, potentially signifying differences in arousal. Report the number of removed epochs.

E. General cleaning. Report approaches used for noise removal. For ICA, include the description of the algorithm, types of identified components (e.g., ocular, muscle), and rejection criteria. Cleaning algorithms improve reproducibility, but require manual checks.

Step 3 – Cardiac Field Artifact Removal (CFA)

A. Rejection of cardiac ICs (if applicable). State whether ICA was applied to epoched or continuous data. ICA on epoched data might improve CFA detectability. Describe rejection criteria, number of removed cardiac-related components, and explained variance.

B. Other CFA removal methods. Detailed description of other approaches used to remove CFA. For example, regressing out ECG from M/EEG data. The HER signal is likely never artifact-free.

C. Limiting analysis time window (optional). Avoid time ranges that are prone to cardiac artifact contamination, e.g., limit analysis to time after T- and before P-wave.

Step 4 – HER computation

A. <u>ROI in time and space.</u> Describe start/end times & preselected sensors as well as the motivation for the selection, also for cluster-based permutation tests.

B. <u>Baseline correction</u>. Clearly state if omitted or performed. We recommend avoiding baseline-correction, and testing if the baseline window shows differences between groups or conditions of interest. If performed, include information on the method and time window.

C. <u>Number of epochs</u>. Describe how many epochs were averaged or otherwise analyzed to obtain HER per participant, group, and condition. The number of epochs affects the signal-to-noise ratio, and imbalance may result in spurious findings.

D. <u>Visualization</u>. Show figures with HER results as evoked responses and as topographies.

Step 5 – Statistical analysis

A. <u>Statistical power.</u> Ensure that the analysis has sufficient statistical power to detect expected results reliably. E.g., simulations that can be combined with resampling pilot data.

B. <u>Regions of practical equivalence.</u> For both Bayesian and frequentist statistics, define the smallest effect of interest to test if observed effects are negligibly small (i.e., "absent") and to avoid highlighting practically irreproducible (too small) effects.

Step 6 – Controls

A. Surrogate heartbeat analysis. Confirms that observed effects are genuinely locked to the heartbeat but has to be carefully designed. To be valid, they should match real heartbeat properties (e.g. IBI, time to stimuli, number of trials, etc.). Describe how it was performed and how exactly it addressed potential confounding effects.

B. Task overlap controls. For example, pseudotrial correction can help isolate heartbeat-driven effects from task-related activity. Describe how it was performed and how exactly it addressed potential confounding effects.

C. ECG control analysis. Assesses whether differences in ECG exist that might confound the HERs via cardiac artifacts, e.g. by repeating statistical analysis on the ECG channel or regressing out ECG amplitudes. Use multiple leads if possible.

D. Effect of baseline. It is advisable to show the HER results without baseline correction and test if the baseline itself shows significant effects. When performing cluster-based testing, include the baseline time window into analysis.

E. Effect of CFA removal approach. It is advisable to report the HER results without CFA removal to show that they are not explained by different CFA removal efficacy across groups or conditions.

F. Other cardiac parameters. Should be included in statistical analysis. Examples: heart rate, heart rate variability, blood pressure.

Step 7 – After publication

A. Data sharing. Publicly shared data enable other researchers to reproduce current findings, develop new research questions, and, importantly, further HER methods. This facilitates more reliable, faster, cost-effective research.

B. <u>Code sharing</u>. Allows for the highest level of transparency in understanding the analysis steps.

Abbreviations: HERs – heartbeat-evoked responses, ECG – electrocardiography, IBI – inter-beat interval, M/EEG – magneto- or electroencephalography, ICA – independent component analysis, ICs – independent components, ROI – region of interest

4. Discussion

This systematic review aimed to comprehensively characterize the diverse set of methods

used in non-invasive heartbeat-evoked responses research using M/EEG. Our analysis of 132

publications revealed substantial heterogeneity across many processing steps, including data

acquisition, data preprocessing, and statistical analyses. While some predominant

methodological patterns emerge, such as a higher percentage of EEG compared to MEG

studies and the frequent use of the R-peak as a cardiac reference marker, most other steps

show considerable variability, as indicated by large entropy scores. However, for the majority

of preprocessing steps, studies typically use only one of the many possible options.

Apart from documenting large variability in methodological choices, this review also reveals a

large proportion (40-80%) of unreported decisions for processing steps that are critical for

HER analysis such as the number of rejected CFA-related independent components (ICs), the

number of epochs averaged to obtain HERs per participant and condition, or the baseline

window approach used. Taken together with a worrying estimation that 67-99% of studies

might be underpowered for detecting meta-level HER effects, it is critical that the community

takes more decisive steps towards transparency and reliability of findings.

To alleviate reporting ambiguity and improve the robustness of experimental designs and

analyses, we compiled a checklist containing key points that should be considered when

conducting and reporting HER studies (Box 1). Since HERs are a subclass of evoked

responses, many potential problems are general to the field of M/EEG research and are
already summarized elsewhere (e.g. Keil et al., 2014). Furthermore, potential problems
relating to the definitions of cardiac phases, a closely related heart-brain topic, are also
summarized in a recent review (Caparco et al., 2025). However, some important
considerations are unique to HER analysis and require special attention to ensure the quality
and reproducibility of results. While previous reviews already highlight some HER-specific best
reporting practices (Azzalini et al., 2019; Park & Blanke, 2019), a comprehensive list of
reporting guidelines could be useful for experienced and new HER researchers, as well as
reviewers of HER studies. Our checklist provides a broad overview of existing approaches for
tackling artifacts (mainly, the CFA) and other confounders (e.g., overlap with task-evoked
activity).

Despite the overall lack of consensus in methodological approaches, our results highlight the
most common choices for each processing step. However, it is important to note that the
correctness and suitability of each choice cannot be judged solely based on its occurrence
frequency, especially when most studies only use one choice for each processing step. To
better understand the impact of the various methodological choices on the observed effects,
it is crucial to either perform simulation-based studies with known ground truth derived from
single cell recordings (De Falco et al., 2024), intracranial measurements (Kern et al., 2013),
or to perform multiverse analyses on real data (Steegen et al., 2016).

Thus, we provide an overview of the currently used methods that could serve as a starting
point for novel experimental designs or systematic method comparisons. Based on this
overview and previous literature, we have also compiled a tentative HER pipeline intended to
provide initial guidance and some unification for HER analysis, rather than to describe strict
rules (Figure 9). In addition, we highlight several processing steps which are likely to affect
HERs, but their influence has not been systematically studied yet (Box 2). Below, we reiterate
the motivation for investigating these steps in more detail.

Box 2. Outstanding methodological questions in HER research.

> To further HER research, the development of methodology is required. Effects of the following processing steps, as well as their interactions, on HER results require further systematic evaluation:
>
> 1. High- and low-pass filter parameters
>
> 2. ECG recording and subsequent ECG-based CFA controls
>
> 3. CFA removal with ICA or other spatial filtering techniques with varying settings
>
> 4. Influence of individual differences in cardiac parameters on HER
>
> 5. Ocular, muscular, and respiratory artifact influence and cleaning
>
> 6. Baseline correction
>
> 7. Variability of HER epochs and required number of trials

The high likelihood of being confounded by cardiac artifacts is a major factor that distinguishes
HERs from other ERPs. Importantly, the complex three-dimensional nature of cardiac electric
fields, with dynamic changes in scalp potential patterns across the heart cycle, poses a critical
challenge (Dirlich et al., 1997; Pérez et al., 2005). Decomposition methods, such as ICA, which
aim to represent the data in lower-dimensional components, often struggle with time-
dependent artifacts. Although the majority of studies use ICA to remove artifacts from the data,
there is no consensus on the best approach for identifying CFA-related ICs. While different
approaches lead to varying results in artifact suppression, no direct comparison between
methods has been conducted to date. It is unlikely, however, that any of the current artifact
removal methods result in the perfect suppression of CFA in M/EEG data (Park & Blanke,
2019). Therefore, the HER time courses and topographies observed in the literature represent
a complex mixture of residual CFA and neural activity.

A prominent example of this mixture is the deflection in HER epochs reminiscent of the R-
peak shape in ECG. It is often observed around 0 ms when the epochs are time-locked to the

R-peak. This deflection likely reflects residual CFA in the EEG because it occurs too early to

reflect a genuine neural reaction to the R-peak. Interestingly, in some studies, it does not occur

exactly at 0 ms. This puzzling observation remains to be resolved in the future, but there may

be several hypothetical explanations. Firstly, systematic errors in the identification of R-peaks

may result in this shift. It is the responsibility of researchers to ensure that R-peaks are

identified precisely. This can be checked by epoching the ECG signal to detected R-peaks, to

ensure that the R-peak aligns with 0 ms. Secondly, EEG preprocessing (e.g., downsampling

and filtering, CFA cleaning) may lead to these shifts. This can be checked by epoching the

raw EEG to detected R-peaks. Thirdly, the shifts may stem from CFA spread and EEG

electrode orientation characteristics (Myllylä et al., 2013). Modeling and recording ECG closer

to the brain (e.g., neck) might inform this hypothesis.

In addition, groups with differences in cardiac function (e.g., patients vs. controls) might show

different contributions of CFA to the M/EEG signal. While this makes artifact removal

especially important, it can also complicate the CFA detection and removal. Importantly,

cardiac artifact removal itself can induce condition differences. For example, if patients and

controls are compared, the ground-truth level of cardiac artifacts might be the same, but due

to differences in overall SNR between the groups, CFA removal is more efficient in controls

than in patients. Hence, more residual CFA remains in patients, potentially leading to spurious

HER differences. One useful control is to verify that results remain overall the same without

cardiac artifact correction. However, if results disappear without correction, this does not

necessarily indicate that the original findings were driven by inadequate artifact removal, it

may also reflect that CFA correction was necessary to reveal the underlying neural signal. A

potential proxy for assessing CFA influence on the data is the magnitude of the residual R-

peak artifact in the M/EEG recording. Since this artifact results from volume conduction, it

likely contains no contribution from neuronal activity. A significant difference in R-peak artifact

amplitude between groups or conditions would therefore be cause for concern, as it indicates

differential CFA contributions that could confound the HER.

Thus, CFA identification and removal pose a significant challenge. On the one hand,

simulations are needed to evaluate the influence and removal efficacy of cardiac artifacts on

the measured signal. On the other hand, high-density ECG recordings with sophisticated

forward models offer a promising tool for a better separation of CFA and brain activity in real

data. The complex nature of the CFA is also affecting ECG-based control analyses, which aim

to confirm that HER effects reflect differences in brain activity, rather than volume-conducted

CFA. Although low-density ECG derivations are generally sufficient for identifying R- or T-

peaks as reference events, they only capture a portion of the complex cardiac electrical field.

Since experimental conditions can induce changes in ECG that might be detectable at some

recording sites but not at others (Gray et al. 2007), several ECG electrodes covering all three

dimensions of the ECG are necessary to show that HER effects are not volume-conducted

CFA.

Even though the removal of cardiac artifacts plays an important role in HER research, artifact

correction also carries the risk of removing genuine parts from the HER. Hence, many

researchers analyze predefined time ranges for their analysis, which show low influence of

the CFA (Dirlich et al., 1997). While early HER studies aimed to identify spatiotemporal points

where the influence of cardiac-related artifacts is minimal (Dirlich et al., 1997), our review finds

little progress in recent years toward identifying and validating these optimal time ranges.

Therefore, it remains essential to ensure that changes in HER are not accompanied by

changes in cardiac parameters.

Our review supports the idea of considerable temporal variability with predominantly

frontocentral HER effects across M/EEG studies. We observe a wide range of chosen time

windows, with a preference for limiting analyses to a time range of approximately 200 to 600

ms after the R-peak. This time range is in agreement with intracranial recordings, revealing

heartbeat-related brain activity to emerge primarily between 200 and 400 ms after the R-peak

(De Falco et al., 2024; Kern et al., 2013). However, even in these invasive studies, there is

considerable variability regarding the onset, duration, and anatomical location of cardiac-

related neural responses. This spatiotemporal variability in observed effects suggests that there is no unique empirically observed manifestation of HER (Gautier et al., 2025), but that it reflects a mixture of responses to different parts of the cardiac cycle, or even a sustained, ongoing tracking signal related to blood pressure and heart rate variations. Overall, those pieces of evidence point to a wide range of cardiac-locked modulations of neuronal activity depending on the specific task, cognitive state, experimental condition, or pathology.

In line with this, another potential source of variability in observed locations of HER effects that obscures comparability is the time-locking reference for HER epochs. While most studies choose R-peak as a reference (123/132, 93.2%), some use T-peak (6/132, 4.5%), and only a few compare both within a single study (3/132, 2.3%). Due to interindividual variability in R-T intervals, the two approaches may result in different spatiotemporal locations of effects and are not directly comparable. Because only a few studies choose both R- and T-peak locking, it is difficult to judge how often the results would be similar or different and why. Ideally, more studies investigate this, and in the future, more mechanistic explanations are proposed. It would also be informative to see larger HER epochs in more studies: how the effects look when we consider a period spanning several heartbeats, e.g., from -2 to 2 seconds around the R-peak, and how the results depend on the point of reference within the cardiac cycle.

The spatiotemporal variability of the HER affects the suitable analysis options. Due to the inconsistency of results across studies, there is currently no clear spatiotemporal region that shows robust HER effects. Hence, many researchers use hypothesis-free cluster-based permutation testing. While this approach is convenient to circumvent the variable nature of the HER, it is important to keep in mind that cluster-based permutation tests do not allow for exact spatial or temporal interpretation of effects (Maris & Oostenveld, 2007; Sassenhagen & Draschkow, 2019). In addition, we observe that many studies use a two-step procedure, where the region of interest and time range are first defined based on primary cluster-based permutation analysis. The resulting cluster is then used in further secondary analyses. However, if the analysis used for data selection is not independent of the secondary analysis,

circularity may become a concern (Kriegeskorte et al., 2009). While this does not mean that

the results are incorrect, it underscores the need for independent replication and careful

evaluation of the statistical validity of such two-step analyses. Ideally, cluster-based

permutation analyses with a more exploratory nature are followed by clear confirmatory

studies with predefined regions of interest on separate datasets, allowing for more

straightforward power estimation and tests for the absence of effects as well as facilitating

replicability. However, in studies that aim to establish the presence of HER differences, e.g.,

between conditions or groups of patients, without the focus on exact spatiotemporal locations,

conventional cluster-based analysis remains a viable option (Rousselet, 2025). Moreover, the

cluster-based tests are not the only way to correct for multiple comparisons when performing

mass univariate tests on spatiotemporal points in neurophysiological data: other ways include

traditional conservative approaches such as Bonferroni (Armstrong, 2014) or Benjamini-

Hochberg (Benjamini & Hochberg, 1995) corrections, as well as novel applications of Bayesian

generalized additive multilevel models (Nalborczyk & Bürkner, 2025) or cluster-depth tests

(Frossard & Renaud, 2022).

Careful experimental design and control analyses are essential to isolate genuine HER activity

from other sources of variability. When HERs are studied in task settings, as is the case in

most of the reviewed studies, task-related evoked responses may overlap with internal

responses like the HER. This can become problematic if, for example, experimental conditions

systematically differ in the temporal alignment between heartbeats and task triggers (e.g.,

presented stimuli), as the resulting HERs might capture different parts of the stimulus-evoked

responses, leading to spurious HER differences. This opens additional challenges. On the one

hand, researchers have to make sure that potential HER effects are not a result of temporally

overlapping task-related differences in neural activity. Currently, we find that there are two

approaches used in the literature to deal with this situation: surrogate heartbeat analyses (Park

et al., 2014) and pseudotrial correction (Steinfath et al., 2025). On the other hand, if effects in

task-evoked activity are observed, researchers have to make sure that these are not a result

of overlapping cardiac-related activity, given that heartbeats may align to task triggers (e.g.,

Kunzendorf et al., 2019). Overall, disentangling overlapping effects might also be approached

by regression-based ERP modelling (Ehinger & Dimigen, 2019). However, the studies

included in our review indicate that, so far, these methods have not gained wider use in the

HER research community. Further work is therefore required to show in which scenarios the

different methods work best and where the limitations are. Such limitations include tasks

where a tight time-locking between HERs and external stimuli is present (e.g., tones presented

synchronously to the heartbeat), because current correction methods depend on jitter to

delineate HER activity from task-evoked responses.

Furthermore, the mere presence of surrogate analysis or pseudotrial correction does not

guarantee the proper control for potential confounders. Depending on the experimental

paradigm and the confounder in question, specific time windows or numbers of simulated

epochs need to be carefully chosen. If a group difference in HERs arises from unequal SNR

due to imbalanced numbers of trials, it is important that the number of surrogate heartbeats is

preserved in the control analysis. Generating surrogate heartbeats with an arbitrary number

of epochs would alter the variance and hence the statistical power of the surrogate test,

undermining its validity. Likewise, the temporal relationship between HER and confounding

factors has to be preserved. For example, if heartbeats occur at different delays relative to

task stimuli across conditions, the resulting HER epochs capture different portions of the task-

evoked ERP (Steinfath et al., 2025). For the surrogate analysis to properly control for this, the

surrogate trials must maintain the temporal distribution of heartbeats relative to the task events

within each condition. When the heartbeats are shuffled randomly, without respecting the

original timings, the overlap with the task-evoked ERP would be disrupted, thus invalidating

the generated null distribution. As the guidelines on the robust design of surrogate control

analysis depend on the individual experimental design, we advise that the authors clearly state

how exactly they performed it, what problem they were trying to address, and how the

surrogate method solved it. This level of transparency enables the readers to make judgments

about the quality of surrogate analysis, beyond the fact that it was performed.

Adding to these challenges, the heart does not operate in isolation and is coupled to several

other physiological systems (Hall & Hall, 2021). This coupling can complicate the separation

of genuine HER effects from other correlated sources of variability, such as eye movements

(Ohl et al., 2016) or respiration (Yasuma & Hayano, 2004; Zaccaro et al., 2024). Consequently,

careful experimental design and control analyses are required to delineate heartbeat-related

effects from other influences. Importantly, our review revealed that, across HER studies, a

broad range of different control variables has been included in the analyses. While eye

movements or respiration are not among the most frequent choices, some experimental

designs may be more prone to these specific confounds, which warrant explicit care. As we

did not assess the adequacy of control analyses across the reviewed studies, some results,

especially spatiotemporal locations of detected HER effects, are likely to be affected by traces

of various artifacts and other sources of activity (task, physiology) and should be interpreted

with caution. We encourage the readers to apply stricter exclusion criteria (e.g., presence of

ECG or respiration controls and surrogate analysis) in the web application we provide

(https://paulsteinfath.shinyapps.io/her-systematic-review/), which offers updated, dynamically

generated　　versions　　of　　the　　figures　　presented　　in　　this　　review.

Pipeline Step	Choice
Acquisition	● Brain data (EEG/MEG...) ▲ Respiration (Park et al., 2014) ▲ (Eye)-Movements (Ohl et al., 2016) ▲ Multiple-lead ECG (Babo-Rebelo et al., 2016)
Bad Channels & Epochs	▲ (semi)automatic pipeline with visual inspection (Luck, 2014)
Referencing	● Linked mastoids or Cz during recording ●▲ Average reference during analysis (Candia-Rivera et al., 2021)
Filtering	●▲ High-pass: 0.1–0.5 Hz (Tanner et al., 2015) ●▲ Low-pass: 30–40 Hz (Luck, 2014)
ICA	▲ (semi)automatic component rejection with visual inspection (Campos et al., 2009; Pion-Tonachini et al., 2019) ●▲ Remove ocular components (Engelen et al., 2023)
CFA Removal	●▲ ICA-based cardiac artifact removal (Buot et al., 2021; Cataldi et al., 2025) ●▲ Minimal IBI threshold (Engelen et al., 2023)
Epoching	● R- or T-peak alignment ● Epoch window: -200–800 ms ▲ Baseline correction is not advised (Azzalini et al., 2019)
Statistical Test	● Cluster-based ● Average over pre-defined ROI (electrodes & time points) ▲ Test for the absence of effects (Bayesian/equivalence testing) (Lakens et al., 2018; Wagenmakers et al., 2018)
Control Analyses	▲ Control for ECG artifact across several ECG leads (Babo-Rebelo et al., 2016) ●▲ Surrogate heartbeat control (Steinfath et al., 2025) ●▲ Physiological covariates (HR, HRV, breathing, eye movements) (Park et al., 2014) ▲ Compare number of epochs between groups or conditions (Luck, 2014) ▲ Repeat analysis without CFA removal to check for cleaning biases (Al et al., 2020)

●Common choice based on Review　▲Based on Literature

1010

Figure 9. Tentative HER pipeline. Future methodological work is required to determine the

best combinations of preprocessing steps and methods for HER research, and the quality of

processing choices cannot be judged solely by the frequency of their occurrence in our review.

However, to promote comparability and provide practical guidance, we present a tentative

pipeline based on suggestions in the literature and common choices among the reviewed

studies. Based on the current knowledge, this tentative pipeline can serve as a solid starting

point for HER analysis rather than a decisive guideline. See Box 1 for detailed considerations

regarding each processing step.

1019

The current review has several limitations. Since only information about methods was extracted, we cannot compare and recommend any methods solely based on how often they are used. Furthermore, methods used by particular HER research laboratories might be overrepresented in the overall results. In fact, we observed that 20% of studies in the reviewed sample (27/132) have the same three last authors (max. 12 studies per last author). However, we added a 'dominance indicator' represented by a diamond to all histograms to foster transparency. Most of the processing choices we reviewed do not represent research practices of any particular group of authors, except for ECG recording and preprocessing. Other determinants, such as the study sample characteristics (patients or controls, age groups) and research topics, may contribute to the reported variability in methods and findings, but an in-depth analysis of these factors is beyond the scope of this review. However, our interactive resource contains the relevant information for further inquiries. Another limitation is that we did not extract the effect sizes per reviewed study or relate them to the employed methods due to the substantial heterogeneity in research questions. Therefore, an objective comparison of methodological choices for HER analysis on the same dataset is a crucial next step for the advancement of HER methodology. Additionally, we did not extensively extract all non-significant control analyses, which might also bias the representativeness of some of the results. It is critical to remember that any review is subject to publication bias when it comes to the included studies (Thornton & Lee, 2000). Overall, we find that a stunning 93% of reviewed studies reported statistically significant findings. Given that we also observe that a large proportion of studies might be underpowered, the small number of studies with non-significant findings is another evidence of a possible publication bias (Button et al., 2013).

The problem of publication bias, however, can be remedied in the future of the HER field by pursuing more transparent practices such as preregistrations and registered reports, and reporting the significance of null findings through Bayesian approaches (Wagenmakers et al., 2018) or frequentist equivalence testing (Lakens et al., 2018). We suggest that, just as for other research domains, preregistrations and especially registered reports can increase the

quality of HER research and are a key solution to many of the issues outlined in this review

(Chambers & Tzavella, 2022; Hardwicke & Wagenmakers, 2023; Soderberg et al., 2021). In

such a growing field, it might be difficult to navigate the multiplicity of methodological choices

across the many processing and analysis steps, and prepare a robust research plan in

advance. This is further complicated by variability in reported HER findings. Therefore, based

on this review, literature, and our current knowledge, we propose a tentative HER pipeline,

which, along with a broader quality-control checklist (Box 1), could serve as a solid starting

point for future investigations (Figure 9). Characterizing the temporal, spatial, and spectral

components of HER across contexts is the key direction of future research in the field, but this,

first and foremost, warrants the advancement of understanding how to analyze HERs in the

most robust way and heavily relies on transparent and reliable findings. This review and future,

much-needed, methodology-oriented HER studies will shed light on these decisions and help

researchers in preparing their HER studies. Ultimately, accumulation of robust evidence

through reliable research with transparent methodology is the way to understanding what HER

is, and thereby, how the brain and body interact.

Our review provides a set of reasonable candidate methods to consider for the preprocessing

and analysis of HERs, and the extracted information is published with the article. To make the

results more accessible, we also provide an easy-to-use interactive interface for inspecting

the dataset. We hope that this review will serve as a foundation for improving the

methodological consistency across HER studies.

Statement on gender composition within HER research and citation diversity

Recent evidence suggests that publications with female names at prominent positions in the

author list are cited less often in the field of neuroscience than should be expected (Dworkin

et al., 2020; Hefter et al., 2025). In light of recent discussions by journals and publishers (e.g.,

"Citation Diversity Statement", 2023; Fulvio et al., 2021; Sweet, 2021), we analyzed gender

approximations across the HER literature included in this review to guide the benchmarking

of future citation practices.

As done by Dworkin et al. (2020), gender estimates (F = female; M = male) were extracted for

the names of the first and last authors across all references (n = 132) through an automated

service (https://gender-api.com) and the cleanBib toolbox (D. Zhou et al., 2022). In case of

low assignment probability (<.80), publicly available pronouns were manually searched for.

Following the exclusion of those for which this failed (n = 2), the author pairs were classified

as 18% (n = 24) FF (first author/last author), 10% (n = 13) MF, 31% (n = 40) FM, and 41% (n

= 53) MM. There were no publications by singular authors. Four publications categorized as

FM were authored by our lab.

A median split by publication year showed that gender balance improved over time (since

October 2020 [Δ before]: 23% FF [+9.2%], 14% MF [+7.7%], 28% FM [-6.2%], 35% MM [-

11%]). The observed pattern resembles the gender distribution (18% FF, 14% MF, 29% FM,

39% MM) that resulted from large-scale analyses by Hefter et al. (2025). For more detailed

analysis see Section E of the Supplementary Material.

To note, the applied procedure falls short of some aspects: 1) shared first and/or last

authorship were not considered, 2) gender estimates and openly accessible pronouns may

not have been, in full, reflective of the respective individuals' gender identities, and 3) gender

was taken as dichotomous variable only, and thus, the procedure does not allow for monitoring

gender representation in its full variety. Lastly, some clustering of HER publications (using a

threshold of ≥5) was observed, with 27 studies (20%) last authored by the same three names

(2 F, 1 M), and 15 studies (11%) first authored by another set of three names (1 F, 2 M).

Lastly, we examined our own citation behavior in the main text (n = 111). After excluding four

references with undetermined author gender, the remaining were classified as 15% (n = 16)

FF, 12% (n = 13) MF, 23% (n = 25) FM, and 50% (n = 53) MM. These proportions include nine

single-authored studies (1 FF, 8 MM) and five from our lab (2 MM; 3 FM, with two amongst the reviewed studies). In total, 35 of the 132 studies included in the review (26.5%) were also cited in the main text.

Studies included in the review

(Al et al., 2020, 2021, 2023; Alkhachroum et al., 2024; Azzalini et al., 2021; Babo-Rebelo et al., 2019; Babo-Rebelo, Richter, et al., 2016; Babo-Rebelo, Wolpert, et al., 2016; Banellis & Cruse, 2020; Baranauskas et al., 2017; Billeci et al., 2021; Birba et al., 2022; Bogdány et al., 2022; Buot et al., 2021; Callara et al., 2023; Cambi et al., 2024; Candia-Rivera, Catrambone, et al., 2021; Canales-Johnson et al., 2015; Candia-Rivera, Annen, et al., 2021; Candia-Rivera et al., 2023; Candia-Rivera & Machado, 2023a, 2023b; Couto et al., 2014, 2015; de la Fuente et al., 2019; Di Bernardi Luft & Bhattacharya, 2015; Dirlich et al., 1997, 1998; Elkommos et al., 2023; Engelen, Buot, et al., 2023; Fittipaldi et al., 2020; Flasbeck et al., 2020, 2021, 2024; Fouragnan et al., 2024; Fukushima et al., 2011; Gao et al., 2023; García-Cordero et al., 2017, 2016; Gentsch et al., 2019; Giusti et al., 2024; Gray et al., 2007; Hermann et al., 2024; Hodossy et al., 2021; Immanuel et al., 2014; Ito et al., 2019; Judah et al., 2018; Kamp et al., 2021, 2023; Katkin et al., 1991; Kato et al., 2020; Khoshnoud et al., 2022, 2024; Kim et al., 2019; Koreki, 2024; Kritzman et al., 2022; Kumral et al., 2022; Lechinger et al., 2015; Legaz et al., 2022; Leopold & Schandry, 2001; Liu et al., 2022, 2023; Liuzzi et al., 2024; Lutz et al., 2019; MacKinnon et al., 2013; Mai et al., 2018; Maister et al., 2017; Majeed et al., 2022; Marshall et al., 2017, 2018, 2019, 2020, 2022; McCraty et al., 2004; Melo et al., 2022; Migeot et al., 2023; Montoya et al., 1993; Müller et al., 2015; Ono et al., 2024; Pang et al., 2019; Park et al., 2014, 2016; Pelentritou et al., 2024; Pérez et al., 2005; Perogamvros et al., 2019; Petzschner et al., 2019; Pollatos et al., 2005, 2016; Pollatos & Schandry, 2004; Poppa et al., 2022; Rapp et al., 2023; Ren et al., 2022; Richter et al., 2021; Richter & Ibáñez, 2021; Salamone et al., 2018, 2020, 2021; Schandry et al., 1986; Schandry & Montoya, 1996; Schandry & Weitkunat, 1990; Schmitz et al., 2020, 2021; Schulz et al., 2013, 2018, 2020;

Schulz, Ferreira de Sá, et al., 2015; Schulz, Köster, et al., 2015; Sel et al., 2016; Shao et al., 2011; Simor et al., 2021; Solcà et al., 2020; Stoupi et al., 2024; Tanaka et al., 2023; Terhaar et al., 2012; Todd et al., 2021; Ventura-Bort & Weymar, 2024; Verdonk et al., 2021, 2024; Villena-González et al., 2017, 2023; Wei et al., 2016; Weijs et al., 2023; Yoris et al., 2017, 2018, 2024; Yuan et al., 2007; Zaccaro et al., 2022, 2024; Zhang et al., 2023; H. Zhou et al., 2022; M. Zhou et al., 2024; Zwienenberg et al., 2023)

CRediT

P.S.; M.A.; N.K.: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Data Curation, Writing – original draft, Writing – review & editing, Visualization, Project administration. T. L.: Data Curation, Writing – review & editing. L.E.: Gender Citation Diversity Statement, Writing – review & editing. V.N.; A.V.: Writing – review & editing, Supervision, Funding acquisition.

Data and code availability statement

The full data table containing all relevant information extracted from the literature that is used in this systematic review can be accessed at https://osf.io/znrbm. In addition, we provide an interactive web application that allows browsing the dataset: https://paulsteinfath.shinyapps.io/her-systematic-review/. All analysis scripts are available at https://github.com/PaulSteinfath/systematic-hep-review. Citation diversity code along with gender approximation datasets can be accessed at https://github.com/liobaenk/citation-diversity-tracking.

Acknowledgement

This project was supported by the Max Planck Society. T.L. received funding from the Ministry of Health and Social Security of the Government of Luxembourg. We thank the reviewers for

1147 their valuable comments and the authors of reviewed studies for moving forward the HER

1148 field.

Declaration of generative AI in scientific writing

1150 During the preparation of this work the authors used OpenAI chatGPT (GPT-4o), Anthropic

1151 Claude 4, and Grammarly in order to proofread the text, and assist during analysis, code

1152 generation and documentation. After using these tools, the authors reviewed and edited the

1153 content as needed and take full responsibility for the content of the publication.

Declaration of competing interest

1155 The authors declare no conflicts of interest.

References

Al, E., Iliopoulos, F., Forschack, N., Nierhaus, T., Grund, M., Motyka, P., Gaebler, M.,

Nikulin, V. V., & Villringer, A. (2020). Heart-brain interactions shape

somatosensory perception and evoked potentials. *Proceedings of the National*

Academy of Sciences of the United States of America, *117*(19), 10575–10584.

https://doi.org/10.1073/pnas.1915629117

Al, E., Iliopoulos, F., Nikulin, V. V., & Villringer, A. (2021). Heartbeat and somatosensory

perception. *NeuroImage*, *238*(February), 118247.

https://doi.org/10.1016/j.neuroimage.2021.118247

Al, E., Stephani, T., Engelhardt, M., Haegens, S., Villringer, A., & Nikulin, V. V. (2023).

Cardiac activity impacts cortical motor excitability. *PLOS Biology*, *21*(11),

e3002393. https://doi.org/10.1371/journal.pbio.3002393

Alkhachroum, A., Fló, E., Manolovitz, B., Cohan, H., Shammassian, B., Bass, D., Aklepi, G.,

Monexe, E., Ghamasaee, P., Sobczak, E., Samano, D., Saavedra, A. B.,

Massad, N., Kottapally, M., Merenda, A., Cordeiro, J. G., Jagid, J., Kanner, A.

M., Rundek, T., … Sitt, J. D. (2024). Resting-State EEG Signature of Early

Consciousness Recovery in Comatose Patients with Traumatic Brain Injury.

Neurocritical Care, *41*(3), 855–865. https://doi.org/10.1007/s12028-024-02005-2

Armstrong, R. A. (2014). When to use the Bonferroni correction. *Ophthalmic and*

Physiological Optics, *34*(5), 502–508. https://doi.org/10.1111/opo.12131

Azzalini, D., Buot, A., Palminteri, S., & Tallon-Baudry, C. (2021). Responses to heartbeats in

ventromedial prefrontal cortex contribute to subjective preference-based

decisions. *Journal of Neuroscience*, *41*(23), 5102–5114.

https://doi.org/10.1523/JNEUROSCI.1932-20.2021

Azzalini, D., Rebollo, I., & Tallon-Baudry, C. (2019). Visceral Signals Shape Brain Dynamics

and Cognition. *Trends in Cognitive Sciences*, *23*(6), 488–509.

https://doi.org/10.1016/j.tics.2019.03.007

Babo-Rebelo, M., Buot, A., & Tallon-Baudry, C. (2019). Neural responses to heartbeats

distinguish self from other during imagination. *NeuroImage*, *191*, 10–20.

https://doi.org/10.1016/j.neuroimage.2019.02.012

Babo-Rebelo, M., Richter, C. G., & Tallon-Baudry, C. (2016). Neural responses to

heartbeats in the default network encode the self in spontaneous thoughts.

Journal of Neuroscience, *36*(30), 7829–7840.

https://doi.org/10.1523/JNEUROSCI.0262-16.2016

Babo-Rebelo, M., Wolpert, N., Adam, C., Hasboun, D., & Tallon-Baudry, C. (2016). Is the

cardiac monitoring function related to the self in both the default network and

right anterior insula? *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *371*(1708). https://doi.org/10.1098/rstb.2016.0004

Banellis, L., & Cruse, D. (2020). Skipping a beat: Heartbeat-evoked potentials reflect

predictions during interoceptive-exteroceptive integration. *Cerebral Cortex*

Communications, 1–14. https://doi.org/10.1093/texcom/tgaa060

Baranauskas, M., Grabauskaitė, A., & Griškova-Bulanova, I. (2017). Brain responses and

self-reported indices of interoception: Heartbeat evoked potentials are inversely

associated with worrying about body sensations. *Physiology and Behavior*, *180*,

1–7. https://doi.org/10.1016/j.physbeh.2017.07.032

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society:*

Series B (Methodological), *57*(1), 289–300. https://doi.org/10.1111/j.2517-

6161.1995.tb02031.x

Billeci, L., Faraguna, U., Santarcangelo, E. L., d'Ascanio, P., Varanini, M., & Sebastiani, L.

(2021). Heartbeat-Evoked Cortical Potential during Sleep and Interoceptive

Sensitivity: A Matter of Hypnotizability. *Brain Sciences*, *11*(8), 1089.

https://doi.org/10.3390/brainsci11081089

Birba, A., Santamaría-García, H., Prado, P., Cruzat, J., Ballesteros, A. S., Legaz, A.,

Fittipaldi, S., Duran-Aniotz, C., Slachevsky, A., Santibañez, R., Sigman, M.,

García, A. M., Whelan, R., Moguilner, S., & Ibáñez, A. (2022). Allostatic-

Interoceptive Overload in Frontotemporal Dementia. *Biological Psychiatry*, *92*(1),

54–67. https://doi.org/10.1016/j.biopsych.2022.02.955

Bogdány, T., Perakakis, P., Bódizs, R., & Simor, P. (2022). The heartbeat evoked potential

is a questionable biomarker in nightmare disorder: A replication study.

NeuroImage: Clinical, *33*. https://doi.org/10.1016/j.nicl.2021.102933

Botvinik-Nezer, R., & Wager, T. D. (2023). Reproducibility in Neuroimaging Analysis:

Challenges and Solutions. *Biological Psychiatry. Cognitive Neuroscience and*

Neuroimaging, *8*(8), 780–788. https://doi.org/10.1016/j.bpsc.2022.12.006

Boudewyn, M. A., Luck, S. J., Farrens, J. L., & Kappenman, E. S. (2018). How many trials

does it take to get a significant ERP effect? It depends. *Psychophysiology*, *55*(6),

e13049. https://doi.org/10.1111/psyp.13049

Buot, A., Azzalini, D., Chaumon, M., & Tallon-Baudry, C. (2021). Does stroke volume

influence heartbeat evoked responses? *Biological Psychology*, *165*, 108165.

https://doi.org/10.1016/j.biopsycho.2021.108165

Button, K. S., Ioannidis, J. P. A., Mokrysz, C., Nosek, B. A., Flint, J., Robinson, E. S. J., & Munafò, M. R. (2013). Power failure: Why small sample size undermines the reliability of neuroscience. *Nature Reviews. Neuroscience*, *14*(5), 365–376. https://doi.org/10.1038/nrn3475

Callara, A. L., Fontanelli, L., Belcari, I., Rho, G., Greco, A., Zelič, Ž., Sebastiani, L., & Santarcangelo, E. L. (2023). Modulation of the heartbeat evoked cortical potential by hypnotizability and hypnosis. *Psychophysiology*, *60*(9), e14309. https://doi.org/10.1111/psyp.14309

Cambi, S., Solcà, M., Micali, N., & Berchio, C. (2024). Cardiac interoception in Anorexia Nervosa: A resting-state heartbeat-evoked potential study. *European Eating Disorders Review*, *32*(3), 417–430. https://doi.org/10.1002/erv.3049

Campos Viola, F., Thorne, J., Edmonds, B., Schneider, T., Eichele, T., & Debener, S. (2009). Semi-automatic identification of independent components representing EEG artifact. *Clinical Neurophysiology*, *120*(5), 868–877. https://doi.org/10.1016/j.clinph.2009.01.015

Canales-Johnson, A., Silva, C., Huepe, D., Rivera-Rei, Á., Noreika, V., Del Carmen Garcia, M., Silva, W., Ciraolo, C., Vaucheret, E., Sedeño, L., Couto, B., Kargieman, L., Baglivo, F., Sigman, M., Chennu, S., Ibáñez, A., Rodríguez, E., & Bekinschtein, T. A. (2015). Auditory feedback differentially modulates behavioral and neural markers of objective and subjective performance when tapping to your heartbeat. *Cerebral Cortex*, *25*(11), 4490–4503. https://doi.org/10.1093/cercor/bhv076

Candia-Rivera, D., Annen, J., Gosseries, O., Martial, C., Thibaut, A., Laureys, S., & Tallon-Baudry, C. (2021). Neural responses to heartbeats detect residual signs of consciousness during resting state in postcomatose patients. *Journal of Neuroscience*, *41*(24), 5251–5262. https://doi.org/10.1523/JNEUROSCI.1740-20.2021

Candia-Rivera, D., Catrambone, V., & Valenza, G. (2021). The role of electroencephalography electrical reference in the assessment of functional

brain–heart interplay: From methodology to user guidelines. *Journal of Neuroscience Methods*, *360*, 109269. https://doi.org/10.1016/j.jneumeth.2021.109269

Candia-Rivera, D., & Machado, C. (2023a). Multidimensional assessment of heartbeat-evoked responses in disorders of consciousness. *European Journal of Neuroscience*, *58*(4), 3098–3110. https://doi.org/10.1111/ejn.16079

Candia-Rivera, D., & Machado, C. (2023b). Reduced Heartbeat-Evoked Responses in a Near-Death Case Report. *Journal of Clinical Neurology*, *19*(6), 581. https://doi.org/10.3988/jcn.2022.0415

Candia-Rivera, D., Raimondo, F., Pérez, P., Naccache, L., Tallon-Baudry, C., & Sitt, J. D. (2023). Conscious processing of global and local auditory irregularities causes differentiated heartbeat-evoked responses. *eLife*, *12*, e75352. https://doi.org/10.7554/eLife.75352

Caparco, A., Lopez-Martin, G., & Galvez-Pol, A. (2025). Phase confusion: How inconsistent cardiac labeling obscures interoception research. *Biological Psychology*, *199*, 109078. https://doi.org/10.1016/j.biopsycho.2025.109078

Cataldi, J., Pelentritou, A., Schwartz, S., & De Lucia, M. (2025). Impact of the pulse artifact on evoked activity in human wakefulness and sleep. *bioRxiv*, 2025.11.20.689416. https://doi.org/10.1101/2025.11.20.689416

Chambers, C. D., & Tzavella, L. (2022). The past, present and future of Registered Reports. *Nature Human Behaviour*, *6*(1), 29–42. https://doi.org/10.1038/s41562-021-01193-7

Champely, S., Ekstrom, C., Dalgaard, P., Gill, J., Weibelzahl, S., Anandkumar, A., Ford, C., Volcic, R., & Rosario, H. D. (2020). *pwr: Basic Functions for Power Analysis* (Version 1.3-0) [Computer software]. https://cran.r-project.org/web/packages/pwr/index.html

Chaumon, M., Bishop, D. V. M., & Busch, N. A. (2015). A practical guide to the selection of independent components of the electroencephalogram for artifact correction.

Journal of Neuroscience Methods, *250*, 47–63.

https://doi.org/10.1016/j.jneumeth.2015.02.025

Citation diversity statement. (2023). *Nature Reviews Bioengineering*, *1*(4), 227–227.

https://doi.org/10.1038/s44222-023-00061-5

Clayson, P. E., Baldwin, S. A., Rocha, H. A., & Larson, M. J. (2021). The data-processing

multiverse of event-related potentials (ERPs): A roadmap for the optimization

and standardization of ERP processing and reduction pipelines. *NeuroImage*,

245(November), 118712. https://doi.org/10.1016/j.neuroimage.2021.118712

Coll, M.-P., Hobson, H., Bird, G., & Murphy, J. (2021). Systematic review and meta-analysis

of the relationship between the heartbeat-evoked potential and interoception.

Neuroscience and Biobehavioral Reviews, *122*, 190–200.

https://doi.org/10.1016/j.neubiorev.2020.12.012

Couto, B., Adolfi, F., Velasquez, M., Mesow, M., Feinstein, J., Canales-Johnson, A., Mikulan,

E., Martínez-Pernía, D., Bekinschtein, T., Sigman, M., Manes, F., & Ibanez, A.

(2015). Heart evoked potential triggers brain responses to natural affective

scenes: A preliminary study. *Autonomic Neuroscience: Basic & Clinical*, *193*,

132–137. https://doi.org/10.1016/j.autneu.2015.06.006

Couto, B., Salles, A., Sedeño, L., Peradejordi, M., Barttfeld, P., Canales-Johnson, A., Dos

Santos, Y. V., Huepe, D., Bekinschtein, T., Sigman, M., Favaloro, R., Manes, F.,

& Ibanez, A. (2014). The man who feels two hearts: The different pathways of

interoception. *Social Cognitive and Affective Neuroscience*, *9*(9), 1253–1260.

https://doi.org/10.1093/scan/nst108

Criscuolo, A., Schwartze, M., & Kotz, S. A. (2022). Cognition through the lens of a body–

brain dynamic system. *Trends in Neurosciences*, *45*(9), 667–677.

https://doi.org/10.1016/j.tins.2022.06.004

De Falco, E., Solcà, M., Bernasconi, F., Babo-Rebelo, M., Young, N., Sammartino, F.,

Tallon-Baudry, C., Navarro, V., Rezai, A. R., Krishna, V., & Blanke, O. (2024).

Single neurons in the thalamus and subthalamic nucleus process cardiac and

respiratory signals in humans. *Proceedings of the National Academy of Sciences*, *121*(11), e2316365121. https://doi.org/10.1073/pnas.2316365121

de la Fuente, A., Sedeño, L., Vignaga, S. S., Ellmann, C., Sonzogni, S., Belluscio, L., García-Cordero, I., Castagnaro, E., Boano, M., Cetkovich, M., Torralva, T., Cánepa, E. T., Tagliazucchi, E., Garcia, A. M., & Ibañez, A. (2019). Multimodal neurocognitive markers of interoceptive tuning in smoked cocaine. *Neuropsychopharmacology*, *44*(8), 1425–1434. https://doi.org/10.1038/s41386-019-0370-3

Di Bernardi Luft, C., & Bhattacharya, J. (2015). Aroused with heart: Modulation of heartbeat evoked potential by arousal induction and its oscillatory correlates. *Scientific Reports*, *5*(March), 1–11. https://doi.org/10.1038/srep15717

Dirlich, G., Dietl, T., Vogl, L., & Strian, F. (1998). Topography and morphology of heart action-related EEG potentials. *Electroencephalography and Clinical Neurophysiology*, *108*(3), 299–305. https://doi.org/10.1016/s0168-5597(98)00003-3

Dirlich, G., Vogl, L., Plaschke, M., & Strian, F. (1997). Cardiac field effects on the EEG. *Electroencephalography and Clinical Neurophysiology*, *102*(4), 307–315. https://doi.org/10.1016/s0013-4694(96)96506-2

Dworkin, J. D., Linn, K. A., Teich, E. G., Zurn, P., Shinohara, R. T., & Bassett, D. S. (2020). The extent and drivers of gender imbalance in neuroscience reference lists. *Nature Neuroscience*, *23*(8), 918–926. https://doi.org/10.1038/s41593-020-0658-y

Ehinger, B. V., & Dimigen, O. (2019). Unfold: An integrated toolbox for overlap correction, non-linear modeling, and regression-based EEG analysis. *PeerJ*, *2019*(10), 1–33. https://doi.org/10.7717/peerj.7838

Elkommos, S., Martin-Lopez, D., Koreki, A., Jolliffe, C., Kandasamy, R., Mula, M., Critchley, H. D., Edwards, M. J., Garfinkel, S., Richardson, M. P., & Yogarajah, M. (2023). Changes in the heartbeat-evoked potential are associated with functional

seizures. *Journal of Neurology, Neurosurgery & Psychiatry*, *94*(9), 769–775.

https://doi.org/10.1136/jnnp-2022-330167

Elson, M. (2019). Examining Psychological Science Through Systematic Meta-Method

Analysis: A Call for Research. *Advances in Methods and Practices in*

Psychological Science, *2*(4), 350–363.

https://doi.org/10.1177/2515245919863296

Engelen, T., Buot, A., Grèzes, J., & Tallon-Baudry, C. (2023). Whose emotion is it?

Perspective matters to understand brain-body interactions in emotions.

NeuroImage, *268*, 119867. https://doi.org/10.1016/j.neuroimage.2023.119867

Engelen, T., Solcà, M., & Tallon-Baudry, C. (2023). Interoceptive rhythms in the brain.

Nature Neuroscience. https://doi.org/10.1038/s41593-023-01425-1

Feldman, M. J., Bliss-Moreau, E., & Lindquist, K. A. (2024). The neurobiology of

interoception and affect. *Trends in Cognitive Sciences*, *28*(7), 643–661.

https://doi.org/10.1016/j.tics.2024.01.009

Fittipaldi, S., Abrevaya, S., Fuente, A. de la, Pascariello, G. O., Hesse, E., Birba, A.,

Salamone, P., Hildebrandt, M., Martí, S. A., Pautassi, R. M., Huepe, D.,

Martorell, M. M., Yoris, A., Roca, M., García, A. M., Sedeño, L., & Ibáñez, A.

(2020). A multidimensional and multi-feature framework for cardiac interoception.

NeuroImage, *212*, 116677. https://doi.org/10.1016/j.neuroimage.2020.116677

Flasbeck, V., Bamberg, C., & Brüne, M. (2021). Short-Term Fasting and Ingestion of Caloric

Drinks Affect Heartbeat-Evoked Potentials and Autonomic Nervous System

Activity in Males. *Frontiers in Neuroscience*, *15*.

https://www.frontiersin.org/articles/10.3389/fnins.2021.622428

Flasbeck, V., Jungilligens, J., Lemke, I., Beckers, J., Öztürk, H., Wellmer, J., Seliger, C.,

Juckel, G., & Popkirov, S. (2024). Heartbeat evoked potentials and autonomic

arousal during dissociative seizures: Insights from electrophysiology and

neuroimaging. *BMJ Neurology Open*, *6*(1), e000665.

https://doi.org/10.1136/bmjno-2024-000665

Flasbeck, V., Popkirov, S., Ebert, A., & Brüne, M. (2020). Altered interoception in patients with borderline personality disorder: A study using heartbeat-evoked potentials. *Borderline Personality Disorder and Emotion Dysregulation*, 7(1), 24. https://doi.org/10.1186/s40479-020-00139-1

Fouragnan, E. F., Hosking, B., Cheung, Y., Prakash, B., Rushworth, M., & Sel, A. (2024). Timing along the cardiac cycle modulates neural signals of reward-based learning. *Nature Communications*, 15(1), 2976. https://doi.org/10.1038/s41467-024-46921-5

Frossard, J., & Renaud, O. (2022). The cluster depth tests: Toward point-wise strong control of the family-wise error rate in massively univariate tests with application to M/EEG. *NeuroImage*, 247, 118824. https://doi.org/10.1016/j.neuroimage.2021.118824

Fukushima, H., Terasawa, Y., & Umeda, S. (2011). Association between interoception and empathy: Evidence from heartbeat-evoked brain potential. *International Journal of Psychophysiology*, 79(2), 259–265. https://doi.org/10.1016/j.ijpsycho.2010.10.015

Fulvio, J. M., Akinnola, I., & Postle, B. R. (2021). Gender (Im)balance in Citation Practices in Cognitive Neuroscience. *Journal of Cognitive Neuroscience*, 33(1), 3–7. https://doi.org/10.1162/jocn_a_01643

Galvez-Pol, A., Virdee, P., Villacampa, J., & Kilner, J. (2022). Active tactile discrimination is coupled with and modulated by the cardiac cycle. *eLife*, 11, e78126. https://doi.org/10.7554/eLife.78126

Gao, J., Sun, R., Leung, H. K., Roberts, A., Wu, B. W. Y., Tsang, E. W., Tang, A. C. W., & Sik, H. H. (2023). Increased neurocardiological interplay after mindfulness meditation: A brain oscillation-based approach. *Frontiers in Human Neuroscience*, 17, 1008490. https://doi.org/10.3389/fnhum.2023.1008490

García-Cordero, I., Esteves, S., Mikulan, E. P., Hesse, E., Baglivo, F. H., Silva, W., García, M. del C., Vaucheret, E., Ciraolo, C., García, H. S., Adolfi, F., Pietto, M., Herrera,

E., Legaz, A., Manes, F., García, A. M., Sigman, M., Bekinschtein, T. A., Ibáñez,
A., & Sedeño, L. (2017). Attention, in and out: Scalp-level and intracranial EEG
correlates of interoception and exteroception. *Frontiers in Neuroscience*,
11(JUL). https://doi.org/10.3389/fnins.2017.00411

García-Cordero, I., Sedeno, L., Slachevsky, A., Forno, G., Klein, F., Lillo, P., Ferrari, J.,
Rodriguez, C., Bustin, J., Torralva, T., Baez, S., Yoris, A., Esteves, S., Melloni,
M., Salamone, P., Huepe, D., Manes, F., Garcıa, A. M., & Ibanez, A. (2016).
Feeling, learning from and being aware of inner states: Interoceptive dimensions
in neurodegeneration and stroke. (371).

Garfinkel, S. N., Minati, L., Gray, M. A., Seth, A. K., Dolan, R. J., & Critchley, H. D. (2014).
Fear from the Heart: Sensitivity to Fear Stimuli Depends on Individual
Heartbeats. *The Journal of Neuroscience*, *34*(19), 6573–6582.
https://doi.org/10.1523/JNEUROSCI.3507-13.2014

Gautier, R., Latinus, M., & Briend, F. (2025). Characterizing the Heartbeat-Evoked Potential:
A Two-Component Model of Cardiac Signal Processing? *Psychophysiology*,
62(12), e70206. https://doi.org/10.1111/psyp.70206

Gentsch, A., Sel, A., Marshall, A. C., & Schütz-Bosbach, S. (2019). Affective interoceptive
inference: Evidence from heart-beat evoked brain potentials. *Human Brain*
Mapping, *40*(1), 20–33. https://doi.org/10.1002/hbm.24352

Gibney, K. D., Kypriotakis, G., Cinciripini, P. M., Robinson, J. D., Minnix, J. A., & Versace, F.
(2020). Estimating statistical power for event-related potential studies using the
late positive potential. *Psychophysiology*, *57*(2), e13482.
https://doi.org/10.1111/psyp.13482

Giusti, G., Zelič, Ž., Callara, A. L., Sebastiani, L., & Santarcangelo, E. L. (2024).
Interoception as a function of hypnotizability during rest and a heartbeat counting
task. *Psychophysiology*, *61*(6), e14535. https://doi.org/10.1111/psyp.14535

Gray, M. A., Taggart, P., Sutton, P. M., Groves, D., Holdright, D. R., Bradbury, D., Brull, D.,
& Critchley, H. D. (2007). A cortical potential reflecting cardiac function.

Proceedings of the National Academy of Sciences of the United States of

America, *104*(16), 6818–6823. https://doi.org/10.1073/pnas.0609509104

Hall, J. E., & Hall, M. E. (2021). *Guyton and Hall textbook of medical physiology* (14th

edition). Elsevier.

Hardwicke, T. E., & Wagenmakers, E.-J. (2023). Reducing bias, increasing transparency

and calibrating confidence with preregistration. *Nature Human Behaviour*, *7*(1),

15–26. https://doi.org/10.1038/s41562-022-01497-2

Hedges, L. V., & Olkin, I. (1985). *Statistical Methods for Meta-Analysis*. Academic Press.

Hefter, K., Teich, E. G., & Bassett, D. S. (2025). *Gender imbalance in citations most*

pronounced in high-impact neuroscience journals (p. 2025.02.19.638905).

bioRxiv. https://doi.org/10.1101/2025.02.19.638905

Hermann, B., Candia-Rivera, D., Sharshar, T., Gavaret, M., Diehl, J., Cariou, A., &

Benghanem, S. (2024). Aberrant brain–heart coupling is associated with the

severity of post cardiac arrest brain injury. *Annals of Clinical and Translational*

Neurology, *11*(4), 866–882. https://doi.org/10.1002/acn3.52000

Hodossy, L., Ainley, V., & Tsakiris, M. (2021). How do we relate to our heart?

Neurobehavioral differences across three types of engagement with cardiac

interoception. *Biological Psychology*, *165*, 108198.

https://doi.org/10.1016/j.biopsycho.2021.108198

Immanuel, S. A., Pamula, Y., Kohler, M., Martin, J., Kennedy, D., Nalivaiko, E., Saint, D. A.,

& Baumert, M. (2014). Heartbeat evoked potentials during sleep and daytime

behavior in children with sleep-disordered breathing. *American Journal of*

Respiratory and Critical Care Medicine, *190*(10), 1149–1157.

https://doi.org/10.1164/rccm.201405-0920OC

Ito, Y., Shibata, M., Tanaka, Y., Terasawa, Y., & Umeda, S. (2019). Affective and temporal

orientation of thoughts: Electrophysiological evidence. *Brain Research*, *1719*,

148–156. https://doi.org/10.1016/j.brainres.2019.05.041

Jammal Salameh, L., Bitzenhofer, S. H., Hanganu-Opatz, I. L., Dutschmann, M., & Egger, V. (2024). Blood pressure pulsations modulate central neuronal activity via mechanosensitive ion channels. *Science*, *383*(6682), eadk8511. https://doi.org/10.1126/science.adk8511

Judah, M. R., Shurkova, E. Y., Hager, N. M., White, E. J., Taylor, D. L., & Grant, D. M. M. (2018). The relationship between social anxiety and heartbeat evoked potential amplitude. *Biological Psychology*, *139*(December 2017), 1–7. https://doi.org/10.1016/j.biopsycho.2018.09.013

Kamp, S.-M., Buntić, N., Amtmann, J., Scharpf, A., Schönen, A., Wagner, L., & Schulz, A. (2023). Reduced concentration performance and heartbeat-evoked potential in individuals with a history of a SARS-CoV-2 infection. *Neuroscience Letters*, *814*, 137466. https://doi.org/10.1016/j.neulet.2023.137466

Kamp, S.-M., Schulz, A., Forester, G., & Domes, G. (2021). Older adults show a higher heartbeat-evoked potential than young adults and a negative association with everyday metacognition. *Brain Research*, *1752*(January), 147238. https://doi.org/10.1016/j.brainres.2020.147238

Katkin, E. S., Cestaro, V. L., & Weitkunat, R. (1991). Individual differences in cortical evoked potentials as a function of heartbeat detection ability. *The International Journal of Neuroscience*, *61*(3–4), 269–276. https://doi.org/10.3109/00207459108990745

Kato, Y., Takei, Y., Umeda, S., Mimura, M., & Fukuda, M. (2020). Alterations of Heartbeat Evoked Magnetic Fields Induced by Sounds of Disgust. *Frontiers in Psychiatry*, *11*, 683. https://doi.org/10.3389/fpsyt.2020.00683

Keil, A., Debener, S., Gratton, G., Junghöfer, M., Kappenman, E. S., Luck, S. J., Luu, P., Miller, G. A., & Yee, C. M. (2014). Committee report: Publication guidelines and recommendations for studies using electroencephalography and magnetoencephalography. *Psychophysiology*, *51*(1), 1–21. https://doi.org/10.1111/psyp.12147

Kern, M., Aertsen, A., Schulze-Bonhage, A., & Ball, T. (2013). Heart cycle-related effects on

event-related potentials, spectral power changes, and connectivity patterns in the

human ECoG. *NeuroImage*, *81*, 178–190.

https://doi.org/10.1016/j.neuroimage.2013.05.042

Khoshnoud, S., Alvarez Igarzábal, F., & Wittmann, M. (2022). Brain–Heart Interaction and

the Experience of Flow While Playing a Video Game. *Frontiers in Human*

Neuroscience, *16*, 819834. https://doi.org/10.3389/fnhum.2022.819834

Khoshnoud, S., Leitritz, D., Çinar Bozdağ, M., Alvarez Igarzábal, F., Noreika, V., &

Wittmann, M. (2024). When the Heart Meets the Mind: Exploring the Brain–Heart

Interaction during Time Perception. *The Journal of Neuroscience*, *44*(34),

e2039232024. https://doi.org/10.1523/JNEUROSCI.2039-23.2024

Kim, J., & Jeong, B. (2019). Heartbeat induces a cortical theta-synchronized network in the

resting state. *eNeuro*, *6*(4), 1–11. https://doi.org/10.1523/ENEURO.0200-

19.2019

Kim, J., Park, H. D., Kim, K. W., Shin, D. W., Lim, S., Kwon, H., Kim, M. Y., Kim, K., &

Jeong, B. (2019). Sad faces increase the heartbeat-associated interoceptive

information flow within the salience network: A MEG study. *Scientific Reports*,

9(1), 1–15. https://doi.org/10.1038/s41598-018-36498-7

Koreki, A. (2024). Aberrant heartbeat-evoked potential in schizophrenia. *Progress in*

Neuropsychopharmacology.

Kriegeskorte, N., Lindquist, M. A., Nichols, T. E., Poldrack, R. A., & Vul, E. (2010).

Everything You Never Wanted to Know about Circular Analysis, but Were Afraid

to Ask. *Journal of Cerebral Blood Flow & Metabolism*, *30*(9), 1551–1557.

https://doi.org/10.1038/jcbfm.2010.86

Kriegeskorte, N., Simmons, W. K., Bellgowan, P. S. F., & Baker, C. I. (2009). Circular

analysis in systems neuroscience: The dangers of double dipping. *Nature*

Neuroscience, *12*(5), 535–540. https://doi.org/10.1038/nn.2303

Kritzman, L., Eidelman-Rothman, M., Keil, A., Freche, D., Sheppes, G., & Levit-Binnun, N.

(2022). Steady-state visual evoked potentials differentiate between internally and

externally directed attention. *NeuroImage*, *254*(March), 119133.

https://doi.org/10.1016/j.neuroimage.2022.119133

Kruschke, J. K. (2021). Bayesian Analysis Reporting Guidelines. *Nature Human Behaviour*,

5(10), 1282–1291. https://doi.org/10.1038/s41562-021-01177-7

Kumral, D., Al, E., Cesnaite, E., Kornej, J., Sander, C., Hensch, T., Zeynalova, S.,

Tautenhahn, S., Hagendorf, A., Laufs, U., Wachter, R., Nikulin, V., & Villringer,

A. (2022). Attenuation of the Heartbeat-Evoked Potential in Patients With Atrial

Fibrillation. *JACC: Clinical Electrophysiology*, *8*(10), 1219–1230.

https://doi.org/10.1016/j.jacep.2022.06.019

Kunzendorf, S., Klotzsche, F., Akbal, M., Villringer, A., Ohl, S., & Gaebler, M. (2019). Active

information sampling varies across the cardiac cycle. *Psychophysiology*, *56*(5),

e13322. https://doi.org/10.1111/psyp.13322

Lakens, D., Scheel, A. M., & Isager, P. M. (2018). Equivalence Testing for Psychological

Research: A Tutorial. *Advances in Methods and Practices in Psychological*

Science, *1*(2), 259–269. https://doi.org/10.1177/2515245918770963

Larson, M. J., & Carbine, K. A. (2017). Sample size calculations in human electrophysiology

(EEG and ERP) studies: A systematic review and recommendations for

increased rigor. *International Journal of Psychophysiology, Rigor and*

Replication: Towards Improved Best Practices in Psychophysiological Research,

111, 33–41. https://doi.org/10.1016/j.ijpsycho.2016.06.015

Lechinger, J., Heib, D. P. J., Gruber, W., Schabus, M., & Klimesch, W. (2015). Heartbeat-

related EEG amplitude and phase modulations from wakefulness to deep sleep:

Interactions with sleep spindles and slow oscillations. *Psychophysiology*, *52*(11),

1441–1450. https://doi.org/10.1111/psyp.12508

Legaz, A., Yoris, A., Sedeño, L., Abrevaya, S., Martorell, M., Alifano, F., García, A. M., &

Ibañez, A. (2022). Heart–brain interactions during social and cognitive stress in

hypertensive disease: A multidimensional approach. In *European Journal of*

Neuroscience (Vol. 55, Issues 9–10). https://doi.org/10.1111/ejn.14979

Leopold, C., & Schandry, R. (2001). *The heartbeat-evoked brain potential in patients*

suffering from diabetic neuropathy and in healthy control persons. 112, 674–682.

Liu, H., Liang, H., Yu, X., Wang, G., Han, Y., Yan, M., Li, S., & Wang, W. (2023). Enhanced

external counterpulsation modulates the heartbeat evoked potential. *Frontiers in*

Physiology, 14, 1144073. https://doi.org/10.3389/fphys.2023.1144073

Liu, H., Yu, X., Wang, G., Han, Y., & Wang, W. (2022). Effects of 24-h acute total sleep

deprivation on physiological coupling in healthy young adults. *Frontiers in*

Neuroscience, 16, 952329. https://doi.org/10.3389/fnins.2022.952329

Liuzzi, P., Cassioli, T., Secci, S., Hakiki, B., Scarpino, M., Burali, R., di Palma, A., Toci, T.,

Grippo, A., Cecchi, F., Frosini, A., & Mannini, A. (2024). A neurophysiological

profiling of the heartbeat-evoked potential in severe acquired brain injuries: A

focus on unconsciousness. *The European Journal of Neuroscience, 60*(3),

4201–4216. https://doi.org/10.1111/ejn.16394

Luck, S. J. (2014). *An introduction to the event-related potential technique* (Second edition).

The MIT Press.

Luck, S. J., & Gaspelin, N. (2017). How to Get Statistically Significant Effects in Any ERP

Experiment (and Why You Shouldn't). *Psychophysiology, 54*(1), 146–157.

https://doi.org/10.1111/psyp.12639

Lutz, A. P. C., Schulz, A., Voderholzer, U., Koch, S., van Dyck, Z., & Vögele, C. (2019).

Enhanced cortical processing of cardio-afferent signals in anorexia nervosa.

Clinical Neurophysiology, 130(9), 1620–1627.

https://doi.org/10.1016/j.clinph.2019.06.009

MacKinnon, S., Gevirtz, R., McCraty, R., & Brown, M. (2013). Utilizing Heartbeat Evoked

Potentials to Identify Cardiac Regulation of Vagal Afferents During Emotion and

Resonant Breathing. *Applied Psychophysiology and Biofeedback, 38*(4), 241–

255. https://doi.org/10.1007/s10484-013-9226-5

Mai, S., Wong, C. K., Georgiou, E., & Pollatos, O. (2018). Interoception is associated with

heartbeat-evoked brain potentials (HEPs) in adolescents. *Biological Psychology*,

137, 24–33. https://doi.org/10.1016/j.biopsycho.2018.06.007

Maister, L., Tang, T., & Tsakiris, M. (2017). Neurobehavioral evidence of interoceptive

sensitivity in early infancy. *eLife*, *6*, 1–12. https://doi.org/10.7554/eLife.25318

Majeed, U., Aftab, M. F., Baloch, D. M., Ahmed, S., Yusuf, I. M., Hasan, M. A., & Qureshi, M.

S. (2022). Modulation of Heart and Brain Function by Surah Al-Rehman

Recitation Among Distressed Diabetic Patients in Pakistan. *Journal of Religion*

and Health, *61*(5), 3852–3865. https://doi.org/10.1007/s10943-021-01431-2

Maris, E., & Oostenveld, R. (2007). Nonparametric statistical testing of EEG- and MEG-data.

Journal of Neuroscience Methods, *164*(1), 177–190.

https://doi.org/10.1016/j.jneumeth.2007.03.024

Marshall, A. C., Gentsch, A., Blum, A. L., Broering, C., & Schütz-Bosbach, S. (2019). I feel

what I do: Relating interoceptive processes and reward-related behavior.

NeuroImage, *191*(November 2018), 315–324.

https://doi.org/10.1016/j.neuroimage.2019.02.032

Marshall, A. C., Gentsch, A., Jelinčić, V., & Schütz-Bosbach, S. (2017). Exteroceptive

expectations modulate interoceptive processing: Repetition-suppression effects

for visual and heartbeat evoked potentials. *Scientific Reports*, *7*(1), 1–15.

https://doi.org/10.1038/s41598-017-16595-9

Marshall, A. C., Gentsch, A., Schröder, L., & Schütz-Bosbach, S. (2018). Cardiac

interoceptive learning is modulated by emotional valence perceived from facial

expressions. *Social Cognitive and Affective Neuroscience*, *13*(7), 677–686.

https://doi.org/10.1093/SCAN/NSY042

Marshall, A. C., Gentsch, A., & Schütz-Bosbach, S. (2020). Interoceptive cardiac

expectations to emotional stimuli predict visual perception. *Emotion*, *20*(7),

1113–1126. https://doi.org/10.1037/emo0000631

Marshall, A. C., Gentsch-Ebrahimzadeh, A., & Schütz-Bosbach, S. (2022). From the inside out: Interoceptive feedback facilitates the integration of visceral signals for efficient sensory processing: Interoceptive feedback facilitates sensory processing. *NeuroImage*, *251*(July 2021), 119011. https://doi.org/10.1016/j.neuroimage.2022.119011

McCraty, R., Atkinson, M., & Bradley, R. T. (2004). Electrophysiological Evidence of Intuition: Part 2. A System-Wide Process? *The Journal of Alternative and Complementary Medicine*, *10*(2), 325–336. https://doi.org/10.1089/107555304323062310

Melo, E., Fiel, J., Milhomens, R., Ribeiro, T., Navegantes, R., Gomes, F., Duarte Gomes, B., & Pereira, A. (2022). Dynamic coupling between the central and autonomic cardiac nervous systems in patients with refractory epilepsy: A pilot study. *Frontiers in Neurology*, *13*, 904052. https://doi.org/10.3389/fneur.2022.904052

Migeot, J., Hesse, E., Fittipaldi, S., Mejía, J., Fraile, M., García, A. M., García, M. D. C., Ortega, R., Lawlor, B., Lopez, V., & Ibáñez, A. (2023). Allostatic-interoceptive anticipation of social rejection. *NeuroImage*, *276*, 120200. https://doi.org/10.1016/j.neuroimage.2023.120200

Moher, D., Liberati, A., Tetzlaff, J., & Altman, D. G. (2009). Systematic Reviews and Meta-Analyses: The PRISMA Statement. *Annulas of Internal Medicine*, *151*(4), 264–269. https://doi.org/10.1371/journal.pmed1000097

Montoya, P., Schandry, R., & Müller, A. (1993). Heartbeat evoked potentials (HEP): Topography and influence of cardiac awareness and focus of attention. *Electroencephalography and Clinical Neurophysiology/ Evoked Potentials*, *88*(3), 163–172. https://doi.org/10.1016/0168-5597(93)90001-6

Müller, L. E., Schulz, A., Andermann, M., Gäbel, A., Gescher, D. M., Spohn, A., Herpertz, S. C., & Bertsch, K. (2015). Cortical representation of afferent bodily signals in borderline personality disorder: Neural correlates and relationship to emotional

dysregulation. *JAMA Psychiatry*, *72*(11), 1077–1086.

https://doi.org/10.1001/jamapsychiatry.2015.1252

Myllylä, T. S., Vihriälä, E. V., Korhonen, V. O., & Sorvoja, H. S. S. (2013). *Case study of ECG signal used as a reference signal in optical pulse transit time measurement of blood flow: The effect of different electrode placements on pulse transit time* (V. V. Tuchin, E. A. Genina, V. L. Derbov, & I. V. Meglinski, Eds.; p. 869903). https://doi.org/10.1117/12.2016682

Nalborczyk, L., & Bürkner, P. (2025). *Precise temporal localisation of M/EEG effects with Bayesian generalised additive multilevel models* (p. 2025.08.29.672336). bioRxiv. https://doi.org/10.1101/2025.08.29.672336

Niazy, R. K., Beckmann, C. F., Iannetti, G. D., Brady, J. M., & Smith, S. M. (2005). Removal of FMRI environment artifacts from EEG data using optimal basis sets. *NeuroImage*, *28*(3), 720–737. https://doi.org/10.1016/j.neuroimage.2005.06.067

Niizeki, K., Kawahara, K., & Miyamoto, Y. (1993). Interaction among cardiac, respiratory, and locomotor rhythms during cardiolocomotor synchronization. *Journal of Applied Physiology*, *75*(4), 1815–1821. https://doi.org/10.1152/jappl.1993.75.4.1815

Ohl, S., Wohltat, C., Kliegl, R., Pollatos, O., & Engbert, R. (2016). Microsaccades are coupled to heartbeat. *Journal of Neuroscience*, *36*(4), 1237–1241. https://doi.org/10.1523/JNEUROSCI.2211-15.2016

Ono, K., Mizuochi, R., Yamamoto, K., Sasaoka, T., & Ymawaki, S. (2024). Exploring the neural underpinnings of chord prediction uncertainty: An electroencephalography (EEG) study. *Scientific Reports*, *14*(1), 4586. https://doi.org/10.1038/s41598-024-55366-1

Pan, J., & Tompkins, W. J. (1985). A Real-Time QRS Detection Algorithm. *IEEE Transactions on Biomedical Engineering*, *BME-32*(3), 230–236. https://doi.org/10.1109/TBME.1985.325532

Pang, J., Tang, X., Li, H., Hu, Q., Cui, H., Zhang, L., Li, W., Zhu, Z., Wang, J., & Li, C.

(2019). Altered Interoceptive Processing in Generalized Anxiety Disorder—A

Heartbeat-Evoked Potential Research. *Frontiers in Psychiatry*, *10*(September),

1–9. https://doi.org/10.3389/fpsyt.2019.00616

Park, H.-D., Bernasconi, F., Bello-Ruiz, J., Pfeiffer, C., Salomon, R., & Blanke, O. (2016).

Transient modulations of neural responses to heartbeats covary with bodily self-

consciousness. *Journal of Neuroscience*, *36*(32), 8453–8460.

https://doi.org/10.1523/JNEUROSCI.0311-16.2016

Park, H.-D., & Blanke, O. (2019). Heartbeat-evoked cortical responses: Underlying

mechanisms, functional roles, and methodological considerations. *NeuroImage*,

197(April), 502–511. https://doi.org/10.1016/j.neuroimage.2019.04.081

Park, H.-D., Correia, S., Ducorps, A., & Tallon-Baudry, C. (2014). Spontaneous fluctuations

in neural responses to heartbeats predict visual detection. *Nature Neuroscience*,

17(4), 612–618. https://doi.org/10.1038/nn.3671

Pelentritou, A., Pfeiffer, C., Schwartz, S., & De Lucia, M. (2024). Cardio-audio

synchronization elicits neural and cardiac surprise responses in human

wakefulness and sleep. *Communications Biology*, *7*(1), 226.

https://doi.org/10.1038/s42003-024-05895-2

Perakakis, P. (2019). *HEPLAB: A Matlab graphical interface for the preprocessing of the

heartbeat-evoked potential* (Version v1.0.0) [Computer software]. Zenodo.

https://doi.org/10.5281/ZENODO.2649943

Pérez, J. J., Guijarro, E., & Barcia, J. A. (2005). Suppression of the cardiac electric field

artifact from the heart action evoked potential. *Medical and Biological

Engineering and Computing*, *43*(5), 572–581.

https://doi.org/10.1007/BF02351030

Perogamvros, L., Park, H.-D., Bayer, L., Perrault, A. A., Blanke, O., & Schwartz, S. (2019).

Increased heartbeat-evoked potential during REM sleep in nightmare disorder.

NeuroImage: Clinical, *22*(December 2018), 101701.

https://doi.org/10.1016/j.nicl.2019.101701

Petzschner, F. H., Weber, L. A., Wellstein, K. V., Paolini, G., Do, C. T., & Stephan, K. E.

(2019). Focus of attention modulates the heartbeat evoked potential.

NeuroImage, *186*(November 2018), 595–606.

https://doi.org/10.1016/j.neuroimage.2018.11.037

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated

electroencephalographic independent component classifier, dataset, and

website. *NeuroImage*, *198*(May), 181–197.

https://doi.org/10.1016/j.neuroimage.2019.05.026

Pollatos, O., Herbert, B. M., Mai, S., & Kammer, T. (2016). Changes in interoceptive

processes following brain stimulation. *Philosophical Transactions of the Royal*

Society B: Biological Sciences, *371*(1708).

https://doi.org/10.1098/rstb.2016.0016

Pollatos, O., Kirsch, W., & Schandry, R. (2005). Brain structures involved in interoceptive

awareness and cardioafferent signal processing: A dipole source localization

study. *Human Brain Mapping*, *26*(1), 54–64. https://doi.org/10.1002/hbm.20121

Pollatos, O., & Schandry, R. (2004). Accuracy of heartbeat perception is reflected in the

amplitude of the heartbeat-evoked brain potential. *Psychophysiology*, *41*(3),

476–482. https://doi.org/10.1111/1469-8986.2004.00170.x

Pontifex, M. B., Gwizdala, K. L., Parks, A. C., Billinger, M., & Brunner, C. (2017). Variability

of ICA decomposition may impact EEG signals when used to remove eyeblink

artifacts. *Psychophysiology*, *54*(3), 386–398. https://doi.org/10.1111/psyp.12804

Poppa, T., Benschop, L., Horczak, P., Vanderhasselt, M.-A., Carrette, E., Bechara, A.,

Baeken, C., & Vonck, K. (2022). Auricular transcutaneous vagus nerve

stimulation modulates the heart-evoked potential. *Brain Stimulation*, *15*(1), 260–

269. https://doi.org/10.1016/j.brs.2021.12.004

R Core Team. (2025). *R: A Language and Environment for Statistical Computing* [Computer

software]. R Foundation for Statistical Computing.

Rapp, L., Mai-Lippold, S. A., Georgiou, E., & Pollatos, O. (2023). Elevated EEG heartbeat-

evoked potentials in adolescents with more ADHD symptoms. *Biological*

Psychology, *184*, 108698. https://doi.org/10.1016/j.biopsycho.2023.108698

Ren, Q., Marshall, A. C., Kaiser, J., & Schütz-Bosbach, S. (2022). Response inhibition is

disrupted by interoceptive processing at cardiac systole. *Biological Psychology*,

170(August 2021), 108323. https://doi.org/10.1016/j.biopsycho.2022.108323

Richter, F., García, A. M., Rodriguez Arriagada, N., Yoris, A., Birba, A., Huepe, D., Zimmer,

H., Ibáñez, A., & Sedeño, L. (2021). Behavioral and neurophysiological

signatures of interoceptive enhancements following vagus nerve stimulation.

Human Brain Mapping, *42*(5), 1227–1242. https://doi.org/10.1002/hbm.25288

Richter, F., & Ibáñez, A. (2021). Time is body: Multimodal evidence of crosstalk between

interoception and time estimation. *Biological Psychology*, *159*(January).

https://doi.org/10.1016/j.biopsycho.2021.108017

Rousselet, G. A. (2025). Using cluster-based permutation tests to estimate MEG/EEG

onsets: How bad is it? *European Journal of Neuroscience*, *61*(1), e16618.

https://doi.org/10.1111/ejn.16618

Salamone, P. C., Esteves, S., Sinay, V. J., García-Cordero, I., Abrevaya, S., Couto, B.,

Adolfi, F., Martorell, M., Petroni, A., Yoris, A., Torquati, K., Alifano, F., Legaz, A.,

Cassará, F. P., Bruno, D., Kemp, A. H., Herrera, E., García, A. M., Ibáñez, A., &

Sedeño, L. (2018). Altered neural signatures of interoception in multiple

sclerosis. *Human Brain Mapping*, *39*(12), 4743–4754.

https://doi.org/10.1002/hbm.24319

Salamone, P. C., Legaz, A., Sedeño, L., Moguilner, S., Fraile-Vazquez, M., Campo, C. G.,

Fittipaldi, S., Yoris, A., Miranda, M., Birba, A., Galiani, A., Abrevaya, S., Neely,

A., Caro, M. M., Alifano, F., Villagra, R., Anunziata, F., Okada De Oliveira, M.,

Pautassi, R. M., … Ibañez, A. (2021). Interoception Primes Emotional

Processing: Multimodal Evidence from Neurodegeneration. *The Journal of Neuroscience*, *41*(19), 4276–4292. https://doi.org/10.1523/JNEUROSCI.2578-20.2021

Salamone, P. C., Sedeño, L., Legaz, A., Bekinschtein, T., Martorell, M., Adolfi, F., Fraile-Vazquez, M., Rodríguez Arriagada, N., Favaloro, L., Peradejordi, M., Absi, D. O., García, A. M., Favaloro, R., & Ibáñez, A. (2020). Dynamic neurocognitive changes in interoception after heart transplant. *Brain Communications*, *2*(2). https://doi.org/10.1093/braincomms/fcaa095

Sassenhagen, J., & Draschkow, D. (2019). Cluster-based permutation tests of MEG/EEG data do not establish significance of effect latency or location. *Psychophysiology*, *56*(6), e13335. https://doi.org/10.1111/psyp.13335

Schandry, R., & Montoya, P. (1996). Event-related brain potentials and the processing of cardiac activity. *Biological Psychology*, *42*(1–2), 75–85. https://doi.org/10.1016/0301-0511(95)05147-3

Schandry, R., Sparrer, B., & Weitkunat, R. (1986). From the heart to the brain: A study of heartbeat contingent scalp potentials. *International Journal of Neuroscience*, *30*(4), 261–275. https://doi.org/10.3109/00207458608985677

Schandry & Weitkunat. (1990). Enhancement of heartbeat-related brain potentials through cardiac awareness training. *International Journal of Neuroscience*, *53*(2–4), 243–253. https://doi.org/10.3109/00207459008986611

Schmitz, M., Müller, L. E., Schulz, A., Kleindienst, N., Herpertz, S. C., & Bertsch, K. (2020). Heart and brain: Cortical representation of cardiac signals is disturbed in borderline personality disorder, but unaffected by oxytocin administration. *Journal of Affective Disorders*, *264*, 24–28. https://doi.org/10.1016/j.jad.2019.11.139

Schmitz, M., Müller, L. E., Seitz, K. I., Schulz, A., Steinmann, S., Herpertz, S. C., & Bertsch, K. (2021). Heartbeat evoked potentials in patients with post-traumatic stress

disorder: An unaltered neurobiological regulation system? *European Journal of Psychotraumatology*, *12*(1). https://doi.org/10.1080/20008198.2021.1987686

Schulz, A., Ferreira de Sá, D. S., Dierolf, A. M., Lutz, A., van Dyck, Z., Vögele, C., & Schächinger, H. (2015). Short-term food deprivation increases amplitudes of heartbeat-evoked potentials. *Psychophysiology*, *52*(5), 695–703. https://doi.org/10.1111/psyp.12388

Schulz, A., Köster, S., Beutel, M. E., Schächinger, H., Vögele, C., Rost, S., Rauh, M., & Michal, M. (2015). Altered Patterns of Heartbeat-Evoked Potentials in Depersonalization/Derealization Disorder: Neurophysiological Evidence for Impaired Cortical Representation of Bodily Signals. *Psychosomatic Medicine*, *77*(5), 506. https://doi.org/10.1097/PSY.0000000000000195

Schulz, A., Rost, S., Flasinski, T., Dierolf, A. M., Lutz, A. P. C., Münch, E. E., Mertens, V.-C., Witthöft, M., & Vögele, C. (2020). Distinctive body perception mechanisms in high versus low symptom reporters: A neurophysiological model for medically-unexplained symptoms. *Journal of Psychosomatic Research*, *137*, 110223. https://doi.org/10.1016/j.jpsychores.2020.110223

Schulz, A., Stammet, P., Dierolf, A. M., Vögele, C., Beyenburg, S., Werer, C., & Devaux, Y. (2018). Late heartbeat-evoked potentials are associated with survival after cardiac arrest. *Resuscitation*, *126*, 7–13. https://doi.org/10.1016/j.resuscitation.2018.02.009

Schulz, A., Strelzyk, F., Ferreira de Sá, D. S., Naumann, E., Vögele, C., & Schächinger, H. (2013). Cortisol rapidly affects amplitudes of heartbeat-evoked brain potentials— Implications for the contribution of stress to an altered perception of physical sensations? *Psychoneuroendocrinology*, *38*(11), 2686–2693. https://doi.org/10.1016/j.psyneuen.2013.06.027

Sel, A., Azevedo, R. T., & Tsakiris, M. (2016). Heartfelt Self: Cardio-Visual Integration Affects Self-Face Recognition and Interoceptive Cortical Processing. *Cerebral Cortex*, cercor;bhw296v1. https://doi.org/10.1093/cercor/bhw296

Shannon, C. E. (1948). A mathematical theory of communicatio. *The Bell System Technical Journal, 27*(3), 379–423. https://doi.org/10.1002/j.1538-7305.1948.tb01338.x.

Shao, S., Shen, K., Wilder-Smith, E. P. V., & Li, X. (2011). Effect of pain perception on the heartbeat evoked potential. *Clinical Neurophysiology, 122*(9), 1838–1845. https://doi.org/10.1016/j.clinph.2011.02.014

Sheppard, J. P., Barker, T. A., Ranasinghe, A. M., Clutton-Brock, T. H., Frenneaux, M. P., & Parkes, M. J. (2011). Does modifying electrode placement of the 12 lead ECG matter in healthy subjects? *International Journal of Cardiology, 152*(2), 184–191. https://doi.org/10.1016/j.ijcard.2010.07.013

Simor, P., Bogdány, T., Bódizs, R., & Perakakis, P. (2021). Cortical monitoring of cardiac activity during rapid eye movement sleep: The heartbeat evoked potential in phasic and tonic rapid-eye-movement microstates. *Sleep, 44*(9), 1–12. https://doi.org/10.1093/sleep/zsab100

Skora, L. I., Livermore, J. J. A., & Roelofs, K. (2022). The functional role of cardiac activity in perception and action. *Neuroscience & Biobehavioral Reviews, 137*, 104655. https://doi.org/10.1016/j.neubiorev.2022.104655

Soderberg, C. K., Errington, T. M., Schiavone, S. R., Bottesini, J., Thorn, F. S., Vazire, S., Esterling, K. M., & Nosek, B. A. (2021). Initial evidence of research quality of registered reports compared with the standard publishing model. *Nature Human Behaviour, 5*(8), 990–997. https://doi.org/10.1038/s41562-021-01142-4

Solcà, M., Park, H.-D., Bernasconi, F., & Blanke, O. (2020). Behavioral and neurophysiological evidence for altered interoceptive bodily processing in chronic pain. *NeuroImage, 217*(April). https://doi.org/10.1016/j.neuroimage.2020.116902

Steegen, S., Tuerlinckx, F., Gelman, A., & Vanpaemel, W. (2016). Increasing Transparency Through a Multiverse Analysis. *Perspectives on Psychological Science, 11*(5), 702–712. https://doi.org/10.1177/1745691616658637

Steinfath, P., Herzog, N., Fourcade, A., Sander, C., Nikulin, V., & Villringer, A. (2025). Validating genuine changes in Heartbeat Evoked Potentials using Pseudotrials

and Surrogate Procedures. *Imaging Neuroscience*.

https://doi.org/10.1162/IMAG.a.30

Stoupi, N. A., Weijs, M. L., Imbach, L., & Lenggenhager, B. (2024). Heartbeat-evoked

potentials following voluntary hyperventilation in epilepsy patients: Respiratory

influences on cardiac interoception. *Frontiers in Neuroscience*, *18*, 1391437.

https://doi.org/10.3389/fnins.2024.1391437

Sweet, D. J. (2021). New at Cell Press: The Inclusion and Diversity Statement. *Cell*, *184*(1),

1–2. https://doi.org/10.1016/j.cell.2020.12.019

Tallon-Baudry, C., Campana, F., Park, H. D., & Babo-Rebelo, M. (2018). The neural

monitoring of visceral inputs, rather than attention, accounts for first-person

perspective in conscious vision. *Cortex*, *102*, 139–149.

https://doi.org/10.1016/j.cortex.2017.05.019

Tanaka, Y., Ito, Y., Terasawa, Y., & Umeda, S. (2023). Modulation of heartbeat-evoked

potential and cardiac cycle effect by auditory stimuli. *Biological Psychology*, *182*,

108637. https://doi.org/10.1016/j.biopsycho.2023.108637

Tanner, D., Morgan-Short, K., & Luck, S. J. (2015). How inappropriate high-pass filters can

produce artifactual effects and incorrect conclusions in ERP studies of language

and cognition. *Psychophysiology*, *52*(8), 997–1009.

https://doi.org/10.1111/psyp.12437

Tarvainen, M. P., Niskanen, J. P., Lipponen, J. A., Ranta-aho, P. O., & Karjalainen, P. A.

(2014). Kubios HRV - Heart rate variability analysis software. *Computer Methods*

and Programs in Biomedicine, *113*(1), 210–220.

https://doi.org/10.1016/j.cmpb.2013.07.024

Terhaar, J., Viola, F. C., Bär, K. J., & Debener, S. (2012). Heartbeat evoked potentials mirror

altered body perception in depressed patients. *Clinical Neurophysiology*,

123(10), 1950–1957. https://doi.org/10.1016/j.clinph.2012.02.086

Thornton, A., & Lee, P. (2000). Publication bias in meta-analysis: Its causes and

consequences. *Journal of Clinical Epidemiology*, *53*(2), 207–216.

https://doi.org/10.1016/S0895-4356(99)00161-4

Todd, J., Cardellicchio, P., Swami, V., Cardini, F., & Aspell, J. E. (2021). Weaker implicit

interoception is associated with more negative body image: Evidence from

gastric-alpha phase amplitude coupling and the heartbeat evoked potential.

Cortex, *143*(xxxx), 254–266. https://doi.org/10.1016/j.cortex.2021.07.006

Trübutschek, D., Yang, Y.-F., Gianelli, C., Cesnaite, E., Fischer, N. L., Vinding, M. C.,

Marshall, T. R., Algermissen, J., Pascarella, A., Puoliväli, T., Vitale, A., Busch, N.

A., & Nilsonne, G. (2024). EEGManyPipelines: A Large-scale, Grassroots Multi-

analyst Study of Electroencephalography Analysis Practices in the Wild. *Journal*

of Cognitive Neuroscience, *36*(2), 217–224.

https://doi.org/10.1162/jocn_a_02087

Umeda, S., Tochizawa, S., Shibata, M., & Terasawa, Y. (2016). Prospective memory

mediated by interoceptive accuracy: A psychophysiological approach.

Philosophical Transactions of the Royal Society B: Biological Sciences,

371(1708), 20160005. https://doi.org/10.1098/rstb.2016.0005

Ventura-Bort, C., & Weymar, M. (2024). Transcutaneous auricular vagus nerve stimulation

modulates the processing of interoceptive prediction error signals and their role

in allostatic regulation. *Human Brain Mapping*, *45*(3), e26613.

https://doi.org/10.1002/hbm.26613

Verdonk, C., Teed, A. R., White, E. J., Ren, X., Stewart, J. L., Paulus, M. P., & Khalsa, S. S.

(2024). Heartbeat-evoked neural response abnormalities in generalized anxiety

disorder during peripheral adrenergic stimulation. *Neuropsychopharmacology*,

49(8), 1246–1254. https://doi.org/10.1038/s41386-024-01806-5

Verdonk, C., Trousselard, M., Di Bernardi Luft, C., Medani, T., Billaud, J. B., Ramdani, C.,

Canini, F., Claverie, D., Jaumard-Hakoun, A., & Vialatte, F. (2021). The

heartbeat evoked potential does not support strong interoceptive sensibility in

trait mindfulness. *Psychophysiology, 58*(10), 1–13.

https://doi.org/10.1111/psyp.13891

Villena-González, M., Moënne-Loccoz, C., Lagos, R. A., Alliende, L. M., Billeke, P., Aboitiz,

F., López, V., & Cosmelli, D. (2017). Attending to the heart is associated with

posterior alpha band increase and a reduction in sensitivity to concurrent visual

stimuli. *Psychophysiology*, *54*(10), 1483–1497.

https://doi.org/10.1111/psyp.12894

Villena-González, M., Rojas-Thomas, F., Morales-Torres, R., & López, V. (2023).

Autonomous sensory meridian response is associated with a larger heartbeat-

evoked potential amplitude without differences in interoceptive awareness.

Psychophysiology, *60*(6), e14277. https://doi.org/10.1111/psyp.14277

Villringer, A., Nikulin, V. V., & Gaebler, M. (2025). Brain–body states as a link between

cardiovascular and mental health. *Trends in Neurosciences*.

https://doi.org/10.1016/j.tins.2025.08.004

Wagenmakers, E.-J., Marsman, M., Jamil, T., Ly, A., Verhagen, J., Love, J., Selker, R.,

Gronau, Q. F., Šmíra, M., Epskamp, S., Matzke, D., Rouder, J. N., & Morey, R.

D. (2018). Bayesian inference for psychology. Part I: Theoretical advantages and

practical ramifications. *Psychonomic Bulletin & Review*, *25*(1), 35–57.

https://doi.org/10.3758/s13423-017-1343-3

Wei, Y., Ramautar, J. R., Colombo, M. A., Stoffers, D., Gómez-Herrero, G., Van Der

Meijden, W. P., Te Lindert, B. H. W., Van Der Werf, Y. D., & Van Someren, E. J.

W. (2016). I Keep a Close Watch on This Heart of Mine: Increased Interoception

in Insomnia. *Sleep*, *39*(12), 2113–2124. https://doi.org/10.5665/sleep.6308

Weijs, M. L., Daum, M. M., & Lenggenhager, B. (2023). Cardiac interoception in infants:

Behavioral and neurophysiological measures in various emotional and self-

related contexts. *Psychophysiology*, *60*(12), e14386.

https://doi.org/10.1111/psyp.14386

Yasuma, F., & Hayano, J. (2004). Respiratory Sinus Arrhythmia. *Chest, 125*(2), 683–690.

https://doi.org/10.1378/chest.125.2.683

Yoder, N. (2026). *Peakfinder(x0, sel, thresh, extrema, includeEndpoints, interpolate)*

[Computer software]. MATLAB Central File Exchange.

https://www.mathworks.com/matlabcentral/fileexchange/25500-peakfinder-x0-

sel-thresh-extrema-includeendpoints-interpolate

Yoris, A., Abrevaya, S., Esteves, S., Salamone, P., Lori, N., Martorell, M., Legaz, A., Alifano,

F., Petroni, A., Sánchez, R., Sedeño, L., García, A. M., & Ibáñez, A. (2018).

Multilevel convergence of interoceptive impairments in hypertension: New

evidence of disrupted body–brain interactions. *Human Brain Mapping, 39*(4),

1563–1581. https://doi.org/10.1002/hbm.23933

Yoris, A. E., Cira, L. F., Luque-Casado, A., Salvotti, C., Tajadura-Jiménez, A., Avancini, C.,

Zarza-Rebollo, J. A., Sanabria, D., & Perakakis, P. (2024). Delving into the

relationship between regular physical exercise and cardiac interoception in two

cross-sectional studies. *Neuropsychologia, 198*, 108867.

https://doi.org/10.1016/j.neuropsychologia.2024.108867

Yoris, A., García, A. M., Traiber, L., Santamaría-García, H., Martorell, M., Alifano, F., Kichic,

R., Moser, J. S., Cetkovich, M., Manes, F., Ibáñez, A., & Sedeño, L. (2017). The

inner world of overactive monitoring: Neural markers of interoception in

obsessive–compulsive disorder. *Psychological Medicine, 47*(11), 1957–1970.

https://doi.org/10.1017/S0033291717000368

Yuan, H., Yan, H.-M., Xu, X.-G., Han, F., & Yan, Q. (2007). Effect of heartbeat perception on

heartbeat evoked potential waves. *Neuroscience Bulletin, 23*(6), 357–362.

https://doi.org/10.1007/s12264-007-0053-7

Zaccaro, A., Della Penna, F., Mussini, E., Parrotta, E., Perrucci, M. G., Costantini, M., &

Ferri, F. (2024). Attention to cardiac sensations enhances the heartbeat-evoked

potential during exhalation. *iScience, 27*(4), 109586.

https://doi.org/10.1016/j.isci.2024.109586

Zaccaro, A., Perrucci, M. G., Parrotta, E., Costantini, M., & Ferri, F. (2022). Brain-heart interactions are modulated across the respiratory cycle via interoceptive attention. *NeuroImage*, *262*, 119548. https://doi.org/10.1016/j.neuroimage.2022.119548

Zhang, Yihui, Zhang, J., Xie, M., Ding, N., Zhang, Yang, & Qin, P. (2023). Dual interaction between heartbeat-evoked responses and stimuli. *NeuroImage*, *266*(December 2022). https://doi.org/10.1016/j.neuroimage.2022.119817

Zhou, D., Stiso, J., Bertolero, M., Cornblath, E., Teich, E., Blevins, A. S., Murphy, K., Oudyk, K., Michael, C., Urai, A., Matelsky, J., Virtualmario, Camp, C., Castillo, R. A., Saxe, R., Dworkin, J., & Bassett, D. (2022). *dalejn/cleanBib: V1.1.3* [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.7375250

Zhou, H., Zou, H., Dai, Z., Zhao, S., Hua, L., Xia, Y., Han, Y., Yan, R., Tang, H., Huang, Y., Du, Y., Wang, X., Yao, Z., & Lu, Q. (2022). Interoception Dysfunction Contributes to the Negative Emotional Bias in Major Depressive Disorder. *Frontiers in Psychiatry*, *13*(April), 1–13. https://doi.org/10.3389/fpsyt.2022.874859

Zhou, M., Cheng, L., Zhou, Y., Zhu, S., Zhang, Y., Kendrick, K. M., & Yao, S. (2024). Intranasal Oxytocin Improves Interoceptive Accuracy and Heartbeat-Evoked Potentials During a Cardiac Interoceptive Task. *Biological Psychiatry: Cognitive Neuroscience and Neuroimaging*, S245190222400137X. https://doi.org/10.1016/j.bpsc.2024.05.004

Zwienenberg, L., van Dijk, H., Enriquez-Geppert, S., van der Vinne, N., Gevirtz, R., Gordon, E., Sack, A. T., & Arns, M. (2023). Heartbeat-Evoked Potential in Major Depressive Disorder: A Biomarker for Differential Treatment Prediction between Venlafaxine and rTMS? *Neuropsychobiology*, 1–10. https://doi.org/10.1159/000529308