The RNA-Binding Protein HuR Regulates Splicing Programs in Osteocytes

Ziqiu Fan[1],
Aseel Marahleh DDS, PhD[1,2]*,
Hideki Kitaura PhD[1],
Jiayi Ren PhD[3],
Abdulrahman Mousa MS[4,5],
Fumitoshi Ohori PhD[1],
Kuniyasu Niizuma MD, PhD [4,5],
Sherif Rashad MD, PhD[4,5],
Hiroyasu Kanetaka PhD[1]

*Corresponding author email: aseel.mahmoud.suleiman.marahleh.e6@tohoku.ac.jp
Corresponding author mailing address: 4-1 Seiryo machi, Aoba ku, Sendai 980-8575, Japan

1 Orthodontics and Dentofacial Orthopedics Department, Graduate School of Dentistry, Tohoku University, Japan.
2 Frontier Research Institute for Interdisciplinary Sciences, Tohoku University, Japan.
3 Center for Environmental Response and Aging, Institute of Development, Aging and Cancer, Tohoku University.
4 Department of Neurosurgical Engineering, Graduate School of Biomedical Engineering, Tohoku University
5 Department of Translational Neuroscience, Graduate School of Medicine, Tohoku University

Data availability statement
Raw sequencing data is deposited in Sequence Read Archive under the project ID PRJNA1262104

Funding statement
This work is supported by The FRIS Creative Interdisciplinary Collaboration Program to AM and JST SPRING to ZF (No. JPMJSP2114).

Conflict of interest disclosure
The authors report no conflict of interest.

Abstract

Post-transcriptional gene regulation plays a crucial role in maintaining cellular homeostasis. Alternative splicing (AS) is one such post-transcriptional gene regulatory mechanisms. In this study we used RNA binding protein (RBP) motif enrichment analysis of RNA-seq data to identify enriched RBP positions around alternatively spliced exons in primary osteocytes cultured in high glucose conditions (HG). We identified the RNA-binding protein human antigen R (HuR) as a top candidate regulator of AS in our datasets. HuR mRNA levels were reduced under hyperglycemic conditions in both MLO-Y4 cells and mice long bones, while protein expression remained unchanged. We investigated the mechanistic role of HuR in osteocyte adaptation to high glucose using HuR knockdown (KD) in MLO-Y4 cells. HuR KD led to profound differential gene expression and differential mRNA splicing changes. Functional splicing analysis revealed enrichment in pathways related to stress response, translation, and insulin signaling. Mechanistic studies confirmed AS analysis; HuR KD cells showed decreased mitochondrial mass, impaired respiratory function, reduced ATP production, and increased sensitivity to oxidative stress under challenging conditions. HuR KD cells exhibited reduced viability specifically under HG but not osmotic stress conditions. Additionally, HuR KD displayed dysregulation in key translational signals, including activation of mTOR kinase, decreased MNK1 activity and decreased in eIF4E phosphorylation while preserving global protein output. Moreover, HuR KD cells exhibited reduced insulin sensitivity as evident by reduced AKT phosphorylation in insulin stimulated cells. These findings establish HuR as a central post-transcriptional regulator of osteocyte survival and metabolic adaptation under high glucose stress, with potential implications for hyperglycemic bone fragility.

Keywords: HuR/*Elavl1*, RNA binding proteins, alternative splicing, hyperglycemia, osteocytes, high fat diet, translation, mTOR.

Introduction

Post-transcriptional gene regulation modulates gene expression at the RNA level through mechanisms such as RNA splicing, modifications, localization, translation and turnover.[1-6] Alternative splicing (AS) is a process where pre-mRNA transcripts undergo differential inclusion of exons, alternative inclusion of 3' or 5' splice sites and intron splicing, generating transcript variants that contribute to proteomic diversity, which plays a key role in stress adaptation optimizing protein synthesis to promote cell survival.[7] Remarkably, over 90% of multi-exon genes undergo AS with splicing patterns varying across cell types, tissues, developmental stages, and disease states.[8,9]

RNA binding proteins (RBPs) regulate AS by binding near splice sites or structured RNA motifs.[8,10-12] Beyond splicing, RBPs control nearly every aspect of RNA metabolism, including pre-mRNA processing, transport, stability, localization and translation.[10] RBPs contribute to tissue-specific disease phenotypes, often reflecting the expression pattern of their RNA targets and cofactors as well as the RBP's own abundance.[13] Alterations in RBPs or their targets are associated with diverse pathologies, including neurodegenerative, muscular, autoimmune, and metabolic diseases driven by impaired RNA transport, splicing, or clearance, highlighting the critical role of RNA-RBPs interactions in determining disease outcomes.[8,13,14]

HuR ['human antigen R', also known as ELAVL1 (embryonic lethal abnormal vision-like 1)] is a ubiquitously expressed RNA-binding protein involved in post transcriptional gene regulation.[15] It has been extensively studied as a cancer drug target[16] and its global deletion results in embryonic lethality highlighting its essential role in development.[17] Conditional knockouts affect post-natal tissue development including skeletal muscles, bones, osteogenesis and skeletal extracellular matrix (ECM) organization, and immune cell differentiation.[18-21] Beyond development, HuR influences metabolic homeostasis; tissue-specific knockdown disrupts glucose and lipid metabolism linking it to obesity and insulin resistance.[22-24]

In ovariectomy-induced osteoporosis, HuR expression is reduced, and its overexpression alleviated bone loss, whereas silencing impaired osteoblast differentiation.[25] In contrast, diabetic mouse models showed increased HuR mRNA levels, and its knockdown improved glucose metabolism and bone microarchitecture.[26] Similarly, in MC3T3-E1 cells exposed to high glucose, HuR expression increased, and silencing reduced apoptosis while promoting osteogenic differentiation.[26] These studies reveal varying changes in HuR expression and effects under metabolic stress different across models.

Osteocytes are the most abundant bone cells, critical for sensing mechanical stress, maintaining skeletal homeostasis through orchestrating bone resorption and bone formation as well as acting as an endocrine and signaling cell.[27-30] To date, little is known about AS events in osteocytes or the RNA-binding proteins that regulate them. Additionally, there have been no reports on the function of HuR in osteocytes. Therefore, this work provides a hypothesis-free splicing and RNA binding motif analysis of primary osteocytes obtained from Dmp1-topaz mice cultured in high glucose conditions. We measured HuR levels in MLO-Y4 cells and the long bones of hyperglycemic mice and we investigated the effect of HuR depletion on the transcriptome including focusing on its regulation of AS. Using an HuR KD MLO-Y4 cell line, we examined how HuR depletion influenced translation signaling pathways, mitochondrial function and cellular homeostasis under both normal and stress conditions. These findings will provide critical mechanistic insights into the role of HuR in maintaining cellular homeostasis and suggest broader implications in metabolic bone diseases.

Materials and Methods

1. Primary cells and cell lines

Primary osteocytes were isolated from neonatal calvariae of C57BL/6-Tg (Dmp1-Topaz)1Ikal/J using serial enzymatic digestion with collagenase and EDTA followed by fluorescent activated cell sorting (FACS) as described previously.[31] After isolation, cells were cultured at standard conditions (37 ⬜⬜, 5% CO_2) overnight in alpha-minimum essential medium (⬜-MEM; Wako Pure Chemical Industries, Osaka, Japan) containing 10% fetal bovine serum (FBS; Biowest, Nuaillé, France), 100IU/mL penicillin G, 100 µg/mL streptomycin (1% P/S) containing 5.5 mM glucose (standard medium). After that, the media was replaced with fresh 5.5mM glucose (normal glucose, NG), 25 mM glucose (Sigma-Aldrich, St. Louis, MO, USA) (high glucose, HG) containing media or 5.5 mM glucose supplemented with 19.5 mM mannitol (osmotic control, M) (Sigma-Aldrich) for 3 days. MLO-Y4 cells (AddexBio Technologies) were maintained at 37 ⬜⬜, 5% CO_2 in standard medium and kept at 70-80% confluency and passaged as needed. Standard medium was changed to NG, HG or M media for subsequent experiments.

2. Mice and feeding experiments

12-week-old male C57BL/6J mice were purchased from CLEA Japan (Tokyo, Japan). Mice were acclimated for 4 weeks in specific pathogen-free (SPF) conditions under a 12-h light/dark cycle with ad libitum feeding (Research Diet, D12450H, Japan) and at 16 weeks the mice were randomly allocated to a 10% Kcal from fat termed normal diet (ND) (Research Diet, D12450H, Japan) or 45% Kcal from fat termed high fat diet (HFD)

(Research Diet, D12451, Japan) for 16 weeks to induce chronic hyperglycemia. The mice were tested for fasting blood glucose and glucose tolerance at 4 and 8 weeks of diet and one day before collecting the bones at 16 weeks of diet with 5 mice per group. After sacrifice, the long bones (femur, tibia, humerus) were excised. The muscles and soft tissues were carefully removed, and the bones were scraped using a scalpel. The epiphyses were cut and the bone marrow flushed in phosphate-buffered saline (PBS) x1 at 9000 xg, 4 ☐☐ three times. The bones were immediately frozen in liquid nitrogen and stored at -80 ☐☐ for RNA and protein extraction. All procedures involving animals were performed in accordance with the ARRIVE guidelines and were approved by the Regulations for Animal Experiments and Related Activities at Tohoku University (2021DnLMO-007-04).

3.   Blood glucose measurement and glucose tolerance test

Fasting blood glucoseand glucose tolerance were assessed after a 6-hour fast (8:00 AM–2:00 PM) using a rodent specific glucometer with compatible LG sensors (LabGluco Funakoshi., Tokyo, Japan). Approximately, 5 µL or less of blood was drawn from a tail tip clip at 0, 15, 30, 60, and 90 minutes.

4.   Short hairpin RNA knockdown

Two short hairpin RNA (shRNA) sequences targeting mouse HuR/ELAVL1 mRNA were designed using VectorBuilder's shRNA Target Design tool. An empty backbone sequence was used as a non-targeting control. The shRNA oligonucleotides were ordered and synthesized by FASMAC Co., Ltd. (Kanagawa, Japan). Oligos were annealed and ligated and cloned into pLKO.1 puro vector (Addgene, Plasmid# 8453). Plasmids were then extracted using Nucleospin plasmid DNA purification kit (Macherey-Nagel, Düren, Germany) followed by validating the insertions using colony-PCR confirming sequences by Sanger Sequencing (Eurofins Genomics, Tokyo, Japan). HEK293T cells were co-transfected with shRNA plasmids, lentiviral packaging plasmid psPAX2 (Addgene, Plasmid#12260) and VSV-G envelope expressing plasmid pMD2.G (Addgene, Plasmid#12259) using Lipofectamine 3000 (Thermo Fisher Scientific, Waltham, MA, USA). Viral particles were collected after 48 and 72 hours, centrifuged at 1500 xg for 5 minutes and supernatant was filtered through a 0.45 µm PVDF filter. Viral particles were stocked at 4 ☐☐ for immediate use or -80 ☐☐ for long term storage. MLO-Y4 cells were grown to 80% confluency in standard medium, then media was changed to transduction medium containing standard medium and lentiviral particles containing supernatant at 4:1 (by volume) and 8 µg/mL polybrene (Sigma-Aldrich, St. Louis, MO, USA) for 24 hours. Transduction medium was then replaced with standard medium and incubated for 6 hours then puromycin was added at (2 µg/mL) and transfected cells were selected for 7 days before passaging. Construct sequences of control Mock

knockdown (Mock[KD]), construct A knockdown (YA[KD]) and construct B knockdown (YB[KD]) in Supplementary Table 5. A and B denote 2 separate constructs and (Y) denotes that MLO-Y4 cells were used for knockdown.

5.   RNA sequencing

Total RNA was extracted using Qiagen RNeasy kit (QIAGEN, Hilden, Germany) with a DNase digestion step (QIAGEN) as per manufacturer's instructions. The samples were processed by Azenta, Japan as follows: samples were screened and quantified on an Agilent 4200 Tapestation with RNA ScreenTape. The resultant RIN (RNA integrity number) was 9.9 or 10 for all samples. mRNA was enriched using NEBNext Poly(A) mRNA Magnetic Isolation Module (NEB) and RNA library preparation was done NEBNext Ultra II Directional RNA Library Prep Kit (Illumina E7760). Samples were pooled on an Illumina Novaseq to obtain paired end 150bp reads with a target depth of 50 million reads per sample. FASTQ files were then used for bioinformatics analysis. All sequencing experiments were done in triplicates.

6.   RNA sequencing analysis

Quality control of raw FASTQ files was conducted using FastQC. Adapters and low-quality reads were removed using Trimmomatic.[32] Trimmed reads were aligned to the mouse reference genome (M35, GRCm39) using the splice-aware aligner HISAT2.[33] The resulting BAM files of mapped reads were quantified against gene features using FeatureCounts with default parameters.[34] Differentially expressed genes (DEGs) were identified using Limma-Voom,[35] applying TMM normalization and defining DEGs as those with $|\log_2$ fold change$| \geq 1$ and adjusted p value < 0.05.

7.   AS and RBP motif analysis

Aligned BAM files were used to conduct AS analysis using the rMATs-turbo suite.[36] Statistical significance of AS events was considered at FDR < 0.05 and $|\Psi| > 0.2$ to filter in differentially spliced genes (DSGs). Splice motif analysis was conducted using rMAPs2.[37]

8.   Gene ontology and functional enrichment analysis

DSGs were used for functional enrichment analyses. The output of rMATs was piped into rMAPs to generate RBP motif maps.[37] DEGs and DSGs were analyzed for gene ontology (GO) and functional enrichment analysis using either David GO functional annotation tools or eVITTA's pre-ranked gene set enrichment analysis (GSEA) or overrepresentation analysis (ORA) toolbox.[38] Figures in this publication were made using bespoke scripts in Rstudio 4.4.0 ggplot or ggplot2 packages and extensions.

9.   Cell viability assay

In a 96-well plate, 1×10^4 Mock[KD], YA[KD] or YB[KD] were seeded in each well. Cells were cultured in each well with 200 µL of α-MEM supplemented with 10% FBS and 1% P/S. Cell viability was measured at 0, 24, 48, and 72 hours. The plate was then incubated for 2 hours at 37°C with each well containing 100 µL of α-MEM and 10 µL of Cell Counting Kit-8 solution (Dojindo Laboratories, Kumamoto, Japan). Absorbance was measured at 450 nm using a microplate reader (Sunrise™ Remote, Tecan, Männedorf, Switzerland).

10.   Reactive oxygen species detection assay

Mock[KD] or YB[KD] (2.5×10^4 cells per well) were seeded in a black, clear bottom 96-well microplate overnight. Intracellular reactive oxygen species (ROS) levels were measured using the DCFDA/H2DCFDA-Cellular ROS Assay Kit (Abcam, Cambridge, UK) according to manufacturer's instructions. Cells were stained with 20 µM DCFDA in HBSS for 45 minutes at 37°C. Mock[KD] or YB[KD] cells treated with 50µM TBHP for 4 hours were used as positive control. DCF fluorescence intensity was measured using a fluorescence plate reader (Flex Station 3, Molecular Devices, San Jose, CA, USA) with excitation at 485 nm and emission at 535 nm.

11.   Insulin response tests

Mock[KD] or YB[KD] cells were seeded in α-MEM supplemented with 10% FBS and 1% P/S until reaching 80% confluency. The media were then replaced with α-MEM containing 1% P/S and 1% FBS for 3 hours followed a switch to the same media with or without 100 nM insulin (Sigma-Aldrich). After a 30-minute incubation period, cells were harvested and protein extracted using radioimmunoprecipitation (RIPA) assay buffer (Millipore, Burlington, MA, USA) containing 1% protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific) on ice for 20 minutes; insoluble material was separated by centrifugation at 14000 xg. Total protein was collected for western blot analysis.

12.   Real-time polymerase chain reaction (qPCR)

RNeasy kit (QIAGEN) was used to extract total RNA from cells according to manufacturer's instructions and Trizol-chloroform was used to extract RNA from bones. 1 femur was immersed in 1 mL Trizol and pulverized 5 times at 5000 RPM speed setting for 60 seconds each in TOMY Micro Smash MS 100 Cell Disruptor. Total RNA was used to synthesize cDNA using the SuperScript® IV First-Strand Synthesis System (Invitrogen, Carlsbad, CA, USA) according to manufacturer's instructions. mRNA expression values were measured using the CFX96 Touch Real-Time PCR Detection System (Bio-Rad Laboratories, Hercules, CA, USA) using the ΔCq method. Target gene expression was

then measured as a function of housekeeping gene expression glyceraldehyde-3-phosphate dehydrogenase (GAPDH). Primers sequences are in Supplementary Table 5.

13. Western blot

Cells were lysed using RIPA assay buffer (Millipore, Burlington) containing 1% protease and phosphatase inhibitor cocktail (Thermo Fisher Scientific) on ice for 20 minutes; insoluble material was separated by centrifugation 14000 xg. Protein from bones was extracted in RIPA buffer and processed similar to RNA extraction above. Total protein was quantified using Pierce™ BCA protein assay kit (Thermo Fisher Scientific). To prepare for SDS-PAGE gel electrophoresis, the protein was treated with β-Mercaptoethanol (Bio-Rad Laboratories) and Laemmli sample buffer (BioRad Laboratories) 3:1 and denatured at 95°C for 5 minutes. Equal protein amounts were loaded onto 4–15% Mini-PROTEAN TGX Precast Gels (Bio-Rad Laboratories) and transferred to a Trans-Blot Turbo Transfer System (Bio-Rad Laboratories) and then blocked with Block-Ace (DS Pharma Biomedical, Osaka, Japan) 1 hour at room temperature (RT). Primary antibodies were incubated overnight at 4°C. The membranes were incubated with horseradish peroxidase-conjugated anti-rabbit antibody (Cell Signaling Technology, Danvers, MA, USA) at a dilution of 1:5,000 or anti-mouse antibody (GE Healthcare, Chicago, IL, USA) at a dilution of 1:10,000 for 1 hour at RT after being washed in tris-buffered saline with Triton X-100 (TBS-T). Bound antibodies were detected with SuperSignalWest Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific) and a FUSION-FX6 EDGE Chemiluminescence Imaging System (Vilber Lourmat, Collégien, France). Primary antibodies are in Supplementary Table 5.

14. Confocal microscopy

1×10^5 Mock^{KD}, YA^{KD} or YB^{KD} cells were seeded in 35mm glass bottom dish overnight. Cells were fixed with 4% formaldehyde for 15 minutes, then washed 3 times with x1 PBS. Cells were permeabilized using 0.1% Triton X-100 for 10 minutes and blocked with 1.5% bovine serum albumin (BSA) for 30 minutes at RT. Cells were incubated with Alexa Fluor 568 Phalloidin (Invitrogen) for 1 hour at RT. Afterwards, cell nuclei were stained with 4′,6-diamidino-2-phenylindole, dihydrochloride (DAPI) (ThermoFisher Scientific) for 5 minutes at RT. For mitochondrial staining, 5×10^4 live cells were incubated with the MitoTracker™ Red CMXRos FM (Invitrogen) in a glass bottom dish for 45 minutes. Cells were washed with x1 PBS 3 times and fixed with 4% formaldehyde for 15 minutes then washed by x1 PBS. Cells were incubated for 10 minutes in x1 PBS containing 0.1% Triton X-100 and stained with DAPI for 5 minutes at RT. Confocal images were acquired using a Zeiss LSM800 confocal laser scanning microscope (Carl Zeiss, Oberkochen, Germany). Confocal micrographs acquired with 10× and 20× objectives for phalloidin staining, and 40x for MitoTracker™ as a single optical slice.

15. Seahorse mitochondrial mito-stress test

Oxygen consumption rate (OCR) was measured using the Seahorse XF96 analyser (Agilent Technologies, Santa Clara, CA, USA) and Mito-stress test kit (Agilent Technologies) along with Seahorse XFe96/XF Pro FluxPak (Agilent Technologies) and Seahorse XF DMEM medium (without Phenol Red/pH 7.4/with HEPES/500 mL, Agilent Technologies). Initially, Mock[KD] and YB[KD] cells were seeded at 25,000 cells/well in 96-well plates on day one, while the XFe96 sensor cartridge hydrated in 200 μL calibration medium. On the next day, the cell medium was replaced with Seahorse XF DMEM, and cells were incubated at 37°C in a CO_2-free incubator for 1 hour. During this time, 1 μM oligomycin, 0.5 μM FCCP, antimycin A, and rotenone were loaded into the drug ports of the cartridge. After loading, the sensor plate was calibrated in the analyser. Following calibration, the cell culture plate was loaded, and the analysis initiated with the program: set: Mixture 2 minutes, Measure 3 minutes, 4 sets in all steps. Ports: (A: Oligomycin / B: FCCP / C: Antimycin/Rotenone).

16. Puromycin incorporation assay

Nascent protein synthesis was evaluated by puromycin incorporation assay. Cells were treated with 10 μg/mL puromycin (Sigma-Aldrich) for 30 minutes prior to harvest. Cell lysates were prepared and analysed by western blot as previously described.

17. Statistical analysis

All experiments were done in 3 or more replicates. Statistical analysis was performed using GraphPad Prism 9.0. Analysis of two group comparisons was done using student's t test and multiple group comparison was done using one-way ANOVA followed by Tukey's post hoc test. p values of less than 0.05 were considered significant.

Results

High glucose drives AS changes and modulates RBP motif enrichment in osteocytes

Primary osteocytes obtained from Dmp1-Topaz mice were cultured in HG or NG for 72 hours. RNA-seq analysis showed minimal gene expression changes (Figure 1A). To evaluate AS changes we analyzed RNA-seq data using rMATs for the 5 canonical splicing patterns; skipped exon (SE), mutually exclusive exons (MXE), retained introns (RI), alternative 5' splice site (A5SS), and alternative 3' splice site (A3SS).[36] Of the total AS events, 2% were significantly different between the two conditions with SE and A3SS exhibiting the highest change (Figure 1B). We used FDR < 0.05 and |Ψ| > 0.2 to extract DSGs. GO Biological Processes (GOBP) and pathway enrichment analysis of SE DSGs revealed enrichment in cytoskeletal remodeling and protein regulation via

ubiquitination, and several pathways involved in matrix maintenance and repair (Figure 1C, Supplementary Table 1). MXE DSGs were enriched in pathways involved in collagen synthesis and organization, skeletal tissue development and pathways involved in cell growth and regulation of fluid balance (Figure 1D, Supplementary Table 1). A5SS DSGs were predominated by pathways related to post transcriptional and translational regulation including ribosome targeting, rRNA processing and quality control of mRNA translation. A3SS DSGs were involved in RNA splicing, several signaling pathways involved in cell growth and differentiation and parathyroid hormone action. DSGs in RI showed cell cycle related terms, ribosome and translation regulation as well as ubiquitination (Supplementary Fig 1A-C and Supplementary Table 1).

To further understand cellular response to HG we looked at post-transcriptional RNA regulators. We used rMAPs2 to visualize RBP motif enrichment around differentially regulated exons from RNA-seq data.[37] We filtered the top up enriched and down enriched motif positions from all 5 AS datasets based on p values < 0.05 (t test). Around upregulated splice events, HNRNPL and HuR were the top enriched in SE (Figure 2A, Supplementary Table 2), in MXE, ZC3H14 and HuR. In A5SS, HNRNPL and HuR, in A3SS, FUS and HNRNPH2 and in RI we observed no enriched RBP motifs at a cut off value of p <0.05 (Supplementary Fig 2A-D and Supplementary Table 2). Around downregulated splice events, KHDRBS1 and ZC3H14 were the top enriched in SE (Figure 2B), in MXE, ZC3H14 and Tra2-beta, in A5SS, HNRPLL and HuR showed the highest enrichment. In A3SS, RBM47 and SRp20 were the most enriched while in RI RBMS3 and PABPN1 were dominant (Supplementary Fig 3A-D and Supplementary Table 2).

The analysis showed that several HuR motifs were enriched in upregulated splice events. Figure 2C shows an example of HuR binding motifs around SE differentially spliced exons. Our bioinformatics analysis points to HuR as a top regulatory RBP in AS, therefore we measured HuR mRNA and protein levels in MLO-Y4 cells exposed to HG. mRNA expression was downregulated as measured by qPCR while protein expression did not change as measured by western blot (Figure 3A-C). Similarly, HuR mRNA also decreased in the long bones of hyperglycemic mice (Supplementary Fig 4) that were fed a moderately HFD for 16 weeks as measured by qPCR while protein levels did not change compared to ND (Figure 3D- F).

HuR KD reveals transcriptomic and phenotypic changes in MLO-Y4 cells

We knocked down HuR in MLO-Y4 cells (Y[KD]) using two independent shRNA constructs, both of which effectively reduced HuR protein levels (Figure 4A). Despite comparable knockdown efficiency, we observed differences between the two constructs. Construct A (YA[KD]) induced severe cytotoxicity, including a significant loss of cell viability (Figure

4B) and dramatic morphological changes characterized by a severely reduced cytoplasmic volume and near complete loss of dendrites (Figure 4C, D). These phenotypic changes are consistent with apoptotic processes and suggest potential off-target effects. Construct B (YB^{KD}) experienced reduced cell viability and morphological changes though to a lesser extent than YA^{KD}, and preserved cell dendrites (Figure 4C, D). To ensure that our results were not confounded by potential off-target effects from construct A, we used YB^{KD} for further analysis, hereafter, the terms HuR KD and YB^{KD} are used interchangeably. RNA-seq analysis of YB^{KD} revealed 129 differentially expressed genes at |Log2FC|≥ 1 and adjusted p value < 0.05 compared to Mock^{KD} including HuR downregulation (Figure 4E). The top 10 enriched pathways in pre-ranked gene set enrichment analysis showed downregulation of extracellular matrix organization and remodeling -including several collagens, matrix metalloproteinases and cathepsins- growth and development pathways, fatty acid and cholesterol biosynthesis pathways, in addition to cell signaling pathways involved in inflammation and cellular homeostasis. Upregulated pathways showed a clear trend towards enrichment in pathways related to quality control of mRNA processing, translation and ribosomal terms. In addition to pathways related to immunity and pathways related to cytoskeleton organization (Figure 4F and Supplementary Table 3). Overall, the transcriptomic profile of HuR KD points to a complex role for HuR in controlling several aspects of cellular homeostasis including cytoskeletal organization, extracellular matrix proteins and a role in regulating mRNA processing and translation.

HuR regulates splicing programs in MLO-Y4 cells

We further compared alternative mRNA splicing patterns between YB^{KD} and Mock^{KD}. We summarized the output of the AS analysis from junction and exon body counts (JCEC), the total number of JCEC events was 24 thousand events, with 3% showing significance between YB^{KD} and Mock^{KD}. Among the AS patterns detected, SE represented the most frequently regulated events, while RI events were the least regulated (Figure 5A). This observation aligns with the earlier HuR motif analysis in primary osteocytes (Supplementary Fig 2 and 3), where RI events showed minimal to no enrichment in HuR motifs around differentially spliced introns. DSGs (Figure 5B, Supplementary Table 4) were used to perform gene ontology (GO) and over representation functional annotation (ORA) following HuR KD. The top 10 scoring cellular anatomical structures are shown in Figure 5C, including ribonucleoprotein complexes (RNPs), ribosomes and mitochondria. ORA analysis revealed a strong enrichment in RNA terms such as rRNA processing, RNA splicing, and mRNA surveillance pathways, including nonsense-mediated decay (NMD). Pathways related to the translation machinery, including ribosomal units, rRNAs, and translational

regulation, were also enriched, consistent with HuR's role as an RBP. Furthermore, pathways associated with autophagy and stress response such as the HIF-1α and metabolic terms such as insulin signaling pathways, were enriched (Figure 5D, Supplementary Table 4).

HuR KD disrupts translational signaling while preserving global protein output in MLO-Y4 cells

Our DSGs analysis revealed significant enrichment in pathways related to translational control, therefore, we investigated several key translational signals. We assessed the phosphorylated status of the mechanistic target of rapamycin (mTOR), a central mediator of mRNA translation and protein synthesis. mTOR phosphorylates two key immediate targets, the p70S6 kinase (p70S6K) and eukaryotic initiation factor 4E-binding protein 1 (4E-BP1).[39] Total levels of p70S6K and 4E-BP1 were reduced in HuR KD cells. However, their phosphorylated forms, p-p70S6K and p-4E-BP1, increased, resulting in increased phosphorylated-to-total protein ratios (Figure 6A, B). We next examined mitogen activated protein kinase-interacting serine/threonine kinase 1 (MAPK-MNK1) axis, which predominantly regulates cap dependent translation initiation, the predominant form of translation in eukaryotic cells.[40] Phosphorylated MNK1 decreased, whereas total MNK1 levels remained unchanged (Figure 6C). Consistently, upstream of MNK1, the extracellular signal-regulated kinases1/2 (ERK1/2) showed reduced levels of both phosphorylated and total protein (Figure 6D). Furthermore, these pathways converge on the eukaryotic initiation factor 4E (eIF4E) – a direct substrate of MNK1 and 4E-BP1. In HuR KD cells, both total and phosphorylated levels were reduced, leading to a lower p-eIF4E/eIF4E ratio (Figure 6E). To determine whether these molecular changes affected overall protein translation, we performed a puromycin incorporation assay to measure nascent protein synthesis rate. Despite the pronounced alterations in these translational regulators, no significant differences in global protein synthesis rates were detected suggesting maintenance of translational output through the compensatory activation of mTOR1 (Figure 6F).

HuR KD induces mitochondrial dysfunction and sensitizes cells to oxidative cell death in MLO-Y4 cells

DSG analysis pointed to mitochondrial terms enrichment, therefore we examined the effect of HuR KD on mitochondrial function and mass. To quantify mitochondrial density, we used Mitotracker CMXRos Red, a selective mitochondrial dye. Mitochondria localized around the nuclei and extended to dendrites, and we observed reduced mean fluorescence intensity in YB[KD] cells compared to Mock[KD] (Figure 7A, B). Next, we measured mitochondrial function using Seahorse XF Cell Mito-Stress test. YB[KD] cells exhibited reduced mitochondrial respiration (Figure 7C, D). Basal OCR, maximal

respiration, spare respiratory capacity, coupling efficiency and ATP production were reduced. Proton leak remained unchanged, suggesting that the mitochondrial membrane is intact. The observed reduction in OCR also extended to non-mitochondrial respiration, suggesting that the decline in oxygen consumption is partly attributable to reduced activity of non-mitochondrial oxygen-consuming enzymatic reactions (Figure 7D). Additionally, given that mitochondrial respiration is a major source of ROS, and reduced mitochondrial mass and respiration were observed in HuR KD cells, we next examined whether these changes affected redox homeostasis and cellular susceptibility to oxidative stress. We performed DCFDA assay which measures ROS levels, the results indicated a decrease in ROS production under baseline culture conditions in YB^{KD}, however, upon treating cells with TBHP, a ROS generating compound, YB^{KD} accumulated ROS double that of Mock^{KD} indicating that the baseline reduction in ROS in YB^{KD} cells may be due to a decrease in oxygen consumption by mitochondrial and non-mitochondrial activities (Figure 7E). This reduction in oxygen consumption likely limits the generation of ROS under normal conditions, as mitochondria are a primary source of intracellular ROS and the doubling of ROS upon treatment with TBHP may indicate a compromised antioxidant defense system. Since our initial analysis pointed to HuR as a top regulator of RNA splicing in HG conditions and HG is known to induce oxidative stress, we next assed the effect of HG on YB^{KD} cells. We cultured cells with NG or HG and M as osmotic pressure control, we used H$_2$O$_2$ as a positive control. HG induced cell death only in YB^{KD} cells but not in Mock^{KD} after 48 and 72 hours indicating that HuR KD sensitizes cells to oxidative damage. Hydrogen peroxide induced oxidative cell death in both cell lines; however, the extent of cell death was observed to be more in YB^{KD} cells compared to Mock^{KD} (Figure 7F). Those result indicate that HuR KD induces mitochondrial dysfunction and sensitizes cells to oxidative stress and oxidative cell death suggesting reduced cellular capacity to adapt to oxidative challenge.

HuR regulates insulin sensitivity in MLO-Y4 cells

Insulin conveys an anabolic signal; hence, it is tightly coupled to protein translation and mitochondrial activity, and our DSGs analysis revealed enrichment in insulin signaling pathway (Figure 5D). Therefore, we assessed whether HuR KD alters insulin responsiveness. HuR KD exhibited a reduced response to insulin stimulation, as evidenced by decreased AKT phosphorylation (p-AKT) in Figure 8A. qPCR analysis showed a modest but statistically significant increase in insulin receptor expression in YB^{KD} cells suggesting transcriptional compensatory feedback (Figure 8B).

Discussion

AS is a process by which a gene can produce multiple mRNA transcripts, contributing to the diversity and complexity of the protein pool. AS modulates gene function under both physiological and pathological conditions; consequently, its dysregulation has been linked to various diseases, including developmental, neurological, metabolic disorders, and cancer.[8] RBPs modulate RNA metabolism including but not limited to AS.[8] HuR, is a ubiquitous RBP member of Hu/ELAV RBP family.[15] HuR has emerged as a key modulator of metabolic disease, but its role appears to vary across tissues and disease models. In adipose tissue, HuR levels are decreased in diabetic and HFD-fed mice, and adipose-specific HuR knockout predisposes mice to HFD-induced obesity, insulin resistance, and glucose intolerance.[22] In the liver, HuR expression is similarly reduced in obesity-associated steatosis, yet its knockout aggravated hepatic steatosis in some models[41] and in others worsened insulin resistance and glucose intolerance.[23] Conversely, increased HuR expression has been observed in diabetic human heart and rat retina, suggesting context-dependent upregulation.[42,43]

These context-dependent discrepancies also manifest in the skeletal system. In ovariectomy-induced osteoporosis, HuR levels were decreased, and its overexpression mitigated bone loss, while silencing impaired osteoblastic differentiation in MC3T3-E1 cells.[25] In contrast, a diabetic mouse model exhibited elevated HuR mRNA levels, and knockdown improved glucose homeostasis and bone structure.[26] Similarly, high glucose exposure increased HuR expression in MC3T3-E1 cells, where silencing attenuated apoptosis and enhanced osteogenic differentiation.[26] Collectively, these reports underscore that HuR responds to metabolic perturbations in a tissue- and condition-specific manner, possibly due to the effect being enacted by the target RNA affected rather than HuR abundance itself.

Our data shows that in osteocytes exposed to high glucose, mRNA splicing regulated a wide array of cellular programs including ECM proteins, translation, and cytoskeletal organization. To explore RBPs contributing to these splicing alterations, we performed motif enrichment analysis, which identified several RBPs including HuR, ZC3H14, and HNRNPL as recurrently enriched around differentially spliced exons. HuR emerged as a prominent candidate, appearing in multiple positions across various splicing events. Therefore, we examined HuR expression at both the mRNA and protein levels in osteocytic MLO-Y4 cells cultured in HG and in the bones of HFD mice. Despite a significant reduction in HuR mRNA, protein levels remained unchanged, suggesting post-transcriptional buffering. Interestingly, mannitol-treated cells used to control osmotic stress exhibited a similar pattern, with decreased HuR mRNA but stable protein levels, further implicating regulatory mechanisms that maintain protein abundance under stress. Previous studies have shown that HuR protein levels

remained stable under hypotonic stress, while its subcellular localization shifted from the nucleus to the cytoplasm.[44] Although we did not directly assess HuR localization in our study, it is conceivable that similar stress-induced relocalization occurred in our cells, potentially contributing to the observed effect on mRNA and protein levels. Our observation is also consistent with evidence that mRNA abundance does not reliably predict protein levels, due to translational control and buffering mechanisms.[45] Given HuR's central role in maintaining cellular homeostasis, its expression may be preserved through enhanced translational efficiency or altered localization to sustain functional output. However, this hypothesis remains speculative and warrants further investigation using approaches such as ribosome profiling or spatial proteomics.

The dynamics of HuR expression under hyperglycemic stress prompted us to investigate its role in osteocytes. HuR KD in MLO-Y4 cells exhibited profound differential gene expression, including reduced expression of genes involved in extracellular matrix remodeling consistent with its reported role in maintaining several ECM proteins including collagens and matrix metalloproteinases.[46] DEGs also showed downregulation in lipid metabolism and cytoskeletal organization reflecting cellular morphological changes. Consistent with that phalloidin staining showed rounding and loss of dendrites after HuR KD consistent with GSEA. Enriched pathways were clearly biased towards mRNA quality control and mRNA translation.

DSGs in HuR KD cells showed enrichment in ribosomal and mitochondrial terms as well as terms related to the translational machinery, such as rRNA processing, RNA splicing, and translation- dependent mRNA surveillance pathways, including NMD, indicating an increased aberrant mRNAs and mRNA degradation.[47] These findings highlight that HuR KD induces substantial splicing alterations that correlate well with transcriptional alterations. Furthermore, AS analysis revealed significant enrichment in stress-related pathways including autophagy and HIF-1α signaling as well as key metabolic pathways such as insulin signaling and non-alcoholic fatty liver disease. Notably, these pathways were not detected through differential gene expression analysis, highlighting AS as a critical post-transcriptional mechanism of gene regulation in MLO-Y4 cells. These findings emphasize the added resolution provided by splicing analyses, enabling the identification of regulatory changes that are not apparent at gene abundance levels alone.

Given the enrichment of AS events in pathways related to mRNA translation and protein synthesis, we examined whether these changes were functionally reflected in translational signaling. Eukaryotic translation can be initiated via two mechanisms, cap-dependent and cap independent translation. Cap dependent translation is the predominant form and cap dependent translation initiation is the rate limiting step in

translation which depends on eIF4E availability.[48] eIF4E binds to the m^7G cap located at the 5' end of most mRNAs where it recruits other initiation factors preparing the mRNA for ribosomal loading and scanning for the translation initiation codon.[48] The activity of eIF4E is regulated via two mechanisms. First, it is sequestered by the inhibitory 4E-BP1 which releases eIF4E upon its phosphorylation by mTORC1 kinase.[39] Second, available eIF4E is phosphorylated by MNKs which serves to modulate selective mRNA translation. Notably, phosphorylation of eIF4E by MNKs is not required for global mRNA translation, conversely, mTORC1 inhibition leads to near complete inhibition of mRNA translation.[49,50] This aligns well with our data where HuR KD preserved global protein output possibly due to enhanced mTORC1 activity and inhibition of 4E-BP1 which serves to release eIF4E for translation initiation.

Our data revealed reduced total levels of p70S6K and 4E-BP1, yet their phosphorylated forms were elevated, resulting in increased phospho-to-total ratios—suggesting relative activation of mTORC1 kinase. In contrast, the MNK1-eIF4E axis was suppressed: phosphorylated MNK1 and its upstream kinase ERK1/2 were reduced, and both total and phosphorylated eIF4E levels were diminished. Despite these alterations, global protein synthesis rates remained unchanged. The preservation of global protein output despite disrupted signaling in key translational pathways suggests a shift toward biased mRNA translation, where the cell prioritizes the synthesis of specific mRNAs. Such reprogramming may represent an adaptive response to preserve proteostasis under energetic and metabolic stress. Although we did not measure selective mRNA translation it is plausible that a shift towards increased translation of mRNAs related to cell survival has taken place in HuR KD cells.

HuR KD increased the sensitivity of osteocytic cells to oxidative stress. Cells exposed to HG for 72 hours exhibited reduced viability, an effect not observed under mannitol treatment, indicating that the stress response is glucose-specific rather than due to osmotic pressure. Moreover, HuR KD cells displayed elevated ROS levels following exposure to oxidative agents (TBHP, H_2O_2) supporting a protective role for HuR in maintaining redox balance. Interestingly, under baseline conditions, HuR KD led to a reduction in ROS levels. Given that mitochondria are the primary source of cellular ROS, we assessed mitochondrial mass and function. HuR KD reduced mitochondrial content, as estimated by Mitotracker staining, and impaired mitochondrial respiration, evidenced by decreased OCR, maximal respiration, and coupling efficiency. ATP production was also diminished, indicating reduced oxidative phosphorylation (OxPhos), while proton leak remained unchanged, suggesting that mitochondrial membrane integrity was preserved. The decrease in ROS under baseline conditions may be explained by reduced mitochondrial activity and total oxygen consumption by

cells reducing ROS generating capacity. However, the heightened vulnerability to exogenous oxidative stress under HG and ROS-inducing conditions highlights a failure of adaptive antioxidant response in HuR KD cells. Collectively, these findings underscore HuR's role in maintaining cell viability under HG conditions and maintaining redox homeostasis and mitochondrial function. Concomitant with these results, HuR KD altered insulin signaling, as evidenced by reduced AKT phosphorylation in response to insulin stimulation. These observations suggest that HuR plays a key role in maintaining an anabolic environment that promotes cell survival and that HuR has a protective role in preventing decreased insulin sensitivity which is a key feature of diabetes affected bones. The enrichment of insulin signaling-related genes in the differential splicing analysis further supports this notion.

In conclusion, our study elucidates the role of HuR in osteocyte biology, particularly in regulating AS, translation, mitochondrial function and redox homeostasis. HuR seems to play a protective role against high glucose and oxidative cell death. Our findings reveal that HuR is indispensable for maintaining cellular and metabolic homeostasis in osteocytes, particularly under HG conditions. Our data provides a mechanistic understanding of HuR's role in cellular homeostasis and has broader implications in metabolic bone diseases.

References

1.   Moore MJ, Proudfoot NJ. Pre-mRNA Processing Reaches Back toTranscription and Ahead to Translation. *Cell*. 2009;136(4):688-700. doi:10.1016/j.cell.2009.02.001
2.   Maniatis T, Tasic B. Alternative pre-mRNA splicing and proteome expansion in metazoans. *Nature*. 2002/07/01 2002;418(6894):236-243. doi:10.1038/418236a
3.   Rodriguez AJ, Czaplinski K, Condeelis JS, Singer RH. Mechanisms and cellular roles of local protein synthesis in mammalian cells. *Current Opinion in Cell Biology*. 2008/04/01/ 2008;20(2):144-149. doi:https://doi.org/10.1016/j.ceb.2008.02.004
4.   Qiu L, Jing Q, Li Y, Han J. RNA modification: mechanisms and therapeutic targets. *Mol Biomed*. Aug 24 2023;4(1):25. doi:10.1186/s43556-023-00139-x
5.   Sonenberg N, Hinnebusch AG. Regulation of Translation Initiation in Eukaryotes: Mechanisms and Biological Targets. *Cell*. 2009;136(4):731-745. doi:10.1016/j.cell.2009.01.042
6.   Houseley J, Tollervey D. The Many Pathways of RNA Degradation. *Cell*. 2009;136(4):763-776. doi:10.1016/j.cell.2009.01.019

7.	Nilsen TW, Graveley BR. Expansion of the eukaryotic proteome by alternative splicing. *Nature*. Jan 28 2010;463(7280):457-63. doi:10.1038/nature08909

8.	Tao Y, Zhang Q, Wang H, Yang X, Mu H. Alternative splicing and related RNA binding proteins in human health and disease. *Signal Transduction and Targeted Therapy*. 2024/02/02 2024;9(1):26. doi:10.1038/s41392-024-01734-2

9.	Wang ET, Sandberg R, Luo S, et al. Alternative isoform regulation in human tissue transcriptomes. *Nature*. 2008/11/01 2008;456(7221):470-476. doi:10.1038/nature07509

10.	Dreyfuss G, Kim VN, Kataoka N. Messenger-RNA-binding proteins and the messages they carry. *Nature Reviews Molecular Cell Biology*. 2002/03/01 2002;3(3):195-205. doi:10.1038/nrm760

11.	Hentze MW, Castello A, Schwarzl T, Preiss T. A brave new world of RNA-binding proteins. *Nature Reviews Molecular Cell Biology*. 2018/05/01 2018;19(5):327-341. doi:10.1038/nrm.2017.130

12.	Cereda M, Pozzoli U, Rot G, et al. RNAmotifs: prediction of multivalent RNA motifs that control alternative splicing. *Genome Biology*. 2014/01/31 2014;15(1):R20. doi:10.1186/gb-2014-15-1-r20

13.	Gerstberger S, Hafner M, Tuschl T. A census of human RNA-binding proteins. *Nature Reviews Genetics*. 2014/12/01 2014;15(12):829-845. doi:10.1038/nrg3813

14.	Cooper TA, Wan L, Dreyfuss G. RNA and Disease. *Cell*. 2009/02/20/ 2009;136(4):777-793. doi:https://doi.org/10.1016/j.cell.2009.02.011

15.	Ma W-J, Cheng S, Campbell C, Wright A, Furneaux H. Cloning and Characterization of HuR, a Ubiquitously Expressed Elav-like Protein (*). *Journal of Biological Chemistry*. 1996;271(14):8144-8151. doi:10.1074/jbc.271.14.8144

16.	Majumder M, Chakraborty P, Mohan S, Mehrotra S, Palanisamy V. HuR as a molecular target for cancer therapeutics and immune-related disorders. *Adv Drug Deliv Rev*. Sep 2022;188:114442. doi:10.1016/j.addr.2022.114442

17.	Katsanou V, Milatos S, Yiakouvaki A, et al. The RNA-binding protein Elavl1/HuR is essential for placental branching morphogenesis and embryonic development. *Mol Cell Biol*. May 2009;29(10):2762-76. doi:10.1128/mcb.01393-08

18.	Ghosh M, Aguila HL, Michaud J, et al. Essential role of the RNA-binding protein HuR in progenitor cell survival in mice. *The Journal of Clinical Investigation*. 12/01/ 2009;119(12):3530-3543. doi:10.1172/JCI38263

19.	Figueroa A, Cuadrado A, Fan J, et al. Role of HuR in skeletal myogenesis through coordinate regulation of muscle differentiation genes. *Mol Cell Biol*. Jul 2003;23(14):4991-5004. doi:10.1128/mcb.23.14.4991-5004.2003

20.	Kota SK, Lim ZW, Kota SB. Elavl1 Impacts Osteogenic Differentiation and mRNA Levels of Genes Involved in ECM Organization. *Front Cell Dev Biol*. 2021;9:606971. doi:10.3389/fcell.2021.606971

21.	Osma-Garcia IC, Capitan-Sobrino D, Mouysset M, et al. The RNA-binding protein HuR is required for maintenance of the germinal centre response. *Nature Communications*. 2021/11/12 2021;12(1):6556. doi:10.1038/s41467-021-26908-2

22.   Li J, Gong L, Liu S, et al. Adipose HuR protects against diet-induced obesity and insulin resistance. *Nature Communications*. 2019/05/30 2019;10(1):2375. doi:10.1038/s41467-019-10348-0

23.   Zhang Z, Zong C, Jiang M, et al. Hepatic HuR modulates lipid homeostasis in response to high-fat diet. *Nature Communications*. 2020/06/16 2020;11(1):3067. doi:10.1038/s41467-020-16918-x

24.   Mynatt RL, Noland RC, Elks CM, et al. The RNA binding protein HuR influences skeletal muscle metabolic flexibility in rodents and humans. *Metabolism*. 2019/08/01/ 2019;97:40-49. doi:https://doi.org/10.1016/j.metabol.2019.05.010

25.   Liu Z, Li B, Hu H, Li X, Zhang X. Potential of RNA-binding protein human antigen R as a driver of osteogenic differentiation in osteoporosis. *J Orthop Surg Res*. Apr 12 2022;17(1):234. doi:10.1186/s13018-022-03073-w

26.   Ren Y, Yang M, Wang X, Xu B, Xu Z, Su B. ELAV-like RNA binding protein 1 regulates osteogenesis in diabetic osteoporosis: Involvement of divalent metal transporter 1. *Molecular and Cellular Endocrinology*. 2022/04/15/ 2022;546:111559. doi:https://doi.org/10.1016/j.mce.2022.111559

27.   Bonewald LF. The amazing osteocyte. *J Bone Miner Res*. Feb 2011;26(2):229-38. doi:10.1002/jbmr.320

28.   Marahleh A, Kitaura H, Ohori F, Noguchi T, Mizoguchi I. The osteocyte and its osteoclastogenic potential. *Front Endocrinol (Lausanne)*. 2023;14:1121727. doi:10.3389/fendo.2023.1121727

29.   Delgado-Calle J, Bellido T. The osteocyte as a signaling cell. *Physiol Rev*. Jan 1 2022;102(1):379-410. doi:10.1152/physrev.00043.2020

30.   Dallas SL, Prideaux M, Bonewald LF. The osteocyte: an endocrine cell ... and more. *Endocr Rev*. Oct 2013;34(5):658-90. doi:10.1210/er.2012-1026

31.   Au - Marahleh A, Au - Kitaura H, Au - Ogawa S, et al. *JoVE*. 2020/06/02/ 2020;(160):e61513. doi:doi:10.3791/61513

32.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics*. Aug 1 2014;30(15):2114-20. doi:10.1093/bioinformatics/btu170

33.   Kim D, Langmead B, Salzberg SL. HISAT: a fast spliced aligner with low memory requirements. *Nat Methods*. Apr 2015;12(4):357-60. doi:10.1038/nmeth.3317

34.   Liao Y, Smyth GK, Shi W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics*. Apr 1 2014;30(7):923-30. doi:10.1093/bioinformatics/btt656

35.   Law CW, Chen Y, Shi W, Smyth GK. voom: Precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*. Feb 3 2014;15(2):R29. doi:10.1186/gb-2014-15-2-r29

36.   Shen S, Park JW, Lu ZX, et al. rMATS: robust and flexible detection of differential alternative splicing from replicate RNA-Seq data. *Proc Natl Acad Sci U S A*. Dec 23 2014;111(51):E5593-601. doi:10.1073/pnas.1419161111

37.   Hwang JY, Jung S, Kook TL, Rouchka EC, Bok J, Park JW. rMAPS2: an update of the RNA map analysis and plotting server for alternative splicing regulation. *Nucleic Acids Res*. Jul 2 2020;48(W1):W300-w306. doi:10.1093/nar/gkaa237

38.   Cheng X, Yan J, Liu Y, Wang J, Taubert S. eVITTA: a web-based visualization and inference toolbox for transcriptome analysis. *Nucleic Acids Res*. Jul 2 2021;49(W1):W207-w215. doi:10.1093/nar/gkab366

39.   Saxton RA, Sabatini DM. mTOR Signaling in Growth, Metabolism, and Disease. *Cell*. 2017;168(6):960-976. doi:10.1016/j.cell.2017.02.004

40.   Yang X, Zhong W, Cao R. Phosphorylation of the mRNA cap-binding protein eIF4E and cancer. *Cell Signal*. Sep 2020;73:109689. doi:10.1016/j.cellsig.2020.109689

41.   Tian M, Wang J, Liu S, et al. Hepatic HuR protects against the pathogenesis of non-alcoholic fatty liver disease by targeting PTEN. *Cell Death & Disease*. 2021/03/04 2021;12(3):236. doi:10.1038/s41419-021-03514-0

42.   Jeyabal P, Thandavarayan RA, Joladarashi D, et al. MicroRNA-9 inhibits hyperglycemia-induced pyroptosis in human ventricular cardiomyocytes by targeting ELAVL1. *Biochemical and Biophysical Research Communications*. 2016/03/18/ 2016;471(4):423-429. doi:https://doi.org/10.1016/j.bbrc.2016.02.065

43.   Amadio M, Bucolo C, Leggio GM, Drago F, Govoni S, Pascale A. The PKCβ/HuR/VEGF pathway in diabetic retinopathy. *Biochemical Pharmacology*. 2010/10/15/ 2010;80(8):1230-1237. doi:https://doi.org/10.1016/j.bcp.2010.06.033

44.   Lema I, Amazit L, Lamribet K, et al. HuR-Dependent Editing of a New Mineralocorticoid Receptor Splice Variant Reveals an Osmoregulatory Loop for Sodium Homeostasis. *Scientific Reports*. 2017/07/06 2017;7(1):4835. doi:10.1038/s41598-017-04838-8

45.   Wang Z-Y, Leushkin E, Liechti A, et al. Transcriptome and translatome co-evolution in mammals. *Nature*. 2020/12/01 2020;588(7839):642-647. doi:10.1038/s41586-020-2899-z

46.   Pan H, Strickland A, Madhu V, et al. RNA binding protein HuR regulates extracellular matrix gene expression and pH homeostasis independent of controlling HIF-1α signaling in nucleus pulposus cells. *Matrix Biol*. Apr 2019;77:23-40. doi:10.1016/j.matbio.2018.08.003

47.   Nickless A, Bailis JM, You Z. Control of gene expression through the nonsense-mediated RNA decay pathway. *Cell & Bioscience*. 2017/05/19 2017;7(1):26. doi:10.1186/s13578-017-0153-7

48.   Merrick WC. Cap-dependent and cap-independent translation in eukaryotic systems. *Gene*. 2004/05/12/ 2004;332:1-11. doi:https://doi.org/10.1016/j.gene.2004.02.051

49.   Ueda T, Watanabe-Fukunaga R, Fukuyama H, Nagata S, Fukunaga R. Mnk2 and Mnk1 are essential for constitutive and inducible phosphorylation of eukaryotic initiation factor 4E but not for cell growth or development. *Mol Cell Biol*. Aug 2004;24(15):6539-49. doi:10.1128/mcb.24.15.6539-6549.2004

50.   Thoreen CC, Chantranupong L, Keys HR, Wang T, Gray NS, Sabatini DM. A unifying model for mTORC1-mediated regulation of mRNA translation. *Nature*. May 2 2012;485(7396):109-13. doi:10.1038/nature11083

Figure legends

Figure 1: High glucose induces minimal differential gene expression and extensive AS changes in primary osteocytes.

(A) The Volcano plot comparing the log2 FC and -log10 adjusted P value of RNA-seq data in primary osteocytes with genes showing the highest changes labelled.

(B) The parent pie chart shows significant AS events (2%) of the total events at target and junction reads (JCEC) and the secondary pie chart shows the percentage of significant AS events of each type (SE, A5SS, A3SS, MXE, RI) from total significant events in primary osteocytes.

(C-G) Volcano plots and overrepresentation analysis of DSGs around AS events showing inclusion level (Δ⍰) difference comparing 5.5mM and 25mM glucose at FDR < 0.05 and 0.2 < Δ⍰ < -0.2 in primary osteocytes. Overrepresentation analysis was done through the eVITTA web-based toolbox.

All analysis was done on n=3.
Also, see Supplementary Table 6.

Figure 2: RBPs motifs significantly enriched around AS events.

(A-B) The x axis (Attribute) represents the position (R) of each splicing event in 5' → 3'. RBP binding motifs are described in the y axis. The panels show SE RBP motifs around up and down regulated DSGs and bubble size and color denote the − log10 P value.

(C) Example positional distribution map of the HuR binding motif around target SE exon. Positional distribution maps were generated using rMAPs2. Intronic regions are 250-bp flanking sequences, and exon regions are 50-bp sequences from the start or end of the exon where the RBP binds around the spliced exon (green) and constitutive exons (grey). The solid red and blue lines are the motif enrichment scores, and the dotted lines are the p values in up and down enriched motifs, respectively. Black line refers to background RBPs distribution.

Also, see Supplementary Table 7.

Figure 3: HuR mRNA is reduced in MLO-Y4 cells cultured in HG and in the long bones of HFD fed mice, while maintaining HuR protein levels.

(A) HuR mRNA level in MLO-Y4 cells under normal glucose (NG), high glucose (HG) and osmotic control mannitol (M). Data are presented as mean ± SD, n=4. *p<0.05, ****p<0.0001.

(B) HuR protein level in MLO-Y4 cells in normal glucose (NG), high glucose (HG) and osmotic control mannitol (M). Data are presented as mean ± SD, n=7.

(C) HuR mRNA level in the long bones of high fat diet (HFD) and normal diet (ND) fed mice. Data are presented as mean ± SD, n=4. *p<0.05.

(D) HuR protein level in the long bones of high fat (HF) and normal diet (ND) fed mice. Data are presented as mean ± SD, n=4. *p<0.05.

Figure 4: HuR KD induces transcriptional changes in MLO-Y4 cells.

(A) Western blot analysis of HuR protein levels after shRNA-mediated knockdown in MLO-Y4 cells. Data are presented as mean ± SD, n=3. **p<0.01.

(B) CCK cell viability assay. Data are presented as mean ± SD, n=4. **p<0.01.

(C) Confocal microscopic images of phalloidin staining, Scale bars: 200 μm for the 10× image and 100 μm for the 20× image.

(D) Cytoplasm to nucleus ratio. The mean of 5 analysed regions (corners and center) of a 35 mm glass bottom dish were used. Data are presented as mean ± SD, n=4. ****p<0.0001.

(E) Volcano plot comparing the log2 FC and -log10 adjusted P value of RNA-seq data (DEGs) after YB^{KD} in MLO-Y4 cells, red arrow =HuR. n=3.

(F) Gene set enrichment analysis using eVITTA toolbox after YB^{KD} of DEGs at LogFC 1 and -log10 adjusted p<0.05.

Figure 5: HuR KD induces AS changes in MLO-Y4 cells

(A) The parent pie chart shows significant AS events (3%) of the total events at target and junction reads (JCEC) and secondary pie chart showing the percentage of significant AS events of each type (SE, A5SS, A3SS, MXE, RI) from total significant events.

(B) Scatter plot showing DSGs around AS events in 5 patterns FDR < 0.05 and 0.2 < Δ⍰ < -0.2.

(C) Gene Ontology Cellular Component (GOCC) enrichment analysis of DSGs performed using the DAVID Functional Annotation Tool.

(D) ORA of DSGs after YB^{KD}.

Also, see Supplementary Table 8.

Figure 6: HuR knockdown induces translation dysregulation and preserves global protein synthesis rate

(A)	Western blot of p-p70S6K and p70S6K protein levels. n=4.

(B)	Western blot of p-4E-BP1 and 4E-BP1 protein levels. n=4.

(C)	Western blot of p-MNK1 and MNK1 protein levels. n=7.

(D)	Western blot of p-ERK1/2 and ERK1/2 protein levels. n=4.

(E)	Western blot of p-eIF4E and eIF4E protein levels. n=4.

(F)	Western blot of puromycin incorporation assay. n=4.

Data are presented as mean ± SD of 2 or more independent blots (*p<0.05, **p<0.01 , ****p<0.0001)

Figure 7: HuR knockdown induces mitochondrial dysfunction

(A)	Fluorescent microscopic images of Mitotracker CMXRos Red staining mitochondria.

(B)	Mean fluorescent intensity (MFI) of Mitotracker CMXRos Red staining mitochondria. Data are presented as mean ± SD, n=4. ***p<0.001.

(C)	Seahorse XF Cell Mito-Stress showing oxygen consumption rate (OCR) in pmol/min.

(D)	Parameters of mitochondrial respiration. Data are presented as mean ± SD, n=4. *p<0.05, **p<0.01)

(E)	Reactive oxygen species (ROS) measurement. Fold change is calculated after treating cells with TBHP to induce ROS formation. Fold change represents TBHP stimulation/ basal line culture conditions. Data are presented as mean ± SD (n=5 per group; *p<0.05, ***p<0.001)

(F)	Cell viability assay in YB^{KD} and Mock^{KD} cells in 5.5mM glucose (NG), 25 mM glucose (HG), mannitol (M) and H_2O_2 stimulated cells. Data are presented as mean ± SD, n=4. *p<0.05, **p<0.01,***p<0.001, ****p<0.0001.

Figure 8: HuR knockdown induces decreased insulin sensitivity.

(A)	AKT/ actin and p-AKT/AKT protein levels in YB^{KD} and Mock^{KD} cells. Data are presented as mean ± SD, n=4. *p<0.05.

(B)   Insulin receptor mRNA expression level in YB^{KD} and Mock^{KD} cells. Data are presented as mean ± SD, n=3. *p<0.05.

Supplementary Fig 1: High glucose induces extensive AS changes in primary osteocytes.

(A-C) Volcano plots and overrepresentation analysis of DSGs around AS events showing inclusion level (Δᵈ) difference comparing 5.5mM and 25mM glucose at FDR < 0.05 and 0.2 < Δᵈ < -0.2 in primary osteocytes. Overrepresentation analysis was done through the eVITTA web-based toolbox. All analyses were done on n=3.

Supplementary Fig 2: Significantly up-enriched RBPs motifs around AS events.

(A-D) The x axis (Attribute) represents the position (R) of each splicing event in 5' → 3'. RBP binding motifs are described in the y axis. The panels show MXE, A5SS, A3SS and RI RBP motifs around up and down regulated DSGs and bubble size and color denote the − log10 P value.

Supplementary Fig 3: Significantly down-enriched RBPs motifs around AS events.

(A-D) The x axis (Attribute) represents the position (R) of each splicing event in 5' → 3'. RBP binding motifs are described in the y axis. The panels show MXE, A5SS, A3SS and RI RBP motifs around up and down regulated DSGs and bubble size and color denote the − log10 P value.

Supplementary Fig 4: Weight and metabolic tests of mice fed a high fat diet for 16 weeks.

(A)   Six-hour fasted blood glucose levels of mice fed a ND or HFD. Data are presented as mean ± SD, n=5.

(B)   Glucose tolerance test area under the curve (AUC) calculated from individual glucose tolerance tests of mice fed a ND or HFD. Data are presented as mean ±SD, n=5.

(C)   Weight in grams of mice fed a ND or HFD Data are presented as mean ± SD, n=5.

Supplementary Table 1: Over representation analysis of DSGs in primary osteocytes cultured in HG conditions.

Supplementary Table 2: Top 20 RNA binding motifs around differentially spliced transcripts in primary osteocytes cultured in HG conditions.

Supplementary Table 3: Over representation analysis of DEGS in after HuR KD.

Supplementary Table 4: DSGs, over representation analysis of DSGs and GOCC enrichment analysis after HuR KD.

Supplementary Table 5: Primers sequences, shRNA construct designs and antibodies used for Western blotting.

Supplementary Table 6: rMATS output from primary osteocytes cultured in HG conditions.

Supplementary Table 7: rMAPS output from primary osteocytes cultured in HG conditions.

Supplementary Table 8: rMATS output after HuR KD.

A
RBP motifs enriched around up SE DSGs

B
RBP motifs enriched around SE down DSGs

C

Motif MAP: HuR-TT[AT]GTTT

A % of sig. AS events by pattern

248398   3%

14   12.3   11.3   17.3   45

Total JCEC   SE   A3SS   MXE   A5SS   RI

B

Changes in inclusion levels

Farsb Rtkn Cd63 4931428F04Rik Bace1 Rptor Blnc1 Ankrd13d Rpp40
Rpl32 Klhl5 Rfc3 Zfp266 Sar1a Exosc7

Kirrel Get1 Rfc3 Tanc1 Syt8
Tom1l2 C230035I16Rik Pafah1b2 Itsn1 Ppil3 2610021A01Rik

A3SS   A5SS   MXE   RI   SE

event
A3SS
A5SS
MXE
RI
SE

C

cytosolic ribosome
ribosome
ribonucleoprotein complex
postsynapse
nucleolus
mitochondrion
nucleoplasm
cytosol
nucleus
cytoplasm

-Log10 FDR

% Genes
30
20
10

% Genes
10
20
30

D Pathway Enrichment of DSG after HuR KD

Major_pathway_of_rRNA_processing_in_the_nucleolus_and_cytosol rRNA_processing

Ribosome

Eukaryotic_Translation_Initiation
Cap-dependent_Translation_Initiation
(NMD)_enhanced_by_the_Exon_Junction_Complex_(EJC) Nonsense-Mediated_Decay_(NMD)

(NMD)_independent_of_the_Exon_Junction_Complex_(EJC)
cytoplasmic_translation

RNA_splicing
Phagosome
chromatin_organization
autophagy Spliceosome
rRNA_processing
translation_at_postsynapse
Non-alcoholic_fatty_liver_disease
HIF-1_signaling_pathway
translation_at_presynapse
Insulin_signaling_pathway

Gene counts in Geneset

padj value (log 10)

log10_padj

Geneset size
50
90
180

A

p-AKT/AKT
*
Mock^{KD}   YB^{KD}

AKT/β-actin
ns
Mock^{KD}   YB^{KD}

Mock KD   YBKD

P-AKT

AKT

β-actin

B

InsR mRNA expression level (InsR/GAPDH)
*
Mock^{KD}   YB^{KD}

A
RBP motifs enriched around down MXE DSGs

B
RBP motifs enriched around down A5SS DSGs

C
RBP motifs enriched around down A3SS DSGs

D
RBP motifs enriched around down RI DSGs

A FBG 16W

B AUC GTT 16W

C Weights 16W