Comparative analysis of the EF-1α Intergenic Region in *Babesia divergens* isolates: Insights
into TA Repeat Variation and Potential Regulatory Implications

Sezayi Ozubek[1], Alejandro Sanchez-Flores[2*], Estrella Montero[3*], Heba Alzan[4,5], Carlos E.
Suarez[5], Ricardo Grande[2], Aitor Gil-Iglesias[3], Munir Aktas[1], Luis Miguel Gonzalez[3]

[1] Department of Parasitology, Faculty of Veterinary Medicine, University of Firat, Elazig,
Turkiye

[2] Unidad Universitaria de Secuenciación Masiva y Bioinformatica, Institute of Biotechnology,
UNAM, Cuernavaca, México.

[3] Laboratorio de Referencia e Investigacion en Parasitologıa, Centro Nacional de Microbiología,
ISCIII Majadahonda, Madrid, Spain

[4] Parasitology and Animal Diseases Department, National Research Center, Dokki, Giza, Egypt

[5] Department of Veterinary Microbiology and Pathology, College of Veterinary Medicine,
Washington State University, Pullman, WA, USA

*Correspondence: alejandro.sanchez@ibt.unam.mx; estrella.montero@isciii.es.

Short title:

The EF-1α locus variation in *Babesia divergens* and other species

Abstract:

Babesia divergens, a zoonotic tick-borne pathogen, causes bovine and human babesiosis in

Europe. The Elongation Factor 1 alpha (EF-1α) protein is important in many cellular processes

and has emerged as a possible target for subunit vaccine development against parasitic infections,

and its intergenic region (IG) is an important tool for genetic manipulation of *Babesia* parasites.

While the EF-1α locus of *B. divergens* has been described, structural variation between isolates

was poorly defined. In order to fill this gap, we performed a comparative analysis of the EF-1α-

IG region in *B. divergens* human (Rouen 87, and Spanish sample), and bovine (Turkiye) host

isolates. Our findings revealed both conserved and variable elements, particularly in TA nucleotide

repeat numbers and IG sequence length. The Spanish isolate exhibited the highest TA repeat

expansion, whereas the Rouen 87 strain had the shortest IG region. Given the known role of repeat-

rich promoter elements in gene regulation, these differences may influence EF-1α transcription.

Additionally, these findings provide insights into the evolutionary divergence of *B. divergens* and

its host adaptation mechanisms. This study establishes a foundation for future gene editing and

transfection strategies, where selecting intergenic sequences with varying TA repeats could

optimize transfection efficiency and explain phenotypic differences between isolates from

different hosts or regions.

Keywords: *Babesia divergens*, comparative analysis, Elongation Factor 1 alpha (EF-1α), host

range

Introduction

Elongation factor 1 alpha (EF-1α) is among the most abundant proteins ever reported to be

expressed by eukaryotic cells and is highly conserved [1,2,3]. Traditionally, EF-1α is chiefly

acknowledged for its contribution to the translation machinery, facilitating the binding of

aminoacyl-tRNA to the ribosomal A site in a GTP-dependent manner [4]. Nevertheless, recent

studies have revealed how it functions in various cellular processes such as microtubule severing,

ubiquitin dependent proteolysis of N-terminal blocked proteins and cytoskeletal rearrangements

[3,5]. In addition, EF-1α plays a crucial role in fundamental cellular events including of

programmed apoptosis in which is up-regulated [3,6,7]. Further studies have also highlighted the

important role of EF-1α in other apicomplexan parasites, including *Cryptosporidium parvum* and

Toxoplasma gondii, for the entry process of these parasites into host cells. This suggest that EF-

1α could be a promising target for vaccine development against cryptosporidiosis and

toxoplasmosis [8,9]. Furthermore, results from studies on vaccination against avian coccidiosis

indicate that EF-1α is a promising candidate antigen for eliciting cross-protective immunity against

this disease in poultry, highlighting its potential for veterinary use [10].

Babesia are *T. gondii* and *Plasmodium*-related apicomplexan tick-borne parasites responsible for

acute and persistent infections in vertebrate hosts, including cattle and humans. Advancements in

gene editing and transfection systems are essential for unraveling the molecular intricacies of

Babesia parasites and enhancing vaccine development strategies [11,12,13]. Among the crucial

components, facilitating these advancements is the EF-1α, which plays a pivotal role in various

cellular processes. In both *Plasmodium* and *Babesia* parasites, the *EF-1α* gene exhibits a

distinctive organization, featuring two identical head-to-head genes, each separated by intergenic

regions [11,14]. The EF-1α intergenic region (EF-1α-IG) hosts robust promoters capable of driving

the expression of both *EF-1α* genes efficiently. This unique characteristic of the EF-1α-IG region

also enables it to effectively promote the expression of foreign genes in transiently and stably

transfected *Babesia bovis* and *Babesia bigemina* parasites [11,12,15]. The utilization of these

strong promoters within the EF-1α-IG region has dramatically accelerated the development of

transfection and gene editing systems in both *Plasmodium* and *Babesia* parasites [11,16].

Researchers have successfully exploited these promoters to introduce foreign genes into parasite

genomes, facilitating precise genetic manipulations, the development of novel vaccines, and

sophisticated studies on parasite biology. This breakthrough has significantly improved gene

function analysis, advanced our understanding of the molecular mechanisms underlying parasite

pathogenesis and host-parasite interactions [13,17,18], and became an important tool for

developing novel vectored vaccines [19,20,21,22].

Babesia divergens, one of the three major species of *Babesia* causing babesiosis in cattle, is

particularly significant as a zoonotic pathogen playing a role in human babesiosis in Europe, in

addition to its impact on cattle [23,24]. Compared to *B. bovis* and *B. bigemina*, limited information is available about the infectivity and pathogenicity of *B. divergens*, leaving many questions unanswered regarding important processes. Understanding the gaps of knowledge on gene function and other mechanisms involved in the biology of *B. divergens* will likely require the identification of novel promoters that can aid in the development of gene editing and transfection systems. Thus far, only one study has drawn attention in this regard, describing a stable transfection system for *B. divergens* based on the promoter of the *elongation factor Tu GTP binding domain family protein* (*ef-tgtp*) gene [25]. Unlike other *Babesia* species found in cattle, *B. divergens* has a broader host spectrum, including humans. We hypothesize that the EF-1α locus of *B. divergens,* a parasite able to infect bovine and human hosts, exhibits structural and regulatory variations among different isolates, which may be linked to its host adaptability, evolutionary divergence, and geographic distribution. Investigating these differences will enhance our understanding of *B. divergens* biology and provide insights into potential mechanisms of gene regulation and parasite-host interactions.

Here, based on existing genome data and NGS and Sanger sequencing, we focused on conducting a comparative analysis of the EF-1α locus from *B. divergens* isolates obtained from humans and cattle in different geographical regions, using the *B. divergens* Rouen 87 (R87) genome as a reference [26,27]. Our approach focused on conducting a comparative analysis of the EF-1α locus, which serves as a pivotal component in transfection investigations, among *B. divergens* isolates sourced from infected bovines in Turkiye and from patients with severe babesiosis in Spain [28]. The data generated in this study implies the possible influence of evolutionary selective forces shaping key sequence and structural differences in the EF-1α locus between these two isolates.

Results

EF-1α locus organization and synteny in *Babesia divergens*

In apicomplexan parasites, the *EF-1α* gene locus predominantly resides between the

Ribonucleoside diphosphate reductase and *Glutamyl tRNA* genes as demonstrated by conserved

synteny between *Theileria* and *Babesia* species. Similar to *B. bovis*, *B. bigemina*, *Babesia* sp.

Xinjiang and *Theileria equi*, the *B. divergens* R87 and Spanish and Turkiye isolates also possess

two identical *EF-1α* genes located within the EF-1α locus (Figure 1). The biological significance

of remarkable conserved synteny among all these apicomplexan parasites remains uninvestigated.

Figure 1: Schematic representation of gene localization and synteny maps of the *EF-1α* gene

locus.

EF-1α coding sequence conservation across *Babesia divergens* isolates

Comparative analysis of the EF-1α locus in *B. divergens* isolates (R87, Spanish, and Turkiye

isolates) revealed that the *EF-1α* gene sequences (1347 bp) were 100% identical across all isolates.

Each of the two EF-1α ORFs from three isolates of *B. divergens* encodes identical proteins of 448

amino acids, with an estimated molecular mass of 49.39 kDa, which is the same size as those

encoded by *B. bovis* (XP_001610983.1), *B. bigemina* (XP_012766720.1), *Babesia caballi*

(GIX62688.1), *Babesia gibsoni* (KAK1443338.1), *Babesia duncani* (KAK2195768.1), *Babesia*

ovis (GFE54231.1), *Babesia* sp. Xinjiang (XP_028869924.1), *Theileria parva* (XP_766247.1),

Theileria annulata (XP_954051.1), and *Theileria orientalis* (UKK00547.1). The EF-1α *B.*

divergens R87, Spanish, and Turkiye ORFs show identical sequences (100% identity) and share

significant similarity with homologous sequences found in *B. gibsoni* (97.99% identity), *B. caballi*

(97.54% identity), *B. bigemina* (96.65% identity), *Babesia* sp. Xinjiang (96.43% identity), *B. bovis*

(94.87% identity), *B. ovis* (94.20% identity), *B. duncani* (93.97% identity), *T. orientalis* (92.41%

identity), *T. parva* (89.51% identity), *T. annulata* (89.06% identity), and *B. microti* (86.58%

identity) (Supplementary Figure 1). Notably, key regions involved in guanosine 50-triphosphate

(GTP) binding are conserved across these organisms, as depicted in Supplementary Figure 2. A

phylogenetic analysis using the EF-1α protein sequence of R87, Spanish, and Turkiye isolates,

along with other related piroplasma species, showed that both isolates clustered within the same

clade as *B. gibsoni* (Figure 2a). Similarly, in the phylogenetic analysis of *B. divergens* R87,

Spanish (MG944238.1), and Turkiye (PP663285.1) *18S rRNA* gene sequences, it was observed

that they were in the same clade with *B. divergens* isolates obtained from different hosts from

GenBank (Figure 2b).

Figure 2: a) Phylogenetic analyses of piroplasm EF-1α protein sequences by maximum likelihood method based on the Le_Gascuel_2008 model [29]. Sequences were obtained from the GenBank database. The EF-1α protein sequence of *Plasmodium falciparum* was utilized as an outgroup. b) Phylogenetic analyses of piroplasm *18S rRNA* gene sequences by maximum likelihood method based on the Tamura-Nei model [30]. The *18S rRNA* gene sequence of *P. falciparum* was utilized as an outgroup. *Babesia divergens* R87, Spanish, and Turkiye isolates of EF-1α are highlighted in bold.

Features, sequence alignment and nucleotide variability in IG region

The EF-1α-IG regions of the three distinct *B. divergens* isolates analyzed hereby are not related in

sequence to the IG regions of other related *Babesia* parasites. However, the *B. divergens* EF-1α-

IG regions do have structural similarities with other *Babesia* parasites, including the presence of

terminal inverted repeat (IR) regions of 237 bp, which were conserved among all isolates. In

addition, similar to what was found on *B. bovis* and *B. bigemina* [11,12], the IR regions harbor a

153 bp intron, maintaining sequence integrity across isolates (Figure 3).

However, the comparative analysis of the EF-1α locus sequences in the three *B. divergens* isolates

(R87, Spanish, and Turkiye isolates) revealed sequence differences in the IG region. While the

EF-1α gene sequences were found to be 100% identical among all isolates, the IG region length

varied across the isolates, with the R87 strain containing 5 TA repeats and measuring 1380 bp, the

Spanish isolate harboring 24 TA repeats with an IG region of 1418 bp, and the Turkiye isolate

exhibiting 8 TA repeats with an IG region of 1386 bp (Figure 3). Overall, these findings indicate

notable differences in the IG region between *B. divergens* isolates originating from different

geographical regions and host sources.

Figure 3: Schematic representation of the EF-1α-IG region between *B. divergens* isolates. The

terminal inverted repeat (IR) regions (red), intron (green, 153 bp), and TA repeats (blue) are

highlighted. The IG region length varies due to differences in TA repeat numbers, with 5 TA

repeats in R87 (1380 bp), 24 in the Spanish isolate (1418 bp), and 8 in the Turkiye isolate (1386

bp).

Multiple sequence alignment of the IG region across the three *B. divergens* isolates further

confirmed variations in TA repeat numbers and additional nucleotide substitutions. The TA repeat

sequences were located around position ~513–516 bp across all isolates but varied in number.

Besides TA repeat variations, additional sequence polymorphisms were identified across the IG

region (Figure 4). The results were consistent and validated using Nanopore and Sanger

sequencing technologies.

Figure 4: Multiple sequence alignment of the IG region within the EF-1α locus between *B.*

divergens isolates. Nucleotide variations are highlighted with black boxes, while differences in TA

repeat numbers are also marked. In addition to *B. divergens*, the intergenic region was examined

in other zoonotic *Babesia* species, including *Babesia* MO1 and *B. duncani*. While the EF-1α locus

in *B. duncani* lacked TA repeats, the *Babesia* MO1 genome contained a short stretch of 3 TA

repeats. These findings indicate that the presence and expansion of TA repeats may be specific to

B. divergens (Supplementary Figure 3).

Validation of TA repeat variations in the IG region

To confirm the sequence differences in the TA repeat region among *B. divergens* isolates, a

combination of PCR amplification, Sanger sequencing, and NGS approaches was employed. For

the R87 isolate, the IG region was amplified using the Bdiv-Ef-For1/IGRev1 primer pair, yielding

an amplicon of approximately 2.8 kb, which was subsequently validated by Sanger sequencing.

The repeat structure identified in the sequencing data confirmed the presence of 5 TA repeats,

consistent with initial findings. For the Turkiye isolate, the IG region was amplified in two

overlapping fragments using the Bdiv-Ef-For1/Sez-For1 and Sez-For1-Rev/GlutamylR1 primer

sets. These amplicons were subjected to NGS analysis, which verified the 8 TA repeats and

confirmed the overall sequence integrity of the IG region. For the Spanish isolate, initial attempts

to amplify the full IG region using Bdiv-Ef-For1/IGRev1 primers were unsuccessful, likely due to

sequence divergence at the primer binding sites. An alternative primer pair, IGLMFR1/IGRev1,

was used to amplify ~1 kb fragment, which was validated through Sanger sequencing. The results

confirmed the presence of 24 TA repeats, aligning with the Nanopore sequencing data.

Figure 5: Multiple sequence alignment of the TA repeat region across *B. divergens* isolates using Nanopore, Sanger, and NGS sequencing methods. The black-highlighted regions indicate TA repeats.

Discussion

Since the discovery of *Babesia*, the availability of transfection and gene editing methods arguably emerged as the most significant developments, after *Babesia in vitro* cultures. Transfection technologies (together with related gene editing methods), which involves the introduction of foreign genetic material into cells, has revolutionized the study of various diseases, including those caused by *Babesia* parasites [15]. This technique has enabled researchers to manipulate cellular processes, elucidate pathogenic mechanisms, and explore potential therapeutic interventions. Thus, transfection represents a crucial tool in advancing our understanding of *Babesia* infections and developing strategies for their prevention and treatment [13,17]. EF-1α-IG region contains bi-directional promoters capable of efficiently driving expression of *EF-1α* genes. Their utilization has significantly advanced transfection systems in *Babesia* [11,15]. In this study, we explored the structural and sequence variability of the EF-1α-IG regions across three *B. divergens* isolates from humans and cattle to better understand its potential role in gene regulation and parasite adaptation. The findings indicated that while the EF-1α coding sequences remain highly conserved, there is variability in the IG region, specifically in TA repeat numbers and the length of the terminal IR

sequences, suggesting that these elements could contribute to transcriptional regulation and genomic stability.

The high level of conservation of the *EF-1α* gene observed in different strains of *B. divergens* and its similarity to the EF-1α protein of the two other pathogenic bovine babesiosis species, *B. bovis* and *B. bigemina*, highlights the potential use of EF-1α in subunit vaccines. These findings revealed that EF-1α protein has potential to be a suitable antigen for vaccine development, as it was previously suggested for other important protozoa such as *T. gondii* [9] and *C. parvum* [8], which are also members of the apicomplexan phylum. Additionally, the broad applicability of the vaccine in suppressing different parasitic infections was further verified by the existing cross-protective immunity against *Eimeria tenella* and *Eimeria maxima* in broiler chickens vaccinated with recombinant EF-1α protein [10]. Thus, EF-1α protein appears as a good candidate for the development of vaccines against babesiosis, taking advantage of its potential to induce cross-immunity against different species.

Comparisons of the IG regions among the three *B. divergens* isolates (R87, Spanish, and Turkiye isolates) provided novel insights into the structural organization of this region. This study challenged a previously established assumption that the IG regulatory regions of the EF-1α loci are fully conserved among *Babesia* spp. and parasite isolates, both in terms of sequence and structural organization. Despite having identical *EF-1α* gene ORFs organized in a typical head-to-head fashion, *B. divergens* isolates exhibited striking differences in their IG regions, despite sharing identical *18S rRNA* sequences. One of the most unexpected findings of this study was the variability in TA repeat numbers within the IG region. The Spanish isolate exhibited the highest number of TA repeats (n=24), while the Turkiye isolate contained 8, and the R87 strain had only 5. Given that repeat sequences in intergenic regions can affect promoter activity, transcription

factor binding, and chromatin accessibility [31], these differences may influence *EF-1α* gene

expression. The conserved 237 bp terminal IR region and the 153 bp intron within the untranslated

region suggest that the IG region may have multiple regulatory roles. Inverted repeats (IRs) are

often associated with promoter function, transcriptional regulation, while intronic sequences

within untranslated regions can influence alternative splicing, mRNA stability, or post-

transcriptional control. The conservation of this structure across isolates suggests its functional

importance in *EF-1α* gene regulation.

As mentioned, the differences within the IG regions among *B. divergens* isolates analyzed hereby

predominantly occur in the initial segment of the putative promoter regions. Although

investigations into EF-1α-IG region variations across different strains remain limited, examination

of these regions from *B. bovis* strains in Argentina, Mexico, and Australia suggests a notable

degree of conservation among them [11]. The distinctions between *B. bovis* and *B. divergens*,

despite both being etiological agents of babesiosis in cattle, are evident in their geographical

distribution, different tick vector preferences, and the clinical manifestations of babesiosis [23,32].

Moreover, the broader host spectrum of *B. divergens*, including its zoonotic potential, highlights

its adaptability and significance beyond the confines of bovine hosts. It is thus logical to speculate

that the discrepancies observed in the EF-1α-IG region could be attributed to the differing host

preferences of *B. divergens*. This is particularly noteworthy in our study where R87, Spanish, and

Turkiye isolates, whose elongation loci we compared, were obtained from humans (R87, and

Spanish isolates) and cattle (Turkiye) respectively. Surprisingly, the EF-1α-IG region from

parasites that infect humans presented differences that could also be related to virulence or

adaptation to laboratory conditions. Such findings raise intriguing questions regarding factors

involved in parasite evolution, and the potential influence of host-specific factors on the genetic

makeup and regulatory mechanisms of *Babesia* species, warranting further investigation into the

intricate interplay between pathogen genetics and host environments. As they conflict with a

previously established paradigm, the substantial differences found in the IG regions of the *EF-1α*

genes between three different strains of *B. divergens*, as confirmed by *18S rRNA gene* analysis are

found for the first time among piroplasmid parasites. Moreover, as previously noted, these

differences may reflect important adaptive events affecting the development of the parasites

among distinct invertebrate and/or vertebrate hosts.

Surprisingly, upon expanding our comparative analysis of the EF-1α IG regions found in other

zoonotic *Babesia* species, *B. duncani* and *Babesia* MO1, striking differences were also noted in

their EF-1α IG regions. *Babesia duncani*, on the other hand, contained no TA repeats, while only

3 TA motifs were present in the MO1 isolate. This suggests that TA repeat expansion within the

EF-1α-IG region may represent a species-specific regulatory feature rather than a common trait of

all zoonotic *Babesia*. Further comparative studies across more isolates will be essential to

determine whether these differences contribute to host tropism, gene expression variability, or

virulence mechanisms.

Conclusion

This study provides new insights into the structural variability of the EF-1α-IG region in *B.*

divergens isolates from different hosts, revealing differences in TA repeat numbers that may

contribute to gene regulation and parasite adaptation. While our findings highlight the potential

regulatory role of this region, functional validation through promoter activity assays, gene

expression analysis, and transfection studies is still needed to determine its precise impact on EF-

1α expression. Given the zoonotic nature of *B. divergens*, future research should explore whether

similar variations in the IG region exist in *B. divergens* isolates from cattle and humans.

Additionally, the gerbil model could be used to investigate potential intergenic region variations

across different vertebrate hosts to better understand the role of this region in host adaptation.

Further investigation of the functional implication of these variations may provide valuable

insights into transcriptional control across diverse parasite strains, provide valuable information

on the mechanisms driving *B. divergens* evolution and improve transfection strategies for

functional genomic studies. Overall, this study lays the groundwork for future research into gene

regulation and host-pathogen interactions in zoonotic *Babesia* species.

Methods

Babesia divergens **- Rouen 87 Strain (R87)**

Babesia divergens R87 is a reference strain originally isolated from a human case of babesiosis in

France [33]. This strain has been widely used in molecular and genomic studies due to its well-

characterized genome sequence [27]. The EF-1α locus data from this strain were utilized in this

study, obtained from the *B. divergens* genome assembly described by [27].

Babesia divergens-**Spanish sample**

Babesia divergens Spanish strain was isolate from a blood sample of elderly patient from Asturias,

Spain who suffers a fulminant fatal babesiosis [28]. The blood sample was cultured in vitro in

human A+ RBCs and RPMI 1640 (Gibco-Life Technology, 11875093) supplemented with 10%

human serum (Gibco, 11021037), 7.5% (w/v) sodium bicarbonate solution (Lonza Group Ltd,

Basel, Switzerland, 144-55-8), and 100 μmol/L hypoxanthine (Sigma-Aldrich Corporation, St

Louis, MO, H9377) at a pH of 7.3. Cells were cultured at 37°C in a humidified atmosphere of 5%

CO_2. The culture medium was replaced every 24 h, and parasitemia was monitored by examining

Giemsa-stained blood smears by a light microscope.

Babesia divergens-Turkiye

Babesia divergens-Turkiye isolate was obtained from a 3-year-old cow exhibiting severe babesiosis symptoms, including high fever, anemia, and hemoglobinuria, in Bartin province of Turkiye in 2023. Microscopic examination revealed a parasitemia level of 1.5%. Following the collection of blood stabilate from the animal, treatment was administered using imidocarb dipropionate [34]. To determine the species of *Babesia*, Reverse Line Blot (RLB) analysis was conducted. Total DNA extraction from blood samples was performed using the PureLink™ Genomic DNA Mini Kit (Invitrogen Corporation, Carlsbad, United States) according to the manufacturer's instructions. The hypervariable V4 region of the piroplasm *18S rRNA* gene was then amplified using primers RLB-F2 and RLB-R2-biotin [35] specifically for the RLB assay. RLB was carried out on the PCR product, following the previously described method [36]. After amplification, 20 µL of all PCR products obtained from each DNA sample were diluted to a final volume of 150 µL with 2X SSPE/0.1% SDS buffer. For RLB hybridization, the samples were heated at 95–100°C for 10 minutes in a Thermal Cycler and rapidly cooled on ice. Subsequently, the PCR products were hybridized with probes specific to the genera and species of *Babesia* and *Theileria*, which were linked to an RLB membrane [36].

Babesia MO1 and Babesia duncani

In this study, we used the EF-1α locus data of these species obtained from the *Babesia* MO1 [37] and *B. duncani* [38,39] assembly. Specifically, Bioproject PRJNA1032622 for *Babesia MO1* and Bioproject PRJNA821606 and locus OL804102 for *B. duncani.* To assess the presence of TA repeats in *Babesia* MO1 and *B. duncani*, we retrieved the EF-1α locus from *Babesia* MO1 and *B. duncani* genome assemblies (PRJNA1032622 and PRJNA821606, respectively). The intergenic regions were identified by locating the flanking genes (*ribonucleoside diphosphate reductase* and

glutamyl tRNA synthetase) and were subsequently aligned to detect potential tandem repeat

regions. The presence or absence of TA repeats was manually inspected by multiple sequence

alignment.

Nanopore sequencing

The total DNA from *Babesia* samples was used to prepare Oxford Nanopore libraries with the

SQK-LSK114 Ligation Sequencing Kit for 24 h in the MinION Platform with an R10.4 flow cell

following the manufacturer's protocol (Oxford Nanopore Technologies, Oxford, United

Kingdom). The base calling was performed using the software Dorado v0.8.3 with the duplex

option. The sequencing and basecalling was carried out at the Unidad Universitaria de

Secuenciación Masiva y Bioinformática, Instituto de Biotecnología, Universidad Nacional

Autónoma de México (Cuernavaca, Mexico).

Designing of primers for EF-1α locus analysis and PCR

Primers were designed utilizing sequences from *Ribonucleoside diphosphate reductase*

(Bdiv_023390c) and *Glutamyl tRNA* (Bdiv_023400) genes obtained from the Piroplasma DB

(https://piroplasmadb.org) and nanopore sequencing data to amplify the EF-1α locus. All primers

were designed using Primer Quest™ Tool (https://www.idtdna.com/pages/tools/primerquest)

(Supplementary Table 1, and Supplementary Figure 4). PCR amplification was carried out using

Phusion® High-Fidelity PCR Master Mix with GC Buffer (M0532S; NEB). The PCR was

performed in a total reaction volume of 20 μL containing 10 μL of 2X Phusion Master Mix, 1 μL

of each forward and reverse primer, 1 μL of template DNA, 7 μL of nuclease-free water.

Illumina sequencing and *de novo* assembly

Sequencing library was prepared using Nextera XT DNA Library Preparation Kit and sequencing

was performed by Illumina Miseq platform as paired end (PE) 2x150 bases reads. Raw NGS reads

(FASTQ) were quality checked by FASTQC [40] and trimmed by Trimmomatic v0.32 [41].

Demultiplexing and low-quality read filtering were performed via CLC Genomics Workbench

(Qiagen, Hilden, Germany). The *de novo* assembly constructed by CLC Genomic's de novo

assembly module with the parameters of minimum contig size as 200 bp, mismatch cost is 2, insert

cost is 3, deletion cost is 3, length fraction is 0.5, similarity fraction is 0.8, and paired read input

as minimum distance of 180.

Sanger sequencing

To validate the assembled EF-1α-IG region, Sanger sequencing was performed on PCR amplified

IG regions from the *B. divergens* isolates. PCR products were purified using the mi-Gel Extraction

Kit (Metabion international AG, Steinkirchen Germany) and sequenced using an ABI PRISM

3730XL DNA Analyzer (Applied Biosystems, San Francisco, CA, USA). The resulting

chromatograms were analyzed using manually inspected, corrected, and edited using Chromas Pro

program (McCarthy, Queensland, Australia) and Laser Gene 12.1 program (DNAStar, Madison,

WI, USA), and sequences were aligned against the de novo assembled IG region to confirm

accuracy and resolve potential misassembles.

Phylogenetic analysis

Phylogenetic relationships were inferred using Molecular Evolutionary Genetics Analysis (MEGA

11) [42]. The maximum likelihood (ML) method was employed to construct phylogenetic trees

based on EF-1α protein sequences and *18S rRNA* gene sequences from various piroplasm species.

Bootstrap support values were calculated with 1,000 replicates to assess tree robustness.

Data availability

All data is provided within the manuscript or supplementary information files.

Ethical statement

B. divergens DNA samples from the cattle were obtained as part of routine veterinary diagnostic

procedures from a clinically infected animal. No experimental infection or additional intervention

was performed for research purposes. Therefore, Institutional Animal Care and Use Committee

(IACUC) or Ethics Committee approval was not required. The *B. divergens* Spanish strain was

obtained as part of routine diagnostic procedures, no additional ethics approval or informed

consent from hospitals or patients was required. Therefore, our study does not involve human

subjects, identifiable human tissue, or personal data, in accordance with the Declaration of

Helsinki.

References

1. Slobin, L. I. (1980). The Role of Eucaryotic Elongation Factor Tu in Protein Synthesis: The

Measurement of the Elongation Factor Tu Content of Rabbit Reticulocytes and Other Mammalian

Cells by a Sensitive Radioimmunoassay. European Journal of Biochemistry 110, 555–563. doi:

10.1111/j.1432-1033.1980.tb04898.x

2. Merrick, W. C. (1992). Mechanism and regulation of eukaryotic protein synthesis. Microbiol

Rev 56, 291–315. doi: 10.1128/mr.56.2.291-315.1992

3. Sasikumar, A. N., Perez, W. B., and Kinzy, T. G. (2012). The many roles of the eukaryotic

elongation factor 1 complex. WIREs RNA 3, 543–555. doi: 10.1002/wrna.1118

4. Riis, B., Rattan, S. I., Clark, B. F., and Merrick, W. C. (1990). Eukaryotic protein elongation

factors. Trends in biochemical sciences 15, 420–424.

5. Stapulionis, R., Kolli, S., and Deutscher, M. P. (1997). Efficient mammalian protein synthesis requires an intact F-actin system. Journal of Biological Chemistry 272, 24980–24986.

6. Chen, E., Proestou, G., Bourbeau, D., and Wang, E. (2000). Rapid up-regulation of peptide elongation factor EF-1α protein levels is an immediate early event during oxidative stress-induced apoptosis. Experimental cell research 259, 140–148.

7. Lamberti, A., Longo, O., Marra, M., Tagliaferri, P., Bismuto, E., Fiengo, A., et al. (2007). C-Raf antagonizes apoptosis induced by IFN-α in human lung cancer cells by phosphorylation and increase of the intracellular content of elongation factor 1A. Cell Death & Differentiation 14, 952–962.

8. Matsubayashi, M., Teramoto-Kimata, I., Uni, S., Lillehoj, H. S., Matsuda, H., Furuya, M., et al. (2013). Elongation factor-1α is a novel protein associated with host cell invasion and a potential protective antigen of Cryptosporidium parvum. Journal of Biological Chemistry 288, 34111–34120.

9. Wang, S., Zhang, Z., Wang, Y., Gadahi, J. A., Xu, L., Yan, R., et al. (2017). Toxoplasma gondii elongation factor 1-alpha (TgEF-1α) is a novel vaccine candidate antigen against toxoplasmosis. Frontiers in microbiology 8, 168.

10. Lin, R.-Q., Lillehoj, H. S., Lee, S. K., Oh, S., Panebra, A., and Lillehoj, E. P. (2017). Vaccination with Eimeria tenella elongation factor-1α recombinant protein induces protective immunity against E. tenella and E. maxima infections. Veterinary Parasitology 243, 79–84.

11. Suarez, C. E., Norimine, J., Lacy, P., and McElwain, T. F. (2006). Characterization and gene expression of Babesia bovis elongation factor-1α. International journal for parasitology 36, 965–973.

12. Silva, M. G., Knowles, D. P., Mazuz, M. L., Cooke, B. M., and Suarez, C. E. (2018). Stable

transformation of Babesia bigemina and Babesia bovis using a single transfection plasmid.

Scientific reports 8, 6096.

13. Hakimi, H., Asada, M., and Kawazu, S. (2021). Recent advances in molecular genetic tools

for babesia. Veterinary Sciences 8, 222.

14. Vinkenoog, R., Sperança, M. A., van Breemen, O., Ramesar, J., Williamson, D. H., Ross-

MacDonald, P. B., et al. (1998). Malaria parasites contain two identical copies of an elongation

factor 1 alpha gene. Molecular and biochemical parasitology 94, 1–12.

15. Suarez, C. E., and McElwain, T. F. (2010). Transfection systems for Babesia bovis: a review

of methods for the transient and stable expression of exogenous genes. Veterinary parasitology

167, 205–215.

16. Fernandez-Becerra, C., Azevedo, M. de, Yamamoto, M. M., and Portillo, H. del (2003).

Plasmodium falciparum: new vector with bi-directional promoter activity to stably express trans

genes. Available at: https://www.cabidigitallibrary.org/doi/full/10.5555/20033124083 (Accessed

March 13, 2025).

17. Suarez, C. E., Alzan, H. F., Silva, M. G., Rathinasamy, V., Poole, W. A., and Cooke, B. M.

(2019). Unravelling the cellular and molecular pathogenesis of bovine babesiosis: is the sky the

limit? International journal for parasitology 49, 183–197.

18. Johnson, W. C., Hussein, H. E., Capelli-Peixoto, J., Laughery, J. M., Taus, N. S., Suarez, C.

E., et al. (2022). A transfected Babesia bovis parasite line expressing eGFP is able to complete the

full life cycle of the parasite in mammalian and tick hosts. Pathogens 11, 623.

19. Laughery, J. M., Knowles, D. P., Schneider, D. A., Bastos, R. G., McElwain, T. F., and Suarez,

C. E. (2014). Targeted surface expression of an exogenous antigen in stably transfected Babesia

bovis. PLoS One 9, e97890.

20. Oldiges, D. P., Laughery, J. M., Tagliari, N. J., Leite Filho, R. V., Davis, W. C., da Silva Vaz

Jr, I., et al. (2016). Transfected Babesia bovis expressing a tick GST as a live vector vaccine. PLoS

neglected tropical diseases 10, e0005152.

21. Suarez, C. E., Bishop, R. P., Alzan, H. F., Poole, W. A., and Cooke, B. M. (2017). Advances

in the application of genetic manipulation methods to apicomplexan parasites. International journal

for parasitology 47, 701–710.

22. Mazuz, M. L., Laughery, J. M., Lebovitz, B., Yasur-Landau, D., Rot, A., Bastos, R. G., et al.

(2021). Experimental infection of calves with transfected attenuated Babesia bovis expressing the

Rhipicephalus microplus Bm86 Antigen and eGFP Marker: Preliminary studies towards a dual

anti-tick/Babesia vaccine. Pathogens 10, 135.

23. Ozubek, S., Bastos, R. G., Alzan, H. F., Inci, A., Aktas, M., and Suarez, C. E. (2020). Bovine

babesiosis in Turkey: Impact, current gaps, and opportunities for intervention. Pathogens 9, 1041.

24. Hildebrandt, A., Zintl, A., Montero, E., Hunfeld, K.-P., and Gray, J. (2021). Human babesiosis

in Europe. Pathogens 10, 1165.

25. Cubillos, E. F., Snebergerova, P., Borsodi, S., Reichensdorferova, D., Levytska, V., Asada,

M., et al. (2023). Establishment of a stable transfection and gene targeting system in Babesia

divergens. Frontiers in Cellular and Infection Microbiology 13. Available at:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC10753763/ (Accessed April 30, 2024).

26. Cuesta, I., González, L. M., Estrada, K., Grande, R., Zaballos, Á., Lobo, C. A., et al. (2014).

High-Quality Draft Genome Sequence of Babesia divergens , the Etiological Agent of Cattle and

Human Babesiosis. Genome Announc 2, e01194-14. doi: 10.1128/genomeA.01194-14

27. Gonzalez, L. M., Estrada, K., Grande, R., Jiménez-Jacinto, V., Vega-Alvarado, L., Sevilla, E.,

et al. (2019). Comparative and functional genomics of the protozoan parasite Babesia divergens

highlighting the invasion and egress processes. PLoS neglected tropical diseases 13, e0007680.

28. Asensi, V., González, L. M., Fernández-Suárez, J., Sevilla, E., Navascués, R. Á., Suárez, M.

L., et al. (2018). A fatal case of Babesia divergens infection in Northwestern Spain. Ticks and tick-

borne diseases 9, 730–734.

29. Le, S. Q., and Gascuel, O. (2008). An improved general amino acid replacement matrix.

Molecular biology and evolution 25, 1307–1320.

30. Tamura, K., and Nei, M. (1993). Estimation of the number of nucleotide substitutions in the

control region of mitochondrial DNA in humans and chimpanzees. Molecular biology and

evolution 10, 512–526.

31. Quilez, J., Guilmatre, A., Garg, P., Highnam, G., Gymrek, M., Erlich, Y., et al. (2016).

Polymorphic tandem repeats within gene promoters act as modifiers of gene expression and DNA

methylation in humans. Nucleic acids research 44, 3750–3762.

32. Schnittger, L., Ganzinelli, S., Bhoora, R., Omondi, D., Nijhof, A. M., and Florin-Christensen,

M. (2022). The Piroplasmida Babesia, Cytauxzoon, and Theileria in farm and companion animals:

Species compilation, molecular phylogeny, and evolutionary insights. Parasitology research 121,

1207–1245.

33. Gorenflot, A., Brasseur, P., Precigout, E., L'Hostis, M., Marchand, A., and Schrevel, J. (1991).

Cytological and immunological responses toBabesia divergens in different hosts: Ox, gerbil, man.

Parasitol Res 77, 3–12. doi: 10.1007/BF00934377

35. Georges, K., Loria, G. R., Riili, S., Greco, A., Caracappa, S., Jongejan, F., et al. (2001).

Detection of haemoparasites in cattle by reverse line blot hybridisation with a note on the

distribution of ticks in Sicily. Veterinary parasitology 99, 273–286.

34. Mosqueda, J., Olvera-Ramírez, A., Aguilar-Tipacamú, G., and Cantó, G. (2012). Current

Advances in Detection and Treatment of Babesiosis. Curr Med Chem 19, 1504–1518. doi:

10.2174/092986712799828355

36. Aktas, M., and Ozubek, S. (2015). Molecular and parasitological survey of bovine piroplasms

in the Black Sea region, including the first report of babesiosis associated with Babesia divergens

in Turkey. Journal of medical entomology 52, 1344–1350.

37. Singh, P., Vydyam, P., Fang, T., Estrada, K., Gonzalez, L. M., Grande, R., et al. (2024).

Insights into the evolution, virulence and speciation of Babesia MO1 and Babesia divergens

through multiomics analyses. Emerging Microbes & Infections 13, 2386136. doi:

10.1080/22221751.2024.2386136

38. Wang, S., Li, D., Chen, F., Jiang, W., Luo, W., Zhu, G., et al. (2022). Establishment of a

transient and stable transfection system for babesia duncani using a homologous recombination

strategy. Frontiers in Cellular and Infection Microbiology 12, 844498.

39. Singh, P., Lonardi, S., Liang, Q., Vydyam, P., Khabirova, E., Fang, T., et al. (2023). Babesia

duncani multi-omics identifies virulence factors and drug targets. Nature microbiology 8, 845–

859.

40. Bioinformatics, B. (2018). Babraham Bioinformatics-FastQC a quality control tool for high

throughput sequence data.

41. Bolger, A. M., Lohse, M., and Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics 30, 2114–2120.

42. Tamura, K., Stecher, G., and Kumar, S. (2021). MEGA11: molecular evolutionary genetics

analysis version 11. Molecular biology and evolution 38, 3022–3027.

Acknowledgements

502 We thank "Centro de Transfusiones de la Comunidad de Madrid", which provided the human A+

503 blood from healthy volunteer donors. This work was funded by grants from Health Institute Carlos

504 III (ISCIII; PI20CIII-00037 to E.M. and L.G.M) and Comunidad Autónoma de Madrid (CAM;

505 TEC-2024/BIO-66/SALAINDEC-CM to EM and L.G.M.).

Author Contrubitions

507 Conceptualization: SO, ASF, EM and CES; Methodology and experiments: OS, LMG, HA, AGI,

508 RGB, MA and RG; Sequencing and bioinformatic analysis: SO, RG, LMG and ASF, Formal

509 analysis and data interpretation SO, EM, ASF and LMG; Writing original draft SO; Writing,

510 reviewing and editing ASF, EM, LMG. All authors read and approved of the final manuscript.

Competing interests

512 The authors declare no competing interests.

Ethics statement

Babesia divergens Spanish isolate were cultured using human A+ blood obtained from healthy

volunteer donors. The blood was sourced from the Blood Transfusion Center of Madrid, Spain,

adhering to approved protocols and in compliance with the relevant institutional guidelines and

regulations. *Babesia divergens* Turkiye isolate was obtained from a clinically diagnosed babesiosis

case in a cow. The sample was submitted for diagnostic purposes to the Department of

Parasitology, Faculty of Veterinary Medicine, Fırat University. The study was conducted in

accordance with institutional guidelines and regulations.

Additional information

Supplementary Information

Supplementary Figure 1: Percentage of identity of amino acid sequence of EF-1α among

different piroplasm species.

Supplementary Figure 2: Comparison of EF-1α protein sequences among *B. divergens* isolates

(R87, Spanish sample, and Turkiye isolates) and various piroplasm species. Three conserved EF-

1α regions with known Guanosine 5'-triphosphate (GTP)-Binding Site Function (GTP BS 1, 2, and

3) highlighted in boxes.

Supplementary Figure 3: Comparative alignment of the EF-1α-IG region across zoonotic

Babesia species. The TA repeat motifs (highlighted in black) show notable variability among *B.*

divergens isolates (R87, Spanish, and Turkiye), while *Babesia* MO1 contains only 3 TA motifs,

and *B. duncani* lacks any TA repeats.

Supplementary Figure 4: Schematic representation of the *B. divergens* EF-1α locus showing the

primers used for IG region amplification. IR (inverted repeat) and IG (intergenic region) are

highlighted, with primer positions indicated by arrows.

a)

B.divergens-Human-Rouen 1987 genome

100 ┐ **B.divergens-Cattle-Turkiye**

91 └ **B.divergens-Human-Spanish sample**

B.gibsoni-KAK1443338.1

B.bigemina-XP 012766720.1

82 ┌ 56 B.caballi-GIX62688.1

Babesiasp.Xinjiang-XP 028869924.1

96 ┌ 62 B.ovis-GFE54231.1

B.bovis-XP 001610983.1

B.duncani-KAK2195768.1

T.orientalis-UKK00547.1

84 ┌ T.parva-XP 766247.1

100 └ T.annulata-XP 954051.1

B.microti-XP 012649309.1

P.falciparum-XP 001350281.1

0.050

b)

75 ┐ B.divergens-Red-deer-GQ304524.1-France

B.divergens-Cattle-KP745627.1-Turkiye

82 **B.divergens-Human-MG944238.1-Spain**

B.divergens-Cattle-PP663285.1-Turkiye

B.divergens-Cattle-LC477140.1-Ireland

B.divergens-Red-deer-MG344781.1-Czechia

56 B.divergens-Human-LC477139.1-Ireland

B.divergens-Cattle-LC477143.1-Ireland

51 **B.divergens-Human-Rouen 1987 genome-France**

99 B.divergens-Cattle-FJ944825.1-France

B.divergens-Ixodes-persulcatus-LC279018.1-Japan

B.divergens-Roe-deer-AY572456.1-Slovenia

B.major-Cattle-EU622907.1-France

91 ┌ B.ovata-Cattle-MN900525.1-China

90 └ B.ovata-Cattle-AY081192.1-Korea

B.noakii-Cattle-MF576177.1-Bangladesh

60 85 ┌ B.bigemina-Cattle-DQ785311.1-Spain

97 B.bigemina-Cattle-KP745623.1-Turkiye

B.bigemina-Cattle-KM046917.1-Switzerland

B.occultans-Hyalomma-marginatum-HQ331479.1-Tunisia

100 B.occultans-Cattle-KP745626.1-Turkiye

B.occultans-Zebu-cattle-OK314934.1-Turkiye

B.bovis-Cattle-KP710223.1-China

100 ┌ B.bovis-Cattle-KP745628.1-Turkiye

100 └ B.bovis-Cattle-AY150059.1-Portugal

Plasmodium-falciparum-JQ627151.1

0.050

```
B. divergens R87            TTTGATGAGGTCTAATAATAACTAATCTGCAAAGGGTTAGTGAGGCCAATGGAAACACAA  60
B. divergens Spanish sample TTTGATGAGGTCTAATAATAACTAATCTGCAAAGGGTTAGTGAGGCCAATGGAAACACAA  60
B. divergens Turkiye        TTTGATGAGGTCTAATAATAACTAATCTGCAAAGGGTTAGTGAGGCCAATGGAAACACAA  60
                            ************************************************************

B. divergens R87            TGGGCAACACCACAACCGAATTAACAACATGGAAATATATATTTATAATGTTATGCCCCT  120
B. divergens Spanish sample TGGGCAACACCACAACCGAATTAACAACATGGAAATATATATTTATAATGTTATGCCCCT  120
B. divergens Turkiye        TGGGCAACACCACAACCGAATTAACAACATGGAAATATATATTTATAATGTTATGCCCCT  120
                            ************************************************************

B. divergens R87            ACGACACTTCTCATAATGACACAGCAGAACGTAGAGATCACGACACCCACACCACTAACA  180
B. divergens Spanish sample ACGACACTTCTCATAATGACACAGCAGAACGTAGAGATCACGACACCCACACCACTAACA  180
B. divergens Turkiye        ACGACACTTCTCATAATGACACAGCAGAACGTAGAGATCACGACACCCACACCACTAACA  180
                            ************************************************************

B. divergens R87            TCAAAAAACCAAACAAACCAAAACAGTTTGACAAAACATACCTTTTGTTTAGCGAATGAA  240
B. divergens Spanish sample TCAAAAAACCAAACAAACCAAAACAGTTTGACAAAACATACCTTTTGTTTAGCGAATGAA  240
B. divergens Turkiye        TCAAAAAACCAAACAAACCAAAACAGTTTGACAAAACATACCTTTTGTTTAGCGAATGAA  240
                            ************************************************************

B. divergens R87            TTTTTAGGCACGTCAGTAGTGTAAACCCTAAGTTGAAAATAGAGTGTTCCTACATATAGA  300
B. divergens Spanish sample TTTTTAGGCACGTCAGTAGTGTAAACCCTAATTTGAAAATAGAGTGTTCCTACATATAGA  300
B. divergens Turkiye        TTTTTAGGCACGTCAGTAGTGTAAACCCTAATTTGAAAATAGAGTGTTCCTACATATAGA  300
                            ******************************* ****************************

B. divergens R87            GCACTAATATATATGGGAGGATAATACTGTGGGGTTTCGTGGAAGCCACGCTAGACAATA  360
B. divergens Spanish sample GCACTAATATATATGGGAGGATAATACTGTGGGGTTTCGTGGAAGCCACGCTAGACAATA  360
B. divergens Turkiye        GCACTAATATATATGGGAGGATAATACTGTGGGGTTTCGTGGAAGCCACGCTAGACAATA  360
                            ************************************************************

B. divergens R87            AGTTAGCGAAACGGTTGAAGATATTATGGTCTGAAGCATACTCATGCACATCGTCTCATG  420
B. divergens Spanish sample AGTTAGCGAAACGGTTGAAGATATTATGGTCTGAAGCATACTCATGCACATCGTCTCATG  420
B. divergens Turkiye        AGTTAGCGAAACGGTTGAAGATATTATGGTCTGAAGCATACTCATGCACATCGTCTCATG  420
                            ************************************************************

B. divergens R87            TTACAGCAATTTTAATTAATTCGGTAGCTTGTGAATATGTTGTTATGCTGGCATCATATA  480
B. divergens Spanish sample TTACAGCAATTTTAATTAATTCGGTAGCTTGTGAATATGTTGTTATGCTGGCATCATATA  480
B. divergens Turkiye        TTACAGCAATTTTAATTAATTCGGTAGCTTGTGAATATGTTGTTATGCTGGCATCATATA  480
                            ************************************************************

B. divergens R87            AACTGCTATTATTTATCTATATCTATCTTTC                             513
B. divergens Spanish sample AACTGCTATTATTTATCTATATCTATCTTTCTATATATATATATATATATATATATATAT  540
B. divergens Turkiye        AACTGCTATTATTTATCTATATCTATCTTTC                             516
                            ******************************

B. divergens R87            ---------TATATATAT-GAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT  562
B. divergens Spanish sample ATATATATATATATATAT-GAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT  600
B. divergens Turkiye        ---TATATATATATATAT-GAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT  568
                                           ********************************************

B. divergens R87            ACATTGATGTTATCTACAAAGCAGCTGAGGGTATTGTATCACTGTGGCTTATGGCATGAT  622
B. divergens Spanish sample ACATTGATGTTATCTACAAAGCAGCTGAGGGTATTGTATCACTGTGGCTTATGGCATGAT  660
B. divergens Turkiye        ACATTGATGTTATCTACAAAGCAGCTGAGGGTATTGTATCACTGTGGCTTATGGCATGAT  628
                            ************************************************************

B. divergens R87            CATTCAATTCTATCACAGTGTGGGCGTTTGCTCTAGGTCGAGTTACATGTTCTAATTTT  682
B. divergens Spanish sample CATTCAATTCTATCACAGTGTGGGCGTTTGCTCTAGGTCGAGTTACATGTTCTAATTTT  720
B. divergens Turkiye        CATTCAATTCTATCACAGTGTGGGCGTTTGCTCTAGGTCGAGTTACATGTTCTAATTTT  688
                            **********************************************************

B. divergens R87            ATCTCATCTGAGAACTACAGTCTATATAACTGTGCGGGTATCTTATGTTACCTTGAGTTT  742
B. divergens Spanish sample ATCTCATCTGAGAACTACAGTCTATATAACTGTGCGGGTATCTTATGTTACCTTGAGTTT  780
B. divergens Turkiye        ATCTCATCTGAGAACTACAGTCTATATAACTGTGCGGGTATCTTATGTTACCTTGAGTTT  748
                            ************************************************************

B. divergens R87            AGAATATTTGTACGCTTATCCGTATAATTAACATCGTCGCGTTTAGTGGCTTAAGGCACC  802
B. divergens Spanish sample AGAATATTTGTACGCTTATCCGTATAATTAACATCGTCGCGTTTAGTGGCTTAAGGCACC  840
B. divergens Turkiye        AGAATATTTGTACGCTTATCCGTATAATTAACATCGTCGCGTTTAGTGGCTTAAGGCACC  808
                            ************************************************************

B. divergens R87            TATATCTATGTGTACTTGCGCAAATCAAGTTGATATTTCGAACGTGATAATTGATGAGAT  862
B. divergens Spanish sample TATATCTATGTGTACTTGCGCAAATCAAGTTGATATTTCGAACGTGATAATTGATGAGAT  900
B. divergens Turkiye        TATATCTATGTGTACTTGCGCAAATCAAGTTGATATTTCGAACGTGATAATTGATGAGAT  868
                            ************************************************************

B. divergens R87            AAGCCACATATTCAATGAAAATAATGTAAAATGCAACACACATGTGACTCAATGTAAACT  922
B. divergens Spanish sample AAGCCACATATTCAATGAAAATAATGTAAAATGCAACACACATGTGACTCAATGTAAACT  960
B. divergens Turkiye        AAGCCACATATTCAATGAAAATAATGTAAAATGCAACACACATGTGACTCAATGTAAACT  928
                            ************************************************************

B. divergens R87            ATGCATAAATTAATGGCACATTGTATTGCACATAGGCAAAAATAGTATTAATAACTGACA  982
B. divergens Spanish sample ATGCATAAATTAATGGCACATTGTATTGCACATAGGCAACAATAGTATTAATAACTGACA  1020
B. divergens Turkiye        ATGCATAAATTAATGGCACATTGTATTGCACATAGGCAAAAATAGTATTAATAACTGACA  988
                            ***************************************  *******************

B. divergens R87            TAATTAATGGTTTATTGTCTAATGTGTTGGAACCAAATTATGAGGCATTGTCCATTCCTA  1042
B. divergens Spanish sample TAATTAATGGTTTATTGTCTAATGTGTTGGAACCAAATTATGAGGCATTGTCCATTCCTA  1080
B. divergens Turkiye        TAATTAATGGTTTATTGTCTAATGTGTTGGAACCAAATTATGAGGCATTGTCCATTCCTA  1048
                            ************************************************************

B. divergens R87            GCCGCACCACAATTGTTAGATTGAGAGTGACTACTAATATTGTCATCTGTTTTAAGCACA  1102
B. divergens Spanish sample GCCGCACCACAATTGTTAGATTGAGAGTGACTACTAATATTGTCATCTGTTTTAAGCACA  1140
B. divergens Turkiye        GCCGCACCACAATTGTTAGATTGAGAGTGACTACTAATATTGTCATCTGTTTTAAGCACA  1108
                            ************************************************************

B. divergens R87            CGTATAGCGATTAACTATACGAGTTGTATTTCTTCCGTTATATTCTCTAAACAAAAGGTA  1162
B. divergens Spanish sample CGTATAGCGATTAACTATACGAGTTGTATTTCTTCCGTTATATTCTCTAAACAAAAGGTA  1200
B. divergens Turkiye        CGTATAGCGATTAACTATACGAGTTGTATTTCTTCCGTTATATTCTCTAAACAAAAGGTA  1168
                            ************************************************************

B. divergens R87            TGTTTTGTCAAACTGTTTTGGTTTGTTTGGTTTTTTGATGTTAGTGGTGTGGGTGTCGTG  1222
B. divergens Spanish sample TGTTTTGTCAAACTGTTTTGGTTTGTTTGGTTTTTTGATGTTAGTGGTGTGGGTGTCGTG  1260
B. divergens Turkiye        TGTTTTGTCAAACTGTTTTGGTTTGTTTGGTTTTTTGATGTTAGTGGTGTGGGTGTCGTG  1228
                            ************************************************************

B. divergens R87            ATCTCTACGTTCTGCTGTGTCATTATGAGAAGTGTCGTAGGGGCATAACATTATAAATAT  1282
B. divergens Spanish sample ATCTCTACGTTCTGCTGTGTCATTATGAGAAGTGTCGTAGGGGCATAACATTATAAATAT  1320
B. divergens Turkiye        ATCTCTACGTTCTGCTGTGTCATTATGAGAAGTGTCGTAGGGGCATAACATTATAAATAT  1288
                            ************************************************************

B. divergens R87            ATATTTCCATGTTGTTAATTCGGTTGTGGTGTTGCCCATTGTGTTTCCATTGGCCTCACT  1342
B. divergens Spanish sample ATATTTCCATGTTGTTAATTCGGTTGTGGTGTTGCCCATTGTGTTTCCATTGGCCTCACT  1380
B. divergens Turkiye        ATATTTCCATGTTGTTAATTCGGTTGTGGTGTTGCCCATTGTGTTTCCATTGGCCTCACT  1348
                            ************************************************************

B. divergens R87            AACCCTTTGCAGATTAGTTATTATTAGACCTCATCAAA  1380
B. divergens Spanish sample AACCCTTTGCAGATTAGTTATTATTAGACCTCATCAAA  1418
B. divergens Turkiye        AACCCTTTGCAGATTAGTTATTATTAGACCTCATCAAA  1386
                            **************************************
```

B. *divergens* R87 Nanopore	AACTGCTATTATTTATCTATATCTATCTTTC------------------------------------TATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	562
B. *divergens* R87 Sanger	AACTGCTATTATTTATCTATATCTATCTTTC------------------------------------TATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	562
B. *divergens* Spanish sample Nanopore	AACTGCTATTATTTATCTATATCTATCTTTCTATATATATATATATATATATATATATATATATATATATATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	600
B. *divergens* Spanish sample Sanger	AACTGCTATTATTTATCTATATCTATCTTTCTATATATATATATATATATATATATATATATATATATATATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	600
B. *divergens* Turkiye Nanapore	AACTGCTATTATTTATCTATATCTATCTTTC-----------------------------TATATATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	568
B. *divergens* Turkiye NGS	AACTGCTATTATTTATCTATATCTATCTTTC-----------------------------TATATATATATATATGAAATTATCAGCTTATACCGGAATTATGTATTAGAAATAT	568