Inhibiting the right dorsolateral prefrontal cortex selectively enhances unsupervised statistical learning

Orsolya Pesthy*[1,2], Zsuzsanna Viktória Pesthy*[2,3,4], Teodóra Vékony[5], Dániel Fabó[6,7,8], Dezso Nemeth[1,6,10]

1 Centre de Recherche en Neurosciences de Lyon, INSERM, CNRS, Université Claude Bernard Lyon 1, CRNL U1028 UMR5292, Bron, France

2 Brain, Memory and Language Research Group, Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary

3 Doctoral School of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary

4 Institute of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary

5 Gran Canaria Cognitive Research Center, Atlántico Medio University, Las Palmas de Gran Canaria, Spain

6 Flash Clinic, Budapest, Hungary

7 Department of Neurology, University of Szeged, Szeged, Hungary

8 Department of Behavioral Sciences, Semmelweis University, Budapest, Hungary

9 Dept of Voice, Speech and Swallowing, Semmelweis University, Budapest, Hungary

10 BML-NAP Research Group, Institute of Psychology, Eötvös Loránd University & Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary

*These authors contributed equally to this article

Abstract

The brain must balance the automatic extraction of environmental regularities with top-down cognitive control, yet the causal neural mechanisms governing this interplay are debated. In particular, the hemispheric contributions of the dorsolateral prefrontal cortex (DLPFC) remain unresolved. Here, we applied inhibitory repetitive transcranial magnetic stimulation (rTMS) to the left, right, or bilateral DLPFC in 95 healthy adults during a probabilistic sequence learning task. We found that inhibiting the right and bilateral DLPFC significantly enhanced statistical learning compared to sham and left DLPFC stimulation. Contrary to the hypothesis that this competition is mediated by the suppression of existing knowledge, episodic memory performance was unaffected across all stimulation groups. These findings challenge a direct episodic gating mechanism and suggest the DLPFC's influence on statistical learning is hemisphere-dependent. As an alternative, we propose that right DLPFC inhibition shifts cognitive processing toward a more exploratory information-sampling strategy, a view supported by our finding of significantly greater reaction time variability in the right and bilateral stimulation groups. Together, our results provide causal evidence for a right-lateralized suppressive influence of the DLPFC on unsupervised learning and suggest its modulatory role is linked to information processing style rather than direct competition with episodic memory systems.

Keywords: statistical learning, unsupervised learning, episodic retrieval, dorsolateral prefrontal cortex, competition theory, transcranial magnetic stimulation

Introduction

The brain resembles an ecosystem, wherein multiple cognitive processes interact and compete for influence over behaviour[1,2]. Adaptation to novel environments requires neural plasticity, long attributed to supervised or reinforcement learning, optimized by model-based processes. However, mounting evidence suggests that neural plasticity can also emerge through passive, unrewarded exposure to structured input, highlighting a fundamental role for unsupervised learning, which is in turn optimized by model-free processes[3,4]. Building on the hypothesis that unsupervised and supervised mechanisms shape representations jointly, we examine a critical instance of this process in humans: statistical learning (the acquisition of probability-based regularities without feedback or reward), and its balance with model-based memory retrieval. According to the competition hypothesis, model-free (such as statistical learning) and model-based (such as executive functions or episodic retrieval) processes rely on overlapping neural resources and may therefore inhibit each other, with the dorsolateral prefrontal cortex (DLPFC) serving as a key hub in controlling their balance[1,5–9]. One proposed mechanism is that the DLPFC acts as a gate to long-term memory □□ thus, its inhibition hinders retrieval while simultaneously supporting the encoding of novel statistical structure[9]. To test this hypothesis, we applied inhibitory repetitive transcranial magnetic stimulation (rTMS) to the left, right, or bilateral DLPFC in healthy participants, assessing its impact on both statistical learning and episodic memory retrieval.

The antagonistic relationship between DLPFC activity and the model-free process of statistical learning has been well-documented. Specifically, reduced DLPFC engagement has been consistently associated with enhanced statistical learning performance. This has been observed in various groups with reduced DLPFC activity, such as children (compared to adults)[10–12], individuals under hypnosis[13] or depletion[2,14,15]. Importantly, studies using inhibitory non-invasive brain stimulation on the bilateral DLPFC have shown better learning[9], and retrieval[16] of probabilistic information in the DLPFC-inhibited group compared to the controls. These findings together suggest an inverse relationship between DLPFC activity and statistical learning, demonstrated through both correlational and causal evidence (although see ref[17]) □□ however, this alone does not clarify which mechanism drives the competition.

The hemisphere-dependent involvement of the DLPFC in statistical learning may offer valuable insight into its underlying mechanisms. Neuropsychological studies have shown a right hemisphere dominance in statistical learning, highlighting that statistical learning remains intact when solely the right hemisphere remains available in split-brain and unilaterally lesioned patients, whereas the left hemisphere supports systematic hypothesis-testing - essentially, model-based functioning[18,19]. This asymmetry seems to extend to the role of the DLPFC, where improved statistical learning has been observed following inhibition of the left or facilitation of the right DLPFC[14,15,17]. However, findings remain inconsistent, with some studies reporting the opposite pattern[20–22] or null results[23,24]. A proposed mechanistic explanation is rooted in hemispheric differences in decision strategies: the right hemisphere favors selecting the most probable option each time (frequency-maximizing strategy), while the left hemisphere engages in choosing each option in proportion to its occurrence frequency

(frequency-matching strategy)[19]. While the latter supports causal inference, statistical learning, in which frequency is the key information source, may benefit more from frequency-maximizing strategies[4]. Although this account is compelling, the lateralization of statistical learning remains unresolved due to inconsistencies in the literature. To move beyond descriptive accounts and clarify the role of DLPFC asymmetry, we applied repetitive transcranial magnetic stimulation (rTMS) over the left, right, and bilateral DLPFC, and examined its effects on both statistical learning and episodic memory retrieval.

In the domain of episodic retrieval, most findings align with the competition hypothesis and the DLPFC's proposed gatekeeping role in regulating access to long-term memory. A positive relationship between DLPFC functioning and episodic retrieval performance has been shown, although null results are common as well (for a review, see ref[25]). This pattern suggests that the DLPFC may modulate memory access, possibly through a working memory involvement[26]. However, it remains unclear whether downregulating episodic retrieval facilitates the encoding of new statistical regularities[9]. To directly test this, we included an episodic memory task in our study, allowing us to examine whether DLPFC modulation affects retrieval performance and whether this, in turn, relates to statistical learning outcomes. Hemispheric asymmetries offer further insight. According to the classic Hemispheric Encoding/Retrieval Asymmetry (HERA) model[27,28], the left prefrontal cortex plays a greater role in encoding, while the right is more engaged during retrieval. Additional factors, such as stimulus modality (verbal stimuli are more left-lateralized, visual more right-lateralized), emotional valence (with mixed findings), and familiarity (with the DLPFC particularly engaged in the recall of novel stimuli), further shape lateralization effects[25]. To control for these variables, our memory task used novel, emotionally neutral stimuli incorporating both verbal and visual elements. Crucially, if increased statistical learning is observed in tandem with impaired episodic retrieval □□ especially following right DLPFC stimulation □□ this would lend support to the hypothesis that the DLPFC acts as a cognitive gate to long-term memory. When this gate is weakened, the system may become more receptive to encoding novel statistical patterns.

We aimed to address open questions about the effects of DLPFC inhibition on statistical learning and episodic memory. We applied 1 Hz (inhibitory) rTMS over the left, right, or bilateral Brodman 9 area to target the DLPFC. In this sham-controlled study, stimulation was on in the breaks of task sessions when participants performed on a statistical learning and an episodic retrieval task in a randomized order. Furthermore, we re-tested the participants after a 24h delay. We had the following expectations regarding the intervention effect:

1) In this study, we go beyond the findings of Ambrus et al.[9] by addressing a key open question: the hemispheric asymmetry in statistical learning. Specifically, we aim to identify which hemisphere plays a more critical role in this process. If 1 Hz inhibitory rTMS over the left hemisphere enhances statistical learning performance, it would support right-hemisphere dominance in statistical learning, emphasizing the role of different cognitive strategies as a background mechanism[19]. Conversely, if altered excitability of the right

dorsolateral prefrontal cortex leads to improved statistical learning, it would raise the possibility of a right-hemisphere gating mechanism, in line with the classic HERA model.

2) Our study was designed to test competing hypotheses about the DLPFC's gatekeeping function—its role in controlling access to the long-term memory systems that store prior knowledge and internal models. Based on this hypothesis, we predict enhanced statistical learning will be accompanied by impaired episodic retrieval in the stimulation groups. If, however, DLPFC inhibition affects statistical learning independently of episodic memory, it would suggest that the DLPFC's influence on statistical learning is mechanistically distinct from its role in accessing the model-based priors required for episodic retrieval.

Methods

Participants

We recruited 104 healthy adults to our experiment with normal or corrected-to-normal vision. They were recruited through a university course and received course credits for their participation. Inclusion criteria were no history of neurological or psychiatric diagnoses and no contraindication for non-invasive brain stimulation (epilepsy in the participant or their first-degree relatives, metal implants in the head or neck, pacemaker, migraine, history of severe head injury or concussion). All of them provided written informed consent. We excluded eight participants due to technical issues, and one participant who did not return for the second session. Thus, in the analyses, we entered the data of 95 participants. The experiment was conducted in accordance with the Declaration of Helsinki, with ethical approval granted by the Ethics Committee of Eötvös Loránd University (approval number: 2019/282).

As detailed in the Procedure section below, participants were randomly assigned to one of four experimental groups: (1) active right DLPFC stimulation, (2) active left DLPFC stimulation, (3) active bilateral DLPFC stimulation, or (4) sham stimulation (control group). Demographic information and complex working memory (counting span, CSPAN) of the participants are shown in Table 1. There were no significant differences between the groups in terms of gender, age, or CSPAN performance.

		Group			Group comparison		
	Sham	Right DLPFC	Left DLPFC	Bilateral DLPFC	Statistics (χ^2/H/F)	p	BF_{10}
Gender (f/m)	21/5	18/4	18/5	18/6	0.939	0.942	0.020

Age (years)	21.85 (SD = 3.77)	21.73 (SD = 5.74)	21 (SD = 1.81)	22.48 (SD = 4.13)	1.636	0.651	0.101
CSPAN	3.74 (SD = 0.97)	3.71 (SD = 0.84)	3.52 (SD = 0.83)	3.86 (SD = 0.8)	0.62	0.606	0.112

Table 1. Demographic characteristics and counting span (CSPAN) of the experimental groups.

Tasks

Alternating Serial Reaction Time Task

To measure statistical learning, we employed the Alternating Serial Reaction Time (ASRT) task[29]. In this task, four black circles were displayed horizontally on a white background of the computer screen, with dog head stimuli appearing one at a time inside these circles (see Figure 1, panel A). Each circle corresponded to a specific response key (Y, C, B, M) on a Hungarian QWERTZ keyboard layout. Participants were instructed to press the appropriate key as quickly and accurately as possible after the stimulus appeared, using their index and middle fingers of both hands. The stimulus remained visible until the correct key was pressed, after which the next stimulus appeared following a 120 ms response-to-stimulus interval.

The task was organized into ~1-minute-long blocks. Each block began with five random stimuli (for practice purposes), which were excluded from later analyses. Following these, an eight-element sequence was repeated ten times, resulting in a total of 85 trials per block. Within this repeating sequence, every first element was a random stimulus, while the alternating elements followed a fixed, predetermined sequence (e.g., 2–r–4–r–3–r–1–r, where numbers denoted the fixed pattern elements, and "r" represented a randomly selected position from the four locations). Importantly, participants were unaware of the embedded sequential regularity. The alternating pattern of fixed and random stimuli generated triplets - runs of three elements formed by a fixed structure of stimulus transitions - in a sliding window manner. Based on the example above, triplets such as 2_X_4, 4_X_3, 3_X_1, and 1_X_2 in the above example (where "X" denoted the middle stimulus, regardless of whether it was a random or fixed pattern element) occurred with high probability, as they can be formed both by two sequence elements comprising a random one or two random ones comprising a sequence element, thus, their last elements could be predicted either from the fixed pattern or from the combination of pattern and random elements. In contrast, triplets like 1_X_3 or 4_X_1 were considered low-probability, as they could only occur when both the first and last elements were random. In total, there were 16 possible high-probability triplets and 48 low-probability triplets.

As a result, the last elements of high-probability triplets were more predictable, and thus responses to them tended to be faster and more accurate. Each high-probability triplet

appeared approximately five times more frequently than each low-probability one. Specifically, high-probability triplets occurred with a ~4% likelihood per trial and 62.5% (16 × 4%) per block, while low-probability triplets appeared with ~0.8% per trial and 37.5% (48 × 0.8%) per block (see Figure 1). The difference in reaction time (RT) and/or accuracy between the last elements of high- and low-probability triplets was taken as a behavioral index of statistical learning. For analytic purposes, data were segmented into epochs, each consisting of 5 blocks, that is, a total of 85 × 5 trials.

Figure 1. Structure of the Alternating Serial Reaction Time Task (ASRT) A. Participants were instructed to press the button corresponding to the location of the stimulus (a dog's head). The order of stimuli followed a hidden alternating pattern: every first element was randomly selected (blue), appearing in any of the four circles, and every second element followed a fixed sequence (green), referred to as the pattern element. This resulted in an eight-element alternating sequence that was repeated 10 times within each block. B. Due to this alternating structure, certain three-element combinations of stimuli—called triplets—occurred with different probabilities. High-probability triplets (orange) can be formed when the third stimulus follows the hidden pattern, and may start with either a random or a pattern element. These make up 62.5% of all triplet occurrences. In contrast, low-probability triplets (purple) can only occur in random-pattern-random configurations and thus represent 37.5% of the total.

Paired Associate Learning Task

The Paired Associate Learning Task (PALT), developed for the Hungarian language by[30], is a computerized paradigm designed to assess declarative memory and is based on the revised version of a task by Cohn & Moscovitch[31]. During the task, two drawings – depicting either objects or animals – were simultaneously presented on the screen. The task consisted of several phases. In the initial learning phase, participants were asked to name each pair of images aloud as they appeared sequentially, without any instruction to memorize them. After naming the pictures, the experimenter advanced the display to the next image pair. A total of 23 image pairs were shown during this phase.

Subsequent retrieval phases involved the presentation of image pairs again, and participants were asked to indicate whether they had previously seen each item during the learning phase. Four possible types of image pairs were presented: old-old (both previously seen), new-new (none of the images were previously seen), old-new, and new-old. In cases where participants responded that both items had been presented before, participants were further asked to determine whether the pair had originally appeared together ("original" order) in the learning phase or had been recombined ("rearranged" order). Although the images did not repeat within the retrieval phase, participants were specifically instructed to give their responses solely based on the learning phase. Responses were recorded by the experimenter to avoid confusion due to the complexity of response options, and therefore, reaction time was not measured.

Unlike the original implementation by Nagy et al.[30], in our study, participants were not informed in the learning phase that they would later be tested on the retrieval of the images following the naming task. As such, they did not expect a memory test, allowing for an incidental learning design. Additionally, in contrast to the original continuous task format, we divided the task into discrete retrieval blocks: three blocks during the first session and one block during the second session. Each block included eight unique image pairs, with no repetition across blocks to prevent interference with consolidation processes.

Stimulation protocol

rTMS was administered using a Deymed DuoMAG XT stimulator over the DLPFC, targeting Brodmann area 9, corresponding to the F3 and F4 positions of the international 10–20 EEG system, which we located using a standard EEG cap (which has been shown to be a reasonably reliable method of neuronavigation[32]). To establish the motor threshold, single TMS pulses were delivered over the participant's right primary motor cortex (M1) as localized by the C4 electrode of a 10-20 system EEG cap, on gradually increasing intensity, while recording electromyographic activity from the left hand muscles. The motor threshold was defined as the minimum stimulation intensity that elicited visible muscle twitches in the hand or evoked motor potentials (MEPs) with an amplitude of at least 100 μV.

Stimulation during the experimental session was applied at 100% of each participant's individual resting motor threshold in the active stimulation groups. A 1 Hz inhibitory stimulation protocol was used, delivering 300 pulses in each stimulation session over 5 minutes to either the right or left hemisphere in the active right and active left groups, respectively. In the active bilateral group, stimulation was applied consecutively to each hemisphere for 2.5-2.5 minutes. In the sham condition, the stimulation coil was held perpendicular to the scalp (following the configuration described by [9]), preventing effective stimulation. However, the coil was still positioned on the participant's head, allowing them to believe they were receiving stimulation. This approach was used to control for potential placebo effects.

Procedure

Participants completed an online screening questionnaire designed to exclude individuals with health-related conditions (e.g., neurological or psychiatric disorders) that

would preclude participation. Only those meeting the inclusion criteria were enrolled in the study, which consisted of two experimental sessions held 24 hours apart. Prior to the first session, participants were randomly assigned to one of the groups (active left, right, bilateral DLPFC stimulation, or sham stimulation) and a counterbalanced task order, determining whether they would first complete the ASRT or the PALT task. This assignment remained fixed throughout the entire experiment for each participant.

After providing written informed consent, the location of Brodmann area 9 and the M1 were identified using a 10-20 system EEG cap, followed by the assessment of each participant's resting motor threshold. Participants then completed a familiarization phase of the ASRT task (two blocks of stimuli without embedded sequences) and the learning phase of the PALT task, with task order determined by prior randomization. Participants received a 5-minute rTMS session targeting either the right DLPFC, left DLPFC, or sham stimulation. In the bilateral stimulation group, 2.5 minutes of stimulation was delivered to each hemisphere consecutively - the order of the hemispheres was also randomized across participants. Immediately thereafter, participants completed the first epoch of the sequence-containing ASRT task (comprising five blocks, approximately five minutes in total) and the first three retrieval phases of the PALT task in a randomized order. This rTMS–ASRT–PALT cycle was repeated five times, with the order of the ASRT and PALT tasks alternating based on the assigned condition. During the second and fourth cycles, the PALT task was replaced with short breaks. In the second session, held 24 hours later, participants again completed one epoch of the ASRT task and a retrieval phase of the PALT tasks in the same randomized order as in the first session. The experimental procedure is illustrated in Figure 2. To ensure that baseline cognitive differences between groups did not confound the results (e.g., one group performing better due to inherently higher cognitive capacity), we administered the CSPAN [33], measuring complex working memory tasks to detect any such pre-existing group-level advantages (see Table 1).

Figure 2. Study design. The experiment consisted of two sessions separated by a 24-hour delay. At the beginning of the first session, participants were randomly assigned to one of four stimulation groups (sham, right DLPFC, left DLPFC, bilateral DLPFC). Task order was also randomized: each participant was assigned to begin the session with either the ASRT task (represented by a dog's head icon) or the PALT task, and this order was kept consistent throughout all subsequent blocks. Following this, participants completed a familiarization

phase of the ASRT task (which was two blocks of randomized trials, following no sequence) and the learning phase of the PALT task. These were followed by the first rTMS stimulation (1 Hz, 5 minutes, at 100% of the participant's individual motor threshold). Stimulation was applied to the assigned site: for bilateral stimulation, 2.5 minutes were applied to each hemisphere. After the first stimulation, five blocks of task performance alternated with five stimulation sessions. Notably, only the ASRT task was administered following the second and fourth stimulation blocks. In total, participants completed five task–stimulation cycles during the first session. After a 24-hour delay, participants returned for the second session, during which they completed a sixth epoch of the ASRT task and the retrieval phase of the PALT task.

Statistical analysis

All statistical analyses were conducted in RStudio (Version 2024.12.0) or JASP (Version 0.19.3)[34,35]. Linear mixed-effects models were fitted using the *mixed* function from the *afex* package[36]. We used the default method (Kenward-Roger) to approximate degrees of freedom for fixed effects in all models. Post-hoc contrasts were computed using the *emmeans* package[37], using Tukey correction for multiple comparisons. All figures and data preprocessing were conducted in Python (Version 3.13) using the *pandas*[38], *numpy*[39], *os*, *io*[40], *seaborn*[41], and *matplotlib*[42] libraries. In all analyses, we determined an alpha level of 0.05.

Trials were categorized into high- and low-probability in a sliding window manner based on whether they were the last element of a high- or a low-probability triplet (henceforth simply referred to as high- and low-probability triplets). We excluded the first five trials of each block as they were random and served as practice trials, and trials that were the last element of repetitions (e.g., 1-1-1, 2-2-2) or trills (e.g., 1-2-1, 2-3-2), as participants may show a pre-existing tendency of faster motor responses to such occurrences[43]. Furthermore, from the analyses of RT, inaccurate trials were excluded.

To investigate group differences in statistical learning on RTs, we conducted a linear mixed-effect model to analyze RTs as a continuous outcome variable and a generalized linear mixed model to analyze accuracy, with responses coded as 0 (incorrect) and 1 (correct). The fixed effects included Epoch (1–6), Triplet Type (high- vs. low-probability), Group (sham, right, left, and bilateral stimulation groups), and their interactions. The models allowed for participant-specific intercepts and correlated slopes for within-subject variables. We used a top-down approach in model selection, starting from the most complex model (allowing for participant-specific intercept and correlated slopes for the interaction of within-subject variables) and simplified it gradually until we got a converging model. The fixed effects of the generalized linear mixed model were tested using Type III Wald chi-square tests with the *anova()* function from the car package, and results are reported in Table X. In addition, full model coefficients, including all interaction terms, are reported using the *tab_model()* function from the *sjPlot* package (for both the reaction time and accuracy models)[44].

To compare the groups in episodic retrieval performance, we conducted a one-way analysis of variance (ANOVA), as all linear mixed-effect models failed to converge on these

data. The task yields three outcome measures. Three learning indices were derived from participants' responses. Item memory was calculated by subtracting the false alarm rate (incorrect Old–Old responses to New–New pairs) from the hit rate for rearranged Old–Old pairs. The association learning index was defined as the difference between the hit rate for original and rearranged Old–Old pairs. The recollection index was computed by subtracting the false alarm rate (incorrect original responses to rearranged Old–Old pairs) from the hit rate for original Old–Old pairs. Prior to the analysis, we tested whether the data were normally distributed within each group using the Shapiro–Wilk test. In cases where the assumption of normality was violated, we applied the non-parametric Kruskal–Wallis test instead of ANOVA. If normality was not violated, we proceeded to test the assumption of homogeneity of variances using Levene's test. If this assumption was violated, we applied the Welch correction to the ANOVA to account for unequal variances. To test null effects, we applied Bayesian ANOVAs on the PALT scores. The Bayes Factor $_{10}$ (BF_{10}) was calculated using JASP to compare the null and alternative hypotheses. BF_{10} expresses how much more likely the observed data are under the alternative hypothesis (H_1) than under the null hypothesis (H_0). Following the interpretation guidelines by Wagenmakers et al.[45], values around 1 indicate no substantial evidence for either hypothesis. BF_{10} values between 1 and 3 indicate anecdotal evidence for H_1, while values between 3 and 10 suggest moderate evidence. Conversely, values between 0.33 and 1 indicate anecdotal evidence for H_0, and values between 0.1 and 0.33 suggest moderate support for H_0. A BF_{10} below 0.1 is considered strong evidence in favor of the null hypothesis, indicating that the groups likely do not differ.

Lastly, we performed some exploratory analysis that may help us understand our results more. First, as an indicator of exploratory behaviour in the task, we tested the RT variability on the ASRT task by conducting a Levene's test, using Group as the independent variable. Post hoc analyses were corrected for multiple comparisons using the Holm method. In addition, as working memory is a relevant factor in the DLPFC-statistical learning relationship[21], we investigated whether working memory capacity is related to statistical learning performance, we ran a correlation between total learning score (calculated as the difference between high- and low-probability triplets across all sequential epochs) and baseline counting span scores, using Pearson's correlation. Please note that these analyses were exploratory and not part of our predetermined analysis plan, and they serve to further explain the results of the main analyses; for justification, see the Discussion section.

Results

Is there a difference between the groups in statistical learning? Reaction time.

The linear mixed-effect model on RT revealed a significant main effect of Epoch, indicating that participants became faster throughout the task, regardless of triplet types (F(5, 183988.17) = 728.44, $p < .001$). This general improvement differed between the groups, as reflected by a significant Group × Epoch interaction (F(15, 183988.17) = 4.02, $p < .001$). The post hoc comparisons showed that the sham and the right group improved in RT in all epochs, except between the 4th and 5th epochs (p < 0.027 in all consecutive epoch pairs

besides 4-5th, where p = 0.539), where they stagnated. The left group showed a similar pattern, except that they did not improve between the 3rd and 4th epochs (p = 0.795), but did so between all others (p < 0.001 in all other consecutive epoch pairs). The bilateral group improved during the whole task (p < 0.044). A significant main effect of Triplet Type was also observed (F(1, 183988.40) = 486.98, p < .001), with faster responses to high-probability triplets than to low-probability ones indicating statistical learning. The difference between the Triplet Types changed over time, shown by the Epoch × Triplet Type interaction (F(5, 183988.32) = 17.54, p < .001), signifying learning improved over time. There was no significant Group main effect, indicating that the groups showed similar overall RT if we disregard triplet types (F(3, 91.13) = 0.88, p = .453). Most importantly, we found a significant Group × Triplet Type interaction (F(3, 183988.40) = 3.71, p = .011), showing that the magnitude of the statistical learning effect (i.e., the RT difference between high- and low-probability triplets) varied across groups. The right stimulation group demonstrated significantly greater statistical learning than the sham group (p = .005, SE = 1.14) and the left

stimulation group (p = .032, SE = 1.17). Additionally, the bilateral stimulation group outperformed the sham group (p = .015, SE = 1.12). A trend-level difference was also observed between the bilateral and left stimulation groups, with the bilateral group showing better learning (p = .079, SE = 1.15). However, this difference between the groups in statistical learning did not change over time, as the Epoch × Triplet Type × Group interaction was not significant (F(15, 183988.32) = 0.53, p = .927; see Figure 3). The model's explanatory power was modest (Marginal R^2 = 0.029, Conditional R^2 = 0.171). The analyses of the accuracy data are reported in the Supplementary Materials.

Figure 3. Reaction times to high- and low-probability triplets in sham, right, left, and bilateral stimulation groups. The figure displays reaction times (in milliseconds) for high-probability (purple) and low-probability (orange) triplets across six epochs (x-axis). The panels (arranged from top right to bottom left) correspond to the sham, right, left, and bilateral DLPFC stimulation groups. The difference in reaction times between high- and low-probability triplets indicates statistical learning. The dashed grey line between the 5th and 6th epochs marks the 24-hour interval between the two experimental sessions. Error bars show the standard error of the mean.

Do the groups differ in episodic retrieval?

Due to violations of the normality assumption for all three scores, we conducted Kruskal-Wallis tests to examine group differences. No significant differences were found between the groups in item memory ($H(3) = 3.96$, $p = 0.265$, $Rank\ \eta^2 = 0.003$, $BF_{10} = 0.064$), or association learning index ($H(3) = 4.56$, $p = 0.207$, $Rank\ \eta^2 = 0.004$, $BF_{10} = 0.065$) and recollection index ($H(3) = 2.19$, $p = 0.533$, $Rank\ \eta^2 = 0.000$, $BF_{10} = 0.024$; see Figure 4).

Figure 4. Results of Kruskal-Wallis tests for group differences in PALT. Bar plots showing mean performance (± standard error) on three outcome measures of the Paired Association Learning Task (PALT) across the four stimulation groups (sham, left, right, bilateral). The three panels represent item memory (left), association learning index (middle), and recollection index (right).

Exploratory analyses: individual differences

Does the variance of reaction times differ due to the stimulation?

To test whether participants in the stimulation groups showed a different variability of reaction times, we performed a Levene's test with group (sham/right/left/bilateral DLPFC stimulation) as the independent variable. Levene's test indicated significant differences in variance across groups [$F(3, 91) = 270.65$, $p < .001$]. The right and the bilateral DLPFC stimulated groups showed significantly higher variances compared to the sham and the left DLPFC stimulated groups ($p_{Holm} < 0.001$ in each case). For easier interpretability, we report the standard deviations (rather than the variance) in Table 2.

	Triplet type		
group	**high**	**low**	**total**
sham	80.47	79.25	80.21
right	90.15	92.91	91.05
left	76.29	77.09	76.61
bilateral	92.40	90.94	92.10

Table 2. RT standard deviations of the groups, by triplet type.

Does working memory correlate differently with statistical learning across groups?

We tested the correlation between statistical learning (defined as the RT difference of high- and low-probability triplets throughout the whole task), separately for each group. We found no significant correlation in any of the groups, although the bilateral DLPFC stimulated group showed a trend-level positive correlation [sham: $r(24) = -0.068$, $p = 0.741$; right: $r(20) = -0.223$, $p = 0.296$; left: $r(21) = -0.089$, $p = 0.686$; bilateral: $r(22) = 0.369$, $p =$

0.076], see Figure 5.

Figure 5. Statistical learning as a function of baseline working memory capacity. The y-axis represents the RT difference of low- minus high-probability triplets (calculated for the sake of this analysis alone). The x-axis represents the Counting span (CSPAN) performance. Dots indicate individual data, lines represent a linear regression trend line, and bands show the 95% confidence interval of the regression estimate.

Discussion

We aimed to causally assess the role of the dorsolateral prefrontal cortex (DLPFC) in statistical learning—an unsupervised form of learning critical for extracting regularities from sensory input, acquiring new skills, and constructing predictive representations—by applying inhibitory transcranial magnetic stimulation (rTMS) over the left, right, or bilateral DLPFC. We found that inhibition of the right and bilateral DLPFC significantly enhanced statistical learning performance, as indicated by faster responses to predictable compared to unpredictable stimuli. To examine the functional specificity of this effect, we included an episodic memory task as a cognitive control measure. Episodic memory retrieval remained unaffected across all stimulation groups, suggesting that the observed enhancement in statistical learning cannot be attributed to a general facilitation or disruption of cognitive performance. Together, these findings demonstrate a right-lateralized influence of the DLPFC on statistical learning, suggesting that this effect may operate through mechanisms distinct from episodic retrieval as measured here.

Our findings in the bilateral stimulation group are in line with the only prior rTMS study on the acquisition phase of statistical learning using bilateral DLPFC inhibition, which similarly reported enhanced learning of probabilistic regularities[9]. One additional study has applied bilateral inhibitory stimulation during the retrieval phase and found improved retrieval of statistical information[16]. These findings provide converging causal evidence for the involvement of the DLPFC in statistical learning, contributing to a growing literature suggesting an inverse relationship between DLPFC-mediated functions □□ such as executive control □□ and statistical learning performance[7,8,15,46,47]. This inverse relationship supports the competition theory, which posits that cognitive control processes □□ although beneficial for goal-directed, model-based processes □□ interfere with the automatic, model-free mechanisms underpinning statistical learning. Evidence from functional connectivity and developmental studies further supports this interpretation: reduced functional coupling between the DLPFC and other brain regions has been associated with improved statistical learning[48,49]; and the learning advantage observed in children may stem from the immaturity of the DLPFC[10–12,50]. Collectively, these findings suggest that reduced prefrontal engagement, whether through development or neuromodulation, can facilitate statistical learning by releasing the system from top-down constraints.

We sought to investigate whether the apparent competition between DLPFC functioning and statistical learning could be explained by the DLPFC's hypothesised control over the access to long-term memory (i.e., its gating role) [9]. Working memory is integral to accessing long-term memory and is often conceptualized as a state of activated long-term memory[51]. Since our 1 Hz rTMS stimulation targeted a major working memory hub[52,53], it is reasonable to assume that we also influenced access to long-term memory. Based on prior theoretical proposals[9], we hypothesized that disrupting DLPFC activity may impair access to pre-existing internal models stored in long-term memory, thereby facilitating the acquisition of novel statistical regularities. While our findings on statistical learning support this hypothesis, we did not reveal a disruption of episodic retrieval following DLPFC inhibition. Thus, we did not find evidence that this effect was mediated by episodic memory

interference. Alternative gat pathways remain plausible, or gating may happen through different memory systems, like semantic or working memory. Overall, while our results do not provide support for a gating mechanism based on episodic memory interference, they leave open the possibility that the DLPFC plays a modulatory role through alternative routes - a hypothesis that warrants further investigation.

Our hemispheric findings offer a more specific interpretation. A compelling explanation for our results lies in the modality-specific lateralization of prefrontal cortex function. Our use of a visuomotor statistical learning task aligns perfectly with findings that right DLPFC disruption primarily affects visual learning[21,22]. This framework also helps reconcile seemingly conflicting results in the literature, which have often reported enhanced *verbal* statistical learning following *left* DLPFC disruption[15,46]. This right-lateralized effect is also broadly consistent with the Hemispheric Encoding/Retrieval Asymmetry (HERA) model[27,28]. This model posits a primary role for the right DLPFC in memory retrieval. From this perspective, inhibiting the right DLPFC may have impaired access to pre-existing models, shifting the cognitive system toward more data-driven processing and thereby facilitating statistical learning. The results on the right hemisphere-specific effect is consistent with the idea that our episodic memory task lacked the sensitivity to detect subtle group differences (see limitations), and that statistical learning was nonetheless improved via an episodic gating mechanism[9] □□ future studies are warranted to test this speculation. Taken together, our results provide causal evidence that right DLPFC inhibition enhances visuomotor statistical learning, likely by attenuating top-down, memory-based constraints. Future studies using systematic, modality-sensitive designs are needed to further elaborate this relationship.

To further interpret our findings, it is essential to consider which cognitive strategies and information processing styles are known to support statistical learning. While frequency-maximizing and frequency-matching strategies were discussed in the introduction[4,19], they are unlikely to explain our results, as such strategies would predict enhanced learning following left DLPFC inhibition, which we did not observe. An alternative potential mechanism draws from reinforcement learning research: trying new, undiscovered patterns (that is, exploration as opposed to exploitation) and a broader, more distributed attention to various environmental signals (that is, broader information sampling) can facilitate the early acquisition of probabilistic regularities[54,55], as frequent re-sampling could lead to a deeper knowledge of statistical structures that are inherently probabilistic (for a discussion, see ref[56]). Although our study was not specifically designed to assess differences in exploration/exploitation or their causal relationship with statistical learning, we propose that right and bilateral DLPFC inhibition fostered a shift toward a more exploratory information-sampling strategy. This interpretation is supported by our finding of significantly greater RT variability □□ a well-established behavioural proxy for exploration □□ in the same groups that showed enhanced learning. We speculate that the observed benefits after right and bilateral DLPFC inhibition in statistical learning may reflect a shift toward a broader, more exploratory information processing style.

The observed shift toward broader, more exploratory information sampling may be caused by the disruption of working memory: lower DLPFC functioning and thus lower working memory have been linked to such a strategy[60,61]. Individuals with reduced working memory capacity (such as children or participants undergoing DLPFC inhibition in our study) tend to re-sample their environment more frequently, as previously acquired information is more susceptible to decay[60]. This explanation is compelling because our stimulation target, the Brodmann 9 area, is known for its role in working memory[53,62], and it is consistent with extensive evidence that working memory exerts a modulatory influence on statistical learning[7,8,21]. This framework may also reconcile the above-mentioned conflicting findings regarding the lateralization of statistical learning. Inhibiting verbal working memory, which shows a well-established left-lateralization[63–66], may result in a broader information sampling strategy in the verbal domain, benefiting verbal statistical learning tasks[15,46]. But in our visuomotor statistical learning task, verbal working memory (and thus, broader verbal information sampling) could have little effect, hence the null result in the left stimulation group. On the other hand, right stimulation may have been effective because it influenced visual working memory (known to be right-lateralized[62–66], thus shifting visual processing to be more explorative. While our study was not designed to directly test this hypothesis, exploratory analyses offer tentative support. Specifically, we observed a trend-level positive correlation between working memory capacity and statistical learning in the bilateral group, suggesting that DLPFC inhibition may enhance statistical learning, particularly in individuals with higher baseline working memory capacity □□ showing the baseline working memory's relation to the stimulation effectiveness (see Figure 5). Taken together, we speculate that DLPFC inhibition may reduce working memory maintenance of prior samples, promoting increased environmental exploration, which facilitates statistical learning of probabilistic regularities □□ future studies should be designed specifically to probe this mechanism.

A key limitation of our study is our choice of task for episodic memory retrieval. Although it is not without example to find no DLPFC-related effect on episodic memory, the majority of studies were able to show a relation to the DLPFC[25]. This may have multiple explanations. As we could see earlier, DLPFC lateralization greatly depends on the modality of the task in many cognitive functions, and episodic memory is no exception[27,28,67]. Our task had both visual and verbal elements, enabling participants to use their preferred strategy, either verbal or visual. Thus, this task was a suboptimal choice for a study that wanted to explore lateral effects. However, compensation from the non-inhibited strategy (verbal in the case of left inhibition, visual in the case of right inhibition) does not account for our results in the bilateral group: when both hemispheres are inhibited, we should have seen the effect if it were there. Furthermore, the PALT measures a specific form of associative memory. It is therefore possible that the DLPFC's gating function does not act through this declarative retrieval but rather through other forms of memory that are more integral to the online processing of statistical information. For instance, DLPFC inhibition may have disrupted acces to semantic representations or the short-term maintenance of statistical priors within working memory itself □□ cognitive processes not captured by our episodic task. Future

studies should systematically explore the effect of modality, and test various memory systems in relation to statistical learning to clarify the precise nature of the DLPFC's modulatory role.

In summary, our study provides novel causal evidence for a right-lateralized role of the DLPFC in modulating statistical learning. These findings offer support for the competition theory by demonstrating that DLPFC-mediated functions, particularly those localized to the right hemisphere, can suppress model-free learning processes. Notably, our results do not support the interpretation that this effect arises from reduced interference by episodic memory systems. Instead, they suggest that, if such a gating mechanism exists, it operates independently of episodic retrieval. It raises the possibility of a long-term memory gating mechanism specifically for unsupervised statistical information. Furthermore, we propose an alternative account in which DLPFC inhibition alters information sampling strategies, promoting a broader, more exploratory mode of learning. The observed increase in RT variability, alongside the hemispheric specificity of the effects, supports this view and highlights a potential modality-dependent component in the prefrontal contribution to statistical learning. Future research should more directly investigate the mechanisms linking working memory capacity, information sampling style, and statistical learning. Our study provides insight and proposes novel hypotheses that will be critical for advancing our understanding of the DLPFC's interaction with statistical learning across both typical and altered brain states.

Acknowledgement

This work was supported by the French National Grant Agency (ANR-22-CPJ1-0042-01 and ANR-24-CE37-5807) (to D.N.); the National Brain Research Program project NAP2022-I-2/2022 (to D.N.). The Authors thank Dóra Osztényi, Noémi Papp, and Ádám Bicski for their work in the data collection.

Data and code availability

Data and codes for analysis are available on the following link: https://osf.io/e2rzs/?view_only=88125069a7ad418f804374378bacc441

References

1. Poldrack, R. A. & Packard, M. G. Competition among multiple memory systems: Converging evidence from animal and human brain studies. *Neuropsychologia* **41**, 245–251 (2003).

2. Borragan, G., Slama, H., Destrebecqz, A. & Peigneux, P. Cognitive fatigue facilitates procedural sequence learning. *Front. Hum. Neurosci.* **10**, 179603 (2016).

3. Zhong, L. *et al.* Unsupervised pretraining in biological neural networks. *Nat. 2025* 1–8

(2025) doi:10.1038/s41586-025-09180-y.

4. Thompson-Schill, S. L., Ramscar, M. & Chrysikou, E. G. Cognition without control: When a little frontal lobe goes a long way. **18**, 259–263 (2009).

5. Sherman, B. E. & Turk-Browne, N. B. Statistical prediction of the future impairs episodic encoding of the present. *PNAS* **117**, 22760–22770 (2020).

6. Daw, N. D., Niv, Y. & Dayan, P. Uncertainty-based competition between prefrontal and dorsolateral striatal systems for behavioral control. *Nat. Neurosci.* **8**, 1704–1711 (2005).

7. Virag, M. *et al.* Competition between frontal lobe functions and implicit sequence learning: evidence from the long-term effects of alcohol. *Exp. Brain Res.* **233**, 2081–2089 (2015).

8. Pedraza, F. *et al.* Evidence for a competitive relationship between executive functions and statistical learning. *bioRxiv* 2023.01.19.524710 (2023) doi:10.1101/2023.01.19.524710.

9. Ambrus, G. G. *et al.* When less is more: Enhanced statistical learning of non-adjacent dependencies after disruption of bilateral DLPFC. *J. Mem. Lang.* **114**, 104144 (2020).

10. Tóth-Fáber, E., Nemeth, D. & Janacsek, K. Lifespan developmental invariance in memory consolidation: evidence from procedural memory. *PNAS Nexus* **2**, (2023).

11. Nemeth, D., Janacsek, K. & Fiser, J. Age-dependent and coordinated shift in performance between implicit and explicit skill learning. *Front. Comput. Neurosci.* **7**, 147 (2013).

12. Juhasz, D., Nemeth, D. & Janacsek, K. Is there more room to improve? The lifespan trajectory of procedural learning and its relationship to the between- and within-group differences in average response times. *PLoS One* **14**, e0215116 (2019).

13. Nemeth, D., Janacsek, K., Polner, B. & Kovacs, Z. A. Boosting Human Learning by Hypnosis. *Cereb. Cortex* **23**, 801–805 (2013).

14. Smalle, E. H. M., Muylle, M., Duyck, W. & Szmalec, A. Less Is More: Depleting

Cognitive Resources Enhances Language Learning Abilities in Adults. *J. Exp. Psychol. Gen.* **150**, 2423–2434 (2021).

15. Smalle, E. H. M., Daikoku, T., Szmalec, A., Duyck, W. & Möttönen, R. Unlocking adults' implicit statistical learning by cognitive depletion. *PNAS* **119**, (2022).

16. Szücs-Bencze, L. *et al.* Enhancing retrieval capacity of the predictive brain through dorsolateral prefrontal cortex intervention. *Cereb. Cortex* **35**, (2025).

17. Janacsek, K., Ambrus, G. G., Paulus, W., Antal, A. & Nemeth, D. Right Hemisphere Advantage in Statistical Learning: Evidence From a Probabilistic Sequence Learning Task. *Brain Stimul.* **8**, 277–282 (2015).

18. Roser, M. E., Fiser, J., Aslin, R. N. & Gazzaniga, M. S. Right hemisphere dominance in visual statistical learning. *J. Cogn. Neurosci.* **23**, 1088–1099 (2011).

19. Wolford, G., Miller, M. B. & Gazzaniga, M. The left hemisphere's role in hypothesis formation. *J. Neurosci.* **20**, RC64–RC64 (2000).

20. Greeley, B. & Seidler, R. D. Differential effects of left and right prefrontal cortex anodal transcranial direct current stimulation during probabilistic sequence learning. *J. Neurophysiol.* **121**, 1906–1916 (2019).

21. Smittenaar, P., FitzGerald, T. H. B., Romei, V., Wright, N. D. & Dolan, R. J. Disruption of Dorsolateral Prefrontal Cortex Decreases Model-Based in Favor of Model-free Control in Humans. *Neuron* **80**, 914–919 (2013).

22. Galea, J. M., Albert, N. B., Ditye, T. & Miall, R. C. Disruption of the Dorsolateral Prefrontal Cortex Facilitates the Consolidation of Procedural Skills. *J. Cogn. Neurosci.* **22**, 1158–1164 (2010).

23. Savic, B., Müri, R. & Meier, B. High Definition Transcranial Direct Current Stimulation Does Not Modulate Implicit Task Sequence Learning and Consolidation. *Neuroscience* **414**, 77–87 (2019).

24. Savic, B., Cazzoli, D., Müri, R. & Meier, B. No effects of transcranial DLPFC stimulation on implicit task sequence learning and consolidation. *Sci. Rep.* **7**, 9649 (2017).

25. Balconi, M. Dorsolateral prefrontal cortex, working memory and episodic memory processes: Insight through transcranial magnetic stimulation techniques. *Neurosci. Bull.* **29**, 381–389 (2013).

26. Ruchkin, D. S., Grafman, J., Cameron, K. & Berndt, R. S. Working memory retention systems: A state of activated long-term memory. *Behav. Brain Sci.* **26**, 709–728 (2003).

27. Habib, R., Nyberg, L. & Tulving, E. Hemispheric asymmetries of memory: The HERA model revisited. *Trends Cogn. Sci.* **7**, 241–245 (2003).

28. Tulving, E., Kapur, S., Craik, F. I. M., Moscovitch, M. & Houle, S. Hemispheric encoding/retrieval asymmetry in episodic memory: positron emission tomography findings. *PNAS* **91**, 2016–2020 (1994).

29. Howard, J. H. & Howard, D. V. Age differences in implicit learning of higher order dependencies in serial patterns. *Psychol. Aging* **12**, 634–656 (1997).

30. Nagy, M., Kónya, A. & Király, I. Az Asszociatív Emlékezet Fejlődése 6–10 Éves Kor És Fiatal Felnőttkor Között. *Pszichológia* **33**, 169–184 (2013).

31. Cohn, M. & Moscovitch, M. Dissociating measures of associative memory: Evidence and theoretical implications. *J. Mem. Lang.* **57**, 437–454 (2007).

32. Herwig, U., Satrapi, P. & Schönfeldt-Lecuona, C. Using the International 10-20 EEG System for Positioning of Transcranial Magnetic Stimulation. *Brain Topogr.* **16**, 95–99 (2003).

33. Case, R., Kurland, D. M. & Goldberg, J. Operational efficiency and the growth of short-term memory span. *J. Exp. Child Psychol.* **33**, 386–404 (1982).

34. JASP Team. JASP (Version 0.19.3.0). at https://jasp-stats.org/ (2024).

35. RStudio Team. RStudio: Integrated Development Environment for R. at https://posit.co/products/open-source/rstudio/ (2020).

36. Singmann, H. *et al.* afex: Analysis of Factorial Experiments. at https://cran.r-project.org/web/packages/afex/index.html (2024).

37.   Lenth, R. V. emmeans: Estimated Marginal Means, aka Least-Squares Means. at https://cran.r-project.org/package=emmeans (2024).

38.   Reback, J. *et al.* pandas-dev/pandas: Pandas 1.0.3. (2020) doi:10.5281/ZENODO.3715232.

39.   Harris, C. R. *et al.* Array programming with NumPy. *Nat. 2020 5857825* **585**, 357–362 (2020).

40.   Shih, D. J. H. io: A Unified Framework for Input-Output Operations in R. *CRAN Contrib. Packag.* (2014) doi:10.32614/CRAN.PACKAGE.IO.

41.   Waskom, M. *et al.* seaborn v0.8.1. at https://doi.org/10.5281/ZENODO.883859 (2017).

42.   Hunter, J. D. Matplotlib: A 2D graphics environment. *Comput. Sci. Eng.* **9**, 90–95 (2007).

43.   Soetens, E., Melis, A. & Notebaert, W. Sequence learning and sequential effects. *Psychol. Res.* **69**, 124–137 (2004).

44.   Lüdecke, D. Data Visualization for Statistics in Social Science [R package sjPlot version 2.9.0]. *CRAN Contrib. Packag.* (2025) doi:10.32614/CRAN.PACKAGE.SJPLOT.

45.   Wagenmakers, E. J., Wetzels, R., Borsboom, D. & van der Maas, H. L. J. Why Psychologists Must Change the Way They Analyze Their Data: The Case of Psi: Comment on Bem (2011). *J. Pers. Soc. Psychol.* **100**, 426–432 (2011).

46.   Smalle, E. H. M., Panouilleres, M., Szmalec, A. & Möttönen, R. Language learning in the adult brain: disrupting the dorsolateral prefrontal cortex facilitates word-form learning. *Sci. Rep.* **7**, 1-9. (2017).

47.   Filoteo, J. V., Lauritzen, S. & Maddox, W. T. Removing the frontal lobes: The effects of engaging executive functions on perceptual category learning. *Psychol. Sci.* **21**, 415–423 (2010).

48.   Tóth, B. *et al.* Dynamics of EEG functional connectivity during statistical learning.

Neurobiol. Learn. Mem. **144**, 216–229 (2017).

49. Park, J., Janacsek, K., Nemeth, D. & Jeon, H. A. Reduced functional connectivity supports statistical learning of temporally distributed regularities. *Neuroimage* **260**, 119459 (2022).

50. Zwart, F. S., Vissers, C. T. W. M., Kessels, R. P. C. & Maes, J. H. R. Procedural learning across the lifespan: A systematic review with implications for atypical development. *Journal of Neuropsychology* vol. 13 149–182 at https://doi.org/10.1111/jnp.12139 (2019).

51. Ruchkin, D. S., Grafman, J., Cameron, K. & Berndt, R. S. Working memory retention systems: A state of activated long-term memory. *Behav. Brain Sci.* **26**, 709–728 (2003).

52. Smith, E. E. *et al.* Spatial versus Object Working Memory: PET Investigations. *J. Cogn. Neurosci.* **7**, 337–356 (1995).

53. Wager, T. D., Insler, R. & Smith, E. E. The neural bases of distracter-resistant working memory. *Nat. Preced. 2011* 1–1 (2011) doi:10.1038/npre.2011.5526.1.

54. Frank, S. M. *et al.* Fundamental Differences in Visual Perceptual Learning between Children and Adults. *Curr. Biol.* **31**, 427-432.e5 (2021).

55. Divjak, D. & Milin, P. Exploring and Exploiting Uncertainty: Statistical Learning Ability Affects How We Learn to Process Language Along Multiple Dimensions of Experience. *Cogn. Sci.* **44**, e12835 (2020).

56. Pesthy, O. *et al.* Intact predictive processing in autistic adults: evidence from statistical learning. *Sci. Reports 2023 131* **13**, 1–10 (2023).

57. Wilson, R. C., Bonawitz, E., Costa, V. D. & Ebitz, R. B. Balancing exploration and exploitation with information and randomization. *Curr. Opin. Behav. Sci.* **38**, 49 (2020).

58. Feng, S. F., Wang, S., Zarnescu, S. & Wilson, R. C. The dynamics of explore–exploit decisions reveal a signal-to-noise mechanism for random exploration. *Sci. Reports 2021 111* **11**, 1–15 (2021).

59.	Wu, C. M., Schulz, E., Pleskac, T. J. & Speekenbrink, M. Time pressure changes how people explore and respond to uncertainty. *Sci. Reports 2022 121* **12**, 1–14 (2022).

60.	Oberauer, K. Working memory and attention - A conceptual analysis and review. *J. Cogn.* **2**, (2019).

61.	Olivers, C. N. L., Meijer, F. & Theeuwes, J. Feature-based memory-driven attentional capture: Visual working memory content affects visual attention. *J. Exp. Psychol. Hum. Percept. Perform.* **32**, 1243–1265 (2006).

62.	Jonides, J. *et al.* Spatial working memory in humans as revealed by PET. *Nat. 1993 3636430* **363**, 623–625 (1993).

63.	Fried, P. J., Rushmore, R. J., Moss, M. B., Valero-Cabré, A. & Pascual-Leone, A. Causal evidence supporting functional dissociation of verbal and spatial working memory in the human dorsolateral prefrontal cortex. *Eur. J. Neurosci.* **39**, 1973–1981 (2014).

64.	Lycke, C., Specht, K., Ersland, L. & Hugdahl, K. An fMRI study of phonological and spatial working memory using identical stimuli. *Scand. J. Psychol.* **49**, 393–401 (2008).

65.	Reuter-Lorenz, P. A. *et al.* Age differences in the frontal lateralization of verbal and spatial working memory revealed by PET. *J. Cogn. Neurosci.* **12**, 174–187 (2000).

66.	Smith, E. E., Jonides, J. & Koeppe, R. A. Dissociating verbal and spatial working memory using PET. *Cereb. Cortex* **6**, 11–20 (1996).

67.	Sandrini, M., Cappa, S. F., Rossi, S., Rossini, P. M. & Miniussi, C. The role of prefrontal cortex in verbal episodic memory: rTMS evidence. *J. Cogn. Neurosci.* **15**, 855–861 (2003).