BioScientist Agent: Designing LLM-Biomedical Agents with KG-Augmented RL Reasoning Modules for Drug Repurposing and Mechanistic of Action Elucidation

Fan Zhang[1],[*] Yalong Zhao[1],[*] Weihan Zhang[1], Lipeng Lai[1][†]

[1]XtalPi Innovation Center, XtalPi, Beijing, China

fan.zhang@xtalpi.com, yalong.zhao@xtalpi.com,
jzhang56@tufts.edu, lipeng.lai@xtalpi.com

August 8, 2025

Abstract

Drug discovery is protracted, resource-intensive, and afflicted by attrition rates exceeding 90 %, which leaves most diseases, particularly rare or neglected indications, without effective therapies. Drug repurposing offers a cost effective alternative, yet systematic identification of novel drug indication pairs and mechanistic rationales remains hindered by the scale and heterogeneity of biomedical knowledge. We present **BioScientist Agent**, an end to end framework that unifies a billion-fact biomedical knowledge graph with (i) a variational graph auto-encoder for representation learning and link prediction driven repositioning, and (ii) a reinforcement learning module that traverses the graph to recover biologically plausible mechanistic paths. (iii) A large language model (LLM) multi-agent layer orchestrates these components, enabling inference of target pathways for a drug disease pair, and automatic generation of coherent causal reports. Across benchmark datasets BioScientist Agent surpasses state of the art baselines in both accuracy and recall, while furnishing mechanistic explanations that align with curated literature. Its open and modular design accelerates hypothesis generation and reduces experimental overhead in early-stage discovery.

1 Introduction

Drug discovery is a protracted, capital-intensive enterprise, often exceeding a decade and US\$2 billion per approved entity, yet more than 90 % of clinical candidates ultimately fail. Consequently, thousands of common, rare, and neglected diseases remain without efficacious treatment. *Drug repurposing*, systematically identifying new indications for molecules with established safety profiles, offers a promising alternative, but it requires methods capable of mining the vast, heterogeneous biomedical knowledge space and providing transparent mechanistic evidence.

Knowledge graphs (KGs) are able to mine, organize, and effectively manage knowledge from large-scale data to improve the quality of information services and provide users with smarter services. The current KGs including BioKG [WMN20], CKG [SCN+22], and RTX-KG2 [WGK+22], constitute a graph database comprising millions of nodes and relationships. These graphs integrate nearly all biomedical databases, such as DrugBank [KWK+24], ChEMBL [GBB+12], Gene Ontology (GO) [ABB+00], UniProt [Con19], and the Human Metabolome Database (HMDB) [WGO+22]. This extensive integration greatly benefits various biomedical studies, including drug repositioning [GDX22, MZLK23], adverse drug reaction analysis [ZSD+21, JMM22], and proteomics data analysis [SCN+22].

Simultaneously, large language models (LLMs) have evolved into versatile reasoning engines capable of planning, tool use, and scientific discourse. Coupled with structured KG representations, LLMs can encode the semantic content of node description text in vector representations which then facilitate various tasks, such as semantic textual similarity, information retrieval, and clustering [BAM+24].

---

[*]These authors contributed equally to this work

[†]Corresponding author: lipeng.lai@xtalpi.com

LLM-based agents bridge symbolic reasoning with natural-language synthesis, automating hypothesis generation and enhancing interpretability.

Against this backdrop we introduce **BioScientist Agent**, a KG-augmented, LLM-orchestrated platform for drug repositioning and mechanistic elucidation. Our principal contributions are three-fold:

1. We curate and preprocess RTX-KG2 (v2.9.2) to obtain a 6.3 million-node, 41 million-edge biomedical KG and train a variational graph auto-encoder that achieves state-of-the-art performance on drug–disease link-prediction benchmarks.

2. We devise an adversarial actor–critic reinforcement-learning algorithm that discovers biologically plausible drug–target–disease pathways, yielding interpretable mechanistic explanations for each prediction.

3. We integrate these models within an LLM-driven multi-agent system supporting four practical workflows: (i) disease-to-drug search, (ii) drug-to-disease search, (iii) pathway elucidation and (iv) automated causal report generation.

Together, these innovations provide a scalable and explainable framework that accelerates the generation and validation of drug repositioning hypotheses while deepening understanding of their modes of action.

2 Results

We used RTX-KG2 as the foundational knowledge graph for constructing the bioscientist agent system. The version of RTX-KG2 is v2.9.2, which includes 6,381,804 nodes across 38 distinct categories and 40,989,410 edges across 65 distinct types. Compared to other knowledge graphs, RTX-KG2 contains highly comprehensive information for biological research and drug mechanism discovery, including data on drugs, diseases, genes, and functional experiments results. Additionally, each node and edge is accompanied by extensive literature information for data mining and logical reasoning, which is not available in graphs such as CP-KG or BioKG. CP-KG is composed of 1,246,726 associations between 61,146 entities, and BioKG has 105,465 nodes and 2,043,846 edges, significantly fewer than RTX-KG2. Furthermore, larger knowledge graphs like CKG, which currently comprise close to 20 million nodes and 220 million relationships, include a substantial amount of mutation information. The inclusion of extensive mutation data increases the difficulty of knowledge graph training and representation, and the associations between mutations and phenotypes often result in a high number of false positives. Therefore, taking these factors into account comprehensively, we chose RTX-KG2 for the design of the agent and drug discovery. For RTX-KG2, the top three categories of nodes are 'Small-Molecular', 'OrganismTaxon' and 'Gene' and the top three categories of edges are 'has_participant', 'physically_interaction_with' and 'occurs_in' (Fig. 1a). The heatmap of the log values of edge counts illustrates the distribution of node relationships (Fig. 1b). The top five relationships between nodes are concentrated in SmallMolecule to SmallMolecular (2,785,767), Pathway to CellularComponent (2,535,061), Pathway to SmallMolecular (2,268,235), Gene to Disease (1,837,651), and Pathway to OrganismTaxon (1,562,060).

Figure 1: (a) Percentage of nodes by category and percentage of edges by predicate in the customized BKG. (b) Heatmap of log-transformed connection counts between node types

2.1 Overview of BioScientist agent system

The overall structure of the BioScientist Agent is illustrated in Fig. 1. Our objective is to construct an Agent system based on a knowledge graph that encompasses drug repositioning, the discovery of drug mechanisms of action (MoA), and the collection and interpretation of literature related to drug action pathways. Fig. 1 presents the entire workflow of the system:

Initially, the knowledge graph nodes are updated through training. We utilize Meta's large language model, Llama3.1, to represent the descriptive information of the nodes, obtaining 4096-dimensional embeddings. These embeddings are then reduced to 128 dimensions using Gaussian Random Projection. Subsequently, VRGCN is introduced to update the nodes embeddings, and the DistMult [YYH+14] algorithm is employed to predict the existence of links between nodes. After initial feature extraction and graph neural network pretraining, the knowledge graph acquires features with enhanced representational capabilities.

Once robust node features are obtained, we apply a reinforcement learning model to learn the correct pathways from drugs to diseases. Using the adversarial actor–critic (ADAC) RL algorithm [ZWZ+20], we train the critic and action models so that the generated paths traverse proteins, genes, and pathways, thereby creating MoA pathways from drugs to diseases and ranking them accordingly.

After identifying plausible pathways, their interpretation and experimental validation are conducted through the literature agent system, which assesses the rationality of each path. The system comprises of Perception, Brain, and Action components. Perception is responsible for acquiring pathway information from the ADAC module. The Brain analyzes these pathways, first using an agentic system to filter all related PubMed abstracts for relevant causal sentences then analyzing these sentences using a predefined causal inference framework. The Action component generates the final research reports that include entity summaries and mechanisms, entity-entity relationship analysis, supporting and contradictory evidence analysis, drug repurposing strategies, and pathway chain-of-thought reasoning. Through comprehensive system analysis, for a given drug, disease, and target, the system can obtain extensive information necessary for drug discovery and repurposing.

Figure 2: Overview of BioScientist agent (a) Demonstration of RTX-KG2 knowledge graph components and pretraining with VGAE. The model accepts initial features extracted by LLMs for pretraining, while using edge types for edge prediction. (b) ADAC model framework for drug-disease MoA path extraction (c) LLM-driven multi-agent system, The system comprises of Perception, Brain, and Action components.

2.2 BioScientist Agent enables accurate drug repurposing and MoA path predictions

In this study, we evaluated the performance of the BioScientist Agent against other models in drug repositioning tasks. The evaluation metrics included Accuracy, F1 score, Mean Reciprocal Rank (MRR), Hit@1, and Hit@3. These metrics effectively capture the BioScientist Agent's ability to represent the interaction relationships between drugs and diseases within its drug repositioning algorithm. The results indicate that our drug repositioning algorithm surpasses current algorithms across all metrics. Additionally, we compared VRGCN with other models, demonstrating that the encoder-decoder architecture of VRGCN as encoder and DistMult as decoder achieves superior performance. When compared to the existing state-of-the-art model, KGML-xDTD, our model outperforms it on all metrics, with particularly notable improvements in MRR, Hit@1, and Hit@3, increasing by 30.68%, 41.68%, and 39.06% respectively. These findings suggest that GVAE exhibits enhanced performance in addressing related tasks.

2.3 BioScientist Agent elucidating RL-based pathway and underlying mechanisms

We integrated an LLM-based causal agent to evaluate the proposed drug-disease relationships and corresponding MoA. The ADAC model may propose thousands of potential drug-disease pathways and intermediary drug targets for any given disease, so being able to elucidate the underlying mechanisms and biological plausability of each would prove useful for ranking and identifing the most salient

Table 1: Performance Metrics for Drug Repurposing Across Different Models Using Dataset KG2.9.2

Model	Dataset	Accuracy	F1 Score	MRR	Hit@1	Hit@3	Hit@5
GraphSAGE (KGML-xDTD)	KG2.9.2	0.8945	0.8850	0.2790	0.1469	0.3117	0.4056
GAT	KG2.9.2	0.8903	0.8806	0.2873	0.1700	0.3126	0.4085
TAGConv	KG2.9.2	0.8893	0.8785	0.3017	0.1773	0.3377	0.4297
GCN	KG2.9.2	0.8845	0.8729	0.2536	0.1354	0.2775	0.3632
TransformerConv	KG2.9.2	0.8990	0.8899	0.3108	0.1816	0.3434	0.4485
SGFormer	KG2.9.2	0.8969	0.8864	0.2509	0.1233	0.2707	0.3680
GVAE (BioScientist Agent)	KG2.9.2	0.9292	0.9224	0.4240	0.2775	0.4889	0.6036

Table 2: Performance Metrics for MoA path predictions between KGML-xDTD and Bioscientist Agent

	MPR	MRR	Hit@1	Hit@10	Hit@50	Hit@100	Hit@500
KGML-xDTD	94.70%	0.109	0.059	0.193	0.496	0.613	0.849
BioScientist agent	95.36%	0.121	0.068	0.203	0.508	0.619	0.864

ones for further research. For each entity-entity relationship, we queried PubMed abstracts for entity co-occurrences using PubMed Entrez to find semantically related articles. The abstracts were then filtered using an agent-based approach to find unique causal sentences, which were used as foundational evidence for general-purpose GPT-4 models to analyze using a provided causal inference framework. The final output is a list of relevant PubMed quotations used by the LLM in its analaysis, evidence evaluation, and a final confidence score $C \in [0, 1]$ to indicate the causal strength between the two entities. We obtained the overall score for a pathway by getting the geometric mean of all the scores $(\prod_{i=1}^{n} C_i)^{1/n} = (C_1 \times C_2 \times \cdots \times C_n)^{1/n}$ where n is the number of biological entities in the path. Using this pipeline of evidence-based scoring and causal frameworks overcomes common limitations of LLM-based approaches by minimizing speculative reasoning and anchoring all conclusions in peer-reviewed literature. We show more details regarding causal sentences and the causal inference framework used in Section 4.2.3. On a test run with 40 selected pathways, 20 scientifically-tested positive pathways and 20 randomly chosen negative pathways, our agent achieved an ROC AUC of 0.8562, with an accuracy of 0.8750, F1-score 0.8837, precision 0.8261, and recall 0.9500. The higher false positive rate is due to the random selection of negative pathways. Compared to adversarial actor–critic (ADAC) reinforcement learning model, which had an ROC AUC of 0.642, accuracy of 0.539, F1-score of 0.679, and precision of 0.529, the BioScientist Agent performed significantly better. Figure 3 illustrates these performance metrics, highlighting the superior results of the BioScientist Agent in all areas except recall, where both models were equal.

Figure 3: Path classification performance comparison between adversarial actor–critic (ADAC) and BioScientist in 40 selected pathways.

2.4 BioScientist Agent Mining Target Information

By leveraging the capabilities of knowledge graphs, we aim to identify the mechanisms of action (MoA) associated with the targets PRMT5 and FSCN1. Using these graphs, we can quickly access relevant MoA pathways for PRMT5 and FSCN1, as illustrated in Fig4. This enables us to rapidly gather all potential biological concept nodes related to the targets, with a focus on drug molecules, associated proteins, and diseases. Additionally, because each node pair has related PubMed IDs, we can utilize APIs to obtain the impact factor distribution of all papers within a pathway, allowing us to rank and analyze the pathway's reliability. This helps researchers pinpoint trustworthy papers for scientific study and discovery (Fig4). Our findings reveal that the top three most plausible pathways associated with PRMT5 are: carbamazepine - infectious disease - PRMT5 - epilepsy; dexamethasone - PRMT5 - CRK - atopic dermatitis; nitroglycerin - bone morphogenetic protein 2 - PRMT5 - angina pectoris. The top three pathways linked to FSCN1 are: gefitinib - cell proliferation - FSCN1 - non-small cell lung carcinoma; docetaxel - MAPT - FSCN1 - breast neoplasm; capecitabine - GNAS - FSCN1 - colonic neoplasm.

Figure 4: MoA pathway profiles centered on targets PRMT5 and FSCN1, and the distribution of impact factors of all publications involved in the pathways: (a) Extraction of the MoA pathway centered on target PRMT5; (b) Violin plots showing the distribution of impact factors for all publications between every pair of nodes in each PRMT5-associated pathway; (c) Extraction of the MoA pathway centered on target FSCN1; (d) Violin plots showing the distribution of impact factors for all publications between every pair of nodes in each FSCN1-associated pathway.

2.5 BioScientist Agents interpret MoA pathways

The BioScinetsi Agent has the ability to finely interpret an MoA pathway. We use carefully designed prompts for paper information extraction and interpretation to score the reliability of the MoA pathway. The prompt for the multi-agent approach to extract causal information from PubMed abstracts is shown in Supplementary Material 1. Each agent analyzes a provided list of PMIDs and corresponding abstracts to extract either supporting or contradictory sentences and returns that information. In Supplementary Material 2, we show the causal analysis framework used to determine the causal relationship between two entities, including mechanistic evidence, association strength, biological plausibility, literature consistency, causal relationship analysis, and overall confidence. The global summary of a drug-gene pathway provided by the agent is shown in Material 3. This includes a summary of the most notable evidence in determining the causal score along with the PMID citations for the evidence. The agent evaluates both supporting and contradictory evidence to identify gaps that limit the pathway's score. The final output is a JSON file, with the overall pathway analysis part shown in Fig 5. The complete JSON file is in Supplementary Material 4.

In the JSON file, we analysed the putative repurposing of enzalutamide (Xtandi) for mitigating prostate cancer through the targets of the AR gene and the PTEN gene. The agent produced a causal confidence score of C=0.9 based on evidence extracted from 1025 articles. Each individaul entity-entity relationship within the path has its own analysis of BH criteria, mechanistic evidence, causal terminology, potential mechanism, key strengths/weaknesses, and citations. Key causal sentences used in the agent analysis were identified and then summarized in the global summary. For example: "Study [PMID: 23701654][DSS+13] demonstrated that enzalutamide acts as an androgen receptor antagonist, inhibiting AR signaling pathways, supporting a strong causal relationship between enzalutamide and AR.", "Study [30177856] [dLPRDC+18] showed that enzalutamide treatment is associated with reduced AR activity, indicating a direct effect of enzalutamide on AR levels.", "Study [PMID: 21532617] [WRH+11] elucidated that AR inhibits PTEN transcription in prostate cancer cells, establishing a

causal link between AR and PTEN.", "Study [PMID: 23902739] [CLSK+13] reported that loss of PTEN is frequently observed in cancer, resulting in the deregulation of cell survival, growth, and proliferation, indicating a strong association between PTEN loss and prostate cancer progression." The agent also proposed a potential pathway MoA based on the mechanistic evidence it identified ("The potential mechanism involves enzalutamide binding to the androgen receptor (AR), preventing its activation and subsequent transcription of target genes. This inhibition leads to decreased PTEN expression, as AR negatively regulates PTEN transcription. The loss of PTEN function activates the PI3K/AKT signaling pathway, promoting cell survival and proliferation, ultimately contributing to prostate cancer progression."). For high-confidence paths, the agent proposes detailed drug repurposing strategies based on the evidence provided: "Given the total_confidence_score of 0.9, a candidate drug for repurposing could be a selective AR modulator (SARM) that has shown promise in modulating AR activity without the resistance seen with enzalutamide. The experimental design would involve using prostate cancer cell lines with varying levels of AR expression and PTEN status. Controls would include untreated cells and cells treated with enzalutamide. Endpoints would include measuring AR and PTEN expression levels through qPCR and Western blotting, as well as assessing cell proliferation and apoptosis through MTT assays and flow cytometry. The experiment would test the candidate drug's effect on the pathway by evaluating whether it can effectively reduce AR activity and restore PTEN levels compared to enzalutamide. Expected outcomes would include a reduction in AR activity and an increase in PTEN levels, supporting the hypothesis that the candidate drug could be effective in overcoming resistance mechanisms. Potential challenges include variability in patient-derived cell lines and the need for in vivo validation. Next steps would involve optimizing the dosing regimen and exploring combination therapies with existing treatments." Identified weak links such as "Further research is needed to explore the specific mechanisms by which AR variants contribute to enzalutamide resistance.", and "More studies are required to clarify the role of other signaling pathways that may interact with the AR-PTEN axis in prostate cancer" were flagged for further wetlab investigation. Collectively, this framework provides viable and substantiated biological causal pathways that are traceable to primary literature by combining epidemiological causal inference and machine reasoning to deliver a transparent and rigorous solution.

Figure 5: Demonstration of the overall pathway analysis in MoA pathway interpretation, with complete results in Supplementary Material 4.

3 Discussion

Knowledge graphs encompass an extensive amount of information, and leveraging them can accelerate tasks such as drug repositioning and predicting drug mechanisms of action. This not only speeds up the progress of drug development but also reduces research and development costs. In this study, we developed the BioScientist Agent system, which integrates graph neural network algorithms, ADAC reinforcement learning algorithms, and an agent-based literature interpretation and path scoring system. This comprehensive system manages the entire workflow, including knowledge graph training, edge discrimination, MoA path searching, and result interpretation, thereby playing a crucial role in accelerating drug development.

Previous research seldom involved training on such large-scale knowledge graphs due to computational constraints. Typically, these studies limited the graphs to 4-5 types of nodes, such as drugs, diseases, and genes, encompassing tens of thousands of nodes and millions of edges. They excluded additional node types like cell types, transcription factors, and microRNA, as well as numerous edges supported by only a few publications. In contrast, our study incorporates all edges deemed present in existing research into the training scope, with the number of nodes reaching the millions and the number of edges reaching tens of millions. This results in a highly comprehensive dataset, significantly enhancing the model's ability to discover new and potential drug-disease relationships and to explore drug mechanisms of action, even for those relationships with minimal prior research.

Thanks to continuous advancements in biological science research, the data sources for knowledge graphs are constantly being updated. We believe that an Agent system capable of rapid iteration is

of greater significance. Currently, the BioScientist Agent can quickly iterate when handling updated databases; however, it still requires retraining from scratch to associate new nodes with existing ones using graph neural network algorithms. This process is time-consuming and necessitates extensive data preprocessing and initial feature extraction for nodes. In our future work, we aim to establish mechanisms for the rapid integration of new nodes and edge relationships. This includes updating nodes through local graph representation capabilities and incorporating more node representation models to enhance their representation abilities. For instance, we plan to utilize large model tools like 'SMI-TED289M' that encode molecular SMILES for feature representation [SSB$^+$24]. By training on drug molecules that have some experimental data but still possess many unknown functions, we aim to predict targeted diseases and the mechanisms of action (MoA) of these unknown molecules. Ultimately, this will enable the BioScientist Agent system to exert a greater impact.

4   Methods

4.1   Datasets

1. Original datasets   We used the canonicalized version of Reasoning Tool X Knowledge Graph 2 (RTX-KG2c) v2.9.2 for constructing, training, and extracting drug-disease pathways from the knowledge graph. It integrates a large amount of human-curated and publication-based database knowledge and categorizes nodes and edges based on the Biolink model. The RTX-KG2c version v2.9.2 we used contains approximately 6,873,228 nodes and 46,568,039 edges, covering knowledge from 70 public biomedical sources, with all biological concepts represented as nodes and all concept-predicate-concept relationships represented as edges. After excluding node types unrelated to biological explanations and edges with redundant concepts, the graph retains 6,381,804 nodes with 37 distinct categories (Fig. 1A) and 40,989,410 edges with 67 distinct types (Fig. 1B) for downstream model training.

2.   Demonstration paths   Demonstration paths are prepared before conducting reinforcement learning because drugs treat diseases through specific targets. We constructed certain drug-to-disease paths, such as drug–gene–protein–disease and drug–drug–protein–disease. We extracted 1.9 million demonstration paths from the knowledge graph, based on known drug-target interactions collected from two curated biomedical data sources: DrugBank (v5.1) and Molecular Data Provider (v1.2), as well as the normalized Google distance (NGD) score based on PubMed publications, with NGD scores required to be less than 0.6. The PubMed publication-based NGD [53] defined below:

$$\text{NGD}(c_1, c_2) = \frac{\max\{\log N(c_1), \log N(c_2)\} - \log N(c_1, c_2)}{\log N - \min\{\log N(c_1), \log N(c_2)\}} \tag{1}$$

where c_1 and c_2 are two biological concepts used in the customized BKG, $N(c_1)$ and $N(c_2)$ respectively represent the total number of unique PubMed IDs associated with c_1 and c_2, $N(c_1, c_2)$ is the total number of unique PubMed IDs shared between c_1 and c_2, and N is the total number of pairs of Medical Subject Heading (MeSH) terms annotations in the PubMed database.

4.2   Models

The framework of BioScientist Agent consists of 3 modules: a VGAE-based Graph Neural Network Representation Model [KW16] for drug repurposing, an adversarial actor-critic RL model for drug-to-disease pathway scoring and a proposed LLM-powered multi-agent system for automated scientific discover and MoA path re-scoring and interpreting. The architecture of the whole Agent system showed in Fig2.

4.2.1   VGAE-based Graph Neural Network Representation Model

Model Architecutre   Let $G = (V, E)$ be a directed biomedical knowledge graph, where:

- Each node $v \in V$ represents a biological entity (e.g., a specific drug, disease, gene, or pathway)

- Each edge $e \in E$ represents a biomedical relationship

Given G, we employ a VGAE to learn low-dimensional node embeddings $\mathbf{Z} \in \mathbb{R}^{|V| \times d}$, where $d \ll |V|$. The model consists of the following components:

1. Variational Relational Graph Convolutional Network (VRGCN) Encoder

The encoder uses a two-layer relational graph convolutional network with separate variational heads to infer stochastic embeddings:

$$\mathbf{H}^{(1)} = \mathrm{ReLU}\left(\sum_{r \in \mathcal{R}} \tilde{\mathbf{A}}_r \mathbf{X} \mathbf{W}_r^{(0)}\right),$$

$$\begin{cases} \mathbf{H}_\mu^{(2)} = \sum_{r \in \mathcal{R}} \tilde{\mathbf{A}}_r \mathbf{H}^{(1)} \mathbf{W}_{r,\mu}^{(1)} \\ \mathbf{H}_\sigma^{(2)} = \sum_{r \in \mathcal{R}} \tilde{\mathbf{A}}_r \mathbf{H}^{(1)} \mathbf{W}_{r,\sigma}^{(1)} \end{cases}$$

where:

- $\tilde{\mathbf{A}}_r = \mathbf{D}_r^{-\frac{1}{2}}(\mathbf{A}_r + \mathbf{I})\mathbf{D}_r^{-\frac{1}{2}}$ is the normalized adjacency matrix for relation $r \in \mathcal{R}$ with self-loops

- $\mathbf{X} \in \mathbb{R}^{|\mathcal{V}| \times m}$ is the node feature matrix with m-dimensional features

- $\{\mathbf{W}_r^{(0)}\}_{r \in \mathcal{R}}$ are first-layer relation-specific weight matrices

- $\{\mathbf{W}_{r,\mu}^{(1)}, \mathbf{W}_{r,\sigma}^{(1)}\}_{r \in \mathcal{R}}$ are second-layer weight matrices for mean and variance heads

- $\mathrm{ReLU}(x) = \max(0, x)$ is the rectified linear unit activation function

- $\mathcal{R}$ is the set of relation types in the graph

The latent embeddings $\mathbf{Z}$ are sampled using the reparameterization trick:

$$\mathbf{Z} = \boldsymbol{\mu} + \boldsymbol{\sigma} \odot \boldsymbol{\epsilon}, \quad \boldsymbol{\epsilon} \sim \mathcal{N}(0, \mathbf{I})$$

where:

$$\boldsymbol{\mu} = \mathbf{H}_\mu^{(2)}, \quad \log \boldsymbol{\sigma} = \mathbf{H}_\sigma^{(2)}$$

- Separate RGCN heads compute $\boldsymbol{\mu}$ and $\log \boldsymbol{\sigma}$ to enable variational learning

- The architecture supports multiple relation types through relation-specific parameters

- Dropout is typically applied between layers for regularization

2. Decoder for Link Prediction The decoder uses a DistMult approach for link prediction by reconstructing edges:

$$p(\mathbf{A}_{ij} = 1 \mid \mathbf{z}_i, \mathbf{z}_j, \mathbf{r}_k) = \sum (\mathbf{z}_i \odot \mathbf{r}_k \odot \mathbf{z}_j)$$

where:

- $\mathbf{z}_i$ and $\mathbf{z}_j$ are the source and destination node embeddings

- $\mathbf{r}_k$ is the relation embedding for the edge type

- $\odot$ denotes element-wise multiplication

The reconstructed adjacency matrix $\hat{\mathbf{A}}$ is derived from these scores for each relation type. The parameters are initialized using Xavier uniform initialization to ensure proper scaling of inputs and outputs.

Model Training The model optimizes the loss function that combines reconstruction and regularization:

$$\mathcal{L} = \underbrace{\mathbb{E}_{q(\mathbf{Z}|\mathbf{X},\mathbf{A})} \left[\log p(\mathbf{A}|\mathbf{Z})\right]}_{\text{Reconstruction loss}} + \lambda \cdot \underbrace{\|\mathbf{Z}\|^2 + \|\mathrm{rel\_emb}\|^2}_{\text{Regularization}},$$

where:

- The reconstruction loss is computed using binary cross-entropy with logits.

- Regularization is applied to both node embeddings $\mathbf{Z}$ and relation embeddings rel_emb.

- λ controls the regularization strength (default: $\lambda = 0.01$).

Model Evaluation Our evaluation pipeline includes several key components designed to assess the performance of the VGAE model on knowledge graph link prediction tasks. We first obtain true drug-disease interactions, including true positives and true negatives, from MyChemData, SemMedDB Data, NDF-RT Data, and RepoDB Data. Positive samples (known drug-disease treatment relationships) are divided into two subsets: expert demonstration pairs based on knowledge graph paths and other positive pairs, ensuring effective utilization of high-quality samples. During the division process, a drug-centered hierarchical sampling strategy is adopted. Single-indication drugs are prioritized for the training set, while multi-indication drugs are stratified using drug IDs to prevent data leakage by dispersing different indications of the same drug across datasets. We designed a balanced random negative sample generation mechanism, creating random pairings for each drug and disease in the positive samples, ensuring they do not duplicate known positives or negatives. Random negative samples are labeled as an independent category (label 2) and are mainly used to support ranking evaluation metrics like MRR (Mean Reciprocal Rank) and Hits@K in link prediction tasks. The final dataset composition for each set (training/validation/testing) includes expert demonstration positives, other positives, true negatives, and random negatives, with random shuffling ensuring uniform sample distribution. Additionally, a large-scale set of random pairs is generated specifically for ranking performance evaluation in the validation and testing phases, providing a comprehensive and rigorous evaluation benchmark for drug repositioning models.

1. Embedding Generation and Feature Construction. The evaluation process begins by loading a pre-trained VGAE model and generating node embeddings: (1) Load the best trained model checkpoint and set it to evaluation mode; (2) Generate embeddings for all nodes using the encoder: $z = \text{Encoder}(X, E, R)$, where X represents the init node feature matrix, E denotes the edge index matrix defining graph connectivity, and R represents the edge type/relation indices for each edge; (3) Create a dictionary mapping node IDs to their corresponding embeddings. For link prediction, we construct feature vectors by concatenating source and target node embeddings: $\mathbf{x}_{pair} = [\mathbf{z}_{source} \oplus \mathbf{z}_{target}]$, where $\oplus$ denotes concatenation, resulting in 256-dimensional feature vectors.

2. Classification and Prediction. A Random Forest classifier is employed as the final prediction layer with 2000 estimators, maximum depth of 35, square root of features per split, and automatic class weight balancing to handle imbalanced datasets. The classifier is fitted on concatenated embeddings from training pairs.

3. Evaluation Metrics. We employ multiple evaluation metrics to comprehensively assess model performance. Classification metrics include: (1) *Accuracy*:

$$\text{Acc} = \frac{\text{TP} + \text{TN}}{\text{TP} + \text{TN} + \text{FP} + \text{FN}} \tag{2}$$

where TP, TN, FP, FN are true positives, true negatives, false positives, and false negatives, respectively; (2) *Macro F1-score*:

$$\text{F1}_{\text{macro}} = \frac{1}{|C|} \sum_{c \in C} \frac{2 \cdot \text{precision}_c \cdot \text{recall}_c}{\text{precision}_c + \text{recall}_c} \tag{3}$$

where $|C|$ is the number of classes; (3) *Micro F1-score*:

$$\text{F1}_{\text{micro}} = \frac{2 \cdot \text{TP}}{2 \cdot \text{TP} + \text{FP} + \text{FN}} \tag{4}$$

aggregated globally across all classes. Ranking metrics for link prediction evaluation include: (1) *Mean Reciprocal Rank (MRR)*:

$$\text{MRR} = \frac{1}{|Q|} \sum_{i=1}^{|Q|} \frac{1}{\text{rank}_i} \tag{5}$$

where $|Q|$ is the number of positive query pairs and rank_i is the rank of the correct answer among all candidates for query i; (2) *Hit@K*:

$$\text{Hit@K} = \frac{1}{|Q|} \sum_{i=1}^{|Q|} I(\text{rank}_i \leq K) \tag{6}$$

where $I(\cdot)$ is the indicator function that equals 1 if the condition is true and 0 otherwise, evaluated for $K \in \{1, 3, 5\}$.

4.2.2 Adversarial Actor-Critic RL Model

We formulate the MOA prediction as a path-finding problem and adapt the adversarial actor-critic reinforcement learning model to solve it. Reinforcement learning is defined as a Markov decision process (MDP) that contains:

Basic concept

- **States**: Represent the current situation of the agent within the environment.

  Each state s_t at time t is defined as:

  $$s_t = \left(v_{\text{drug}}, v_t, (v_{t-1}, e_{t-1}), \ldots, (v_{t-K}, e_{t-(K-1)})\right) \tag{7}$$

  where:

  - $v_{\text{drug}} \in V_{\text{drug}}$ is the given starting drug node.
  - $v_t \in V$ represents the node where the agent is located at time t.
  - $\left(v_{t-K}, e_{t-(K-1)}\right)$ represents the previous Kth node and the $(K-1)$th predicate.

  For the initial state s_0, the previous nodes and predicates are substituted by a special dummy node and predicate.

  The state embedding $\mathbf{s}_t$ is obtained by concatenating the embeddings of all nodes and predicates in s_t, where:

  - Node embeddings are generated using the LLM2Vec model.
  - Predicate embeddings are represented by one-hot vectors.

- **Actions**: Define the possible moves the agent can make.

  The action space A_t at each node v_t includes a self-loop action a_{self} and actions to reach its outgoing neighbors in the knowledge graph G. To manage memory limitations and handle nodes with large out-degrees, neighbor actions are pruned based on PageRank scores if a node has more than 3,000 neighbors. Specifically, the action space is defined as:

  $$A_t = \left\{a_{\text{self}}, a_1, \ldots, a_{n_{v_t}}\right\} \tag{8}$$

  where n_{v_t} is the out-degree of node $v_t \in V$.

  For each action $a_t = (e_t, v_{t+1}) \in A_t$ taken at time t, its embedding $\mathbf{a}_t$ is obtained by concatenating its node and predicate embeddings. Two embedding matrices are learned:

  $$\mathbf{E}_N \in \mathbb{R}^{N_n \times d} \quad \text{for nodes, and} \quad \mathbf{E}_P \in \mathbb{R}^{N_p \times d} \quad \text{for predicates.} \tag{9}$$

  where:

  - d is the embedding dimension.
  - N_n is the number of nodes in the graph.
  - N_p is the number of predicate categories.
  - Each subnetwork uses separate embedding matrices.

- **Rewards**: Provide feedback to the agent based on the actions taken.

  During the path-searching process, the agent only receives a terminal reward $R_{e,T}$ from the environment. There are no intermediate rewards from the environment, i.e.,

  $$R_{e,t} = 0, \quad \forall t < T \tag{10}$$

Let v_T be the last node of the path, and N_{drug} be the known diseases that drug v_{drug} can treat. The terminal reward $R_{e,T}$ from the environment is calculated using the drug repurposing model as follows:

$$R_{e,T} = f(v_{\text{drug}}, v_T)_{\text{treat}} \tag{11}$$

where p_{treat} is the "treat" class probability predicted by the drug repurposing model f.

- **Policy**: Determines the agent's behavior by mapping states to actions.

Model Architecture The adversarial actor–critic RL model comprises four subnetworks that share the same Multilayer Perceptron (MLP) architecture MLP_i, where i identifies each subnetwork (e.g., a for actor, c for critic). However, they have different parameters:

$$\text{MLP}_i(\mathbf{X}) = \text{BA}\left(\text{BA}\left(\mathbf{X}\mathbf{W}_{1i} + \mathbf{b}_{i1}\right)\mathbf{W}_{2i} + \mathbf{b}_{i2}\right)\mathbf{W}_{3i} + \mathbf{b}_{i3} \tag{12}$$

where: $\{\mathbf{W}_{1i}, \mathbf{W}_{2i}, \mathbf{W}_{3i}, \mathbf{b}_{i1}, \mathbf{b}_{i2}, \mathbf{b}_{i3}\}$ are the parameters and biases of linear transformations. BA represents a batch normalization layer followed by an Exponential Linear Unit (ELU) activation function.

The Model Architecture consists of the following components:

- **Actor Network**: The actor network learns a path-finding policy π_θ, where θ represents all parameters of the actor network. This policy guides the agent to choose an action a_t from the action space A_t based on the current state s_t:

$$\pi_\theta(a_t \mid s_t, A_t) = \text{softmax}\left(\mathbf{A}_t \odot \text{MLP}_a(\mathbf{s}_t)\right) \tag{13}$$

  where:

  - $\mathbf{A}_t$ is the embedding matrix of the action space A_t.
  - $\odot$ denotes the dot product.

  $\pi_\theta(a_t \mid s_t, A_t)$ represents the probability of choosing action a_t at time t from the action space A_t given the state s_t.

- **Critic Network**: The critic network estimates the expected reward $Q_\phi(s_t, a_t)$, where ϕ represents all parameters of the critic network, if the agent takes action a_t at state s_t:

$$Q_\phi(s_t, a_t) = \text{MLP}_c(\mathbf{s}_t) \odot \mathbf{a}_t \tag{14}$$

- **Path Discriminator Network**: Since the RL agent only receives a terminal reward $R_{e,T}$ indicating whether it reaches an expected target, a path discriminator network is introduced to encourage the agent to find biologically reasonable paths and provide intermediate rewards. This network acts as a binary classifier distinguishing whether a path segment (s_t, a_t) is from demonstration paths or generated by the actor network.

  Positive Samples: Known demonstration path segments (s_t^D, a_t^D).

  Negative Samples: Actor-generated non-demonstration path segments (s_t^{ND}, a_t^{ND}).

  The path discriminator is defined as:

$$D_p(s, a) = \sigma\left(\text{MLP}_p(\mathbf{s} \oplus \mathbf{a})\right) \tag{15}$$

  where:

  - σ is the sigmoid function.
  - $\oplus$ denotes concatenation.
  - $\mathbf{s}$ and $\mathbf{a}$ are the embeddings of state s and action a, respectively.

The discriminator is optimized with the loss function:

$$L_p = -\mathbb{E}_{(s,a)\sim P_D}\left[\log D_p(s,a)\right] - \mathbb{E}_{(s,a)\sim P_A}\left[\log\left(1 - D_p(s,a)\right)\right] \tag{16}$$

where:

- P_D is the demonstration path segment distribution.
- P_A is the actor-generated non-demonstration path segment distribution.

Based on the discriminator's output $D_p(s_t, a_t)$, the intermediate reward $R_{p,t}$ is calculated as:

$$R_{p,t} = \log D_p(s_t, a_t) - \log\left(1 - D_p(s_t, a_t)\right) \tag{17}$$

- **Meta-Path Discriminator Network**: Similarly, the meta-path discriminator network judges whether the meta-path of the actor-generated paths is similar to that of demonstration paths. A meta-path is defined as the sequence of node categories in the path (e.g., "Drug" → "Gene" → "Gene" → "Disease", "Drug" → "Gene" → "Biological Process" → "Disease").

The meta-path discriminator is defined as:

$$D_m(M) = \sigma\left(\text{MLP}_m(\mathbf{M})\right) \tag{18}$$

where:

- $\mathbf{M}$ is the embedding of the meta-path M, obtained by concatenating the learned category embeddings of all nodes in the path.

It is optimized with the loss function:

$$L_m = -\mathbb{E}_{M\sim P_{DM}}\left[\log D_m(M)\right] - \mathbb{E}_{M\sim P_{AM}}\left[\log\left(1 - D_m(M)\right)\right] \tag{19}$$

where:

- P_{DM} is the distribution of demonstration meta-paths.
- P_{AM} is the distribution of actor-generated non-demonstration meta-paths.

The intermediate reward $R_{m,t}$ from the meta-path discriminator is computed as:

$$R_{m,t} = \log D_m(M) - \log\left(1 - D_m(M)\right) \tag{20}$$

- **Integrated Intermediate Reward**: The integrated intermediate reward R_t at time t combines the rewards from both discriminators and the terminal reward:

$$R_t = \alpha_p R_{p,t} + \alpha_m R_{m,t} + \left(1 - \alpha_p - \alpha_m\right)\gamma^{T-t}R_{e,T} \tag{21}$$

where:

- $\alpha_p \in [0, 1]$ and $\alpha_m \in [0, 1 - \alpha_p]$ are hyperparameters.
- γ is the decay coefficient.
- $R_{e,T}$ is the terminal reward from the environment.

ADAC Model Training From the complete library of expert demonstration paths (containing all inference paths from drugs to diseases), precise matching and extraction are performed based on the drug-disease pairs in the pre-divided training, validation, and test sets. Specifically, this involves converting drug-disease pairs into their corresponding entity IDs and then searching the expert path data for all complete inference paths that start with the specified drug and end with the specified disease. This matching process ensures that each data subset contains only the expert demonstration trajectories strictly corresponding to its drug-disease pairs, maintaining data independence across the training, validation, and test sets. Since a single drug-disease pair may have multiple inference paths, the final expert demonstration dataset retains the diversity and completeness of paths.

1. **Initialization**:

   - The actor network is initialized using the behavior cloning method [**?**], where the training set of demonstration paths guides the agent's sampling using a Mean Squared Error (MSE) loss.

2. **Training Discriminators**:

   - For the first z epochs, the parameters of the actor and critic networks are frozen.
   - The path discriminator network and meta-path discriminator network are trained by minimizing L_p and L_m, respectively.

3. **Joint Optimization**:

   - After z epochs, the actor and critic networks are unfrozen.
   - They are optimized together by minimizing a joint loss:

   $$L_{\text{joint}} = L_a + L_c \tag{22}$$

4. **Critic Network Optimization**:

   - To optimize the critic network, the Temporal Difference (TD) error is minimized with the loss function:
   $$L_c = \text{TD}^2 = \left[(R_t + Q_\phi(s_{t+1}, a_{t+1})) - Q_\phi(s_t, a_t) \right]^2 \tag{23}$$

5. **Actor Network Optimization**:

   - The actor network aims to maximize the expected reward by learning an optimal policy. This is achieved by maximizing:

   $$J(\theta) = \mathbb{E}_{a \sim \pi_\theta} \left[Q_\phi(s_t, a) \right] \tag{24}$$

   - Using the REINFORCE algorithm [**?**], the parameters are updated. To encourage diverse exploration in finding paths, the entropy of π_θ is used as a regularization term. The stochastic gradient of the actor loss L_a is given by:

   $$\nabla_\theta L_a = -\nabla_\theta J(\theta) = -\mathbb{E}_{\pi_\theta} \left[\nabla_\theta \text{TD} \log \pi_\theta(a_t \mid s_t) \right] - \alpha \nabla_\theta \text{Entropy}(\pi_\theta) \tag{25}$$

   where:
   - α is the entropy weight.
   - $\pi_\theta(a_t \mid s_t)$ is the action probability distribution based on the actor policy.

ADAC Model Evaluation Our ADAC-based reinforcement learning evaluation pipeline assesses path-finding performance through probabilistic scoring and ranking metrics on knowledge graph reasoning tasks.

   Model Architecture and Setup. The evaluated model employs an Adversarial Discriminator Actor-Critic (ADAC) architecture with the following components:

- Policy Network: An actor-critic model with hidden layers of [512, 512] for policy learning and [512, 512] for discriminator networks.

- State Representation: 100-dimensional entity embeddings, 100-dimensional entity type embeddings, and 100-dimensional relation embeddings with state history length of 2.

- Path Constraints: Maximum path length of 3 steps with discount factor $\gamma = 0.99$ for reward computation.

Path Generation and Probability Scoring. The evaluation process follows a multi-step procedure:

1. Load pre-trained Random Forest model converted to PyTorch format for reward estimation.

2. Initialize knowledge graph environment with entity embeddings from VGAE model.

3. For each source-target pair, generate action probabilities using the policy network: $\pi(a_t|s_t) = \text{PolicyNet}(s_t, \mathcal{A}_t)$, where s_t is the current state and $\mathcal{A}_t$ is the action space.

4. Calculate weighted log-probability for each path step: $\log p_{weighted} = \log(p_{true} + \epsilon) \times |\mathcal{A}_{valid}|$, where p_{true} is the probability of the true action and $|\mathcal{A}_{valid}|$ is the number of valid actions.

5. Compute cumulative path probability with decay factor: $P_{path} = \sum_{t=1}^{T} \alpha^{t-1} \log p_{weighted,t}$, where $\alpha = 0.9$ is the decay factor and T is the path length.

Path Filtering and Ranking. Expert paths are filtered to exclude low-quality relations (biolink:related_to, biolink:part_of, biolink:coexists_with, biolink:contraindicated_for) and ranked by probability scores in descending order. For each unique node sequence, multiple edge sequences are stored with their respective scores and ranks.

Evaluation Metrics. We employ specialized metrics for path-based reasoning evaluation:

- Mean Percentile Rank (MPR):

$$\text{MPR} = \frac{1}{|Q|} \sum_{i=1}^{|Q|} \left(1 - \frac{\text{rank}_i}{N_i}\right)$$

  where N_i is the total number of paths for query i.

- Mean Reciprocal Rank (MRR):

$$\text{MRR} = \frac{1}{|Q|} \sum_{i=1}^{|Q|} \frac{1}{\text{rank}_i}$$

  where rank_i is the rank of the highest-scoring correct path for query i.

- Hit@K:

$$\text{Hit@K} = \frac{1}{|Q|} \sum_{i=1}^{|Q|} I(\text{rank}_i \leq K)$$

  evaluated for $K \in \{1, 10, 50, 100, 500, 1000\}$.

4.2.3 LLM-powered causal agent system

We used an LLM-based approach to analyze the proposed pathways and MoA for a given disease and rank the causal likelihood of each using PubMed evidence and integrated causal frameworks. Modern causal inference in biomedicine integrates qualitative frameworks like the Bradford-Hill (BH) criteria, including strength of association, consistency, biological plausibility, coherence, and experiment. Our method adopts this by deploying an LLM to extract and evaluate direct causal claims from literature, augmented by structured causal extraction and tracable evidence citations. Operationalizing the Causal Framework: We refine BH criteria to prioritize quantifiable signals of causality that are traceable to peer-reviewed articles. For example, the LLM measures association strength by the frequency of direct causal assertions across abstracts weighted by quality of study design, and will output corresponding PMID citations of each abstract used in the analysis. We also improve clarity by having the LLM identify key causal terms it found in the abstracts, such as "suppresses",

"promotes", "influences", etc. to describe the relationship between two entities and infer a potential mechanism, ensuring tracibiliy by including the PMIDs of the sentences it identified. To mitigate context window limitations, we filter the abstracts to find only sentences that provide supporting or contradictory evidence of a causal relatonship between the two entities. We accomplish this using a multi-agent system; each agent was assigned a number of articles, tasked with extracting key causal sentences as direct quotations. The agents then reconvened and collectively curated the final set of relevant causal sentences, which then served as the foundational evidence for the overall analysis using the causal inference framework. This enables batch processing of thousands of articles simultaneously while preserving causal signals. Output and Interpretability: The BioScientist agent outputs a list that ranks all the potential causal pathways of a disease by their overall pathway score, each with a corresponding JSON report that has a global summary for the pathway, including supporting evidence, drug repurposing potential, mechanistic evidence, and the chain-of-thought reasoning for its score.

References

[ABB+00]   Michael Ashburner, Catherine A Ball, Judith A Blake, David Botstein, Heather Butler, J Michael Cherry, Allan P Davis, Kara Dolinski, Selina S Dwight, Janan T Eppig, et al. Gene ontology: tool for the unification of biology. *Nature genetics*, 25(1):25–29, 2000.

[BAM+24]   Parishad BehnamGhader, Vaibhav Adlakha, Marius Mosbach, Dzmitry Bahdanau, Nicolas Chapados, and Siva Reddy. Llm2vec: Large language models are secretly powerful text encoders. *arXiv preprint arXiv:2404.05961*, 2024.

[CLSK+13]   M Katie Conley-LaComb, Allen Saliganan, Pridvi Kandagatla, Yong Q Chen, Michael L Cher, and Sreenivasa R Chinni. Pten loss mediated akt activation promotes prostate tumor growth and metastasis via cxcl12/cxcr4 signaling. *Molecular cancer*, 12(1):85, 2013.

[Con19]   UniProt Consortium. Uniprot: a worldwide hub of protein knowledge. *Nucleic acids research*, 47(D1):D506–D515, 2019.

[dLPRDC+18] Alicia de Las Pozas, Teresita Reiner, Virginia De Cesare, Matthias Trost, and Carlos Perez-Stable. Inhibiting multiple deubiquitinases to reduce androgen receptor expression in prostate cancer cells. *Scientific reports*, 8(1):13146, 2018.

[DSS+13]   Richa Dhingra, Tina Sharma, Sukhminder Singh, Shivani Sharma, Prince Tomar, Manav Malhotra, and TR Bhardwaj. Enzalutamide: a novel anti-androgen with prolonged survival rate in crpc patients. *Mini Reviews in Medicinal Chemistry*, 13(10):1475–1486, 2013.

[GBB+12]   Anna Gaulton, Louisa J Bellis, A Patricia Bento, Jon Chambers, Mark Davies, Anne Hersey, Yvonne Light, Shaun McGlinchey, David Michalovich, Bissan Al-Lazikani, et al. Chembl: a large-scale bioactivity database for drug discovery. *Nucleic acids research*, 40(D1):D1100–D1107, 2012.

[GDX22]   Zhenxiang Gao, Pingjian Ding, and Rong Xu. Kg-predict: A knowledge graph computational framework for drug repurposing. *Journal of biomedical informatics*, 132:104133, 2022.

[JMM22]   Pratik Joshi, V Masilamani, and Anirban Mukherjee. A knowledge graph embedding based approach to predict the adverse drug reactions using a deep neural network. *Journal of biomedical informatics*, 132:104122, 2022.

[KW16]   Thomas N Kipf and Max Welling. Variational graph auto-encoders. *arXiv preprint arXiv:1611.07308*, 2016.

[KWK+24]   Craig Knox, Mike Wilson, Christen M Klinger, Mark Franklin, Eponine Oler, Alex Wilson, Allison Pon, Jordan Cox, Na Eun Chin, Seth A Strawbridge, et al. Drugbank 6.0: the drugbank knowledgebase for 2024. *Nucleic acids research*, 52(D1):D1265–D1275, 2024.

[MZLK23]   Chunyu Ma, Zhihan Zhou, Han Liu, and David Koslicki. Kgml-xdtd: a knowledge graph–based machine learning framework for drug treatment prediction and mechanism description. *GigaScience*, 12:giad057, 2023.

[SCN+22]   Alberto Santos, Ana R Colaço, Annelaura B Nielsen, Lili Niu, Maximilian Strauss, Philipp E Geyer, Fabian Coscia, Nicolai J Wewer Albrechtsen, Filip Mundt, Lars Juhl Jensen, et al. A knowledge graph to interpret clinical proteomics data. *Nature biotechnology*, 40(5):692–702, 2022.

[SSB+24]   Eduardo Soares, Victor Shirasuna, Emilio Vital Brazil, Renato Cerqueira, Dmitry Zubarev, and Kristin Schmidt. A large encoder-decoder family of foundation models for chemical language. *arXiv preprint arXiv:2407.20267*, 2024.

[WGK+22]   EC Wood, Amy K Glen, Lindsey G Kvarfordt, Finn Womack, Liliana Acevedo, Timothy S Yoon, Chunyu Ma, Veronica Flores, Meghamala Sinha, Yodsawalai Chodpathumwan, et al. Rtx-kg2: a system for building a semantically standardized knowledge graph for translational biomedicine. *BMC bioinformatics*, 23(1):400, 2022.

[WGO+22]   David S Wishart, AnChi Guo, Eponine Oler, Fei Wang, Afia Anjum, Harrison Peters, Raynard Dizon, Zinat Sayeeda, Siyang Tian, Brian L Lee, et al. Hmdb 5.0: the human metabolome database for 2022. *Nucleic acids research*, 50(D1):D622–D631, 2022.

[WMN20]   Brian Walsh, Sameh K Mohamed, and Vít Nováček. Biokg: A knowledge graph for relational learning on biological data. In *Proceedings of the 29th ACM International Conference on Information & Knowledge Management*, pages 3173–3180, 2020.

[WRH+11]   Y Wang, T Romigh, X He, MH Tan, MS Orloff, RH Silverman, WD Heston, and C Eng. Differential regulation of pten expression by androgen receptor in prostate and breast cancers. *Oncogene*, 30(42):4327–4338, 2011.

[YYH+14]   Bishan Yang, Wen-tau Yih, Xiaodong He, Jianfeng Gao, and Li Deng. Embedding entities and relations for learning and inference in knowledge bases. *arXiv preprint arXiv:1412.6575*, 2014.

[ZSD+21]   Fei Zhang, Bo Sun, Xiaolin Diao, Wei Zhao, and Ting Shu. Prediction of adverse drug reactions based on knowledge graph embedding. *BMC Medical Informatics and Decision Making*, 21:1–11, 2021.

[ZWZ+20]   Kangzhi Zhao, Xiting Wang, Yuren Zhang, Li Zhao, Zheng Liu, Chunxiao Xing, and Xing Xie. Leveraging demonstrations for reinforcement recommendation reasoning over knowledge graphs. In *Proceedings of the 43rd international ACM SIGIR conference on research and development in information retrieval*, pages 239–248, 2020.

5   Supplementary Materials

You will be provided a list of tuples, where each tuple's first entry index corresponds to a PubMed article's PMID, and the second entry corresponds to that article's abstract. You must analyse each abstract of each tuple, and for EACH sentence of the abstract, determine if it provides either:

    1) Supporting evidence for a causal relationship between {entity1} and {entity2}
    2) Contradictory evidence against a causal relationship between {entity1} and {entity2}
    3) No clear evidence either way

Take into account that biological entities may have different aliases in different contexts

The PMIDs and corresponding abstracts: "{articles}"

You must return a list of tuples just like the originally provided list of tuples. However, each entry of your output list of tuples should be a PMID and a string that is a shortened version of the PMID's abstract with only sentences that either provide supporting evidence to the causal relationship between {entity1} and {entity2} or contradictory evidence to the causal relationship between {entity1} and {entity2}. If there are no such sentences, you can omit that PMID and abstract tuple from your output list. Do not wrap your output in backticks. This should be a limited selection that includes only sentences that indicate some sort of causal relationship between the two entities. These sentences must be direct quotations from the abstract. Do not output anything other than these direct quotation sentences. If a sentence is unrelated to supporting nor contradictory evidence of a causal relationship, don't include it.

Supplementary Material 1: The multi-agent approach to extract causal information from PubMed abstracts. Each agent analyzes a provided list of PMIDs and corresponding abstracts to extract either supporting or contradictory sentences from each and returns that information.

Analyze the causal relationship between {entity1_name} ({entity1_type}) and {entity2_name} ({entity2_type}) using the provided causal inference frameworks from biology and epidemiology.

The analysis must only consider the causal relationship from {entity1_name} to {entity2_name} and not the other way around. You should take into account the fact that entities have different names, synonyms, or closely related elements in the PubMed articles and adjust accordingly if you see an alias for an entity.

ENTITY 1 {entity1_name}:

ENTITY 2 {entity2_name}:

SCIENTIFIC EVIDENCE: {articles_text}

Return valid JSON with the following structure. You must fill in the blanks and categories and all scores must be justified by evidence:
{{
"counterfactual_statement": "Describe the causal question using potential outcomes (e.g., 'If {entity1_name} were [intervention] vs. [control], how would {entity2_name} change, holding other factors constant?')",

"causal_evidence_framework": {{
    "mechanistic_evidence": {{
        "score":,
        "reasoning": "Evidence of biological mechanism or pathway connecting the entities",
        "citations": [full PMIDs]
    }},
    "association_strength": {{
        "score":,
        "reasoning": "Strength of reported associations in the literature",
        "association_type": "strong/moderate/weak/conflicting/insufficient",
        "citations": [full PMIDs]
    }},
    "biological_plausibility": {{

```
            "score":,
            "reasoning": "How well the relationship fits established biological knowledge",
            "plausibility_level":"well_established/plausible/uncertain/implausible",
            "citations": [full PMIDs]
        }},
        "literature_consistency": {{
            "score":,
            "reasoning": "Consistency of findings across different studies",
            "consistency_level": "consistent/mostly_consistent/mixed/contradictory/insufficient",
            "citations": [full PMIDs]
        }},
        "causal_terminology": {{
            "score":,
            "reasoning": "Presence and strength of causal language in abstracts",
            "causal_terms_found": [],
            "citations": [full PMIDs]
        }}
}},

"causal_relationship_analysis": {{
        "causal_sentence_examples": [], // Here you must describe the causal sentences or lack thereof
            that were important factors to your final score. These must be direct quotations from the
            provided PubMed abstracts. List as many causal sentences as you can that are relevant and
            suggest a causal relationship between the two entities.
        "PMID_citations": [List the full PubMed PMIDs as the citations for the sentences you listed
            above],
        "identified_causal_terms": [],
        "contradictory_evidence": {{
            "contradictory_sentences": [], // List direct quotations from abstracts that provide evidence
                against a causal relationship or suggest an opposing effect.
            "PMID_citations": [List the full PubMed PMIDs as the citations for the contradictory
                sentences above],
            "contradiction_analysis": "Analyze the impact of the contradictory evidence on the overall
                assessment. Discuss how this evidence affects the confidence in a causal relationship, and
                whether it suggests alternative explanations or weakens the proposed mechanism."
        }},
        "term_justification": "Based on the scientific evidence, identify 1-3 causal terms that best
            describe the relationship between the entities. Each term should be supported by specific
            evidence from the provided abstracts. Take into account the official interaction category for
            this pair of entities is biolink_rel. If there is no indicated causal relationship, then say 'Unlikely
            to be causal'",
        "relationship_description": "Describe how the identified causal terms apply to the specific
            relationship between these two entities based on the evidence.",
        "potential_mechanism": ""
        }},
"overall_confidence": {{
        "score": // Give a nuanced score based on the evidence provided and your analysis,
        "score_interpretation": "Explain the score you gave and why you gave it. Remember that this is
            the causal strength score between the two entities. Consider the following guidelines: 1) High
            scores (0.9-1.0) require strong mechanistic evidence and clear causal language in the literature.
            2) Scores 0.65-0.89 indicate likely influence but uncertain causality, often due to indirect
            evidence or mixed findings. 3) Moderate scores (0.5-0.65) suggest plausible relationships with
            limited direct evidence. 4) Low scores (0.2-0.49) indicate weak or indirect associations. 5) Very
            low scores (0-0.19) suggest no meaningful causal relationship. Base your explanation on
            specific evidence from the provided abstracts and mechanistic reasoning.",
        "key_strengths": [],
        "key_weaknesses": []
        }}
}}
```

INSTRUCTIONS:

1. **Evidence Requirements**:
- High scores require there to be a plausible biological pathway based on the evidence provided. You should evaluate the relationship based on the scientific evidence provided and provide a score to judge the causal strength of the relationship accordingly.
- For all citations, you must provide the full PMID of the articles you obtained the evidence from.

2. **Critical Checks**:
- Evaluate contradictory evidence and determine if it should reduce the proportionality score.
- Do not repeat the placeholder instructions - generate original content based on your analysis.
- You must give specific examples for the scores you provide. For example, don't just say something general like "The articles indicate a causal relationship between the two entities" - instead, you must give specific quotes from the abstracts that support your score.
3. **Formatting**:
- Do not include "'json or any other code block markers before or after your JSON output. Return only the JSON object.

Supplementary Material 2: The causal analysis framework used to determine the causal relationship between two entities. The agent is provided the names of two biological entities and the filtered abstract information from Fig X. The agent then determines scores for various metrics based on the provided literature to justify a set of causal terms and a final causal score between the two entities.

Analyze the overall causal pathway from {start_entity.name} ({start_entity.entity_type}) to {end_entity.name} ({end_entity.entity_type}) using PubMed-based evidence.

PATHWAY: {start_entity.name} → {' → '.join(intermediate_names)} → {end_entity.name}

STEP-BY-STEP ANALYSIS SUMMARY:
{path_relationships}

Provide a comprehensive overall pathway analysis in JSON format:
```
{{
"pathway_summary": {{
    "start_entity": {{
        "name": "{start_entity.name}",
        "type": "{start_entity.entity_type}"
      }},
    "intermediate_entities": [{intermediate_json}],
    "end_entity": {{
        "name": "{end_entity.name}",
        "type": "{end_entity.entity_type}"
      }},
    "pathway_length": {len(intermediate_entities) + 1}
    }},
"step_by_step_evidence": {{
    "confidence_scores": {step_scores},
    "supporting_evidence": [
    "List of sentences describing evidence that supports the causal relationship. Each sentence
      should include the PMID, specific findings, type of evidence, and relationship type observed,
      but can follow any natural sentence structure. An example could be 'Study [PMID]
      demonstrated [specific finding] and [additional finding] with [evidence] to support this, and
      observed a [relationship type] relationship between the two entities'."
    ]
    "contradictory_evidence": [
    "List of sentences describing evidence that contradict the causal relationship. Each sentence
      should include the PMID, specific findings, type of evidence, and observations, but can follow
      any natural sentence structure. An example could be 'Study [PMID] demonstrated [specific
      finding] and [additional finding] with [evidence] to contradict that there is a causal relationship
      between the two entities'.
    ]
```

```
        }},
    "evidence_integration": {{
        "evidence_consistency": "high/medium/low - how consistent are findings across studies",
        "mechanistic_evidence": [
        "List the sentences describing mechanistic evidence between the pathways of the entities. Each
          sentence should include the PMID and specific findings but can follow any natural sentence
          structure. Study [PMID] elucidated the mechanism by [specific mechanistic finding] and
          [additional mechanistic detail] with [type of evidence], revealing how the entities interact"
        ],
        "gaps_in_evidence": [
        "Specific gaps where more research is needed, such as [specific gap] which was not addressed in
          the reviewed literature. List here as many gaps as you think would be useful to researchers
          studying this causal relationship."
        ],
        "potential_mechanism": "Describe the potential mechanism of the causal pathway, if any. Do
          not just state a general statement, but rather state a specific biological mechanism of the path
          based on evidence from the literature. Be as quantitative as possible and include details such
          as numbers, sequences, and specific scientific terminology",
        "drug_repurposing_potential": "If the total_confidence_score ≥ 0.5, then using chain-of-thought
          reasoning, describe in great detail a hypothetical experiment that could lead to drug
          repurposing for this pathway. Your response should include: (1) the rationale for selecting a
          candidate drug based on the evidence, (2) the experimental design (including model systems,
          controls, endpoints, and methods), (3) how the experiment would test the drug's effect on the
          pathway, (4) expected outcomes and how they would support or refute repurposing, and (5)
          potential challenges and next steps. Be as specific and technical as possible, integrating
          evidence from the literature and the pathway analysis. Make sure to return this part as a
          single string."
        }},
    "overall_evidence_assessment": {{
        "total_confidence_score": {overall_confidence},
        "confidence_reasoning": "How confident are we in the overall pathway based on individual
          steps?",
        "evidence_quality": "high/medium/low/insufficient"
        }}
}}

INSTRUCTIONS:
1. You must return valid JSON format
2. Do not repeat any of the placeholder instructions - instead, generate original content based on your
analysis.
3. Integrate evidence from all pathway steps
4. Assess biological plausibility of the complete pathway
5. Be honest about evidence limitations and knowledge gaps
6. For supporting evidence and contradictory evidence, you do not need to follow the exact sentence
structure, but you should still provide the critical information, including the PMID's, specific finding,
type of evidence, and relationship type observed.
```

Supplementary Material 3: The global summary of a drug-gene pathway provided by the agent. This includes a summary of the most notable evidence in determining the causal score along with the PMID citations for the evidence. The agent evaluates both supporting and contradictory evidence to determine gaps in evidence that is limiting the score of the pathway. The agent also proposes potential mechanisms and drug repurposing potentials for high-scoring pathways (overall score ≥ 0.5).

```
"association_strength": {
  "score": 0.9,
  "reasoning": "The association between enzalutamide administration and inhibition of
  AR activity is strong and consistently reported across preclinical and clinical studies.
  Enzalutamide is FDA-approved for CRPC based on its ability to inhibit AR signaling,
  reduce PSA levels, and improve survival (PMID: 23701654). Multiple studies show that
  enzalutamide reduces AR activity, cell proliferation, and tumor growth in AR-positive
  models (PMID: 26932782, 28512248, 25713333). However, resistance mechanisms (e.g.,
  AR-V7 upregulation) can attenuate this effect, but the initial association is strong.",
  "association_type": "strong",
  "citations": [
    "23701654",
    "26932782",
    "28512248",
    "25713333",
    "25719830",
    "29572225",
    "30177856"
    ]
  },
"biological_plausibility": {
  "score": 1.0,
  "reasoning": "The relationship is extremely well established in biological and clinical
  literature. Enzalutamide is designed to target AR, and its mechanism of action is to
  inhibit AR signaling. This is the canonical pathway for anti-androgen therapy in prostate
  cancer and is supported by decades of research.",
  "plausibility_level": "well_established",
  "citations": [
    "23701654",
    "25719830",
    "29572225",
    "27765852",
    "26932782"
    ]
  },
"literature_consistency": {
  "score": 0.85,
  "reasoning": "The literature is highly consistent in showing that enzalutamide inhibits
  AR activity and signaling. However, there is consistent reporting of resistance mech-
  anisms (e.g., AR-V7, AR amplification, alternative splicing) that can restore AR sig-
  naling despite enzalutamide treatment (PMID: 25048254, 27486973, 32989253). These
  do not contradict the primary effect but indicate that the relationship can be context-
  dependent.",
  "consistency_level": "mostly_consistent",
  "citations": [
    "23701654",
    "25048254",
    "27486973",
    "32989253",
    "25719830",
    "29572225"
    ]
  },
"causal_terminology": {
  "score": 0.9,
  "reasoning": "Many abstracts use strong causal language such as 'inhibits', 'blocks',
  'antagonizes', 'induces degradation', and 'suppresses' when describing the effect of enza-
  lutamide on AR. For example, PMID: 29572225 states 'enzalutamide-induced AR degra-
  dation and attenuated AR downstream activity', and PMID: 23701654 says 'enzalutamide
  showed...reduction in the serum prostate specific antigen (PSA) level', which is a direct
```

```
              AR target.",
              "causal_terms_found": [
                "inhibits",
                "blocks",
                "antagonizes",
                "induces degradation",
                "suppresses"
                ],
              "citations": [
                "29572225",
                "23701654",
                "27765852",
                "25713333",
                "26932782"
                ]
            }
        },
      "causal_relationship_analysis": {
        "causal_sentence_examples": [
          """AR antagonist enzalutamide-induced AR degradation and attenuated AR downstream
          activity in sunitinib-resistant cells, also indicated by decreased secretion of human
          kallikrein 2." (PMID: 29572225)",
          """Abiraterone and enzalutamide are novel endocrine treatments that abrogate andro-
          gen receptor (AR) signalling in castration-resistant prostate cancer (CRPC)." (PMID:
          25719830)",
          """Enzalutamide is an approved drug that inhibits AR activity and increases overall sur-
          vival." (PMID: 30177856)",
          """Pretreatment with enzalutamide (Enza) decreased the tumor-initiating capacity of
          TNBC cells and reduced tumor volume and viability..." (PMID: 28512248)",
          """Enzalutamide reduced clonogenic potential and cell growth in a 3D matrix in AR-
          positive cells. In addition, enzalutamide also inhibited cell migration and invasion in an
          AR-dependent manner." (PMID: 26932782)",
          """In phase III clinical trial, enzalutamide showed that 160 mg once daily oral adminis-
          tration is well tolerated and significantly enhanced overall survival in men with CRPC
          after chemotherapy, demonstrated by reduction in the serum prostate specific antigen
          (PSA) level and increased survival rate by 4.8 months." (PMID: 23701654)",
          """We show that VPC-13566 inhibits the growth of various prostate cancer cell lines,
          including an enzalutamide-resistant cell line, and reduces the growth of AR-dependent
          prostate cancer xenograft tumors in mice. Furthermore, we used this AR-BF3-directed
          small molecule to demonstrate that inhibition of AR activity through the BF3 function-
          ality can block translocation of the receptor into the nucleus." (PMID: 27765852)"
          ],
        "pmid_citations": [
          "29572225",
          "25719830",
          "30177856",
          "28512248",
          "26932782",
          "23701654",
          "27765852"
          ],
        "identified_causal_terms": [
          "inhibits",
          "blocks",
          "antagonizes",
          "induces degradation"
          ],
        "contradictory_evidence": {
          "contradictory_sentences": [
            """Antiandrogen treatment also altered the expression of multiple enzymes potentially
```

involved in steroid metabolism. Identical to clinical CRPC, the expression levels of the full-length AR (twofold, P ¡ 0.05) and the AR splice variants 1 (threefold, P ¡ 0.05) and 7 (threefold, P ¡ 0.01) were further increased in the antiandrogen-treated tumors." (PMID: 29126837)",

""Enzalutamide resistance was accompanied by increased AR and AR-V7 mRNA and protein expression as well as AR gene amplification, while no additional AR mutations have been identified." (PMID: 27486973)",

""Clinical evidence confirms that resistance to the next-generation antiandrogen, enzalutamide, may be mediated to a large extent by alternative splicing of the androgen receptor to generate constitutively active splice variants such as AR-V7." (PMID: 28729398)",

""Both drugs were designed to further suppress androgen receptor (AR) signaling, which is restored following first-line androgen deprivation therapies. Resistance to enzalutamide and abiraterone, however, is again marked by a return of AR signaling, indicating a remarkable 'addiction' of prostate cancer cells to the AR pathway. Among these, the androgen receptor splice variants (AR-Vs), particularly variant 7 (AR-V7), have been implicated in resistance to enzalutamide and abiraterone in preclinical studies, and they cannot be targeted by currently available AR-directed drugs." (PMID: 25048254)"
],
"pmid_citations": [
"29126837",
"27486973",
"28729398",
"25048254"
],
"contradiction_analysis": "The contradictory evidence does not refute the direct inhibitory effect of enzalutamide on AR, but highlights the development of resistance mechanisms, such as upregulation of AR, AR gene amplification, and especially the emergence of AR splice variants (notably AR-V7) that are not inhibited by enzalutamide. These findings indicate that while enzalutamide causally inhibits AR signaling, cancer cells can adapt by increasing AR expression or producing resistant AR variants, leading to restored AR activity and clinical resistance. This context-dependence slightly reduces the overall confidence in a universal, persistent causal effect but does not negate the primary mechanistic relationship."
},
"term_justification": "The terms 'inhibits', 'antagonizes', and 'induces degradation' best describe the relationship, as supported by direct mechanistic and functional evidence (e.g., enzalutamide-induced AR degradation, inhibition of AR nuclear translocation, and suppression of AR signaling). The official interaction category 'physically_interacts_with' is consistent with enzalutamide binding to AR and blocking its activity. There is overwhelming evidence for a direct inhibitory causal effect, with resistance mechanisms as a secondary phenomenon.",
"relationship_description": "Enzalutamide directly binds to the androgen receptor (AR), antagonizes its activity by preventing nuclear translocation and DNA binding, and can induce AR degradation. This leads to inhibition of AR-dependent transcription and downstream signaling, resulting in reduced proliferation and survival of AR-positive cancer cells. However, resistance mechanisms such as AR amplification and AR-V7 splice variant expression can restore AR signaling despite enzalutamide treatment.",
"potential_mechanism": "Enzalutamide acts as a competitive antagonist at the AR ligand-binding domain, preventing androgen binding, inhibiting AR nuclear translocation, DNA binding, and transcriptional activation of AR target genes. It can also promote AR degradation via the proteasome pathway. Resistance can develop via increased AR expression, gene amplification, or the emergence of AR splice variants (e.g., AR-V7) that lack the ligand-binding domain and are not inhibited by enzalutamide."
},
"overall_confidence": {
"score": 0.92,
"score_interpretation": "A score of 0.92 reflects very high confidence in a direct, causal, and mechanistically well-understood inhibitory effect of enzalutamide on AR. This is supported

by strong mechanistic evidence, consistent and strong associations, well-established biological plausibility, and frequent use of causal terminology in the literature. The only limitation is the context-dependent emergence of resistance mechanisms (e.g., AR-V7), which can attenuate the effect in some settings but do not negate the primary causal relationship.",
            "key_strengths": [
               "Robust mechanistic evidence for direct inhibition of AR by enzalutamide",
               "Consistent findings across preclinical, clinical, and mechanistic studies",
               "Strong causal language and functional outcomes (e.g., reduced AR activity, tumor growth inhibition)",
               "FDA approval and clinical efficacy based on AR inhibition"
               ],
            "key_weaknesses": [
               "Resistance mechanisms (AR-V7, AR amplification) can restore AR signaling despite enzalutamide treatment",
               "Some studies report increased AR or AR-V7 expression after prolonged enzalutamide exposure, indicating adaptive feedback"
               ]
            }
         }
      },
   {
      "relationship": {
         "cause": {
            "name": "AR",
            "type": "Gene"
            },
         "effect": {
            "name": "PTEN",
            "type": "Gene"
            }
         },
      "relation_analysis": {
         "counterfactual_statement": "If AR (androgen receptor) expression or activity were increased versus decreased (or inactivated), how would PTEN (phosphatase and tensin homolog) expression or activity change, holding other factors constant?",
         "causal_evidence_framework": {
            "mechanistic_evidence": {
               "score": 0.8,
               "reasoning": "There is direct evidence that AR can regulate PTEN transcriptionally, with fine-mapping of AR-binding motifs within the PTEN promoter in prostate and breast cancer cells. Specifically, AR inhibits PTEN transcription in prostate cancer cells (PMID: 21532617). Additional studies show that AR and PTEN physically interact, with AR activity being suppressed by PTEN, but also that AR can directly affect PTEN levels. In breast cancer, AR upregulates PTEN, but in prostate cancer, AR downregulates PTEN. This organ-specific regulation is mechanistically supported by promoter binding and transcriptional regulation. There is also evidence that AR inactivation modifies PTEN deletion-induced pathology in uterine tissue (PMIDs: 21532617, 26285813, 28439009, 23418309).",
               "citations": [
                  "21532617",
                  "26285813",
                  "28439009",
                  "23418309"
                  ]
               },
            "association_strength": {
               "score": 0.7,
               "reasoning": "Multiple studies report an inverse correlation between AR and PTEN transcript expression in prostate cancer (PMID: 21532617), and that AR inactivation reduces PTEN deletion-induced pathology (PMID: 26285813). There are also studies showing

mechanistically direct, involving promoter binding and transcriptional regulation. There is also evidence of physical interaction between AR and PTEN proteins, supporting the 'physically_interacts_with' category.",
"potential_mechanism": "AR, as a nuclear hormone receptor and transcription factor, binds to specific motifs in the PTEN promoter region, leading to repression of PTEN transcription in prostate cancer cells. This may involve recruitment of co-repressors or chromatin remodeling factors. In breast cancer, AR binding to the PTEN promoter leads to transcriptional activation, possibly via different co-regulators or chromatin context. The direct protein-protein interaction between AR and PTEN may also modulate AR nuclear translocation and stability, further influencing PTEN expression."
},
"overall_confidence": {
"score": 0.8,
"score_interpretation": "The evidence supports a strong, mechanistically plausible, and mostly consistent causal relationship from AR to PTEN, particularly in prostate and breast cancer contexts. The presence of direct mechanistic evidence (promoter binding, transcriptional regulation), strong causal language, and multiple studies reporting the relationship justifies a high score. The main limitation is the context-dependent (tissue-specific) directionality and some correlative findings, but these do not undermine the existence of a causal effect. Therefore, a score of 0.8 is appropriate.",
"key_strengths": [
"Direct mechanistic evidence of AR binding to PTEN promoter and regulating transcription",
"Strong causal language in multiple abstracts",
"Consistent findings in prostate and breast cancer models",
"Evidence of physical interaction between AR and PTEN proteins"
],
"key_weaknesses": [
"Tissue-specific directionality (inhibition in prostate, activation in breast)",
"Some findings are correlative rather than strictly causal",
"Potential co-regulation or influence by other factors in certain pathological states"
]
}
}
},
{
"relationship": {
"cause": {
"name": "PTEN",
"type": "Gene"
},
"effect": {
"name": "prostate cancer",
"type": "Disease"
}
},
"relation_analysis": {
"counterfactual_statement": "If PTEN were inactivated or deleted (intervention) versus retained and functional (control), how would the incidence, progression, and aggressiveness of prostate cancer change, holding other genetic and environmental factors constant?",
"causal_evidence_framework": {
"mechanistic_evidence": {
"score": 1.0,
"reasoning": "There is overwhelming evidence for a direct mechanistic link between PTEN loss and prostate cancer development and progression. PTEN is a tumor suppressor that negatively regulates the PI3K/AKT/mTOR pathway, and its loss leads to increased cell proliferation, survival, and tumorigenesis. Multiple studies using mouse models with prostate-specific PTEN deletion show that loss of PTEN alone is sufficient to induce prostate intraepithelial neoplasia and invasive carcinoma. Additional studies demonstrate that PTEN loss cooperates with other genetic alterations (e.g., TP53, ERG,

SOX9, RUNX2, etc.) to accelerate tumor progression, metastasis, and therapy resistance. Mechanistic details include activation of AKT, mTORC1/2, metabolic reprogramming, immune evasion, and changes in cell cycle regulation.",
    "citations": [
      "29309058",
      "16079851",
      "19185849",
      "21840483",
      "33186519",
      "19399032",
      "11175773",
      "11175795",
      "23727861",
      "19396168",
      "29335545",
      "28255082",
      "32385075",
      "24866151",
      "22777769",
      "29581176",
      "23902739",
      "23670050",
      "9661880",
      "15994948",
      "10363971",
      "17616663",
      "15930280",
      "24480624",
      "29706654",
      "29765153",
      "28402859",
      "25176644",
      "25818291",
      "31213464",
      "21556061",
      "29142193",
      "31551414",
      "30733438",
      "26975529",
      "26379078",
      "33155366",
      "35773413",
      "26028029"
    ]
  },
  "association_strength": {
    "score": 0.95,
    "reasoning": "The association between PTEN loss and prostate cancer is strong and well-documented. PTEN mutations, deletions, or reduced expression are frequently observed in prostate cancer cell lines, primary tumors, and metastatic lesions. PTEN loss is associated with higher Gleason scores, advanced stage, increased risk of recurrence, metastasis, and poor prognosis. Quantitative associations include hazard ratios for recurrence and metastasis (e.g., HR 2.25 and 3.90 in PMID: 27477529). The frequency of PTEN loss in prostate cancer is high (up to 60"association_type": "strong",
    "citations": [
      "9661880",
      "10485474",
      "27477529",
      "15994948",
      "26379078",

```
      "causal_terms_found": [
        "induces",
        "drives",
        "triggers",
        "promotes",
        "required for",
        "leads to",
        "caused by",
        "initiates",
        "suppresses",
        "prevents"
        ],
      "citations": [
        "16079851",
        "19185849",
        "11175795",
        "23670050",
        "10363971",
        "33105713",
        "30733438",
        "26379078",
        "27477529",
        "22674438",
        "29309058",
        "21840483",
        "33186519",
        "19399032",
        "23727861",
        "19396168",
        "29335545",
        "28255082",
        "32385075",
        "24866151",
        "22777769",
        "29581176",
        "23902739"
        ]
      }
    },
  "causal_relationship_analysis": {
    "causal_sentence_examples": [
      "Complete Pten inactivation in the prostate triggers non-lethal invasive prostate cancer
      after long latency.",
      "We find that Rictor is a haploinsufficient gene and that deleting one copy protects
      Pten heterozygous mice from prostate cancer. Finally, we show that the development
      of prostate cancer caused by Pten deletion specifically in prostate epithelium requires
      mTORC2...",
      "Inactivation of the tumor-suppressor gene PTEN and lack of p27(KIP1) expression have
      been detected in most advanced prostate cancers. PTEN activity leads to the induction
      of p27(KIP1) expression, which in turn can negatively regulate the transition through
      the cell cycle. Here we show that the concomitant inactivation of one Pten allele and one
      or both Cdkn1b alleles accelerates spontaneous neoplastic transformation and incidence
      of tumors of various histological origins. Moreover, Pten(+/-)/Cdkn1b(-/-) mice develop
      prostate carcinoma at complete penetrance within three months from birth.",
      "Prostate-specific inactivation of Zbtb7a leads to a marked acceleration of Pten loss-
      driven prostate tumorigenesis through bypass of Pten loss-induced cellular senescence
      (PICS).",
      "We find that prostate cancer specimens containing the TMPRSS2-ERG rearrangement
      are significantly enriched for loss of the tumor suppressor PTEN. In concordance with
      these findings, transgenic overexpression of ERG in mouse prostate tissue promotes
```

marked acceleration and progression of high-grade prostatic intraepithelial neoplasia (HG-PIN) to prostatic adenocarcinoma in a Pten heterozygous background.",

"Conditional disruption of Pten in the mouse prostate leads to tumorigenesis and increased phosphorylation of PTK6 Y342, and disruption of Ptk6 impairs tumorigenesis. In human prostate tumor tissue microarrays, loss of PTEN correlates with increased PTK6 PY342 and poor outcome. These data suggest PTK6 activation promotes invasive prostate cancer induced by PTEN loss.",

"Loss of the tumor suppressor PTEN (phosphatase and tensin homolog) is frequently observed in cancer, resulting in the deregulation of cell survival, growth, and proliferation. We hypothesize that loss of PTEN and subsequent activation of Akt, frequent occurrences in prostate cancer, regulate the CXCL12/CXCR4 signaling axis in tumor growth and bone metastasis.",

"Loss of the tumor suppressor phosphatase and tensin homolog (PTEN) occurs frequently in prostate cancers. Preclinical evidence suggests that activation of PI3K/AKT signaling through loss of PTEN can result in resistance to hormonal treatment in prostate cancer.",

"Genetic inactivation of PTEN through either gene deletion or mutation is common in metastatic prostate cancer, leading to activation of the phosphoinositide 3-kinase (PI3K-AKT) pathway, which is associated with poor clinical outcomes.",

"Loss of chromosome 10q is a frequently observed genetic defect in prostate cancer. Recently, the PTEN/MMAC1 tumor suppressor gene was identified and mapped to chromosome 10q23.3. In five samples (PC3, PC133, PCEW, PC295, and PC324), homozygous deletions of the PTEN gene or parts of the gene were detected. The high frequency (60"PTEN is frequently mutated or deleted in both prostate cancer cell lines and primary prostate cancers. Thus, loss of PTEN protein is correlated with pathological markers of poor prognosis in prostate cancer.",

"Loss of PTEN expression in paraffin-embedded primary prostate cancer correlates with high Gleason score and advanced stage.",

"Loss of PTEN expression is an important factor in progression towards metastatic disease and could potentially serve as an early prognostic marker for prostate cancer metastasis."
],
"pmid_citations": [
"16079851",
"19185849",
"11175795",
"23727861",
"19396168",
"29142193",
"23902739",
"25454616",
"23670050",
"9661880",
"10485474",
"17163422"
],
"identified_causal_terms": [
"induces",
"drives",
"triggers",
"promotes",
"required for",
"leads to",
"caused by",
"initiates",
"suppresses",
"prevents"
],
"contradictory_evidence": {
"contradictory_sentences": [
"Our results suggest either that mutation of PTEN is a late event in prostate tumorigenesis, or that another tumour suppressor gene important in prostate cancer may lie

close to PTEN in 10q23.",
"Our results showing a low frequency of alterations of PTEN/MMAC1 in pT2 and pT3 prostate cancers suggest that this gene plays an insignificant role in the development of most low stage carcinomas of the prostate."
],
"pmid_citations": [
"9582022",
"9788441"
],
"contradiction_analysis": "These sentences suggest that PTEN mutation may not be the initiating event in all prostate cancers, or that its role may be less significant in early-stage or low-grade tumors. However, the bulk of evidence indicates that PTEN loss is a major driver in progression, aggressiveness, and poor prognosis, especially in advanced and metastatic disease. The contradictory evidence does not negate the causal role of PTEN loss but suggests that its impact may be context-dependent and that other tumor suppressors may also contribute, particularly in early tumorigenesis."
},
"term_justification": "The terms 'induces', 'drives', and 'promotes' best describe the relationship, as supported by multiple studies showing that PTEN loss is sufficient to induce prostate cancer in mouse models, drives progression and metastasis, and promotes therapy resistance and poor outcomes in human disease. The official interaction category 'gene_associated_with_condition' is supported, but the evidence here is much stronger and supports a direct causal role.",
"relationship_description": "PTEN loss or inactivation causally induces prostate cancer initiation, drives progression to more aggressive and metastatic disease, and promotes resistance to therapy. This is mediated through well-established mechanisms including activation of the PI3K/AKT/mTOR pathway, increased cell proliferation, survival, metabolic reprogramming, immune evasion, and cooperation with other oncogenic events. The relationship is robustly supported by genetic, molecular, animal model, and clinical outcome data.",
"potential_mechanism": "PTEN is a lipid phosphatase that antagonizes the PI3K/AKT/mTOR signaling pathway. Loss of PTEN function leads to constitutive activation of AKT, resulting in increased cell proliferation, survival, metabolic changes (e.g., Warburg effect), resistance to apoptosis, and enhanced metastatic potential. PTEN loss also cooperates with other genetic alterations (e.g., TP53, ERG, SOX9, RUNX2) to further accelerate tumorigenesis and progression. Additional mechanisms include modulation of the tumor microenvironment, immune evasion (e.g., PD-L1 upregulation), and therapy resistance."
},
"overall_confidence": {
"score": 0.98,
"score_interpretation": "The evidence for a causal relationship from PTEN loss to prostate cancer is extremely strong. There is direct mechanistic evidence from animal models, strong and consistent associations in human studies, well-established biological plausibility, and widespread use of causal terminology in the literature. The only minor weaknesses are context-dependence in early-stage disease and the possibility of other tumor suppressors contributing in some cases. However, these do not substantially detract from the overwhelming evidence that PTEN loss is a major causal driver of prostate cancer initiation, progression, and poor prognosis.",
"key_strengths": [
"Direct mechanistic evidence from genetically engineered mouse models showing PTEN loss induces prostate cancer.",
"Strong, consistent association between PTEN loss and advanced/aggressive prostate cancer in human studies.",
"Well-established biological pathway (PI3K/AKT/mTOR) linking PTEN loss to oncogenesis.",
"Causal language and experimental interventions demonstrating sufficiency and necessity.",
"Evidence for cooperation with other oncogenic events, explaining heterogeneity in disease."
],

```
                "key_weaknesses": [
                    "Some evidence suggests PTEN mutation may be a late event or less significant in
                    early/low-grade tumors.",
                    "Other tumor suppressors may also contribute to prostate cancer initiation in some con-
                    texts.",
                    "Heterogeneity in PTEN loss frequency across populations and tumor stages."
                    ]
                }
            }
        }
    ],
    "overall_pathway_analysis": {
        "pathway_summary": {
            "start_entity": {
                "name": "enzalutamide",
                "type": "SmallMolecule"
            },
            "intermediate_entities": [
                {
                    "name": "AR",
                    "type": "Gene"
                },
                {
                    "name": "PTEN",
                    "type": "Gene"
                }
            ],
            "end_entity": {
                "name": "prostate cancer",
                "type": "Disease"
            },
            "pathway_length": 3
        },
        "step_by_step_evidence": {
            "confidence_scores": {
                "enzalutamide -¿ AR": 0.92,
                "AR -¿ PTEN": 0.8,
                "PTEN -¿ prostate cancer": 0.98
            },
            "supporting_evidence": [
                "Study 29572225 demonstrated that enzalutamide-induced AR degradation and attenuated
                AR downstream activity in sunitinib-resistant cells, with mechanistic evidence showing a
                direct inhibitory relationship between enzalutamide and AR.",
                "Study 25719830 reported that enzalutamide abrogates androgen receptor (AR) signalling in
                castration-resistant prostate cancer (CRPC), providing clinical and mechanistic evidence for
                a strong inhibitory effect.",
                "Study 30177856 found that enzalutamide inhibits AR activity and increases overall survival
                in clinical settings, supporting a direct causal relationship.",
                "Study 28512248 showed that pretreatment with enzalutamide decreased tumor-initiating ca-
                pacity and reduced tumor volume and viability in AR-positive models, indicating a functional
                inhibitory relationship.",
                "Study 26932782 observed that enzalutamide reduced clonogenic potential and cell growth
                in AR-positive cells, and inhibited cell migration and invasion in an AR-dependent manner,
                supporting a strong mechanistic link.",
                "Study 23701654 reported that enzalutamide administration led to a reduction in serum PSA
                levels and increased survival in men with CRPC, confirming clinical efficacy via AR inhibi-
                tion.",
                "Study 27765852 demonstrated that inhibition of AR activity through the BF3 functionality
                can block translocation of the receptor into the nucleus, providing mechanistic evidence for
                enzalutamide's action.",
```

"Study 21532617 showed that AR inhibits PTEN transcription in prostate cancer cells, with fine-mapping of AR-binding motifs within the PTEN promoter, providing direct mechanistic evidence for AR's negative regulation of PTEN.",

"Study 23418309 found that AR upregulates PTEN transcription in breast cancer cells, but in prostate cancer, AR acts as a repressor, highlighting tissue-specific regulation with mechanistic promoter binding evidence.",

"Study 26285813 demonstrated that AR inactivation mediated protection against PTEN knockout-induced uterine pathology, indicating a functional interaction between AR and PTEN in vivo.",

"Study 28439009 reported that ERβ down-regulates AR signaling and up-regulates PTEN, supporting the regulatory axis between AR and PTEN.",

"Study 16079851 showed that complete Pten inactivation in the prostate triggers non-lethal invasive prostate cancer after long latency in mouse models, providing direct genetic evidence for PTEN's tumor suppressor role.",

"Study 19185849 found that deleting one copy of Rictor protects Pten heterozygous mice from prostate cancer, and that prostate cancer caused by Pten deletion requires mTORC2, supporting mechanistic links between PTEN loss and tumorigenesis.",

"Study 11175795 demonstrated that concomitant inactivation of one Pten allele and one or both Cdkn1b alleles accelerates spontaneous neoplastic transformation and incidence of prostate carcinoma, providing genetic evidence for PTEN's causal role.",

"Study 23727861 reported that prostate-specific inactivation of Zbtb7a accelerates Pten loss-driven prostate tumorigenesis, supporting the sufficiency of PTEN loss for cancer development.",

"Study 19396168 found that ERG overexpression in mouse prostate tissue promotes progression of high-grade prostatic intraepithelial neoplasia to adenocarcinoma in a Pten heterozygous background, indicating cooperation between PTEN loss and other oncogenic events.",

"Study 23902739 showed that conditional disruption of Pten in the mouse prostate leads to tumorigenesis and increased phosphorylation of PTK6 Y342, with loss of PTEN correlating with poor outcome in human prostate tumors.",

"Study 23670050 demonstrated that loss of PTEN and subsequent activation of Akt regulate the CXCL12/CXCR4 signaling axis in tumor growth and bone metastasis, providing mechanistic evidence for PTEN's role in cancer progression.",

"Study 25454616 found that PTEN loss can result in resistance to hormonal treatment in prostate cancer, supporting its role in therapy resistance.",

"Study 9661880 identified a high frequency (60"Study 10485474 and 17163422 reported that PTEN is frequently mutated or deleted in prostate cancer cell lines and primary tumors, with loss of PTEN protein correlated with poor prognosis.",

"Study 27477529 quantified that PTEN loss is associated with higher Gleason scores and increased risk of recurrence and metastasis, providing strong clinical association evidence."
],
"contradictory_evidence": [
"Study 29126837 found that antiandrogen treatment increased expression levels of full-length AR and AR splice variants in antiandrogen-treated tumors, suggesting adaptive resistance mechanisms that can restore AR signaling despite enzalutamide treatment.",

"Study 27486973 reported that enzalutamide resistance was accompanied by increased AR and AR-V7 mRNA and protein expression as well as AR gene amplification, indicating that resistance mechanisms can attenuate the inhibitory effect of enzalutamide on AR.",

"Study 28729398 and 25048254 described that resistance to enzalutamide may be mediated by alternative splicing of AR to generate constitutively active variants such as AR-V7, which are not inhibited by current AR-directed drugs.",

"Study 21532617 observed an inverse correlation between AR and PTEN transcript expression in prostate cancer tissues, but a positive correlation in breast cancer, highlighting tissue-specific directionality and context dependence.",

"Study 33920045 found decreased PTEN and AR gene and protein expression in proliferative inflammatory atrophy (PIA) compared to normal samples, suggesting co-regulation or influence by other factors in certain pathological states.",

"Study 9582022 suggested that mutation of PTEN may be a late event in prostate tumorigenesis or that another tumor suppressor gene near PTEN may be important in prostate cancer.",

"Study 9788441 reported a low frequency of PTEN alterations in pT2 and pT3 prostate

cancers, suggesting PTEN may play a less significant role in early/low-stage carcinomas."
            ]
        },
    "evidence_integration": {
        "evidence_consistency": "high",
        "mechanistic_evidence": [
            "Study 29572225 elucidated the mechanism by which enzalutamide induces AR degradation and attenuates AR downstream activity, providing direct evidence of enzalutamide's antagonistic effect on AR.",
            "Study 27765852 demonstrated that inhibition of AR activity through the BF3 functionality blocks AR nuclear translocation, revealing the molecular basis for enzalutamide's inhibition of AR signaling.",
            "Study 21532617 mapped AR-binding motifs within the PTEN promoter and showed that AR inhibits PTEN transcription in prostate cancer cells, providing mechanistic insight into AR's regulation of PTEN.",
            "Study 23418309 showed that AR upregulates PTEN transcription in breast cancer cells via promoter activation, but in prostate cancer, AR acts as a repressor, highlighting the context-dependent mechanistic regulation.",
            "Study 16079851 used genetically engineered mouse models to show that complete Pten inactivation in the prostate triggers invasive prostate cancer, directly linking PTEN loss to tumorigenesis.",
            "Study 19185849 demonstrated that the development of prostate cancer caused by Pten deletion requires mTORC2, providing mechanistic detail on downstream signaling pathways.",
            "Study 23670050 revealed that loss of PTEN and subsequent activation of Akt regulate the CXCL12/CXCR4 signaling axis, contributing to tumor growth and metastasis.",
            "Study 25454616 provided evidence that PTEN loss leads to activation of PI3K/AKT signaling, resulting in resistance to hormonal treatment in prostate cancer."
            ],
        "gaps_in_evidence": [
            "Limited direct evidence for the quantitative impact of enzalutamide-induced AR inhibition on PTEN expression in prostate cancer cells in vivo.",
            "Insufficient studies directly linking changes in PTEN expression following AR inhibition to downstream effects on prostate cancer progression in clinical settings.",
            "Context-dependent regulation of PTEN by AR (inhibition in prostate, activation in breast) is not fully understood at the molecular level, especially regarding co-regulators and chromatin context.",
            "Mechanistic details of how resistance mechanisms (e.g., AR-V7 splice variants) affect the AR-PTEN axis and subsequent impact on prostate cancer progression are not fully elucidated.",
            "Potential compensatory pathways that may bypass PTEN loss or AR inhibition in prostate cancer are not comprehensively mapped."
            ],
        "potential_mechanism": "Enzalutamide acts as a competitive antagonist at the androgen receptor (AR) ligand-binding domain, preventing androgen binding and inhibiting AR nuclear translocation, DNA binding, and transcriptional activation of AR target genes. In prostate cancer cells, AR functions as a transcriptional repressor of PTEN by binding to specific motifs in the PTEN promoter, leading to decreased PTEN expression. Loss or inhibition of AR activity (e.g., by enzalutamide) may relieve this repression, potentially increasing PTEN expression, though this effect is context-dependent and may be attenuated by resistance mechanisms. PTEN is a lipid phosphatase that antagonizes the PI3K/AKT/mTOR signaling pathway; its loss leads to constitutive activation of AKT, promoting cell proliferation, survival, metabolic reprogramming, and tumorigenesis. The pathway thus links enzalutamide-mediated AR inhibition to modulation of PTEN expression and downstream suppression of prostate cancer progression, though resistance mechanisms and tissue-specific regulation may alter these effects.",
        "drug_repurposing_potential": "Given the high total confidence score (0.9) and robust mechanistic evidence linking enzalutamide to AR inhibition, AR to PTEN regulation, and PTEN loss to prostate cancer progression, a rational drug repurposing experiment could focus on enhancing PTEN expression or function in prostate cancer, particularly in cases with AR-driven PTEN repression. Rationale: Since enzalutamide inhibits AR and AR represses PTEN in prostate cancer, combining enzalutamide with a PTEN-activating agent could synergistically suppress tumor growth. Experimental design: Use AR-positive, PTEN-intact and PTEN-deficient prostate can-

Supplementary Material 4: Bioscientist full path analysis example (Enzalutamide $\rightarrow$ AR $\rightarrow$ PTEN$\rightarrow$ prostate cancer). For each entity-entity relationship, the agent analysis causal sentences from the provided PubMed abstracts and scores the relationship on various causal metrics, providing direct quotatoins and citations to justify its score. The agent also extracts important causal sentences that were important in the determination of the final score of the relationship. The path analyses are summarized in the global summary, which includes key supporting and contradictory evidence, evidence integration, gaps in evidence, potential mechanisms, and drug repurposing potential for high scoring paths.

a

Distribution of Node Categories (Top 10)

- biolink:SmallMolecule 41.4%
- biolink:OrganismTaxon 20.2%
- biolink:Gene 5.5%
- biolink:ChemicalEntity 4.6%
- biolink:Protein 4.3%
- biolink:Transcript 3.8%
- biolink:MolecularActivity 3.6%
- biolink:Drug 2.0%
- biolink:AnatomicalEntity 1.9%
- biolink:MolecularMixture 1.9%
- others 8.5%

Distribution of Edge Categories (Top 10)

- biolink:has_participant 13.9%
- biolink:physically_interacts_with 12.4%
- biolink:occurs_in 11.1%
- biolink:affects 10.7%
- biolink:related_to 9.2%
- biolink:subclass_of 8.4%
- biolink:located_in 6.7%
- biolink:coexists_with 4.6%
- biolink:gene_associated_with_condition 3.6%
- biolink:has_part 3.6%
- others 15.9%

b

Log Number of Connections Between Categories

a **Graph training module**

RTX-KG2 Database

VRGCN Encoder

LLM based embedding

μ

ε

σ

Node embedding

Edge embedding

DistMult Decoder

$\otimes$ $\otimes$ = Σ

Genetic network

Molecular network

Organism taxon

Population genetics and Diseases

Cellular network

Organ networks

b **ADAC training module**

Policy

TD error

State → Critic → Actor

reward

Action

Meta-path discriminator

+

Path discriminator

Environment

c **Agent interperter**

Perception

inputs

Drug A -> Gene G -> Bio-Process T -> Disease D

Drug A -> Gene N, Gene T -> Disease D

Analysis all this pathway from Drug to disease, which one is the most posiable path for drug-disease, why?

Brain

Filter causal sentences

Causal framework

PubMed quotes and cite

Evidence analysis and gaps in evidence

Drug repurposing strategies

I will show these results: Analysis of relationships in pathways; Target analysis and drug repurposing strategies; Extended evidence analysis; Extended mechanism analysis; Summary, confidence reasoning, evidence gaps, and so on.

Action

1. Node summary and mechanism
2. Evidence between nodes
3. Evaluate evidence strength
4. Global summary:
5. Global conclusion
6. Global logical chain
7. Global evidence strength