A mechanistic framework for the recognition of chemically diverse brassinosteroids by BRI1-

family receptor kinases.

Alberto Caregnato[1,*], Houming Chen[1,*], Miroslav Kvasnica[2], Ulrich Hohmann[1,3], Jana Oklestkova[2],

Karoll Ferrer[2], Larissa Broger[1], Ludwig A. Hothorn[4], Miroslav Strnad[2], Michael Hothorn[1,#]

[1]Structural Plant Biology Laboratory, Department of Plant Science, University of Geneva, 1211

Geneva, Switzerland

[2]Laboratory of Growth Regulators, Faculty of Science, Palacký University & Institute of

Experimental Botany, The Czech Academy of Sciences, CZ-77900, Olomouc, Czech Republic

[3]Present address: Institute of Molecular Biotechnology of the Austrian Academy of Sciences

(IMBA) & Research Institute of Molecular Pathology (IMP), Vienna BioCenter (VBC), 1030,

Vienna, Austria

[4]retired from Leibniz University Hannover, Germany, present address: Im Grund 12, 31867

Lauenau, Germany

[*]These authors contributed equally to this work.

[#]Correspondence: michael.hothorn@unige.ch

Abstract

Brassinosteroids (BRs) are chemically diverse plant steroid hormones produced via a branched

biosynthetic pathway. The potent BR brassinolide is sensed by the membrane receptor kinase BRI1

and a SERK co-receptor, but the physiological functions of other abundant BRs remain to be

characterized. Here, we present quantitative binding kinetics for four *Arabidopsis thaliana* BR

receptors and fifteen BRs, which define the key chemical features required for high affinity receptor

binding, ligand positioning, and co-receptor recognition. BRI1, BRL1, and BRL3 share overlapping

ligand preferences, whereas BRL2 preferentially binds C_{28} BRs with moderate affinity. Structural

analyses of BR-bound BRI1 and BRL3 ectodomains, combined with extensive in vitro and in vivo

mutagenesis studies reveal a high structural plasticity of the hormone binding pocket. Functional

assays using structure-based BR agonists and antagonists uncover that BR receptor - co-receptor

signaling complexes can recognize chemically diverse BRs, introducing an additional, intriguing

layer of BR signaling regulation.

(149 words)

 Introduction

Brassinosteroids (BRs) are a class of lipophilic, polyhydroxylated steroid hormones first
isolated from rapeseed pollen and later recognized for their growth-promoting roles in plants[1]. BRs
are synthesized via three main biosynthetic branches, producing C_{27}, C_{28} or C_{29} compounds. While
all share a steroidal ring core, they differ in the structure of their aliphatic side chains[2] (Fig. 1a). The
C_{27} branch, derived from cholesterol, yields 28-norbrassinolide[3,4], whereas the C_{29} branch, from
sitosterol, produces 28-homocastasterone and 28-homobrassinolide (Fig. 1b). The C_{28} pathway
originates from campesterol and leads to brassinolide, one of the most bioactive BRs[2] (Fig. 1b). The
C_{28} pathway includes two routes: early and late C_{22} oxidation (Fig. 1b). In the early pathway,
DWARF4 (DWF4), a cytochrome P450 monooxygenase, converts campesterol to (22*S*)-22-
hydroxycampesterol[3,5,6]. Further steps involve the P450 enzymes CPD[7,8] and ROT3[9], and the steroid-
5α-reductase DET2[10–12], producing 6-deoxocastasterone (Fig. 1b). The enzymes BR6ox1 and
BR6ox2 then convert this intermediate to castasterone[13], and BR6ox2 can further catalyze formation
of a seven-membered lactone ring from the C_6 ketone, yielding brassinolide[13] (Fig. 1b).

In the late C_{22} oxidation pathway, campestanol is processed into cathasterone and
typhasterol[14] (Fig. 1b). BR6ox2 can convert typhasterol into 2-deoxybrassinolide[15], or alternatively,
typhasterol may be converted to castasterone via secasterone, a 2,3-epoxy intermediate[16] (Fig. 1b).
The different branches of BR biosynthesis are likely interconnected[2]. BR homeostasis is regulated
by several catabolic enzymes, including the P450 enzyme BAS1[17], the UDP-glycosyltransferase
UGT73C5[18], and sulfotransferases[19].

BR biosynthesis occurs, at least in part, within the endoplasmic reticulum[20,21]. Local BR
transport is facilitated via plasmodesmata[22]. The ATP-binding cassette transporters ABCB1 and
ABCB19 are responsible for translocating BRs across the plasma membrane from the cytosol into
the apoplast[23], where BR perception takes place[24].

Brassinosteroids (BRs) are perceived by the plasma membrane-localized receptor kinases
BRASSINOSTEROID INSENSITIVE 1 (BRI1)[24–28], BRL1[29–31] and BRL3[29], which share a
conserved extracellular leucine-rich repeat (LRR) ligand binding domain. BRI1 is broadly
expressed, and *bri1* loss-of-function mutants display a characteristic dwarf phenotype[25]. BRL1 and
BRL3 are present in vascular tissues and can rescue the growth defects of the weak *bri1-301*
allele[32], when expressed from the *BRI1* promoter[29]. Expression of the sequence-related BRL2 could
not rescue the *bri1-301* mutant phenotype[29], or a *bri1 brl1 brl3* triple mutant[33]. BRL2 is also known
as VASCULAR HIGHWAY1 (VH1) and has been implicated in leaf vein patterning[34].

BRs bind to a steroid-binding pocket formed by the LRR core and a ~70 amino acid island domain within BRI1 or BRL1[26,27,31]. BR binding enables BRI1 to recruit SOMATIC EMBRYOGENESIS RECEPTOR KINASEs (SERKs) to form an active signaling complex[26,35,36]. In the absence of BRs, SERKs are sequestered by BIR pseudokinases[37,38]. Assembly of the BRI1-BR-SERK complex activates the cytoplasmic kinase domains[39], triggering a downstream signaling cascade that leads to dephosphorylation of the BR transcription factors BES1 and BZR1[40,41].

Bioassays with synthetic, exogenously applied brassinosteroids (BRs) have been performed across various plant species. In Arabidopsis, nanomolar levels of brassinolide promote hypocotyl elongation in both wild-type and *det2* mutant seedlings[42]. Brassinolide or 24-epibrassinolide applied at high nanomolar to micromolar concentrations inhibits root growth[43], whereas low concentrations - high picomolar to low nanomolar - of 24-epibrassinolide or 24-epicastasterone stimulate root growth[44].

Initial BR-binding assays used microsomal fractions from transgenic Arabidopsis expressing BRI1-GFP and [³H]-brassinolide, yielding estimated dissociation constants (K_d) of ~10-15 nM for brassinolide and 40-75 nM for castasterone[24]. BRL1 and BRL3 bound brassinolide with K_d values of ~5 nM and ~50 nM, respectively, while BRL2 showed no detectable binding[29]. Grating-coupled interferometry (GCI) quantified binding of unlabeled brassinolide to BRI1 with a K_d of ~10 nM[28]. Brassinolide-bound BRI1 enables the binding of the SERK3/BAK1 LRR ectodomain, with K_d values of ~0.2-1 μM, as determined by GCI and isothermal titration calorimetry[28,36]. Mutation of Gly644→Asp (*bri1-6*/*bri1-119*)[45] in the BR island domain[26] reduces BR binding in vitro[28,46]. The corresponding *bri1-6* allele exhibits intermediate growth defects[45]. Mutation of neighboring Gly643→Glu in the gain-of-function BRI1[sud1] allele[47], enhances the structural stability of the island domain[26]. Notably, five simultaneous point mutations in the BR-binding pocket were required to completely abolish brassinolide binding in vitro, and to mimic the severe growth phenotype of *bri1* null alleles in vivo[46].

The natural occurrence of different BRs has been characterized using high-performance liquid chromatography coupled with gas chromatography-mass spectrometry. In Arabidopsis shoots, castasterone, 6-deoxocastasterone, and typhasterol were identified[11,48,49]. Like in other species, brassinolide accumulates more prominently in Arabidopsis seeds[50].

Brassinolide, the most bioactive BR[2], is widely considered the primary signaling ligand for BRI1, BRL1, and BRL3. Consequently, prior structural and biochemical investigations have focused on brassinolide[24,26-29,31,36,46]. Given the chemical diversity of BRs from the C_{27}, C_{28}, and C_{29} biosynthetic branches that accumulate in plants, we set out to systematically characterize their

binding kinetics to the ectodomains of the *Arabidopsis thaliana* BR receptors BRI1, BRL1, BRL2,

and BRL3. By integrating these measurements with structural and mutational analyses of the BRI1

binding pocket - both in vitro and in planta - as well as targeted BR bioassays, we define the ligand-

binding specificities of each receptor and uncover the chemical features in BRs required for

receptor binding and co-receptor recruitment to form an active signaling complex.

Results

<u>BRI1, BRL1 and BRL3 bind chemically diverse BRs with high affinity</u>

To investigate the ligand-binding specificity of Arabidopsis BR receptors, we obtained the

LRR ectodomains of BRI1, BRL1, and BRL3 by secreted expression in baculovirus-infected insect

cells, as described[26,28,46]. Quantitative GCI kinetic binding assays confirmed[28,29] that BRI1, BRL1

and BRL3 bind brassinolide with nanomolar affinity (Fig. 2a, Extended Data Figs. 1,2, Extended

Data Table 1). We then determined the crystal structure of the BRL3 ectodomain in complex with

brassinolide at 2.45 Å resolution (Fig. 2b and Supplementary Fig. 1 and Supplementary Table 1).

BRI1[26,27], BRL1[31] and BRL3 share a highly conserved ligand binding pocket (root mean square

deviation, r.m.s.d. is ~0.7 Å comparing 180 corresponding C_α atoms, Supplementary Fig. 1a),

formed by conserved hydrophobic residues along the inner LRR surface and island domain (Fig.

2b). The brassinolide C_{23} hydroxyl forms a hydrogen bond with a main chain amino group in the

island domain of all three receptors (Fig. 2b and Supplementary Fig. 1a). Few receptor-specific

polar contacts with the ligand are observed. In BRI1, Tyr597 forms hydrogen bonds with the

brassinolide C_{22} and C_{23} hydroxyls via a water molecule (Fig. 2b), whereas in BRL1 and BRL3,

Tyr597 is replaced by phenylalanine. In the BRL3 complex, Arg588 (corresponding to Lys601 in

BRI1) and Tyr627 form hydrogen bonds with the C_6 hydroxyl group of brassinolide (Fig. 2b).

BRL3 Gln666 (Phe681 in BRI1) coordinates the C_2 hydroxyl via a water molecule (Fig. 2b). Thus,

BRI1, BRL1, and BRL3 share a structurally conserved BR-binding pocket with limited receptor-

specific features, and all bind brassinolide with high affinity.

Next, we quantified the interaction of BRI1, BRL1, and BRL3 with 12 natural BRs from the

three biosynthetic branches, along with synthetic compounds 3-deoxycastasterone, 22-

deoxycastasterone, and 23-deoxycastasterone (Fig. 2c,d). BRI1 displayed binding constants ranging

from low micromolar (red in k_a/k_d plots, Fig. 2c) to low nanomolar (blue in Fig. 2c, Extended Data

Figs. 1,2, Extended Data Table 1). Low-affinity ligands include cathasterone (an early C_{28}

precursor), 24-epibrassinolide and 24-epicastasterone (late C_{28} branch), and the C_{29} products 28-

homocastasterone and 28-homobrassinolide (Fig. 1b, Fig. 2c,d). 28-norbrassinolide (C_{27} branch), as

well as brassinolide and castasterone (late C_{28} branch) represent high affinity ligands (Fig. 1b and Fig. 2c,d). Additional high-affinity binders include typhasterol, secasterone, and 2-deoxybrassinolide (late C_{22} oxidation), as well as 6-deoxocastasterone (early C_{22} oxidation), all within the C_{28} branch (Fig. 1b, Fig. 2c,d). BRL1 and BRL3 share highly similar ligand binding specificity profiles with BRI1 (Fig. 2c, Extended Data Figs. 1,2, Extended Data Table 1), consistent with their structurally conserved BR-binding pockets (Fig. 2b, Supplementary Fig. 1) and their ability to rescue *bri1-301* mutant phenotypes[29].

We next sought to define the chemical features of BRs that contribute to high-affinity receptor binding. Side chain modifications - such as the C_{24} ethyl substitution in 28-homocastasterone or 28-homobrassinolide (Fig. 1a,b) - were associated with ~10- to 30-fold and ~10- to 35-fold reductions in binding affinity, respectively, compared to castasterone and brassinolide (Fig. 2c,d). Similarly, the brassinolide / castasterone enantiomers 24-epibrassinolide and 24-epicastasterone (Fig. 1a,b) showed ~10- to 20-fold and ~10- to 40-fold reduced affinities (Fig. 2c,d). Absence of a hydroxyl group at C_{23} - as in cathasterone - correlated with weak receptor binding (Fig. 1b, Fig. 2c,d).

To further assess the role of side chain hydroxylation, we synthesized 23-deoxycastasterone (Supplementary Note 1). 23-deoxycastasterone bound BRI1 with a dissociation constant of ~3 μM, representing a 100-fold reduction compared to castasterone (Fig. 2c,d). In contrast, 22-deoxycastasterone - lacking the C_{22} hydroxyl group present in castasterone and brassinolide (Fig. 1b) - still bound BRI1 with medium to high affinity (K_d ~200 nM, Fig. 2c,d). These results are consistent with our structural data showing that the C_{23}, but not the C_{22} hydroxyl group forms conserved hydrogen bonds with the island domains of BRI1, BRL1, and BRL3 (Fig. 2b, Supplementary Fig. 1a).

Ring modifications were generally well tolerated by the BRI1, BRL1, and BRL3 ectodomains (Fig. 2c,d). Lactonization of the B-ring had minimal impact on binding, as brassinolide and castasterone showed nearly identical dissociation constants (Fig. 2c,d). Although C_6 oxidation is a key biosynthetic step towards brassinolide[13,51], 6-deoxocastasterone bound all three receptors with high affinity (Fig. 2c,d). Supporting this, the complex structures of BRI1, BRL1, and BRL3 show no conserved interactions with the C_6 ketone[26,27,31] (Fig. 2b, Supplementary Fig. 1a). The 2α and 3α hydroxyl groups in the A-ring also appear dispensable for receptor binding, as ligands lacking either - such as 2-deoxybrassinolide, 3-deoxycastasterone, typhasterol, and secasterone - bound BRI1, BRL1, and BRL3 with nanomolar affinity (Fig. 2c,d).

164   Overall, our quantitative binding data demonstrate that BR receptors recognize a broad
165   range of structurally diverse BRs with high affinity. High-affinity receptor binding is driven by the
166   configuration and size of the aliphatic side chain - most critically, the presence of a C_{23} hydroxyl
167   group and the correct C_{24} stereochemistry.
168
High structural plasticity of the hormone binding pocket enables BRI1/BRLs to sense different BRs
To clarify the structural basis of ligand-binding specificity in BR receptors, we co-
crystallized the BRI1 ectodomain with various BRs. Crystals grown with cathasterone, 24-
epicastasterone, or 28-homocastasterone showed no interpretable difference electron density in the
BR-binding pocket, while 6-deoxocastasterone yielded poorly diffracting crystals. A 2.1 Å structure
of the BRI1 - castasterone complex revealed a ligand conformation nearly identical to that of
brassinolide (Fig. 3a,b, Supplementary Table 1), consistent with its high binding affinity (Fig. 2c,d),
and confirming that B-ring lactonization is not essential for receptor recognition. Similarly, a 2.7 Å
structure of the BRI1 - typhasterol complex showed that the absence of a C_2 hydroxyl group does
not affect ligand coordination by the receptor (Fig. 3a,b). In contrast, the 2.4 Å BRI1 - 24-
epibrassinolide structure exhibited weak electron density for the BR side chain, consistent with its
reduced binding affinity (Fig. 2c,d, Fig. 3d). Finally, the 2.4 Å BRI1 - 28-homobrassinolide
complex revealed that the receptor can in principle accommodate a C_{24} ethyl substitution in its
binding pocket (Fig. 3a,b).
We next resolved the crystal structures of the BRL3 ectodomain bound to castasterone and
typhasterol at 2.6 Å and 2.3 Å, respectively. In both structures, the ligands adopt conformations
closely resembling that of brassinolide (Fig. 3a,b). A 3.2 Å structure of the BRL3 - 6-
deoxocastasterone complex shows that, despite lacking B-ring modifications, the ligand is
accommodated in the binding pocket, albeit in a shifted conformation (Fig. 3a,b). Notably, in all BR
receptor - ligand complexes analyzed, the BR C_{23} hydroxyl group consistently forms a conserved
hydrogen bond with the main chain amino group of Ser647 in BRI1 or Met632 in BRL1/BRL3
(Fig. 3a).
Several catabolic enzymes contribute to BR homeostasis, including BAS1, which
hydroxylates side chain C_{26}[17], UGT73C5, which glucosylates the hydroxyl group at C_{23}[18], and
ST1/ST4a, which sulfonylates the C_{22} hydroxyl[19]. Structural analysis of our receptor–ligand
complexes indicates that such tail modifications are not compatible with the BR-binding pocket
(Supplementary Fig. 2). Together, the BRI1 and BRL3 ligand-bound structures rationalize the BR

binding kinetics data, emphasizing the importance of the BR side chain for high-affinity receptor

binding, while showing that modifications to the steroid ring core are generally tolerated.

To evaluate the physiological relevance of our biochemical and structural experiments, we

selected ligands representing high, medium, and low-affinity binders for root growth inhibition

assays, as previously described[43]. Brassinolide (K_d ~20 nM) and 28-norbrassinolide (K_d ~80 nM)

were tested as high-affinity binders, 28-homobrassinolide (K_d ~380 nM) as medium-affinity, and

23-deoxycastasterone (K_d ~2800 nM) as a low-affinity binder (Fig. 2d). Root growth inhibition

assays were performed using each compound at a final concentration of 100 nM. A strong trend for

increasing root length with increasing dissociation constant (K_d) was observed in treated wild-type

seedlings, and a weaker trend in the *bsk3-1* control (Fig. 3c, Extended Data Table 2), suggesting that

in vivo BR bioactivities correlate well with their receptor binding affinity in vitro. Notably, even the

low-affinity binder 23-deoxycastasterone significantly inhibited root growth of wild-type seedlings

at a concentration of 100 nM (Fig. 3c).

We next correlated in vitro BR receptor binding affinities with BR levels in planta. We

quantified BR levels in 10-d-old Arabidopsis plants using UHPLC-MS/MS (see Methods). In whole

wild-type seedlings, brassinolide was detected at low levels (~0.6 pmol/g fresh weight), while

castasterone levels were ~20-fold higher (Fig. 3d). 28-norbrassinolide was also present at low

concentrations (~0.2 pmol/g FW), whereas 28-homocastasterone was abundant (~140 pmol/g FW)

(Fig. 3d). Correlating these endogenous BR levels with their respective in vitro binding affinities

(Fig. 2c,d), suggests that, in addition to brassinolide, other BRs - such as castasterone, 28-

norbrassinolide, and 28-homocastasterone - may act as physiologically relevant ligands for BRI1,

BRL1, and BRL3.

Constitutive activation or inactivation of BRI1 modulates BR homeostasis

We next compared BR levels in wild-type and *bri1* null mutant seedlings[47]. In the *bri1*

mutant, brassinolide and castasterone levels were elevated by ~7-fold and ~3-fold, respectively,

compared to wild type (Fig. 3d), as previously described[45]. This aligns with earlier findings that BR

metabolic and catabolic gene expression is altered in *bri1* loss-of-function mutants[52], partly

mediated by the BR-regulated transcription factor BZR1[40,53]. To further investigate, we performed

RNA-seq on untreated 10-d-old Col-0 seedlings, a *bri1* null mutant[47], and in *bri1* complementation

lines expressing either a wild-type BRI1 receptor[47], the BRI1[sud1] gain-of-function allele[47,36], or a

constitutively active, ligand-independent oBIR3-iBRI1 receptor chimera[54]. We found many BR

biosynthesis genes to be mildly induced in *bri1* mutant seedlings and to be repressed in the

constitutively active oBIR3-iBRI1[54] line (Fig. 3e and Extended Data Table 3). BR biosynthesis

genes were up-regulated in *bri1* and down-regulated in oBIR3-iBRI1 seedlings (Fig. 3e, Extended

Data Table 3). Conversely, several BR catabolic genes were repressed in *bri1* and induced in

oBIR3-iBRI1 plants (Fig. 3e). ABCB1 and ABCB19, which function as BR transporters required

for signaling[23], also showed differential expression in *bri1* and oBIR3-iBRI1 seedlings (Fig. 3e).

Notably, *BRL1*, *BRL2*, and *BRL3* transcript levels were reduced in oBIR3-iBRI1 and partially

induced in *bri1* (Fig. 3e). Expression of a wild-type BRI1 receptor restored gene expression in *bri1*

mutants to wild-type levels (Fig. 3e). The gain-of-function allele BRI1^{sud1} mimicked the effects of

the constitutively active oBIR3-iBRI1 receptor chimera, albeit with less pronounced effects (Fig.

3e). Together, these experiments indicate that BRI1 signaling activity shapes BR homeostasis by

modulating the expression of BR metabolic, catabolic, and transport genes.

<u>BRL2 is a medium-affinity BR receptor</u>

The structural flexibility of the BR-binding pockets in BRI1, BRL1, and BRL3, along with

altered BRL2 expression in BRI1 signaling mutants (Fig. 3e), prompted us to re-examine BRL2's

ligand-binding properties. BRI1 and BRL2 share ~40% sequence identity in their extracellular

domains. We expressed and purified the BRL2 ectodomain from insect cells and solved its apo

crystal structure to 2.6 Å resolution (see Methods). The N-glycosylated BRL2 adopts a superhelical

architecture with 21 LRRs (compared to 25 in BRI1) with a more open conformation

(Supplementary Fig. 3a). BRL2's island domain is partially disordered and participates in a domain-

swapped crystallographic dimer (Supplementary Fig. 3a).

The ligand-binding pocket of BRL2 exhibits a high degree of structural similarity with the

BR-binding sites of BRI1, BRL1, and BRL3 (Fig. 4a, Supplementary Fig. 3b). Using GCI assays,

we found that BRL2 can bind BRs, but it has a distinct ligand binding profile compared to BRI1,

BRL1 and BRL3 (Fig 4b,c, Fig. 2c,d). BRL2 bound brassinolide and castasterone with K_d values of

~200 nM and ~400 nM, respectively (Fig. 4b,c; Extended Data Fig. 3, Extended Data Table 4) - 10-

30-fold weaker than BRI1 (Fig. 2c,d). Unlike BRI1, which binds typhasterol and 28-norbrassinolide

with nanomolar affinity, BRL2 bound these ligands only weakly (Fig. 4b,c). BRL2 showed no

detectable binding to 28-homocastasterone, 24-epicastasterone, 24-epibrassinolide, or to

dehydroepiandrosterone (DHEA), triolone, lupeol, or betulin (Fig. 4b,c). Our experiments suggest

that BRL2 functions as a medium-affinity receptor for C_{28} BRs (Fig. 1b). In line with this, the BR-

binding pocket of BRI1, BRL1/3 and BRL2 share a high degree of structural and sequence

conservation. The only notable exception is Glu614 in BRL2, which corresponds to Met657 and Ile642 in BRI1 and BRL3, respectively (Fig. 4a, Supplementary Fig. 3b).

BRL2/VH1 has been previously linked to leaf vein patterning[34]. To assess its expression, we generated a *BRL2prom*::NLS-3xmVenus reporter line in Col-0 background and found *BRL2* to be expressed in vascular tissues across different developmental stages (Fig. 4d). To investigate its function, we analyzed the *brl2-1* loss-of-function mutant alongside a *bri1* null mutant[47]. Unlike *bri1*, *brl2-1* showed no dwarf phenotype when grown in soil (Fig. 4e). A second loss-of-function allele, *brl2-2*, created by CRISPR/Cas9 gene editing (Supplementary Fig. 3c), also showed no visible growth defects (Fig. 4e). Although prior work reported vein patterning defects in *brl2-1*[34], a quantitative analysis of leaf vein patterns did not reveal statistically significant differences compared to the Col-0 wild-type control (Supplementary Fig. 3d). *brl2-1* mutants also showed wild-type responses in hypocotyl growth assays (Supplementary Fig. 3e) and normal BES1 dephosphorylation in response to brassinolide (Supplementary Fig. 3f).

We have previously reported that receptor chimeras, in which the extracellular and transmembrane domains of BIR3 (oBIR3) are fused to the cytoplasmic kinase domain of BRI1 (iBRI1), trigger constitutive, ligand-independent BR responses[38,54]. To assess the signaling capacity of BRL2, we expressed a *BRL2prom*::oBIR3-iBRL2 chimera in Col-0 wild-type background. These transgenics displayed elongated, slender petioles and curly leaves, similar to BRI1[sud1] gain-of-function mutants[24,47] (Fig. 4e). However, their rosette phenotypes were milder than those of *BRI1prom*::oBIR3-iBRI1 chimeras[54] (Fig. 4e). Overall, BRL2 is a vascular-localized, medium-affinity BR receptor that, like BRL1 and BRL3[29,30], does not exhibit strong loss-of-function phenotypes.

A mutational map of the BR-binding pocket rationalizes the specificity and plasticity of BR receptors

Given the high structural conservation of the BR-binding site in BRI1, BRL2, BRL1 and BRL3 (Supplementary Fig. 1b, Supplementary Fig. 4a), we examined the role of individual pocket-lining residues in BR perception. Based on our BRI1 - BR complex structures (Fig. 3a), we mutated eleven residues in the LRR core and island domain (Fig. 5a) and measured brassinolide binding kinetics for each mutant ectodomain (Fig. 5b,c, Extended Data Fig. 4, and Extended Table 5). All mutants behaved as well-folded monomers in solution (Supplementary Fig. 4b).

Consistent with the BR-binding site's structural plasticity, mutation of the conserved Thr646, Tyr597, or Lys601 from the island domain had no detectable effect on brassinolide binding (Fig. 5a-

c). Substitution of Tyr599 or Tyr642 with alanine moderately reduced binding affinity (K_d ~100 nM

and ~200 nM, respectively), and replacing the invariant Leu615 with phenylalanine or tyrosine

likewise had little effect on BR binding (Fig. 5a–c). In contrast, mutation of Trp564 → His caused a

~100-fold drop in BR binding (K_d ~2 μM) (Fig. 5a–c). Notably, mutating Met657 (Glu614 in BRL2,

see Supplementary Fig. 3b) to either alanine or glutamate did not affect brassinolide binding (Fig.

5a-c). Consistently, a mild growth phenotype has been previously reported for the BRI1[M657E]

mutant[46]. These findings underscoring the presence of a conserved BR-binding site in

BRI1/BRL1/BRL3 and in BRL2 (Fig. 4a). As previously shown[46], the *bri1-6*[45] mutation

(Gly644→Asp) strongly reduces brassinolide binding (Fig. 5a-c). In contrast, the neighboring

Gly643→Glu mutation in the BRI1*sud1* gain-of-function allele[47] stabilizes the island domain[36] and

enhances brassinolide binding by ~13-fold over wild type (Fig. 5a–c). Similarly, mutating LRR core

Phe681 to alanine increases binding by ~7-fold. These results further highlight the BR-binding site's

structural plasticity and identify Trp564 from the LRR core as essential for ligand recognition.

To assess the functional impact of these mutations, we expressed mutant BRI1 receptors

under the native BRI1 promoter in a *bri1* null background (Fig. 5d). T3 lines with wild-type-like

BRI1 expression (Fig. 5e and Supplementary Fig. 4c) and proper plasma membrane localization

(Supplementary Fig. 4d) were selected. High-affinity mutants (Tyr597→Phe, Leu615→Phe,

Thr646→Ile) restored wild-type rosette and root growth phenotypes (Fig. 5d-f), and BR responses

(Supplementary Fig. 4e,f and Extended Data Table 6). Intermediate phenotypes were observed in

lines with Tyr597→Met, Leu615→Tyr, or Met657→Ala, consistent with their moderate in vitro

binding affinities (Fig. 5a–c). The Gly644→Asp mutant displayed medium-to-strong defects, while

Trp564→His, which showed ~100-fold reduced binding, phenocopied the *bri1* null at both seedling

and adult stages (Fig. 5d–f).

Conversely, Gly643→Glu and Phe681→Ala conferred gain-of-function rosette phenotypes,

matching their enhanced binding affinities compared to wild type (Fig. 5a-d). Three mutants

(Tyr599→Ala, Lys601→Leu, Tyr642→Ala) displayed severe growth phenotypes despite retaining

strong brassinolide binding in vitro (Fig. 5a-f, see below). Taken together, we observe a strong

correlation between the in vitro ligand binding affinities and in planta growth phenotypes of BRI1

mutant receptors, yielding novel loss-of-function (Trp564-His) and gain-of-function (Phe681-Ala)

alleles.

 BR antagonists target the interaction of BR receptors with SERK co-receptors

We previously showed that the BR-binding site comprises the LRR core and island domain

of BRI1, and the LRR ectodomain of a SERK co-receptor[26,36,28] (Fig. 6a). SERK1 Phe61 stacks with

the A/B rings of brassinolide, while His62 forms a conserved hydrogen bond with the C_2 hydroxyl

group of the ligand[36] (Fig. 6a). Mutating the corresponding Phe60 and His61 in SERK3/BAK1 to

alanine abolishes both co-receptor interaction in vitro and BR signaling in planta[28].

To test whether this activation mechanism is conserved across BR receptors, we

immobilized BRI1, BRL1, BRL2, and BRL3 ectodomains on a GCI chip and examined SERK1

binding in the presence of 100 nM brassinolide. BRL1 and BRL3, like BRI1, bound SERK1 with

high affinity, while BRL2 displayed much weaker SERK1 binding (K_d ~35 μM), even in the

presence of 2 μM brassinolide (Fig. 6b, Extended Data Fig. 5, Extended Data Table 7).

We next quantified the interaction of SERK1 with BRI1 in the presence of different BRs. We

used an avi-tagged BRI1 ectodomain to minimize steric hindrance between the GCI chip and the

SERK1 co-receptor (see Methods), and measured dissociation constants of ~100 nM for SERK1

association to either brassinolide- or castasterone-bound BRI1 (Fig. 6c,d, Extended Data Fig. 5,

Extended Data Table 7). However, SERK1 binding was ~15-fold weaker with typhasterol- vs.

castasterone-bound BRI1 (Fig. 6c,d), despite these ligands having similar receptor binding affinities

(Fig. 2c,d). This reveals that the C_2 hydroxyl coordinated by SERK1 His62 is key for receptor – co-

receptor complex formation (Fig. 6a,e). Similarly, 2-deoxybrassinolide reduced SERK1 binding

~10-fold, while 3-deoxycastasterone, lacking the C_3 hydroxyl, still supported high-affinity SERK1

binding (Fig. 6c–e). Secasterone - with a bulky 2,3-α-epoxy group - impaired complex formation, as

did 6-deoxocastasterone, despite possessing both C_2 and C_3 hydroxyls (Fig. 6c–e), likely due to its

distorted binding mode in the BR pocket (Fig. 3a,b).

Our data identify the BR C_2 hydroxyl as essential for SERK recruitment. While C_6 oxidation

on the B-ring is not required for receptor binding, its coordination by BRI1 island domain residues

Tyr599, Tyr642, and Lys601 may help orient the ligand for optimal co-receptor binding (Fig. 6a,c–

e). Indeed, SERK1 binding is partially or severely compromised in the BRI1[Y599A], BRI1[K601L] and

BRI1[Y642A] mutants in the presence of high, saturating concentrations of brassinolide (Fig. 6f). This

results rationalize the dwarf phenotypes of the BRI1 Tyr599→Ala, Lys601→Leu and Tyr642→Ala

mutants (Figs. 5d–f, 6a), despite their ability to bind brassinolide with high affinity (Fig. 5c).

As BRs that bind BRI1 but block SERK association can act as receptor antagonists, we

performed seedling root growth inhibition assays in the presence of 100 nM brassinolide and with

different antagonist candidates from our biochemical screen (Fig. 6d). We found that secasterone

applied on its own did not trigger root growth inhibition in a concentration-dependent manner (Fig. 6g, Extended Data Table 8). However, applying increasing concentrations of secasterone significantly reversed brassinolide-induced root growth inhibition (Fig. 6g). 2-deoxybrassinolide acted as a weak BR agonist when applied on its own, but had a significant, dose-dependent antagonistic effect on brassinolide-induced root growth inhibition (Fig. 6g). In contrast, 3-deoxycastasterone acted as a weak BR agonist both in the absence and presence of brassinolide in the assay (Fig. 6g). We conclude from these experiments that the BRs lacking a C_2 but not C_3 hydroxyl group behave as receptor antagonist (Fig. 6a,c-e,g). A bulky head group, such as in secasterone is also effective in blocking receptor activation in vitro and in planta (Fig. 6a,c-e,g). We speculate that the agonistic effect of 2-deoxybrassinolide may be caused by this metabolite entering the biosynthetic pathway, generating other, bioactive BRs (Fig. 6g and Fig. 1b). Taken together, BRs that block BRI1 – SERK1 complex formation (Fig. 6c-e) in vitro without compromising isolated receptor binding (Fig. 2c,d), can partially reverse brassinolide-induced root growth inhibition in a dose-dependent manner (Fig. 6g), underscoring the importance of A-ring features in BR receptor activation[36,28].

Discussion

Genetic and metabolite studies have revealed a complex BR biosynthetic network in plants[2,14]. In this study, we provide quantitative binding kinetics for 15 distinct BRs from this network, evaluated against the four BR receptors in Arabidopsis. Our results highlight that hydroxylation at C_{23} and C_{24} stereochemistry in the BR aliphatic tail are critical for receptor binding (Fig. 2b–d) and BR bioactivity (Fig. 3d and Fig. 7). Consistent with this, C_{23} hydroxylase mutants display severe dwarfism[9,51]. Lack of a C_6 ketone is tolerated by the isolated receptors (Fig. 2b-d) but impairs proper ligand positioning and SERK co-receptor recruitment (Fig. 6c-f and Fig. 7). The presence of a B-ring lactone is not required for high affinity receptor binding (Fig. 2b-d and Fig. 3a,b), in agreement with the moderate growth phenotype of the *br6ox2* mutant[13,51]. Notably, C_2 hydroxylation is essential for SERK recognition[28,36] (Fig. 6c-e) but not for receptor binding (Fig. 2b-d and Fig. 3a,b), providing a mechanistic basis for known BR antagonists[55,56] (Fig. 7). 2-deoxybrassinolide and secasterone act as receptor antagonists (Fig. 6e,g), supporting the idea that these abundant[48] biosynthetic intermediates may modulate BR signaling by competing with agonists for receptor binding (Fig. 1b, Fig. 6g). This could prevent formation of active BR signaling complexes, in a similar way to the known BIR receptor pseudo-kinases[37,38].

Importantly, our quantitative BR binding assay identifies BRL2 as a functional BR receptor
with a distinct ligand binding spectrum (Fig. 2c,d, Fig. 4b,c). BRL2 shares a conserved BR binding
mechanism (Supplementary Fig. 3b) and weakly interacts with SERK co-receptors (Fig. 6b). The
gain-of-function phenotype of oBIR3-iBRL2 receptor chimera (Fig. 4e) further supports the notion
that BRL2 shares the SERK-based receptor activation mechanism[54] with BRI1[28,35,36], BRL1 and
BRL3 (Fig. 6b). We speculate that the inability of BRL2 to replace BRI1 in *bri1* mutants[29,33] is
caused by its distinct BR ligand binding spectrum (Fig. 4b,c), while its receptor activation
mechanism and cytoplasmic signaling pathway appear to be conserved[57,58]. Taken together, BRL2,
which emerged early on in land plant evolution, may represents an ancestral BR receptor[59].

Mutational analysis of the BR-binding pocket in BRI1 reveals that the receptor can tolerate
various missense mutations (Fig. 5 and Fig. 7), in agreement with previous mutational studies[46,60].
We identified single point mutation causing a strong BR loss-of-function phenotype (Trp564His),
two residues (Tyr599 and Tyr642) critically involved in the proper positioning of the ligand in the
BR binding pocket, and a novel gain-of-function allele (Phe681Ala) (Fig. 5d-f and Fig. 7). The
structural plasticity of the hormone binding pocket supports the receptor's capacity to recognize
chemically diverse BRs, while providing sufficient selectivity to discriminate between BRs, other
phytosterols and triterpenoids. Notably, the BR-binding site is conserved among the four
Arabidopsis receptors and their orthologs in crop species, implying that BR receptors in crops likely
share a similar BR ligand binding spectrum (Supplementary Fig. 4a).

Based on binding profiles (Fig. 2c,d, Fig. 6c,d), bioactivity data, and in vivo BR levels[11,48,50]
(Fig. 3d), we propose that brassinolide and castasterone (C_{28}), 28-norbrassinolide (C_{27}), and possibly
28-homocastasterone (C_{29}) act as physiological BR ligands in Arabidopsis. Such a broad ligand
spectrum has been previously reported for other small molecule[61] and peptide[62] hormone receptors
in plants. Feedback regulation of BR biosynthesis likely fine-tunes BR levels (Fig. 3e) in different
tissues[51]. We would like to note that our in vitro binding assays with the isolated LRR domains
probably underestimate the actual BR binding affinities of intact BRI1 - SERK receptor complexes
embedded in the plasma membrane, as plants can respond to BRs applied externally at picomolar to
low nanomolar concentrations[44,51].

Taken together, distinct chemical features in BRs required for (1) high affinity receptor
binding, (2) proper ligand orientation in the receptor hormone binding pocket, and (3) high affinity
co-receptor binding (Fig. 7) have now been defined and can be exploited for rational BR agonist
and antagonist design.
(main text: 4,866 words, 61 citations)

Online Material and methods

<u>Protein expression and purification</u>

The ectodomains from AtBRI1 (https://www.uniprot.org UniProt-ID O22476, residues 1-788), AtBRL1 (Q9ZWC8, 1-770), AtBRL2 (Q9ZPS9, 1-752), AtBRL3 (Q9LJF3, 1-770) and AtSERK1 (Q94AG2 , 23-270) ectodomains were amplified from *Arabidopsis thaliana* cDNA (ecotype Col-0). AtBRI1, AtBRL1, AtBRL2 and AtBRL3 were cloned in vector pBB2 providing tobacco etch virus (TEV) protease cleavable C-terminal Twin-StrepII and 9xHis affinity tags. AtSERK1 was cloned in the same pBB2 vector containing an N-terminal azurocidin signal peptide[63,28]. For grating-coupled interferometry assays, AtBRI1 was also cloned in vector pBB2 containing a C-terminal Avitag followed by TEV cleavable C-terminal Twin-StrepII and 10xHis affinity tags.

All proteins were expressed in *Trichoplusia ni* (Tnao cells, Invitrogen) using the baculovirus system[64]. Tnao cells in SF900 medium (Gibco) were grown to a density of 2.0 x 10^6 cells/mL and subsequently infected with 10 mL of P2 virus. Cells were then incubated for 24 h at 28 ºC and for an additional 48 h at 21 ºC while shaking at 110 rpm. Cells were then pelleted at 4'000 x g for 20 min and the supernatant containing the secreted protein was filtrated using 0.45 µm filters (Durapore Merck). The cleared medium was then loaded onto an Ni^{2+} metal affinity column (HisTrap Excel 5 mL, Cytiva) equilibrated in buffer A (50 mM KPi pH 7.8, 250mM NaCl, 1 mM β-mercaptoethanol). The column was washed with 25 mL of buffer A and subsequently eluted onto a Strep-Tactin XT 4flow 5mL column (IBA) in buffer A supplemented with 250 mM imidazole (pH 8.0). The Strep-Tactin affinity column was further washed with 25 mL buffer (20 mM Tris pH 8.0, 250 mM NaCl and 1 mM EDTA), and the protein was eluted from the Strep-Tactin column with buffer C (10 mM Tris pH 8.0, 150 mM NaCl, 1 mM EDTA, 50 mM biotin, 1xBTX, IBA). For protein crystallization, the affinity purified proteins were incubated overnight at 4 ºC with TEV protease using a molar receptor:protease ration of ~10:1. The tag free protein was separated from the tag and from TEV protease by a second Ni^{2+} affinity chromatography step. Finally, the untagged protein was applied to a HiLoad Superdex 200 16/200 pg column (Cytiva) equilibrated in 20 mM sodium citrate pH 5.0, 150 mM NaCl. Fractions containing the purified protein were concentrated and stored at -80 ºC.

Receptor biotinylation for grating-coupled interferometry was achieved by incubating Avi-tagged BRI1 with His-tagged BirA[65]. Briefly, BRI1 at a final concentration of 20 µM was incubated for 1 h at 30 °C with BirA, biotin, ATP and $MgCl_2$ at final concentrations of 85 µM, 150 µM, 2 mM and 5 mM respectively. The BirA enzyme was subsequently removed using a Ni^{2+} affinity

chromatography step. Biotinylated BRI1 was concentrated and loaded onto a HiLoad Superdex 200

16/200 pg column (Cytiva) equilibrated in 20 mM sodium citrate pH 5.0, 150 mM NaCl.

Chemical synthesis and analysis of BRs

The chemical synthesis of typhasterol, 3-deoxycastasterone and 2-deoxybrassinolide are

described in Supplementary Note 1.

Grating Coupled Interferometry (GCI)

GCI experiments were performed using the Creoptix WAVE system (Malvern Panalytical).

Binding of BRs to the isolated BRI1, BRL1, BRL2 and AtBRL3 ectodomains was measured using

4PCH WAVEchips. Borate buffer (100 mM sodium borate, pH 9.0, 1 M NaCl, Xantec) injections

were used to condition the chips. For the interaction screening between BRI1, BRL1, BRL2, BRL3

or BRI1^{mut} with BRs, the purified ectodomains were immobilized using standard amine-coupling.

Chips were activated using a 1:1 mix of 400 mM N-(3-dimethylaminopropyl)-N'-ethylcarbodiimide

hydrochloride and 100 mM N-hydroxysuccinimide (Xantec). Protein Ligands (in 10 mM sodium

acetate, pH 5.0) were injected to the desired chip surface at the concentration indicated in Extended

Data Tables 1, 4 and 5. In order to minimize non-specific binding of analytes, a second injection of

BSA (0.5% [w/v] in 10 mM sodium acetate, pH 5.0) was used to passivate the surface. Finally, the

chip was then quenched by injecting 1 M ethanolamine. All kinetic analyses were performed at

room temperature in citrate buffer (20 mM sodium citrate pH 5.0, 150 mM NaCl) supplemented

with 0.05 % (v/v) Tween and 0.025 % (v/v) dimethyl sulfoxide to reduce unspecific binding of the

respective BR analyte. Analyte concentrations were varied in 1:3 dilution series (compare Extended

Data Tables 1, 4, 5). All experiments were done in duplicates, blank injections were used for double

referencing and DMSO calibration was performed for bulk correction.

Binding of the co-receptor SERK1 to brassinolide-bound wild-type and mutant receptors

was done by amine-coupling the BRI1, BRL1, BRL2 and BRL3 ectodomains onto 4PCP

WAVEchips and using the purified SERK1 ectodomain as analyte (Extended Data Table 7) in 20

mM sodium citrate pH 5.0, 150 mM NaCl) supplemented with 0.05 % (v/v) Tween and 0.025 %

(v/v) dimethyl sulfoxide and brassinolide to a final concentration of 1 μM. Binding of SERK1 to

BRI1 in the presence of different BRs was performed by immobilizing Avi-tagged and biotinylated

BRI1 by neutravidin capturing. After chip activation, neutravidin (100 μg/mL in 10 mM sodium

acetate, pH 5.0) was injected. To passivate and quench the chip surface, a further injection of 0.5%

(w/v) BSA followed by an injection of 1 M ethanolamine were performed, respectively.

Biotinylated BRI1 (100 µg/mL in citrate buffer) was then injected to the desired surface

concentration (Extended Data Table 7), followed by an injection of biotinylated 0.5% (w/v) BSA in

order to passivate the surface. All the binding assays between BRI1 and SERK1 were done at room

temperature in citrate buffer supplemented with 0.05% Tween and the respective BR at a final

concentration of of 1 µM. SERK1 injections were done in 1:3 dilution series (Extended Data Table

7).

Data analysis was performed with the Creoptix WAVEcontrol software (version: 4.7.2) and

data were fitted using one to one binding models or mass transport model, using bulk correction.

Plotting and data representation were performed using R (https://www.r-project.org, version: 4.1.2).

Crystallization and data collection

BRL3 (in 20 mM citric acid pH 5.0, 250 mM NaCl) at a concentration of ~1 mg/mL was

incubated with the respective BR (10 mM stock solution in 100 % [v/v] dimethyl sulfoxide

[DMSO]) to a final concentration of 50 µM for 4 h at room temperature and then concentrated to

~6-7 mg/mL using a Amicon Ultra concentrator (Merck Millipore) with a molecular weight cut-off

of 30 kDa. Triclinic crystals developed in hanging drops composed of 1.0 µL of protein solution and

1.0 µL of crystallization buffer (23-25 % [w/v] PEG 3,350, 0.2 M Li_2SO_4, 0.1 M citric acid pH 4.0)

suspended over 1.0 mL of the latter as reservoir solution. Crystals were cryoprotected by serial

transfer into crystallization buffer containing ethylene glycol to a final concentration of 25 % (v/v),

and snap frozen in liquid N_2. BRI1 (in 20 mM citric acid pH 5.0, 250 mM NaCl) - BR (at a final

concentration of 50 µM) complex crystals developed in the same crystallization condition. Crystals

of apo BRL2 were obtained using micro-seeding protocols in hanging drops containing 1.0 µL of

protein solution (BRL2 at 6 mg/mL in 20 mM citric acid pH 5.0, 250 mM NaCl) and 1.0 µL of

crystallization buffer (23 % [w/v] PEG 3,350, 0.2 M MgCl2 · 6H_2O, 0.1 M Tris pH 9.0) over 1 mL

of reservoir solution. Tetragonal crystals were cryoprotected by serial transfer in crystallization

buffer supplemented with 20% (v/v) glycerol. All datasets were collected at beam line X06DA of

the Swiss Light Source, Villigen, Switzerland (Supplementary Table 1). Data processing and scaling

was done with XDS[66] (version: June 2024).

Crystallographic structure solution and refinement

The structures of the BRL3 - brassinolide complex and of apo BRL2 were solved by

molecular replacement in PHASER[67], with BRI1 ectodomain structure[26] (PDB-ID pdb_00003rj0) as

search model. For BRL3, the solution revealed a dimer in the asymmetric unit, closely resembling

the previously reported BRL1 dimer[31]. The BRL2 structure was resolved at 2.6 Å resolution in
space group $P \; 4_1 \; 2_1 \; 2$, containing a single molecule per asymmetric unit (Supplementary Table 1).
Crystals of the BRI1–BR complex shared the same crystal form as the previously described apo
BRI1[26]. Restraints for the BR ligand were generated from SMILES strings using phenix.elbow[68]. All
structures were completed through iterative cycles of manual model building in COOT[69] and
restrained TLS refinement in phenix.refine[70]. Analysis of the refined structures with
phenix.molprobity[71] revealed good to excellent stereochemistry. Omit maps of the different BR
ligands were generated with phenix.polder[72]. Data collection and refinement statistics are provided
in Supplementary Table 1. Structural figures were prepared with ChimeraX[73], chemical structures
were drawn with PubChem sketcher[74].

<u>Plant material</u>
Arabidopsis thaliana ecotype Col-0, T-DNA insertion mutants *bsk3-1* (SALK_096500C)[75],
bri1 null (GABI_134E10)[47] and *brl2-1* (SALK_016024)[76] were obtained from the Nottingham
Arabidopsis Stock Center, alongside the previously reported *BRI1prom*::BRI1-mCitrine / *bri1* and
BRI1prom::BRI1*sud1*-mCitrine / *bri1*[36] and *BRI1prom*::oBIR3-iBRI1-mCitrine / *bri1*[54] transgenic
lines. Genotyping primers for the mutants are listed in Supplementary Table 2.

<u>Molecular cloning and generation of transgenic lines</u>
To generate BRL2 expression reporter lines, a 3.6 kb fragment upstream of the BRL2 ORF
was cloned into the entry vector pUC57- L4_EcoRV-XbaI-BamHI_R1 via Gibson assembly to
produce L4-*BRL2prom*-R1[77]. Multi-site Gateway cloning (Thermo Fisher Scientific) with Gateway
LR reaction (11791020; Thermo Fisher Scientific) was used to assemble the following fragments
L4-BRL2prom-R1, L1-NLS-3xmVenus-L2[78] and R2-tHSP18.2-L3[79] into the destination vector
pB7m34GW[80], to generate *BRL2prom*::NLS-3xmVenus. The BRL2 chimera (*BRL2prom*::oBIR3-
iBRL2-mCitrine) were generated using a construct in which the N-terminal part of AtBIR3
(residues 1-246, containing the native signal peptide, extracellular domain, and transmembrane
helix) and the C-terminal region of BRL2 (residues 779-1143, comprising the juxtamembrane and
kinase domains) were joined using two-fragment Gibson assembly, as previously described[54]. For
the BRI1 complementary assays, the BRI1 coding sequence lacking a stop codon was cloned into
entry vector pDONR221[36] using the Gateway BP reaction (11789020; Thermo Fisher Scientific).
The single point mutations were introduced by the site-directed mutagenesis by overlap extension[81]
on the entry vector. A 1690 bp upstream fragment of BRI1 (*BRI1prom*) and a mCitrine coding

sequence with the stop codon was cloned into pDONR P4-P1R and pDONR P2R-P3, respectively, as previously reported[54]. The final binary vectors *BRI1prom*::BRI1mu-mCitrine were generated by the multi-site Gateway technology (Thermo Fisher Scientific) with destination vector pB7m34GW[80]. Cloning primers are list in Supplementary Table 3.

The *brl2-2* allele was generated using a CRISPR/Cas9[82] system[83]. Two 20 bp sgRNA target sequences, designed using CRISPR-P 2.0[84], were cloned into shuttle vectors pEF004/005 via mutagenesis PCR[85]. These cassettes were subsequently assembled into the pBay-Bar-CRmCherry vector using Gibson assembly and the final construct was transformed into Col-0 wild-type plants using the floral dip method[86]. Primary transformants (T1) were selected on $^{1/2}$MS medium containing 15 mg/L phosphinothricin; Cas9-free mutant lines (T2/T3) were identified by screening for loss of the mCherry marker (Leica Stereo Microscope MSV269, Wetzlar, Germany).

All constructs were transferred into *Agrobacterium tumefaciens* strain GV3101 by electroporation and transgenic lines were generated by floral dip[86]. Transformants were selected on half-strength Murashige and Skoog ($^{1/2}$MS) media containing 1% (w/v) sucrose, 0.8% (w/v) agar and 15 mg/L phosphinothricin.

BR root growth inhibition and competition assays

For BR-mediated root growth inhibition assays[43,44], surface sterilized seeds were stratified at 4°C for 2 d and plated on $^{1/2}$MS media plates containing 1% (w/v) sucrose and 0.8% (w/v) agar, supplemented with 100 nM of the indicated BR from 10 mM stock solutions prepared in 100 % (v/v) DMSO (Carl ROTH) or, with 0.01% (v/v) DMSO as control. The previously reported *bsk3-1* T-DNA insertion mutant (SALK_096500C)[75] was used as negative control. For the BR antagonist competition assays, 100 nM brassinolide and 100, 500 or 1000 nM of antagonist candidate were added to $^{1/2}$MS plates containing 1% (w/v) sucrose and 0.8% (w/v) agar. DMSO to a final concentration of 0.02% (v/v) was used as control treatment. In all cases, plates were moved to a Percival (SE-41AR3; CLF PlantClimatics, settings: 70% humidity, 16 h light period at 22°C with 250 µmol m^{-2} s^{-1} light intensity and 8 h dark period at 18°C), vertically placed for 7 d. The plates were scanned using a flat-bed scanner (Epson perfection V600 Photo, Epson). The primary root length of 30-35 individual seedlings was then determined using Fiji[87].

Statistical analysis

Multiple comparisons of different treatments vs. control(s) in one-way layouts were performed using a two-sided Dunnett[88] test as implemented in the R package multcomp[89] . The relationship

between root length and dissociation constant were tested using a Tukey trend test[90,91], i.e. a multiple

regression model approach using arithmetic, ordinal or logarithmic scores. We chose the linear

regression model and represented the corresponding slope b (positive slopes were interpreted as

antagonistic; negative slopes as agonistic) and associated p-values. Data were plotted in R[92] (version

4.2.2), the associated raw data can be found in Extended Data Tables 2, 6 and 8.

<u>BR in vivo quantification</u>

The extraction and quantification of endogenous BRs were carried out as previously

described[93]. In brief, five biological replicates, each consisting of approximately 10 mg fresh weight

(FW) of plant material, were extracted using ice-cold 60% (v/v) acetonitrile. To each sample, 25

pmol of deuterium-labelled BR internal standards (Olchemim LtD, Olomouc, Czech Republic) were

added. After a 12h incubation at 4°C, samples were centrifuged, and the resulting supernatants were

purified using 50 mg Discovery DPA-6S solid-phase extraction cartridges (Supelco, Bellefonte, PA,

USA). The eluates were then evaporated to dryness and reconstituted in 40 µL of methanol. BR

quantification was performed using ultra-high-performance liquid chromatography coupled with

tandem mass spectrometry (UHPLC-MS/MS), employing the ACQUITY UPLC I-Class System

(Waters, Milford, MA, USA) and the Xevo TQ-S triple quadrupole mass spectrometer (Waters MS

Technologies, Manchester, UK), using the previously reported UHPLC-MS/MS parameters[93,94].

<u>RNA-seq experiments</u>

10-d-old Arabidopsis plants grown on $^{1/2}$MS media plates containing 1% (w/v) sucrose and

0.8% (w/v) agar were harvested. 15 plants were pooled per individual biological replicate and all

samples were ground. RNA was extracted using a RNeasy Plant Mini Kit (74904; Qiagen) and total

RNA concentration was estimated using a Qubit™ 4 Fluorometer (Q33238; Thermo Fisher

Scientific) and a 2100 Bioanalyzer system (Agilent Technologies). Library preparation and

sequencing (100 bp single-end with Novaseq 6000 sequencing system; Illumina) was performed at

the iGE3 Genomic Platform at the Faculty of Medicine, University of Geneva,, Switzerland

(https://ige3.genomics.unige.ch/). Data analysis was conducted within the Galaxy EU platform

(https://usegalaxy.eu/) using the following tools: quality control and adaptor trimming by

MultiQC[95], reads mapping to the reference genome (Arabidopsis TAIR10 reference genome from

TAIR; https://www.arabidopsis.org/) by HISAT2[96], transcript assembly and FPKM estimates by

Cufflinks[97], reads counting by FeatureCounts[98] and differentially gene expression analysis by

DESeq2[99]. Visualization was done using SRplot[100] (https://www.bioinformatics.com.cn/).

 Confocal microscopy

To detect BRL2 promoter activity with *BRL2prom*::NLS-3xmVenus (in T2 generation) in the

early vasculature of embryos, SCRI Renaissance 2200 staining was performed according to the

protocol[83,101]. The embryos were dissected from the siliques in the flowering 5-week-old plants

under a Leica S6D stereo microscope (Leica S6D, Wetzlar, Germany) and incubated in an SR2200

solution containing 0.1% (v/v) SR2200, 1% (v/v) DMSO, and 0.5% (v/v) Triton X-100, 5% (v/v)

glycerol, and 4% (v/v) paraformaldehyde in phosphate-buffered saline buffer (pH 8.0) for five min.

Then, the embryos were washed once with H_2O and mounted in 5% (v/v) glycerol for imaging. The

cell wall signal stained by SR2200 was obtained using a 405 nm laser and a detection wavelength

ranging from 415 to 475 nm. To image roots, propidium iodide (PI) staining was used to label the

cell wall. The primary root of 10-d-old seedlings previously grown on $^{1/2}$MS plates containing 1%

(w/v) sucrose and 0.8% (w/v) agar was incubated in a 10 mg/L PI solution for 10 minutes. Then, it

was washed three times with water and incubated in a 5% (v/v) glycerol solution as the imaging

buffer. PI fluorescence was detected from 571 nm to 656 nm upon a 561 nm wavelength laser

excitation. The mVenus signal was detected using a laser with a 514 nm wavelength laser for

excitation and detectors with an emission range between 517 and 570 nm. To detect the mCitrine

signal in the roots of T2 generation *BRI1prom*::BRI1[mut]-mCitrine complementation lines, primary

root tips from 10-d-old seedlings growing on $^{1/2}$MS plates containing 1% (w/v) sucrose and 0.8%

(w/v) agar were sampled. The mCitrine signal was detected by a laser with 514 nm wavelength for

excitation, detectors recorded emission between 517 nm and 570 nm. All imaging was performed

with a LSM 780 NLO microscope (Zeiss) at the Photonic Bioimaging Center, University of

Geneva, Switzerland and data were analyzed using ZEN 2.0 blue edition and Fiji[87].

Analysis of leaf vein patterning

Leaf vein patterns were analyzed as described[76]. Cotyledons from 2-week-old seedlings

grown on $^{1/2}$MS plates containing 1% (w/v) sucrose and 0.8% (w/v) agar were dissected and cleared

in 95% ethanol, hydrated in an ethanol dilution series to water, and mounted on slides in 10% (v/v)

glycerol for imaging on a Nikon binocular microscope (Nikon SMZ18).

Hypocotyl growth assay

Surface-sterilized Col-0 wild-type and *brl2-1* mutant[76] seeds were cold-stratified at 4 °C for 48 h

and sown on $^{1/2}$MS medium containing 0.8% (w/v) agar. The growth medium was supplemented

with either: 1 μM brassinazole (BRZ)[102] prepared from a 10 mM stock solution in 100% DMSO, or

0.1% (v/v) DMSO as control. After a 1 h light treatment to promote germination, the plates were

wrapped in aluminium foil and maintained in complete darkness at 22 °C for 5 d. Seedling where

then scanned using a flat-bed scanner (Epson Perfection V600 Photo scanner, Epson).

<u>BES1 and BRI1-mCitrine immunoblotting</u>

Arabidopsis seeds (Col-0 wild type, *bri1*[47], *brl2-1*[76] and T3 generation *BRI1prom*::BRI1[mut]-

mCitrine complementation lines) were surface-sterilized and grown on $^{1/2}$MS plates containing 1%

(w/v) sucrose and 0.8% (w/v) agar for 10 or 14 d, respectively. Around 250 mg of tissue samples

were collected for each line in 1.5 ml tubes containing glass beads and immediately frozen with

liquid nitrogen. The samples were ground with Silamat S6 (Ivolar Vivodent) and then mixed with

extraction buffer (50 mM Tris-HCl pH 7.5, 150 mM NaCl, 1 mM EDTA, 1% [v/v] Triton X-100, 1x

cOmplete EDTA-free protease inhibitor cocktail [11873580001; Roche], 1x plant protease inhibitor

cocktail [P9599; Sigma-Aldrich] and 1x PhosStop [4906845001; Roche]). After protein

solubilization for 30 min on ice, cell debris was removed by centrifugation (14,000 xg, 2x 15 min,

4˚C) and the supernatant of each sample was transferred to a fresh tube. Protein concentration

determination with Pierce BCA Protein Assay Kit (23225; Thermo Fisher Scientific). For BRI1-

mCitrine fusion protein detection, an anti-GFP antibody coupled to horseradish peroxidase (anti-

GFP-HRP, 130-091-833; Miltenyi Biotec) at a dilution of 1:5,000 was applied. Alternatively, BRI1

was detected using a primary BRI1 antibody[103] (1:5000 dilution) and a secondary Anti-Rabbit IgG–

Peroxidase antibody produced in goat (1:10000 dilution, A6154; Sigma-Aldrich). BES1

dephosphorylation levels were estimated using a primary anti-BES1 antibody[41] at a dilution of

1:2000 and a secondary Anti-Rabbit IgG–Peroxidase antibody produced in goat (1:10000 dilution,

A6154; Sigma-Aldrich). SuperSignal West Pico PLUS Chemiluminescent Substrate (34577;

Thermo Fisher Scientific) was used for chemiluminescence detection on an Amersham Imager 680

(GE). ReadyBlue Protein Gel Stain (RSB-1L; Sigma-Aldrich) was used for the total protein input

detection.

<u>Analytical size-exclusion chromatography</u>

The oligomeric state of recombinantly expressed and purified wild-type and point-mutant

BRI1 ectodomains was characterized using analytical size-exclusion chromatography. 500 µL

sample at 1.0 mg/mL was loaded onto a Superdex 200 Increase 10/300 GL column (Cytiva) pre-

equilibrated in 20 mM citrate buffer (pH 5.0), 150 mM NaCl. Protein elution at a flow rate of

0.7 mL/min was monitored by UV absorption at 280 nm.

<u>Quantitative real-time PCR</u>

Expression of the BR marker genes CPD and DWF4 was estimated by qRT-PCR, using ~ 30

individual 8-d-old seedlings per each T3 generation transgenic line. Samples were homogenized by

Silamat S6 (Ivolar Vivodent) and RNA was extracted using a RNeasy Plant Mini Kit (74904;

Qiagen). Subsequently, cDNA synthesis was performed with the SuperScript™ II Reverse

Transcriptase (18064014; Invitrogen), following the manufacturer's protocol. Quantitative Real-

time PCR was conducted on a QuantStudio 5 thermo-cycler (Applied Biosystems) using SYBR™

Select Master Mix for CFX (4472942; Applied Biosystems). The relative expression (n=3) was

quantified with the $2^{-\Delta\Delta Ct}$ method. ACT2 was used as the internal reference gene. qPCR Primers

are included in Supplementary Table 4.

Figure legends

Fig. 1 | The branched architecture of the BR biosynthesis pathway. a, Chemical structure of

brassinolide with labeled steroidal rings (letters) and carbon atoms (numbers). **b**, Schematic

representation of the BR biosynthesis pathway, highlighting the relative positions of the BRs

discussed in this study. The C_{28}, C_{29}, and C_{27} branches are shaded from light to dark gray,

respectively. Key chemical differences between branches are indicated in blue, while red marks

chemical modifications along each biosynthetic route. Solid lines denote single-step enzymatic

reactions; dotted lines indicate multi-step conversions. Known enzymes involved in the canonical

C_{28} pathway are shown alongside.

Fig. 2 | BRI1, BRL1 and BRL3 bind chemically diverse BRs. a, GCI analysis of ligand-binding

kinetics between BRI1, BRL1, BRL3 and brassinolide. Shown are sensorgrams (n=2, red) with

global fits (black) and equilibrium dissociation constants (K_d) indicated. **b,** BR-binding pockets of

BRI1 (left; PDB-ID: pdb_00003rj0) and BRL3 (right) shown as ribbon diagrams. LRRs in blue,

island domains in red, brassinolide in yellow. Pocket residues and hydrogen bonds (dotted lines)

shown; conserved C_{23}-OH hydrogen bond highlighted in purple. **c,** Isoaffinity plots of association

(k_a) and dissociation (k_d) rates for receptor - BR pairs. Each point (n = 2) reflects a global fit;

diagonal lines denote K_d. High- and low-affinity interactions are in blue and red, respectively. **d,**

Structures of tested BRs, numbered as in **c**, with BRI1 K_d values. Red highlights indicate

modifications that reduce binding, while blue highlights indicate modifications that have a minimal

effect.

Fig. 3 | The hormone binding pocket can accommodate structurally diverse BRs. a, Structural superposition of different BRI1 - BR (top, r.m.s.d. ~0.4 Å comparing 186 corresponding C_α atoms) and BRL3 - BR complexes (bottom, ~r.m.s.d. ~0.3 Å comparing 190 corresponding C_α atoms). The receptors are shown in combined ribbon diagram (LRR domain in light blue, island domain in light red) and molecular surface views. BRs are in bonds representation color-coded according to their receptor binding affinities, ranging from low (light red) to high affinity (purple; BRI1 - brassinolide complex PDB-ID: pdb_00003rj0). Dashed lines highlight key polar interactions among the ligand and the protein receptor binding pockets. A conserved hydrogen bond between brassinolide and the island domains of BRI1, BRL1, and BRL3 is marked in magenta. **b,** Structures of BRs (in bonds representation, color-coded as in **a** with BRI1 (top) or BRL3 (bottom) and including final $mF_o–DF_c$ omit electron density maps (gray mesh, contoured at the indicated σ level). The respective chemical structures are shown above, with modifications relative to brassinolide marked in red (for changes that reduce binding affinity), or blue (neutral). **c,** Root growth inhibition assay of 7-d-old Col-0 wild-type and *bsk3-1*[75] mutant seedlings (n=31-35) grown on $^{1/2}$MS plates supplemented with the indicated BR (brassinolide, BL; 28-norbrassinolide, 28norBL; 28-homobrassinolide, 28hBL; 23-deoxycastasterone, 23dCS, 0.1% [v/v] DMSO as control) at a final concentration of 100 nM (scale bar = 1 cm). Quantifications (n=30) are shown as box plots (centre line, median; box, interquartile range IQR; whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data shown a gray dots). Multiple comparison of the different treatments vs. the DMSO control were performed using a Dunnett[88] test as implemented in the R package multcomp[89] (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). The relationship between the estimated dissociation constant (K_d) for each ligand and root length were tested using a Tukey trend test[90,91]. The blue line represent the linear trend with slope (b) and the associated test p-value (*$p < 0.05$; **$p < 0.01$; ***$p < 0.001$). **d,** BR quantification in 10 d old Col-0 and *bri1* (GABI_134E10[47]) seedlings (FW, fresh weight). **e,** RNA-seq derived expression profile heat map of BR biosynthesis, catabolism, transport and receptor genes in 10 d old seedlings in untreated BR signaling mutants and the Col-0 wild type. Expression values were transformed into Z-scores based on the normalized counts, three biological replicates are shown in three horizontal rows, the color scale indicates expression level (red, up-regulation, blue, down-regulation).

Fig. 4 | BRL2 is a medium-affinity BR receptor. a, Structural superposition of the BRI1 ectodomain (ribbon diagram, in gray; PDB-ID: pdb_00003rj0) with the AlphaFold2 model of the apo BRL2 ectodomain (in blue, AlphaFoldDB-ID: AF-Q9ZPS9-F1, r.m.s.d. ~ 0.8 Å comparing 163

C_α atoms). Brassinolide is in bonds representation, interacting residue are shown as ball-and-sticks.

b, GCI-derived binding kinetics of the BRL2 ectodomain vs. brassinolide, castasterone and

typhasterol. Red traces represent experimental sensorgrams (n=2), and black lines show the global

fits. **c,** Dissociation constant (K_d) table of the BRL2 ectodomain vs. different BRs, non-BR steroids

and   triterpenpoids (n.d., no binding detected). **d,** Confocal images of a representative

BRL2prom::NLS-3xmVenus reporter (in green) line in different stages: heart-stage embryo, mature

embryo (cell wall stained with Renaissance SR2200, gray, scale bar = 20 µm), cotyledon (scale bar

= 200 µm) and root from a 10-d-old seedling (cell wall stained with propidium iodide, magenta,

scale bar = 30 µm). **e,** Rosette phenotypes of 5-week-old plants, germinated on $^{1/2}$MS for 2 weeks

before transferring to soil for additional 3 weeks (scale bar = 2 cm).

Fig. 5 | Conserved aromatic residues in the hormone binding pocket are critical for BR

perception. a, Ribbon diagram of BRI1 (LRR domain in blue, island domain in red) in complex

with brassinolide (in bonds representation, PDB-ID: pdb_00003rj0). Mutated residues are displayed

as ball-and-sticks. **b,** GCI-derived binding kinetics for BRI1^{W564H}, BRI1^{G644D}, BRI1^{M657A} and

BRI1^{F681A} vs. brassinolide. Red traces represent experimental data (n=2), black lines show the global

fits. **c,** Dissociation constant (K_d) table for BRI1 point mutants vs. brassinolide. **d,** Rosette

phenotypes of representative, 5-week-old transgenic T3 *BRI1prom*::BRI1-Citrine^{mut}

complementation lines in *bri1* null[47] background compared to the Col-0 wild-type control (scale bar

= 2 cm). **e,** Quantification of BRI1^{mut}-mCitrine (theoretical molecular mass ~ 158.6 kDa) protein

levels by western blot from 10-d-old seedling protein extracts, immuno-blotted with an anti-GFP

antibody. ReadyBlue-stained RuBisCO is shown below as input loading control. **f,** Root length

phenotypes of 8-d-old seedlings (n=30) shown as box plots (centre line, median; box, interquartile

range IQR; whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data

shown a gray dots). Multiple comparison of the different treatments vs. the indicated controls were

performed using a Dunnett[88] test as implemented in the R package multcomp[89] (multiple

comparison against *bri1*: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; against Col-0 # $p < 0.05$; ## $p < 0.01$;

$p < 0.001$).

Fig. 6 | BR C_2 hydroxylation and C_6 oxidation are critical for SERK co-receptor binding. a,

Close-up view of the SERK1-bound (in blue) BRI1 (in gray) - brassinolide (in bonds representation,

in yellow) complex (PDB-ID: 00004lsx). Receptor and co-receptor amino acids involved in BR-

binding are highlighted as ball-and-sticks in gray and blue, respectively. Polar interactions between

the brassinolide C_2 hydroxyl group and His62, or the C_6 oxygen with Tyr642 and Lys601 are shown

as dashed lines (in black). **b,** GCI binding kinetics of amine-coupled and brassinolide-bound BRI1,

BRL1, BRL2, and BRL3 ectodomains vs. SERK1. Shown are sensorgrams with red traces

indicating experimental data and black lines representing global fits (n=1). Binding affinities (K_d)

are reported alongside. **c,** Representative GCI traces showing the interaction between SERK1 and

biotinylated, Avi-tagged BRI1 ectodomain in the presence of either brassinolide, typhasterol, 3-

deoxybrassinolide, or 6-deoxocastasterone. **d,** Table summaries of the receptor - BR - co-receptor

interaction dissociation constants. **e,** Chemical structures of selected BRs tested in **c**,**d**. Structural

modifications relative to brassinolide that reduce co-receptor binding are highlighted in red.

Modifications that have little effect on complex formation are shown in blue. **f,** GCI binding

kinetics of amine-coupled wild-type BRI1, BRI1^{Y597F}, BRI1^{Y599A}, BRI1^{K601L}, and BRI1^{Y642A}

ectodomains vs. SERK1 in the presence of a high, saturating concentration of brassinolide. Shown

are sensorgrams with red traces indicating experimental data and black lines representing global fits

(n=2). Binding affinities (K_d) are reported alongside. **g,** BR seedling root growth inhibition

antagonist assay (n=30-35). Brassinolide (BL) at a fixed concentration of 100 nM was used in the

different treatments, vs. 0.02% (v/v) DMSO as control (in yellow). The antagonist candidate

compounds secasterone, 2-deoxybrassinolide and 3-deoxycastasterone were tested for their ability

to inhibit seedling root growth by themselves (in blue), or to reverse BL-induced root growth

inhibition (in orange). Shown are boxplots with centre lines (medians), box limits indicate the 25th

and 75th percentiles, whiskers extend to the full data range. The relationship between antagonist

concentration and root length were tested using a Tukey trend test[90,91]. The blue line represent the

linear trend with slope b (negative values indicate agonistic behavior, positive values indicate

antagonistic behavior) and the associated test p-value ($*p < 0.05$; $**p < 0.01$; $***p < 0.001$).

Representative seedling images are shown above (scale bar = 0.5 cm). The DMSO and BL 100 nM

controls are same in each panel, indicated by a white vertical line.

Fig 7 | Strong receptor binding, correct positioning within the BR binding site, and interaction

with the co-receptor are essential for BR bioactivity. Schematic overview of brassinosteroid-

induced receptor - co-receptor complex formation and activation. A BR ligand binds to the BRI1

receptor ectodomain (in gray), creating a binding surface for a SERK co-receptor kinase (in blue).

This brings the kinase domains of receptor and co-receptor in close proximity, leading to trans-

phosphorylation and intracellular signaling. The BR-binding pocket is formed by the receptor (in

gray) and co-receptor (in blue). Chemical features of the BR ligand (in black) involved in co-

receptor binding are highlighted in dark-blue, key carbon atoms critical for receptor or co-receptor interaction are numbered, chemical features required for receptor and/or co-receptor binding are shown in red. The approximate positions of residues critically involved in BR binding, positioning and co-receptor binding are indicated.

Supplementary Fig. 1 | BRI1, BRL1 and BRL3 share a structurally conserved BR-binding pocket. a, Right: Ribbon diagram of the BRL3 ectodomain bound to brassinolide (yellow, bonds representation). Left: Close-up of a structural superposition of the BR-binding pockets of BRL3 (blue), BRI1 (purple; r.m.s.d. ~ 0.7 Å over 172 C_α atoms), and BRL1 (yellow; r.m.s.d. ~ 0.6 Å over 166 C_α atoms). Key residues involved in brassinolide coordination are shown in ball-and-stick representation. The conserved hydrogen bond between the C_{23} hydroxyl group of brassinolide and the island domain backbone is indicated with a black dotted line. **b,** Multiple sequence alignment of BR-binding pocket regions from *Arabidopsis thaliana* BRI1, BRL1, and BRL3, along with orthologs from rice (OsBRI1, https://www.uniprot.org/ UniProt-ID Q942F3), maize (ZmBRI1, A0A3L6FIW6), and tomato (SlBRI1, F2XYF6), generated using MUSCLE[104]. Secondary structure elements based on AtBRI1 and calculated with DSSP[105] are shown above the alignment. BR-contacting residues (compare Fig. 2b) are highlighted in gray; residues mutated in this study (compare Fig. 4a) are shown in red.

Supplementary Fig 2. | BR catabolic enzyme modify BRs at positions critical for receptor and co-receptor binding. Left panel: Molecular surface view of the BRI1 hormone binding pocket (in gray) occupied by brassinolide (in bonds representation, in yellow; PDB-ID: pdb_00003rj0). The A, B, C, and D rings that make up the hormone's steroid core are labeled and carbon atoms targeted for catabolic modifications are highlighted. Right panel: Chemical structure of brassinolide with carbon numbering. Red, bold numbers indicate sites of BR catabolic reactions: C_{26} hydroxylation by BAS1[17], C_{23} O-glucosylation by UGT73C5[18], C_{22} O-sulfonation by AtST1/ST4a[19], C_6 reduction by BEN1[106] and C_3 O-laurylation by PIZZA/BAT1/DRL1[107–109].

Supplementary Fig. 3 | BRL2 harbors a conserved BR-binding pocket but has not detectable function in BR growth signaling. a, Structural superposition of the BRI1 ectodomain (ribbon diagram, in orange; PDB-ID: pdb_00003rj0) with the apo BRL2 crystal structure (in blue, r.m.s.d. ~ 0.8 Å comparing 314 C_α atoms). **b,** Close-up view of the BRL2 BR-binding pocket (in blue, AlphaFoldDB-ID: AF-Q9ZPS9-F1) superimposed with BRL3 (right panel, in green, brassinolide in

bonds representation, r.m.s.d. ~ 0.7 Å comparing 162 C$_\alpha$ atoms). Residues lining the BR binding

site are shown in ball-and-stick representation, with corresponding residues in BRL2 shown in blue.

Non-synonymous residues among BRI1, BRL2 and BRL3 are labeled. **c,** Schematic diagram of

brl2-2 CRISPR/Cas9 deletion allele. The BRL2 genomic locus is shown, including the translation

start codon (black arrow), exon (black rectangle) and 5' and 3- primer untranslated regions (UTRs,

gray blocks). The putative binding sites of the designed guide RNAs are indicated in red. **d,**

Cotyledon vein patterning analysis performed using 2-week-old Col-0 and *brl2-1* seedlings. Leaf

patterns are categorized in seven groups, indicated 4 (4 distinct regions are formed by the

intersection of the leaf veins, blue), 3+1 (3 distinct regions formed with 1 free branch, beige), 3 (3

distinct regions formed, lemon green), 2+2 (2 distinct regions formed with 2 free branches, light

purple), 2+1 (2 distinct regions formed with 1 free branch, canary), 2 (2 distinct regions formed,

gray) and gap (no distinct region formed, earthy yellow). **e,** Hypocotyl growth assay of dark-grown

brassinazole (BRZ)[102] treated vs untreated control seedlings. Seeds of Col-0 and *brl2-1* were

germinated and grown on $^{1/2}$MS plates supplemented with 1 μM BRZ or 0.1% (v/v) DMSO (scale

bar = 1 cm). f, BES1 (theoretical molecular mass ~ 39.2 kDa) western blot. 14-d-old Col-0, *brl2-1*

and *bri1* seedlings were grown on $^{1/2}$MS plates with 1% (w/v) sucrose, transferred to $^{1/2}$MS liquid

medium containing 1% (w/v) sucrose and either 1 μM brassinolide (BL) or 0.01 % DMSO (as

control) for 90 min, and probed with an anti-BES1 antibody[41] by western blotting.

Supplementary Fig. 4 | Mutations of conserved residues in the BRI1 hormone binding pocket

alter BR signaling without compromising the plasma-membrane localization of the mutant

receptor. a, Structural superposition of the brassinolide-bound AtBRI1 hormone binding pocket

(*Arabidopsis thaliana*, in orange, PDB-ID: pdb_00003rj0) with the corresponding regions in

ZmBRI1 (*Zea mays*, in blue, AlphaFoldDB-ID: AF-A0A3L6FIW6-F1, r.m.s.d ~ 0.7 Å between 175

C$_\alpha$ atoms), SlBRI1 (*Solanum lycopersicum*, in purple, AlphaFoldDB-ID: AF-F2XYF6-F1, r.m.s.d ~

0.6 Å between 178 C$_\alpha$ atoms), and OsBRI1 (*Oryza sativa*, in green, AlphaFoldDB-ID: AF-Q942F3-

F1, r.m.s.d ~ 0.7 Å between 170 C$_\alpha$ atoms). Mutated residues in AtBRI1 are displayed as ball-and-

stick models, the corresponding amino acid side-chains from ZmBRI1, SlBRI1, and OsBRI1 are

included. **b,** Analytical size-exclusion chromatography elution profiles for recombinant wild-type

(WT) and point-mutant BRI1 ectodomains used in the GCI assays. **c,** BRI1 protein level in Col-0,

bri1 and *BRI1prom*::BRI1-mCitrine / *bri1* lines, detected with a primary anti-BRI1 antibody[103].

ReadyBlue-stained RuBisCO is shown as loading control below. **d,** Confocal imaging of T2

generation *BRI1prom*::BRI1^{mut}-mCitrine signal in the mersistematic zone of 10-d-old primary roots

mounted for confocal imaging (scale bar = 20 μm). **e,** Brassinolide (BL)-stimulated BES1 dephosphorylation. 14-d-old seedlings of Col-0 wild-type, *bri1* and *BRI1prom*::BRI1[mut]-mCit T3 complementation lines were treated with 0.1% (v/v) DMSO or 100 nM BL and 1000 nM BL for 60 min before protein extraction and anti-BES1 antibody[41] immuno-blotting. The upper band represents phosphorylated BES1, the lower band (more) dephosphorylated BES1. ReadyBlue-stained RuBisCO is shown as loading control below. **f,** qRT-PCR analysis of the BR marker genes CPD and DWF4 in Col-0 wild-type, *bri1* and *BRI1prom*::BRI1[mut]-mCit T3 complementation, with ACT2 used as internal reference gene. Each dot represents a biological replicate (n=3).

Supplementary Table 1 | Crystallographic data collection and refinement statistics

Supplementary Table 2 | List of genotyping primers used in this study

Supplementary Table 3 | Cloning primers used in this study

Supplementary Table 4 | qPCR primers for BR marker genes CPD and DWF4

Extended Data Fig. 1 | Complete set of GCI sensorgrams for the experiments described in Fig. 2a, c, d. Grating Coupling Interferometry (GCI)-derived binding kinetics for BRI1, BRL1, and BRL3 in response to different BRs. Raw sensorgrams are shown with experimental traces in red, and global fits as black lines. CATH: cathasterone, 23dCS: 23-deoxycastasterone, 24eCS: 24-epicastasterone, 28hCS: 28-homocastasterone, 24eBL: 24-epibrassinolide, 28hBL: 28-homobrassinolide, 22dCS: 22-deoxycastasterone, 6dCS: 6-deoxocastasterone, NorBL: 28-norbrassinolide, 3dCS: 3-deoxycastasterone, TYPH: typhasterol, CS: castasterone, SEC: secasterone, 2dBL: 2-deoxybrassinolide, BL: brassinolide.

Extended Data Fig. 2 | Complete set of GCI kinetic parameters for the experiments described in Fig. 2a, c, d. a, Schematic illustration of the Grating Coupling Interferometry (GCI) setup used to screen receptor - brassinosteroid interactions. **b,** Kinetic parameters obtained for BRI1, BRL1, and BRL3 in response to the 15 BRs analyzed in this study.

Extended Data Fig. 3 | Complete set of GCI sensorgrams and derived kinetic parameters for the experiments described in Fig. 4b, c. a, Grating Coupling Interferometry (GCI)-derived

binding kinetics for BRL2 in response to different BRs and other plant steroids. Raw sensorgrams are shown with experimental traces in red (n=2), and global fits as black lines (n.d., no detectable binding). BL: brassinolide, 2dBL: 2-deoxybrassinolide, CS: castasterone, TYPH: typhasterol, 28norBL: 28-norbrassinolide, 28hBL: 28-homobrassinolide, 28hCS: 28-homocastasterone, CATH: cathasterone, 24eCS: 24-epicastasterone, 24eBL: 24-epibrassinolide, DHEA: dehydroepiandrosterone. **b,** Kinetic parameters obtained for BRL2 in response to the different BR and non-BR ligand candidates analyzed in this study. **c,** Chemical structures of dehydroepiandrosterone (DHEA), triolone, lupeol and betulin[110–113].

Extended Data Fig. 4 | Complete set of GCI sensorgrams and derived kinetic parameters for the experiments described in Fig. 5b,c. a, Grating Coupling Interferometry (GCI)-derived binding kinetics for different BRI1 ectodomains carrying point mutations in the BR-binding pocket vs. brassinolide. Raw sensorgrams are shown with experimental traces in red, and global fits as black lines. **b,** Table summaries of kinetic parameters obtained for the experiments shown in **a**.

Extended Data Fig. 5 | Complete set of GCI sensorgrams and derived kinetic parameters for experiments shown in Fig. 6b-f. a, Table summaries of kinetic parameters obtained for the experiments shown in Fig. 6b. **b,** Schematic overview of the GCI receptor capturing and SERK1 binding assay. For Avitag based coupling, streptavidin (SA, shown as a tetramer) was immobilized using the amine-coupling method on the GCI chip (black line). Next, the biotinylated avi-tagged BR receptor ectodomain was captured by streptavidin, followed by passivation and quenching of the surface with BSA (in light blue). Then, in the presence of a constant concentration of brassinolide in the analyte buffer (in yellow), SERK1 (in dark blue) association and dissociation at different protein concentrations was performed. **c,** Grating Coupling Interferometry (GCI)-derived binding kinetics for the avi-tagged BRI1 ectodomain vs. different antagonist candidate ligands including 6dCS: 6-deoxocastasterone, SEC: secasterone, TYPH: typhasterol, 3dCS: 3-deoxocastasterone, CS: castasterone, BL: brassinolide. Raw sensorgrams are shown with experimental traces in red, and global fits as black lines. **d,** Table summaries of kinetic parameters obtained for the experiments shown in Fig. 6c,d. **e,** Schematic overview of the GCI receptor capturing and SERK1 binding assay to analyze the effect of BR binding site missense mutations in BRI1 for SERK1 co-receptor binding (related to Fig. 6f). BRI1[WT], BRI1[Y597F], BRI1[Y599A], BRI1[K601L] and BRI1[Y642A] were immobilized by amine-coupling on the GCI chip (black line). SERK1 binding to the receptors (in gray) was then quantified in the presence of a constant, high concentration of brassinolide (in yellow) in the

running buffer and by injecting the co-receptor at different concentrations (in blue). **f,** Table

summaries of kinetic parameters obtained for the experiments shown in Fig. 6f.

Extended Data Table 1 | Immobilized receptor density and analyte concentrations used for the

GCI assays described in Fig. 2a,c,d.

Extended Data Table 2 | BR-mediated root growth inhibition raw data corresponding to Fig.

3c.

Extended Data Table 3 | RNA-seq raw counts used to calculate the heat map shown in Fig. 3e.

Extended Data Table 4 | Immobilized receptor density and analyte concentrations used for the

GCI assays described in Fig. 4b,c.

Extended Data Table 5 | Immobilized receptor density and analyte concentrations used for the

GCI assays described in Fig. 5.b,c.

Extended Data Table 6 | Wild-type and BRI1^{mut} seedling root length raw data corresponding

to Fig. 5f.

Extended Data Table 7 | Immobilized receptor density and analyte concentrations used for the

GCI assays described in Fig. 6b,c,d and g.

Extended Data Table 8 | BR-mediated root growth inhibition antagonist assay raw data

corresponding to Fig. 6g.

Data availability

Raw RNA-seq reads have been deposited with the NCBI sequence read archive (SRA) with

identifier PRJNA1299306 (https://www.ncbi.nlm.nih.gov/sra/PRJNA1299306). BRI1, BRL2 and

BRL3 complex structures have been deposited with the Protein Data Bank (http://www.rcsb.org)

with accession numbers pdb_00009S80 (BRL3 – brassinolide), pdb_00009S87 (BRI1 –

castasterone), pdb_00009S8S (BRI1 – typhasterol), pdb_00009S8V (BRI1 – 24-epibrassinolide),

pdb_00009S8Z (BRI1 – 28-homobrassinolide), pdb_00009S90 (BRL3 – castasterone),

pdb_00009S96 (BRL3 – typhasterol), pdb_00009S9A (BRL3 – 6-deoxocastasterone),
pdb_00009S9C (apo BRL2).

Acknowledgments

We thank Jenny Russinova for providing us with the BES1 antibody, Danuše Tarkowská for
help with UHPLC-MS/MS analyses, Adam Přibylka for the measurement of HRMS spectra,
Samuele Meschini for help with the root growth inhibition assays and Philippe Rieu, Roman Ulm
and Dorothea Fiedler for providing feedback on the manuscript. We acknowledge the Paul Scherrer
Institute, Villigen, Switzerland for provision of synchrotron radiation beamtime at beamline X06DA
of the SLS. This work was supported by Grant 310030_205201 from the Swiss National Science
Foundation (to M.H.) and IGA grant IGA_PrF_2025_019 from Palacky University (to M.S.).

Author contributions

A.C. and M.H. conceived the project. A.C expressed and purified proteins, performed GCI
assays and crystallized proteins. H.C. generated transgenic lines with the help of L.B., performed
plant phenotyping and BR-response assays, confocal microscopy, transcriptomics and western
blotting. M.K. and K.F. synthesized BRs. U.H. crystallized the BRL3 – brassinolide complex. J.O.
quantified endogenous BR levels. A.C., U.H. and M.H. solved and refined structures. A.C., H.C.,
M.K., J.O., L.H., M.S. and M.H. analyzed data. A.C. and M.H. wrote the manuscript, with input
from the other authors. M.H. and M.S. acquired funding and M.H. supervised the project.

References

1. Grove, M. D. *et al.* Brassinolide, a plant growth-promoting steroid isolated from Brassica napus pollen. *Nature* **281**, 216–217 (1979).

2. Fujioka, S. & Yokota, T. Biosynthesis and Metabolism of Brassinosteroids. *Annual Review of Plant Biology* **54**, 137–164 (2003).

3. Fujioka, S., Takatsuto, S. & Yoshida, S. An Early C-22 Oxidation Branch in the Brassinosteroid Biosynthetic Pathway. *Plant Physiol* **130**, 930–939 (2002).

4. Joo, S.-H. *et al.* Biosynthesis of a cholesterol-derived brassinosteroid, 28-norcastasterone, in Arabidopsis thaliana. *Journal of Experimental Botany* **63**, 1823–1833 (2012).

5. Fujita, S. *et al.* Arabidopsis CYP90B1 catalyses the early C-22 hydroxylation of C27, C28 and C29 sterols. *The Plant Journal* **45**, 765–774 (2006).

6. Choe, S. *et al.* The DWF4 gene of Arabidopsis encodes a cytochrome P450 that mediates multiple 22alpha-hydroxylation steps in brassinosteroid biosynthesis. *Plant Cell* **10**, 231–243 (1998).

7. Ohnishi, T. *et al.* CYP90A1/CPD, a Brassinosteroid Biosynthetic Cytochrome P450 of Arabidopsis, Catalyzes C-3 Oxidation. *J Biol Chem* **287**, 31551–31560 (2012).

8. Szekeres, M. *et al.* Brassinosteroids Rescue the Deficiency of CYP90, a Cytochrome P450, Controlling Cell Elongation and De-etiolation in Arabidopsis. *Cell* **85**, 171–182 (1996).

9. Ohnishi, T. *et al.* C-23 Hydroxylation by Arabidopsis CYP90C1 and CYP90D1 Reveals a Novel Shortcut in Brassinosteroid Biosynthesis. *Plant Cell* **18**, 3275–3288 (2006).

10. Chory, J., Nagpal, P. & Peto, C. Phenotypic and Genetic Analysis of det2, a New Mutant That Affects Light-Regulated Seedling Development in Arabidopsis. *Plant Cell* **3**, 445–459 (1991).

11. Fujioka, S. *et al.* The Arabidopsis deetiolated2 mutant is blocked early in brassinosteroid biosynthesis. *Plant Cell* **9**, 1951–1962 (1997).

12. Noguchi, T. *et al.* Arabidopsis det2 Is Defective in the Conversion of (24R)-24-Methylcholest-4-En-3-One to (24R)-24-Methyl-5α-Cholestan-3-One in Brassinosteroid Biosynthesis. *Plant Physiol* **120**, 833–840 (1999).

13. Shimada, Y. *et al.* Brassinosteroid-6-Oxidases from Arabidopsis and Tomato Catalyze Multiple C-6 Oxidations in Brassinosteroid Biosynthesis. *Plant Physiol* **126**, 770–779 (2001).

14. Noguchi, T. *et al.* Biosynthetic pathways of brassinolide in Arabidopsis. *Plant Physiol* **124**, 201–209 (2000).

15. Katsumata, T. *et al.* Arabidopsis CYP85A2 Catalyzes Lactonization Reactions in the Biosynthesis of 2-Deoxy-7-oxalactone Brassinosteroids. *Bioscience, Biotechnology, and Biochemistry* **72**, 2110–2117 (2008).

16. Antonchick, A. P., Schneider, B., Zhabinskii, V. N., Konstantinova, O. V. & Khripach, V. A. Biosynthesis of 2,3-epoxybrassinosteroids in seedlings of Secale cereale. *Phytochemistry* **63**, 771–776 (2003).

17. Neff, M. M. *et al.* BAS1: A gene regulating brassinosteroid levels and light responsiveness in Arabidopsis. *Proc Natl Acad Sci U S A* **96**, 15316–15323 (1999).

18. Poppenberger, B. *et al.* The UGT73C5 of Arabidopsis thaliana glucosylates brassinosteroids. *Proc Natl Acad Sci U S A* **102**, 15253–15258 (2005).

19. Marsolais, F. *et al.* Molecular and biochemical characterization of two brassinosteroid sulfotransferases from Arabidopsis, AtST4a (At2g14920) and AtST1 (At2g03760). *Planta* **225**, 1233–1244 (2007).

20. Kim, H. B. *et al.* The Regulation of DWARF4 Expression Is Likely a Critical Mechanism in Maintaining the Homeostasis of Bioactive Brassinosteroids in Arabidopsis. *Plant Physiology* **140**, 548–557 (2006).

21. Northey, J. G. B. *et al.* Farnesylation mediates brassinosteroid biosynthesis to regulate abscisic acid responses. *Nat Plants* **2**, 16114 (2016).

22. Wang, Y. *et al.* Plasmodesmata mediate cell-to-cell transport of brassinosteroid hormones. *Nat Chem Biol* **19**, 1331–1341 (2023).

23. Ying, W. *et al.* Structure and function of the Arabidopsis ABC transporter ABCB19 in brassinosteroid export. *Science* **383**, eadj4591 (2024).

24. Wang, Z. Y., Seto, H., Fujioka, S., Yoshida, S. & Chory, J. BRI1 is a critical component of a plasma-membrane receptor for plant steroids. *Nature* **410**, 380–383 (2001).

25. Li, J. & Chory, J. A putative leucine-rich repeat receptor kinase involved in brassinosteroid signal transduction. *Cell* **90**, 929–938 (1997).

26. Hothorn, M. *et al.* Structural basis of steroid hormone perception by the receptor kinase BRI1. *Nature* **474**, 467–471 (2011).

27. She, J. *et al.* Structural insight into brassinosteroid perception by BRI1. *Nature* **474**, 472–476 (2011).

28. Hohmann, U. *et al.* Mechanistic basis for the activation of plant membrane receptor kinases by SERK-family co-receptors. *Proc Natl Acad Sci U S A* **115**, 3488–3493 (2018).

29. Caño-Delgado, A. *et al.* BRL1 and BRL3 are novel brassinosteroid receptors that function in vascular differentiation in Arabidopsis. *Development* **131**, 5341–5351 (2004).

30. Zhou, A., Wang, H., Walker, J. C. & Li, J. BRL1, a leucine-rich repeat receptor-like protein kinase, is functionally redundant with BRI1 in regulating Arabidopsis brassinosteroid signaling. *Plant J* **40**, 399–409 (2004).

31. She, J., Han, Z., Zhou, B. & Chai, J. Structural basis for differential recognition of brassinolide by its receptors. *Protein Cell* **4**, 475–482 (2013).

32. Zhang, X. *et al.* A Temperature-Sensitive Misfolded bri1-301 Receptor Requires Its Kinase Activity to Promote Growth. *Plant Physiology* **178**, 1704–1719 (2018).

33. Blanco-Touriñán, N. *et al.* The brassinosteroid receptor gene BRI1 safeguards cell-autonomous brassinosteroid signaling across tissues. *Sci Adv* **10**, eadq3352 (2024).

34. Clay, N. K. & Nelson, T. VH1, a provascular cell-specific receptor kinase that influences leaf cell patterns in Arabidopsis. *Plant Cell* **14**, 2707–2722 (2002).

35. Gou, X. *et al.* Genetic evidence for an indispensable role of somatic embryogenesis receptor kinases in brassinosteroid signaling. *PLoS Genet* **8**, e1002452 (2012).

36. Santiago, J., Henzler, C. & Hothorn, M. Molecular mechanism for plant steroid receptor activation by somatic embryogenesis co-receptor kinases. *Science* **341**, 889–892 (2013).

37. Imkampe, J. *et al.* The Arabidopsis Leucine-Rich Repeat Receptor Kinase BIR3 Negatively Regulates BAK1 Receptor Complex Formation and Stabilizes BAK1. *Plant Cell* **29**, 2285–2303 (2017).

38. Hohmann, U., Nicolet, J., Moretti, A., Hothorn, L. A. & Hothorn, M. The SERK3 elongated allele defines a role for BIR ectodomains in brassinosteroid signalling. *Nat Plants* **4**, 345–351 (2018).

39. Wang, X. *et al.* Sequential transphosphorylation of the BRI1/BAK1 receptor kinase complex impacts early events in brassinosteroid signaling. *Dev Cell* **15**, 220–235 (2008).

40. Wang, Z. Y. *et al.* Nuclear-localized BZR1 mediates brassinosteroid-induced growth and feedback suppression of brassinosteroid biosynthesis. *Dev Cell* **2**, 505–513 (2002).

41. Yin, Y. *et al.* BES1 accumulates in the nucleus in response to brassinosteroids to regulate gene expression and promote stem elongation. *Cell* **109**, 181–191 (2002).

42. Li, J., Nagpal, P., Vitart, V., McMorris, T. C. & Chory, J. A role for brassinosteroids in light-dependent development of Arabidopsis. *Science* **272**, 398–401 (1996).

43. Li, J., Nam, K. H., Vafeados, D. & Chory, J. BIN2, a new brassinosteroid-insensitive locus in Arabidopsis. *Plant Physiol* **127**, 14–22 (2001).

44. Müssig, C., Shin, G.-H. & Altmann, T. Brassinosteroids Promote Root Growth in Arabidopsis. *Plant Physiol* **133**, 1261–1271 (2003).

45. Noguchi, T. *et al.* Brassinosteroid-Insensitive Dwarf Mutants of Arabidopsis Accumulate Brassinosteroids. *Plant Physiol* **121**, 743–752 (1999).

46. Neubus Claus, L. A. *et al.* BRASSINOSTEROID INSENSITIVE1 internalization can occur independent of ligand binding. *Plant Physiol* **192**, 65–76 (2023).

47. Jaillais, Y., Belkhadir, Y., Balsemão-Pires, E., Dangl, J. L. & Chory, J. Extracellular leucine-rich repeats as a platform for receptor/coreceptor complex formation. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 8503–8507 (2011).

48. Nomura, T. *et al.* Accumulation of 6-deoxocathasterone and 6-deoxocastasterone in Arabidopsis, pea and tomato is suggestive of common rate-limiting steps in brassinosteroid biosynthesis. *Phytochemistry* **57**, 171–178 (2001).

49. Fujioka, S. *et al.* Identification of castasterone, 6-deoxocastasterone, typhasterol and 6-deoxotyphasterol from the shoots of Arabidopsis thaliana. *Plant Cell Physiol* **37**, 1201–1203 (1996).

50. Fujioka, S., Noguchi, T., Yokota, T., Takatsuto, S. & Yoshida, S. Brassinosteroids in *Arabidopsis thaliana*. *Phytochemistry* **48**, 595–599 (1998).

51. Vukašinović, N. *et al.* Local brassinosteroid biosynthesis enables optimal root growth. *Nat. Plants* **7**, 619–632 (2021).

52. Tanaka, K. *et al.* Brassinosteroid Homeostasis in Arabidopsis Is Ensured by Feedback Expressions of Multiple Genes Involved in Its Metabolism. *Plant Physiology* **138**, 1117–1125 (2005).

53. He, J.-X. *et al.* BZR1 is a transcriptional repressor with dual roles in brassinosteroid homeostasis and growth responses. *Science* **307**, 1634–1638 (2005).

54. Hohmann, U. *et al.* Constitutive Activation of Leucine-Rich Repeat Receptor Kinase Signaling Pathways by BAK1-INTERACTING RECEPTOR-LIKE KINASE3 Chimera. *Plant Cell* **32**, 3311–3323 (2020).

55. Back, T. G., Janzen, L., Pharis, R. P. & Yan, Z. Synthesis and bioactivity of C-2 and C-3 methyl ether derivatives of brassinolide. *Phytochemistry* **59**, 627–634 (2002).

56. Muto, T. & Todoroki, Y. Brassinolide-2,3-acetonide: a brassinolide-induced rice lamina joint inclination antagonist. *Bioorg Med Chem* **21**, 4413–4419 (2013).

57. Kim, S. *et al.* Chemical control of receptor kinase signaling by rapamycin-induced dimerization. *Mol Plant* **14**, 1379–1390 (2021).

58. Zheng, B. *et al.* Pan-brassinosteroid signaling revealed by functional analysis of NILR1 in land plants. *New Phytol* **235**, 1455–1469 (2022).

59. Ferreira-Guerra, M., Marquès-Bueno, M., Mora-García, S. & Caño-Delgado, A. I. Delving into the evolutionary origin of steroid sensing in plants. *Curr Opin Plant Biol* **57**, 87–95 (2020).

60. Li, W. *et al.* Two Conserved Amino Acids Characterized in the Island Domain Are Essential for the Biological Functions of Brassinolide Receptors. *International Journal of Molecular Sciences* **23**, 11454 (2022).

61. Hothorn, M., Dabi, T. & Chory, J. Structural basis for cytokinin recognition by Arabidopsis thaliana histidine kinase 4. *Nat Chem Biol* **7**, 766–768 (2011).

62. Okuda, S. *et al.* Molecular mechanism for the recognition of sequence-divergent CIF peptides by the plant receptor kinases GSO1/SGN3 and GSO2. *Proc Natl Acad Sci U S A* **117**, 2693–2703 (2020).

63. Olczak, M. & Olczak, T. Comparison of different signal peptides for protein secretion in nonlytic insect cell system. *Anal. Biochem* **359**, 45–53 (2006).

64. Hashimoto, Y., Zhang, S. & Blissard, G. W. Ao38, a new cell line from eggs of the black witch moth, Ascalapha odorata (Lepidoptera: Noctuidae), is permissive for AcMNPV infection and produces high levels of recombinant proteins. *BMC Biotechnol* **10**, 50 (2010).

65. Fairhead, M. & Howarth, M. Site-specific biotinylation of purified proteins using BirA. *Methods Mol. Biol.* **1266**, 171–184 (2015).

66. Kabsch, W. Automatic processing of rotation diffraction data from crystals of initially unknown symmetry and cell constants. *J. Appl. Crystallogr.* **26**, 795–800 (1993).

67. McCoy, A. J. *et al.* Phaser crystallographic software. *J. Appl. Crystallogr.* **40**, 658–674 (2007).

68. Moriarty, N. W., Grosse-Kunstleve, R. W. & Adams, P. D. electronic Ligand Builder and Optimization Workbench (eLBOW): a tool for ligand coordinate and restraint generation. *Acta Crystallogr D Biol Crystallogr* **65**, 1074–1080 (2009).

69. Emsley, P. & Cowtan, K. Coot: model-building tools for molecular graphics. *Acta Cryst D* **60**, 2126–2132 (2004).

70. Afonine, P. V. *et al.* Towards automated crystallographic structure refinement with phenix.refine. *Acta Cryst D* **68**, 352–367 (2012).

71. Davis, I. W. *et al.* MolProbity: all-atom contacts and structure validation for proteins and nucleic acids. *Nucleic Acids Res* **35**, W375-383 (2007).

72. Liebschner, D. *et al.* Polder maps: improving OMIT maps by excluding bulk solvent. *Acta Crystallogr D Struct Biol* **73**, 148–157 (2017).

73. Goddard, T. D. *et al.* UCSF ChimeraX: Meeting modern challenges in visualization and analysis. *Protein Sci* **27**, 14–25 (2018).

74. Ihlenfeldt, W. D., Bolton, E. E. & Bryant, S. H. The PubChem chemical structure sketcher. *J Cheminform* **1**, 20 (2009).

75. Tang, W. *et al.* BSKs mediate signal transduction from the receptor kinase BRI1 in Arabidopsis. *Science* **321**, 557–560 (2008).

76. Ceserani, T., Trofka, A., Gandotra, N. & Nelson, T. VH1/BRL2 receptor-like kinase interacts with vascular-specific adaptor proteins VIT and VIK to influence leaf venation. *Plant J* **57**, 1000–1014 (2009).

77. Han, J.-P. *et al.* GWAs reveals SUBER GENE1-mediated suberization via Type One Phosphatases. 2025.05.06.652434 Preprint at https://doi.org/10.1101/2025.05.06.652434 (2025).

78. Vermeer, J. E. M. *et al.* A spatial accommodation by neighboring cells is required for organ initiation in Arabidopsis. *Science* **343**, 178–183 (2014).

79. Kalmbach, L. *et al.* Putative pectate lyase PLL12 and callose deposition through polar CALS7 are necessary for long-distance phloem transport in Arabidopsis. *Curr Biol* **33**, 926-939.e9 (2023).

80. Karimi, M., De Meyer, B. & Hilson, P. Modular cloning in plant cells. *Trends Plant Sci* **10**, 103–105 (2005).

81. Ho, S. N., Hunt, H. D., Horton, R. M., Pullen, J. K. & Pease, L. R. Site-directed mutagenesis by overlap extension using the polymerase chain reaction. *Gene* **77**, 51–59 (1989).

82. Jinek, M. *et al.* A programmable dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* **337**, 816–821 (2012).

83. Chen, H. *et al.* Phosphorylation-dependent activation of the bHLH transcription factor ICE1/SCRM promotes polarization of the Arabidopsis zygote. *New Phytol* **245**, 1029–1039 (2025).

84. Liu, H. *et al.* CRISPR-P 2.0: An Improved CRISPR-Cas9 Tool for Genome Editing in Plants. *Mol Plant* **10**, 530–532 (2017).

85. Wu, R. *et al.* An efficient CRISPR vector toolbox for engineering large deletions in Arabidopsis thaliana. *Plant Methods* **14**, 65 (2018).

86. Clough, S. J. & Bent, A. F. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J.* **16**, 735–743 (1998).

87. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods* **9**, 676–682 (2012).

88. Dunnett, C. W. A Multiple Comparison Procedure for Comparing Several Treatments with a Control. *Journal of the American Statistical Association* **50**, 1096–1121 (1955).

89. Hothorn, T., Bretz, F. & Westfall, P. Simultaneous inference in general parametric models. *Biom J* **50**, 346–363 (2008).

90. Schaarschmidt, F., Ritz, C. & Hothorn, L. A. The Tukey Trend Test: Multiplicity Adjustment Using Multiple Marginal Models. *Biometrics* **78**, 789–797 (2022).

91. Tukey, J. W., Ciminera, J. L. & Heyse, J. F. Testing the statistical certainty of a response to increasing doses of a drug. *Biometrics* **41**, 295–301 (1985).

92. R Core Team. *R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna, Austria. 2013*. (ISBN 3-900051-07-0, 2014).

93. Oklestkova, J. *et al.* Immunoaffinity chromatography combined with tandem mass spectrometry: A new tool for the selective capture and analysis of brassinosteroid plant hormones. *Talanta* **170**, 432–440 (2017).

94. Tarkowská, D., Novák, O., Oklestkova, J. & Strnad, M. The determination of 22 natural brassinosteroids in a minute sample of plant tissue by UHPLC-ESI-MS/MS. *Anal Bioanal Chem* **408**, 6799–6812 (2016).

95. Ewels, P., Magnusson, M., Lundin, S. & Käller, M. MultiQC: summarize analysis results for multiple tools and samples in a single report. *Bioinformatics* **32**, 3047–3048 (2016).

96. Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome alignment and genotyping with HISAT2 and HISAT-genotype. *Nat Biotechnol* **37**, 907–915 (2019).

97. Trapnell, C. *et al.* Transcript assembly and quantification by RNA-Seq reveals unannotated transcripts and isoform switching during cell differentiation. *Nat Biotechnol* **28**, 511–515 (2010).

98. Liao, Y., Smyth, G. K. & Shi, W. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. *Bioinformatics* **30**, 923–930 (2014).

99. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

100. Tang, D. *et al.* SRplot: A free online platform for data visualization and graphing. *PLoS One* **18**, e0294236 (2023).

101. Musielak, T. J., Schenkel, L., Kolb, M., Henschen, A. & Bayer, M. A simple and versatile cell wall staining protocol to study plant reproduction. *Plant Reprod* **28**, 161–169 (2015).

102. Asami, T. *et al.* Characterization of brassinazole, a triazole-type brassinosteroid biosynthesis inhibitor. *Plant Physiol* **123**, 93–100 (2000).

103. Bojar, D. *et al.* Crystal structures of the phosphorylated BRI1 kinase domain and implications for brassinosteroid signal initiation. *Plant J* **78**, 31–43 (2014).

104. Edgar, R. C. MUSCLE: a multiple sequence alignment method with reduced time and space complexity. *BMC Bioinformatics* **5**, 113 (2004).

105. Kabsch, W. & Sander, C. Dictionary of protein secondary structure: pattern recognition of hydrogen-bonded and geometrical features. *Biopolymers* **22**, 2577–2637 (1983).

106. Yuan, T. *et al.* BEN1, a gene encoding a dihydroflavonol 4-reductase (DFR)-like protein, regulates the levels of brassinosteroids in Arabidopsis thaliana. *Plant J* **51**, 220–233 (2007).

107. Schneider, K. *et al.* Arabidopsis PIZZA Has the Capacity to Acylate Brassinosteroids. *PLOS ONE* **7**, e46805 (2012).

108. Choi, S. *et al.* BAT1, a putative acyltransferase, modulates brassinosteroid levels in Arabidopsis. *Plant Journal* **73**, 380–391 (2013).

109. Zhu, W. *et al.* Homeostasis of Brassinosteroids Regulated by DRL1, a Putative Acyltransferase in *Arabidopsis*. *Molecular Plant* **6**, 546–558 (2013).

110. Oktay, C. *et al.* Arabidopsis thaliana accumulates dehydroepiandrosterone after infection with phytopathogenic fungi - Effects on plants and fungi. *Plant Physiol Biochem* **221**, 109570 (2025).

111. Herrera, J. B., Bartel, B., Wilson, W. K. & Matsuda, S. P. Cloning and characterization of the Arabidopsis thaliana lupeol synthase gene. *Phytochemistry* **49**, 1905–1911 (1998).

112. Huang, L. *et al.* Molecular characterization of the pentacyclic triterpenoid biosynthetic pathway in Catharanthus roseus. *Planta* **236**, 1571–1581 (2012).

113. Shiko, G. *et al.* Occurrence and conversion of progestogens and androgens are conserved in land plants. *New Phytol* **240**, 318–337 (2023).

Fig. 1 | The branched architecture of the BR biosynthesis pathway. a, Chemical structure of brassinolide with labeled steroidal rings (letters) and carbon atoms (numbers). **b**, Schematic representation of the BR biosynthesis pathway, highlighting the relative positions of the BRs discussed in this study. The C_{28}, C_{29}, and C_{27} branches are shaded from light to dark gray, respectively. Key chemical differences between branches are indicated in blue, while red marks chemical modifications along each biosynthetic route. Solid lines denote single-step enzymatic reactions; dotted lines indicate multi-step conversions. Known enzymes involved in the canonical C_{28} pathway are shown alongside.

Fig. 2 | BRI1, BRL1 and BRL3 bind chemically diverse BRs. a, GCI analysis of ligand-binding kinetics between BRI1, BRL1, BRL3 and brassinolide. Shown are sensorgrams (n=2, red) with global fits (black) and equilibrium dissociation constants (K_d) indicated. **b,** BR-binding pockets of BRI1 (left; PDB-ID: pdb_00003rj0) and BRL3 (right) shown as ribbon diagrams. LRRs in blue, island domains in red, brassinolide in yellow. Pocket residues and hydrogen bonds (dotted lines) shown; conserved C_{23}-OH hydrogen bond highlighted in purple. **c,** Isoaffinity plots of association (k_a) and dissociation (k_d) rates for receptor - BR pairs. Each point (n = 2) reflects a global fit; diagonal lines denote K_d. High- and low-affinity interactions are in blue and red, respectively. **d,** Structures of tested BRs, numbered as in **c**, with BRI1 K_d values. Red highlights indicate modifications that reduce binding, while blue highlights indicate modifications that have a minimal effect.

Fig. 3 | The hormone binding pocket can accommodate structurally diverse BRs. a, Structural superposition of different BRI1 - BR (top, r.m.s.d. ~0.4 Å comparing 186 corresponding C_α atoms) and BRL3 - BR complexes (bottom, ~r.m.s.d. ~0.3 Å comparing 190 corresponding C_α atoms). The receptors are shown in combined ribbon diagram (LRR domain in light blue, island domain in light red) and molecular surface views. BRs are in bonds representation color-coded according to their receptor binding affinities, ranging from low (light red) to high affinity (purple; BRI1 - brassinolide complex PDB-ID: pdb_00003rj0). Dashed lines highlight key polar interactions among the ligand and the protein receptor binding pockets. A conserved hydrogen bond between brassinolide and the island domains of BRI1, BRL1, and BRL3 is marked in magenta. **b,** Structures of BRs (in bonds representation, color-coded as in **a** with BRI1 (top) or BRL3 (bottom) and including final mF_o–DF_c omit electron density maps (gray mesh, contoured at the indicated σ level). The respective chemical structures are shown above, with modifications relative to brassinolide marked in red (for changes that reduce binding affinity), or blue (neutral). **c,** Root growth inhibition assay of 7-d-old Col-0 wild-type and $bsk3-1$[75] mutant seedlings (n=31-35) grown on $^{1/2}$MS plates supplemented with the indicated BR (brassinolide, BL; 28-norbrassinolide, 28norBL; 28-homobrassinolide, 28hBL; 23-deoxycastasterone, 23dCS, 0.1% [v/v] DMSO as control) at a final concentration of 100 nM (scale bar = 1 cm). Quantifications (n=30) are shown as box plots (centre line, median; box, interquartile range IQR; whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data shown a gray dots). Multiple comparison of the different treatments vs. the DMSO control were performed using a Dunnett[89] test as implemented in the R package multcomp[89] (*$p <$ 0.05; **$p < 0.01$; ***$p < 0.001$). The relationship between the estimated dissociation constant (K_d) for each ligand and root length were tested using a Tukey trend test[90,91]. The blue line represent the linear trend with slope (b) and the associated test p-value (*$p <$ 0.05; **$p < 0.01$; ***$p < 0.001$). **d,** BR quantification in 10 d old Col-0 and $bri1$ (GABI_134E10[47]) seedlings (FW, fresh weight). **e,** RNA-seq derived expression profile heat map of BR biosynthesis, catabolism, transport and receptor genes in 10 d old seedlings in untreated BR signaling mutants and the Col-0 wild type. Expression values were transformed into Z-scores based on the normalized counts, three biological replicates are shown in three horizontal rows, the color scale indicates expression level (red, up-regulation, blue, down-regulation).

Fig. 4 | BRL2 is a medium-affinity BR receptor. a, Structural superposition of the BRI1 ectodomain (ribbon diagram, in gray; PDB-ID: pdb_00003rj0) with the AlphaFold2 model of the apo BRL2 ectodomain (in blue, AlphaFoldDB-ID: AF-Q9ZPS9-F1, r.m.s.d. ~ 0.8 Å comparing 163 C_α atoms). Brassinolide is in bonds representation, interacting residue are shown as ball-and-sticks. **b,** GCI-derived binding kinetics of the BRL2 ectodomain vs. brassinolide, castasterone and typhasterol. Red traces represent experimental sensorgrams (n=2), and black lines show the global fits. **c,** Dissociation constant (K_d) table of the BRL2 ectodomain vs. different BRs, non-BR steroids and triterpenpoids (n.d., no binding detected). **d,** Confocal images of a representative *BRL2prom*::NLS-3xmVenus reporter (in green) line in different stages: heart-stage embryo, mature embryo (cell wall stained with Renaissance SR2200, gray, scale bar = 20 μm), cotyledon (scale bar = 200 μm) and root from a 10-d-old seedling (cell wall stained with propidium iodide, magenta, scale bar = 30 μm). **e,** Rosette phenotypes of 4-week-old plants, germinated on $^{1/2}$MS for 2 weeks before transferring to soil for additional 2 weeks (scale bar = 2 cm).

Fig. 5 | Conserved aromatic residues in the hormone binding pocket are critical for BR perception. a, Ribbon diagram of BRI1 (LRR domain in blue, island domain in red) in complex with brassinolide (in bonds representation, PDB-ID: pdb_00003rj0). Mutated residues are displayed as ball-and-sticks. **b,** GCI-derived binding kinetics for BRI1[W564H], BRI1[G644D], BRI1[M657A] and BRI1[F681A] vs. brassinolide. Red traces represent experimental data (n=2), black lines show the global fits. **c,** Dissociation constant (K_d) table for BRI1 point mutants vs. brassinolide. **d,** Rosette phenotypes of representative, 5-week-old transgenic T3 *BRI1prom*::BRI1-Citrine[mut] complementation lines in *bri1* null[47] background compared to the Col-0 wild-type control (scale bar = 2 cm). **e,** Quantification of BRI1[mut]-mCitrine (theoretical molecular mass ~ 158.6 kDa) protein levels by western blot from 10-d-old seedling protein extracts, immuno-blotted with an anti-GFP antibody. ReadyBlue-stained RuBisCO is shown below as input loading control. **f,** Root length phenotypes of 8-d-old seedlings (n=30) shown as box plots (centre line, median; box, interquartile range IQR; whiskers, lowest/highest data point within 1.5 IQR of the lower/upper quartile, raw data shown a gray dots). Multiple comparison of the different treatments vs. the indicated controls were performed using a Dunnett[88] test as implemented in the R package multcomp[89] (multiple comparison against *bri1*: *$p < 0.05$; **$p < 0.01$; ***$p < 0.001$; against Col-0 # $p < 0.05$; ## $p < 0.01$; ### $p < 0.001$).

Fig. 6

Fig. 6 | BR C$_2$ hydroxylation and C$_6$ oxidation are critical for SERK co-receptor binding. a, Close-up view of the SERK1-bound (in blue) BRI1 (in gray) - brassinolide (in bonds representation, in yellow) complex (PDB-ID: 00004lsx). Receptor and co-receptor amino acids involved in BR-binding are highlighted as ball-and-sticks in gray and blue, respectively. Polar interactions between the brassinolide C$_2$ hydroxyl group and His62, or the C$_6$ oxygen with Tyr642 and Lys601 are shown as dashed lines (in black). **b,** GCI binding kinetics of amine-coupled and brassinolide-bound BRI1, BRL1, BRL2, and BRL3 ectodomains vs. SERK1. Shown are sensorgrams with red traces indicating experimental data and black lines representing global fits (n=1). Binding affinities (K$_d$) are reported alongside. **c,** Representative GCI traces showing the interaction between SERK1 and biotinylated, Avi-tagged BRI1 ectodomain in the presence of either brassinolide, typhasterol, 3-deoxybrassinolide, or 6-deoxocastasterone. **d,** Table summaries of the receptor - BR - co-receptor interaction dissociation constants. **e,** Chemical structures of selected BRs tested in **c,d**. Structural modifications relative to brassinolide that reduce co-receptor binding are highlighted in red. Modifications that have little effect on complex formation are shown in blue. **f,** GCI binding kinetics of amine-coupled wild-type BRI1, BRI1^{Y597F}, BRI1^{Y599A}, BRI1^{K601L}, and BRI1^{Y642A} ectodomains vs. SERK1 in the presence of a high, saturating concentration of brassinolide. Shown are sensorgrams with red traces indicating experimental data and black lines representing global fits (n=2). Binding affinities (K$_d$) are reported alongside. **g,** BR seedling root growth inhibition antagonist assay (n=30-35). Brassinolide (BL) at a fixed concentration of 100 nM was used in the different treatments, vs. 0.02% (v/v) DMSO as control (in yellow). The antagonist candidate compounds secasterone, 2-deoxybrassinolide and 3-deoxycastasterone were tested for their ability to inhibit seedling root growth by themselves (in blue), or to reverse BL-induced root growth inhibition (in orange). Shown are boxplots with centre lines (medians), box limits indicate the 25th and 75th percentiles, whiskers extend to the full data range. The relationship between antagonist concentration and root length were tested using a Tukey trend test[90,91]. The blue line represent the linear trend with slope b (negative values indicate agonistic behavior, positive values indicate antagonistic behavior) and the associated test p-value (* $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$). Representative seedling images are shown above (scale bar = 0.5 cm). The DMSO and BL 100 nM controls are same in each panel, indicated by a white vertical line.

Fig 7 | Strong receptor binding, correct positioning within the BR binding site, and interaction with the co-receptor are essential for BR bioactivity. Schematic overview of brassinosteroid-induced receptor - co-receptor complex formation and activation. A BR ligand binds to the BRI1 receptor ectodomain (in gray), creating a binding surface for a SERK co-receptor kinase (in blue). This brings the kinase domains of receptor and co-receptor in close proximity, leading to trans-phosphorylation and intracellular signaling. The BR-binding pocket is formed by the receptor (in gray) and co-receptor (in blue). Chemical features of the BR ligand (in black) involved in co-receptor binding are highlighted in dark-blue, key carbon atoms critical for receptor or co-receptor interaction are numbered, chemical features required for receptor and/or co-receptor binding are shown in red. The approximate positions of residues critically involved in BR binding, positioning and co-receptor binding are indicated.

a

b

Supplementary Fig. 1 | BRI1, BRL1 and BRL3 share a structurally conserved BR-binding pocket. a, Right: Ribbon diagram of the BRL3 ectodomain bound to brassinolide (yellow, bonds representation). Left: Close-up of a structural superposition of the BR-binding pockets of BRL3 (blue), BRI1 (purple; r.m.s.d. ~ 0.7 Å over 172 C_α atoms), and BRL1 (yellow; r.m.s.d. ~ 0.6 Å over 166 C_α atoms). Key residues involved in brassinolide coordination are shown in ball-and-stick representation. The conserved hydrogen bond between the C_{23} hydroxyl group of brassinolide and the island domain backbone is indicated with a black dotted line. **b,** Multiple sequence alignment of BR-binding pocket regions from *Arabidopsis thaliana* BRI1, BRL1, and BRL3, along with orthologs from rice (OsBRI1, https://www.uniprot.org/ UniProt-ID Q942F3), maize (ZmBRI1, A0A3L6FIW6), and tomato (SlBRI1, F2XYF6), generated using MUSCLE[104]. Secondary structure elements based on AtBRI1 and calculated with DSSP[105] are shown above the alignment. BR-contacting residues (compare Fig. 2b) are highlighted in gray; residues mutated in this study (compare Fig. 4a) are shown in red.

Supplementary Fig 2. | BR catabolic enzyme modify BRs at positions critical for receptor and co-receptor binding. Left panel: Molecular surface view of the BRI1 hormone binding pocket (in gray) occupied by brassinolide (in bonds representation, in yellow; PDB-ID: pdb_00003rj0). The A, B, C, and D rings that make up the hormone's steroid core are labeled and carbon atoms targeted for catabolic modifications are highlighted. Right panel: Chemical structure of brassinolide with carbon numbering. Red, bold numbers indicate sites of BR catabolic reactions: C_{26} hydroxylation by BAS1[17], C_{23} O-glucosylation by UGT73C5[18], C_{22} O-sulfonation by AtST1/ST4a[19], C_6 reduction by BEN1[106] and C_3 O-laurylation by PIZZA/BAT1/DRL1[107–109].

Supplementary Fig. 3 | BRL2 harbors a conserved BR-binding pocket but has not detectable function in BR growth signaling. a, Structural superposition of the BRI1 ectodomain (ribbon diagram, in orange; PDB-ID: pdb_00003rj0) with the apo BRL2 crystal structure (in blue, r.m.s.d. ~ 0.8 Å comparing 314 C_α atoms). **b,** Close-up view of the BRL2 BR-binding pocket (in blue, AlphaFoldDB-ID: AF-Q9ZPS9-F1) superimposed with BRL3 (right panel, in green, brassinolide in bonds representation, r.m.s.d. ~ 0.7 Å comparing 162 C_α atoms). Residues lining the BR binding site are shown in ball-and-stick representation, with corresponding residues in BRL2 shown in blue. Non-synonymous residues among BRI1, BRL2 and BRL3 are labeled. **c,** Schematic diagram of *brl2-2* CRISPR/Cas9 deletion allele. The BRL2 genomic locus is shown, including the translation start codon (black arrow), exon (black rectangle) and 5' and 3- primer untranslated regions (UTRs, gray blocks). The putative binding sites of the designed guide RNAs are indicated in red. **d,** Cotyledon vein patterning analysis performed using 2-week-old Col-0 and *brl2-1* seedlings. Leaf patterns are categorized in seven groups, indicated 4 (4 distinct regions are formed by the intersection of the leaf veins, blue), 3+1 (3 distinct regions formed with 1 free branch, beige), 3 (3 distinct regions formed, lemon green), 2+2 (2 distinct regions formed with 2 free branches, light purple), 2+1 (2 distinct regions formed with 1 free branch, canary), 2 (2 distinct regions formed, gray) and gap (no distinct region formed, earthy yellow). **e,** Hypocotyl growth assay of dark-grown brassinazole (BRZ)[102] treated vs untreated control seedlings. Seeds of Col-0 and *brl2-1* were germinated and grown on ½MS plates supplemented with 1 µM BRZ or 0.1% (v/v) DMSO (scale bar = 1 cm). f, BES1 (theoretical molecular mass ~ 39.2 kDa) western blot. 14-d-old Col-0, *brl2-1* and *bri1* seedlings were grown on ½MS plates with 1% (w/v) sucrose, transferred to ½MS liquid medium containing 1% (w/v) sucrose and either 1 µM brassinolide (BL) or 0.01 % DMSO (as control) for 90 min, and probed with an anti-BES1 antibody[41] by western blotting.

Supplementary Fig. 4 | Mutations of conserved residues in the BRI1 hormone binding pocket alter BR signaling without compromising the plasma-membrane localization of the mutant receptor. a, Structural superposition of the brassinolide-bound AtBRI1 hormone binding pocket (*Arabidopsis thaliana*, in orange, PDB-ID: pdb_00003rj0) with the corresponding regions in ZmBRI1 (*Zea mays*, in blue, AlphaFoldDB-ID: AF-A0A3L6FIW6-F1, r.m.s.d ~ 0.7 Å between 175 C$_\alpha$ atoms), SlBRI1 (*Solanum lycopersicum*, in purple, AlphaFoldDB-ID: AF-F2XYF6-F1, r.m.s.d ~ 0.6 Å between 178 C$_\alpha$ atoms), and OsBRI1 (*Oryza sativa*, in green, AlphaFoldDB-ID: AF-Q942F3-F1, r.m.s.d ~ 0.7 Å between 170 C$_\alpha$ atoms). Mutated residues in AtBRI1 are displayed as ball-and-stick models, the corresponding amino acid side-chains from ZmBRI1, SlBRI1, and OsBRI1 are included. **b,** Analytical size-exclusion chromatography elution profiles for recombinant wild-type (WT) and point-mutant BRI1 ectodomains used in the GCI assays. **c,** BRI1 protein level in Col-0, *bri1* and *BRI1prom*::BRI1-mCitrine / *bri1* lines, detected with a primary anti-BRI1 antibody[103]. ReadyBlue-stained RuBisCO is shown as loading control below. **d,** Confocal imaging of T2 generation *BRI1prom*::BRI1^{mut}-mCitrine signal in the mersistematic zone of 10-d-old primary roots mounted for confocal imaging (scale bar = 20 μm). **e,** Brassinolide (BL)-stimulated BES1 dephosphorylation. 14-d-old seedlings of Col-0 wild-type, *bri1* and *BRI1prom*::BRI1^{mut}-mCit T3 complementation lines were treated with 0.1% (v/v) DMSO or 100 nM BL and 1000 nM BL for 60 min before protein extraction and anti-BES1 antibody[41] immuno-blotting. The upper band represents phosphorylated BES1, the lower band (more) dephosphorylated BES1. ReadyBlue-stained RuBisCO is shown as loading control below. **f,** qRT-PCR analysis of the BR marker genes CPD and DWF4 in Col-0 wild-type, *bri1* and *BRI1prom*::BRI1^{mut}-mCit T3 complementation, with ACT2 used as internal reference gene. Each dot represents a biological replicate (n=3).

Surface Mass (pg/mm^2)

Time (s)

Extended Data Fig. 1

a

b

Receptor	Analyte	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
BRI1	norbrassinolide	3.5E5	3.0E-2	80
BRI1	22-deoxycastasterone	3.1E4	6.5E-3	210
BRI1	secasterone	2.3E5	7.3E-3	30
BRI1	2-deoxybrassinolide	2.3E5	3.0E-3	13
BRI1	typhasterol	1.6E5	1.3E-2	80
BRI1	castasterone	4.2E5	1.2E-2	30
BRI1	cathasterone	1.7E4	1.5E-1	9000
BRI1	brassinolide	3.1E5	6.4E-3	20
BRI1	28-homocastasterone	8.2E4	7.2E-2	870
BRI1	28-homobrassinolide	1.5E5	5.6E-2	370
BRI1	24-epibrassinolide	1.3E5	6.6E-2	515
BRI1	24-epicastasterone	7.8E4	9.9E-2	1200
BRI1	6-deoxocastasterone	1.4E5	2.7E-2	180
BRI1	3-deoxycastasterone	1.2E5	1.2E-2	100
BRI1	23-deoxycastasterone	3.3E4	9.5E-2	2900
BRL1	secasterone	3.0E5	9.4E-3	30
BRL1	2-deoxybrassinolide	5.2E5	8.7E-3	16
BRL1	typhasterol	1.4E5	7.9E-3	55
BRL1	castasterone	2.8E5	9.5E-3	30
BRL1	cathasterone	3.8E4	8.7E-2	2300
BRL1	brassinolide	3.0E5	7.7E-3	25
BRL1	28-homocastasterone	1.7E5	3.0E-2	180
BRL1	28-homobrassinolide	1.4E5	2.6E-2	180
BRL1	24-epibrassinolide	1.4E5	4.7E-2	340
BRL1	24-epicastasterone	1.6E5	4.4E-2	280
BRL3	typhasterol	2.2E5	7.9E-3	30
BRL1	6-deoxocastasterone	1.4E5	1.2E-2	90
BRL3	22-deoxycastasterone	4.2E4	2.1E-2	500
BRL1	3-deoxycastasterone	2.6E5	1.4E-2	50
BRL3	cathasterone	1.9E4	4.8E-2	2500
BRL3	norbrassinolide	2.0E5	1.2E-1	600
BRL3	secasterone	3.9E5	8.3E-3	20
BRL1	23-deoxycastasterone	5.9E4	6.9E-2	1200
BRL3	brassinolide	5.7E5	8.0E-3	15
BRL3	castasterone	4.2E5	8.5E-3	20
BRL3	2-deoxybrassinolide	3.2E5	7.7E-3	20
BRL3	28-homocastasterone	1.9E5	4.8E-2	250
BRL3	28-homobrassinolide	6.2E4	3.2E-2	520
BRL3	24-epibrassinolide	1.4E5	4.8E-2	330
BRL3	24-epicastasterone	9.8E4	5.7E-2	570
BRL3	6-deoxocastasterone	1.7E5	1.2E-2	70
BRL3	3-deoxycastasterone	3.5E5	9.3E-3	26
BRL3	23-deoxycastasterone	2.3E4	7.5E-2	3300

Extended Data Fig. 2

a

b

Analyte	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
brassinolide	1.6E5	3.5E-2	220
2-deoxybrassinolide	9.3E4	3.9E-2	420
castasterone	7.7E4	5.8E-2	750
typhasterol	4.0E4	9.6E-2	2400
28-norbrassinolide	7.6E4	1.5E-1	1900
28-homobrassinolide	4.8E4	1.2E-1	2400
cathasterone	1.7E4	1.3E-1	8000
28-homocastasterone	4.0E3	1.2E-1	30000
24-epicastasterone	n.d.	n.d.	n.d.
DHEA	n.d.	n.d.	n.d.
triolone	n.d.	n.d.	n.d.
24-epibrassinolide	n.d.	n.d.	n.d.
betulin	n.d.	n.d.	n.d.
lupeol	n.d.	n.d.	n.d.

c

dehydroepiandrosteron (DHEA)   triolone   lupeol   betulin

Extended Data Fig. 3

a

b

BRI1 mutant	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
W564H	3.6E4	7.6E-2	2100
Y597M	1.0E6	1.1E-2	11
Y597F	2.0E5	5.8E-3	30
Y599A	2.6E5	2.7E-2	100
K601L	4.2E5	1.3E-2	30
L615Y	3.8E5	2.9E-2	80
L615F	7.0E4	7.3E-3	100
Y642A	2.6E5	5.4E-2	200
G643E	1.8E6	2.7E-3	1.5
M657A	4.3E5	8.2E-3	20
G644D	6.4E4	3.9E-2	610
M657E	9.6E4	2.9E-3	30
T646I	2.2E5	6.5E-3	30
F681A	4.3E6	7.8E-3	3

Extended Data Fig. 4

a

Receptor	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
BRI1	3.1E4	1.4E-2	440
BRL1	4.6E4	2.6E-2	570
BRL2	3.5E2	1.2E-2	36000
BRL3	3.6E4	2.3E-2	630

b SERK1 injections - constant BR concentration

c

surface mass (pg/mm²)

BRI1-6dCS vs. SERK1

BRI1-SEC vs. SERK1

BRI1-2dBL vs. SERK1

BRI1-TYPH vs. SERK1

BRI1-3dCS vs. SERK1

BRI1-CS vs. SERK1

BRI1-BL vs. SERK1

time (s)

d

brassinosteroid	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
brassinolide	3.1E5	2.3E-2	75
castasterone	4.1E5	2.1E-2	50
typhasterol	1.8E5	1.8E-1	970
2-deoxybrassinolide	1.5E5	1.5E-1	1000
3-deoxycastasterone	3.1E5	5.7E-2	180
secasterone	1.5E5	3.3E-1	2100
6-deoxocastasterone	1.2E5	2.0E-1	1700

e SERK1 injections - 1000 nM brassinolide

f

receptor	k_a (M^{-1}s^{-1})	k_d (s^{-1})	K_d (nM)
BRI1^{WT}	7.4E5	2.7E-2	40
BRI1^{Y597F}	4.7E5	4.0E-2	10
BRI1^{Y599A}	2.8E4	1.5E-1	5500
BRI1^{K601L}	3.7E5	6.2E-2	170
BRI1^{Y642A}	4.2E4	7.1E-2	1700

Extended Data Fig. 5

General methods
The NMR spectra were taken on a JEOL JNM-ECA 500 (JEOL, Tokyo, Japan; [1]H, 500 MHz; [13]C, 125 MHz) spectrometer equipped with a 5mm JEOL Royal probe. [1]H NMR and [13]C NMR chemical shifts (δ) were calibrated using tetramethylsilane (TMS, [1]H δ=0 ppm) or solvents: CDCl$_3$ ([1]H δ=7.27 ppm, [13]C δ=77.00 ppm) or DMSO-d$_6$ ([1]H δ=2.46 ppm, [13]C δ=40.00 ppm). Chemical shifts are given in ppm (δ-scale), coupling constants (J) in Hz. All values were obtained by first order analysis. The NMR data were processed using the ACD/NMR Processor Academic Edition (ver. 12.01). For ESI or APCI HRMS analysis, the samples were dissolved in chloroform or methanol. Samples were analyzed with 1290 Infinity II Liquid Chromatographer (Agilent) with HPLC column infinityLab Poroshell 120 EC-C18 (4.6×50 mm; 2.7 µm). For API HRMS analysis, the samples were dissolved in chloroform (or chloroform:methanol; 1:1; v/v, in the case of hydroxylated compounds) to a concentration 10 µg·mL^{-1}. The ASAP (Atmospheric Solids Analysis Probe) was dipped into the sample solution, placed into the ion source and analyzed in full scan mode. The source of the Synapt G2-Si Mass Spectrometer (Waters, Manchester, UK) was operated in positive ionization mode (ASAP+) and, if not stated otherwise, at source temperature of 120 °C. The Corona needle current was kept at 5 µA and the collision energy at value 4. The probe temperature was ramped up from 50 °C to 600 °C in 3 min. Data were acquired from 50 to 1000 Da with 1.0 s scan time in High Resolution Mode and processed using the Masslynx 4.1 software (Waters). Mass accuracy of 1 ppm or less was achieved with the described instrumentation for all compounds.

Merck silica gel Kieselgel 60 (230–400 mesh) was used for column chromatography. The HPLC system consisted of a Waters semi-preparative HPLC system including quaternary pump, liquid handler, UV-VIS and ELSD detectors. The HPLC semi preparative column was filled with silica gel. Reagents and solvents were purchased from Sigma–Aldrich and were not purified.

Scheme 1: Synthesis of 3-deoxycastasterone (**7**), typhasterol (**8**), and 2-deoxybrassinolide (**11**). **a**: (*S*)-2-(3-methylbutan-2-yl)-1,3-dithiane/n-BuLi/THF, r.t. then -78 °C, 1h; **b**: NaH$_2$PO$_4$/2-methyl-2-butene/ NaClO$_2$/H$_2$O/ethanol, 0 °C, 10 min; **c**: TBSOTf/2,6-lutidine/CH$_2$Cl$_2$, 0 °C, 1

h; **d**: DIBAL-H/toluen, -78 °C, 1h; **e**: TBAF/THF, r.t., 1h; **f**: 1, BH_3/THF, r.t., 6h; 2, $NaOH/H_2O_2/H_2O$/THF, r.t., 12h; **g**: MCPBA/EtOAc, r.t., 4h; **h**: $LiAlH_4$/THF/Et_2O, 40 °C, 3h; **i**: HCl/H_2O/THF, 40 °C, 4h; **j**: TFPAA/CH_2Cl_2, r.t., 4h; **k**: pyridine/Ac_2O, 50 °C, 12h; **l**: MeONa/MeOH/THF, r.t., 1h. Note: absolute configuration for carbons 8, 9, 13, and 17 is shown only for compound **1** but is identical for all compounds prepared.

Synthesis of (22R)-6,6-ethylenedioxy-23,23-(trimethylenedithio)-5α-campest-2-en-22-ol (2)

1.6 M solution of *n*-BuLi in hexane (3.0 mL, 4.8 mmol) was added to a solution of (*S*)-2-(3-methylbutan-2-yl)-1,3-dithiane[1] (790 mg, 4.0 mmol) in THF (15 mL) at room temperature. The mixture was stirred for 30 minutes and then cooled to -78 °C. A solution of aldehyde **1**[2] (1.0 g, 2.69 mmol) in THF (3 mL) was added dropwise to the reaction mixture. The solution was stirred at -78 °C for 30 min, quenched with saturated aqueous NH_4Cl (20 mL), and warmed to room temperature. The product was extracted with ethyl acetate. The organic layer was washed with water followed by drying over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (3-5% ethyl acetate in cyclohexane) to give alcohol **2** (1.22 g, 81 %) as a viscous oil. [1]H NMR (500 MHz, DMSO-d_6): δ 5.63 (m, 1H, H-3), 5.51 (m, 1H, H-2), 4.50 (d, 1H, *J* = 6.4 Hz, H-22), 3.92-3.79 (m, 3H, H_{ketal}), 3.69 (q, 1H, *J* = 6.6 Hz, H_{ketal}), 2.83-2.60 (m, 5H), 2.12-1.69 (m, 11H), 1.62-1.23 (m, 8H), 1.15-0.75 (m, 4H), 1.11 (d, 3H, *J* = 6.4 Hz, CH_3), 0.99 (d, 3H, *J* = 7.0 Hz, CH_3), 0.89 (d, 3H, *J* = 6.7 Hz, CH_3), 0.86 (d, 3H, *J* = 6.7 Hz, CH_3), 0.80 (s, 3H, CH_3), 0.66 (s, 3H, CH_3). [13]C NMR (125 MHz, DMSO-d_6): 125.45, 124.86, 108.99, 73.86, 65.27, 63.78, 63.56, 55.49, 53.79, 52.49, 47.57, 43.26, 42.13, 40.80, 40.62, 39.46, 38.13, 35.37, 32.99, 27.73, 27.25, 26.33, 25.51, 24.75, 24.39, 23.73, 21.07, 20.49, 18.44, 14.58, 13.53, 11.59, 9.87. HRMS (ESI+) calculated for $C_{33}H_{55}O_3S_2$ ([M + H]$^+$) 563.3593, found 563.3597.

Synthesis of (22R)-6,6-ethylenedioxy-22-hydroxy-5α-campest-2-en-23-one (3)

A solution of NaH_2PO_4 (511 mg, 4.26 mmol) in water (11 mL), 2-methyl-2-butene (2.25 mL, 21.3 mmol) and $NaClO_2$ (80%, 1.44 g, 16.0 mmol) were sequentially added at 0 °C to a stirred solution of compound **2** (1.2 g, 2.13 mmol) in ethanol (100 mL). The reaction mixture was stirred for 10 min and quenched with aqueous saturated $Na_2S_2O_3$ (mL). The mixture was diluted with ethyl acetate and extracted with water. The organic layer was dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (10% ethyl acetate in cyclohexane) to give hydroxyketone 3 (736 mg, 73%) as a viscous oil. [1]H NMR (500 MHz, CDCl$_3$): δ 5.66 (m, 1H, H-3), 5.54 (m, 1H, H-2), 4.33 (dd, 1H, *J* = 5.5, 1.5 Hz, H-22), 4.01-3.90 (m, 3H, H_{ketal}), 3.80 (q, 1H, *J* = 6.6 Hz, H_{ketal}), 3.31 (d, 1H, *J* = 5.5 Hz, OH), 2.45 (p, 1H, *J* = 7.3 Hz, H-24), 2.13-1.60 (m, 12H), 1.57-0.77 (m, 9H), 1.08 (d, 3H, *J* = 7.3 Hz, CH_3), 0.94 (d, 3H, *J* = 6.7 Hz, CH_3), 0.892 (s, 3H, CH_3), 0.887 (d, 3H, *J* = 6.7 Hz, CH_3), 0.75 (s, 3H, CH_3), 0.74 (d, 3H, *J* = 6.7 Hz, CH_3). [13]C NMR (125 MHz, CDCl$_3$): 216.23, 125.64, 124.78, 110.00, 77.31, 65.61, 64.09, 55.86, 53.27, 52.52, 48.11, 48.02, 42.35, 41.15, 41.17, 39.47, 38.18, 35.85, 33.43, 28.70, 28.25, 24.00, 21.50, 21.41, 20.80, 19.03, 15.18,

13.59, 12.52, 11.91. HRMS (ESI+) calculated for $C_{30}H_{49}O_4$ ([M + H]$^+$) 473.3631, found 473.3631.

Synthesis of (22*R*)-6,6-ethylenedioxy-22-((*tert*-butyldimethylsilyl)oxy)-5α-campest-2-en-23-one (4)

TBSOTf (670 µl, 2.92 mmol) was added at 0 °C to a stirred solution of hydroxyketone **3** (700 mg, 1.48 mmol) and 2,6-lutidine (850 µL, 7.3 mmol) in CH_2Cl_2 (10 mL). The reaction mixture was stirred for 1 hour at 0 °C and then quenched with saturated aqueous NaHCO$_3$ (10 mL). The mixture was diluted with ethyl acetate and extracted with water. The organic layer was dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (5% ethyl acetate in cyclohexane) to give silyl ether **4** (811 mg, 94%) as a viscous oil. ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.67 (m, 1H, H-3), 5.55 (m, 1H, H-2), 4.31 (d, 1H, *J* = 0.9 Hz, H-22), 4.02-3.90 (m, 3H, H$_{ketal}$), 3.80 (q, 1H, *J* = 6.6 Hz, H$_{ketal}$), 2.38 (dq, 1H, *J* = 8.6, 7.0 Hz, H-24), 2.14-1.60 (m, 11H), 1.57-0.75 (m, 10H), 1.02 (d, 3H, *J* = 7.0 Hz, CH$_3$), 0.93 (s, 9H, 3×CH$_3$), 0.92 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.89 (s, 3H, CH$_3$), 0.83 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.81 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.73 (s, 3H, CH$_3$), 0.06 (s, 3H, CH$_3$), -0.01 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 214.65, 125.65, 124.82, 110.03, 79.17, 65.62, 64.10, 56.17, 53.26, 52.37, 48.49, 48.08, 42.37, 41.20, 41.15, 39.62, 38.53, 35.88, 33.41, 29.11, 28.56, 26.89, 25.88 (3×C), 24.13, 21.56, 21.45, 20.79, 19.39, 15.94, 13.59, 12.87, 11.74, -4.22, -5.00. HRMS (ESI+) calculated for $C_{36}H_{63}O_4Si$ ([M + H]$^+$) 587.4496, found 587.4491.

Synthesis of (22*R*,23*R*)-6,6-ethylenedioxy-22-((*tert*-butyldimethylsilyl)oxy)-5α-campest-2-en-23-ol (5)

1M Solution of DIBAL-H in toluene (6.4 mL, 6.4 mmol) was added to a stirred solution of compound **4** (750 mg, 1.28 mmol) in dry toluene (30 mL) at -78 °C. The reaction mixture was stirred at the same temperature for 1 hour and then quenched with MeOH (6 mL). Saturated aqueous solution of sodium potassium tartrate (30 mL) was added and the mixture was diluted with ethyl acetate. The aqueous layer was separated and extracted twice with ethyl acetate. The combined organic layers were washed with water, dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (2-5% ethyl acetate in cyclohexane) to give alcohol **5** (715 mg, 95%) as a viscous oil (containing 12% of inseparable 23*S*-isomer). ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.56 (m, 1H, H-3), 5.53 (m, 1H, H-2), 3.99-3.88 (m, 3H, H$_{ketal}$), 3.78 (q, 1H, *J* = 6.6 Hz, H$_{ketal}$), 3.58 (d, 1H, *J* = 7.5 Hz, H-23), 3.49 (dd, 1H, *J* = 7.5, 2.9 Hz, H-22), 2.11-1.85 (m, 6H), 1.78-1.37 (m, 9H), 1.24-0.75 (m, 8H), 0.94 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.92 (s, 9H, 3×CH$_3$), 0.90 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.89 (d, 3H, *J* = 6.7 Hz, CH$_3$), 0.87 (s, 3H, CH$_3$), 0.83 (d, 3H, *J* = 7.0 Hz, CH$_3$), 0.68 (s, 3H, CH$_3$), 0.12 (s, 3H, CH$_3$), 0.10 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 125.67, 124.80, 110.02, 76.09, 73.02, 65.59, 64.09, 56.18, 53.33, 52.21, 48.07, 42.54, 41.21, 41.15, 40.91, 39.75 (2×C), 35.88, 33.34, 30.79, 28.55, 26.14 (3×C), 24.21, 21.44, 21.04, 20.81, 19.88, 18.60, 13.60, 12.91, 11.61, 10.00, -3.57, -4.09. HRMS (APCI+) calculated for $C_{36}H_{65}O_4Si$ ([M + H]$^+$) 589.4652, found 589.4651.

Synthesis of (22R,23R)-6,6-ethylenedioxy-5α-campest-2-ene-22,23-diol (6)

1M Solution of TBAF (6.5 mL, 6.5 mmol) was added to a solution of compound **5** (700 mg, 1.19 mmol) in THF (30 mL). The reaction mixture was stirred at room temperature for 1 hour. Then, it was diluted with ethyl acetate and washed with saturated solution of NH_4Cl (20 mL). The combined organic layers were washed with water, dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (25% ethyl acetate in cyclohexane) to give diol **6** (524 mg, 93%) as white solid. ^1H NMR (500 MHz, $CDCl_3$): δ 5.67 (m, 1H, H-3), 5.55 (m, 1H, H-2), 4.01-3.90 (m, 3H, H_{ketal}), 3.79 (q, 1H, $J = 6.6$ Hz, H_{ketal}), 3.73 (dd, 1H, $J = 8.4$, 2.0 Hz, H-23), 3.57 (dd, 1H, $J = 8.4$, 1.1 Hz, H-22), 2.12-1.91 (m, 5H), 1.81-1.37 (m, 10H), 1.31-0.76 (m, 7H), 0.98 (d, 3H, $J = 6.7$ Hz, CH_3), 0.96 (d, 3H, $J = 6.7$ Hz, CH_3), 0.91 (d, 3H, $J = 6.7$ Hz, CH_3), 0.86 (s, 3H, CH_3), 0.86 (d, 3H, $J = 7.0$ Hz, CH_3), 0.71 (s, 3H, CH_3). ^{13}C NMR (125 MHz, $CDCl_3$): 125.65, 124.78, 110.03, 74.87, 73.42, 65.59, 64.08, 55.87, 53.36, 52.40, 48.03, 42.36, 41.18 (2×C), 39.95, 39.79, 36.79, 35.86, 33.40, 30.75, 27.71, 23.99, 21.40, 20.86 (2×C), 20.77, 13.59, 11.88 (2×C), 10.12. HRMS (ESI+) calculated for $C_{30}H_{51}O_4$ ($[M + H]^+$) 475.3787, found 475.3787.

Synthesis of (22R,23R)-2α,22,23-trihydroxy-5α-campestan-6-one (3-deoxycastasterone, 7) and (22R,23R)-3α,22,23-trihydroxy-5α-campestan-6-one (typhasterol, 8)

1M Solution of borane in THF (1 mL, 1.0 mmol) was added dropwise to the solution of olefin **6** (50 mg, 0.1 mmol) in dry THF (2 mL). The reaction mixture was stirred at room temperature for 6 h. Then, the reaction mixture was quenched by water (2 mL) and cooled to 10 °C. Aqueous solution of NaOH (10%, 1 mL) was added followed by dropwise addition of 30% hydrogen peroxide (2 mL). This reaction mixture was stirred at room temperature overnight. It was then poured into a mixture of sodium sulfite (500 mg) in water (5 mL) and acetic acid (0.5 mL) and stirred for additional 30 min. After then, the reaction mixture was diluted with ethyl acetate and extracted twice with water. The organic layer was dried over $MgSO_4$ and evaporated under reduced pressure. A crude mixture of products was dissolved in a mixture of THF and acetone (5:1, 3 mL) and 5% aqueous HCl (1 mL) was added. The reaction mixture was heated under reflux for 3 h. Then it was diluted with ethyl acetate and extracted twice with water. The organic layer was again dried over $MgSO_4$ and evaporated under reduced pressure. The residue was purified by HPLC (75% ethyl acetate in cyclohexane) to give two alcohols, **7** (19 mg, 40%) and **8** (15 mg, 32%), as white solids.

Compound **7** (3-deoxycastasterone): ^1H NMR (500 MHz, $CDCl_3$): δ 3.78 (m, 1H, H-2β), 3.73 (dd, 1H, $J = 8.4$, 1.8 Hz, H-23), 3.56 (dd, 1H, $J = 8.4$, 1.2 Hz, H-22), 2.30 (dd, $J = 13.0$, 4.4 Hz, H-7a), 2.17 (dd, $J = 12.1$, 2.9 Hz, H-5α), 2.09-1.94 (m, 3H), 1.85-1.46 (m, 11H), 1.38-1.09 (m, 11H), 0.97 (d, 3H, $J = 6.7$ Hz, CH_3), 0.95 (d, 3H, $J = 6.7$ Hz, CH_3), 0.91 (d, 3H, $J = 6.7$ Hz, CH_3), 0.85 (d, 3H, $J = 7.0$ Hz, CH_3), 0.76 (s, 3H, CH_3), 0.69 (s, 3H, CH_3). ^{13}C NMR (125 MHz, $CDCl_3$): 211.91, 74.68, 73.50, 67.06, 57.91, 56.58, 54.09, 52.25, 47.07, 46.63, 43.06, 42.75,

40.00, 39.43, 37.67, 36.76, 34.60, 30.73, 27.61, 23.79, 21.28, 20.86, 20.73, 19.48, 14.17, 11.91, 11.88, 10.09. HRMS (ESI+) calculated for $C_{28}H_{49}O_4$ ([M + H]$^+$) 449.3631, found 449.3627.

Compound **8** (typhasterol): ^{1}H NMR (500 MHz, CDCl$_3$): δ 4.18 (m, 1H, H-3β), 3.73 (bd, 1H, J = 8.3 Hz, H-23), 3.57 (bd, 1H, J = 8.3, 1.2 Hz, H-22), 2.74 (t, 1H, J = 7.9 Hz, H-5α), 2.31 (dd, J = 13.1, 4.6 Hz, H-7a), 2.04-1.94 (m, 3H), 1.84-1.47 (m, 15H), 1.41-1.19 (m, 6H), 1.10 (m, 1H), 0.97 (d, 3H, J = 6.7 Hz, CH$_3$), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.92 (d, 3H, J = 6.7 Hz, CH$_3$), 0.85 (d, 3H, J = 6.9 Hz, CH$_3$), 0.74 (s, 3H, CH$_3$), 0.69 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 212.74, 74.72, 73.50, 65.42, 56.68, 53.74, 52.28, 51.65, 46.80, 42.77, 41.55, 40.01, 39.51, 38.02, 36.78, 31.64, 30.74, 28.14, 27.66, 27.64, 23.79, 21.08, 20.87, 20.75, 12.32, 11.92, 11.88, 10.01. HRMS (APCI+) calculated for $C_{28}H_{49}O_4$ ([M + H]$^+$) 449.3631, found 449.3626.

Synthesis of (22*R*,23*R*)-2α,3α-epoxy-6,6-ethylenedioxy-5α-campestane-22,23-diol (9)

3-Chloroperoxybenzoic acid (77%, 375 mg, 1.68 mmol) was added to a solution of olefin **6** (200 mg, 0.42 mmol) in ethyl acetate (10 mL) and the reaction mixture was stirred at room temperature for 4 hours. The reaction was quenched with saturated aqueous solution of Na$_2$SO$_3$ (5 mL) and NaHCO$_3$ (5 mL). The organic layer was separated, washed with water, dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (25% ethyl acetate in cyclohexane) to give epoxide **9** (186 mg, 90%) as an off-white solid. ^{1}H NMR (500 MHz, CDCl$_3$): δ 3.99-3.89 (m, 3H, H$_{ketal}$), 3.79 (m, 1H, H$_{ketal}$), 3.72 (dd, 1H, J = 8.6, 1.5 Hz, H-23), 3.56 (d, 1H, J = 8.4 Hz, H-22), 3.25 (m, 1H), 3.10 (dd, 1H, J = 5.8, 4.0 Hz), 2.14 (ddd, 1H, J = 14.8, 4.1, 1.2 Hz), 2.09-1.96 (m, 2H), 1.97 (dt, 1H, J = 12.7, 3.3 Hz), 1.92 (dd, 1H, J = 14.8, 6.1 Hz), 1.81 (m, 1H), 1.75 (dd, 1H, J = 13.1, 3.4 Hz), 1.68-0.67 (m, 17H), 0.97 (d, 3H, J = 6.7 Hz, CH$_3$), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.92 (d, 3H, J = 6.7 Hz, CH$_3$), 0.97 (d, 3H, J = 6.7 Hz, CH$_3$), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.90 (d, 3H, J = 6.7 Hz, CH$_3$), 0.89 (s, 3H, CH$_3$), 0.85 (d, 3H, J = 6.9 Hz, CH$_3$), 0.69 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 109.78, 74.83, 73.39, 65.47, 64.08, 55.71, 53.06, 52.83, 52.35, 50.46, 43.22, 42.25, 40.91, 39.97, 39.68, 39.59, 36.80, 34.73, 33.10, 30.74, 27.69, 23.99, 20.87, 20.81, 20.77, 20.07, 14.73, 11.89, 11.84, 10.13. HRMS (ESI+) calculated for $C_{30}H_{51}O_5$ ([M + H]$^+$) 491.3736, found 491.3739.

Synthesis of (22*R*,23*R*)-6,6-ethylenedioxy-5α-campestane-3α,22,23-triol (10)

1M Solution of LiAlH$_4$ in diethylether (1 mL, 1.0 mmol) was added dropwise to a solution of compound **9** (130 mg, 0.27 mmol) in dry THF (6 mL). The reaction mixture was stirred at 40 °C for 3 hours and then quenched with MeOH (3 mL). Saturated aqueous solution of sodium potassium tartrate (10 mL) was added and the mixture was diluted with ethyl acetate. The aqueous layer was separated and extracted with ethyl acetate (2×). The combined organic layers were washed with water, dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (90% ethyl acetate in cyclohexane) to give triol **10** (114 mg, 87%) as a viscous oil. ^{1}H NMR (500 MHz, CDCl$_3$): δ 4.13 (m, 1H, H-3β), 4.02-3.92 (m, 3H, H$_{ketal}$), 3.75-3.71 (m, 2H, H-23, H$_{ketal}$), 3.55 (d, 1H, J = 8.5 Hz, H-22), 2.08-1.93 (m, 3H), 1.79 (dd, 1H, J = 13.1, 3.7 Hz), 1.70-1.05 (m, 22H), 0.97 (d, 3H, J = 6.7 Hz, CH$_3$), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.93 (s, 3H, CH$_3$), 0.90 (d, 3H, J = 6.7 Hz, CH$_3$), 0.85 (d, 3H, J = 6.9 Hz, CH$_3$), 0.71 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 110.59,

75.44, 72.53, 65.68, 65.44, 64.05, 56.22, 53.85, 52.66, 45.05, 42.56, 40.64, 40.09, 39.60, 37.66, 37.17, 33.82, 33.35, 30.87, 27.86, 27.51, 27.18, 24.11, 20.92, 20.87, 20.66, 13.04, 11.93, 11.87, 10.28. HRMS (ESI+) calculated for $C_{30}H_{53}O_5$ ($[M + H]^+$) 493.3893, found 493.3895.

Synthesis of typhasterol (8) from compound 10

Aqueous solution of HCl (5%, 0.5 mL) was added to a solution of compound **10** (80 mg, 0.16 mmol) in THF (5 mL) and the reaction mixture was stirred at 40 °C for 4 hours and then diluted with ethyl acetate. Organic layer was washed twice with water, dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (90% ethyl acetate in cyclohexane) to give typhasterol (**8**) (75 mg, 90%) as a white solid. The spectroscopic data were entirely consistent with those reported for typhasterol above.

Synthesis of (22R,23R)-3α,22,23-trihydroxy-7a-homo-7-oxa-5α-campestan-6-one (2-deoxybrassinolide, 11)

Freshly prepared trifluoroperoxyacetic acid (4 mL) was added to a solution of typhasterol (**8**) (60 mg, 0.13 mmol) in dichloromethane (10 mL) and the reaction mixture was stirred at room temperature for 4 hours and quenched with saturated aqueous solutions of Na_2SO_3 (5 mL) and $NaHCO_3$ (5 mL). The aqueous layer was separated and extracted twice with dichloromethane. The combined organic layers were washed with water, dried over $MgSO_4$, filtrated and evaporated under reduced pressure to give inseparable crude mixture of isomeric lactones (65 mg) with compound **11** as the major isomer (ratio approx. 3:1). For separation the mixture had to be acetylated. Crude lactones were dissolved in pyridine (6 mL) and acetic anhydride (1 mL) was added. The reaction mixture was stirred at 50 °C for 12 hours. Then, ethanol (5 mL) and ethyl acetate (10 mL) were added and the mixture was stirred for additional 30 min. The mixture was washed twice with water, dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by HPLC (30% ethyl acetate in cyclohexane) to give two triacetates, **12** (50 mg, 63%) and **12a** (15 mg, 19%), as a viscous oil.

Compound **12** (40 mg, mmol) was dissolved in THF (5 mL) and 5M methanolic solution of sodium methoxide (0.1 mL) was added. The reaction mixture was stirred at room temperature for 1 hour and quenched with 5% aqueous solution of HCl (1 mL). The mixture was diluted with ethyl acetate, washed twice with water, dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by HPLC (ethyl acetate) to give lactone **11** (28 mg, 89%) as a white solid.

Compound **11** (2-deoxybrassinolide): ^1H NMR (500 MHz, $CDCl_3$): δ 4.17 (m, 1H, H-3β), 4.11-4.08 (m, 2H, H-7aα, H-7aβ), 3.73 (dd, 1H, $J = 8.4$, 1.2 Hz, H-23), 3.55 (d, 1H, $J = 8.4$ Hz, H-22), 3.19 (dd, 1H, $J = 12.2$, 4.6 Hz, H-5α), 2.14 (ddd, 1H, $J = 14.9$, 12.5, 2.3 Hz), 2.02-1.96 (m, 3H), 1.81-1.19 (m, 19H), 0.90 (m, 1H), 0.97 (d, 3H, $J = 6.7$ Hz, CH_3), 0.95 (d, 3H, $J = 6.7$ Hz, CH_3), 0.91 (d, 3H, $J = 6.7$ Hz, CH_3), 0.90 (s, 3H, CH_3), 0.85 (d, 3H, $J = 6.9$ Hz, CH_3), 0.72 (s, 3H, CH_3). ^{13}C NMR (125 MHz, $CDCl_3$): 176.78, 74.59, 73.53, 70.39, 64.88, 58.29, 52.25, 51.37, 42.44, 41.75, 40.02, 39.50, 36.79, 36.28, 32.45, 30.73, 29.69, 28.23, 27.57, 24.71, 22.15,

20.85, 20.74, 14.55, 14.12, 11.82, 11.72, 10.10. HRMS (APCI+) calculated for $C_{28}H_{49}O_5$ ([M + H]$^+$) 465.3580, found 465.3574.

Compound **12** ((22R,23R)-7a-homo-7-oxa-6-oxo-5α-campestan-3α,22,23-triyl triacetate): ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.33 (dd, 1H, J = 8.9, 1.2 Hz, H-23), 5.15 (d, 1H, J = 8.9 Hz, H-22), 5.09 (m, 1H, H-3β), 4.12-4.03 (m, 2H, H-7aα, H-7aβ), 3.02 (dd, 1H, J = 12.2, 4.3 Hz, H-5α), 2.17 (ddd, 1H, J = 15.0, 12.5, 2.5 Hz), 2.10-1.96 (m, 3H), 2.07, 2.01, 2.00 (all s, 1H, CH$_3$), 1.85-1.13 (m, 17H), 1.01 (d, 3H, J = 6.7 Hz, CH$_3$), 0.96 (d, 3H, J = 6.7 Hz, CH$_3$), 0.94 (d, 3H, J = 6.7 Hz, CH$_3$), 0.91 (d, 3H, J = 6.7 Hz, CH$_3$), 0.90 (s, 3H, CH$_3$), 0.73 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 176.04, 170.56, 170.51, 170.30, 75.67, 74.05, 70.29, 68.41, 58.37, 52.29, 51.42, 42.55, 42.46, 39.78, 39.56, 39.43, 36.97, 36.11, 33.67, 30.34, 29.73, 27.98, 25.14, 24.69, 22.12, 21.37, 20.93, 20.86 (2×C), 20.31, 14.61, 12.71, 11.59, 11.05. HRMS (APCI+) calculated for $C_{34}H_{55}O_8$ ([M + H]$^+$) 591.3897, found 591.3896.

Compound **12a** ((22R,23R)-7a-homo-6-oxa-7-oxo-5α-campestan-3α,22,23-triyl triacetate): ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.33 (dd, 1H, J = 8.9, 1.5 Hz, H-23), 5.15 (d, 1H, J = 8.9 Hz, H-22), 5.14 (m, 1H, H-3β), 4.46 (dd, 1H, J = 11.2, 5.5 Hz, H-5α), 2.53 (d, 1H, J = 13.3 Hz, H-7aβ), 2.45 (dd, 1H, J = 13.3, 11.3 Hz, H-7aα), 2.12-1.95 (m, 3H), 2.07, 2.01, 2.00 (all s, 1H, CH$_3$), 1.90-1.09 (m, 17H), 1.01 (d, 3H, J = 6.7 Hz, CH$_3$), 0.96 (d, 3H, J = 6.7 Hz, CH$_3$), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.914 (s, 3H, CH$_3$), 0.908 (d, 3H, J = 6.7 Hz, CH$_3$), 0.73 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 174.74, 170.53 (2×C), 170.16, 79.67, 75.61, 74.03, 69.56, 58.02, 55.45, 52.62, 42.52, 39.79, 39.64, 39.58, 38.03, 36.97, 34.80, 32.93, 31.84, 30.36, 27.65, 25.14, 24.83, 22.18, 21.31, 20.94, 20.86 (2×C), 20.31, 12.70, 11.64, 11.59, 11.06. HRMS (APCI+) calculated for $C_{34}H_{55}O_8$ ([M + H]$^+$) 591.3897, found 591.3888.

Scheme 2: Synthesis of 23-deoxycastasterone (**15**), 22-deoxycastaterone (**20**). **a**: Ra-Ni/EtOH, r.t., 6h; **b**: HCl/H$_2$O/THF, 40 °C, 4h; **c**: OsO$_4$/K$_2$CO$_3$/K$_2$[Fe(CN)$_6$]/CH$_3$SO$_2$NH$_2$/HQ-4-ClBz/t-BuOH/H$_2$O, r.t., 10 h; **d**: p-TsCl/pyridine, r.t., 3 days; **e**: LiAlH$_4$/THF/Et$_2$O, r.t., 3h; **f**: HCl/H$_2$O/THF, 40 °C, 2h; **g**: OsO$_4$/K$_2$CO$_3$/K$_2$[Fe(CN)$_6$]/CH$_3$SO$_2$NH$_2$/HQ-4-ClBz/t-BuOH/H$_2$O, r.t., 10 h.

Synthesis of (22*S*)-6,6-ethylenedioxy-5α-campest-2-en-22-ol (13)

Raney-Nickel catalyst 2400 (Merck, 1.4g) was added to a solution of compound **2** (200 mg, mmol) in ethanol (20 mL). The resulting suspension was shaken vigorously for 6 hours and then the liquid was decanted. Catalyst was washed with ethanol (2× 10 mL) and the combined ethanolic fractions were evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (5-10% ethyl acetate in cyclohexane) to give alcohol **13** (85 mg, 52%) as a viscous oil. ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.67 (m, 1H, H-3), 5.55 (m, 1H, H-2), 4.01-3.89 (m, 3H, H$_{ketal}$), 3.81-3.75 (m, 2H, H-22, H$_{ketal}$), 2.11-1.90 (m, 4H), 1.81-1.70 (m, 3H), 1.63-0.76 (m, 18H), 0.895 (d, 3H, J = 6.7 Hz, CH$_3$), 0.889 (s, 3H, CH$_3$), 0.880 (d, 3H, J = 6.7 Hz, CH$_3$), 0.84 (d, 3H, J = 6.7 Hz, CH$_3$), 0.82 (d, 3H, J = 6.7 Hz, CH$_3$) 0.71 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 125.67, 124.80, 110.05, 71.66, 65.59, 64.08, 55.89, 53.41, 52.58, 48.05, 42.46, 41.21 (2×C), 39.79, 39.35, 39.29, 35.87, 35.29, 33.40, 32.00, 27.71, 24.08, 21.42, 20.87, 19.98, 17.80, 15.76, 13.60, 11.92, 11.22. HRMS (ESI+) calculated for C$_{30}$H$_{51}$O$_3$ ([M + H]$^+$) 459.3838, found 459.3834.

Synthesis of (22*S*)-22-hydroxy-5α-campest-2-en-6-one (14)

Aqueous solution of HCl (5%, 0.5 mL) was added to a solution of compound **13** (70 mg, 0.15 mmol) in THF (5 mL) and the reaction mixture was stirred at 40 °C for 4 hours and then diluted with ethyl acetate. Organic layer was washed twice with water, dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (10% ethyl acetate in cyclohexane) to give compound **14** (55 mg, 87%) as a white solid. ^{1}H NMR (500 MHz, CDCl$_3$): δ 5.68 (m, 1H, H-3), 5.56 (m, 1H, H-2), 3.67 (td, J = 6.8, 1.1 Hz, H-22), 2.37-2.34 (m, 2H), 2.25 (m, 1H), 2.06-1.92 (m, 4H), 1.74 (m, 1H), 1.62-0.96 (m, 17H), 0.89 (d, 3H, J = 6.7 Hz, CH$_3$), 0.87 (d, 3H, J = 6.7 Hz, CH$_3$), 0.82 (d, 3H, J = 6.7 Hz, CH$_3$), 0.80 (d, 3H, J = 7.0 Hz, CH$_3$), 0.71 (s, 3H, CH$_3$), 0.68 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 212.06, 124.87, 124.46, 71.48, 56.59, 53.74, 53.27, 52.39, 46.92, 42.66, 39.99, 39.41, 39.31, 39.26, 39.22, 37.70, 35.22, 31.96, 27.57, 23.83, 21.64, 21.07, 19.92, 17.77, 15.71, 13.45, 11.82, 11.16. HRMS (ESI+) calculated for C$_{28}$H$_{47}$O$_2$ ([M + H]$^+$) 415.3576, found 415.3578.

Synthesis of (22*S*)-2α,3α,22-trihydroxy-5α-campestan-6-one (23-deoxycastasterone, 15)

Solution of osmium tetroxide in *t*-butanol (0.2M, 0.15 mL, 0.03 mmol) was added to a solution of compound **14** (40 mg, 0.1 mmol), hydroquinidine 4-chlorobenzoate (25 mg; 0.05 mmol), methanesulfonamide (5 mg; 0.05 mmol), potassium carbonate (100 mg; 0.72 mmol), and potassium ferricyanide (240 mg; 0.73 mmol) in the mixture of *t*-butanol and water (5 mL; 1 : 1 v/v). The reaction mixture was stirred at room temperature for 10 h and quenched with saturated solution of Na$_2$SO$_3$ (3 mL) was then added. After an additional 30 minutes of stirring, the reaction mixture was diluted with ethyl acetate (30 mL) and extracted with water (2 × 20 mL). Organic layer was dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (0.5% isopropanol in ethyl

acetate) to give triol **15** (31 mg, 72%) as a white solid. ^{1}H NMR (500 MHz, CDCl$_3$ + 5% CD$_3$OD): 3.95 (m, 1H, H-3β), 3.73 (m, 1H, H-2β), 3.71 (bt, 1H, J = 6.4 Hz, H-22), 2.65 (dd, 1H, J = 12.4, 2.3 Hz, H-5α), 2.25 (dd, 1H, J = 13.0, 4.6 Hz, H-7β), 2.01-1.21 (m, 21H), 1.06 (m, 1H), 0.85 (d, 3H, J = 6.7 Hz, CH$_3$), 0.83 (d, 3H, J = 6.7 Hz, CH$_3$), 0.78 (d, 3H, J = 7.0 Hz, CH$_3$), 0.77 (d, 3H, J = 7.0 Hz, CH$_3$), 0.71 (s, 3H, CH$_3$), 0.64 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$ + 5% CD$_3$OD): 213.04, 71.30, 67.97, 67.88, 56.48, 53.57, 52.27, 50.71, 46.62, 42.75, 42.54, 39.84, 39.31, 39.13, 39.08, 37.70, 35.14, 31.90, 29.57, 27.50, 23.77, 21.09, 19.84, 17.68, 15.59, 13.42, 11.81, 11.08. HRMS (ESI+) calculated for C$_{28}$H$_{49}$O$_4$ ([M + H]$^+$) 449.3631, found 449.3633.

Synthesis of (22*R*,23*R*)-6,6-ethylenedioxy-23-hydroxy-5α-campest-2-en-22-yl *p*-toluenesulfonate (16) and (22*R*,23*R*)-6,6-ethylenedioxy-22-hydroxy-5α-campest-2-en-23-yl *p*-toluenesulfonate (17)

p-Toluenesulfonyl chloride (200 mg, 1.05 mmol) was added to a solution of compound **6** (100 mg; 0.21 mmol) in pyridine (4 mL). The reaction mixture was stirred at room temperature for 3 days. Then, the mixture was diluted with ethyl acetate and aqueous solution of HCl (5%, 5 mL) was added. The aqueous layer was separated and extracted with ethyl acetate (2×). The combined organic layers were washed with water, dried over MgSO$_4$, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (10% ethyl acetate in cyclohexane). This chromatography did not lead to separation of both tosylates and they were used in the next reaction step as a mixture of isomers. Yield of both isomers 91 mg (69%). Analytical sample was used for HPLC separation (8% ethyl acetate in cyclohexane) to give two tosylates, **16** and **17,** as a viscous oil.

Compound **16**: ^{1}H NMR (500 MHz, CDCl$_3$): 7.85 (d, 2H, J = 8.4 Hz, H$_{Ts}$), 7.34 (d, 2H, J = 8.4 Hz, H$_{Ts}$), 5.67 (m, 1H, H-3), 5.55 (m, 1H, H-2), 4.77 (d, 1H, J = 8.6 Hz, H-22), 4.01-3.91 (m, 4H, H-23, H$_{ketal}$),3.80 (m, 1H, H$_{ketal}$), 2.46 (s, 3H, CH$_3$), 2.31 (d, 1H, J = 4.9 Hz, OH), 2.13-0.68 (m, 22H), 0.95 (d, 3H, J = 6.7 Hz, CH$_3$), 0.94 (d, 3H, J = 6.7 Hz, CH$_3$), 0.90 (d, 3H, J = 7.0 Hz, CH$_3$), 0.88 (d, 3H, J = 7.0 Hz, CH$_3$), 0.87 (s, 3H, CH$_3$), 0.66 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 144.50, 134.67, 129.61 (2×C), 127.62 (2×C), 125.72, 124.69, 109.92, 91.04, 71.41, 65.60, 64.12, 55.87, 53.52, 51.87, 48.08, 42.24, 41.26, 41.19, 40.88, 39.66, 37.72, 35.84, 33.27, 30.59, 27.95, 23.95, 21.70, 21.41, 20.84, 20.76, 20.70, 13.61, 12.57, 11.61, 10.17. HRMS (APCI+) calculated for C$_{37}$H$_{57}$O$_6$S ([M + H]$^+$) 629.3876, found 629.3865.

Compound **17**: ^{1}H NMR (500 MHz, CDCl$_3$): 7.83 (d, 2H, J = 8.3 Hz, H$_{Ts}$), 7.34 (d, 2H, J = 8.3 Hz, H$_{Ts}$), 5.66 (m, 1H, H-3), 5.55 (m, 1H, H-2), 5.04 (d, 1H, J = 8.6 Hz, H-22), 4.00-3.89 (m, 3H, H$_{ketal}$), 3.81-3.75 (m, 2H, H-22, H$_{ketal}$), 2.45 (s, 4H, OH, CH$_3$), 2.13-0.76 (m, 22H), 1.00 (d, 3H, J = 6.7 Hz, CH$_3$), 0.94 (d, 3H, J = 6.7 Hz, CH$_3$), 0.888 (s, 3H, CH$_3$), 0.8875 (d, 3H, J = 7.0 Hz, CH$_3$), 0.84 (d, 3H, J = 7.0 Hz, CH$_3$), 0.70 (s, 3H, CH$_3$). ^{13}C NMR (125 MHz, CDCl$_3$): 144.50, 134.44, 129.65 (2×C), 127.58 (2×C), 125.64, 124.81, 110.03, 89.23, 72.97, 65.59, 64.09, 55.79, 53.32, 52.06, 48.03, 42.29, 41.18, 41.16, 40.57, 39.73, 37.32, 35.86, 33.41, 29.76, 27.65, 23.95, 21.65, 21.42, 21.14, 20.86, 20.66, 13.60, 11.96, 11.77, 11.15. HRMS (APCI+) calculated for C$_{37}$H$_{57}$O$_6$S ([M + H]$^+$) 629.3876, found 629.3868.

Synthesis of (23*S*)-6,6-ethylenedioxy-5α-campest-2-en-23-ol (18) and (22*S*)-6,6-ethylenedioxy-5α-campest-2-en-22-ol (13)

1M Solution of LiAlH₄ in diethylether (1 mL, 1.0 mmol) was added dropwise to a solution of compounds **16** and **17** (50 mg, 0.08 mmol) in dry THF (5 mL). The reaction mixture was stirred at room temperature for 3 hours and then quenched with MeOH (3 mL). Saturated aqueous solution of sodium potassium tartrate (10 mL) was added and the mixture was diluted with ethyl acetate. The aqueous layer was separated and extracted with ethyl acetate (2×). The combined organic layers were washed with water, dried over MgSO₄, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (5-10% ethyl acetate in cyclohexane) to give crude mixture of compounds **13** and **18** (32 mg). Both compounds were separated by HPLC (8% ethyl acetate in cyclohexane) to afford two alcohols **13** (11 mg, 31%) and **18** (14 mg, 39%) as a viscous oil. The spectroscopic data for compound **13** were entirely consistent with those reported above.

Compound **18**: ^{1}H NMR (500 MHz, CDCl₃): 5.67 (m, 1H, H-3), 5.55 (m, 1H, H-2), 4.01-3.88 (m, 4H, H-23, H$_{ketal}$), 3.79 (q, 1H, J = 6.6 Hz, H$_{ketal}$), 2.12-1.86 (m, 5H), 1.80-0.75 (m, 20H), 0.98 (d, 3H, J = 6.7 Hz, CH₃), 0.96 (d, 3H, J = 6.7 Hz, CH₃), 0.93 (d, 3H, J = 6.7 Hz, CH₃), 0.89 (s, 3H, CH₃), 0.84 (d, 3H, J = 7.0 Hz, CH₃), 0.70 (s, 3H, CH₃). ^{13}C NMR (125 MHz, CDCl₃): 125.68, 124.81, 110.05, 70.98, 65.58, 64.09, 57.00, 55.93, 53.42, 48.07, 42.61, 42.45, 42.06, 41.22, 41.20, 39.80, 35.88, 33.97, 33.33, 30.43, 28.40, 24.13, 21.43, 21.11, 20.85, 20.49, 19.21, 13.60, 11.97, 9.27. HRMS (APCI+) calculated for C₃₀H₅₁O₃ ([M + H]⁺) 459.3838, found 459.3831.

Synthesis of (23*S*)-23-hydroxy-5α-campest-2-en-6-one (19)

Aqueous solution of HCl (5%, 0.5 mL) was added to a solution of compound **18** (12 mg, 0.03 mmol) in THF (3 mL) and the reaction mixture was stirred at 40 °C for 2 hours and then diluted with ethyl acetate. Organic layer was washed twice with water, dried over MgSO₄, filtrated and evaporated under reduced pressure. The residue was purified by column chromatography on silica gel (10% ethyl acetate in cyclohexane) to give compound **19** (9 mg, 83%) as a viscous oil. ^{1}H NMR (500 MHz, CDCl₃): 5.69 (m, 1H, H-3), 5.57 (m, 1H, H-2), 3.90 (ddd, 1H, J = 7.9, 5.7, 2.6 Hz, H-23), 2.38-2.34 (m, 2H), 2.27 (m, 1H), 2.09-1.89 (m, 5H), 1.75 (m, 1H), 1.66-1.04 (m, 16H), 0.99 (d, 3H, J = 6.7 Hz, CH₃), 0.96 (d, 3H, J = 6.7 Hz, CH₃), 0.93 (d, 3H, J = 6.7 Hz, CH₃), 0.84 (d, 3H, J = 7.0 Hz, CH₃), 0.72 (s, 3H, CH₃), 0.69 (s, 3H, CH₃). ^{13}C NMR (125 MHz, CDCl₃): 212.09, 124.94, 124.51, 70.97, 58.87, 58.67, 53.81, 53.34, 46.97, 42.87, 42.49, 41.98, 40.04, 39.50, 39.32, 37.67, 33.93, 30.41, 28.31, 23.93, 21.70, 21.11, 21.09, 20.48, 19.18, 13.50, 11.91, 9.26. HRMS (APCI+) calculated for C₂₈H₄₇O₂ ([M + H]⁺) 415.3576, found 415.3569.

Synthesis of (23*S*)-2α,3α,23-trihydroxy-5α-campestan-6-one (22-deoxycastasterone, 20)

Solution of osmium tetroxide in *t*-butanol (0.2M, 0.05 mL, 0.01 mmol) was added to a solution of compound **19** (8 mg, 0.02 mmol), hydroquinidine 4-chlorobenzoate (10 mg; 0.02 mmol),

methanesulfonamide (3 mg; 0.03 mmol), potassium carbonate (20 mg; 0.14 mmol), and potassium ferricyanide (45 mg; 0.14 mmol) in the mixture of *t*-butanol and water (3 mL; 1 : 1 v/v). The reaction mixture was stirred at room temperature for 10 h and quenched with saturated solution of Na_2SO_3 (2 mL) was then added. After an additional 30 minutes of stirring, the reaction mixture was diluted with ethyl acetate (20 mL) and extracted with water (2 × 20 mL). Organic layer was dried over $MgSO_4$, filtrated and evaporated under reduced pressure. The residue was purified by HPLC (70% ethyl acetate in cyclohexane) to give triol **20** (6 mg, 69%) as a white solid (lyophilized from *t*-butanol). ^{1}H NMR (500 MHz, $CDCl_3$ + 10% CD_3OD): 4.03 (m, 1H, H-3β), 3.89 (ddd, 1H, J = 8.0, 5.7, 2.6 Hz, H-23), 3.75 (m, 1H, H-2β), 2.68 (dd, 1H, J = 12.2, 3.4 Hz, H-5α), 2.29 (dd, 1H, J = 13.1, 4.6 Hz, H-7β), 2.07-1.05 (m, 22H), 0.97 (d, 3H, J = 6.7 Hz, CH_3), 0.95 (d, 3H, J = 6.7 Hz, CH_3), 0.92 (d, 3H, J = 6.7 Hz, CH_3), 0.83 (d, 3H, J = 6.7 Hz, CH_3), 0.75 (s, 3H, CH_3), 0.67 (s, 3H, CH_3). ^{13}C NMR (125 MHz, $CDCl_3$ + 10% CD_3OD): 212.38, 70.94, 68.33, 68.20, 56.79, 56.57, 53.62, 50.68, 46.72, 42.98, 42.58, 42.43, 41.89, 40.12, 39.37, 37.65, 33.90, 30.38, 28.29, 26.25, 23.90, 21.14, 21.09, 20.47, 19.11, 13.52, 11.96, 9.24. HRMS (APCI+) calculated for $C_{28}H_{49}O_4$ ([M + H]$^+$) 449.3631, found 449.3629.

References

1. Hurski, A. L., Ermolovich, Y. V., Zhabinskii, V. N. & Khripach, V. A. The development and use of a general route to brassinolide, its biosynthetic precursors, metabolites and analogues. *Org. Biomol. Chem*. **13**, 1446-1452 (2015).
2. Eignerova, B., Slavikova, B., Budesinsky, M., Dracinsky, M., Klepetarova, B., Stastna, E., Kotora, M. Synthesis of Fluorinated Brassinosteroids Based on Alkene Cross-Metathesis and Preliminary Biological Assessment. *J. Med. Chem*. **52**, 5753-5757 (2009).

¹H and ¹³C NMR spectra of compounds prepared

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
473.3631	473.3631	0.0	0.0	6.5	C30 H49 O4	13.1	n/a	n/a	30	49		4

MK-441
190205_08 36 (0.725) Cm (36)

05-Feb-2019
1: TOF MS ASAP+
1.75e+004

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	O	Si
587.4491	587.4496	-0.5	-0.9	6.5	C36 H63 O4 Si	55.7	n/a	n/a	36	63	4	1

MK-444
190205_13 32 (0.637) Cm (32)

05-Feb-2019
1: TOF MS ASAP+
1.02e+007

571.4182
587.4491
572.4214
573.4224
570.4098   574.4228
251.1865
281.2334
293.2328
313.2220
343.2688
355.2683
393.3196
437.3449
455.3551
456.3583
487.3622
527.3928 529.3726 530.3749

For Help, press F1

Scan: #820 µS: 4 IT: 0.24 NL: 3.88E8
Type: FTMS + p APCI corona Full lock ms [100.0000-750.0000]

MK-454
190206_02 38 (0.758) Cm (36:47)

06-Feb-2019
1: TOF MS ASAP+
6.38e+007

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
449.3627	449.3631	-0.4	-0.9	4.5	C28 H49 O4	37.6	n/a	n/a	28	49		4

MK-455A
190206_03 31 (0.620) Cm (28:39)

06-Feb-2019
1: TOF MS ASAP+
1.44e+007

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
491.3739	491.3736	0.3	0.6	5.5	C30 H51 O5	81.2	n/a	n/a	30	51		5

MK-458   06-Feb-2019
190206_17 33 (0.674) Cm (33)   1: TOF MS ASAP+
                                 1.16e+007

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
493.3895	493.3893	0.2	0.4	4.5	C30 H53 O5	47.4	n/a	n/a	30	53		5

12

13

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
459.3834	459.3838	-0.4	-0.9	5.5	C30 H51 O3	709.8	n/a	n/a	30	51		3

MK-520
200528_13 41 (0.829) Cm (41)
1: TOF MS ASAP+
4.21e+006

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
415.3578	415.3576	0.2	0.5	5.5	C28 H47 O2	1259.3	n/a	n/a	28	47		2

MK-521
200602_01 38 (0.758) Cm (38:43)

1: TOF MS ASAP+
6.96e+007

14

* residual CHD₂OD, ** water signal

* signal of CD₃OD

Mass	Calc. Mass	mDa	PPM	DBE	Formula	i-FIT	i-FIT Norm	Fit Conf %	C	H	N.	O
449.3633	449.3631	0.2	0.4	4.5	C28 H49 O4	1350.8	n/a	n/a	28	49		4

MK-529
200602_05 35 (0.707) Cm (29:40)

1: TOF MS ASAP+
4.67e+008

15

Scan: #2206 μS: 1 IT: 0.26 NL: 1.78E9
Type: FTMS + p ESI Full ms [250.0000-1000.0000]

16

646.4135

647.4160

629.3865

630.3900

631.3934

648.4192

651.3674

663.4525

667.3420

700.4598

457.3672

458.3705

447.3465

439.3568

18

Scan: #11005 μS: 4 IT: 0.51 NL: 2.70E8
Type: FTMS + p ESI Full lock ms [100.0000-750.0000]

* residual CHD$_2$OD, ** water signal

* signal of CD₃OD

Scan: #24216 μS: 4 IT: 0.26 NL: 3.30E8
Type: FTMS + p APCI corona Full lock ms [250.0000-750.0000]

431.3524

449.3629

282.2792

338.3419

313.2738

341.3050   432.3556   450.3661
359.3157   413.3417
331.2845   531.3687
                        487.3786

Relative Abundance (%)

300   400   500   600   700

m/z

71