Inactivation of Airborne Pathogen Surrogates by Triethylene Glycol

by

Grishma Desai[1], Emanuel Goldman[2], William Jordan[3], Jamie Balarashti[4], Jack Caravanos[5],

Rachel Edgar[6], Etienne Grignard[7], Gurumurthy Ramachandran[8], Gediminas Mainelis[9#]

[1] Grignard Company LLC, Rahway, NJ 07065.

[2] Department of Microbiology, Biochemistry, and Molecular Genetics, Rutgers-New Jersey

Medical School, Newark, NJ 07103.

[3] Former Deputy Director, Programs, Office of Pesticide Programs, Environmental Protection

Agency. William Jordan Consulting, Washington, DC 20016.

[4] President and EOC, Aerosol Research and Engineering Labs, Overland Park, Kansas, 66213.

[5] Clinical Professor of Environmental Public Health Services, New York University, New York,

NY 10012.

[6] Department of Infectious Disease, Imperial College London, London SW7 2AZ, UK. Francis

Crick Institute, 1 Midland Road, London NW1 1AT, UK.

[7] Founder, CEO, Bleu Garde, LLC, Rahway, NJ 07065.

[8] Department of Environmental Health and Engineering, Johns Hopkins Education and Research

Center for Occupational Safety and Health, Bloomberg School of Public Health, Whiting School

of Engineering, The Johns Hopkins University, Baltimore, MD 21218.

[9] Department of Environmental Sciences, School of Environmental and Biological Sciences,

Rutgers, The State University of NJ, New Brunswick, NJ 08901.

Address correspondence to Gediminas Mainelis, mainelis@envsci.rutgers.edu

Keywords:

Airborne transmission; Antimicrobial efficacy; Aerosolized antimicrobials; Triethylene glycol;

Surrogate microorganisms

Running title: Inactivation of Bioaerosols by Triethylene Glycol

Abstract

The COVID-19 outbreak brought to the fore the importance of airborne transmission in spreading human infectious diseases and highlighted the need for sustainable mitigation strategies. Triethylene glycol (TEG) has been documented as having microbicidal capabilities and has been proposed as one such mitigation strategy. Aerosolized TEG exhibits antimicrobial activity against airborne microorganisms; Grignard Pure™ Technology was developed to safely aerosolize TEG for air decontamination. Here we show that this TEG formulation effectively inactivates airborne microorganisms, resulting in 2 to 4.5 net log reduction in concentration of viable bacteria, viruses, and mycobacteria within 30-60 minutes at TEG concentration (aerosol + vapor) of ~0.7 mg/m^3, which is well within the range considered safe for humans. Our data also demonstrate that aerosolizing both the test organisms and the antimicrobial product provides a more accurate and relevant measure of the product's efficacy for indoor usage compared to traditional surface - or solution - based disinfection assays. Accurate evaluation of antimicrobial efficacy is a crucial step in adopting novel interventions and tools to control airborne pathogens that pose a public health risk. Our findings argue that testing protocols must match the intended use of any intervention. Given the safety concerns of aerosolizing human pathogens for direct testing of airborne infectious burden, we also advance an approach for selecting suitable surrogate microorganisms based on their phenotypic and biophysical similarity to corresponding pathogenic species.

Importance

During the COVID-19 pandemic, personal protective equipment, social distancing, and even vaccinations proved sub-optimal in controlling the spread of COVID-19. Public health practice and the hierarchy of controls prefer primary prevention, whereby the pathogen is removed or destroyed before exposure to the public. Triethylene glycol (TEG) has the potential to inactivate airborne pathogens and limit their spread. TEG is designated a "safer chemical" by the US EPA and has been used for decades in aerosol deodorizers and theatrical special effects. This study shows that aerosolized TEG is highly effective at eliminating a wide spectrum of viable airborne pathogen surrogates at concentrations well below the threshold of safety concern and so it may afford significant protection against the transmission of infectious agents with pandemic potential.

INTRODUCTION

Spread of diseases *via* the air significantly affects global health, economies, and societal structures. Due to their global impact, COVID-19, influenza, and tuberculosis (TB) are probably the best-known examples of airborne diseases. To date, COVID-19 has led to over 7 million fatalities globally (https://data.who.int/dashboards/covid19/cases). Seasonal flu affects millions of people each year, with the World Health Organization (WHO) estimating between 290,000 and 650,000 annual deaths (https://www.who.int/news-room/fact-sheets/detail/influenza-(seasonal)). Additionally, TB continues to be among the leading causes of death worldwide, with around 1.25 million deaths reported in 2023 according to WHO data (https://www.who.int/news-room/fact-sheets/detail/tuberculosis).

The COVID-19 pandemic prompted a re-evaluation of the importance of airborne transmission as a driver of infections, and therefore the interventions needed to combat them (1). When an infected person coughs, sneezes, speaks, or breathes, respiratory pathogens are exhaled within droplets on a size continuum ranging from sub-micrometer to millimeters in size (2). There is an emerging consensus that inhalation of these infectious particles can transmit disease over both short and long distances, and that many factors beyond droplet size determine airborne dissemination (3). The airborne route is likely the dominant mode of transmission for many respiratory diseases, rather than direct deposition of large droplets onto the eyes, nose, or mouth of a susceptible individual, and may be accelerated in enclosed spaces where infectious particles tend to accumulate (1). In addition to exhalation, infectious agents can disperse into the air from other sources like toilet flushing (e.g. gastrointestinal pathogens such as norovirus), skin blisters (e.g. measles, chickenpox, *Staphylococcus aureus*), or release of fungal spores (e.g. *Aspergillus*) (4)

Public health strategies to curb the spread of airborne diseases typically include practices like hand hygiene, the use of face masks and other personal protective equipment, and social (physical) distancing. Although these strategies can help, they rely on individual compliance. Vaccination is another pillar of infection control, but development or adaptation of existing vaccines requires time, and the primary goal is to protect the population from severe disease rather than preventing transmission; people vaccinated against COVID-19 who became infected could still efficiently transmit SARS-CoV-2, for example (5). Ultimately, reducing the infectious burden within the environment will reduce infection risk, through improved ventilation, pathogen-neutralizing technologies, or adjusting indoor climate (6). An ideal transmission-blocking intervention should be safe, cost-effective, scalable, environmentally responsible, pan-microbial with minimal chance of resistance development, and compatible with other infection control measures for a layered approach. To have the greatest impact, an intervention should be suitable for long-term proactive use and rapid global deployment early in a pandemic. Small non-toxic glycols meet these criteria, and the effectiveness of triethylene glycol (TEG) against various airborne microbes is the focus of this paper.

Microbicidal properties of TEG were identified over 70 years ago, and TEG was identified as a substance against airborne biological threats (7). The microbicidal effects of TEG occur when a sufficient quantity of TEG vapor condenses on particles or droplets carrying microbes (8), which leads to their quick inactivation (the specific mechanism of action is discussed below in this paper). A concentration as low as $2 - 5$ mg/m^3 of TEG in the air was found to be enough for achieving "maximum germicidal action" against a range of airborne infectious agents (7). These

concentrations are well below the threshold of safety concern (9). Extensive research conducted at the time showed that TEG vapor effectively neutralizes viruses or bacteria responsible for influenza, meningopneumonitis, and psittacosis (10) as well as pneumococci types I, II, and III, beta hemolytic streptococci groups A and C, staphylococci, influenza bacilli, *Escherichia coli*, and *Bacillus aerogenes* (11), and other microbes. However, interest in such primary prevention methods declined as antibiotics became more widely available and effective.

TEG received renewed interest during the COVID-19 pandemic as a potential tool to rapidly inactivate the airborne SARS-CoV-2 virus, thus limiting its spread. In a recent study, we have shown that TEG achieves a net 3-log inactivation within 15 minutes against airborne bacteriophage MS2, a surrogate for viral pathogens, including SARS-CoV-2 (12). Since MS2 is a non-enveloped virus, we speculate that TEG would be even more efficacious against enveloped viruses, including SARS-CoV-2. A related compound, propylene glycol (PG), has also been shown to inactivate several airborne respiratory viruses (13). However, PG requires higher concentrations than TEG, as was observed in the 1940s (8). The antimicrobial activity of various glycols has recently been reviewed (14). The efficacy of TEG against airborne MS2 at different building operational parameters was also recently investigated (15). Thus, evidence indicates that TEG vapor can be used to inactivate airborne microorganisms, thus potentially reducing disease transmission *via* airborne route.

While recent inactivation studies focused on MS2, the application of vaporized TEG as a public health tool for infection control requires this compound to be able to inactivate a variety of airborne microorganisms. This study fills this data gap by demonstrating the broad-spectrum efficacy of TEG against Gram-negative and Gram-positive bacteria, enveloped and non-enveloped viruses, a mycobacterium, and a mold species. The study developed protocols to accurately assess the potential of TEG vapor to inactivate microorganisms in the air, thus enabling a comparison of the microbicidal activity during the intended airborne phase application with conventional surface disinfection testing protocols. The results of our study demonstrate and highlight the need for testing protocols that reflect the 'real world' application of novel infection control measures, and the need for appropriate pathogen surrogates that can be safely used to comprehensively assess the control of airborne transmission. Otherwise, surface-centered testing (as is typically performed) obscures the ability of an airborne phase product like TEG to inactivate microorganisms that are primarily infectious via the airborne route.

MATERIALS AND METHODS

Test microorganisms

The selection of test microbes was guided by biosafety constraints associated with aerosolized pathogen work. Many clinically relevant organisms, such as *Staphylococcus aureus*, require BSL-2 containment for benchtop procedures and BSL-3 or higher for aerosolized studies. As most commercial and academic aerosol testing laboratories, including ours, are equipped with only BSL-2 aerosol chambers, testing in the airborne phase is limited to BSL-1 organisms. Biosafety guidelines, such as the CDC BMBL (6th Edition), explicitly note that additional primary containment and personnel precautions—such as those recommended for BSL-3—should be considered for activities with a high potential for aerosol or droplet production (16). Similarly, the

WHO Laboratory Biosafety Manual (3rd Edition) emphasizes that aerosol-generating procedures may require a higher biosafety level than that indicated by an organism's Risk Group, based on risk assessment (17). Therefore, appropriate BSL-1 surrogates, based on the similarity of the phenotypic traits, were selected to assess the response of corresponding BSL-2 pathogens to treatment by TEG.

This study prioritized diverse microbial species, representing a broad spectrum of potential pathogens. The study aimed to evaluate the antimicrobial agent's broad-spectrum potential under realistic conditions by incorporating organisms with varying susceptibilities and structural differences. The organisms selected for the study included:

- Gram-negative bacteria: *Pseudomonas fluorescens* (ATCC 13525) (American Type Culture Collection, Manassas, VA), *Salmonella typhimurium* (ATCC 53648), and *Klebsiella aerogenes* (ATCC 51697).
- Gram-positive bacteria: Methicillin-resistant *Staphylococcus epidermidis* (ATCC 12228) and *Listeria innocua* (ATCC 33090).
- Viral species: Non-enveloped bacteriophage MS2 (ATCC 15597-B1) and enveloped bacteriophage Phi6 (ATCC 21781-B1).
- Acid-fast Gram-positive bacterium: *Mycobacterium smegmatis* (ATCC 607).
- Mold species: *Aspergillus brasiliensis* (ATCC 16404).

The microbes above were exposed to airborne TEG in the airborne phase, and to liquid TEG while surface-bound to carriers.

In another subset of experiments, we evaluated the inactivation of surface-bound actual pathogenic strains and their Biosafety Level 1 (BSL-1) surrogate counterparts (i.e., *S. epidermidis, K. aerogenes*, *P. fluorescens*) in parallel when exposed to TEG vapor. This test also advanced the selection of appropriate surrogates, providing justification for their use in evaluating airborne antimicrobial efficacy of TEG.
The pathogenic species were:
- Gram-positive bacteria: methicillin-resistant *Staphylococcus aureus* (ATCC 6538)
- Gram-negative bacteria: *Klebsiella pneumoniae* (ATCC 4352), Gram-negative bacteria: *Pseudomonas aeruginosa* (ATCC 15442)

Inoculum preparation

Pure strain seed stocks were sourced from the ATCC. Working stock cultures were prepared under aseptic conditions using a Class II biological safety cabinet (Labconco Purifier Class II Safety Cabinet, Kansas City, MO), following ATCC guidelines. Strains were revived in appropriate growth media and incubated under conditions specific to each organism, following ATCC guidelines. Except for *Listeria innocua*, all other bacterial species were cultured in tryptic soy broth and plated on tryptic soy agar. *Listeria innocua* was cultured in brain heart infusion broth and plated on brain heart agar. *Aspergillus brasiliensis* was used as a dry spore powder and plated on potato dextrose agar. Serial dilutions were performed in sterile phosphate-buffered saline (PBS) supplemented with 0.005% Tween 80 to facilitate homogeneity.

For virus strains, the host bacteria (*E. coli* for MS2 and *P. syringae* for Phi6) were grown
overnight in tryptic soy broth at 37°C for *E. coli* and 28°C for *P. syringae*. During the
logarithmic growth phase, the bacterial suspension was infected with either MS2 or Phi6
bacteriophage. After 24 hours, the infected cells were lysed, and cellular debris was removed by
centrifugation at 3000 rpm for 15 minutes. The resulting supernatant was poured off from the
cellular debris, sampled, serially diluted, and plated to determine the viral concentration.
195
196 The stock concentrations of test microorganisms were verified through colony-forming unit
197 (CFU) or plaque forming unit (PFU) enumeration. Stock dilutions with microorganism
198 concentrations ranging from 10^9 to 10^{10} CFU/ml or PFU/ml were used for aerosolization, as
199 described below.

Sample processing

In the inactivation experiments described below, the airborne test microorganisms were directly
captured into PBS using impingers, while the surface-bound organisms on glass coupons were
eluted into PBS using 50ml conical tubes (Celltreat, Pepperell, MA), followed by enumeration
by culturing in triplicate. If necessary, the collected samples were diluted by a series of ten-fold
dilutions before culturing. Appropriate dilutions of virus samples underwent small-drop plaque
assays technique, while bacterial and mold spore samples were plated using a spread plate
technique. *Listeria innocua* was cultured on Tryptic Soy Agar supplemented with 5% sheep
blood, *Aspergillus brasiliensis* was grown on Potato Dextrose Agar with chloramphenicol, and
Mycobacterium smegmatis was cultured on Middlebrook 7H11 Agar. Tryptic Soy Agar was used
for all other species. All media were supplied by Hardy Diagnostics, Santa Maria, CA. All
samples were plated across a minimum of three dilution ranges, with triplicate plating per
dilution range on pre-labeled, culture plates with species-appropriate agar and organism. Plates
were incubated at species-specific temperatures ranging from 28–37°C for 24–48 hours before
enumeration. Counts between 5–50 colony-forming units (CFU) or plaque-forming units (PFU)
per plate were recorded; counts outside of this range were not counted. This approach ensured
consistent quantification across all tested conditions.

Inactivation of airborne microbes by airborne TEG (aerosol and vapor).

Airborne inactivation studies were conducted using paired trials for each microbial species
discussed in this paper. For every species, a positive control trial and a corresponding
inactivation trial were performed using a single, unified biological stock that was split between
the two conditions. This approach ensured both trials began with identical starting material. All
other trial conditions—such as the nebulizer type, nebulization pressure and duration, hold and
sampling times, and methods for collection, elution, and plating—were kept constant between
the paired trials to ensure consistency.

During the study, airborne concentrations of viable test microbes were measured at different time
points for each paired control with and without exposure to the inactivating agent, triethylene
glycol (TEG). TEG was aerosolized through Grignard Pure™ Technology (Bleu Garde LLC,
Rahway, NJ) developed a few years ago to disperse TEG in aerosol and vapor form.

Inactivation of airborne microbes was investigated in an insulated ~16 m³ stainless steel (SS 304)
bioaerosol chamber with internal dimensions 9.1 ft × 9.1 ft × 7 ft (2.8 x 2.8 x 2.1 m), with a total
volume of just over 16,000 liters. It features three windows and a hermetically sealed door to

maintain environmental integrity. Environmental parameters, including temperature and relative humidity, were continuously monitored using embedded sensors (HT.w Sensors, SensorPush, Brooklyn, NY) within the chamber. During experiments, the average temperature and RH were 22-24°C and 30-40%, respectively. Aerosol samples were collected through two stainless steel probes (3/8-inch inner diameter) located 18 inches from the chamber's sidewalls at a height of 36 inches. Sampling points were positioned at the diagonal corners of the chamber to ensure representative samples. Two interior mixing fans, operating at ~50 cubic feet per minute (cfm) each (equivalent to about 10 internal ACH), were used for the entirety of all trials to provide bulk mixing and to ensure homogeneity of the chamber during the test period. This setup ensured uniform sampling across the chamber's interior for precise and reproducible results (18).

The timeline of inactivation experiments is shown in Figure 1. For most of the organisms tested, Grignard Pure™ solution in its undiluted form (Lot#021722) was aerosolized via the Aura device (Bleu Garde, LLC), which was positioned in the center of the chamber and activated remotely. *A. brasiliensis* was treated with the Clearify device (Bleu Garde, LLC) due to anticipated need of higher TEG concentration because of spores. First, the chamber was pretreated with aerosolized Grignard Pure™ for 30 minutes (from t = -40 min to t = -10 min in Fig. 1), which simulated an indoor environment treated with Grignard Pure™. After this 30-min pretreatment period, while Aura (or Clearify) dispersal device continued to operate, bioaerosols were aerosolized into the chamber over a 20-min period using a 1-inch diameter stainless steel conduit. Bacteria and viruses were aerosolized using 24-jet Collison nebulizer (CH Technologies, Westwood, NJ) operated at 40 psi. *A. brasiliensis* spores were disseminated via a dry powder eductor constructed from stainless steel and operated using in-house filtered compressed air. Since microorganisms interacted with TEG while being introduced, we designated the mid- point of this 20-min aerosolization period as t= 0 min. Once the bioaerosol introduction was completed, this time marked time t = 10 minutes in our treatment timeline, because that is the average duration that the organisms have interacted with airborne TEG (Fig. 1). Bioaerosol samples were started at time points t = 10, 35, 50, and 80 minutes using two AGI-30 impingers (Ace Glass Incorporated, Vineland, NJ), and sampling duration was 10 minutes. Impingers contained 20 mL of sterile PBS supplemented with 0.005% Tween 80 to enhance capture efficiency and promote microorganism dispersion within collection liquid. The impingers operated at a flow rate of 12.5 L/min, provided by 1/3 HP Gast Rotary Vane Vacuum Pumps (Eureka, MO). After each sampling period, the contents of the two impingers were combined to provide a chamber average and transferred to sterile conical tubes (Celltreat, Pepperell, MA) for microbial enumeration. A single low velocity mixing fan (12" Oscillating Black HTF1220B, Honeywell, Charlotte, NC) was used in the chamber during testing to ensure a homogenous microbe distribution throughout the chamber and promote GP and organism mixing. Between samples, impingers were thoroughly cleaned and sterilized by immersion in a 3% bleach solution, followed by rinses with tap water and distilled water before air drying. Following the experiments, the remaining particles and vapor were evacuated from the chamber (19).

The aerosol concentration of Grignard Pure™ was estimated using a calibrated TSI SidePak AM520 (TSI, Inc., Shoreview, MN). The Grignard Pure™ aerosol concentration can be easily converted to TEG aerosol concentration as described in our earlier study (12). The total TEG

concentration (aerosol + vapor) was estimated using an earlier-developed correlation between
AM520 particle concentration data and NIOSH tube sampling method 5523, which determines
the total (aerosol + vapor) airborne concentration of TEG. During the inactivation of most
airborne microbes (except *A. brasiliensis*), Aura device was used, and the average TEG aerosol
and total concentrations were 0.04 ± 0.02 and 0.66 ± 0.35 mg/m^3, respectively. In anticipation of
higher resistance of fungal spores, for testing with *A. brasiliensis*, we used the Clearify device,
and the total TEG concentration was increased to 1.96 mg/m^3, of which 0.3 mg/m^3 was in
aerosol form. Data from this experiment are provided in Table S1 (Supplemental Material).

In control trials for each test microbe, the procedure was exactly the same as described above,
except Grignard Pure™ was not introduced into the chamber. The control trial accounted for the
loss of viable microorganisms due to gravitational settling and natural die-off. The efficacy of
TEG was determined by comparing the airborne concentrations of viable microorganisms -
measured as colony-forming units per m^3 (CFU/m^3) for bacteria and plaque-forming units per m^3
(PFU/m^3) for viruses - collected at the same time points as during air treatment with TEG. The
inactivation efficiency was determined for each sampling point as a percent value and log
reduction relative to microorganism recovery without treatment. The initial microorganism
concentrations with and without treatment by TEG were normalized. The applied mathematical
expressions are presented in our earlier work (12). For all microbes, except MS2, there was a
single positive control trial and a corresponding TEG inactivation trial. For MS2, two sets of
experiments were conducted, and average values are presented.

Neutralization Verification

Neutralization verification was performed using procedures aligned with EPA SOP MB-26-03,
Standard Operating Procedure for Neutralization of Microbicidal Activity Using the Quantitative
Method for Evaluating Bactericidal and Mycobactericidal Activity of Microbicides on Hard,
Non-Porous Surfaces (20) .

These tests confirmed that microbial inactivation attributed to TEG occurred exclusively during
airborne exposure and not due to residual TEG activity post-sampling. To assess this, test
organisms were directly exposed to TEG in liquid at defined concentrations and contact times,
then plated and enumerated. Results were compared to unexposed controls to determine any
residual biocidal activity post-collection.

TEG exposure concentrations were calculated based on known airborne concentrations and
impinger sampling volumes. These values were then multiplied by a safety factor of 10× to
establish the challenge concentration for neutralization testing. A neutralizer solution consisting
of TEG10× + phosphate-buffered saline (PBS) + 0.005% Tween 80 was used; PBS + 0.005%
Tween 80 served as the neutralizer control.

Triplicate 50 mL conical tubes were prepared for both the TEG-neutralizer and control solutions,
each containing 20 mL. Cultures were diluted to $<1.00 \times 10^3$ CFU or PFU/mL before being
added to the tubes. After a 15-minute contact time, samples were serially diluted, plated using
previously described methods, incubated, and enumerated.

Comparisons were made to PBS-only controls and water-diluted controls to verify neutralization
efficacy. ANOVA analysis confirmed that PBS + 0.005% Tween 80 had no statistically significant

impact on microbial recovery for any species tested, validating that reported reductions were due
solely to in-chamber inactivation and not post-collection residual activity.

Inactivation of surface-bound organisms by liquid TEG

Currently, the efficacy of most antimicrobial products is typically evaluated using surface-
focused methods - such as ASTM E1153 (21), ASTM 1053 (22) and the AOAC use-dilution
method (23) – that measure the reduction of microorganisms deposited on hard, non-porous
carriers following exposure to a liquid, spray, or wipe. We posited that products intended for
inactivation of airborne microorganisms must be evaluated using test methods specifically
designed to assess their efficacy against airborne microorganisms, as surface-based tests do not
capture this mode of action. The comparison of the two modes of inactivation (e.g.,
microorganisms in the air vs surface-bound) is crucial to properly determine a product's
suitability to act an airborne antimicrobial. Therefore, inactivation of airborne microorganisms
by Grignard Pure™ in aerosol and vapor form described above was compared to inactivation of
surface-bound microorganisms by liquid Grignard Pure™. The methodology for surface-bound
testing adhered to ASTM E1153 (21) guidelines for bacterial and fungal assessments on non-
porous surfaces and ASTM E1053-20 (22) for virucidal evaluations. While Grignard Pure™ is
not classified as a disinfectant or sanitizer, these standardized protocols provided a structured
framework to assess the efficacy of TEG against microbial targets in both aerosolized and liquid
forms.

Microbial inocula were prepared using the same procedure as described above for aerosol phase
tests; however, to simulate environmental contamination, a soil load consisting of 0.3% bovine
serum albumin, 0.3% mucin, and 0.3% yeast extract was added to the microbial stocks in
accordance with ASTM E2197 and OECD Guideline 13 standards (24).

The timeline for surface-bound microbe inactivation experiments is shown in Figure 1b.
Quadruplicate sets of 304 stainless steel carrier discs (0.8mm thick, 2cm diameter) were
inoculated (covered) with 0.03 mL of prepared microbial suspension, ensuring uniform coverage.
Carriers were dried at 35–39°C for approximately 20 minutes or until visibly dry. For exposure
to TEG, carriers were submerged in 5 mL of the Grignard Pure™ solution for 0, 15, 30, and 60
min. For 0-min exposure, the carriers were dipped into the Grignard Pure™ solution and
immediately pulled out. After exposure, 20 ml of PBS + 0.005% Tween 80 was added to the
container with the test solution and carrier to halt antimicrobial activity. The container was then
vortexed for 2 minutes, followed by shaking for 30 seconds to recover microbes from the carrier
surface. The recovered microbes were then cultured onto species-specific media using a standard
culturing or small plaque assay across various dilutions and incubated at species-specific
temperatures (28–37°C) for 24–48 hours before enumeration. The number of CFU or PFU
recovered from exposed carriers were compared with the number of carriers not exposed to
liquid Grignard Pure™ but otherwise handled in the same way. These control carriers were
exposed to the air for 0, 15, 30, 60 min. The inactivation efficacy was determined as a percentage
and net log reduction relative to the unexposed carriers. Data from this experiment are provided
in Table S2 (Supplemental Material).

Inactivation of surface-bound organisms by TEG vapor

As briefly discussed above, the selection of appropriate surrogate microorganisms is important
because many laboratories are limited to working with airborne BSL-1 organisms only, yet many
regulatory procedures require showing evidence of antimicrobial efficiency against BSL-2
organisms. Therefore, we compared the inactivation of several BSL-1 microorganisms with their
BSL-2 counterparts when exposed to TEG aerosol and vapor while surface-bound. Microbial
cultures and their carriers were prepared as described above for the testing of inactivation by
liquid TEG. The experiments were carried out in the same bioaerosol chamber at an average
temperature of 22-24°C and 30-40% RH. The chamber was then pretreated with aerosolized
Grignard Pure™ for 30 minutes using the Aura device. TEG concentration in this experiment
was 0.54 mg/m^3. Dried carriers in triplicate were placed on a stainless-steel workbench within
the bioaerosol chamber at a distance of approximately 3 feet from the dispersal device. The
chamber was sealed, and the Aura dispersal device was operated according to its specifications
for a total exposure duration of 90 minutes. After exposure, carriers were transferred to labeled
conical tubes containing neutralizing buffer and vortexed for 2 minutes, followed by shaking for
30 seconds to recover microorganisms.

The recovered microorganisms were serially diluted in microcentrifuge tubes containing sterile
PBS as diluent and enumerated as described above. The number of CFUs recovered from
exposed carriers were compared with the number of CFU from carriers not exposed to TEG in
the chamber but handled in the same way. The inactivation efficacy was determined as a
percentage and net log reduction relative to unexposed carriers. Data from this experiment are
provided in Table S3 (Supplemental Material).

RESULTS

Overall inactivation of airborne microorganisms

Inactivation of viable test microorganisms in the air by airborne TEG and on hard surface by
liquid TEG as supplied by Grignard Pure™ is presented in Figures 2 and 3. Figure 2 shows data
for tested Gram-positive and Gram-negative bacteria, while Figure 3 presents inactivation data
for tested bacteriophages, acid-fast microorganism, and mold. The inactivation data are presented
as net log inactivation and also as net percent inactivation for different sampling time points
(Figure 1). The raw data are provided in Table S1 (Supplemental Material).

Substantial inactivation of all tested airborne microorganisms, except mold, was observed at the
10-minute sampling point, with bacterial inactivation efficacy ranging from 50 to 95%. Both
bacteriophages were inactivated with 99% efficacy at the same 10-minute sampling time point.
Inactivation of *A. brasiliensis* was 30% at the same time point. All microbes except *A.*
brasiliensis were inactivated with at least 99% efficacy at the 35-minute sampling mark.
Inactivation of *A. brasiliensis* spores at that time point was ~80%. At later sampling points, the
net inactivation efficacy remained steady or slightly decreased (e.g., Phi6 and *P. fluorescens*) due
to increasing natural die-off of the organisms, which led to a reduced net efficacy because the
initial concentration of airborne microbes turned out not to be high enough. Both Phi6 and *P.*
fluorescens are known to be sensitive to environmental stressors (25, 26), and prohibitively large
airborne concentrations of them are needed to observe net inactivation over longer periods. In
contrast, inactivation of the same organisms on hard surfaces proceeded much more slowly, with

lower inactivation than in the airborne phase at the 30-minute mark for all tested microbes, except Phi6 bacteriophage.

Inactivation of airborne bacteria

Among airborne bacteria, the highest inactivation was achieved for Gram-positive bacteria, where viable *S. epidermidis* was reduced by 4.28 logs at the 35-minute sampling point, and its inactivation was more than 4 logs at 50 and 80 minutes; *L. innocua* showed more than 3 log reduction at 50- and 80-min sampling points (Fig. 2). In contrast, hard surface treatment resulted in slower viability reductions, with *L. innocua* and *S. epidermidis* inactivated by less than 1 log by 30-min treatment and less than 2 logs by 60-min treatment. A similar dynamic was observed for all Gram-negative bacteria. Treatment of airborne organisms again proved superior, with *P. fluorescence* reduced by 3.10 logs at 35 minutes, followed by *S. typhimurium* (2.5 logs) and *K. aerogenes* (2 logs). At later sampling points, their inactivation stayed the same or slightly decreased (e.g., *P. fluorescens*) due to the role played by the natural die-off and settling in inactivation calculation. Viability reduction on hard surface by liquid TEG was notably slower but eventually reached about 2.5 logs for all three bacteria by 60 minutes. These results emphasize the more rapid and significant inactivation of airborne bacteria by airborne TEG compared to surface treatments by liquid TEG.

Inactivation of airborne viruses, mycobacteria, and fungi

Figure 3 shifts focus to the inactivation of bacteriophages. Aerosol treatment was markedly faster for MS2, achieving 2.7-log reduction at 35 minutes and 3.46-log reduction at 50 minutes, before plateauing at 3.42-log by 80 minutes. In contrast, the reduction of MS2 on a hard surface was slower: only a 0.20-log reduction was achieved by 60 minutes. Phi6 showed a net inactivation of 2.1 logs at 35 min, but then the efficiency declined to about 2 logs at 80 min because of its natural die-off. Due to the sensitivity of Phi6 to environmental stressors, its inactivation on the surface was also relatively rapid, reaching 2.2 logs at the 30-minute mark and exceeding that in the airborne state at the 60-minute mark.

Airborne *M. smegmatis* was also rapidly inactivated: by 3-logs at the 35 min mark, and then the inactivation reached a plateau of 3.54 log by 80 minutes. Inactivation of this bacterium on the hard surface proceeded substantially slower, but it reached 2.93-logs by 60 min mark. Its inactivation profile in the air was slower compared to Gram-positive bacteria, but similar to that of some Gram-negative bacteria.

Mold spores are known to be resistant to environmental stress, but airborne TEG achieved 1.5-log inactivation of airborne *A. brasiliensis* after 50 min, and the level stayed the same after 80 minutes of interaction. Inactivation of these spores on the surface was much slower and reached only 0.19-log reduction in 60 minutes. *A. brasiliensis* appears to be more resistant overall compared to tested bacteria, for both airborne and surface testing. It is also worth mentioning that the airborne TEG concentration when inactivating *A. brasiliensis* was almost three times higher than when treating other organisms, confirming the anticipated hardiness of this species. Nevertheless, about 95% of the airborne spores were inactivated after 50-min treatment by airborne TEG.

Summary of airborne inactivation results

Taken together, Figs. 2-3 demonstrate that TEG provided by Grignard Pure™ is highly effective
in reducing the presence of a wide range of viable organisms, particularly in airborne
environments where aerosol application consistently outperforms hard surface treatments by
liquid Grignard Pure™ in both speed and magnitude of reduction. The findings highlight its
robust efficacy against bacteria such as *S. epidermidis* and *S. typhimurium* as well as against
bacteriophages like MS2 and Phi6, while also confirming its effectiveness against more resilient
organisms like *M. smegmatis* and *A. brasiliensis* over time. These results underscore the utility of
Grignard Pure™ in antimicrobial strategies aimed at controlling airborne pathogens in enclosed
diverse settings, while offering complementary benefits for surface decontamination over longer
durations when applied in liquid form.

Comparison of inactivation of pathogens with their corresponding surrogates

The next set of experiments examined the inactivation of a select set of BSL-1 organisms and
their BSL-2 counterparts when first deposited on slides and then exposed to TEG vapor and
droplets for 90 min (Fig. 4). The selected BSL-1 organisms serve as surrogates for their BSL-2
counterparts. Among the BSL-1 surrogates, *K. aerogenes* demonstrated the highest reduction
(0.49 log), closely mirroring its BSL-2 counterpart *K. pneumoniae* (0.46-log reduction). This
similarity confirms the suitability of *K. aerogenes* to be used as a surrogate for pathogenic *K.*
pneumoniae. Likewise, inactivation of *P. fluorescens* (0.18 logs) was similar to that of BSL-2 *P.*
aeruginosa (0.21 logs), further validating its use as a representative model for air testing studies.
Both Gram-positive species, BSL-1 *S. epidermidis* (0.17-log reduction) and BSL-2 *S. aureus*
(0.14-log reduction), showed lower reductions compared to other species but maintained
consistent responses between the surrogates and pathogenic counterparts. Overall, a paired t-test
of all three microorganism pairs showed that there was no statistically significant difference
($p > 0.05$) among inactivation of the BSL-1 bacteria (surrogates) and BSL-2 bacteria (pathogens).
This consistent response between the BSL-1 bacteria and their BSL-2 counterparts demonstrates
the relevance and suitability of selected surrogates in assessing treatment efficacy against higher-
risk BSL-2 pathogens under safe laboratory conditions. Fig. 4 highlights how closely surrogate
organisms replicate the behavior of their pathogenic counterparts under similar exposure
conditions. Notably, Gram-negative *K. aerogenes* and *P. fluorescens* provide reliable models for
airborne pathogen testing due to their similar responses to *K. pneumoniae* and *P. aeruginosa*,
while the same can be said about Gram-positive *S. epidermidis* and *S. aureus*.

DISCUSSION

Efficacy Testing Protocol

Although standardized methods exist for evaluating the efficacy of antimicrobial products in
liquid suspension and on surfaces, there is a notable lack of standardized testing methods for
antimicrobial products intended to be used against pathogens in the air (27). Moreover, even
well-established surface testing methods may not be directly relevant for aerosol-transmitted
pathogens, as they fail to capture the dynamics of airborne exposure and transmission risk. The
exception to this for airborne testing is ASHRAE 185.3 (28) standard that defines the testing
methods for evaluating commercial and industrial in-room air-cleaning devices and systems
specifically for their effectiveness in removing or inactivating microorganism bioaerosols in a

test chamber. Specifically, for airborne antimicrobials to be effective, they must directly contact
pathogens in the air—either as vapor or aerosolized droplets. When a liquid antimicrobial
product is tested for efficacy, it may be sprayed as large droplets or wiped onto a surface
(coupon) or evaluated using suspension-based methods. In these cases, the product typically
comes in contact with either the dried form of the microbe or interacts directly with microbes in
a liquid suspension or wet droplet form. Unlike surface-based applications, where products
interact with dried microbes, airborne delivery involves the dynamic interaction of the agent with
suspended microorganisms. Our findings with TEG highlight that the mode of application
significantly influences inactivation efficacy and time, reinforcing the need to evaluate
antimicrobial products under conditions that reflect their intended use. In addition, aerosolizing
test organisms allows for a more realistic assessment of antimicrobial efficacy by accounting for
environmental factors like humidity and temperature that influence microbial susceptibility.
Compared to surface-bound microbes, aerosolized organisms better represent real-world
exposure to airborne pathogens. Moreover, dried surface samples can artificially shield microbes,
potentially underestimating a product's effectiveness in air.

The collected data across multiple trials in this study consistently demonstrate that TEG
delivered by Grignard Pure™ exhibits significantly greater antimicrobial efficacy in aerosolized
applications compared to hard surface treatments, underscoring the critical importance of
evaluating air treatment products under conditions that closely mimic their intended real-world
use. Across diverse microbial groups—including Gram-positive and Gram-negative bacteria,
mycobacteria, viruses, and fungi—the aerosolized treatment of airborne microbes with TEG
delivered by Grignard Pure^{TM} consistently achieved higher and more rapid reductions in viable
organism concentration than surface treatment by the same product in liquid phase (Figures 2
and 3), where reductions were slower and less pronounced. Treatment of microorganisms on
surfaces by TEG vapor delivered by Grignard Pure™ was even less effective (Figure 4). These
findings are corroborated by testing conducted by the EPA's Office of Research and
Development (EPA ORD), which similarly reported superior log_{10} reductions in airborne
pathogen concentration following aerosolized TEG application compared to surface treatment,
further validating the notion that the efficacy of antimicrobial products intended for air treatment
should be tested in airborne phase (14) and not on surface-bound microorganisms. By the same
token, the data suggest that TEG will not likely eliminate the need for surface sterilization, which
is important to minimize the transfer of fomite-mediated infections. However, the use of a
combination of TEG for air treatment and specific agents, e.g., isopropyl alcohol, for surface
treatment is worth exploring in a separate study for potential synergistic effects.

On the other hand, the antimicrobial products designed for surface inactivation might not be
effective in the airborne phase. A study conducted in 1963 examined the comparative
inactivation of a germicide-resistant strain of phage *S. cremoris* 144F by surface-active
compounds applied as aerosols. The study observed that surface-active compounds such as
quaternary ammonium compounds (QAC) (tested in airborne concentrations of 0.024 ppm and
0.097 ppm) and phosphoric acid wetting agent (PAWA) (tested in airborne concentrations of
0.024 ppm, 0.048 ppm and 0.097 ppm) were ineffective as aerosols, despite their virucidal
activity in aqueous solutions (29). Conversely, in a separate study, PAWA, QAC, hypochlorites,
and iodophors, which are commonly used as sanitizing agents in the dairy industry, were tested
for their virucidal efficacy in aqueous solutions against the same *S. cremoris* 144F. The PAWA
sanitizer at a concentration of 12.5 ppm, as well as QAC at a concentration of 50 ppm

completely inactivated the phage within 15 seconds (30). The inability of these compounds to act
as effective agents in aerosol form can be attributed to the different physical environments under
which they must exert their activity (29).

Thus, the data from the research described in this paper, as well as existing literature, strongly
support the need to test antimicrobial products in the environment they are designed to treat.
Otherwise, extrapolating surface disinfection data to predict airborne efficacy or extrapolating
airborne efficacy to predict efficacy on surfaces might lead to inaccurate, and even diametrically
opposite conclusions.

Another aspect to consider in developing and adopting testing protocols for airborne
disinfectants is the sequence of introducing the inactivating product vs. introducing test
microorganisms. The EPA ORD study showed that the sequence of product introduction —
whether MS2 bacteriophage was introduced into a TEG-saturated chamber or vice versa—
strongly influenced early kill kinetics. Introducing MS2 into a TEG-saturated chamber resulted
in up to an additional 1 $\log_{10}$ reduction within the first 15 minutes, compared to adding TEG to a
virus-laden chamber (27). However, data sets from both test scenarios demonstrated that
aerosolized TEG treatment delivered by Grignard Pure™ at concentrations of 1.2–1.5 mg/m³
achieved 2–3 $\log_{10}$ reductions in airborne MS2 within 30–90 minutes. In our view, if TEG is
eventually approved for use as an aerosolized antimicrobial product, it should be
prophylactically deployed, i.e., people who may be infectious will enter spaces that have an
effective preexisting level of TEG in the air. Based on the earlier mentioned research of ORD
(14), as well as our earlier study (5), prophylactic use would produce greater reductions in the
levels of airborne pathogens than releasing TEG only after infected individuals begin to sneeze
or cough. Thus, we submit efficacy trials should be designed to reflect the real-world use of TEG
and should release the inactivating product into a test chamber before introducing the test
microorganisms.

Thus, an understanding of how various factors might impact the observed product efficacy is
necessary for predicting performance in the real world and is important for developing useful and
reliable standardized test methods (27). Ultimately, the most accurate assessment of an
antimicrobial product's efficacy requires a test methodology that mirrors its real-world
application. For air treatment products, this means introducing both the test mixture and the test
organisms into the air during efficacy studies.

Another critical aspect of developing a proper testing protocol involves selecting appropriate test
microorganisms, especially if a product will be used against BSL-2 microorganisms or higher.
Testing against such microorganisms in the airborne phase represents substantial technical
obstacles, e.g., the need for at least BSL-3 testing environment, which cannot be surmounted by
smaller companies or commercial testing laboratories. Therefore, the use of valid BSL-1
surrogates in such testing is critically important so long as data could be extrapolated to more
pathogenic BSL-2 or even higher counterparts. When properly selected, BSL-1 surrogates
provide a scientifically valid and practical alternative for evaluating airborne antimicrobial
efficacy. Surrogates such as *K. aerogenes* and *P. fluorescens* have demonstrated reduction
patterns on hard surfaces that closely align with those of selected pathogenic BSL-2 counterparts
from the same genus (i.e., *K. pneumoniae* and *P. aeruginosa*), supporting their suitability for
controlled testing in the airborne phase. Similarly, *S. epidermidis* serves as an effective surrogate
for *S. aureus*. These correlations suggest that surrogate-based testing offers a framework for

predicting the efficacy of antimicrobial products in real-world airborne scenarios, while
maintaining safety during laboratory studies conducted at BSL-2 levels.

Additionally, this study uses BSL-1 Gram-positive and Gram-negative organisms as surrogates
for airborne pathogens, but these groups also reflect skin- and enteric-derived contaminants.
While aerosols may not address direct contact transmission, their use could help limit re-
colonization of vulnerable hosts with antibiotic-resistant organisms in clinical settings—a
potential application worth further study.

Although surrogate-based testing involves extrapolating performance from surrogate organisms
to regulatory-recommended pathogens (e.g., US EPA), it provides a more scientifically
defensible basis for assessing airborne efficacy than surface-only data. At the same time, we
recognize that more comparative testing between the suggested surrogates (and perhaps other
candidates) and their pathogenic counterparts might be needed, especially when developing and
validating novel testing protocols, or when selecting surrogate microorganisms for testing of
therapeutic agents. By validating air treatment products through aerosolized trials with
appropriate surrogates, this methodology predicts that efficacy claims are both scientifically
sound and reflective of real-world conditions. This approach addresses a critical limitation in
traditional testing protocols and provides a reliable foundation for evaluating the potential of
antimicrobial products to reduce airborne microbial contamination effectively. Insisting on
airborne testing with microorganisms that cannot be studied in a BSL-2 facility would have the
consequence of discouraging, if not preventing, the development of aerosolized antimicrobial
products that could be effective tools for addressing diseases caused by transmission of airborne
pathogens.

596 *Broad-Spectrum Efficacy of TEG-based product (Grignard Pure^{TM})*

Our testing protocol aimed to evaluate the antimicrobial agent's broad-spectrum potential under
realistic conditions, including the use of Gram-negative and Gram-positive bacteria, non-
enveloped and enveloped bacteriophages, mycobacteria, and mold. We have shown that TEG can
achieve 2 to 4 orders of magnitude reductions in viable microbial counts of airborne bacteria,
viruses, mold, and mycobacteria at total TEG concentrations (aerosol + vapor) ranging from 0.7
to 2.0 mg/m^3 which are well within established safety margins – as discussed below in the
"Safety of TEG" section. We also show that this reduction can occur in as fast as 35 minutes,
with some organisms, like both tested bacteriophages, inactivated by almost two logs in 15
minutes. Two-log (99%) inactivation in 15 minutes or three-log (99.9%) inactivation in 35
minutes is equivalent to 18.4 and 11.8 air changes per hour (ACH), respectively (ACH is
calculated using a first order kinetics equation $C(t) = C(0) \exp(-ACH \times t)$, where $C(0)$ is the
initial microbial concentration, $C(t)$ is the microbial concentration at time t in hours). This is
substantially faster than airborne microorganism removal by ventilation systems operating at 5
ACH, a current CDC recommendation (https://www.cdc.gov/niosh/ventilation/prevention/Aim-
for-5.html). Sultan et al. recently further confirmed that aerosolized TEG delivers consistent
antimicrobial activity against MS2 bacteriophage across a broad range of real-world HVAC
conditions—including temperatures of 22–25 °C, relative humidities of 40–70 %, air
recirculation rates from 0 to 6 ACH, outdoor ventilation from 0.8 to 5 ACH, and filtration
efficiencies ranging from no filter to MERV 14—demonstrating that aerosolized TEG maintains
high inactivation rates even under dynamic ventilation and filtration scenarios (15). Notably,

Sultan et al.'s study used lower total (aerosol + vapor) TEG concentration of 0.44 mg/m^3, which
could explain why some of the inactivation values in their study were lower than those observed
in this study, where somewhat higher TEG concentrations were used.

The data presented by our study also suggest that Gram positive bacteria, both tested
bacteriophages, and mycobacteria are highly susceptible to airborne treatment by TEG, while
Gram-negative bacteria such as *S. typhimurium* and *P. fluorescens* are susceptible but not to the
same extent. Fungal spores of *A. brasiliensis* have proven to be the hardiest, but even they were
inactivated with ~95% efficiency after 35 minutes of exposure.

These results emphasize the importance of testing antimicrobial air treatment products against
diverse microorganisms to ensure comprehensive assessments of their real-world performance in
the airborne phase. Notably, among the organisms tested, viruses and mycobacteria—those most
strongly associated with clinically significant airborne transmission—showed substantial
susceptibility to TEG, highlighting its potential impact where it is most needed. The data also
corroborate the efficacy and utility of a TEG-based reagent (Grignard Pure^{TM}) to decontaminate
an indoor air environment and should provide a high degree of protection against airborne
transmission of a number of different pathogens.

Mechanism of Action

To understand the reason for the higher efficacy of TEG in air treatment compared to surface
treatment, we hypothesize that when certain compounds are atomized and vaporized, they not
only vaporize but also create microdroplets with a high surface-to-volume ratio that supports the
efficient evaporation of active molecules to maintain vapor saturation. The exact mechanism
behind TEG's higher efficacy in air treatment remains under study. Studies suggest that TEG
vapor, due to its high diffusion coefficient, may readily contact airborne microbes. As Puck et al.
demonstrated (3), TEG vapor can condense onto microdroplets carrying microorganisms,
achieving lethal local concentrations more rapidly than liquid-phase treatments. These
observations point toward a vapor- and condensation-driven mode of action. Its hygroscopic
nature may further contribute to microbial inactivation upon condensation (8). In addition to
membrane permeabilization, proposed in earlier literature, TEG may additionally act by altering
water activity, protein hydration and structure, as suggested for inactivation of non-enveloped
viruses (13). Further studies are warranted to confirm the multiple mechanisms of action of TEG
against different classes of microbes.

Public Health Significance

For an airborne antimicrobial to be effective, it must directly contact pathogens in their most
relevant form—suspended in air as vapor or microdroplets. Unlike surface-based methods, which
can artificially shield microbes and misrepresent product efficacy, aerosolized testing captures
the true dynamics of airborne transmission. Our findings demonstrate that TEG performs
optimally under these conditions, inactivating a broad spectrum of pathogens within minutes, and
at concentrations well below the ACGIH (American Conference of Governmental Industrial
Hygienists) safety threshold of 10 mg/m³ (https://www.acgih.org/triethylene-glycol/).

Safety of triethylene glycol has been extensively studied and a comprehensive discussion on the
safety of TEG is presented in Desai et al. (5). TEG exhibits very low toxicity through oral,

dermal, and inhalation exposure routes, with no toxicological endpoints of concern identified (31). As described by Desai et al. (5), the U.S. EPA classifies TEG as a "Safer Chemical" and has exempted it from tolerance requirements for antimicrobial residues on food-contact surfaces. Long-term inhalation studies in animals have shown no systemic adverse effects at concentrations significantly higher than those used in Grignard Pure™. A Margin of Safety analysis conducted by Nelson Labs further concluded that continuous inhalation exposure to TEG from product use presents a low risk of adverse effects, with no expected concerns related to acute, subchronic, or chronic toxicity, genotoxicity, or carcinogenicity. Moreover, TEG has been safely utilized in lighting effects (https://entertainingsafety.com/knowledge-base/understanding_fog_effects/) and air fresheners (https://edu.rsc.org/news/air-fresheners/2020753.article) for decades without significant health issues.

These results carry important implications for public health. During the COVID-19 pandemic, public health responses largely relied on personal behaviors—masking, distancing, and vaccination—despite inconsistent scientific consensus and variable compliance. This approach, though necessary, proved insufficient in stopping airborne spread and contributed to erosion of public trust. According to the hierarchy of controls, as explained by the CDC (https://www.cdc.gov/niosh/hierarchy-of-controls/about/index.html), the most effective strategy is to eliminate or inactivate the hazard itself. Our data demonstrates that TEG rapidly inactivates non-pathogenic airborne surrogates, and this is consistent with historical studies showing that TEG vapors inactivate pathogens such as *Staphylococcus aureus*(8) and influenza viruses (7), indicating its broader relevance for airborne infection control. Further, based on the inactivation of a broad range of BSL-1 surrogates in the air and results from testing surface-bound BSL-2 organisms, we posit that TEG as delivered by Grignard Pure™ will also be efficient against airborne BSL-2 organisms.

Study limitations

Our findings demonstrate the substantial antimicrobial efficacy of triethylene glycol (TEG) against a diverse array of airborne and surface-bound microorganisms; however, several limitations of the study should be acknowledged. Although all organisms, except for MS2, were only tested once, the inactivation kinetics were consistent across organisms, reaching net inactivation of 2-3 logs. Thus, similar results obtained with multiple organisms comprise a "de facto" aggregate replication, demonstrating the efficacy of TEG to inactivate a variety of airborne pathogens. Future work would benefit from repeated trials to establish confidence intervals around log-reduction values and to better observe differences in organism susceptibility. Higher initial concentrations of airborne microorganisms would also allow for more robust determination of time-dependent inactivation kinetics.

Additionally, all studies were conducted within a controlled range of temperature and relative humidity (RH). Both temperature and RH can affect aerosol dynamics, droplet evaporation rates, and microbial viability, which in turn can influence the germicidal activity of vapor-phase agents such as TEG. Previous research has shown that TEG vapor exhibits optimal antimicrobial efficacy at mid-range RH (approximately 50–60%) and moderate temperatures (~22 °C), with reduced performance observed at higher or lower RH or elevated temperatures (15). Moreover, environmental fluctuations in temperature and RH — such as those encountered in real-world indoor settings—might alter aerosol persistence and microbial susceptibility (32, 33). Therefore,

subsequent studies should systematically vary temperature and RH to better assess the germicidal efficacy of TEG against various microorganisms.

Lastly, another limitation of this study is the absence of a representative soil load in the aerosolization medium. In real-world settings, airborne pathogens are often embedded in organic and non-organic matter or biological fluids, which can reduce the effectiveness of antimicrobial agents (34). Future studies should incorporate soil loads, such as proteins or mucins, to better simulate real-world bioaerosols and evaluate TEG's performance under more realistic conditions.

Data availability

Data presented in this manuscript are available upon request.

ACKNOWLEDGMENTS

The authors acknowledge with gratitude helpful discussions and advice from Toni K Choueiri, William Esposito, Andre Fay, Antony Galione, Altaf Lal, and Donald W Schaffner.

Some of the research described in this article was funded by Bleu Garde, LLC, the company that makes Grignard Pure™.

CONFLICT OF INTEREST STATEMENT

Mr. Etienne Grignard is founder and Managing Director of Bleu Garde, LLC, which makes the Grignard Pure™ product investigated in this study. E.G., J.C., G.R., and G.M. are unpaid scientific advisors to the Bleu Garde company. They, and W.J., have also been granted an equity interest in potential future earnings from the Bleu Garde company.

References

1. Airborne transmission of respiratory viruses | Science. https://www.science.org/doi/10.1126/science.abd9149. Retrieved 5 August 2025.

2. Bourouiba L. 2021. Fluid Dynamics of Respiratory Infectious Diseases. Annu Rev Biomed Eng 23:547–577.

3. Organization WH. 2024. Global technical consultation report on proposed terminology for pathogens that transmit through the air. World Health Organization.

4. Reid JP, Ferro AR, Finn A, Lawler JV, Lednicky JA, Löndahl J, Fraenkel C-J, Santarpia JL, Ratnesar-Shumate SA, Wu C-Y. 2024. World Health Organization report removes the aerosol/droplet dichotomy but does not move us forward in infection control strategies. Aerosol Sci Technol 58:1089–1092.

5. Singanayagam A, Hakki S, Dunning J, Madon KJ, Crone MA, Koycheva A, Derqui-Fernandez N, Barnett JL, Whitfield MG, Varro R, Charlett A, Kundu R, Fenn J, Cutajar J, Quinn V, Conibear E, Barclay W, Freemont PS, Taylor GP, Ahmad S, Zambon M, Ferguson NM, Lalvani A, ATACCC Study Investigators. 2022. Community transmission and viral load kinetics of the SARS-CoV-2 delta (B.1.617.2) variant in vaccinated and unvaccinated individuals in the UK: a prospective, longitudinal, cohort study. Lancet Infect Dis 22:183–195.

6. Glas I, David SC. 2023. A new chapter of healthy indoor air: antiviral air treatments. EMBO Mol Med 15:e18710.

7.   Robertson OH, Puck TT, Lemon, H F, Loosli C G. 1943. The Lethal Effect of Triethylene

Glycol Vapor on Air-Borne Bacteria and Influenza Virus. Science 97.

8.   Puck TT. 1947. The Mechanism of Aerial Disinfection by Glycols and Other Chemical

Agents: II. An Analysis of the factors governing the efficiency of Chemical Disinfection of

the air. J Exp Med 85:741–757.

9.   Kaiser. 2014. Provisional Peer-Reviewed Toxicity Values for Triethylene Glycol (CASRN

112-27-6) 53.

10.   Rosebury T, Meiklejohn G, Kingsland LC, Boldt MH. 1947. Disinfection of Clouds of

Meningopneumonitis and Psittacosis Viruses with Triethylene Glycol Vapor. J Exp Med

85:65–76.

11.   Lester W, Kaye S, Robertson OH, Dunklin EW. 1950. Factors of Importance in the Use of

Triethylene Glycol Vapor for Aerial Disinfection. Am J Public Health Nations Health

40:813–820.

12.   Desai G, Ramachandran G, Goldman E, Esposito W, Galione A, Lal A, Choueiri TK, Fay

A, Jordan W, Schaffner DW, Caravanos J, Grignard E, Mainelis G. 2023. Efficacy of

Grignard Pure to Inactivate Airborne Phage MS2, a Common SARS-CoV-2 Surrogate.

Environ Sci Technol 57:4231–4240.

13.   Styles CT, Zhou J, Flight KE, Brown JC, Lewis C, Wang X, Vanden Oever M, Peacock TP,

Wang Z, Millns R, O'Neill JS, Borodavka A, Grove J, Barclay WS, Tregoning JS, Edgar

RS. 2023. Propylene glycol inactivates respiratory viruses and prevents airborne

transmission. EMBO Mol Med 15:e17932.

14.   Duggan K, Ijaz MK, McKinney J, Maillard J-Y. 2024. Reviewing the evidence of

antimicrobial activity of glycols. J Appl Microbiol 135:lxae071.

15.   Sultan Z, Luhung I, Aung NW, Uchida A, Natarajan A, Puramadathil S, Li J, Schuster S,

Schiavon S. 2024. Effectiveness of triethylene glycol disinfection on airborne MS2

bacteriophage under diverse building operational parameters. Indoor Environ 1:100042.

16.   Biosafety in Microbiological and Biomedical Laboratories—6th Edition.

17.   2004. Laboratory biosafety manual3rd ed. World Health Organization, Geneva.

18.   ARE Labs. 2021. 16m3 Bioaerosol Chamber Homogeneity Validation Report.

19.   ASHRAE Standard 241-2023 - Control of Infectious Aerosols.

https://webstore.ansi.org/standards/ashrae/ashraestandard2412023?gad_source=1&gad_cam

paignid=1042027569&gbraid=0AAAAAD_gXFUVFOGrqC79AulYLWs3az-

Gx&gclid=EAIaIQobChMIzveshsL0jgMVjk7_AR0PTDMaEAAYASAAEgI_bfD_BwE.

Retrieved 5 August 2025.

20.   MB-26-03 Neutralization of Microbicidal Activity using the Quantitative Method for

Evaluating Bactericidal and Mycobactericidal Activity of Microbicides Used on Hard, Non-

Porous Surfaces.

21.   E35 Committee. Test Method for Efficacy of Sanitizers Recommended for Inanimate, Hard,

Nonporous Non-Food Contact Surfaces. ASTM International.

22.   E35 Committee. Test Method to Assess Virucidal Activity of Chemicals Intended for

Disinfection of Inanimate, Nonporous Environmental Surfaces. ASTM International.

23.   Standard Operating Procedure for AOAC Use Dilution Method for Testing Disinfectants.

24.   2013. Guidance Document On Quantitative Methods For Evaluating The Activity Of

785   Microbicides Used On Hard Non-Porous Surfaces. OECD.

786   https://www.oecd.org/en/publications/guidance-document-on-quantitative-methods-for-

787   evaluating-the-activity-of-microbicides-used-on-hard-non-porous-surfaces_cb91d642-

788   en.html. Retrieved 27 May 2025.

789   25.   Prussin AJ, Schwake DO, Lin K, Gallagher DL, Buttling L, Marr LC. 2018. Survival of the

790   Enveloped Virus Phi6 in Droplets as a Function of Relative Humidity, Absolute Humidity,

791   and Temperature. Appl Environ Microbiol 84:e00551-18.

792   26.   Scales BS, Dickson RP, LiPuma JJ, Huffnagle GB. 2014. Microbiology, Genomics, and

793   Clinical Significance of the Pseudomonas fluorescens Species Complex, an Unappreciated

794   Colonizer of Humans. Clin Microbiol Rev 27:927–948.

795   27.   Ratliff KM, Oudejans L, Archer J, Calfee W, Gilberry JU, Hook DA, Schoppman WE,

796   Yaga RW, Brooks L, Ryan S. 2023. Impact of test methodology on the efficacy of

797   triethylene glycol (Grignard Pure) against bacteriophage MS2. Aerosol Sci Technol J Am

798   Assoc Aerosol Res 57:1178–1185.

799   28.   ANSI/ASHRAE Standard 185.3-2024 - Method of Testing Commercial and Industrial In-

800   Room Air-Cleaning Devices and Systems for Microorganism Bioaerosol Removal or

801   Inactivation in a Test Chamber.

802   https://webstore.ansi.org/standards/ashrae/ansiashraestandard1852024?gad_source=1&gad_

803   campaignid=1042027569&gbraid=0AAAAAD_gXFU8XflqK9MwTCO8FbDQPWy3L&g

clid=EAIaIQobChMIttu658b5jgMVjVxHAR0c1giDEAAYASAAEgLzW_D_BwE.

Retrieved 7 August 2025.

15. Sing EL, Elliker PR, Sandine WE. 1964. Comparative Destruction of Air-borne Lactic

Bacteriophages by Various Germicides Applied as Aerosols. J Food Prot 27:129–134.

30. Sing EL, Elliker PR, Sandine WE. 1964. Comparative Destruction of Lactic Streptococcus

Bacteriophages on Various Surfaces by Germicidal Aerosols. J Food Prot 27:161–165.

18. Sandars, F. T. Reregistration Eligibility Decision (RED) forTriethylene Glycol. 2005,

https://www3.epa.gov/pesticides/chem_search/reg_actions/reregistration/red_PC-

083501_26-Sep-03.pdf

32. Yang W, Marr LC. 2012. Mechanisms by Which Ambient Humidity May Affect Viruses in

Aerosols. Appl Environ Microbiol 78:6781–6788.

33. Lin K, Marr LC. 2020. Humidity-Dependent Decay of Viruses, but Not Bacteria, in

Aerosols and Droplets Follows Disinfection Kinetics. Environ Sci Technol 54:1024–1032.

34. Noda J, Tomizawa S, Takahashi K, Morimoto K, Mitarai S. 2022. Air pollution and

airborne infection with mycobacterial bioaerosols: a potential attribution of soot. Int J

Environ Sci Technol 19:717–726.

FIGURE LEGENDS

Figure 1

Timeline of experiments investigating the efficacy of triethylene glycol (TEG) against microbes: a) inactivation of airborne microbes and b) inactivation of surface-bound microbes. See text for full description.

Figure 2

Inactivation of Gram-positive and Gram-negative bacteria by triethylene glycol (TEG). Net log and percent reduction in airborne and surface-bound concentrations of viable bacteria due to treatment by triethylene glycol (TEG). Airborne bacteria were treated by aerosolized TEG, while surface-bound bacteria were treated by liquid TEG. Treatment time for airborne bacteria also denotes the start of 10-min sampling by impingers. See Materials and Methods for experimental details. Error bars were propagated using a partial derivative method.

Figure 3

Inactivation of bacteriophages, mycobacteria, and fungal spores by triethylene glycol (TEG). Net log and percent reduction in airborne and surface-borne concentrations of viable bacteriophages, rnycobacteria and fungal spores due to treatment by triethylene glycol (TEG). Airborne microbes were treated by aerosolized TEG, while surface-bound microbes were treated by liquid TEG. Treatment time for airborne microbes also denotes the start of 10-min sampling by impingers. Error bars were propagated using a partial derivative method.

Figure 4

Comparison of inactivation by TEG of bacterial pathogens and their non-pathogenic surrogates. Net log inactivation of surface-bound bacteria by TEG vapor as delivered by Grignard Pure™. The data are mean values of four replicates (carriers) and error bars represent one standard deviation.

Figure 1. Timeline of experiments investigating the efficacy of triethylene glycol (TEG) against microbes: a) inactivation of airborne microbes and b) inactivation of surface-bound microbes.

Inactivation of Gram-positive and Gram-negative bacteria by triethylene glycol (TEG)

Figure 2. Net log and percent reduction in airborne and surface-bound concentrations of viable bacteria due to treatment by triethylene glycol (TEG). Airborne bacteria were treated by aerosolized TEG, while surface-bound bacteria were treated by liquid TEG. Treatment time for airborne bacteria also denotes the start of 10-min sampling by impingers. The propagated error for each data point is provided in Supplemental Materials.

Inactivation of bacteriophages, mycobacteria, and fungal spores by triethylene glycol (TEG)

Figure 3. Net log and percent reduction in airborne and surface-borne concentrations of viable bacteriophages, mycobacteria and fungal spores
due to treatment by triethylene glycol (TEG). Airborne microbes were treated by aerosolized TEG, while surface-bound microbes were treated by liquid TEG.
Treatment time for airborne microbes also denotes the start of 10-min sampling by impingers.The propagated error for each data point is provided in Supplemental Materials.

Fig. 4. Net log inactivation of surface-bound bacteria by TEG vapor as delivered by Grignard Pure^{TM}. The data are mean values of four replicates (carriers) and error bars represent one standard deviation.