ZYS-1 is not an ADAR1 inhibitor

Cassandra N. Smoak[1], Estelle N. Gardner[1], Renee N. Chua[1], Kyle A. Cottrell [1,2]

[1]Department of Biochemistry, Purdue University, West Lafayette, IN, USA

[2]Purdue Institute for Cancer Research, Purdue University, West Lafayette, IN, USA

The authors declare no potential conflicts of interest.

<u>Correspondence:</u>

Kyle A Cottrell, Ph.D.
Department of Biochemistry
Purdue University
201 S University St.
West Lafayette, IN
Email: kacottre@pudue.edu
Telephone: 765-494-6941
16

Shortened title: ZYS-1 is not an ADAR1 inhibitor

Keywords: ADAR1, cancer, RNA

Abstract

Adenosine deaminase acting on RNA 1 (ADAR1) edits double-stranded RNA (dsRNA)

substrates by the deamination of adenosine to inosine in a process known as A-to-I editing.

Modulation of ADAR1 expression and editing activity has previously been described to play a

role in cancer development and progression, with upregulation of ADAR1 being observed in a

range of cancers. Further, depletion of ADAR1 leads to increased sensing of endogenous

dsRNAs by dsRNA sensors in cell lines that require ADAR1 for survival, which are termed

ADAR1-dependent. The activation of these sensors induces downstream production of type I

interferons as well as translational inhibition and apoptosis. Therefore, ADAR1 is a promising

oncologic therapeutic target. Recently, the small molecule ZYS-1 has been developed and

presented as a direct inhibitor of ADAR1. We performed a series of in vitro and cellular

experiments to validate the efficacy and specificity of ZYS-1 as an ADAR1 inhibitor. Evaluating

the effect of ZYS-1 on cell viability revealed it to be equally cytotoxic to both ADAR1-dependent

and ADAR1-independent cell lines, as well as wildtype and ADAR1 knockout cells. Moreover,

ZYS-1 treatment had little effect on activation of PKR or induction of IFN stimulated genes.

Importantly, treatment with ZYS-1 did not reduce cellular A-to-I editing for several known

ADAR1 editing sites, and did not inhibit in vitro A-to-I editing by recombinant ADAR1. Together,

these data indicate that ZYS-1 is not a selective inhibitor of ADAR1.

Introduction

The RNA editing enzyme ADAR1 is an important regulator of double-stranded RNA (dsRNA) sensing pathways (Cottrell et al. 2024a; Heraud-Farlow and Walkley 2020). Adenosine-to-inosine (A-to-I) editing by ADAR1 prevents activation of MDA5 by endogenous dsRNAs (Liddicoat et al. 2015; Pestal et al. 2015; Mannion et al. 2014). Additionally, ADAR1 suppresses activation of the dsRNA sensor PKR in an RNA editing-independent manner (Sinigaglia et al. 2024; Chung et al. 2018; Hu et al. 2023). These functions of ADAR1 are essential in embryonic development and, importantly, in many cancer cell lines (Wang et al. 2004; Hartner et al. 2004; Gannon et al. 2018; Liu et al. 2019; Kung et al. 2021). Depletion of ADAR1 in roughly half of cancer cells causes cell death (Liu et al. 2019; Gannon et al. 2018; Kung et al. 2021). In these ADAR1-dependent cells, depletion of ADAR1 causes activation of PKR – which drives cell death – and activation of MDA5 leading to type I interferon signaling and expression of IFN-stimulated genes (ISGs). The IFN response caused by ADAR1 knockout can overcome resistance to immune checkpoint blockade (Ishizuka et al. 2019). These phenotypes associated with ADAR1 depletion make ADAR1 an important therapeutic target for many cancers, especially those with poor response to immune checkpoint blockade. However, there are currently no FDA-approved ADAR1 inhibitors.

The substrate of ADAR1 is an adenosine residue within duplexed RNA (Bass and Weintraub 1988; Wagner and Nishikura 1988). While the modified adenosine can either be base-paired or not (Wong et al. 2001), ADAR1 has no activity on single-stranded RNA (Bass and Weintraub 1988; Wagner and Nishikura 1988), or on adenosine or adenine alone (Polson et al. 1991). Previously, two small molecule adenosine analogues have been reported to disrupt ADAR1 function – 8-azaadenosine (Ramírez-Moya et al. 2020; Zipeto et al. 2016) and 8-chloroadenosine (Ding et al. 2020) – however, through extensive evaluation, we found that neither are selective inhibitors of ADAR1 (Cottrell et al. 2021). Recently, ZYS-1, another adenosine analogue (Figure 1a), has been reported to inhibit ADAR1(Wang et al. 2025) and has

been used in subsequent studies (Shi et al. 2025). Here we carefully evaluate the efficacy and

specificity of ZYS-1 as an ADAR1 inhibitor. Through a series of cellular and in vitro experiments,

we found no evidence supporting ZYS-1 as an ADAR1 inhibitor.

Results

ZYS-1 is broadly cytotoxic

A selective inhibitor of ADAR1 would be expected to be more cytotoxic towards ADAR1-

dependent cell lines than ADAR1-independent lines. As such, we evaluated the effect of ZYS-1

treatment on cell viability in a panel of ADAR1-dependent and ADAR1-independent cell lines.

We chose two dependent cell lines (HCC1806 and MDA-MB-468) and two independent cell

lines (BT-549 and MDA-MB-453) based on either DepMap data (Broad 2024) (Fig. 1b) or our

previous studies (Kung et al. 2021). Irrespective of the ADAR1-dependency status, ZYS-1 was

strongly cytotoxic towards all cell lines evaluated (Fig. 1c). As measured by an ATP-dependent

luciferase-based assay for cell viability, we found the EC50 of ZYS-1 to vary little across the cell

lines in our panel and to be lowest in the ADAR1-independent cell line BT-549. Crystal violet

staining of cells treated with ZYS-1 supported the robust toxicity of ZYS-1 (Figure 1d-e).

To more rigorously evaluate the specificity of ZYS-1, we assessed its effect on cell

viability in WT and ADAR1 knockout (ADAR1-KO) BT-549 (Fig. 1f). The EC50 for ZYS-1 in WT

and ADAR1-KO BT-549 cells was nearly identical (0.14 and 0.12 µM, respectively (Fig. 1g)).

Importantly, the WT and ADAR1-KO BT-549 were equally viable in the absence of ZYS-1

(Supplemental Figure 1a). We attempted to assess the effect of ZYS-1 by crystal violet staining,

but found that ADAR1-KO BT-549 cells detached easily during the staining process

(Supplemental Fig. S1b). Together with the results above, these data strongly indicate that ZYS-

1 does not cause cell death by affecting ADAR1.

ZYS-1 does not phenocopy ADAR1 depletion

Depletion of ADAR1 in ADAR1-dependent cells produces a viral mimicry phenotype that

includes activation of PKR (indicated by autophosphorylation on Thr446 (Romano et al. 1998))

and MDA5, with the latter driving a type I IFN response (Liu et al. 2019; Gannon et al. 2018).

Both of these phenotypes were observed upon ZYS-1 treatment in the original study reporting

ZYS-1 as an ADAR1 inhibitor (Wang et al. 2025). We evaluated the effect of ZYS-1 on PKR

activation and ISG induction using two ADAR1-dependent cell lines that we have previously

studied (Cottrell et al. 2021; Kung et al. 2021). ZYS-1 treatment of the ADAR1-dependent cell

lines HCC1806 and MDA-MB-468 had no effect on PKR activation (Fig. 2a-b). Previously ZYS-1

has been shown to reduce ADAR1 expression (Wang et al. 2025), however, we observed no

change in ADAR1 expression in ZYS-1-treated HCC1806 and MDA-MB-468 (Fig. 2a and Fig.

2c). Increased expression of ISGs is a phenotype of ADAR1 depletion in ADAR1-dependent

cancer cell lines (Cottrell et al. 2021; Liu et al. 2019). We evaluated the expression of four ISGs

in ZYS-1-treated HCC1806 and MDA-MB-468. ZYS-1 did not affect ISG expression in HCC1806

and caused elevated expression of only one ISG in MDA-MB-468 (Fig. 2d). Together with our

evaluation of PKR activation, these data strongly indicate that ZYS-1 does not phenocopy

ADAR1 depletion.

ZYS-1 does not inhibit A-to-I editing

A-to-I editing by ADAR1 can be easily evaluated by sequencing of cDNA. Since inosine

more readily base-pairs with cytidine, cDNA made from an edited RNA will have a guanosine at

the editing site (for the sense strand of duplex cDNA or a cytidine on the anti-sense strand).

Sanger sequencing is routinely used to evaluate the level of editing at a given site by comparing

the peak heights of A or G (T or C on the antisense stand). We used this approach to evaluate

the effect of ZYS-1 treatment on A-to-I editing in cells and in vitro using purified ADAR1 and a

well-characterized substrate RNA. For editing in cells, we evaluated editing at three previously

characterized editing sites. Editing at all three sites was greatly reduced by knockdown of

ADAR1 in HCC1806 (Fig. 3a-b, Supplemental Figure S2). In ZYS-1-treated cells, editing was

either not affected or increased for all but one site (Fig. 3c-h). For an editing site in the gene

ZDHHC20, we observed significantly reduced editing at the highest concentration of ZYS-1

used in BT-549, but not in the other cell lines evaluated. It should be noted that this reduced

editing occurred with 1 µM ZYS-1 treatment – a dose that is an order of magnitude higher than

the EC50 for cell viability in this cell line.

Since it is known that other proteins can regulate A-to-I editing at specific sites (Hong et

al. 2018; Freund et al. 2020), it is possible that some changes in editing observed in cells upon

ZYS-1 treatment could be the result of off-target effects. To evaluate direct inhibition of ADAR1,

we performed an in vitro A-to-I editing assay with purified ADAR1 and an *in vitro* transcribed

RNA containing a portion of the mRNA coding for 5-hydroxytryptamine receptor 2C (5-HT$_{2C}$,

encoded by *HTR2C*). The 5-HT$_{2C}$ mRNA contains a well-characterized ADAR1 editing site,

known as the 'B-site', and is frequently used in A-to-I editing assays (Burns et al. 1997). The 5-

HT$_{2C}$ B-site was well edited by ADAR1 in DMSO controls (Fig. 3i-j), though we observed

differences in editing based on the lot of ADAR1 used (see Supplemental Fig. S3 for additional

data from a different enzyme lot and additional assay controls). Addition of ZYS-1 to the editing

assay had no effect on A-to-I editing at any dose used (Fig. 3i-j and Supplemental Fig. S3a-b).

Together with the cellular A-to-I editing data above, these data indicate that ZYS-1 is not an

ADAR1 inhibitor.

Discussion

Many studies have now shown the therapeutic potential of targeting ADAR1 to treat a

range of cancers (Liu et al. 2019; Ishizuka et al. 2019; Gannon et al. 2018; Kung et al. 2021). In

addition to the cell-intrinsic effects of disrupting ADAR1, the cell-extrinsic effects, such as

overcoming resistance to immune checkpoint blockade, strengthen the potential of therapies

targeting ADAR1. As such, identifying small molecule inhibitors of ADAR1 is of great

importance. Previously reported ADAR1 inhibitors, the adenosine analogues 8-azaadenosine

and 8-chloroadenosine, have not proven to be effective in subsequent studies (Cottrell et al.

2021). Unfortunately, our data above indicate that the adenosine analogue ZYS-1 is not a

selective inhibitor of ADAR1.

Smoak et al., 2025   6

While we show that ZYS-1 effectively kills many breast cancer cell lines, it does so not

by inhibiting ADAR1. Our data are very different from those reported in the initial publication

describing ZYS-1 as an ADAR1 inhibitor (Wang et al. 2025). There are a few possible

explanations for the differences between these studies. First, the original paper reporting ZYS-1

as an ADAR1 inhibitor, Wang et al., 2025, used a commercial assay for Adenosine Deaminase

(ADA) to assess inhibition of A-to-I editing by ADAR1 (Wang et al. 2025). That assay is not

designed to assess ADAR1 activity and has been reported to be incapable of detecting A-to-I

editing (Zhang et al. 2025). Here we used a conventional in vitro A-to-I editing assay to evaluate

inhibition of ADAR1 by ZYS-1 and observed no inhibition, consistent with the findings of a

recent preprint (Zhang et al. 2025). Second, while Wang et al., 2025 provided evidence that

ZYS-1 inhibited A-to-I editing within cells, they evaluated only one editing site in two cell lines.

We evaluated three editing sites across four cell lines and observed a significant editing

reduction at only one site, in one cell line. Most sites had no observable change, and we even

found some sites to have increased editing. Because A-to-I editing by ADAR1 can be regulated

by other proteins (Freund et al. 2020; Hong et al. 2018), it is possible that a small molecule

could indirectly affect A-to-I editing, which could explain both increased or decreased editing at

specific sites. Third, in Wang et al., 2025, the authors evaluated the effect of ZYS-1 on cell

viability in control and ADAR1 knockout cell lines. They found ZYS-1 to be less effective in the

ADAR1 knockout cells. However, in that experiment, knockout of ADAR1 greatly reduced cell

viability, which could have reduced the efficacy of ZYS-1 (which we suspect is inhibiting DNA

synthesis, see below). In contrast, we performed a similar experiment with control versus

ADAR1 knockout cells, using an ADAR1-independent cell line to ensure that knockout did not

affect cell viability. In our experiment, ZYS-1 efficacy was not affected by ADAR1 knockout.

While ZYS-1 is not a selective ADAR1 inhibitor, it is highly cytotoxic to cancer cells.

ZYS-1 is a derivative of fludarabine. Fludarabine has a long history as a chemotherapeutic used

for multiple leukemias and lymphomas (Gandhi and Plunkett 2002). Its mechanism of action

involves inhibition of DNA synthesis, through both inhibition of DNA polymerase and

ribonucleotide reductase (Gandhi and Plunkett 2002). We suspect that ZYS-1 is functioning

through the same mechanism. This would explain its cytotoxicity.

Finally, it is our opinion that adenosine analogues are unlikely to inhibit ADAR1 (or

ADAR2). Early studies demonstrated that ADAR1 is highly selective for dsRNA(Bass and

Weintraub 1988; Wagner and Nishikura 1988; Polson et al. 1991). Unlike ADA, ADAR1 cannot

be inhibited by coformycin or 8-azanebularine – both of which structurally resemble the hydrated

reaction intermediate of the adenosine deamination reaction (Polson et al. 1991; Mendoza et al.

2023). However, when incorporated into a duplexed RNA, but not single-stranded RNA, 8-

azanebularine potently inhibits ADAR1 with a low nanomolar IC50 (Mendoza et al. 2023). These

findings suggest that additional interactions between ADAR1 and the substrate dsRNA are

required to adjust the conformation of the active site and allow binding to adenosine – a model

that was first suggested over three decades ago (Polson et al. 1991). This model is supported

by structural studies of ADAR1 and ADAR2 indicating multiple interactions between active site

adjacent residues and either the phosphodiester backbone of the dsRNA substrate or the

'orphan' base opposite the edited adenosine (Fisher and Beal 2024; Matthews et al. 2016; Deng

et al. 2025). For ADAR2, the interactions between the deaminase domain and the dsRNA

substrate cause long-distance changes to the deaminase domain structure (Matthews et al.

2016). These prior observations, along with our evaluation of ZYS-1 and other adenosine

analogues (Cottrell et al. 2021), all indicate that selective inhibition of ADAR1 is unlikely to be

accomplished via an adenosine analogue.

Materials and Methods

Cell Culture and Lentiviral Transduction

The cell lines (HEK293T (RRID: CVCL_0063), MDA-MB-453 (RRID: CVCL _0418),

MDA-MB-468 (RRID: CVCL_0063), BT-549 (RRID: CVCL_1092), and HCC1806 (RRID:

CVCL_1258)) were purchased from the American Type Culture Collection and cultured as

described previously (Young et al. 2025). Detailed methods are provided in the Supplemental

Information. For some experiments we used the above cell lines engineered for inducible

expression of Cas9. These iCas9 cell lines were described previously by our laboratory (Young

et al. 2025). The iCas9 lines were used for ADAR1 knockout (Fig. 1e-f, BT-549 iCas9) and for

evaluating ZYS-1 efficacy in the cell lines MDA-MB-453 and MDA-MB-468. ZYS-1 (Fludarabine-

Cl) was purchased from MedChemExpress (Cat# HY-145443).

Lentiviral production and transduction was performed as described previously (Young et

al. 2025). Detailed methods are provided in the Supplemental Information.

Plasmids

Oligos encoding an ADAR1-targeting sgRNA (Table S1) were cloned into lenti-sgRNA-

hygro (a gift from Brett Stringer (Stringer et al. 2019), Addgene #104991;

RRID:Addgene_104991) by ligating annealed and phosphorylated oligos into a digested and

dephosphorylated vector. The lenti-sgC1-hygro, generated previously (Young et al. 2025),

expresses a control sgRNA that targets a control genomic locus (AAVS1) and has been used

previously (Zou et al. 2024; Young et al. 2025). The ADAR1 sgRNA used is from the Avana

library used in DepMap (Broad 2024).

The control and ADAR1 knockdown plasmids (tet-pLKO-shSCR-puro-luc and tet-pLKO-

shADAR1-puro-luc) were generated by cloning of oligonucleotides encoding either shSCR or

shADAR1 into the plasmid tet-pLKO-puro-luc by ligating annealed and phosphorylated oligos

into a digested and dephosphorylated vector, see Supplemental Information for oligonucleotide

sequences. To generate tet-pLKO-puro-luc, first a PstI site was inserted at the end of PuroR in

tet-pLKO-puro (a gift from Dmitri Wiederschain (Wiederschain et al. 2009), Addgene plasmid #

21915, RRID:Addgene_21915) by site directed mutagenesis using PrimeSTAR Max DNA

Polymerase (Takara Bio) and In-Fusion Snap Assembly (Takara). A region of the plasmid

pRG01-U6-DR-crRNA-BsmbI(x2)-6T; EFS-Puro-2A-Fluc-WPRE (a gift from Sidi Chen (Chow et

al. 2019), Addgene plasmid # 123362, RRID:Addgene_123362) encoding PuroR-2A-Fluc was

PCR amplified and digested with BsiWI and PstI prior to ligation into the tet-pLKO-puro-PstI

digested with the same enzymes. See Supplemental Table 1 for all primer sequences. Both the

control and ADAR1 targeting shRNAs used here have been used previously to evaluate ADAR1

function in breast cancer (Kung et al. 2025; Kung et al. 2021; Young et al. 2025; Cottrell et al.

2024b; Cottrell et al. 2021). The sequences for both shRNAs are listed in Table S1.

All plasmids were confirmed by Sanger sequencing and Nanopore whole plasmid

sequencing as well as restriction enzyme digestion.

Genetic Depletion by CRISPR-Cas9 or RNAi

To generate ADAR1 knockout in BT-549, BT-549 iCas9 cell lines were transduced with

lentivirus made with lenti-sgADAR1-hygro or lenti-sgC1-hygro and selected with hygromycin.

The cells were treated with 2 µg/mL doxycycline to induce Cas9 expression for at least seven

days prior to maintenance in media without doxycycline for at least three days before

experimentation. To establish the BT-549 ADAR1 knockout cell line, we screened clonal lines

for ADAR1 knockout by immunoblot. We used the polyclonal sgC1 line as our control.

For knockdown of ADAR1, HCC1806 cells were transduced with either tet-pLKO-

shSCR-puro-luc or tet-pLKO-shADAR1-puro-luc and selected with puromycin. Knockdown of

ADAR1 was initiated by doxycycline treatment at 2 µg/mL.

Immunoblot

Immunoblot analysis was performed as previously described, with between 25 and 40

micrograms of protein loaded per lane (Young et al. 2025). Detailed methods are described in

the Supplemental Information. For ZYS-1-treated cells, cells were treated with DMSO or ZYS-1

for 48 hours prior to harvesting protein. Primary antibodies used include: ADAR1 (Santa Cruz

Biotechnology Cat# sc-73408, RRID:AB_2222767), PKR (Cell Signaling Technology Cat# 3072,

RRID:AB_2277600), PKR Thr-446-P (Abcam Cat# ab32036, RRID:AB_777310), GAPDH

(Santa Cruz Biotechnology Cat# sc-47724, RRID:AB_627678), and beta-tubulin (Abcam Cat#

ab6046, RRID:AB_2210370).

Cell Viability and Proliferation Assessment

To assess cell viability, 2,500 (HCC1806 and BT-549) or 5,000 (MDA-MB-453 and MDA-MB-468) cells were plated in triplicate per condition in 96-well plates. The cells were treated as indicated one day after plating. CellTiter-Glo 2.0 (Promega) was used to determine cell viability per manufacturer's protocol three days after treatment.

To assess cell proliferation and viability by crystal violet staining, 25,000 (HCC1806 and BT-549) or 50,000 (MDA-MB-453 and MDA-MB-468) cells were plated per well of a 6-well dish. The cells were treated as indicated one day following plating. Cells were washed with 1x PBS and fixed in 100% methanol after seven to fourteen days of growth, depending on the cell line. The cells were then stained with 0.005% Crystal Violet solution containing 25% methanol (Sigma-Aldrich), prior to washing with water and imaging using a ChemiDoc Imaging System (Bio-Rad). Percent stained area was measured using Fiji (ImageJ).

Cellular A-to-I Editing Analysis

Cells were treated with DMSO or ZYS-1 for 48 hours prior to RNA harvesting and purification using the Nucleospin RNA kit (Macherey-Nagel). cDNA was made from isolated RNA using LunaScript Supermix (NEB). Sections around the A-to-I editing sites in ZDHHC20, BPNT1, and MRPS16 were amplified using PrimeSTAR Max DNA Polymerase (Takara Bio) and the primers listed in Table S1. PCR products were then resolved by agarose gel electrophoresis and purified by the Monarch Gel Extraction kit (New England Biolabs). Purified products were Sanger sequenced by Azenta with the primers listed in Table S1. The QSVAnalyzer program was used to measure the peak heights of each A-to-I editing site. Percent editing was found by dividing the edited base (G or C) peak height by the sum of the edited and unedited (A or T) peak heights and multiplying by 100. The primer used to sequence the regions around the ZDHHC20 and MRPS16 editing sites yielded the antisense sequence of this region; percent editing for these sites was determined using C (edited) and T (unedited) peak heights.

In Vitro A-to-I Editing Assay

A plasmid encoding a portion of the 5-HT$_{2C}$ mRNA was synthesized by Twist Bioscience.

This pTwist-5HT plasmid was linearized by KpnI restriction enzyme digest and purified by

agarose gel electrophoresis as described above. The 5-HT$_{2C}$ RNA was in vitro transcribed using

HiScribe T7 Quick High Yield RNA (New England Biolabs). The in vitro transcribed RNA was

purified using the Monarch Spin RNA Cleanup kit (New England Biolabs) and the quality of the

RNA was assessed by denaturing polyacrylamide gel electrophoresis. The RNA was annealed

in template annealing buffer (10 mM Tris-HCl pH 7.5, 1 mM EDTA, and 100 mM NaCl) by

heating to 95 °C and cooling slowly to room temperature.

The in vitro A-to-I editing assay was modified from those used by the laboratory of Peter

Beal, University of California Davis (Mendoza et al. 2023). In vitro editing assays were carried

out in 10 µL reactions consisting of 1X ADAR1 reaction buffer (15 mM Tris pH 7.4, 26 mM KCl,

1.5 mM EDTA, 0.003% NP-40, 4% glycerol and 40 mM potassium glutamate), 0.5 mM DTT, 1

µg/mL yeast tRNA (Invitrogen), 0.16 U/µL RNase Inhibitor Murine (New England Biolabs), 4 nM

5-HT$_{2C}$ RNA, 5 µM RT_primer (Table S1), 1 mM dNTPs and water to 8 µL. To this reaction mix,

DMSO or ZYS-1 dissolved in DMSO was added prior to initiating the reaction by addition of 200

nM ADAR1-p150 (BPS Bioscience). The reaction was incubated at 30 °C for 15 minutes for

ADAR1-p150 lot #231010-1 or 1 hr for lot #250218. The reaction was stopped by heating to 95

°C for 5 minutes. Reverse transcription followed immediately by the addition of ProtoScript II

Buffer to 1X final (New England Biolabs), DTT to 10.25 mM final, 200 U ProtoScript II Reverse

Transcriptase (New England Biolabs), 8 U RNase Inhibitor Murine and water to 20 µL total. The

samples were incubated at 25 °C for 5 minutes, 42 °C for 1 hour and 65 °C for 5 minutes. The

cDNA was cleaned using the Monarch PCR and DNA Cleanup kit. PrimeSTAR Max DNA

Polymerase was used to PCR amplify the cDNA using primers EA_F and EA_R, described in

Table S1. The PCR products were purified by agarose gel electrophoresis and Sanger

sequenced using the EA_F primer. A-to-I editing was quantified as described above.

Quantitative PCR

Total RNA isolation and cDNA synthesis was performed as above. Quantitative PCR

(qPCR) was performed with Luna Universal qPCR MasterMix (NEB) on a QuantStudio3 system

(Thermo Scientific). The primers used are listed in Table S1. For each primer, the amplification

efficiency was verified to be within 90-110% allowing 'Fold Change' determination via the ΔΔCt.

The geometric mean Ct of EEF1A1 and HSPA5 was used for normalization.

Data Availability Statement

All analysis scripts are available at (https://github.com/cottrellka/Smoak_et_al_2025).

Dependency data (DepMap_Public_24Q4+Score, Chronos, and Achilles+DRIVE+Marcotte,

DEMETER2) were acquired from the DepMap portal

(https://depmap.org/portal/download/custom/, RRID:SCR_017655).

Acknowledgements:

This work was supported by R00MD016946 (K.A. Cottrell), the Purdue Institute for

Cancer Research (start-up funding) NIH grant P30CA023168, and start-up funding from Purdue

University Department of Biochemistry and College of Agriculture.

Author Contributions

K.A.C. and C.N.S. conceived the project. K.A.C. and C.N.S. designed the experiments.

C.N.S., E.N.G., R.N.C., and K.A.C. performed the experiments. C.N.S. and K.A.C. performed

the data analysis. K.A.C. and C.N.S. wrote the manuscript. All authors edited the manuscript.

Declaration of interests

The authors declare no competing interests.

References

Bass BL, Weintraub H (1988) An unwinding activity that covalently modifies its double-stranded
RNA substrate. Cell 55 (6):1089-1098. doi:10.1016/0092-8674(88)90253-x
Broad D (2024) DepMap 24Q4 Public. doi:10.25452/figshare.plus.27993248.v1
Burns CM, Chu H, Rueter SM, Hutchinson LK, Canton H, Sanders-Bush E, Emeson RB (1997)
Regulation of serotonin-2C receptor G-protein coupling by RNA editing. Nature 387
(6630):303-308. doi:10.1038/387303a0
Chow RD, Wang G, Ye L, Codina A, Kim HR, Shen L, Dong MB, Errami Y, Chen S (2019) In
vivo profiling of metastatic double knockouts through CRISPR-Cpf1 screens. Nat
Methods 16 (5):405-408. doi:10.1038/s41592-019-0371-5
Chung H, Calis JJA, Wu X, Sun T, Yu Y, Sarbanes SL, Dao Thi VL, Shilvock AR, Hoffmann HH,
Rosenberg BR, Rice CM (2018) Human ADAR1 Prevents Endogenous RNA from
Triggering Translational Shutdown. Cell 172 (4):811-824.e814.
doi:10.1016/j.cell.2017.12.038
Cottrell KA, Andrews RJ, Bass BL (2024a) The competitive landscape of the dsRNA world. Mol
Cell 84 (1):107-119. doi:10.1016/j.molcel.2023.11.033
Cottrell KA, Ryu S, Pierce JR, Soto Torres L, Bohlin HE, Schab AM, Weber JD (2024b)
Induction of Viral Mimicry Upon Loss of DHX9 and ADAR1 in Breast Cancer Cells.
Cancer Res Commun 4 (4):986-1003. doi:10.1158/2767-9764.CRC-23-0488
Cottrell KA, Torres LS, Dizon MG, Weber JD (2021) 8-azaadenosine and 8-chloroadenosine are
not selective inhibitors of ADAR. Cancer Res Commun 1 (2):56-64. doi:10.1158/2767-
9764.crc-21-0027
Deng X, Sun L, Zhang M, Basavaraj R, Wang J, Weng YL, Gao Y (2025) Biochemical profiling
and structural basis of ADAR1-mediated RNA editing. Mol Cell 85 (7):1381-1394.e1386.
doi:10.1016/j.molcel.2025.02.017
Ding HY, Yang WY, Zhang LH, Li L, Xie F, Li HY, Chen XY, Tu Z, Li Y, Chen Y, Yang SY (2020)
8-Chloro-Adenosine Inhibits Proliferation of MDA-MB-231 and SK-BR-3 Breast Cancer
Cells by Regulating ADAR1/p53 Signaling Pathway. Cell Transplant
29:963689720958656. doi:10.1177/0963689720958656
Fisher AJ, Beal PA (2024) Structural perspectives on adenosine to inosine RNA editing by
ADARs. Mol Ther Nucleic Acids 35 (3):102284. doi:10.1016/j.omtn.2024.102284
Freund EC, Sapiro AL, Li Q, Linder S, Moresco JJ, Yates JR, Li JB (2020) Unbiased
Identification of trans Regulators of ADAR and A-to-I RNA Editing. Cell Rep 31
(7):107656. doi:10.1016/j.celrep.2020.107656
Gandhi V, Plunkett W (2002) Cellular and clinical pharmacology of fludarabine. Clin
Pharmacokinet 41 (2):93-103. doi:10.2165/00003088-200241020-00002
Gannon HS, Zou T, Kiessling MK, Gao GF, Cai D, Choi PS, Ivan AP, Buchumenski I, Berger
AC, Goldstein JT, Cherniack AD, Vazquez F, Tsherniak A, Levanon EY, Hahn WC,
Meyerson M (2018) Identification of ADAR1 adenosine deaminase dependency in a
subset of cancer cells. Nat Commun 9 (1):5450. doi:10.1038/s41467-018-07824-4
Hartner JC, Schmittwolf C, Kispert A, Müller AM, Higuchi M, Seeburg PH (2004) Liver
Disintegration in the Mouse Embryo Caused by Deficiency in the RNA-editing Enzyme
ADAR1*. Journal of Biological Chemistry 279 (6):4894-4902.
doi:https://doi.org/10.1074/jbc.M311347200
Heraud-Farlow JE, Walkley CR (2020) What do editors do? Understanding the physiological
functions of A-to-I RNA editing by adenosine deaminase acting on RNAs. Open Biol 10
(7):200085. doi:10.1098/rsob.200085
Hong H, An O, Chan THM, Ng VHE, Kwok HS, Lin JS, Qi L, Han J, Tay DJT, Tang SJ, Yang H,
Song Y, Bellido Molias F, Tenen DG, Chen L (2018) Bidirectional regulation of

adenosine-to-inosine (A-to-I) RNA editing by DEAH box helicase 9 (DHX9) in cancer. Nucleic Acids Res 46 (15):7953-7969. doi:10.1093/nar/gky396

Hu SB, Heraud-Farlow J, Sun T, Liang Z, Goradia A, Taylor S, Walkley CR, Li JB (2023) ADAR1p150 prevents MDA5 and PKR activation via distinct mechanisms to avert fatal autoinflammation. Mol Cell 83 (21):3869-3884.e3867. doi:10.1016/j.molcel.2023.09.018

Ihlenfeldt WD, Bolton EE, Bryant SH (2009) The PubChem chemical structure sketcher. J Cheminform 1 (1):20. doi:10.1186/1758-2946-1-20

Ishizuka JJ, Manguso RT, Cheruiyot CK, Bi K, Panda A, Iracheta-Vellve A, Miller BC, Du PP, Yates KB, Dubrot J, Buchumenski I, Comstock DE, Brown FD, Ayer A, Kohnle IC, Pope HW, Zimmer MD, Sen DR, Lane-Reticker SK, Robitschek EJ, Griffin GK, Collins NB, Long AH, Doench JG, Kozono D, Levanon EY, Haining WN (2019) Loss of ADAR1 in tumours overcomes resistance to immune checkpoint blockade. Nature 565 (7737):43-48. doi:10.1038/s41586-018-0768-9

Kung C-P, Terzich ND, Ilagen MXG, Prinsen MJ, Kaushal M, Kladney RD, Weber JH, Mabry AR, Torres LS, Bramel ER, Freeman EC, Sabloak T, Cottrell KA, Ryu S, Weber WM, Maggi L, Shriver LP, Patti GJ, Weber JD (2025) ADAR1 Regulates Lipid Remodeling through MDM2 to Dictate Ferroptosis Sensitivity. bioRxiv:2025.2001.2016.633410. doi:10.1101/2025.01.16.633410

Kung CP, Cottrell KA, Ryu S, Bramel ER, Kladney RD, Bao EA, Freeman EC, Sabloak T, Maggi L, Weber JD (2021) Evaluating the therapeutic potential of ADAR1 inhibition for triple-negative breast cancer. Oncogene 40 (1):189-202. doi:10.1038/s41388-020-01515-5

Liddicoat BJ, Piskol R, Chalk AM, Ramaswami G, Higuchi M, Hartner JC, Li JB, Seeburg PH, Walkley CR (2015) RNA editing by ADAR1 prevents MDA5 sensing of endogenous dsRNA as nonself. Science 349 (6252):1115-1120. doi:10.1126/science.aac7049

Liu H, Golji J, Brodeur LK, Chung FS, Chen JT, deBeaumont RS, Bullock CP, Jones MD, Kerr G, Li L, Rakiec DP, Schlabach MR, Sovath S, Growney JD, Pagliarini RA, Ruddy DA, MacIsaac KD, Korn JM, McDonald ER (2019) Tumor-derived IFN triggers chronic pathway agonism and sensitivity to ADAR loss. Nat Med 25 (1):95-102. doi:10.1038/s41591-018-0302-5

Mannion Niamh M, Greenwood SM, Young R, Cox S, Brindle J, Read D, Nellåker C, Vesely C, Ponting Chris P, McLaughlin Paul J, Jantsch Michael F, Dorin J, Adams Ian R, Scadden ADJ, Öhman M, Keegan Liam P, O'Connell Mary A (2014) The RNA-Editing Enzyme ADAR1 Controls Innate Immune Responses to RNA. Cell Reports 9 (4):1482-1494. doi:https://doi.org/10.1016/j.celrep.2014.10.041

Matthews MM, Thomas JM, Zheng Y, Tran K, Phelps KJ, Scott AI, Havel J, Fisher AJ, Beal PA (2016) Structures of human ADAR2 bound to dsRNA reveal base-flipping mechanism and basis for site selectivity. Nat Struct Mol Biol 23 (5):426-433. doi:10.1038/nsmb.3203

Mendoza HG, Matos VJ, Park S, Pham KM, Beal PA (2023) Selective Inhibition of ADAR1 Using 8-Azanebularine-Modified RNA Duplexes. Biochemistry 62 (8):1376-1387. doi:10.1021/acs.biochem.2c00686

Pestal K, Funk CC, Snyder JM, Price ND, Treuting PM, Stetson DB (2015) Isoforms of RNA-Editing Enzyme ADAR1 Independently Control Nucleic Acid Sensor MDA5-Driven Autoimmunity and Multi-organ Development. Immunity 43 (5):933-944. doi:10.1016/j.immuni.2015.11.001

Polson AG, Crain PF, Pomerantz SC, McCloskey JA, Bass BL (1991) The mechanism of adenosine to inosine conversion by the double-stranded RNA unwinding/modifying activity: a high-performance liquid chromatography-mass spectrometry analysis. Biochemistry 30 (49):11507-11514. doi:10.1021/bi00113a004

Ramírez-Moya J, Baker AR, Slack FJ, Santisteban P (2020) ADAR1-mediated RNA editing is a novel oncogenic process in thyroid cancer and regulates miR-200 activity. Oncogene 39 (18):3738-3753. doi:10.1038/s41388-020-1248-x

Romano PR, Garcia-Barrio MT, Zhang X, Wang Q, Taylor DR, Zhang F, Herring C, Mathews MB, Qin J, Hinnebusch AG (1998) Autophosphorylation in the activation loop is required for full kinase activity in vivo of human and yeast eukaryotic initiation factor 2alpha kinases PKR and GCN2. Mol Cell Biol 18 (4):2282-2297. doi:10.1128/MCB.18.4.2282

Shi Z, Li J, Ding J, Zhang Y, Min W, Zhu Y, Hou Y, Yuan K, Sun C, Wang X, Shen H, Wang L, Liang SQ, Kuang W, Yang P (2025) ADAR1 is required for acute myeloid leukemia cell survival by modulating post-transcriptional Wnt signaling through impairing miRNA biogenesis. Leukemia 39 (3):599-613. doi:10.1038/s41375-024-02500-7

Sinigaglia K, Cherian A, Du Q, Lacovich V, Vukić D, Melicherová J, Linhartova P, Zerad L, Stejskal S, Malik R, Prochazka J, Bondurand N, Sedláček R, O'Connell MA, Keegan LP (2024) An ADAR1 dsRBD3-PKR kinase domain interaction on dsRNA inhibits PKR activation. Cell Rep 43 (8):114618. doi:10.1016/j.celrep.2024.114618

Stringer BW, Day BW, D'Souza RCJ, Jamieson PR, Ensbey KS, Bruce ZC, Lim YC, Goasdoué K, Offenhäuser C, Akgül S, Allan S, Robertson T, Lucas P, Tollesson G, Campbell S, Winter C, Do H, Dobrovic A, Inglis PL, Jeffree RL, Johns TG, Boyd AW (2019) A reference collection of patient-derived cell line and xenograft models of proneural, classical and mesenchymal glioblastoma. Sci Rep 9 (1):4902. doi:10.1038/s41598-019-41277-z

Wagner RW, Nishikura K (1988) Cell cycle expression of RNA duplex unwindase activity in mammalian cells. Mol Cell Biol 8 (2):770-777. doi:10.1128/mcb.8.2.770-777.1988

Wang Q, Miyakoda M, Yang W, Khillan J, Stachura DL, Weiss MJ, Nishikura K (2004) Stress-induced apoptosis associated with null mutation of ADAR1 RNA editing deaminase gene. J Biol Chem 279 (6):4952-4961. doi:10.1074/jbc.M310162200

Wang X, Li J, Zhu Y, Shen H, Ding J, Zeng T, Min W, Liang SQ, Huang L, Shi Z, Huang F, Yuan K, Kuang W, Ji M, Sun C, Hou Y, Wang L, Chen W, Jiang Y, Hao H, Xiao Y, Yang P (2025) Targeting ADAR1 with a small molecule for the treatment of prostate cancer. Nat Cancer 6 (3):474-492. doi:10.1038/s43018-025-00907-4

Wiederschain D, Wee S, Chen L, Loo A, Yang G, Huang A, Chen Y, Caponigro G, Yao YM, Lengauer C, Sellers WR, Benson JD (2009) Single-vector inducible lentiviral RNAi system for oncology target validation. Cell Cycle 8 (3):498-504. doi:10.4161/cc.8.3.7701

Wong SK, Sato S, Lazinski DW (2001) Substrate recognition by ADAR1 and ADAR2. RNA 7 (6):846-858. doi:10.1017/s135583820101007x

Young AA, Juhler IG, Pierce JR, Bohlin HE, Harper HA, Onishile DS, Chua RN, Liu ME, Gardner EN, Elzey BD, Cottrell KA (2025) PACT suppresses PKR activation through dsRNA binding and dimerization, and is a therapeutic target for triple-negative breast cancer. bioRxiv:2025.2005.2027.656288. doi:10.1101/2025.05.27.656288

Zhang M, Deng X, Gao Y, Wang J (2025) Re: Concerns Regarding the Validation of ZYS-1 as a Bona Fide ADAR1 Inhibitor. bioRxiv:2025.2003.2007.641892. doi:10.1101/2025.03.07.641892

Zipeto MA, Court AC, Sadarangani A, Delos Santos NP, Balaian L, Chun HJ, Pineda G, Morris SR, Mason CN, Geron I, Barrett C, Goff DJ, Wall R, Pellecchia M, Minden M, Frazer KA, Marra MA, Crews LA, Jiang Q, Jamieson CHM (2016) ADAR1 Activation Drives Leukemia Stem Cell Self-Renewal by Impairing Let-7 Biogenesis. Cell Stem Cell 19 (2):177-191. doi:10.1016/j.stem.2016.05.004

Zou T, Zhou M, Gupta A, Zhuang P, Fishbein AR, Wei HY, Capcha-Rodriguez D, Zhang Z, Cherniack AD, Meyerson M (2024) XRN1 deletion induces PKR-dependent cell lethality in interferon-activated cancer cells. Cell Rep 43 (2):113600. doi:10.1016/j.celrep.2023.113600

Figure 1: ZYS-1 is broadly cytotoxic to breast cancer cells, irrespective of ADAR1 expression or ADAR1-dependency.

a Structures of adenosine, ZYS-1, and fludarabine, made using PubChem Sketcher V2.4 (Ihlenfeldt et al. 2009). **b** CHRONOS and DEMETER2 ADAR1 dependency scores for relevant TNBC cell lines as evaluated using CRISPR-Cas9 knockout screening or RNAi screening, respectively (Broad 2024). A CHRONOS or DEMETER2 score below –0.5 indicates gene dependency (Broad 2024). **c** Dose-response evaluation of cell viability for ZYS-1 in cell lines given in **b**, as measured using the CellTiter-Glo 2.0 assay. Representative images (**d**) and quantification (**e**) of crystal violet staining following ZYS-1 treatment of cell lines given in **b**. **f** Immunoblot analysis of ADAR1 protein expression in WT and ADAR1-KO BT-549 cells. **g** Dose-response evaluation of cell viability for ZYS-1 in WT and ADAR1-KO BT-549 cells, as measured using the CellTiter-Glo 2.0 assay. For **c** and **g**, large points are the average of three biological replicates, and smaller points are the average of three technical replicates performed for each biological replicate. For **e**, bars indicate the average of three biological replicates. Error bars represent the average ± SD. Statistical analysis for **e** performed using one-way ANOVA with post-hoc Tukey. ** $p < 0.01$, *** $p < 0.001$.

491

Figure 2: Treatment with ZYS-1 does not induce PKR activation or ISG expression.

a Immunoblot analysis of PKR activation (P-PKR) in the ADAR1-dependent cell lines HCC1806

and MDA-MB-468 following treatment with ZYS-1. **b** and **c**, Quantification of the immunoblot

shown in **a**. **d** qPCR analysis of IFN-stimulated gene expression in ZYS-1-treated HCC1806 and

MDA-MB-468 cells. Bars indicate the average of three biological replicates. Error bars represent

the average ± SD.

Figure 3

Figure 3: ZYS-1 treatment does not inhibit A-to-I editing by cellular or recombinant ADAR1.

a Sanger sequencing trace of BPNT1, MRPS16, and ZDHHC20 editing sites in HCC1806 control or ADAR1 knockdown cells. **b** Quantification of percent editing using Sanger sequencing results depicted in **a**. Representative Sanger sequencing trace of BPNT1 (**c**), MRPS16 (**e**), and ZDHHC20 (**g**) editing sites in HCC1806 following treatment with ZYS-1. Percent editing quantification of the BPNT1 (**d**), MRPS16 (**f**), and ZDHHC20 (**h**) transcripts using Sanger sequencing in relevant TNBC cell lines. **i** Sanger sequencing trace of the 5-HT$_{2C}$ B-site following incubation with purified ADAR1 and ZYS-1 as indicated. **j** Quantification of B-site percent editing using Sanger sequencing results depicted in **i**. For **a**, **c**, **e**, **g**, and **i**, sites that are edited by ADAR1 are indicated by the red vertical line. Bars indicate the average of three biological replicates (**b**, **d**, **f**, and **h**) or two experimental replicates (**j**). Error bars represent the average ± SD. Statistical analysis performed using Student's t-test for **b** and one-way ANOVA with post-hoc Tukey for **d**, **f**, and **h**. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$.