MORC2 Mediates Transcriptional Regulation Through

Liquid-Liquid Phase Separation and DNA Binding

Yanshen Zhang[1,2,3,#], Weiya Xu[1,2,#], Wenxiu Duan[1,2,#], Yu Wei[1,2,3], Wenli Jiang[1,2,3], Feng Zhu[1], Chengdong Huang[1,2,*], Chao Wang[1,2,3*], and Yihui Bi[4,*],

[1]Hefei National Research Center for Physical Sciences at the Microscale, Center for Advanced Interdisciplinary Science and Biomedicine of IHM, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[2]Ministry of Education Key Laboratory for Membraneless Organelles and Cellular Dynamics, Biomedical Sciences and Health Laboratory of Anhui Province, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[3]Department of Neurology, the First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[4]Department of Orthopedics, The Second Affiliated Hospital of Anhui Medical University, Hefei, China.

Running title: Multivalent interactions and DNA binding underlies phase separation of MORC2

[#]These authors contributed equally to this work.

[*]Corresponding authors: Chengdong Huang (huangcd@ustc.edu.cn);

Chao Wang (cwangust@ustc.edu.cn);

Yihui Bi (docbyh1992@163.com).

Abstract

MORC2 is a chromatin-associated ATPase implicated in transcriptional silencing and human neuropathies such as Charcot–Marie–Tooth disease and spinal muscular atrophy. However, the molecular mechanisms governing its transcriptional regulatory activity remain elusive. Here, we demonstrate that full-length MORC2 undergoes liquid-liquid phase separation (LLPS) to form nuclear condensates, a process essential for transcriptional repression. Endogenous MORC2 forms dynamic condensates in neurons from EGFP-MORC2 knock-in mice, supporting the physiological relevance of LLPS *in vivo*. The 3.1 Å crystal structure of coiled-coil 3 (CC3) reveals a dimeric scaffold that drives phase separation, while multivalent interactions between the intrinsically disordered region (IDR) and a newly defined IDR-binding domain (IBD) further promote condensate formation. Moreover, LLPS enhances MORC2's ATPase activity in a DNA-dependent manner, indicating a functional coupling between phase separation, DNA binding, and enzymatic regulation. RNA-seq analysis shows that only wild-type, but not LLPS-deficient MORC2, represses core target genes in knockout cells, directly linking condensate formation to transcriptional control. Furthermore, disease-associated MORC2 variants alter condensate dynamics, ATPase activity, and DNA binding, offering mechanistic insights into their pathogenic effects. Together, these findings identify LLPS as a key regulatory mechanism for MORC2 function and provide a molecular framework for understanding its role in transcriptional regulation and human disease.

Introduction

Chromatin dynamically regulates DNA accessibility to orchestrate essential processes such as transcription, replication, and DNA repair[1–5]. These processes are dynamically regulated by chromatin-associated complexes, such as the human silencing hub (HUSH) complex[6], which control the accessibility of genetic material to regulatory machinery. The Microrchidia (MORC) protein family, characterized by a conserved N-terminal GHL-ATPase domain and a structurally diverse C-terminal region, regulates gene expression and chromatin organization[7–10]. These C-terminal domains are thought to contribute to dimerization and functional specificity[11–14] (Fig. 1a).

GHL ATPases, including DNA gyrase, Hsp90, and MutL, rely on ATP-dependent dimerization to drive conformational changes essential for substrate engagement and enzymatic regulation[15–19]. Similar mechanisms have been proposed for MORC family members. In humans, the MORC family comprises of five members—MORC1, MORC2, MORC3, MORC4, and SMCHD1, and different MORC paralogs target distinct genomic loci, including transposable elements, viral DNA, and long-exon genes[7,20,21]. The conserved N-terminal GHL-ATPase domain of MORC family proteins has been extensively characterized, implicating the evolutionarily diversified, multi-domain C-terminal regions as key determinants of chromatin targeting and functional specificity. Despite their structural similarities, the molecular basis for MORC-mediated repression remains poorly understood, particularly due to limited structural insight into its full-length and C-terminal regions.

MORC2 has recently been identified as a key regulator of chromatin architecture and gene silencing, primarily through its recruitment to HUSH-target loci via C-terminal CC2 domain-mediated interactions with TASOR and MPP8[12]. While its silencing activity was initially attributed to the N-terminal ATPase domain, genetic complementation studies have revealed that the CW domain and all three coiled-coil (CC) regions are also indispensable for its function[12,13]. Beyond its structured domains, sequence analyses and structural predictions have uncovered an extended intrinsically

disordered region (IDR) spanning the central to C-terminal regions of the protein,

suggesting potential regulatory mechanisms beyond enzymatic activity[22].

Transcriptional silencing is often spatially organized within membrane-less

nuclear compartments[23,24]. Recent studies suggest that intrinsically disordered regions

(IDRs) can drive liquid-liquid phase separation (LLPS), providing a plausible

mechanism by which chromatin-associated factors form functional condensates[25,26].

However, the molecular mechanisms driving LLPS in chromatin-associated proteins,

and how such phase behavior modulates genome architecture and gene expression,

remain poorly defined.

In this study, we provide direct and comprehensive evidence that MORC2

undergoes LLPS to form nuclear condensates. We define the structural basis for this

behavior, identifying a dimeric CC3 scaffold and multivalent IDR-IBD interactions as

core elements driving phase separation. We further demonstrate that DNA binding

promotes LLPS and that phase separation enhances MORC2's enzymatic activity.

Using both cellular and *in vivo* models, we establish that LLPS is functionally

essential for MORC2-mediated transcriptional repression, offering new mechanistic

insights into how LLPS integrates with DNA binding and enzymatic function to

regulate gene silencing.

Results

Coiled-coil 3 (CC3) mediates dimerization of full-length MORC2

To determine whether MORC2 has N- and C-terminal dimerization interfaces similar to those in other GHL-type ATPases[27], we established an *in vitro* system to purify full-length (FL) human MORC2 (MORC2FL, residues 1-1032) from HEK293F cells, yielding nucleic acid-free, highly pure proteins for structural and functional analyses (Fig. 1a, b). Analytical size-exclusion chromatography coupled with static light scattering (SLS) demonstrated that MORC2FL forms dimers, with a molecular weight of approximately 239 ± 1 kDa under buffer conditions containing 1000 mM NaCl even in the absence of ATP (Fig. 1c). Notably, the addition of 601 DNA resulted in a detectable increase in molecular weight (303 ± 2 kDa), indicative of DNA-protein complex formation, but did not disrupt the dimeric assembly of MORC2 (Fig. 1c).

To elucidate the atomic structure of MORC2FL, we initially employed negative-stain electron microscopy to reveal that MORC2FL particles were poorly resolved and exhibited considerable heterogeneity. Notably, a subset of particles displayed a morphology consistent with dimer formation (Fig. S1a). To further investigate this, we performed structural prediction of MORC2FL using AlphaFold3 and identified a predicted structure that closely resembled the observed reconstruction (Fig. S1b). We further examined the oligomerization properties of various MORC2 constructs. MORC2ΔCC3-IBD (residues 1-900) was found to exist as a monomer, whereas CC3-IBD (residues 901-1032) and CC3 alone (residues 901-1003) exhibited dimeric properties, confirming the critical role of the CC3 domain in mediating MORC2 dimerization (Fig. 1c, Fig. S1c).

To gain further structural insights, we determined the 3.1 Å crystal structure of the CC3 dimer using X-ray diffraction (Table S1). The structure revealed that the dimer interface is stabilized predominantly by hydrophobic interactions. Three key hydrophobic residues—L911, L915, and F922—form distinct layers within the dimer interface (layers 1-3, Fig. 1e, Fig. S1e, f). The side chains of these residues are buried

within the dimerization interface, creating a hydrophobic core critical for maintaining

the stability of the dimer.

We systematically introduced single-point mutations in these hydrophobic

residues to assess their contributions to dimerization. SLS analysis demonstrated that

the wild-type (WT) MORC2 CC3-IBD (residues 901-1032, with Trx-tag) fragment

predominantly exists as a dimer (~61 kDa). In contrast, all three key mutants (L911Q,

L915Q, and F922Q) displayed altered molecular weight distributions indicative of

disrupted dimer formation and enhanced higher-order oligomerization (Fig. S1g).

However, none of these mutations completely disrupted dimerization or converted the

dimer to a monomeric state (Fig. S1g). Further analyses of additional single-site

mutants (e.g., I908Q, Y921A, F951Q, Y954A, L958Q) showed minimal effects on

dimer stability, underscoring the central importance of L915 and its neighboring

hydrophobic layers in stabilizing the CC3 dimer (Fig. S1g). Moreover, we examined

the dimeric status of truncated CC3 constructs (residues 901-975), which confirmed

that the CC3 domain alone is sufficient to drive dimer formation (Fig. 1f). To

delineate the structural basis of these interactions, we mapped the positions of

hydrophobic residues within the CC3 backbone (Fig. S1e). The distances between

interacting residues, such as I908-I908' (3.4 Å), L911-L915' (3.7 Å), F922-F922' (3.9

Å), and L958-L958' (4.0 Å), revealed a tightly packed hydrophobic network. The

robustness of this network ensures the integrity of the CC3 dimer, even under varying

physiological conditions. These findings collectively suggest that the CC3 domain

plays a critical role in MORC2 dimerization. The dimeric assembly of CC3 is

essential for maintaining the structural integrity of the protein, which likely underpins

its functional activities, including interactions with chromatin.

MORC2 forms condensates both *in vivo* and *in vitro*

In the predicted structure of MORC2FL, we noticed a long unstructured region at

its C-terminus (Fig. S1b), a characteristic often associated with proteins capable of

phase separation[25]. To further investigate this, we utilized the IUPred2A tool to

predict disordered regions in MORC2FL. The analysis revealed that the previously

undefined 593-735 region predominantly exhibits disorder, prompting us to classify it

as an intrinsically disordered region (IDR) (Fig. 1a).

Next, we explored whether MORC2FL exhibits phase separation properties.

Lowering the salt concentration to physiological levels is a common strategy for

inducing LLPS[28]. Sedimentation assays showed that when the salt concentration was

reduced to 150 mM, more than half of the EGFP fused MORC2FL

(EGFP-MORC2FL) precipitated into the pellet fraction (Fig. 2a, b). Fluorescence

imaging revealed that at a salt concentration of 150 mM, EGFP-MORC2FL

spontaneously demixed to form well-defined spherical droplets of varying sizes,

which increased in size with rising protein concentration (Fig. 2c, d, and Fig. S2a).

Sedimentation assays further showed that the pellet fraction of EGFP-MORC2FL

after centrifugation was concentration-dependent (Fig. S2b). The

EGFP-MORC2FL-enriched condensates exhibited dynamic behavior, including

droplet fusion (Fig. S2c) and fluorescence recovery after photobleaching (FRAP) (Fig.

2e, f), indicating a liquid-like state.

Using antibody labeling, we observed that endogenous MORC2 forms dense

punctate structures in the nucleus of HeLa cells (Fig. 2g), whereas these structures are

absent in MORC2-knockout HeLa cells (Fig. S2d, e), suggesting that MORC2 may

undergo LLPS in the nucleus. To further investigate this, we transfected HeLa cells

with EGFP-MORC2FL and observed that EGFP-MORC2FL formed dense,

droplet-like condensates in the nucleus (Fig. 2h). Notably, in agreement with antibody

staining results, EGFP-MORC2FL condensates preferentially localize to regions of

reduced chromatin density, as indicated by histone H2A-mCherry signal. This spatial

distribution suggests a potential role for MORC2 condensates in heterochromatin

regulation. FRAP of EGFP-MORC2FL puncta revealed recovery on the timescale of

seconds in these transfected HeLa cells (Fig. 2i, j), consistent with characteristics of

liquid-like condensates with a highly dynamic internal environment.

To determine whether MORC2 forms phase-separated biomolecular condensates

under physiological conditions, we generated mice expressing EGFP-MORC2 protein

from the endogenous locus (Fig. S2f). We focus on neurons in brain and spinal cord as

missense mutations in *MORC2* cause neuropathies including Charcot-Marie-Tooth
and spinal muscular atrophy disease[11,13]. Imaging of NeuN-positive neurons revealed
that EGFP-MORC2FL condensates occur in neurons under endogenous expression
level (Fig. 2k). FRAP analysis in freshly prepared brain slices further demonstrated
rapid and partial fluorescence recovery, supporting the dynamic and liquid-like nature
of endogenous MORC2 assemblies (Fig. 2l). In addition, unlike MORC3, which
shows cell cycle-dependent nuclear puncta formation[29], MORC2 condensates were
absent in mitotic cells (Fig. S2g). Together, these results indicate that MORC2 forms
phase-separated biomolecular condensates under physiological conditions *in vivo*.
202

The MORC2 C-terminal domain (CTD) drives condensate formation

To identify the key determinants of MORC2 LLPS, we divided the protein into
two distinct domains: the N-terminal domain (NTD, residues 1-472), which includes
the ATPase and CC1 domains, and the C-terminal domain (CTD, residues 473-1032),
containing the IDR. A series of EGFP-tagged truncation constructs were expressed in
HeLa cells (Fig. 3a). Deletion of the CTD disrupted puncta formation and resulted in
diffuse localization across the nucleus and cytoplasm (Fig. 3a, Fig. S3a). By contrast,
NTD deletion had little effect on nuclear localization or condensate formation (Fig. 3a,
Fig. S3a). Addition of an exogenous nuclear localization signal (PKKKRKV)[30] to the
NTD redirected it to the nucleus but failed to restore puncta formation (Fig. 3a),
indicating that the CTD is necessary for condensate assembly.

Further dissection of the CTD revealed that deletion of either the CC2 or TCD
domains did not impair its ability to form nuclear puncta (Fig. S3b). In contrast,
removal of the IDR-NLS region (residues 593-761) disrupted both puncta formation
and nuclear localization (Fig. S3b). Notably, even when nuclear localization was
restored by fusing the PKKKRKV to CTDΔIDR-NLS, puncta formation remained
abolished (Fig. S3c), indicating that the IDR is essential for condensate formation.
Deletion of the CC3-IBD segment allowed nuclear localization but also abolished
puncta formation, suggesting that this region is likewise required (Fig. S3b). However,
neither the IDR-NLS fragment alone, the CC3-IBD fragment alone, nor the

PKKKRKV-fused CC3-IBD fragment was sufficient to form puncta (Fig. S3d),

implying a cooperative role between these two regions. The CW domain modulates

condensate size, as its deletion led to enlarged and often fused CTDΔCW puncta. In

the context of MORC2FL, we final conclude that the IDR, CC3, and IBD constitute

the structural core required for condensate formation, while the CW domain restricts

condensate overgrowth or fusion (Fig. 3a, Fig. S3e). Notably, condensates formed

exclusively in the nucleus and not in the cytoplasm of transfected HeLa cells,

suggesting that chromatin-associated nuclear factors, such as DNA, may contribute to

the nucleation or stabilization of MORC2 condensates.

To assess whether the CTD is intrinsically prone to LLPS *in vitro*, we expressed

and purified CTDΔCW from *E. coli*, as the CTD exhibited poor biochemical stability

(Fig. S4a, b). Under 150 mM NaCl conditions, Cy3-labeled CTDΔCW readily formed

condensates in a concentration-dependent manner (Fig. 3b, c), and underwent phase

separation-associated sedimentation (Fig. 3d, e). Fluorescence microscopy and static

light scattering (SLS) confirmed that Cy3 labeling did not affect its phase behavior,

droplet morphology, or molecular weight distribution (Fig. S4c–f).

Multivalent interactions between IDR and CC3-IBD promote MORC2 LLPS

Next, we elucidated the mechanistic basis for MORC2 phase separation

mediated by the IDR and residues CC3-IBD. We hypothesized that multivalent

interactions likely exist between the IDR and CC3-IBD, which are important for

driving LLPS, leading to a condensed phase coexisting with a dilute aqueous phase.

Using nuclear magnetic resonance (NMR) spectroscopy, we investigated the

interaction between the IDR and the CC3-IBD fragment (Fig. 4a-c). The ^{15}N-labeled

CC3 (residues 901-1003) or IBD (residues 1004-1032) samples were prepared for

^{15}N-HSQC assays. Titration experiments with the IDR (residues 593-735) revealed

that the IBD is the primary binding site, whereas the CC3 region showed minimal

interaction (Fig. 4b, c). Sequence analysis confirmed that IBD is enriched in acidic

residues and is highly conserved across species, which may facilitate its interaction

with the basic residues in the IDR (Fig. 4g). Further subdivision of the IDR into three

segments—IDRa (residues 593-643), IDRb (residues 644-694), and IDRc (residues

695-735)—pinpointed IDRa as the key region responsible for interaction with IBD

(Fig. 4a, d-f). NMR titration data indicate that IDRa binds IBD via transient weak

interactions (Fig. 4d, g). The partial compaction of IDRa, which we identify as a

proline-arginine-rich region, together with its weak multivalent association with IBD

(Fig. 4g), may serve as a nucleation mechanism for LLPS—an organizational strategy

that has also been observed in other phase-separating proteins[31–33].

To assess the role of IDRa-IBD multivalent interactions in promoting MORC2

LLPS within the nucleus, we transfected full-length MORC2 constructs with

individual deletions of the IBD, IDRa, IDRb, or IDRc regions. Deletion of the IBD

markedly reduced droplet formation in both HeLa cells (Fig. 3a, Fig. S3e) and

HEK293T cells (Fig. S5). Similarly, deletion of IDRa significantly reduced

condensate formation, resulting in a marked decrease in both droplet area and size

(Fig. 4h, i). In contrast, deletion of IDRb or IDRc had minimal impact on droplet

formation (Fig. 4h, i). Based on these findings, we propose a model in which

multivalent interactions between the IDRa and IBD drive MORC2 intermolecular

assembly, a process further potentiated by CC3-mediated dimerization, which is

essential for LLPS (Fig. 4j).

DNA binding promotes MORC2 phase separation and ATPase activity

Our results indicate that nuclear localization is critical for MORC2 condensate

formation when over-expressed in HeLa cells. As MORC2 has been reported to bind

DNA via its CC1 domain[13], we further assess the role of DNA binding in regulating

its LLPS behavior. To this end, we first purified MORC2FL and several truncated

variants, and assessed their ability to bind 601 DNA using electrophoretic mobility

shift assays (EMSA). Increasing concentrations of MORC2FL progressively retarded

601 DNA migration, confirming direct DNA binding (Fig. 5a). Consistent with

previous reports[13], CC1-containing constructs exhibited robust DNA-binding activity

(Fig. 5b). Notably, the CC2 and IDR domain also displayed strong DNA-binding

activity (Fig. 5c, d). In addition, while the CC3-IBD fragment showed no detectable

effect on 601 DNA migration (Fig. 5f), the TCD-CC3-IBD construct clearly impaired

migration (Fig. 5e), indicating that the TCD also possesses DNA-binding capability.

285   Building on these findings, we tested whether DNA enhances MORC2 LLPS *in*

vitro. At 0.9 μM, MORC2FL alone failed to form droplets; however, the addition of

25 nM 601 DNA induced robust condensate formation, indicating that DNA promotes

MORC2 phase separation (Fig. 5g, h). Moreover, 601 DNA also facilitated droplet

formation by the CTDΔCW construct (Fig. 5i). Together with the observation that the

TCD is dispensable for condensate formation and that the CC3-IBD fragment does

not bind DNA, these results suggest that DNA-induced condensation is primarily

mediated through interactions with the IDR-NLS region. We speculated that this

interaction is driven by nonspecific electrostatic attraction between DNA and the

arginine/lysine-rich sequence within the IDR-NLS (Fig. 4g).

295   We further examined the impact of DNA binding on MORC2 ATPase activity.

The presence of 601 DNA markedly enhanced ATP hydrolysis, indicating that DNA

binding positively modulates MORC2 enzymatic function (Fig. 5j). To confirm that

the observed activity is intrinsic to MORC2, we included the ATP-binding-deficient

mutant N39A as a negative control. N39A exhibited negligible ATP hydrolysis under

both basal and DNA-stimulated conditions, in contrast to the robust and

DNA-enhanced activity observed with the WT protein (Fig. 5j). Together, these

findings reveal a synergistic interplay between phase separation, DNA binding, and

ATPase activity, highlighting their collective importance in coordinating

transcriptional regulation.

LLPS is required for MORC2-mediated transcriptional repression

307   To evaluate whether phase separation contributes to MORC2-dependent

transcriptional regulation, we performed rescue experiments in MORC2-knockout

HeLa cells by expressing EGFP, EGFP-MORC2FL, EGFP-MORC2ΔCC3, or

EGFP-MORC2ΔIDR, respectively. Bulk RNA sequencing was used to assess

transcriptomic changes induced by FL MORC2 and LLPS-deficient mutants (Fig. 6a).

Expression analysis confirmed robust overexpression of EGFP-MORC2FL and

EGFP-MORC2ΔCC3 compared to EGFP controls (424.9-fold and 542.7-fold

increases, respectively), while EGFP-MORC2ΔIDR exhibited only a 2.2-fold increase

(Fig. 6b). Given that transfection conditions were identical and that this low

expression was consistently observed across three replicates, we speculate that IDR

deletion may impair mRNA stability or transcription. Accordingly, the

EGFP-MORC2ΔIDR group was excluded from downstream analyses.

Relative to EGFP, re-expression of EGFP-MORC2FL led to differential

expression of 240 genes (fold change > 1.2 or < 0.83, p < 0.05), including 182

downregulated and 58 upregulated transcripts (Fig. S6a). Gene Ontology (GO)

analysis showed that these genes are primarily localized to nucleus and chromatin

compartments (Fig. S6b), and enriched for functions related to transcriptional

regulation (Fig. S6c), including DNA-binding transcription factors and cytokine genes

(Fig. S6d). These results are consistent with MORC2's known role in transcriptional

silencing.

The LLPS-deficient MORC2ΔCC3 mutant showed markedly reduced regulatory

capacity on these targets (Fig. S6a, e). Notably, among 21 high-confidence MORC2

target genes previously reported in the literature (Fig. 6c), including 18 repressed and

3 activated genes (Fig. 6d), MORC2ΔCC3 failed to repress any of the 18

downregulated targets, while its effect on the upregulated genes was modest (Fig. 6e,

f). These findings indicate that condensate formation is essential for the

transcriptional repression activity of MORC2. Together, these results support a model

in which phase separation underpins MORC2's transcriptional regulatory function by

enabling its engagement with chromatin-associated gene silencing pathways.

Pathogenic MORC2 variants differentially affect phase separation, DNA binding,

and ATPase activity

MORC2 is essential for gene silencing at H3K9me3-marked loci, chromatin

regulation, and retrotransposon repression[6,34–36]. Pathogenic mutations in its

GHL-type ATPase domain have been associated with neuropathies and cancer[37–41].

We examined a panel of disease-associated variants, including those linked to

Charcot–Marie–Tooth disease type 2Z (CMT2Z: E236G, R252W, Q400R, D466N,

colored in red) and spinal muscular atrophy (SMA: S87L, S218L, F256L, R266A,

T424R, colored in blue) (Fig. 7a). In HeLa cells transiently expressing these variants,

none disrupted nuclear condensate formation, likely due to their localization within

the N-terminal ATPase domain (Fig. 7b). However, the CMT2Z-linked E236G and the

SMA-linked T424R mutants led to a significantly higher proportion of cells

exhibiting nuclear condensates, suggesting these mutations promote MORC2 phase

separation (Fig. 7b). *In vitro* ATPase assays revealed that T424R significantly

increases enzymatic activity (Fig. 7c), consistent with its reported role in promoting

ATPase dimer cycling[13]. Additionally, S218L and F256L variants also enhanced ATP

hydrolysis, whereas others, such as Q400R and D466N, showed activity comparable

to WT (Fig. 7c). Due to severe solubility issues, E236G could not be biochemically

characterized *in vitro*.

We next assessed DNA binding affinity of MORC2 variants using fluorescence

polarization (FP) assays. Despite subtle differences among the variants, none

significantly altered DNA-binding capacity compared to wild-type MORC2 (Fig. 7d),

indicating that the disease-associated mutations primarily affect enzymatic and phase

behaviors rather than DNA affinity. Taken together, these results highlight the

functional plasticity of MORC2's N-terminal domain in modulating enzymatic

activity and phase separation, while reinforcing the role of LLPS as a regulatory

mechanism rather than a prerequisite for DNA binding. These findings also provide

mechanistic insight into how specific mutations may contribute to distinct

pathological outcomes.

In summary, our results support a model in which MORC2 enforces

transcriptional silencing via multivalent interactions that promote liquid–liquid phase

separation. The N-terminal ATPase domain undergoes conformational cycling

between open and closed states, while the C-terminal CC3-mediated dimer acts as a

stable scaffold for condensate formation (Fig. 7e). Electrostatic interactions between

the basic regions of MORC2 and chromosomal DNA further potentiate phase

separation and ATPase activation. Together, these properties enable MORC2 to bridge

distal chromatin regions and establish repressive condensates, highlighting its central

role in chromatin organization and genome stability.

Discussion

Our integrative structural and functional analyses uncover a direct, mechanistically validated role for MORC2 phase separation in gene regulation, extending the understanding of chromatin-associated ATPases. The MORC gene family is evolutionarily conserved and plays a central role in epigenetic regulation and genome stability[42-44]. Originally identified in mice, MORC proteins are crucial for spermatid formation and meiotic progression[45,46]. Human MORC proteins share a characteristic domain architecture, but the mechanisms governing their chromatin interactions, especially in their full-length forms, remains unknown owing to the lack of comprehensive structural information.

In this study, we provide evidence that MORC2 undergoes liquid-liquid phase separation (LLPS) both *in vivo* and in *vitro*, forming dynamic nuclear condensates and driven by multivalent interactions between its IDR and IBD domain, and scaffolded by CC3-mediated dimerization. We demonstrate that full-length MORC2, along with its C-terminal domain, forms LLPS droplets exhibiting key characteristics, including salt sensitivity and recovery after photobleaching. Moreover, nucleosome assembly 601 DNA enhances condensate formation. Importantly, we demonstrate that LLPS is essential for the transcriptional repression activity of MORC2, and phase separation-deficient variants of MORC2 markedly reduce its silencing capacity. Notably, under our ATPase assay conditions, several disease-associated MORC2 variants—particularly S218L, F256L, and T424R—exhibited elevated enzymatic activity compared to WT. This trend partially aligns with previous findings for T424R, but differs from reports describing S87L as a loss-of-function mutant. We speculate that differences in ionic strength and assay sensitivity, as well as potential LLPS-mediated local enrichment of MORC2, may contribute to this discrepancy. These results highlight the importance of biophysical context in interpreting the functional consequences of disease mutations and warrant further mechanistic investigation. Together, our findings uncover a previously unrecognized LLPS-based mechanism by which MORC2 regulates chromatin transcriptional activity, and

suggest a potential connection between this process and neurodevelopmental

pathology.

Mutations in human MORC proteins have been linked to diverse pathologies,

including inflammatory and neurodevelopmental disorders, as well as cancer[47–49].

MORC2, in particular, has been implicated in the repression of transgenes at

H3K9me3-enriched loci and in chromatin relaxation following DNA damage[5,12,50].

Missense mutations within its ATPase domain are causally associated with axonal

neuropathies, underscoring the functional importance of ATP hydrolysis *in vivo*.

ATP-dependent dimerization of the MORC2 ATPase domain is essential for its

transcriptional regulatory activity. In addition, MORC2 contributes to the silencing of

retrotransposons and specific gene classes, including LINE-1 elements and intronless

genes, highlighting its broader role in maintaining genome integrity[51,52]. The silencing

function of MORC2 requires its ATPase activity and its three coiled-coil domains.

While the CW and TCD domains are potential chromatin-binding regions, only the

CC1 domain is known to bind DNA. Our findings in this study further reveal that the

intrinsically disordered region (IDR) of MORC2 exhibits strong DNA-binding

capacity, providing new insights into the regulatory mechanisms of MORC2,

emphasized the role of arginine/lysine-rich disordered regions in facilitating

electrostatic DNA binding[53–56]. These findings refine our understanding of MORC2's

function in chromatin architecture and illuminate its mechanistic involvement in

human disease.

In conclusion, this study demonstrates that MORC2 undergoes liquid–liquid

phase separation to form nuclear condensates, regulated by DNA binding and

associated with chromatin condensation and transcriptional silencing. The interplay

between phase separation and ATPase activity in MORC2 reveals a distinct

mechanism of transcriptional repression. These insights enhance our understanding of

MORC2's molecular mechanisms and lay the groundwork for further studies on its

role in human diseases.

Materials and Methods

Plasmid construction, cell culture and transfection

WT MORC2 (UniProt: Q9Y6X9) cDNA was amplified from a cDNA library derived from HeLa cells via reverse transcription. The MORC2 cDNA was then subcloned into the p3xFLAG-Myc-CMV-24 vector between BglII and KpnI restriction sites using seamless cloning. The seamless cloning process was facilitated by the ClonExpress II One-Step Cloning Kit (Vazyme, #C112-02) according to the manufacturer's instructions. An EGFP sequence was subsequently inserted into the BglII site at the N-terminal of MORC2 to generate the EGFP-MORC2 construct. To generate deletion mutant constructs of MORC2 for intracellular studies, specific regions of MORC2 were targeted and deleted using a polymerase chain reaction (PCR)-based approach. An 18- or 21 bp homologous sequence originating from one of the forward or reverse primers was introduced into the other primer, and the resulting PCR products were subjected to homologous recombination cloning, facilitated by the 18-bp homologous sequence. For protein expression and purification, MORC2FL and truncated MORC2 were cloned into prokaryotic expression vectors with MBP or Trx fusion proteins for bacterial expression. In addition, MORC2FL and a series of truncation mutants were purified using a HEK293F eukaryotic expression system for *in vitro* phase separation and static light scattering (SLS) assays. Detailed information on all protein purifications and their experimental applications in this study is provided in Table S2. HeLa, HEK293T cells were cultured in DMEM medium (Biological Industries, DMEM, high re maintained in a humidified incubator at 37 °C with 5% CO_2. Cell transfection was performed using Lipofectamine 3000 (Invitrogen) according to the manufacturer's instructions. Briefly, cells were seeded in appropriate culture plates the day before transfection and allowed to reach 40-60% confluency. The separately diluted Lipofectamine 3000 reagent and plasmids in Opti-MEM reduced serum medium (Gibco) were mixed, and the transfection complexes were added to cells after 20 minutes. After a 6-hour incubation, the transfection medium was replaced with fresh complete DMEM medium.

Recombinant protein expression, purification

Plasmids containing His-tagged genes were transformed into *E. coli* BL21 (DE3)

cells and cultured in Luria-Bertani medium with 50 mg/L ampicillin. When the

culture reached an optical density (OD) of 0.6-0.8 at 600 nm, the temperature was

lowered to 16 °C. Isopropyl β-D-1-thiogalactopyranoside (IPTG) (0.25 mM) was

added to induce protein expression for 18-20 hours. Cells were harvested by

centrifugation at 13,000 × g for 15 minutes, then resuspended in Buffer A (50 mM

Tris-HCl, pH 8.0, 1000 mM NaCl, 10 mM imidazole, 1 mM dithiothreitol (DTT), and

1 mM phenylmethylsulfonyl fluoride (PMSF)). The cell suspension was snap-frozen

and stored at −80 °C for subsequent protein purification. HEK293F

suspension-adapted cells were cultured in conical flasks at 37 °C, 120 rpm, and 5%

CO_2, until a density of $1.0 \times 10^\square$ cells/mL was reached. The cells were transiently

transfected with 1 mg of MORC2-6 × His plasmid using 2 mL of branched

polyethyleneimine (1 mg/mL). After 48 hours of incubation, cells were harvested by

centrifugation at 3,000 × g for 5 minutes, resuspended in Buffer A, snap-frozen, and

stored at −80 °C for further purification.

All purification steps were performed at 4°C to preserve protein activity. Frozen

cell suspensions were rapidly thawed in water, lysed by sonication (200 W for ~0.3

hours), and centrifuged at 20,000 × g for 30 minutes at 4 °C. The supernatant was

filtered through a 0.22 μm syringe filter to obtain soluble material. A His-Trap

Ni-NTA 6FF chromatography column was equilibrated with five column volumes of

Buffer A (Buffer A: 50 mM Tris-HCl, pH 8.0, 1000 mM NaCl). The clarified extract

was loaded onto the column, washed with Buffer A to remove non-specific proteins,

and eluted with a linear gradient of nickel elution buffer (Buffer B: 50 mM Tris-HCl,

pH 8.0, 1 M NaCl, 300 mM imidazole, 1 mM DTT). Eluted fractions were analyzed

by SDS-PAGE and Coomassie staining. The protein of interest was concentrated,

exchanged into Buffer C (50 mM Tris-HCl, pH 8.0, 1000 mM NaCl, 1 mM EDTA, 1

mM DTT) using an Amicon Ultra centrifugal filter unit, following overnight

treatment with Benzonase nuclease to eliminate nucleic acid contamination or HRV

3C Protease to cut His-tag. The purification process was subsequently refined through

gel filtration using a Superdex 6 Increase column (GE Healthcare) for the MORC2FL.

Following this, fragments were conjugated with a GPGS linker, and the resulting

mixture was subjected to a size-exclusion chromatography process (Superdex 200/75

column, GE Healthcare) to yield a final solution comprising 50 mM Tris-HCl, pH 8.0,

500 mM NaCl, 1 mM EDTA, and 1 mM DTT. Peak fractions were analyzed by

SDS-PAGE and Coomassie staining, confirming the successful purification of highly

homogeneous MORC2FL (1-1032) and various fragments (Fig. S7), then flash-frozen

in liquid nitrogen and stored at −80 °C.

Crystallization, data collection and processing

The CC3 domain (901-1003) and various mutations of MORC2constructs were

cloned into a modified pET32M vector (with Trx-tag) using standard PCR-based

methods and confirmed by DNA sequencing. Due to the lack of homologous

structures, protein crystals with heavy atom derivatives are required to determine their

phases. Therefore, we used the methionine-deficient strain B834 to express the target

protein with selenomethionine (SeMet) in an inorganic selenosubstituted medium.

Recombinant proteins were expressed in *E. coli* B834 cells in M9 medium at 16 °C.

MORC2 CC3 fused with a GPGS linker were finally changed by step of

size-exclusion chromatography (Superdex 75 increase column, GE Healthcare) into

50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 1 mM EDTA and 1 mM DTT). The freshly

purified protein was concentrated to 10~20 mg/mL. Crystals were obtained by the

static sitting drop method in 0.1M Citric acid (pH 3.5) and 25% w/v PEG 3350 at

25°C. Glycerol (20%) was added as the cryo-protectant. A 3.1-Å resolution X-ray

dataset was collected at the beamline BL18U1 of the Shanghai Synchrotron Radiation

Facility (SSRF). The diffraction data were processed and scaled by HKL2000

(http://www.hkl-xray.com/). Using the structure of auto-build model by SAD data and

AlpahFold2 predicted model as the search model, the initial structural model was

solved by molecular replacement in PHASER (https://www.ccp4.ac.uk/). Further

manual model adjustment and refinement were completed iteratively using COOT and

PHENIX. The final structure was validated by PISA and MolProbity

(https://www.phenix-online.org/). The final refinement statistics of the complex

structure are listed in Supplementary Table 1. All structure figures were prepared

using the programme PyMOL (http://pymol.sourceforge.net/).

Protein labeling with fluorophore

The Cy3 NHS ester (AAT Bioquest) fluorophores were dissolved in DMSO at 10

mg/mL. Purified proteins were prepared in reaction buffer at a concentration of 10

mg/mL. The fluorophores were added to the protein solution at a 1:1 molar ratio. The

reaction was conducted with shaking at room temperature for 1 hour in the dark. After

the reaction, labeling was quenched with 200 mM Tris-HCl, pH 8.3, and the labeled

proteins were desalted using a HiTrap desalting column (GE Healthcare).

Fluorescence labeling efficiency was assessed using a Nanodrop 2000

spectrophotometer (ThermoFisher #ND-2000). The final labeling efficiency of each

protein was adjusted to 1% by mixing the labeled protein with an excess of unlabeled

protein (1:99 ratio).

Lentiviral mediated H2A-mCherry inducible expression in HeLa cells

Stable expression of H2A-mCherry in HeLa cells was achieved through

lentiviral-mediated transduction. First, H2A-mCherry gene sequence was inserted into

the inducible lentiviral vector pTRIPZ to generate the expression plasmid

pTRIPZ-H2A-mCherry. Lentiviral particles were generated by triple transfection of

HEK293T cells with pTRIPZ-H2A-mCherry, PMD2.G, and PSPAX2 plasmids using

Lipofectamine 3000. The viral supernatant was collected 48 hours after transfection,

filtered with 0.45-μm filter to remove cell debris, and used to transduce HeLa cells.

Following transduction, HeLa cells were selected with 50 μg/mL hygromycin to

establish stable cell lines. H2A-mCherry expression was induced with 1 μg/mL

doxycycline.

Live cell imaging and immunofluorescence staining

For live-cell imaging, HeLa cells were seeded onto 35 mm culture dishes with

glass-bottom plates (Biosharp) and transfected with EGFP-MORC2 or its mutant

constructs. Images were captured 12 hours post-transfection using a Nikon confocal

microscope equipped with an incubator system (37 °C, 5% CO_2). Nuclei were stained

with Hoechst 33342 or visualized using mCherry-labeled H2A fusion protein. For

immunofluorescence, cells were fixed with 4% paraformaldehyde, permeabilized with

0.1% Triton X-100, and blocked with 10% goat serum. Primary antibodies were

incubated overnight at 4 °C, followed by incubation with fluorescence-conjugated

secondary antibodies for 2 hours at room temperature in the dark. Fluorescence

images were acquired using a Nikon confocal microscope.

CRISPR-Cas9 mediated MORC2 knockout cells

MORC2 knockout (KO) HeLa cell lines were generated using the CRISPR-Cas9

system. A CRISPR single-guide RNA (sgRNA) targeting exon 5 of MORC2

(5'-ACACCTGAGTCTACTCAGAT-3') was designed using the online tool

(http://crispor.tefor.net/). Two DNA oligos with complementary sgRNA targeting

sequence and additional bases were designed as previously described[57,58]. These

oligos were annealed and cloned into pLentiCRISPR v2 (Addgene plasmid #52961)

to generate the MORC2 knockout vector. After transfection and puromycin selection,

MORC2-deleted cells were obtained. Individual cells were sorted into separate wells

of a culture plate to establish monoclonal cell lines using flow cytometry. To validate

the knockout cell lines, genomic DNA was extracted from both WT and KO cells. The

specific regions targeted by the sgRNA were amplified using PCR with primers:

5'-CAACATTCTCGAGCTGGACCTACAG-3'   (forward)   and

5'-CCACGACAAGACTGGAAACGTGACTC-3' (reverse). The PCR products were

purified and subjected to Sanger sequencing, revealing the deletion and insertion

mutations in the target sites of the MORC2 allelic genes (Fig. S2d).

Generation of endogenous EGFP-MORC2 chimeric mice

CRISPR-Cas9 was used to generate EGFP-MORC2 chimeric mice. The EGFP

and linker sequence were inserted at the N-terminus of the mouse *MORC2a* gene.

Briefly, a single-guide RNA (sgRNA) was designed to target exon 1 of *MORC2a*,

with the sequence GACTGAAGACTCATTGCTGTCA. A donor DNA fragments

containing the EGFP-linker region flanked by homologous arms was constructed.

Fertilized mouse zygotes were collected, and a mixture of Cas9 mRNA (50 ng/µL),

sgRNA (25 ng/µL), and donor DNA (50 ng/µL) was microinjected into pronucleus of

the zygotes. The injected embryos were transferred into the oviducts of

pseudopregnant recipient female mice. Resulting pups were genotyped by PCR and

Sanger sequencing to confirm transgene integration (Fig. S2f). The genotyping

primers   for   the   EGFP-MORC2   allele   were   as   follows: Left-F1:

GCGGTGGTTGAGTTCCAATTCC; Left-R1: CTTCAGGGTCAGCTTGCCGTAG;

Right-F1:   GAACGGCATCAAGGTGAACTTC;   Right-R1:

TGTATCCAAGGTAATCTCTGTGGTA.

Mice were bred at the Animal Center of the University of Science and

Technology of China (USTC). Animals were housed under controlled conditions

(22 °C, 12-hour light/dark cycle) with free access to food and water ad libitum. All

experimental procedures were approved by the Institutional Animal Care and Use

Committee (IACUC) of the University of Science and Technology of China (Hefei,

China).

Fluorescence recovery after photobleaching (FRAP)

The FRAP assay was performed to investigate the dynamics of protein mobility

within condensates, either in *in vivo* or *in vitro*. Specific regions of interest (ROIs)

approximately 2 µm² in size were selected and subjected to photobleaching using 100%

laser power (488 nm) for approximately 20 ns. One droplet was left unbleached as a

control. Fluorescence recovery within the ROI was monitored for 100-300 seconds.

FRAP was performed using a 63x oil immersion objective mounted on a Nikon

confocal microscope (in cells) or a Zeiss LSM 880 (in vitro). Fluorescence intensity

was normalized to the pre-bleach signal to analyze the recovery kinetics. For brain

slice FRAP, organotypic brain slices were prepared from adult chimeric mice

expressing endogenous EGFP-MORC2, following a slightly modified protocol

previously described[59]. After decapitation, brains were rapidly extracted and placed in

ice-cold dissection medium consisting of Hibernate A, 2% B27 supplement, 2 mM

L-glutamine, and 1% penicillin–streptomycin. The cerebellum and midbrain were

removed, and the remaining cerebral hemispheres were coronally sectioned at

250 µm using a McIlwain tissue chopper. Slices were gently separated in chilled

dissection medium and transferred onto glass-bottom dishes containing culture

medium composed of Neurobasal A, 2% B27 supplement, 2 mM L-glutamine, and 1%

penicillin–streptomycin. Imaging was initiated immediately after brain slice

preparation. FRAP experiments were performed using the Nikon confocal microscope.

Photobleaching was carried out with 5-7 laser pulses (50 µs dwell time), and images

were acquired at 2-s intervals. Post-bleach images were acquired 20 s after

photobleaching.

Electrophoretic mobility shift assays (EMSA)

Widom 601 DNA (147 bp, 25 nM) was amplified by PCR and incubated with

increasing concentrations of MORC2 or its domain constructs in binding buffer (20

mM Tris-HCl, pH 7.5, 45 mM KCl, 4 mM $MgCl_2$, 5% glycerol, 5 mM DTT). The

reaction mixture was loaded onto a 6% polyacrylamide gel and electrophoresed at 120

V for 70 minutes. DNA was visualized using SYBR Gold stain, and images were

captured using the ChemiDoc MP Imaging System.

Size-exclusion chromatography coupled with static light scattering

The analysis was performed using an AKTA FPLC system (GE Healthcare)

coupled to a static light scattering detector (miniDawn, Wyatt) and a differential

refractive index detector (Optilab, Wyatt). Protein samples (70 µM) were filtered

and applied to a Superdex 200 increase or Superose 6 increase column,

pre-equilibrated with a column buffer containing 50 mM Tris-HCl, pH 8.0,

150 mM NaCl or 1000 mM NaCl (MORC2FL), 1 mM EDTA, and 1 mM DTT.

Data were analyzed using ASTRA6 software (Wyatt).

***In vitro* phase separation and fluorescence microscopy imaging**

Droplet formation was monitored using differential interference contrast (DIC)

and fluorescence microscopy. To induce phase separation, protein was added to phase

separation buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM DTT). For

consistency, all MORC2 samples were diluted to a final concentration of 4 μM with

phase separation buffer containing 200 mM NaCl, unless otherwise specified. All

preparation steps were performed at 4 °C to preserve protein activity. The phase

separation mixtures were then equilibrated at room temperature for 20 min. A

5-10 μL aliquot of each sample was placed onto a custom-made glass slide, covered

with an 18 mm diameter coverslip, and incubated at room temperature for 20 min

before imaging.

Protein isotope labeling and NMR Spectroscopy

Isotopically labeled protein samples were expressed in minimal M9/H_2O or

M9/D_2O medium. Uniformly ^{15}N-labeled (or ^{13}C- or ^{15}N-labeled) protein samples

were prepared in M9/H_2O medium supplemented with 1 g/L of $^{15}NH_4Cl$ and 2 g/L

of glucose (or ^{13}C-glucose). The cells were harvested at an OD_{600} of approximately

1.0. Isotope-labeled precursors and amino acids were purchased from Cambridge

Isotope Laboratories (CIL). Isotopically labeled samples were prepared in 50 mM

Tris-HCl, pH 8.0, 100 mM NaCl, and 7% D O. NMR experiments were performed

using Bruker Avance III 600 MHz spectrometers equipped with a cryo-probe. NMR

spectra of MORC2 and its complex were recorded at 283 K (for

CC3-IBD/CC3/IBD). Protein concentrations ranged from 0.5 to 1.0 mM, and

nucleotides were added to a final concentration of 10 mM. All NMR spectra were

processed using NMRPipe and analyzed with NMRViewJ software

(http://www.onemoonscientific.com).

Fluorescence polarization Assay (FP)

Fluorescence polarization (FP) assays were performed at room temperature (25 °C) using a BioTek H1 fluorescence spectrophotometer to evaluate the binding affinity between MORC2 protein and 5-carboxyfluorescein (5-FAM)-labeled 601 DNA (FAM-601) across a concentration range of 0-2000 nM. The 147-bp Widom 601 DNA fragment was generated via PCR amplification using a 5-FAM-labeled primer. The labeled primer was dissolved in 0.1 M NaHCO□ buffer, pH 8.3, and the amplified product was purified by agarose gel extraction to eliminate residual primers and nonspecific amplification products.

Fluorescence measurements were carried out in assay buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, and 10% glycerol) containing purified FAM-601 DNA at a final concentration of 0.4 ng/μL. Excitation and emission wavelengths were optimized according to the instrument manual, with excitation set at 485 nm and emission at 525 nm. Fluorescence polarization values were recorded in triplicate, and control measurements were included to correct for background signals.

Data were analyzed to determine the equilibrium dissociation constant (K_d) for the MORC2-FAM-601 interaction by fitting data to a log(agonist) vs. response model. In parallel, the effect of small molecules on this binding interaction was assessed by introducing compounds into the assay mixture and monitoring changes in fluorescence polarization.

ATPase activity measurements

ATPase activity was measured using 1 μM MORC2 or its mutant variant at 37 °C in 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 8 mM MgCl$_2$, 10% glycerol and 5 mM ATP[60]. All experiments were repeated independently at least three times.

RNA-seq and bioinformatic analysis

MORC2-knockout HeLa cells were seeded at equal density and transfected with 3 μg of plasmid DNA and 12 μg of PEI (polyethylenimine) per dish. After

6 hours, the transfection medium was replaced with fresh complete medium, and
cells were incubated for an additional 48 hours. Cells were then washed twice with
ice-cold PBS and lysed directly in RNA Isolator (R401, Vazyme) on ice. Total RNA
was extracted using the RNeasy Mini Kit (Qiagen, 74004) according to the
manufacturer's protocol. RNA quality was assessed using NanoDrop 2000 (Thermo
Fisher Scientific), Agilent 2100 Bioanalyzer, and LabChip GX (PerkinElmer). For
library preparation, 1 µg of total RNA was used per sample, employing the VAHTS
Universal V6 RNA-seq Library Prep Kit for MGI®. The resulting double-stranded
DNA libraries were denatured, circularized, and digested to obtain single-stranded
circular DNA. DNA nanoballs (DNBs) were generated from these templates through
rolling circle amplification (RCA). Sequencing was performed on the MGI
DNBSEQ-T7 platform (MGI Tech Co., Ltd) using combinatorial Probe-Anchor
Synthesis (cPAS) chemistry.

Before performing differential expression analysis, lowly expressed genes were
filtered out by retaining only those with a raw count of ≥ 10 in at least three samples.
Differential gene expression analysis between sample groups was carried out using
the DESeq2 package (v1.26.0). Genes with an absolute fold change > 1.2 or < 0.83,
and $p < 0.05$ were defined as differentially expressed genes (DEGs). These DEGs
were subsequently subjected to Gene Ontology (GO) enrichment analysis in DAVID
(Knowledgebase v2025_1).

Statistical Analysis

All statistical analyses were performed in GraphPad Prism. All data underwent
normality testing (D'Agostino & Pearson test and Anderson-Darling test) in advance.
Two-tailed unpaired Student's t-test post-hoc Welch's correction and one-way
ANOVA with Tukey's post hoc test was used when data were distributed normally.
The significance level was set as $p < 0.05$. All statistical results are provided as
Source Data.

Acknowledgments

We thank Dr. Xuebiao Yao (University of Science and Technology of China, USTC), Dr. Zhi Qi and Dr. Linyu Zuo (Peking University) for helpful discussion. We thank members of the Shanghai Synchrotron Radiation Facility (SSRF, China; beamlines BL02U1, BL18U1, and BL19U1) for X-ray beam time. This work was supported by grants from the National Key R&D Program of China (2019YFA0508402), the National Natural Science Foundation of China (22122703, 32470808, 32170767, 32471273, 31971144, and T2221005), the Center for Advanced Interdisciplinary Science and Biomedicine of IHM (QYPY20220014), the Strategic Priority Research Program of the Chinese Academy of Sciences (XDB0490000), the Major Frontier Research Project of USTC (LS9100000002), USTC Research Funds of Double First-Class Initiative (YD9100002507 and YD9100002054), the Anhui Medical University Foundation (9101224201), and National Natural Science Foundation Incubation Program of The Second Affiliated Hospital of Anhui Medical University (2021GQFY02).

Author contributions

Y.B., Y.Z., W.X., W.D.,C.H., and C.W. designed the experiments. Y.B., Y.Z., W.X., and W.D. conducted the experiments. Y.Z. determined the crystal structure. Y.W., W.J. and F.Z. assisted with the experiments and reagents. All authors analyzed the data. Y.B., Y.Z., C.H., and C.W. wrote the manuscript with input from all authors. All authors approved the final version of the manuscript. C.H., and C.W supervised the project.

Competing interests

The authors declare that they have no competing interests.

Data and materials availability

All data needed to evaluate the conclusions in the paper are present in the paper

757   and/or the Supplementary Materials. Raw gene expression data are available in

758   Source Data. The atomic coordinates of the MORC2 CC3 domain have been

759   deposited to the Protein Data Bank under the accession code: 9KQL.

References

1. Alabert, C. & Groth, A. Chromatin replication and epigenome maintenance. *Nat Rev Mol Cell Biol* **13**, 153–167 (2012).

2. Ishihara, S. *et al.* Local states of chromatin compaction at transcription start sites control transcription levels. *Nucleic Acids Res* **49**, 8007–8023 (2021).

3. Downs, J. A. & Gasser, S. M. Chromatin remodeling and spatial concerns in DNA double-strand break repair. *Curr Opin Cell Biol* **90**, 102405 (2024).

4. Blackledge, N. P. & Klose, R. J. The molecular principles of gene regulation by Polycomb repressive complexes. *Nat Rev Mol Cell Biol* **22**, 815–833 (2021).

5. Li, D.-Q. *et al.* MORC2 signaling integrates phosphorylation-dependent, ATPase-coupled chromatin remodeling during the DNA damage response. *Cell Rep* **2**, 1657–1669 (2012).

6. Müller, I. & Helin, K. Keep quiet: the HUSH complex in transcriptional silencing and disease. *Nat Struct Mol Biol* **31**, 11–22 (2024).

7. Zhong, Z. *et al.* MORC proteins regulate transcription factor binding by mediating chromatin compaction in active chromatin regions. *Genome Biol* **24**, 96 (2023).

8. Xue, Y. *et al.* Arabidopsis MORC proteins function in the efficient establishment of RNA directed DNA methylation. *Nat Commun* **12**, 4292 (2021).

9. Weiser, N. E. *et al.* MORC-1 Integrates Nuclear RNAi and Transgenerational Chromatin Architecture to Promote Germline Immortality. *Developmental Cell* **41**, 408-423.e7 (2017).

10. Moissiard, G. *et al.* MORC family ATPases required for heterochromatin condensation and gene silencing. *Science* **336**, 1448–1451 (2012).

11. Wang, H., Zhang, L., Luo, Q., Liu, J. & Wang, G. MORC protein family-related signature within human disease and cancer. *Cell Death Dis* **12**, 1112 (2021).

12. Tchasovnikarova, I. A. *et al.* Hyperactivation of HUSH complex function by Charcot-Marie-Tooth disease mutation in MORC2. *Nature Genetics* (2017) doi:10.1038/ng.3878.

13. Douse, C. H. *et al.* Neuropathic MORC2 mutations perturb GHKL ATPase dimerization dynamics and epigenetic silencing by multiple structural mechanisms. *Nat Commun* **9**, 651 (2018).

14. Xie, H.-Y., Zhang, T.-M., Hu, S.-Y., Shao, Z.-M. & Li, D.-Q. Dimerization of MORC2 through its C-terminal coiled-coil domain enhances chromatin dynamics and promotes DNA repair. *Cell Commun Signal* **17**, 160 (2019).

15. Morais Cabral, J. H. *et al.* Crystal structure of the breakage-reunion domain of DNA gyrase. *Nature* **388**, 903–906 (1997).

16. Chen, K. *et al.* Crystal structure of the hinge domain of Smchd1 reveals its dimerization mode and nucleic acid-binding residues. *Sci Signal* **13**, eaaz5599 (2020).

17. Liu, J. *et al.* Cascading MutS and MutL sliding clamps control DNA diffusion to activate mismatch repair. *Nature* **539**, 583–587 (2016).

18. Ortega, J., Lee, G. S., Gu, L., Yang, W. & Li, G.-M. Mispair-bound human MutS-MutL complex triggers DNA incisions and activates mismatch repair. *Cell Res* **31**, 542–553 (2021).

19. Lee, K. *et al.* The structure of an Hsp90-immunophilin complex reveals cochaperone recognition of the client maturation state. *Mol Cell* **81**, 3496-3508.e5 (2021).

20. Pastor, W. A. *et al.* MORC1 represses transposable elements in the mouse male germline. *Nat*

Commun **5**, 5795 (2014).

21. Dopkins, N. *et al.* A field guide to endogenous retrovirus regulatory networks. *Mol Cell* **82**,
3763–3768 (2022).

22. Fendler, N. L. *et al.* Identification and characterization of a human MORC2 DNA binding
region that is required for gene silencing. *Nucleic Acids Res* **53**, gkae1273 (2025).

23. Leidescher, S. *et al.* Spatial organization of transcribed eukaryotic genes. *Nat Cell Biol* **24**,
327–339 (2022).

24. Membrane-less compartments in the nucleus: Separated or connected phases? *Current*
Opinion in Cell Biology **84**, 102215 (2023).

25. Boeynaems, S. *et al.* Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol*
28, 420–435 (2018).

26. Yoshizawa, T. *et al.* Nuclear Import Receptor Inhibits Phase Separation of FUS through
Binding to Multiple Sites. *Cell* **173**, 693-705.e22 (2018).

27. Verba, K. A. *et al.* Atomic structure of Hsp90-Cdc37-Cdk4 reveals that Hsp90 traps and
stabilizes an unfolded kinase. *Science* **352**, 1542–1547 (2016).

28. Alberti, S., Gladfelter, A. & Mittag, T. Considerations and Challenges in Studying
Liquid-Liquid Phase Separation and Biomolecular Condensates. *Cell* **176**, 419–434 (2019).

29. Zhang, Y. *et al.* MORC3 Forms Nuclear Condensates through Phase Separation. *iScience* **17**,
182–189 (2019).

30. Kalderon, D., Roberts, B. L., Richardson, W. D. & Smith, A. E. A short amino acid sequence
able to specify nuclear location. *Cell* **39**, 499–509 (1984).

31. Alderson, T. R. & Kay, L. E. NMR spectroscopy captures the essential role of dynamics in
regulating biomolecular function. *Cell* **184**, 577–595 (2021).

32. Kim, T. H. *et al.* Phospho-dependent phase separation of FMRP and CAPRIN1 recapitulates
regulation of translation and deadenylation. *Science* **365**, 825–829 (2019).

33. Murthy, A. C. *et al.* Molecular interactions underlying liquid–liquid phase separation of the
FUS low-complexity domain. *Nat Struct Mol Biol* **26**, 637–648 (2019).

34. Pandiloski, N. *et al.* DNA methylation governs the sensitivity of repeats to restriction by the
HUSH-MORC2 corepressor. *Nat Commun* **15**, 7534 (2024).

35. Fukuda, K., Okuda, A., Yusa, K. & Shinkai, Y. A CRISPR knockout screen identifies
SETDB1-target retroelement silencing factors in embryonic stem cells. *Genome Res* **28**,
846–858 (2018).

36. Liu, N. *et al.* Selective silencing of euchromatic L1s revealed by genome-wide screens for L1
regulators. *Nature* **553**, 228–232 (2018).

37. Jacquier, A., Roubille, S., Lomonte, P. & Schaeffer, L. Microrchidia CW-Type Zinc Finger 2,
a Chromatin Modifier in a Spectrum of Peripheral Neuropathies. *Front Cell Neurosci* **16**,
896854 (2022).

38. Stafki, S. A. *et al.* The Spectrum of MORC2-Related Disorders: A Potential Link to Cockayne
Syndrome. *Pediatr Neurol* **141**, 79–86 (2023).

39. Chung, H. Y. *et al.* Morc2a variants cause hydroxyl radical-mediated neuropathy and are
rescued by restoring GHKL ATPase. *Brain* **147**, 2114–2127 (2024).

40. Ding, Q.-S. *et al.* Aberrant high expression level of MORC2 is a common character in
multiple cancers. *Hum Pathol* **76**, 58–67 (2018).

41. Zhang, S. *et al.* Oncogenic MORC2 in cancer development and beyond. *Genes Dis* **11**,

861–873 (2024).

42. Li, D.-Q., Nair, S. S. & Kumar, R. The MORC family: new epigenetic regulators of

transcription and DNA damage response. *Epigenetics* **8**, 685–693 (2013).

43. Koch, A. *et al.* MORC Proteins: Novel Players in Plant and Animal Health. *Front Plant Sci* **8**,

1720 (2017).

44. Dong, W. *et al.* MORC Domain Definition and Evolutionary Analysis of the MORC Gene

Family in Green Plants. *Genome Biol Evol* **10**, 1730–1744 (2018).

45. Inoue, N. *et al.* New gene family defined by MORC, a nuclear protein required for mouse

spermatogenesis. *Hum Mol Genet* **8**, 1201–1207 (1999).

46. Wang, G.-L. *et al.* Identification and expression analysis of a novel CW-type zinc finger

protein MORC2 in cancer cells. *Anat Rec (Hoboken)* **293**, 1002–1009 (2010).

47. Guddeti, R. K., Chutani, N. & Pakala, S. B. MORC2 Interactome: Its Involvement in

Metabolism and Cancer. *Biophys Rev* **13**, 507–514 (2021).

48. Tencer, A. H. *et al.* Molecular mechanism of the MORC4 ATPase activation. *Nat Commun* **11**,

5466 (2020).

49. Prasanth, M. I. *et al.* Role of Epigenetic Modulation in Neurodegenerative Diseases:

Implications of Phytochemical Interventions. *Antioxidants (Basel)* **13**, 606 (2024).

50. Zhang, L., Zhang, L. & Li, D.-Q. MORC2 regulates DNA damage response through a

PARP1-dependent pathway. *Nucleic Acids Research* (2019) doi:10.1093/nar/gkz545.

51. Tunbak, H. *et al.* The HUSH complex is a gatekeeper of type I interferon through epigenetic

regulation of LINE-1s. *Nat Commun* **11**, 5387 (2020).

52. Pandiloski, N. *et al.* DNA methylation governs the sensitivity of repeats to restriction by the

HUSH-MORC2 corepressor. *Nat Commun* **15**, 7534 (2024).

53. Vuzman, D. & Levy, Y. Intrinsically disordered regions as affinity tuners in protein-DNA

interactions. *Mol Biosyst* **8**, 47–57 (2012).

54. van der Lee, R. *et al.* Classification of intrinsically disordered regions and proteins. *Chem Rev*

114, 6589–6631 (2014).

55. Ukmar-Godec, T. *et al.* Lysine/RNA-interactions drive and regulate biomolecular

condensation. *Nat Commun* **10**, 2909 (2019).

56. Ukmar-Godec, T. *et al.* Multimodal interactions drive chromatin phase separation and

compaction. *Proceedings of the National Academy of Sciences* **120**, e2308858120 (2023).

57. Shalem, O. *et al.* Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science*

343, 84–87 (2014).

58. Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide libraries for

CRISPR screening. *Nat Methods* **11**, 783–784 (2014).

59. Li, C. H. *et al.* MeCP2 links heterochromatin condensates and neurodevelopmental disease.

Nature **586**, 440–444 (2020).

60. Microtiter assay for glutamine synthetase biosynthetic activity using inorganic phosphate

detection. *Analytical Biochemistry* **327**, 114–118 (2004).

Figure legends

Fig. 1 The CC3 domain serves as the structural basis for full-length MORC2
dimerization.

(a) The domain organization of human MORC2FL, highlighting the ATPase module
(GHL, CC1 and S5 domains, collectively termed NTD), the C-terminal domains
(CTD) with coiled-coil regions (CC2 and CC3), CW-type zinc finger (CW),
intrinsically disordered region (IDR, marked by a fresh green line by IUPRED2
prediction), nuclear localization signal (NLS), Tudor-chromodomain (TCD), and a
30-residues tail domain (IBD).

(b) Purification and characterization of full-length human MORC2 in HEK293F cell
system. SDS-PAGE analysis of the purified protein demonstrates its purity and
distribution.

(c) Static light-scattering (SLS) analysis shows that MORC2 forms dimers in the
absence of ATP, with a molecular weight of 239 ± 1 kDa in standard buffer (mint
green line) and 303 ± 2 kDa in the presence of 601 DNA (sunflower yellow line).

(d) SLS analysis reveals the oligomerization states of MORC2 truncations.
Full-length MORC2 lacking CC3-IBD (1-900) is monomeric, while CC3-IBD
(901-1032) and CC3 alone (901-1003) form stable dimers.

(e) Crystal structural of the CC3 dimer. Hydrophobic residues contributing to dimer
stability (L911, L915, and F922) are highlighted using ball-and-stick representations.
Top view of the hydrophobic core of the CC3 dimer. Layers of hydrophobic contacts
stabilize the dimer interface, illustrated with paired residues (e.g., Layer 1:
L911-L911', Layer 2: L915-L915', and Layer 3: F922-F922').

(f) Additional SLS analysis confirms that fragments spanning residues 537-1032,
744-1032, and Trx-901-975 (CC3 truncation) adopt dimeric conformations.

Fig. 2 MORC2 undergoes liquid-liquid phase separation (LLPS) under

physiological conditions.

(a) SDS-PAGE analysis of EGFP-MORC2FL protein purified from HEK293F cells,

showing protein concentration at 2.5 μM and 5 μM.

(b) Sedimentation assay shows the distribution of EGFP-MORC2FL (7.2 μM)

between supernatant (S) and pellet (P) fractions under 1000 mM (control) and 150

mM NaCl, indicating its propensity for phase separation.

(c) EGFP-MORC2 (7.2 μM) undergoes phase separation into droplets in a buffer

containing 150 mM NaCl, but not in 1000 mM NaCl buffer. Scaler bar: 2 μm.

(d) The proteins were examined in a buffer containing 150 mM NaCl, and phase

separation was assessed by fluorescence microscopy with 488 nm excitation for EGFP.

Quantification of droplet areas from (Fig. S2a), presented as violin plots with box

plots indicating medians. Data are derived from ≥ 3 independent images and reflect

the size distribution of phase-separated droplets. Data are presented as mean ± SEM;

one-way ANOVA with Tukey's post hoc test. ****$p < 0.0001$; *$p < 0.05$.

(e) Time-lapse FRAP imaging of EGFP-MORC2FL droplets formed *in vitro*,

demonstrating fluorescence recovery over time. Scaler bar: 2 μm.

(f) Quantitative analysis of fluorescence recovery from (e), showing the dynamic

properties of *in vitro* MORC2 droplets. Data represent mean ± SEM, with $n ≥ 3$

replicates.

(g) Immunostaining of endogenous MORC2 in HeLa cells revealed punctate localized

within the nucleus, likely representing condensates involved in transcriptional

regulation (white arrows). Nuclei are counterstained with DAPI to visualize

chromatin-rich regions. Scaler bar: 5 μm.

(h) Live-cell imaging of transiently transfected EGFP-MORC2FL showed its

assembly into dispersed, nearly spherical condensates in the nucleus, while stable

expression of H2A-mCherry served to mark chromatin-enriched domains. Scaler bar:

10 μm.

(i) Time-lapse FRAP analysis of EGFP-MORC2FL condensates in the nuclei of

transiently transfected HeLa cells revealed rapid fluorescence recovery within

seconds, indicative of their dynamic, liquid-like nature. Scaler bar: 1 µm.

(j) Quantitative analysis of fluorescence recovery from (i), showing the dynamic

properties of *in vitro* MORC2 droplets. Data represent mean ± SEM, with n ≥ 10

replicates.

(k) Sections of brain and spinal cord from endogenous EGFP-MORC2 chimeric mice

show EGFP-MORC2 condensation distribution in NeuN-positive neurons. Scaler bar:

10 µm.

(l) Representative FRAP images from of FRAP experiments performed on fresh

250-µm brain slices of endogenous EGFP-MORC2 chimeric mice demonstrate the

dynamic nature of endogenous MORC2 condensates. Data are presented as mean ±

SEM, with *n* = 3 droplets analyzed. Scaler bar: 5 µm.

Fig. 3 Multiple domains regulate phase separation of MORC2.

(a) Sequential deletion analysis revealed that the NTD is dispensable for condensate formation, as EGFP-tagged NTD alone failed to form condensates and was excluded from the nucleus. Fusion with a canonical nuclear localization signal ("PKKKRKV") restored nuclear localization but did not rescue condensate formation. In contrast, the CTD alone was sufficient to form nuclear condensates. Deletion of the CW domain resulted in enlarged condensates in a subset of cells, while removal of CC2 or the TCD had no apparent effect on condensate formation. Strikingly, deletion of either the IDR or CC3 completely abolished nuclear condensate assembly. Deletion of the IBD markedly reduced the frequency of condensate formation. Loss of the intrinsic NLS led to cytoplasmic localization of MORC2; however, no condensates were detected in the cytoplasm under transient transfection conditions in HeLa cells. Together, these results define the IDR and CC3 as the minimal and essential elements for MORC2-mediated phase separation, and implicate the CW domain, IBD, and nuclear microenvironment in the fine-tuning of condensate assembly. Scaler bar: 10 μm.

(b) *In vitro* phase separation of Cy3-labeled CTDΔCW at concentrations from 1.25 μM to 20 μM in 150 mM NaCl buffer, visualized as spherical condensates. Scaler bar: 10 μm.

(c) Quantification of droplet areas from (b), displayed as violin plots with box plots indicating median values. Data are presented as mean ± SEM; one-way ANOVA with Tukey's post hoc test. ****$p < 0.0001$, *n.s.* not significant.

(d) SDS-PAGE analysis of Cy3-CTDΔCW protein distribution between supernatant (S) and pellet (P) fractions after centrifugation at increasing concentrations (5 μM to 40 μM) in 150 mM NaCl buffer. At 40 μM, increasing the NaCl concentration to 300 mM or 500 mM reduced droplet formation, indicating salt sensitivity.

(e) Quantification of Cy3-CTDΔCW showing the percentage of soluble protein from densitometric analysis of S/P fractions in panel (d), based on three independent biological replicates.

Fig. 4 Multivalent interactions between IDR and IBD drive MORC2 LLPS.

(a) Domain organization of MORC2 C-terminal regions analyzed by NMR titration, including the IDR, CC3, and IBD.

(b) ^{15}N-labeled CC3 (residues 901–1003) shows no chemical shift upon titration with IDR (residues 593–735), indicating no detectable interaction.

(c) ^{15}N-labeled IBD (residues 1004–1032) exhibits clear chemical shift perturbations (CSPs) upon IDR titration, confirming a direct, specific interaction.

(d-f) NMR titrations of ^{15}N-labeled IBD with IDR subregions: IDRa (593–643), IDRb (644–694), and IDRc (695–735). IDRa induces clear CSPs (d), while IDRb (e) and IDRc (f) show minimal or no shifts, identifying IDRa as the primary IBD-binding segment.

(g) Sequence analysis of IDR subregions reveals that IDRa is enriched in proline and arginine residues, enabling electrostatic interactions. Sequence alignment of IBD across species highlights conserved residues involved in IDRa binding. A summary table lists representative residues from IDRa and IBD exhibiting CSPs in titration assays (marked with asterisks).

(h) Representative confocal images of HeLa cells expressing EGFP-tagged MORC2ΔIDRa, ΔIDRb, or ΔIDRc constructs. Scale bar: 10 μm.

(i) Quantification of droplet area per cell in (h). Deletion of IDRa significantly impairs condensate formation. n = 10 cells per condition. Data are presented as mean ± SEM; statistical analysis: one-way ANOVA with Tukey's post hoc test. ***p < 0.001; *n.s.* not significant.

(j) Working model of MORC2 LLPS. CC3 dimerization serves as a structural scaffold, while weak, transient, but specific multivalent interactions between IDRa and IBD cooperatively promote condensate formation.

Fig. 5 DNA binding enhances MORC2 phase separation and modulates its ATPase activity.

(a) Electrophoretic mobility shift assay (EMSA) showing that MORC2FL binds 25 nM 601 DNA, as evidenced by shifts in DNA mobility corresponding to protein-DNA complexes.

(b) EMSA highlighting the strong DNA-binding affinity of the CC1 domain, corroborating previous reports that identify this region as a key mediator of DNA interaction.

(c-f) EMSA results demonstrating the binding capacities of individual MORC2 domains to 25 nM 601 DNA: CC2 (c), IDR (d), TCD-CC3-IBD (e), and CC3-IBD (f). The IDR exhibits robust DNA-binding activity, CC2 and TCD-CC3-IBD show weaker interactions, and CC3-IBD alone fails to bind DNA. These findings identify CC1, CC2, IDR, and the TCD as DNA-binding domains of MORC2.

(g) Confocal imaging showing that 601 DNA promotes MORC2 phase separation. At 0.9 μM, EGFP-MORC2FL alone does not form droplets, but the addition of 25 nM 601 DNA induces distinct condensate formation. Scaler bar: 5 μm.

(h) Quantification of MORC2 condensate sizes with or without 601 DNA. Violin plots illustrate the distribution of droplet sizes, with box plots indicating median values. Data are presented as mean ± SEM; unpaired student *t*-test. ****$p < 0.0001$.

(i) Fluorescence microscopy showing Cy3-labeled CTDΔCW (1.25 μM) coalescing with 25 nM FAM-labeled 601 DNA. DNA is recruited into MORC2 droplets, forming distinct condensates. Scaler bar: 5 μm.

(j) In the presence of 10 nM 601 DNA, the ATP hydrolysis activity of MORC2FL is significantly enhanced. The N39A mutant, which is deficient in ATP binding, serves as a negative control; nevertheless, a slight but detectable increase in ATPase activity is observed upon DNA binding. ATP hydrolysis activity of MORC2FL is significantly increased in the presence of 10 nM 601 DNA. The N39A mutant, which is deficient in ATP binding, serves as a negative control. Data are presented as mean ± SEM; statistical analysis: one-way ANOVA with Tukey's post hoc test. ****$p < 0.0001$.

Fig. 6 MORC2 phase separation is required for its transcriptional regulatory activity.

(a) Schematic of the rescue experiment strategy. HeLa cells with CRISPR-Cas9-mediated MORC2 knockout were reconstituted with EGFP, EGFP-MORC2FL, or phase separation-deficient mutants EGFP-MORC2ΔCC3 and EGFP-MORC2ΔIDR.

(b) RNA-seq-derived read counts for the MORC2 gene in EGFP, EGFP-MORC2FL, EGFP-MORC2ΔCC3, and EGFP-MORC2ΔIDR expressing cells.

(c-d) Venn diagram illustrating the overlap between differentially expressed genes identified following EGFP-MORC2FL overexpression in this study and previously characterized MORC2-regulated targets.

(e) Volcano plots of RNA-seq read counts for EGFP-MORC2FL and EGFP-MORC2ΔCC3 overexpression compared to EGFP control. Significantly upregulated genes are shown in yellow, and significantly downregulated genes are shown in green, based on three biological replicates (fold change > 1.2 or < 0.83, $p < 0.05$).

(f) LLPS-deficient mutant of MORC2 fails to regulate these 18 downregulated genes despite comparable protein expression levels, supporting a functional role of MORC2 condensates in transcriptional regulation.

Fig. 7 Pathogenic variants of MORC2 alter phase separation, DNA binding, and ATPase activity.

(a) Schematic illustration of pathogenic MORC2 mutations associated with Charcot-Marie-Tooth disease type 2Z (CMT2Z, red) and spinal muscular atrophy (SMA, blue), mapped onto the MORC2 domain structure.

(b) Quantification of HeLa cells transfected with WT MORC2 or nine pathogenic mutations exhibiting nuclear condensates was performed for each field of view. E236G and T424R shows the most pronounced enhancement of phase separation. n ≥ 10 fields. Data are presented as mean ± SEM; one-way ANOVA with Tukey's post hoc test. $****p < 0.0001$, $**p < 0.01$, *n.s.* not significant.

(c) ATPase activity of WT MORC2 and eight pathogenic variants, measured under revised conditions. S218L, F256L, and T424R exhibit significantly elevated activity. We were unable to purify MORC2 constructs bearing the E236G mutation from either HEK293F cells, suggesting that it may cause misfolding of the ATPase module. Data are presented as mean ± SEM; one-way ANOVA with Tukey's post hoc test. $****p < 0.0001$, $**p < 0.01$, *n.s.* not significant.

(d) Fluorescence polarization (FP) analysis of DNA binding affinities to the 601 DNA sequence. All protein variants were evaluated under uniform assay conditions, including those pertinent to phase separation. Dissociation constants (K_d) were determined by fitting the FP data to a log(agonist) vs. response model and are reported as mean ±SEM: WT (98 ± 1 nM), R252W (51 ± 2 nM), Q400R (145 ± 26 nM), D466N (49 ± 6 nM), S87L (107 ± 24 nM), S218L (92 ± 10 nM), F256L (69 ± 2 nM), R266A (63 ± 5 nM), and T424R (242 ± 13 nM). Several variants, including R252W, D466N, F256L, and R266A, demonstrated increased binding affinity relative to WT, while others, such as T424R and Q400R, exhibited markedly reduced binding.

(e) Proposed model summarizing the impact of DNA binding and phase separation on MORC2-mediated chromatin remodeling. DNA promotes MORC2 condensate formation through electrostatic interactions, which enhances its ability to compact chromatin and regulate transcriptional silencing.

Figure 1

1096
1097
1098 **Figure 2**
1099

Figure 3

1104

1105

1106   **Figure 4**

1107

1108

1109

1110   **Figure 5**

1111

1112

1113

1114

1115

Figure 6

1116

1117

1118   **Figure 7**

1119

1120 **Table S1**

1121

1122 **Statistics of X-ray Crystallographic Data Collection and Model refinement**

Data collection	
Data sets	MORC2-CC3
Space group	*I41*
Wavelength (Å)	0.97915
Unit Cell Parameters (Å)	a=130.253, b=130.253, c= 40.895
	α=β=γ=90°
Resolution range (Å)	50-3.05 (3.10-3.05)
No. of unique reflections	6774 (333)
Redundancy	12.6 (10.5)
I/σ	25.897 (2.600)
Completeness (%)	100.0 (100.0)
R_{merge} [a] (%)	12.0 (121.4)
$CC_{1/2}$ (last resolution shell) [b]	0.703

Structure refinement	
Resolution (Å)	32.56-3.10
R_{cryst} [c]/R_{free} [d] (%)	19.81 / 24.80
Rmsd bonds (Å) / angles (°)	0.004 / 0.627
Average B factor (Å^2) [e]	57.55
No. of atoms	
Protein atoms	1543
Water	6
Ligands	0
Ramachandran plot regions [e]	
Favored (%)	98.99
Allowed (%)	1.01
Outliers (%)	0

Numbers in parentheses represent the value for the highest resolution shell.

a. $R_{merge} = Σ |I_i - | / / ΣI_i$, where I_i is the intensity of measured reflection and is the mean intensity of all

symmetry-related reflections.

b. $CC_{1/2}$ were defined by Karplus and Diederichs.

c. $R_{cryst}=Σ||F_{calc}| − |F_{obs}||/ΣF_{obs}$, where F_{obs} and F_{calc} are observed and calculated structure factors.

d. $R_{free}= Σ_T||F_{calc}| − |F_{obs}||/ΣF_{obs}$, where T is a test data set of about 5% of the total unique reflections randomly

chosen and set aside prior to refinement.

e. B factors and Ramachandran plot statistics are calculated using MOLPROBITY.

1131

1133 *Table S2. Summary of MORC2 protein constructs used in this study, including expression systems,*
1134 *extinction coefficients, A260/280 ratios, and corresponding experimental applications.*

MORC2 fragment in vitro	*Expression system*	*Ext. coefficient*	*A260/280 ratios*	*Experimental applications*
MORC2FL (1-1032)	**E. coli**	*88240*	*0.59*	*EMSA*
CTD (473-1032)	*E. coli*	*46410*	*0.71*	*EMSA; LLPS*
CTDΔCW (537-1032)	*E. coli*	*26930*	*0.58*	*EMSA; LLPS; FPLC-MALS*
TCD-CC3-IBD (744-1032)	*E. coli*	*26930*	*0.67*	*EMSA; FPLC-MALS*
CC3-IBD (901-1032)	*E. coli*	*7450*	*0.51*	*EMSA; FPLC-MALS*
CC1 (279-362)	*E. coli*	*1490*	*0.53*	*EMSA*
CC2 (551-603)	*E. coli*	*0*	*ND*	*EMSA*
CC3 (901-1003)	*E. coli*	*5960*	*0.49*	*FPLC-MALS NMR; Structure determination*
IDR (593-735)	*E. coli*	*0*	*ND*	*EMSA*
IBD (1004-1032)	*E. coli*	*1490*	*0.58*	*NMR*
IDRa (593-643)	*E. coli*	*0*	*ND*	*NMR*
IDRb (644-694)	*E. coli*	*0*	*ND*	*NMR*
IDRc (695-735)	*E. coli*	*0*	*ND*	*NMR*
Trx-CC3	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3^{trun}_901-975	*E. coli*	*14105*	*0.58*	*FPLC-MALS*
Trx-CC3-IBD_L915Q	*E. coli*	*21555*	*0.91*	*FPLC-MALS*
Trx-CC3-IBD_F922Q	*E. coli*	*21555*	*0.59*	*FPLC-MALS*
Trx-CC3-IBD_L911Q	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3-IBD_I908Q	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3-IBD_F951Q	*E. coli*	*21555*	*0.54*	*FPLC-MALS*
Trx-CC3-IBD_Y921A	*E. coli*	*20065*	*0.56*	*FPLC-MALS*
Trx-CC3-IBD_F954Q	*E. coli*	*21555*	*0.53*	*FPLC-MALS*
Trx-CC3-IBD_L958Q	*E. coli*	*21555*	*0.58*	*FPLC-MALS*
MORC2FL (1-1032)	**HEK293F**	*88240*	*0.6*	*FPLC-MALS; LLPS; FP; ATPase activity*
MORC2ΔCC3-IBD (1-900)	*HEK293F*	*80790*	*0.56*	*FPLC-MALS*
MORC2_N39A	*HEK293F*	*88240*	*0.79*	*FP; ATPase activity*
MORC2_R252W	*HEK293F*	*93740*	*0.66*	*FP; ATPase activity*
MORC2_Q400R	*HEK293F*	*88240*	*0.65*	*FP; ATPase activity*
MORC2_D466N	*HEK293F*	*88240*	*0.58*	*FP; ATPase activity*
MORC2_S87L	*HEK293F*	*88240*	*0.75*	*FP; ATPase activity*
MORC2_S218L	*HEK293F*	*88240*	*0.72*	*FP; ATPase activity*
MORC2_F256L	*HEK293F*	*88240*	*0.6*	*FP; ATPase activity*
MORC2_R266A	*HEK293F*	*88240*	*0.75*	*FP; ATPase activity*
MORC2_T424R	*HEK293F*	*88240*	*0.55*	*FP; ATPase activity*

EGFP-MORC2FL	HEK293F	110130	0.59	LLPS; FRAP

Supplemental Figures

Fig. S1 Full-length MORC2 is a dimeric protein, with dimerization mediated by its C-terminal CC3 domain.

(a) Negative-stain electron microscopy revealed that MORC2FL particles purified from HEK293F cells appeared poorly resolved and exhibited considerable morphological heterogeneity. Scaler bar: 100 nm.

(b) Structural prediction of MORC2FL using AlphaFold3 under a two-copy modeling condition revealed the presence of an extended disordered region at the C-terminus and suggested a potential dimerization interface within this region.

(c) Summary of theoretical and measured molecular weight of MORC2 constructs, including MORC2FL, MORC2ΔCC3-IBD, CC3-IBD, CC3, and CC3 mutants. Data are derived from recombinant protein purification and static light-scattering (SLS) analysis.

(d) AlphaFold3-based modeling of MORC2 with the N-terminal domain removed revealed structural features of the C-terminus, including extensive disorder and a putative dimerization interface.

(e) Molecular model of the CC3 dimer highlights hydrophobic core residues (I908, L911, L915, F921, F922, F951, Y954). Key intermolecular distances include I908-I908' (3.4 Å), L911-L915' (3.7 Å), F922-F922' (3.9 Å) and L958-L958' (4.0 Å), supporting dimer stability.

(f) Molecular model of the CC3 dimer formation interface show the additional hydrophobic interactions.

(g) Point mutations in key residues (L911, L915, F922) affect dimerization. Additional mutants (I908Q, F951Q, Y921A, Y954A, L958Q) show no significant change in molecular weight compared to WT, confirming their minimal impact on dimer formation. SLS analysis shows molecular weight distributions, with WT CC3-IBD forming dimers (~ 60 kDa) as indicated by red curves, while mutants exhibit higher molecular weights indicative of aggregation. Constructs were tagged with a Trx tag (14 kDa) for solubility.

1167

Fig. S2 In vitro and in vivo evidence that MORC2 undergoes LLPS.

(a) Concentration gradients of EGFP-MORC2FL proteins (0.9, 1.8, 3.6, and 7.2 µM)

were analyzed for in vitro phase separation. The proteins were examined in a buffer

containing 150 mM NaCl, and phase separation was characterized by fluorescence

microscopy with 488 nm excitation for EGFP. Scaler bar: 2 µm.

(b) SDS-PAGE analysis of EGFP-MORC2FL in sedimentation-based assays. Protein

distribution between supernatant (S) and pellet (P) fractions is shown for increasing

concentrations (0.9, 1.8, 3.6, and 7.2 μM).

(c) Time-lapse imaging of EGFP-MORC2FL phase separation over 120 s highlights

dynamic droplet fusion. Scaler bar: 2 μm.

(d) Schematic of CRISPR-Cas9 strategy for generating MORC2 knockout (KO) HeLa

cells. sgRNA targeting sequences and protospacer adjacent motifs (PAM) are

highlighted. Genomic sequencing confirms insertion-deletion mutations introduced at

target sites in MORC2 allelic genes.

(e) Immunostaining images of WT and MORC2 KO HeLa cells validate the specificity

of the custom-made MORC2 antibody. Scaler bar: 5 μm.

(f) Generation of endogenous EGFP-MORC2 chimeric mice. Genomic sequencing

confirms insertion-EGFP mutations introduced at target sites in MORC2 allelic

genes.

(g) Live-cell imaging revealed that EGFP-MORC2FL exhibits dynamic subcellular

localization throughout mitosis, without forming detectable condensates. Scaler bar:

10 μm.

1190

1191

Fig. S3 The C-terminal domain (CTD) of MORC2 mediates nuclear condensate

formation.

(a) EGFP-tagged MORC2 CTD forms nuclear condensates in HeLa cells, whereas the

NTD does not. Scaler bar: 10 μm.

(b) Truncation analysis of EGFP-tagged CTD reveals that both the CC3-IBD and

IDR-NLS regions are important for condensate formation. Scaler bar: 10 μm.

(c) EGFP-tagged CTD constructs lacking the IDR-NLS region but fused to an exogenous

N-terminal nuclear localization signal ("PKKKRKV") successfully localize to the nucleus

but fail to form condensates, indicating that the IDR is essential for phase separation.

Scaler bar: 10 μm.

(d) Targeted expression of IDR-NLS and CC3-IBD fragments in the nucleus failed to

induce condensate formation, indicating that these domains alone are not sufficient to

drive phase separation. Scaler bar: 10 μm.

(e) The statistical data for droplet area in (Fig. 3a) demonstrate that the deletion of IDR

or CC3 is imperative for condensate formation. The data are expressed as the mean ±

standard error of the mean (SEM). n ≥ 15 fields; one-way ANOVA with Tukey's post hoc

test. ****P < 0.0001.

1209

1210

Fig. S4 The biophysical behavior of CTD and CTDΔCW in vitro.

(a) SDS-PAGE analysis of Cy3-labeled CTD showing expected size and degradative

behavior.

(b) Fluorescence microscopy images of Cy3-labeledCTD at increasing protein

concentrations (1.25 μM, 2.5 μM, 5 μM, and 10 μM) in standard phase

separation buffer (50 mM Tris-Cl, pH 8.0, 150 mM NaCl). Despite increasing

concentration, CW-CTD formed only small, dense puncta that did not exhibit robust

size expansion or morphological changes. Scaler bar: 10 μm.

(c) Representative bright-field and Cy3 fluorescence microscopy images of labeled

CTDΔCW protein droplets, showing intact condensate morphology under standard

phase separation conditions. Scaler bar: 5 μm.

(d) SDS-PAGE analysis comparing Cy3-labeled and unlabeled CTDΔCW, indicating

no degradation or mobility shift.

(e) Static light scattering (SLS) profiles of Cy3-labeled and unlabeled CTDΔCW

1225 *proteins reveal similar molecular weight distributions and hydrodynamic behavior.*

1226 *(f) Time-lapse imaging of Cy3-labeled CTDΔCW over 180 s reveals dynamic droplet*

1227 *behavior in vitro. Scaler bar: 2 μm.*

1228

1229
1230

Fig. S5 Improved visualization of MORC2 nuclear condensates in HEK293T cells.

(a) Representative single-cell images of EGFP-MORC2-expressing HEK293T cells,

reformatted from original Fig. 4i to enhance resolution and comparability with Fig. 4j.

Nuclear puncta are clearly visible at this scale. Scaler bar: 10 μm.

(b) Quantification of the percentage of cells showing visible nuclear condensates in

each construct. Data are presented as mean ± SEM; n ≥ 11 replicates. Statistical

significance was assessed using unpaired Student's t-test. ****p < 0.0001.

1238
1239

1240

Fig. S6 MORC2 phase separation regulates gene transcription.

(a) Abundance trajectories of all 240 regulated genes were normalized to their

respective mean values.

(b) GO-based cellular component analysis of the regulated genes mapped in DAVID

(v2025_1) knowledgebase. A substantial number of MORC2-regulated genes are

primarily localized to the nucleus, chromatin, and nucleoplasm.

(c) GO-based biological process analysis. MORC2-regulated genes are

predominantly involved in the biological process of transcriptional regulation.

(d) GO-based molecular function analysis. MORC2 primarily regulates genes

encoding transcription factors with transcriptional regulatory activity. Additionally,

MORC2 modulates the transcription of several cytokine genes.

(e) A selected panel of 12 genes that were regulated by MORC2 in this sequencing

context but lost responsiveness to the LLPS-deficient mutant was shown, including 8

downregulated and 4 upregulated genes.

1255

1256

Fig. S7 Representative SDS-PAGE gels for all purified protein constructs.

(a) MORC2 fragments used for conformational analysis by FPLC-MALS (related to

Fig. 1).

(b) MORC2 CC3 domain fragments used for dimerization analysis by FPLC-MALS

(related to Fig. S1).

(c) MORC2 fragments expressed in E. coli for EMSA-based DNA-binding assays

(related to Fig. 5).

(d) MORC2 fragments expressed in HEK293F cells for ATPase activity measurements

with or without 601 DNA (related to Fig. 5).

(e) Disease-associated MORC2 mutants expressed in HEK293F cells for ATPase and

DNA-binding assays,while E236G mutant exhibited poor biochemical properties,

precluding further analysis in vitro (related to Fig. 7).