MORC2 Mediates Transcriptional Regulation Through Liquid-Liquid Phase Separation

Yanshen Zhang[1,2,3,#], Weiya Xu[1,2,#], Wenxiu Duan[1,2,#], Yu Wei[1,2,3], Wenli Jiang[1,2,3], Feng Zhu[1], Chengdong Huang[1,2,*], Chao Wang[1,2,3*], and Yihui Bi[4*]

[1]Hefei National Research Center for Physical Sciences at the Microscale, Center for Advanced Interdisciplinary Science and Biomedicine of IHM, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[2]Ministry of Education Key Laboratory for Membraneless Organelles and Cellular Dynamics, Biomedical Sciences and Health Laboratory of Anhui Province, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[3]Department of Neurology, the First Affiliated Hospital of USTC, Division of Life Sciences and Medicine, University of Science and Technology of China, Hefei, China.

[4]Department of Orthopedics, The Second Affiliated Hospital of Anhui Medical University, Hefei, China.

Running title: Multivalent interactions and DNA binding underlie phase separation of MORC2

[#]These authors contributed equally to this work.

[*]Corresponding authors: Chengdong Huang (huangcd@ustc.edu.cn);

Chao Wang (cwangust@ustc.edu.cn);

Yihui Bi (yihuibi@ahmu.edu.cn).

Abstract

MORC2 is a chromatin-associated ATPase essential for transcriptional silencing and genome stability, yet the biophysical principles governing its regulatory activity remain elusive. Here, we demonstrate that full-length MORC2 undergoes biomolecular condensation to form dynamic nuclear assemblies, a process fundamentally required for its repressor function. Endogenous MORC2 forms discrete, dynamic condensates in neurons from *EGFP-MORC2* knock-in mice, supporting the physiological relevance of these assemblies *in vivo*. Mechanistically, a 3.1 Å crystal structure of coiled-coil 3 (CC3) identifies a dimeric scaffold that serves as a structural hub, while multivalent 'sticker' interactions between an intrinsically disordered region (IDR) and a newly defined IDR-binding domain (IBD) drive condensation. We show that DNA acts as a molecular scaffold that triggers MORC2 condensation, which in turn allosterically stimulates its ATPase activity. Critically, by employing a "killswitch" strategy to decouple assembly from internal fluidity, we reveal that only dynamic MORC2 condensates—not static aggregates or condensation-deficient mutants—can restore transcriptional repression in MORC2-knockout cells. Furthermore, pathogenic variants linked to CMT2Z and SMA differentially perturb these material properties and enzymatic turnover, providing a mechanistic link between condensate dysregulation and human neuropathies. Together, our findings establish a DNA-templated condensation mechanism for MORC2 and provide a molecular framework for understanding how the material state of chromatin-associated machinery dictates gene regulation and disease pathogenesis

Introduction

Chromatin dynamically regulates DNA accessibility to orchestrate essential processes such as transcription, replication, and DNA repair[1–5]. These processes are dynamically regulated by chromatin-associated complexes, such as the human silencing hub (HUSH) complex[6], which control the accessibility of genetic material to regulatory machinery. The Microrchidia (MORC) protein family, characterized by a conserved N-terminal GHL-ATPase domain and a structurally diverse C-terminal region, regulates gene expression and chromatin organization[7–10]. These C-terminal domains are thought to contribute to dimerization and functional specificity[11–14]. Notably, MORC2 has been identified as a component of the HUSH complex and is required for HUSH-mediated transcriptional repression[12].

GHL ATPases, including DNA gyrase, Hsp90, and MutL, rely on ATP-dependent dimerization to drive conformational changes essential for substrate engagement and enzymatic regulation[15–19]. Similar mechanisms have been proposed for MORC family members. In humans, the MORC family comprises five members—MORC1, MORC2, MORC3, MORC4, and SMCHD1, and different MORC paralogs target distinct genomic loci, including transposable elements, viral DNA, and long-exon genes[7,20,21]. The conserved N-terminal GHL-ATPase domain of MORC family proteins has been extensively characterized, whereas the evolutionarily diversified, multi-domain C-terminal regions are thought to act as key determinants of chromatin targeting and functional specificity. Despite these shared structural features, the molecular basis for MORC-mediated transcriptional regulation remains poorly understood, partly due to limited structural insight into the full-length proteins and C-terminal regions.

MORC2 has recently been identified as a key regulator of chromatin architecture and gene silencing, primarily through its recruitment to HUSH-target loci via C-terminal CC2 domain-mediated interactions with TASOR and MPP8[12]. While its silencing activity was initially attributed to the N-terminal ATPase domain, genetic complementation studies have revealed that the CW domain and all three coiled-coil (CC) regions are also indispensable for its function[12,13]. Beyond its structured domains,

sequence analyses and structural predictions have uncovered an extended intrinsically

disordered region (IDR) spanning the central to C-terminal regions of the protein,

suggesting potential regulatory mechanisms by facilitating multivalent and dynamic

interactions beyond enzymatic activity[22].

Transcriptional silencing is frequently spatially organized within membrane-less

nuclear compartments, often driven by the process of biomolecular condensation[23,24].

Sequence analyses of MORC2 reveal an extended intrinsically disordered region (IDR)

spanning the central to C-terminal domains, suggesting a capacity for the multivalent,

weak interactions that underpin phase separation[25,26]. While IDRs are common in

chromatin-associated proteins, their presence alone does not guarantee condensation;

rather, phase separation often emerges from the coordinated interplay between

disordered segments and structured oligomerization hubs. How such mesoscale

organization modulates MORC2's enzymatic turnover and its ability to remodel the

genome remains a fundamental question in the field.

In this study, we demonstrate that MORC2 undergoes phase separation to form

dynamic nuclear condensates. We define the structural basis for this behavior,

identifying a dimeric CC3 scaffold that, together with multivalent interactions

between the IDR and a newly identified IDR-binding domain (IBD), drives

condensation. We further show that DNA binding acts as a molecular scaffold to

promote MORC2 assembly, which in turn allosterically enhances its ATPase activity.

Using a combination of cellular "killswitch" assays and *in vivo* models, we establish

that the dynamic material properties of MORC2 condensates—rather than mere

protein recruitment—are strictly required for transcriptional regulation. Finally, we

provide mechanistic insights into how neuropathic disease-associated mutations in the

MORC2 N-terminus perturb this biophysical balance, linking condensate

dysregulation to human pathology.

Results

Coiled-coil 3 (CC3) mediates dimerization of full-length MORC2

To determine whether MORC2 has N- and C-terminal dimerization interfaces similar to those in other GHL-type ATPases[27], which previously identified the role of CC3 (previously designated as CC4) in MORC2 dimerization[14], we established an *in vitro* system to purify full-length (FL) human MORC2 (MORC2FL, residues 1-1032) from HEK293F cells, yielding nucleic acid-free, highly pure proteins for structural and functional analyses (Fig. 1a, b). Analytical size-exclusion chromatography coupled with static light scattering (SLS) demonstrated that MORC2FL forms dimers, with a molecular weight of approximately 239 ± 1 kDa under buffer conditions containing 1000 mM NaCl even in the absence of ATP (Fig. 1c). Notably, the addition of 601 DNA resulted in a detectable increase in molecular weight (303 ± 2 kDa), indicative of DNA-protein complex formation, but did not disrupt the dimeric assembly of MORC2 in the same buffer condition (Fig. 1c).

To elucidate the atomic structure of MORC2FL, we initially employed negative-stain electron microscopy to reveal that MORC2FL particles were poorly resolved and exhibited considerable heterogeneity in buffer containing 300 mM NaCl. Notably, a subset of particles displayed a morphology consistent with dimer formation (Fig. S1a). To further investigate this, we performed structural prediction of MORC2FL using AlphaFold3 and identified a predicted structure that closely resembled the observed reconstruction (Fig. S1b). We then examined the oligomerization properties of various MORC2 constructs. MORC2$^{1-900}$ (FLΔCC3-IBD domain) was found to exist as a monomer, whereas MORC2$^{901-1032}$ and MORC2$^{901-1003}$ exhibited dimeric properties, confirming the critical role of the CC3 domain (residues 901-1003) in mediating MORC2 dimerization (Fig. 1c, and Fig. S1c).

To gain further structural insights, we determined the 3.1 Å crystal structure of the CC3 dimer using X-ray diffraction (Table S1). The structure revealed that the dimer interface is stabilized predominantly by hydrophobic interactions. Three key

hydrophobic residues—L911, L915, and F922—form distinct layers within the dimer interface (layers 1-3, Fig. 1e, and Fig. S1e, f). The side chains of these residues are buried within the dimerization interface, creating a hydrophobic core critical for maintaining the stability of the dimer.

We systematically introduced single-point mutations in these hydrophobic residues to assess their contributions to dimerization. SLS analysis demonstrated that the wild-type (WT) MORC2$^{901-1032}$ (with Trx-tag) fragment predominantly exists as a dimer (~61 kDa). In contrast, all three key mutants (L911Q, L915Q, and F922Q) displayed altered molecular weight distributions indicative of disrupted dimer formation and enhanced higher-order oligomerization (more than 61 kDa) (Fig. S1g). However, none of these mutations completely disrupted dimerization or converted the dimer to a monomeric state (Fig. S1g). Further analyses of additional single-site mutants (e.g., I908Q, Y921A, F951Q, Y954A, L958Q) showed minimal effects on dimer stability, underscoring the central importance of L915 and its neighboring hydrophobic layers in stabilizing the CC3 dimer (Fig. S1g). Moreover, we examined the dimeric status of truncated CC3 constructs (residues 901-975), which confirmed that the CC3 N-terminal region is sufficient to drive dimer formation (Fig. 1f). To delineate the structural basis of these interactions, we mapped the positions of hydrophobic residues within the CC3 backbone (Fig. S1e). The distances between interacting residues, such as I908-I908' (3.4 Å), L911-L915' (3.7 Å), F922-F922' (3.9 Å), and L958-L958' (4.0 Å), revealed a tightly packed hydrophobic network. The robustness of this network ensures the integrity of the CC3 dimer, even under varying salt concentration and protein concentration tested *in vitro* conditions. These findings collectively suggest that the CC3 domain plays a critical role in MORC2 dimerization. The dimeric assembly of CC3 is essential for maintaining the structural integrity of the protein, which likely underpins its functional activities, including associations with chromatin[14].

MORC2 forms condensates both *in vivo* and *in vitro*

In the predicted structure of MORC2FL, we observed an extended unstructured

region at the C-terminus of the protein (Fig. S1b). This region is classified as an intrinsically disordered region (IDR), a protein architecture frequently reported to facilitate multivalent interactions and can act as a scaffold for phase separation[25, 28]. Specifically, IUPred2A analysis predicts that the residues 593-735 predominantly exhibit high disorder scores, prompting us to define this segment as the IDR domain (Fig. 1a).

In vitro, sedimentation assays showed that when the salt concentration was reduced to 150 mM, more than half of the EGFP fused MORC2FL (EGFP-MORC2FL) precipitated into the pellet fraction (Fig. 2a, b). Fluorescence imaging revealed that at this near-physiological ionic strength, EGFP-MORC2FL spontaneously demixed to form well-defined spherical droplets (Fig. 2c, d). These droplets exhibited a characteristic polydispersity in size and showed significant coarsening as protein concentration increased (Fig. S2a). Sedimentation assays further showed that the pellet fraction of EGFP-MORC2FL after centrifugation was concentration-dependent (Fig. S2b). The EGFP-MORC2FL-enriched condensates exhibited dynamic behavior, including droplet fusion (Fig. S2c) and fluorescence recovery after photobleaching (FRAP) (Fig. 2e, f), indicating a dynamic state.

Using antibody labeling, we observed that endogenous MORC2 forms discrete, dense punctate structures within the nuclei of HeLa cells (Fig. 2g). The specificity of these puncta was confirmed by their absence in MORC2-knockout (KO) HeLa cells (Fig. S2d, e), suggesting that MORC2 may natively organize into higher-order nuclear condensates under physiological conditions. To further characterize these assemblies, we transfected HeLa cells with EGFP-MORC2FL and observed the formation of prominent, droplet-like nuclear condensates (Fig. 2h). Notably, consistent with the endogenous staining patterns, EGFP-MORC2FL condensates preferentially localized to regions of reduced chromatin density, as visualized by the H2A-mCherry signal. This distinct spatial distribution implies a potential role for MORC2 condensates in organizing or regulating the heterochromatin landscape. FRAP analysis of these EGFP-MORC2FL puncta revealed fluorescence recovery on a timescale of seconds (Fig. 2i, j). Furthermore, these puncta underwent active fusion events (Fig. S2f), a

hallmark of dynamic liquid-like condensates with a highly mobile internal environment.

To determine whether MORC2 forms biomolecular condensates under physiological conditions, we generated mice expressing EGFP-MORC2 protein from the endogenous locus (Fig. S3a). We focus on neurons in the brain and spinal cord as missense mutations in *MORC2* cause neuropathies including Charcot-Marie-Tooth and spinal muscular atrophy disease[11,13]. Imaging of NeuN-positive neurons revealed that EGFP-MORC2FL condensates occur in neurons under endogenous expression level (Fig. 2k). FRAP analysis in freshly prepared brain slices further demonstrated rapid and partial fluorescence recovery, supporting the dynamic nature of endogenous MORC2 assemblies (Fig. 2l). In addition, unlike MORC3, which shows cell cycle-dependent nuclear puncta formation[29], MORC2 condensates were absent in mitotic cells (Fig. S3b). Together, these results indicate that MORC2 forms phase-separated biomolecular condensates under physiological conditions *in vivo* and *in vitro*.

The MORC2 C-terminal domain (CTD) drives condensate formation

To identify the key determinants governing the assembly of MORC2 nuclear condensates, we partitioned the protein into two functional modules based on its domain architecture: the N-terminal domain (NTD, residues 1–472), comprising the ATPase and CC1 motifs, and the C-terminal domain (CTD, residues 473–1032), which encompasses the IDR and multiple coiled-coil regions. A series of EGFP-tagged truncation constructs were then expressed in HeLa cells to systematically map the regions essential for condensate formation (Fig. 3a). Deletion of the CTD completely abolished puncta formation, resulting in a diffuse localization across the nucleus and cytoplasm (Fig. S4a). By contrast, NTD deletion had negligible effects on nuclear localization or the capacity for condensation. To exclude the possibility that the NTD's diffuse pattern was merely due to its loss of nuclear entry, we appended an exogenous nuclear localization signal (NLS, PKKKRKV[30]) to the NTD; however, while this redirected the protein to the nucleus, it failed to restore

puncta (Fig. 3a), confirming that the CTD is the primary driver of MORC2

condensation. To assess whether the CTD is intrinsically prone to undergo

condensation *in vitro*, we purified a CTDΔCW (residues 537–1032) variant from *E.*

coli, as the full CTD exhibited suboptimal biochemical stability (Fig. S5a, b). At 150

mM NaCl, Cy3-labeled CTDΔCW readily formed discrete condensates in a

concentration-dependent manner (Fig. 3b, c), and displayed a sedimentation profile

characteristic of phase-separated assemblies (Fig. 3d, e). Fluorescence microscopy

and static light scattering (SLS) confirmed that Cy3 labeling did not perturb its

condensation behavior, morphology, or molecular weight distribution (Fig. S5c–f).

Further dissection of the CTD revealed that deletion of either the CC2 or TCD

domains did not impair the formation of nuclear puncta (Fig. S4b). In contrast,

removal of the IDR-NLS region (residues 593-761) abolished both puncta formation

and nuclear localization (Fig. S4b). Notably, appending an exogenous NLS

(PKKKRKV) to CTDΔIDR-NLS restored its nuclear entry but failed to rescue

condensate assembly (Fig. S4c), identifying the IDR as a prerequisite for

condensation. Similarly, deletion of the CC3-IBD segment maintained nuclear

localization but eliminated puncta formation, indicating that this region is also

indispensable (Fig. S4b). Intriguingly, neither the IDR-NLS fragment nor the

CC3-IBD fragment alone (even when NLS-fused) was sufficient to trigger

condensation (Fig. S4d), implying a cooperative mechanism between these modules.

Consistent with these cellular findings, *in vitro* assays showed that neither the

MORC2ΔCC3-IBD (residues 1–900) nor the isolated CC3-IBD fragment could

undergo condensation (Fig. S6a-b). However, mixing the IDR with the IBD fragment

reconstituted robust condensate formation at higher concentration (Fig. S6c),

confirming their synergistic role. Finally, the CW domain appears to modulate

condensate morphology, as its deletion led to enlarged and frequently fused puncta.

We conclude that the IDR, CC3, and IBD constitute the minimal structural core

required for MORC2 condensation, while the CW domain serves to restrict

condensate overgrowth and maintain proper morphology (Fig. 3a, and Fig. S4e).

Multivalent interactions between the IDR and IBD promote MORC2 condensation

Next, we elucidated the mechanistic basis for MORC2 assembly mediated by the IDR and CC3-IBD. We hypothesized that multivalent interactions between these modules drive the formation of a condensed phase that coexists with a dilute aqueous phase. Using nuclear magnetic resonance (NMR) spectroscopy, we characterized the interaction between the IDR and the C-terminal segment (residues 901–1032). ^{15}N-HSQC titration assays using the ^{15}N-labeled CC3 (residues 901-1003) or the 1004-1032 fragment revealed that residues 1004-1032 constitute the primary binding site for the IDR, which we termed the IDR-Binding Domain (IBD), whereas the CC3 region exhibited negligible interaction (Fig. 4a-c).

Further subdivision of the IDR into three segments—IDRa (residues 593-643), IDRb (residues 644-694), and IDRc (residues 695-735)—pinpointed IDRa as the key region responsible for interaction with IBD (Fig. 4a, d-f). As is typical for IDR-mediated interactions involved in condensation, the observed chemical shift perturbations (CSPs) were relatively small but highly region-specific. Notably, the CSP profile of IDRa closely mirrors that of the full-length IDR, whereas IDRb and IDRc showed minimal perturbations (Fig. S7). Detailed sequence analysis revealed that the CSPs are clustered around charged residues (D, E, K, R) and their neighboring residues (Y, T, N, I, L, P). This suggests that the association is driven by a concerted action of multiple non-covalent forces: beyond the foundational electrostatic attraction, the involvement of aromatic (Y) and aliphatic (I, L, P) residues points to the contribution of π-π stacking, cation-π interactions, and hydrophobic effects, while polar residues (T, N) likely facilitate intermolecular hydrogen bonding. This hierarchical assembly, characterized by both charge complementarity and hydrophobic-like contributions, aligns with the 'sticker-and-spacer' model. In this framework, these multivalent residues act as 'stickers' that provide the requisite valency to scaffold a robust yet dynamic network, while the intervening sequences serve as 'spacers' to maintain the fluidity necessary for the condensed state (Fig. 4g). Collectively, these quantitative CSP data support a

model of weak, dynamic, and region-specific interactions, a hallmark of IDR-driven

condensation mechanisms[31–33].

To assess the role of IDRa-IBD multivalent interactions in cells, we transfected

HeLa and HEK293T cells with MORC2FL variants harboring individual deletions of

the IBD, IDRa, IDRb, or IDRc regions. Deletion of the IBD abolished condensate

formation in both HeLa cells (Fig. 3a, and Fig. S4e) and HEK293T cells (Fig. S8).

Similarly, deletion of IDRa significantly impaired condensation, resulting in a marked

decrease in both the number and area of nuclear puncta, whereas IDRb or IDRc

deletions had minimal impact (Fig. 4h, i). Based on these findings, we propose a

model in which multivalent 'sticker' interactions between IDRa and the IBD drive

MORC2 intermolecular assembly, a process further potentiated by CC3-mediated

dimerization, which collectively provides the valency threshold required for

condensation (Fig. 4j).

DNA binding promotes MORC2 condensation and stimulates ATPase activity

Notably, MORC2 condensates form exclusively within the nuclei of transfected

HeLa cells, implying that chromatin-associated factors—most likely DNA—serve as

essential cues for their nucleation or stabilization. While the N-terminal CC1 domain

was previously identified as a DNA-binding module[13], we hypothesized that the

full-length MORC2 architecture utilizes additional interfaces to facilitate complex

chromatin interactions.

To test this, we performed a comprehensive domain screen using electrophoretic

mobility shift assays (EMSA). Beyond confirming the robust DNA-binding activity of

the CC1 domain (Fig. 5a, b), our screen revealed that the CC2, IDR, and TCD (Tudor

Chromo-like domain) also possess significant DNA-binding capabilities (Fig. 5c-e).

In contrast, the CC3–IBD fragment showed no affinity for DNA (Fig. 5f). These

results indicate that MORC2 employs a multivalent DNA-binding architecture

distributed across both its N-terminal and C-terminal regions.

Building on this, we investigated whether these multivalent DNA-protein

interactions could scaffold MORC2 condensation *in vitro*. At 0.9 μM, MORC2FL

alone failed to form discrete assemblies; however, the addition of 25 nM 601 DNA

induced robust condensate formation, demonstrating that DNA acts as a molecular

seed to promote MORC2 phase separation (Fig. 5g, h). Interestingly, while the NTD

(residues 1–536) binds DNA, it remains diffuse (Fig. S6d,e), whereas the CTDΔCW

(residues 537–1032) undergoes DNA-driven condensation (Fig. 5i). Systematic

evaluation across a matrix of protein and DNA concentrations yielded a phase

diagram mapping the transitions between clear solutions, liquid-like condensates, and

amorphous precipitates, consistently showing that DNA significantly lowers the

concentration threshold for MORC2 condensation (Fig. S9). This functional

divergence suggests a division of labor within MORC2: while multiple domains

provide DNA affinity, the CTD provides the essential protein-protein interaction

network required for higher-order assembly.

Finally, we explored how this assembly impacts the MORC2 enzymatic core.

The addition of 601 DNA markedly stimulated the ATPase activity of wild-type

MORC2, but not the ATP-binding-deficient N39A mutant (Fig. 5j). This reveals a

synergistic interplay: DNA binding does not merely recruit MORC2 to chromatin; it

acts as a molecular scaffold that facilitates condensation, which in turn may optimize

the productive conformation or local density required for allosteric activation of the

ATPase domain. In conclusion, these findings highlight a coordinated mechanism that

integrates DNA binding, condensation, and enzymatic activity, providing a

biophysical basis for the multivalent assembly and allosteric regulation of MORC2 on

chromatin.

Condensate formation and dynamics are required for MORC2-mediated
transcriptional regulation

Establishing a causal link between biomolecular condensation and biological

function—particularly in transcriptional regulation—remains a central challenge. A

key question is whether MORC2 condensates represent discrete functional entities or

merely inactive byproducts. To address this, we sought to decouple condensate

formation from internal dynamics, allowing for a rigorous interrogation of their

functional relevance. While deletion of the CC3 domain abolishes MORC2

condensation, CC3 is also indispensable for dimerization, confounding functional

interpretations. Furthermore, although deleting IDRa markedly weakens MORC2

condensation, a tool to specifically manipulate the solubility and internal mobility of

these assemblies was previously lacking.

To overcome these limitations, we employed a micropeptide-based 'killswitch'

(KS) strategy fused to the C-terminus of MORC2 (Fig. 6a). This approach selectively

reduces condensate dynamics without disrupting protein expression, folding, or

domain architecture[34]. Unlike the ΔCC3 or ΔIDRa mutants, the MORC2+KS variant

robustly form nuclear condensates but exhibits markedly reduced internal dynamics,

as demonstrated by FRAP analyses showing minimal fluorescence recovery (Fig. 6b,

c). This strategy allowed us to perturb the material properties of the condensates

independently of structural integrity.

To evaluate how the modulation of MORC2 condensates influences its gene

regulatory capacity, we compared the transcriptional consequences of rescuing

MORC2-knockout HeLa cells with MORC2FL, condensation-deficient mutants

(ΔCC3 and ΔIDRa), and the dynamics-defective MORC2+KS (Fig. 6d). Bulk

RNA-seq was performed 48 hours post-transfection. We hypothesized that impairing

either the assembly or the fluidity of the condensates would compromise

MORC2-dependent gene regulation. Notably, despite being expressed at substantially

higher levels than MORC2FL (Fig. 6e), all three mutants showed a striking and

consistent failure to restore the MORC2-dependent transcriptional profile (Fig. 6f–h).

This defect was particularly pronounced for transcriptionally repressed genes,

including two sets of high-confidence MORC2 targets identified in prior studies[12, 22]

(Fig. 6i, and Fig. S10). Gene Ontology analysis revealed that the MORC2-regulated

genes are involved in DNA damage responses, viral infection, and cell cycle

regulation (Fig. 6g). These findings demonstrate that neither increased protein

abundance nor the mere presence of static, condensate-like structures is sufficient to

restore MORC2 function.

Collectively, these data support a model in which the dynamic nature of MORC2

condensates is essential for full transcriptional activity. While soluble MORC2
complexes are likely involved in initial target recognition, our results indicate that
proper condensate formation—and critically, internal fluidity—are required for
effective transcriptional control. The inability of the MORC2+KS mutant to rescue
transcriptional defects, despite forming visible nuclear assemblies, strongly argues
against a model where MORC2 condensates are merely inert byproducts of its
activity.

Pathogenic MORC2 variants differentially modulate condensation, DNA binding,
and ATPase activity

MORC2 is essential for gene silencing at H3K9me3-marked loci and
retrotransposon repression[6,35–37]. Pathogenic mutations in its GHL-type ATPase
domain have been linked to severe neuropathies and cancer[38–42]. To explore whether
these mutations influence the condensation behavior and biochemical activities of the
full-length protein, we examined a panel of disease-associated variants, including
those linked to Charcot–Marie–Tooth disease type 2Z (CMT2Z: E236G, R252W,
Q400R, D466N) and spinal muscular atrophy (SMA: S87L, S218L, F256L, R266A,
T424R) (Fig. 7a).

In HeLa cells, all variants retained the ability to form nuclear puncta. However,
the CMT2Z-linked E236G and, notably, the SMA-linked T424R mutants significantly
increased the proportion of cells exhibiting nuclear condensates (Fig. 7b). These
observations suggest that the N-terminal region plays a nuanced role in modulating
the phase boundary of MORC2. Specifically, previous structural studies indicated that
the T424R mutation has minimal effects on the overall N-terminal architecture but
primarily alters the assembly and dissociation kinetics of the N-terminal dimer[13]. This
suggests that the conformational cycling or dimerization state of the N-terminus
directly contributes to the macroscopic regulation of MORC2 condensates.
Interestingly, while deletion of the N-terminal region did not abolish puncta formation,
FRAP analyses revealed a marked loss of condensate fluidity compared with the
full-length protein (Fig. S11). This highlights the N-terminus as a critical regulator of

the material properties of MORC2 assemblies. Consequently, we evaluated the dynamics of E236G and T424R condensates; FRAP measurements showed substantially reduced recovery for E236G, indicating decreased internal fluidity, whereas T424R exhibited minimal effects (Fig. 7e, f).

In vitro ATPase assays revealed that T424R significantly increases enzymatic activity (Fig. 7c), consistent with its reported role in promoting ATPase dimer cycling[13]. Additionally, S218L and F256L variants enhanced ATP hydrolysis, whereas others, such as Q400R and D466N, showed activity comparable to WT (Fig. 7c). Due to severe solubility issues, E236G could not be biochemically characterized *in vitro* (Fig. S12e).

We next assessed DNA binding affinity of MORC2 variants using fluorescence polarization (FP) assays. Despite subtle differences, none of the variants showed statistically significant changes in DNA-binding capacity (Fig. 7d), indicating that these mutations primarily perturb enzymatic turnover and condensation dynamics rather than DNA recruitment.

In summary, our results support a model in which MORC2 mediates transcriptional regulation through its intrinsic ability to undergo biomolecular condensation. The N-terminal ATPase domain undergoes conformational cycling between open and closed states, while the C-terminal CC3-mediated dimer acts as a stable scaffold for condensate formation (Fig. 7g). MORC2 condensation is further fine-tuned by multivalent interactions between the IBD and IDRa regions, as well as by DNA binding. Together, these features enable MORC2 to bridge distal chromatin regions and establish repressive condensates, underscoring its central role in chromatin organization and genome stability

Discussion

Biophysical Principles of MORC2 Organization

While the MORC family of ATPases is recognized as a central player in epigenetic regulation[43], genome stability[44,45], and meiotic progression[46,47], the biophysical

principles governing their spatiotemporal organization have remained elusive. In this study, we establish a causal link between MORC2 condensation and its transcriptional regulatory function. By integrating high-resolution structural insights with functional assays, we reveal that MORC2 does not act merely as a solitary chromatin remodeler[3,5,48]; instead, it functions as phase separation-driven DNA compaction machinery. This discovery demonstrates how mesoscale material properties—specifically the fluidity and density of protein-DNA assemblies—are directly translated into precise gene regulation.

A Modular Engine for Condensate Assembly

The architectural basis of MORC2 assembly relies on a finely tuned multi-domain interplay. The C-terminal region, comprising the structured CC3-IBD and the intrinsically disordered IDR, provides the mechanistic engine for this process. CC3 serves as a critical structural scaffold (Fig. 1e, and 3a), mediating homodimerization that effectively doubles the molecular valency, thereby facilitating multivalent "sticker" interactions between the IDR and the defined IBD (Fig. 4). We found that mutations selectively abrogating condensate formation, or those that arrest condensate dynamics without blocking assembly, impair MORC2's ability to repress core target genes (Fig. 6). This underscores that high internal fluidity is functionally paramount, ensuring the rapid exchange kinetics required for continuous ATP hydrolysis and dynamic chromatin engagement.

The "Viscoelastic Window" and Molecular Stoichiometry

A defining insight from our study is that the mere presence of nuclear puncta is insufficient for MORC2's repressor activity. Instead, its function is governed by the precise material state of the condensates. We demonstrate that DNA acts as an essential functional scaffold (Fig. 5), driving condensation potentially mediated by nonspecific electrostatic interactions with the arginine/lysine-rich IDR. Crucially, these assemblies are exquisitely sensitive to molecular stoichiometry. This establishes a narrow, functionally permissive "viscoelastic window": Too Diffuse: MORC2 fails

to achieve the local density necessary for mechanical DNA compaction. Too Rigid: If

the system crosses the threshold into a solid-like, arrested state (as seen in the

MORC2 +KS mutant or E236G variant), it loses the fluidity required for enzymatic

turnover and co-repressor engagement. Within the nucleus, the CW domain likely

serves as a regulatory node, orchestrating the compositional regulation of these

condensates by reading specific epigenetic marks to fine-tune local valency and

morphology (Fig. 3 and Fig. S3).

Mechanistic Insights into Pathogenic Variants

Our analysis of disease-associated variants—particularly E236G (CMT2Z) and

T424R (SMA)—highlights how mutations in the ATPase domain can pleiotropically

affect both enzymatic turnover and phase behavior (Fig. 7). While T424R primarily

accelerates the ATPase dimer cycle[13], E236G triggers a transition toward a less

dynamic, "hardened" condensate state. These results emphasize that disease mutations

do not simply "turn off" the protein; they perturb the biophysical context of the

assembly. The observation that some variants exhibit elevated ATPase activity *in vitro*

suggests that in the native cellular environment, condensation-mediated local

enrichment may be required to properly couple ATP hydrolysis to productive

chromatin remodeling.

Our findings clarify previous discrepancies regarding MORC2's DNA binding. By

utilizing the 601 nucleosome-positioning sequence, we demonstrate a much more

robust interaction than previously captured with short, random dsDNA. Furthermore,

the use of full-length MORC2 reveals that C-terminal regulatory domains are

essential for maintaining enzymatic homeostasis—a feature masked in truncated

constructs. Whether MORC2-mediated silencing is achieved through direct

modulation of ATPase cycles or an independent mechanical compaction mechanism

remains an open question.

While our study establishes the essential role of MORC2 condensation in

transcriptional control, this exploration represents an initial step toward deciphering
the complex coordination between enzymatic turnover and phase behavior. Future
research is required to resolve how the ATP-dependent "DNA clamp" cycle is
spatially coupled to the physical compaction of chromatin, and whether additional
co-repressors are selectively partitioned into these dynamic assemblies to maintain
epigenetic stability.
506
507

Materials and Methods

Plasmid construction, cell culture and transfection

WT MORC2 (UniProt: Q9Y6X9) cDNA was amplified from a cDNA library derived from HeLa cells via reverse transcription. The MORC2 cDNA was then subcloned into the p3xFLAG-Myc-CMV-24 vector between BglII and KpnI restriction sites using seamless cloning. The seamless cloning process was facilitated by the ClonExpress II One-Step Cloning Kit (Vazyme, #C112-02) according to the manufacturer's instructions. An EGFP sequence was subsequently inserted into the BglII site at the N-terminal of MORC2 to generate the EGFP-MORC2 construct. To generate deletion mutant constructs of MORC2 for intracellular studies, specific regions of MORC2 were targeted and deleted using a polymerase chain reaction (PCR)-based approach. An 18- or 21 bp homologous sequence originating from one of the forward or reverse primers was introduced into the other primer, and the resulting PCR products were subjected to homologous recombination cloning, facilitated by the 18-bp homologous sequence. For protein expression and purification, MORC2FL and truncated MORC2 were cloned into prokaryotic expression vectors with MBP or Trx fusion proteins for bacterial expression.

In addition, MORC2FL and a series of truncation mutants were purified using a HEK293F eukaryotic expression system for *in vitro* phase separation and static light scattering (SLS) assays. A260/280 measurements, agarose gel analysis, or EMSA information on protein purifications and their experimental applications in this study is provided in Table S2. HeLa, HEK293T cells were cultured in DMEM medium (Biological Industries, DMEM, high re maintained in a humidified incubator at 37°C with 5% CO_2. Cell transfection was performed using Lipofectamine 3000 (Invitrogen) according to the manufacturer's instructions. Briefly, cells were seeded in appropriate culture plates the day before transfection and allowed to reach 40-60% confluency. The separately diluted Lipofectamine 3000 reagent and plasmids in Opti-MEM reduced serum medium (Gibco) were mixed, and the transfection complexes were

added to cells after 20 minutes. After a 6-hour incubation, the transfection medium

was replaced with fresh complete DMEM medium.

Recombinant protein expression, purification

Plasmids containing His-tagged genes were transformed into *E. coli* BL21 (DE3)

cells and cultured in Luria-Bertani medium with 50 mg/L ampicillin. When the

culture reached an optical density (OD) of 0.6-0.8 at 600 nm, the temperature was

lowered to 16°C. Isopropyl β-D-1-thiogalactopyranoside (IPTG) (0.25 mM) was

added to induce protein expression for 18-20 hours. Cells were harvested by

centrifugation at 13,000 × g for 15 minutes, then resuspended in Buffer A (50 mM

Tris-HCl, pH 8.0, 1000 mM NaCl, 10 mM imidazole, 1 mM dithiothreitol (DTT), and

1 mM phenylmethylsulfonyl fluoride (PMSF)). The cell suspension was snap-frozen

and stored at -80°C for subsequent protein purification. HEK293F suspension-adapted

cells were cultured in conical flasks at 37°C, 120 rpm, and 5% CO_2, until a density of

$1.0 \times 10\square$ cells/mL was reached. The cells were transiently transfected with 1 mg of

MORC2-6 × His plasmid using 2 mL of branched polyethyleneimine (1 mg/mL).

After 48 hours of incubation, cells were harvested by centrifugation at 3,000 × g for 5

minutes, resuspended in Buffer A, snap-frozen, and stored at -80 °C for further

purification.

All purification steps were performed at 4°C to preserve protein activity. Frozen

cell suspensions were rapidly thawed in water, lysed by sonication (200 W for ~0.3

hours), and centrifuged at 20,000 × g for 30 minutes at 4°C. The supernatant was

filtered through a 0.22 μm syringe filter to obtain soluble material. A His-Trap

Ni-NTA 6FF chromatography column was equilibrated with five column volumes of

Buffer A (Buffer A: 50 mM Tris-HCl, pH 8.0, 1000 mM NaCl). The clarified extract

was loaded onto the column, washed with Buffer A to remove non-specific proteins,

and eluted with a linear gradient of nickel elution buffer (Buffer B: 50 mM Tris-HCl,

pH 8.0, 1 M NaCl, 300 mM imidazole, 1 mM DTT). Eluted fractions were analyzed

by SDS-PAGE and Coomassie staining. The protein of interest was concentrated,

exchanged into Buffer C (50 mM Tris-HCl, pH 8.0, 1000 mM NaCl, 1 mM EDTA, 1

mM DTT) using an Amicon Ultra centrifugal filter unit, following overnight

treatment with Benzonase nuclease to eliminate nucleic acid contamination or HRV

3C Protease to cut His-tag. The purification process was subsequently refined through

gel filtration using a Superdex 6 Increase column (GE Healthcare) for the MORC2FL.

Following this, fragments were conjugated with a GPGS linker, and the resulting

mixture was subjected to a size-exclusion chromatography process (Superdex 200/75

column, GE Healthcare) to yield a final solution comprising 50 mM Tris-HCl, pH 8.0,

500 mM NaCl, 1 mM EDTA, and 1 mM DTT. Peak fractions were analyzed by

SDS-PAGE and Coomassie staining, confirming the successful purification of highly

homogeneous MORC2FL (1-1032) and various fragments (Fig. S7), then flash-frozen

in liquid nitrogen and stored at -80°C.

Crystallization, data collection and processing

The CC3 domain (901-1003) and various mutations of MORC2constructs were

cloned into a modified pET32M vector (with Trx-tag) using standard PCR-based

methods and confirmed by DNA sequencing. Due to the lack of homologous

structures, protein crystals with heavy atom derivatives are required to determine their

phases. Therefore, we used the methionine-deficient strain B834 to express the target

protein with selenomethionine (SeMet) in an inorganic selenosubstituted medium.

Recombinant proteins were expressed in *E. coli* B834 cells in M9 medium at 16°C.

MORC2 CC3 fused with a GPGS linker were finally changed by step of

size-exclusion chromatography (Superdex 75 increase column, GE Healthcare) into

50 mM Tris-HCl, pH 8.0, 100 mM NaCl, 1 mM EDTA and 1 mM DTT). The freshly

purified protein was concentrated to 10~20 mg/mL. Crystals were obtained by the

static sitting drop method in 0.1M Citric acid (pH 3.5) and 25% w/v PEG 3350 at

25°C. Glycerol (20%) was added as the cryo-protectant. A 3.1-Å resolution X-ray

dataset was collected at the beamline BL18U1 of the Shanghai Synchrotron Radiation

Facility (SSRF). The diffraction data were processed and scaled by HKL2000

(http://www.hkl-xray.com/). Using the structure of auto-build model by SAD data and

AlpahFold2 predicted model as the search model, the initial structural model was

solved by molecular replacement in PHASER (https://www.ccp4.ac.uk/). Further

manual model adjustment and refinement were completed iteratively using COOT and

PHENIX. The final structure was validated by PISA and MolProbity

(https://www.phenix-online.org/). The final refinement statistics of the complex

structure are listed in Supplementary Table 1. All structure figures were prepared

using the programme PyMOL (http://pymol.sourceforge.net/).

Protein labeling with fluorophore

The Cy3 NHS ester (AAT Bioquest) fluorophores were dissolved in DMSO at 10

mg/mL. Purified proteins were prepared in reaction buffer at a concentration of 10

mg/mL. The fluorophores were added to the protein solution at a 1:1 molar ratio. The

reaction was conducted with shaking at room temperature for 1 hour in the dark. After

the reaction, labeling was quenched with 200 mM Tris-HCl, pH 8.3, and the labeled

proteins were desalted using a HiTrap desalting column (GE Healthcare).

Fluorescence labeling efficiency was assessed using a Nanodrop 2000

spectrophotometer (ThermoFisher #ND-2000). The final labeling efficiency of each

protein was adjusted to 1% by mixing the labeled protein with an excess of unlabeled

protein (1:99 ratio).

Lentiviral mediated H2A-mCherry inducible expression in HeLa cells

Stable expression of H2A-mCherry in HeLa cells was achieved through

lentiviral-mediated transduction. First, H2A-mCherry gene sequence was inserted into

the inducible lentiviral vector pTRIPZ to generate the expression plasmid

pTRIPZ-H2A-mCherry. Lentiviral particles were generated by triple transfection of

HEK293T cells with pTRIPZ-H2A-mCherry, PMD2.G, and PSPAX2 plasmids using

Lipofectamine 3000. The viral supernatant was collected 48 hours after transfection,

filtered with 0.45-μm filter to remove cell debris, and used to transduce HeLa cells.

Following transduction, HeLa cells were selected with 50 μg/mL hygromycin to

establish stable cell lines. H2A-mCherry expression was induced with 1 μg/mL

doxycycline.

Live cell imaging and immunofluorescence staining

      For live-cell imaging, HeLa cells were seeded onto 35 mm culture dishes with glass-bottom plates (Biosharp) and transfected with EGFP-MORC2 or its mutant constructs. Images were captured 12 hours post-transfection using a Nikon confocal microscope equipped with an incubator system (37°C, 5% CO_2). Nuclei were stained with Hoechst 33342 or visualized using mCherry-labeled H2A fusion protein. For immunofluorescence, cells were fixed with 4% paraformaldehyde, permeabilized with 0.1% Triton X-100, and blocked with 10% goat serum. Primary antibodies were incubated overnight at 4°C, followed by incubation with fluorescence-conjugated secondary antibodies for 2 hours at room temperature in the dark. Fluorescence images were acquired using a Nikon confocal microscope.

CRISPR-Cas9 mediated MORC2 knockout cells

      MORC2 knockout (KO) HeLa cell lines were generated using the CRISPR-Cas9 system. A CRISPR single-guide RNA (sgRNA) targeting exon 5 of MORC2 (5'-ACACCTGAGTCTACTCAGAT-3') was designed using the online tool (http://crispor.tefor.net/). Two DNA oligos with complementary sgRNA targeting sequence and additional bases were designed as previously described[49,50]. These oligos were annealed and cloned into pLentiCRISPR v2 (Addgene plasmid #52961) to generate the MORC2 knockout vector. After transfection and puromycin selection, MORC2-deleted cells were obtained. Individual cells were sorted into separate wells of a culture plate to establish monoclonal cell lines using flow cytometry. To validate the knockout cell lines, genomic DNA was extracted from both WT and KO cells. The specific regions targeted by the sgRNA were amplified using PCR with primers: 5'-CAACATTCTCGAGCTGGACCTACAG-3' (forward) and 5'-CCACGACAAGACTGGAAACGTGACTC-3' (reverse). The PCR products were purified and subjected to Sanger sequencing, revealing the deletion and insertion mutations in the target sites of the MORC2 allelic genes (Fig. S2d).

Generation of endogenous EGFP-MORC2 chimeric mice

CRISPR-Cas9 was used to generate EGFP-MORC2 chimeric mice. The EGFP and linker sequence were inserted at the N-terminus of the mouse *MORC2a* gene. Briefly, a single-guide RNA (sgRNA) was designed to target exon 1 of *MORC2a*, with the sequence GACTGAAGACTCATTGCTGTCA. A donor DNA fragments containing the EGFP-linker region flanked by homologous arms was constructed. Fertilized mouse zygotes were collected, and a mixture of Cas9 mRNA (50 ng/μL), sgRNA (25 ng/μL), and donor DNA (50 ng/μL) was microinjected into pronucleus of the zygotes. The injected embryos were transferred into the oviducts of pseudopregnant recipient female mice. Resulting pups were genotyped by PCR and Sanger sequencing to confirm transgene integration (Fig. S2f). The genotyping primers for the EGFP-MORC2 allele were as follows: Left-F1: GCGGTGGTTGAGTTCCAATTCC; Left-R1: CTTCAGGGTCAGCTTGCCGTAG; Right-F1: GAACGGCATCAAGGTGAACTTC; Right-R1: TGTATCCAAGGTAATCTCTGTGGTA.

Mice were bred at the Animal Center of the University of Science and Technology of China (USTC). Animals were housed under controlled conditions (22 °C, 12-hour light/dark cycle) with free access to food and water ad libitum. All experimental procedures were approved by the Institutional Animal Care and Use Committee (IACUC) of the University of Science and Technology of China (Hefei, China).

Fluorescence recovery after photobleaching (FRAP)

The FRAP assay was performed to investigate the dynamics of protein mobility within condensates, either in *in vivo* or *in vitro*. Specific regions of interest (ROIs) approximately 2 μm² in size were selected and subjected to photobleaching using 100% laser power (488 nm) for approximately 20 ns. One droplet was left unbleached as a control. Fluorescence recovery within the ROI was monitored for 100-300 seconds. FRAP was performed using a 63x oil immersion objective mounted on a Nikon confocal microscope (in cells) or a Zeiss LSM 880 (*in vitro*). Fluorescence intensity

was normalized to the pre-bleach signal to analyze the recovery kinetics. For brain slice FRAP, organotypic brain slices were prepared from adult chimeric mice expressing endogenous EGFP-MORC2, following a slightly modified protocol previously described[49]. After decapitation, brains were rapidly extracted and placed in ice-cold dissection medium consisting of Hibernate A, 2% B27 supplement, 2 mM L-glutamine, and 1% penicillin–streptomycin. The cerebellum and midbrain were removed, and the remaining cerebral hemispheres were coronally sectioned at 250 µm using a McIlwain tissue chopper. Slices were gently separated in chilled dissection medium and transferred onto glass-bottom dishes containing culture medium composed of Neurobasal A, 2% B27 supplement, 2 mM L-glutamine, and 1% penicillin–streptomycin. Imaging was initiated immediately after brain slice preparation. FRAP experiments were performed using the Nikon confocal microscope. Photobleaching was carried out with 5-7 laser pulses (50 µs dwell time), and images were acquired at 2-s intervals. Post-bleach images were acquired 20 s after photobleaching.

Sedimentation assay

To assess concentration-dependent partitioning behavior, purified proteins at the indicated concentrations were incubated under the same buffer conditions used for *in vitro* phase separation assays and equilibrated for 10 min at room temperature. Samples were then centrifuged at $16,000 \times g$ for 10 min to separate the supernatant and pellet fractions. Following centrifugation, the supernatant was carefully collected, and the pellet was resuspended in an equal volume of buffer. Equal volumes of supernatant and pellet fractions were analyzed by SDS–PAGE and quantified by densitometry.

Electrophoretic mobility shift assays (EMSA)

Widom 601 DNA (147 bp, 25 nM) was amplified by PCR and incubated with increasing concentrations of MORC2 or its domain constructs in binding buffer (20 mM Tris-HCl, pH 7.5, 45 mM KCl, 4 mM $MgCl_2$, 5% glycerol, 5 mM DTT). The

reaction mixture was loaded onto a 6% polyacrylamide gel and electrophoresed at 120

V for 70 minutes. DNA was visualized using SYBR Gold stain, and images were

captured using the ChemiDoc MP Imaging System.

Size-exclusion chromatography coupled with static light scattering

The analysis was performed using an AKTA FPLC system (GE Healthcare)

coupled to a static light scattering detector (miniDawn, Wyatt) and a differential

refractive index detector (Optilab, Wyatt). Protein samples (70 μM) were filtered

and applied to a Superdex 200 increase or Superose 6 increase column,

pre-equilibrated with a column buffer containing 50 mM Tris-HCl, pH 8.0,

150 mM NaCl or 1000 mM NaCl (MORC2FL), 1 mM EDTA, and 1 mM DTT.

Data were analyzed using ASTRA6 software (Wyatt).

In vitro **phase separation and fluorescence microscopy imaging**

Droplet formation was monitored using differential interference contrast (DIC)

and fluorescence microscopy. To induce phase separation, protein was added to phase

separation buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM DTT). For

consistency, all MORC2 samples were diluted to a final concentration of 4 μM with

phase separation buffer containing 200 mM NaCl, unless otherwise specified. All

preparation steps were performed at 4 °C to preserve protein activity. The phase

separation mixtures were then equilibrated at room temperature for 20 min. A

5-10 μL aliquot of each sample was placed onto a custom-made glass slide, covered

with an 18 mm diameter coverslip, and incubated at room temperature for 20 min

before imaging.

Protein isotope labeling and NMR Spectroscopy

Isotopically labeled protein samples were expressed in minimal M9/H_2O or

M9/D_2O medium. Uniformly ^{15}N-labeled (or ^{13}C- or ^{15}N-labeled) protein samples

were prepared in M9/H_2O medium supplemented with 1 g/L of $^{15}NH_4Cl$ and 2 g/L

of glucose (or ^{13}C-glucose). The cells were harvested at an OD_{600} of approximately

1.0. Isotope-labeled precursors and amino acids were purchased from Cambridge Isotope Laboratories (CIL). Isotopically labeled samples were prepared in 50 mM Tris-HCl, pH 8.0, 100 mM NaCl, and 7% D₂O. Protein concentrations ranged from 0.5 to 1.0 mM. NMR experiments were performed using Bruker Avance III 600 MHz spectrometers equipped with a cryo-probe. NMR spectra of MORC2 and its complex were recorded at 298 K. Backbone assignments of MORC2[1004-1032] were obtained using SOFAST versions of standard triple-resonance experiments, including HNCACB and CBCACONH, recorded at 298 K. Residual assignment ambiguities were resolved using [15]N-edited HMQC-NOESY-HMQC spectra. All NMR spectra were processed using NMRPipe and analyzed with NMRViewJ software (http://www.onemoonscientific.com). CSPs were calculated in Hz at 600 MHz using the following equation:

$$\Delta v = ((\Delta^1 \mathrm{H} \times 600)^2 + (\Delta^{13}\mathrm{C} \times 150)^2)^{1/2}$$

Fluorescence polarization Assay (FP)

Fluorescence polarization (FP) assays were performed at room temperature (25 °C) using a BioTek H1 fluorescence spectrophotometer to evaluate the binding affinity between MORC2 protein and 5-carboxyfluorescein (5-FAM)-labeled 601 DNA (FAM-601) across a concentration range of 0-2000 nM. The 147-bp Widom 601 DNA fragment was generated via PCR amplification using a 5-FAM-labeled primer. The labeled primer was dissolved in 0.1 M NaHCO₃ buffer, pH 8.3, and the amplified product was purified by agarose gel extraction to eliminate residual primers and nonspecific amplification products.

Fluorescence measurements were carried out in assay buffer (50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 1 mM EDTA, 1 mM DTT, and 10% glycerol) containing purified FAM-601 DNA at a final concentration of 0.4 ng/μL. Excitation and emission wavelengths were optimized according to the instrument manual, with excitation set at 485 nm and emission at 525 nm. Fluorescence polarization values were recorded in triplicate, and control measurements were included to correct for background signals.

Data were analyzed to determine the equilibrium dissociation constant (K_d) for

the MORC2-FAM-601 interaction by fitting data to a log(agonist) vs. response model.

In parallel, the effect of small molecules on this binding interaction was assessed by

introducing compounds into the assay mixture and monitoring changes in

fluorescence polarization.

ATPase activity measurements

ATPase activity was measured[52] using 1 µM MORC2 or its mutant variants at

37°C in 50 mM Tris-HCl, pH 8.0, 150 mM NaCl, 8 mM $MgCl_2$, 10% glycerol and

5 mM ATP. All experiments were repeated independently at least three times.

RNA-seq and bioinformatic analysis

MORC2-knockout HeLa cells were seeded at equal density and transfected with

3 µg of plasmid DNA and 12 µg of PEI (polyethylenimine) per dish. After

6 hours, the transfection medium was replaced with fresh complete medium, and

cells were incubated for an additional 48 hours. Cells were then washed twice with

ice-cold PBS and lysed directly in RNA Isolator (R401, Vazyme) on ice. Total RNA

was extracted using the RNeasy Mini Kit (Qiagen, 74004) according to the

manufacturer's protocol. RNA quality was assessed using NanoDrop 2000 (Thermo

Fisher Scientific), Agilent 2100 Bioanalyzer, and LabChip GX (PerkinElmer). For

library preparation, 1 µg of total RNA was used per sample, employing the VAHTS

Universal V6 RNA-seq Library Prep Kit for MGI®. The resulting double-stranded

DNA libraries were denatured, circularized, and digested to obtain single-stranded

circular DNA. DNA nanoballs (DNBs) were generated from these templates through

rolling circle amplification (RCA). Sequencing was performed on the MGI

DNBSEQ-T7 platform (MGI Tech Co., Ltd) using combinatorial Probe-Anchor

Synthesis (cPAS) chemistry.

Before performing differential expression analysis, lowly expressed genes were

filtered out by retaining only those with a raw count of ≥10 in at least three samples

(baseMean). Differential gene expression analysis between sample groups was carried

out using the DESeq2 package (v1.26.0). Genes with *p-adjust* < 0.05 were defined as

differentially expressed genes (DEGs). These DEGs were subsequently subjected to

Gene Ontology (GO) enrichment analysis in DAVID (Knowledgebase v2025_1).

Statistical Analysis

All statistical analyses were performed in GraphPad Prism. All data underwent

normality testing (D'Agostino & Pearson test and Anderson-Darling test) in advance.

Two-tailed unpaired Student's *t*-test post-hoc Welch's correction and one-way

ANOVA with Tukey's post hoc test was used when data were distributed normally.

The significance level was set as $p < 0.05$. All statistical results are provided as

Source Data.

Acknowledgments

We thank Dr. Xuebiao Yao (University of Science and Technology of China,

USTC), Dr. Zhi Qi and Dr. Linyu Zuo (Peking University) for helpful discussion. We

thank members of the Shanghai Synchrotron Radiation Facility (SSRF, China;

beamlines BL02U1, BL18U1, and BL19U1) for X-ray beam time. This work was

supported by grants from the National Key R&D Program of China

(2019YFA0508402), the National Natural Science Foundation of China (22122703,

32470808, 32170767, 32471273, 31971144, and T2221005), the Center for Advanced

Interdisciplinary Science and Biomedicine of IHM (QYPY20220014), the Strategic

Priority Research Program of the Chinese Academy of Sciences (XDB0490000), the

Major Frontier Research Project of USTC (LS9100000002), USTC Research Funds

of Double First-Class Initiative (YD9100002507 and YD9100002054), the Anhui

Medical University Foundation (9101224201), and National Natural Science

Foundation Incubation Program of The Second Affiliated Hospital of Anhui Medical

University (2021GQFY02).

Author contributions

Y.Z., W.X., W.D., C.H., C.W., and Y.B. designed the experiments. Y.B., Y.Z.,

W.X., and W.D. conducted the experiments. Y.Z. determined the crystal structure.

Y.W., W.J. and F.Z. assisted with the experiments and reagents. All authors analyzed

the data. Y.Z., Y.B., C.H., and C.W. wrote the manuscript with input from all authors.

All authors approved the final version of the manuscript. C.H., and C.W supervised

the project.

Competing interests

842   The authors declare that they have no competing interests.

Data and materials availability

844   All data needed to evaluate the conclusions in the paper are present in the paper

845   and/or the Supplementary Materials. Raw gene expression data are available in

846   Source Data. The atomic coordinates of the MORC2 CC3 domain have been

847   deposited to the Protein Data Bank under the accession code: 9KQL.

References

1. Alabert, C. & Groth, A. Chromatin replication and epigenome maintenance. *Nat Rev Mol Cell Biol* **13**, 153–167 (2012).

2. Ishihara, S. *et al.* Local states of chromatin compaction at transcription start sites control transcription levels. *Nucleic Acids Res* **49**, 8007–8023 (2021).

3. Downs, J. A. & Gasser, S. M. Chromatin remodeling and spatial concerns in DNA double-strand break repair. *Curr Opin Cell Biol* **90**, 102405 (2024).

4. Blackledge, N. P. & Klose, R. J. The molecular principles of gene regulation by Polycomb repressive complexes. *Nat Rev Mol Cell Biol* **22**, 815–833 (2021).

5. Li, D.-Q. *et al.* MORC2 signaling integrates phosphorylation-dependent, ATPase-coupled chromatin remodeling during the DNA damage response. *Cell Rep* **2**, 1657–1669 (2012).

6. Müller, I. & Helin, K. Keep quiet: the HUSH complex in transcriptional silencing and disease. *Nat Struct Mol Biol* **31**, 11–22 (2024).

7. Zhong, Z. *et al.* MORC proteins regulate transcription factor binding by mediating chromatin compaction in active chromatin regions. *Genome Biol* **24**, 96 (2023).

8. Xue, Y. *et al.* Arabidopsis MORC proteins function in the efficient establishment of RNA directed DNA methylation. *Nat Commun* **12**, 4292 (2021).

9. Weiser, N. E. *et al.* MORC-1 Integrates Nuclear RNAi and Transgenerational Chromatin Architecture to Promote Germline Immortality. *Developmental Cell* **41**, 408-423.e7 (2017).

10. Moissiard, G. *et al.* MORC family ATPases required for heterochromatin condensation and gene silencing. *Science* **336**, 1448–1451 (2012).

11. Wang, H., Zhang, L., Luo, Q., Liu, J. & Wang, G. MORC protein family-related signature within human disease and cancer. *Cell Death Dis* **12**, 1112 (2021).

12. Tchasovnikarova, I. A. *et al.* Hyperactivation of HUSH complex function by Charcot-Marie-Tooth disease mutation in MORC2. *Nature Genetics* (2017) doi:10.1038/ng.3878.

13. Douse, C. H. *et al.* Neuropathic MORC2 mutations perturb GHKL ATPase dimerization dynamics and epigenetic silencing by multiple structural mechanisms. *Nat Commun* **9**, 651 (2018).

14. Xie, H.-Y., Zhang, T.-M., Hu, S.-Y., Shao, Z.-M. & Li, D.-Q. Dimerization of MORC2 through its C-terminal coiled-coil domain enhances chromatin dynamics and promotes DNA repair. *Cell Commun Signal* **17**, 160 (2019).

15. Morais Cabral, J. H. *et al.* Crystal structure of the breakage-reunion domain of DNA gyrase. *Nature* **388**, 903–906 (1997).

16. Chen, K. *et al.* Crystal structure of the hinge domain of Smchd1 reveals its dimerization mode and nucleic acid-binding residues. *Sci Signal* **13**, eaaz5599 (2020).

17. Liu, J. *et al.* Cascading MutS and MutL sliding clamps control DNA diffusion to activate mismatch repair. *Nature* **539**, 583–587 (2016).

18. Ortega, J., Lee, G. S., Gu, L., Yang, W. & Li, G.-M. Mispair-bound human MutS-MutL complex triggers DNA incisions and activates mismatch repair. *Cell Res* **31**, 542–553 (2021).

19. Lee, K. *et al.* The structure of an Hsp90-immunophilin complex reveals cochaperone recognition of the client maturation state. *Mol Cell* **81**, 3496-3508.e5 (2021).

20. Pastor, W. A. *et al.* MORC1 represses transposable elements in the mouse male germline. *Nat Commun* **5**, 5795 (2014).

21. Dopkins, N. *et al.* A field guide to endogenous retrovirus regulatory networks. *Mol Cell* **82**,

3763–3768 (2022).

22. Fendler, N. L. *et al.* Identification and characterization of a human MORC2 DNA binding region
that is required for gene silencing. *Nucleic Acids Res* **53**, gkae1273 (2025).

23. Leidescher, S. *et al.* Spatial organization of transcribed eukaryotic genes. *Nat Cell Biol* **24**,
327–339 (2022).

24. Membrane-less compartments in the nucleus: Separated or connected phases? *Current Opinion in*
Cell Biology **84**, 102215 (2023).

25. Boeynaems, S. *et al.* Protein Phase Separation: A New Phase in Cell Biology. *Trends Cell Biol* **28**,
420–435 (2018).

26. Yoshizawa, T. *et al.* Nuclear Import Receptor Inhibits Phase Separation of FUS through Binding to
Multiple Sites. *Cell* **173**, 693-705.e22 (2018).

27. Verba, K. A. *et al.* Atomic structure of Hsp90-Cdc37-Cdk4 reveals that Hsp90 traps and stabilizes
an unfolded kinase. *Science* **352**, 1542–1547 (2016).

28. Alberti, S., Gladfelter, A. & Mittag, T. Considerations and Challenges in Studying Liquid-Liquid
Phase Separation and Biomolecular Condensates. *Cell* **176**, 419–434 (2019).

29. Zhang, Y. *et al.* MORC3 Forms Nuclear Condensates through Phase Separation. *iScience* **17**,
182–189 (2019).

30. Kalderon, D., Roberts, B. L., Richardson, W. D. & Smith, A. E. A short amino acid sequence able
to specify nuclear location. *Cell* **39**, 499–509 (1984).

31. Alderson, T. R. & Kay, L. E. NMR spectroscopy captures the essential role of dynamics in
regulating biomolecular function. *Cell* **184**, 577–595 (2021).

32. Kim, T. H. *et al.* Phospho-dependent phase separation of FMRP and CAPRIN1 recapitulates
regulation of translation and deadenylation. *Science* **365**, 825–829 (2019).

33. Murthy, A. C. *et al.* Molecular interactions underlying liquid–liquid phase separation of the FUS
low-complexity domain. *Nat Struct Mol Biol* **26**, 637–648 (2019).

34. Zhang, Y., Stöppelkamp, I., Fernandez-Pernas, P. et al. Probing condensate microenvironments
with a micropeptide killswitch. *Nature* **643**, 1107–1116 (2025).

35. Pandiloski, N. *et al.* DNA methylation governs the sensitivity of repeats to restriction by the
HUSH-MORC2 corepressor. *Nat Commun* **15**, 7534 (2024).

36. Fukuda, K., Okuda, A., Yusa, K. & Shinkai, Y. A CRISPR knockout screen identifies
SETDB1-target retroelement silencing factors in embryonic stem cells. *Genome Res* **28**, 846–858
(2018).

37. Liu, N. *et al.* Selective silencing of euchromatic L1s revealed by genome-wide screens for
L1regulators. *Nature* **553**, 228–232 (2018).

38. Jacquier, A., Roubille, S., Lomonte, P. & Schaeffer, L. Microrchidia CW-Type Zinc Finger 2, a
Chromatin Modifier in a Spectrum of Peripheral Neuropathies. *Front Cell Neurosci* **16**, 896854
(2022).

39. Stafki, S. A. *et al.* The Spectrum of MORC2-Related Disorders: A Potential Link to Cockayne
Syndrome. *Pediatr Neurol* **141**, 79–86 (2023).

40. Chung, H. Y. *et al.* Morc2a variants cause hydroxyl radical-mediated neuropathy and are rescued
by restoring GHKL ATPase. *Brain* **147**, 2114–2127 (2024).

41. Ding, Q.-S. *et al.* Aberrant high expression level of MORC2 is a common character in multiple
cancers. *Hum Pathol* **76**, 58–67 (2018).

42. Zhang, S. *et al.* Oncogenic MORC2 in cancer development and beyond. *Genes Dis* **11**, 861–873

(2024).

43. Li, D.-Q., Nair, S. S. & Kumar, R. The MORC family: new epigenetic regulators of transcription and DNA damage response. *Epigenetics* **8**, 685–693 (2013).

44. Koch, A. *et al.* MORC Proteins: Novel Players in Plant and Animal Health. *Front Plant Sci* **8**, 1720 (2017).

45. Dong, W. *et al.* MORC Domain Definition and Evolutionary Analysis of the MORC Gene Family in Green Plants. *Genome Biol Evol* **10**, 1730–1744 (2018).

46. Inoue, N. *et al.* New gene family defined by MORC, a nuclear protein required for mouse spermatogenesis. *Hum Mol Genet* **8**, 1201–1207 (1999).

47. Wang, G.-L. *et al.* Identification and expression analysis of a novel CW-type zinc finger protein MORC2 in cancer cells. *Anat Rec (Hoboken)* **293**, 1002–1009 (2010).

48. Tan, W., Park, J., Venugopal, H. et al. MORC2 is a phosphorylation-dependent DNA compaction machine. Nat Commun 16, 5606 (2025).

49. Shalem, O. *et al.* Genome-scale CRISPR-Cas9 knockout screening in human cells. *Science* **343**, 84–87 (2014).

50. Sanjana, N. E., Shalem, O. & Zhang, F. Improved vectors and genome-wide libraries for CRISPR screening. *Nat Methods* **11**, 783–784 (2014).

51. Li, C. H. *et al.* MeCP2 links heterochromatin condensates and neurodevelopmental disease. *Nature* **586**, 440–444 (2020).

52. Microtiter assay for glutamine synthetase biosynthetic activity using inorganic phosphate detection. *Analytical Biochemistry* **327**, 114–118 (2004).

958 **Table S1**

960 **Statistics of X-ray Crystallographic Data Collection and Model refinement**

Data collection	
Data sets	MORC2-CC3
Space group	*I41*
Wavelength (Å)	0.97915
Unit Cell Parameters (Å)	a=130.253, b=130.253, c= 40.895
	α=β=γ=90°
Resolution range (Å)	50-3.05 (3.10-3.05)
No. of unique reflections	6774 (333)
Redundancy	12.6 (10.5)
I/σ	25.897 (2.600)
Completeness (%)	100.0 (100.0)
R_{merge} [a] (%)	12.0 (121.4)
$CC_{1/2}$ (last resolution shell) [b]	0.703

Structure refinement	
Resolution (Å)	32.56-3.10
R_{cryst} [c]/R_{free} [d] (%)	19.81 / 24.80
Rmsd bonds (Å) / angles (°)	0.004 / 0.627
Average B factor (Å^2) [e]	57.55
No. of atoms	
Protein atoms	1543
Water	6
Ligands	0
Ramachandran plot regions [e]	
Favored (%)	98.99
Allowed (%)	1.01
Outliers (%)	0

Numbers in parentheses represent the value for the highest resolution shell.

(**a**) R_{merge} = Σ $|I_i$ - $|$ / $ΣI_i$, where I_i is the intensity of measured reflection and   is the mean intensity of all
symmetry-related reflections.

(**b**) $CC_{1/2}$ were defined by Karplus and Diederichs.

(**c**) R_{cryst}=Σ$||F_{calc}| − |F_{obs}||/ΣF_{obs}$, where F_{obs} and F_{calc} are observed and calculated structure factors.

(**d**) R_{free}= $Σ_T||F_{calc}| − |F_{obs}||/ΣF_{obs}$, where T is a test data set of about 5% of the total unique reflections randomly
chosen and set aside prior to refinement.

(**e**) B factors and Ramachandran plot statistics are calculated using MOLPROBITY.

Table S2. *Summary of MORC2 protein constructs used in this study, including expression systems, extinction coefficients, A260/280 ratios, and corresponding experimental applications.*

MORC2 fragment in vitro	*Expression system*	*Ext. coefficient*	*A260/280 ratios*	*Experimental applications*
MORC2FL (1-1032)	**E. coli**	*88240*	*0.59*	*EMSA, Agarose gel analysis (AGA)*
CTD (473-1032)	*E. coli*	*46410*	*0.71*	*EMSA; Phase separation*
CTDΔCW (537-1032)	*E. coli*	*26930*	*0.58*	*EMSA; Phase separation; FPLC-MALS*
TCD-CC3-IBD (744-1032)	*E. coli*	*26930*	*0.67*	*EMSA; FPLC-MALS*
CC3-IBD (901-1032)	*E. coli*	*7450*	*0.51*	*EMSA; FPLC-MALS, AGA*
CC1 (279-362)	*E. coli*	*1490*	*0.53*	*EMSA*
CC2 (551-603)	*E. coli*	*0*	*ND*	*EMSA*
CC3 (901-1003)	*E. coli*	*5960*	*0.49*	*FPLC-MALS NMR; Structure determination, AGA*
IDR (593-735)	*E. coli*	*0*	*ND*	*EMSA*
IBD (1004-1032)	*E. coli*	*1490*	*0.58*	*NMR*
IDRa (593-643)	*E. coli*	*0*	*ND*	*NMR*
IDRb (644-694)	*E. coli*	*0*	*ND*	*NMR*
IDRc (695-735)	*E. coli*	*0*	*ND*	*NMR*
Trx-CC3	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3^{trun}_901-975	*E. coli*	*14105*	*0.58*	*FPLC-MALS*
Trx-CC3-IBD_L915Q	*E. coli*	*21555*	*0.91*	*FPLC-MALS*
Trx-CC3-IBD_F922Q	*E. coli*	*21555*	*0.59*	*FPLC-MALS*
Trx-CC3-IBD_L911Q	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3-IBD_I908Q	*E. coli*	*21555*	*0.6*	*FPLC-MALS*
Trx-CC3-IBD_F951Q	*E. coli*	*21555*	*0.54*	*FPLC-MALS*
Trx-CC3-IBD_Y921A	*E. coli*	*20065*	*0.56*	*FPLC-MALS*
Trx-CC3-IBD_F954Q	*E. coli*	*21555*	*0.53*	*FPLC-MALS*
Trx-CC3-IBD_L958Q	*E. coli*	*21555*	*0.58*	*FPLC-MALS*
MORC2FL (1-1032)	**HEK293F**	*88240*	*0.6*	*FPLC-MALS; Phase separation; FP; ATPase activity, AGA*
MORC2ΔCC3-IBD (1-900)	*HEK293F*	*80790*	*0.56*	*FPLC-MALS*
MORC2_N39A	*HEK293F*	*88240*	*0.79*	*FP; ATPase activity*
MORC2_R252W	*HEK293F*	*93740*	*0.66*	*FP; ATPase activity*
MORC2_Q400R	*HEK293F*	*88240*	*0.65*	*FP; ATPase activity*
MORC2_D466N	*HEK293F*	*88240*	*0.58*	*FP; ATPase activity*
MORC2_S87L	*HEK293F*	*88240*	*0.75*	*FP; ATPase activity*

MORC2_S218L	HEK293F	88240	0.72	FP; ATPase activity
MORC2_F256L	HEK293F	88240	0.6	FP; ATPase activity
MORC2_R266A	HEK293F	88240	0.75	FP; ATPase activity
MORC2_T424R	HEK293F	88240	0.55	FP; ATPase activity
EGFP-MORC2FL	HEK293F	110130	0.59	Phase separation; FRAP, AGA

Figure Legend

Fig. 1 The CC3 domain serves as the structural basis for full-length MORC2 dimerization.

(*a*) *The domain organization of human MORC2FL, highlighting the ATPase module (GHL, CC1 and S5*

domains, collectively termed NTD), the C-terminal domains (CTD) with coiled-coil regions (CC2 and

CC3), CW-type zinc finger (CW), intrinsically disordered region (IDR, marked by a fresh green line by

IUPRED2 prediction), nuclear localization signal (NLS), Tudor-chromodomain (TCD), and a

30-residues tail domain (IBD).

(*b*) *Purification and characterization of full-length human MORC2 in HEK293F cell system.*

SDS-PAGE analysis of the purified protein demonstrates its purity and distribution.

(*c*) *Static light-scattering (SLS) analysis shows that MORC2 forms dimers in the absence of ATP, with a*

molecular weight of 239 ± 1 kDa in standard buffer (mint green line) and 303 ± 2 kDa in the presence

of 601 DNA (sunflower yellow line).

(*d*) *SLS analysis reveals the oligomerization states of MORC2 truncations. Full-length MORC2 lacking*

CC3-IBD (1-900) is monomeric, while CC3-IBD (901-1032) and CC3 alone (901-1003) form stable

dimers.

(*e*) *Crystal structural of the CC3 dimer. Hydrophobic residues contributing to dimer stability (L911,*

L915, and F922) are highlighted using ball-and-stick representations. Top view of the hydrophobic core

of the CC3 dimer. Layers of hydrophobic contacts stabilize the dimer interface, illustrated with paired

residues (e.g., Layer 1: L911-L911', Layer 2: L915-L915', and Layer 3: F922-F922').

(*f*) *Additional SLS analysis confirms that fragments spanning residues 537-1032, 744-1032, and*

Trx-901-975 (CC3 truncation) adopt dimeric conformations.

Fig. 2 Endogenous MORC2 organizes into dynamic condensates within the nucleus

(*a*) *SDS-PAGE analysis of EGFP-MORC2FL protein purified from HEK293F cells, showing protein*

concentration at 2.5 μM and 5 μM.

(*b*) *Sedimentation assay shows the distribution of EGFP-MORC2FL (7.2 μM) between supernatant (S)*

and pellet (P) fractions under 1000 mM (control) and 150 mM NaCl, indicating its propensity for

phase separation.

(*c*) *EGFP-MORC2 (7.2 μM) undergoes phase separation into droplets in a buffer containing 150 mM*

NaCl, but not in 1000 mM NaCl buffer. Scaler bar: 2 μm.

(*d*) *The proteins were examined in a buffer containing 150 mM NaCl, and phase separation was*

assessed by fluorescence microscopy with 488 nm excitation for EGFP. Quantification of droplet areas

*from (*Fig. S2a*), presented as violin plots with box plots indicating medians. Data are derived from ≥ 3*

independent images and reflect the size distribution of phase-separated droplets. Data are presented as

*mean ± SEM; one-way ANOVA with Tukey's post hoc test. ****p < 0.0001; *p < 0.05.*

(*e*) *Time-lapse FRAP imaging of EGFP-MORC2FL droplets formed in vitro, demonstrating*

fluorescence recovery over time. Scaler bar: 2 μm.

(*f*) *Quantitative analysis of fluorescence recovery from (*e*), showing the dynamic properties of in vitro*

MORC2 droplets. Data represent mean ± SEM, with n ≥ 3 replicates.

(*g*) *Immunostaining of endogenous MORC2 in HeLa cells revealed punctate localized within the*

nucleus, likely representing condensates involved in transcriptional regulation (white arrows). Nuclei

are counterstained with DAPI to visualize chromatin-rich regions. Scaler bar: 5 μm.

(*h*) *Live-cell imaging of transiently transfected EGFP-MORC2FL showed its assembly into dispersed,*

nearly spherical condensates in the nucleus, while stable expression of H2A-mCherry served to mark
chromatin-enriched domains. Scaler bar: 10 μm.
*(**i**) Time-lapse FRAP analysis of EGFP-MORC2FL condensates in the nuclei of transiently transfected*
HeLa cells revealed rapid fluorescence recovery within seconds, indicative of their dynamic, nature.
Scaler bar: 1 μm.
*(**j**) Quantitative analysis of fluorescence recovery from (**i**), showing the dynamic properties of in vitro*
MORC2 droplets. Data represent mean ± SEM, with n ≥ 10 replicates.
*(**k**) Sections of brain and spinal cord from endogenous EGFP-MORC2 chimeric mice show*
EGFP-MORC2 condensation distribution in NeuN-positive neurons. Scaler bar: 10 μm.
*(**l**) Representative FRAP images from of FRAP experiments performed on fresh 250-μm brain slices of*
endogenous EGFP-MORC2 chimeric mice demonstrate the dynamic nature of endogenous MORC2
condensates. Data are presented as mean ± SEM, with n = 3 droplets analyzed. Scaler bar: 5 μm.
1029
Fig. 3 Multiple domains regulate phase separation of MORC2.
*(**a**) Sequential deletion analysis revealed that the NTD is dispensable for condensate formation, as*
EGFP-tagged NTD alone failed to form condensates and was excluded from the nucleus. Fusion with a
canonical nuclear localization signal ("PKKKRKV") restored nuclear localization but did not rescue
condensate formation. In contrast, the CTD alone was sufficient to form nuclear condensates. Deletion
of the CW domain resulted in enlarged condensates in a subset of cells, while removal of CC2 or the
TCD had no apparent effect on condensate formation. Strikingly, deletion of either the IDR or CC3
completely abolished nuclear condensate assembly. Deletion of the IBD markedly reduced the
frequency of condensate formation. Loss of the intrinsic NLS led to cytoplasmic localization of MORC2;
however, no condensates were detected in the cytoplasm under transient transfection conditions in
HeLa cells. Together, these results define the IDR and CC3 as the minimal and essential elements for
MORC2-mediated phase separation, and implicate the CW domain, IBD, and nuclear
microenvironment in the fine-tuning of condensate assembly. Scaler bar: 10 μm.
*(**b**) In vitro phase separation of Cy3-labeled CTDΔCW at concentrations from 1.25 μM to 20 μM in 150*
mM NaCl buffer, visualized as spherical condensates. Scaler bar: 10 μm.
*(**c**) Quantification of droplet areas from (b), displayed as violin plots with box plots indicating median*
*values. Data are presented as mean ± SEM; one-way ANOVA with Tukey's post hoc test. ****p <*
0.0001, n.s. not significant.
*(**d**) SDS-PAGE analysis of Cy3-CTDΔCW protein distribution between supernatant (S) and pellet (P)*
fractions after centrifugation at increasing concentrations (5 μM to 40 μM) in 150 mM NaCl buffer. At
40 μM, increasing the NaCl concentration to 300 mM or 500 mM reduced droplet formation, indicating
salt sensitivity.
*(**e**) Quantification of Cy3-CTDΔCW showing the percentage of soluble protein from densitometric*
*analysis of S/P fractions in panel (**d**), based on three independent biological replicates.*
1054
Fig. 4 Multivalent interactions between IDR and IBD drive MORC2 condensation.
*(**a**) Domain organization of MORC2 C-terminal regions analyzed by NMR titration, including the IDR,*
CC3, and IBD.
*(**b**) ^{15}N-labeled CC3 (residues 901–1003) shows no chemical shift upon titration with IDR (residues*
593–735), indicating no detectable interaction.
*(**c**) ^{15}N-labeled IBD (residues 1004–1032) exhibits clear chemical shift perturbations (CSPs) upon IDR*

titration, confirming a direct, specific interaction.

*(**d-f**) NMR titrations of ^{15}N-labeled IBD with IDR subregions: IDRa (593–643), IDRb (644–694), and*
*IDRc (695–735). IDRa induces clear CSPs (**d**), while IDRb (**e**) and IDRc (**f**) show minimal or no shifts,*
identifying IDRa as the primary IBD-binding segment.

*(**g**) Sequence analysis of IDR subregions reveals that IDRa is enriched in proline and arginine residues,*
enabling electrostatic interactions. Sequence alignment of IBD across species highlights conserved
residues involved in IDRa binding. A summary table lists representative residues from IDRa and IBD
exhibiting CSPs in titration assays (marked with asterisks).

*(**h**) Representative confocal images of HeLa cells expressing EGFP-tagged MORC2ΔIDRa, ΔIDRb, or*
ΔIDRc constructs. Scale bar: 10 μm.

*(**i**) Quantification of droplet area per cell in (h). Deletion of IDRa significantly impairs condensate*
formation. n = 10 cells per condition. Data are presented as mean ± SEM; statistical analysis: one-way
ANOVA with Tukey's post hoc test. ***p < 0.001; n.s. not significant.

*(**j**) Working model of MORC2 condensates. CC3 dimerization serves as a structural scaffold, while*
weak, transient, but specific multivalent interactions between IDRa and IBD cooperatively promote
condensate formation.

1077

Fig. 5 DNA binding enhances MORC2 condensation and modulates its ATPase activity.

*(**a**) Electrophoretic mobility shift assay (EMSA) showing that MORC2FL binds 25 nM 601 DNA, as*
evidenced by shifts in DNA mobility corresponding to protein-DNA complexes.

*(**b**) EMSA highlighting the strong DNA-binding affinity of the CC1 domain, corroborating previous*
reports that identify this region as a key mediator of DNA interaction.

*(**c-f**) EMSA results demonstrating the binding capacities of individual MORC2 domains to 25 nM 601*
*DNA: CC2 (**c**), IDR (**d**), TCD-CC3-IBD (**e**), and CC3-IBD (**f**). The IDR exhibits robust DNA-binding*
activity, CC2 and TCD-CC3-IBD show weaker interactions, and CC3-IBD alone fails to bind DNA.
These findings identify CC1, CC2, IDR, and the TCD as DNA-binding domains of MORC2.

*(**g**) Confocal imaging showing that 601 DNA promotes MORC2 phase separation. At 0.9 μM,*
EGFP-MORC2FL alone does not form droplets, but the addition of 25 nM 601 DNA induces distinct
condensate formation. Scaler bar: 5 μm.

*(**h**) Quantification of MORC2 condensate sizes with or without 601 DNA. Violin plots illustrate the*
distribution of droplet sizes, with box plots indicating median values. Data are presented as mean ±
SEM; unpaired student t-test. ****p < 0.0001.

*(**i**) Fluorescence microscopy showing Cy3-labeled CTDΔCW (1.25 μM) coalescing with 25 nM*
FAM-labeled 601 DNA. DNA is recruited into MORC2 droplets, forming distinct condensates. Scaler
bar: 5 μm.

*(**j**) In the presence of 10 nM 601 DNA, the ATP hydrolysis activity of MORC2FL is significantly*
enhanced. The N39A mutant, which is deficient in ATP binding, serves as a negative control;
nevertheless, a slight but detectable increase in ATPase activity is observed upon DNA binding. ATP
hydrolysis activity of MORC2FL is significantly increased in the presence of 10 nM 601 DNA. The
N39A mutant, which is deficient in ATP binding, serves as a negative control. Data are presented as
mean ± SEM; statistical analysis: one-way ANOVA with Tukey's post hoc test. ****p < 0.0001.

1102

Fig. 6 Dynamic MORC2 condensation is required for its transcriptional regulatory activity.

*(**a**) Schematic diagram showing the fusion of two 17–amino acid killswitch sequences to the C terminus*

of MORC2.

*(**b**) Live-cell FRAP images of HeLa cells overexpressing EGFP–MORC2 full length (FL) and*
EGFP–MORC2–Killswitch (+KS). Scaler bar: 1 µm.

*(**c**) FRAP analysis of EGFP–MORC2 condensates in the nucleus. Data represent mean ± SEM, with n*
= 11 replicates.

*(**d**) Schematic of the rescue experiment strategy. HeLa cells with CRISPR-Cas9-mediated MORC2*
knockout were reconstituted with EGFP, EGFP-MORC2FL, or phase separation-deficient mutants
EGFP-MORC2ΔCC3 and EGFP-MORC2ΔIDR, and dynamics-defective +KS mutant.

*(**e**) RNA-seq-derived read counts for the MORC2 gene in EGFP, EGFP-MORC2FL,*
EGFP-MORC2ΔCC3, EGFP-MORC2ΔIDR, and EGFP-MORC2FL+KS expressing cells.

*(**f**) The effect of MORC2 rescue on the transcriptome. MA plot showing all significantly differentially*
expressed genes between EGFP–MORC2FL and EGFP transfection in MORC2-knockout HeLa cells (n
= 1,074; DESeq2 adjusted p < 0.05), highlighted in red.

*(**g**) GO-based biological process terms enriched among all differentially expressed genes. Data*
represent mean ± SD, with n = 3 replicates.

*(**h**) Abundance trajectories of all regulated genes, normalized to their respective mean values. Shown is*
a selected panel of ten downregulated genes that are influenced by MORC2 rescue but show reduced
responsiveness to all three mutants.

*(**i**) Venn diagram illustrating the overlap between differentially expressed genes identified following*
EGFP-MORC2FL overexpression in this study and previously characterized MORC2-regulated targets.
Fold-change (log₂ fold change) analysis of the 84 overlapping genes showing significant expression
changes upon MORC2FL overexpression, compared with their corresponding fold changes following
overexpression of the three MORC2 mutants. Dashed lines connect the same gene across the four
transfection conditions.

1129

Fig. 7 Pathogenic variants of MORC2 alter condensates behavior, DNA binding, and ATPase
activity.

*(**a**) Schematic illustration of pathogenic MORC2 mutations associated with Charcot-Marie-Tooth*
disease type 2Z (CMT2Z, red) and spinal muscular atrophy (SMA, blue), mapped onto the MORC2
domain structure.

*(**b**) Quantification of HeLa cells transfected with WT MORC2 or nine pathogenic mutations exhibiting*
nuclear condensates was performed for each field of view. E236G and T424R shows the most
pronounced enhancement of phase separation. n ≥ 10 fields. Data are presented as mean ± SEM;
*one-way ANOVA with Tukey's post hoc test. ****p < 0.0001, **p < 0.01, n.s. not significant.*

*(**c**) ATPase activity of WT MORC2 and eight pathogenic variants, measured under revised conditions.*
S218L, F256L, and T424R exhibit significantly elevated activity. We were unable to purify MORC2
constructs bearing the E236G mutation from either HEK293F cells, suggesting that it may cause
misfolding of the ATPase module. Data are presented as mean ± SEM; one-way ANOVA with Tukey's
*post hoc test. ****p < 0.0001, **p < 0.01, n.s. not significant.*

*(**d**) Fluorescence polarization (FP) analysis of DNA binding affinities to the 601 DNA sequence. All*
protein variants were evaluated under uniform assay conditions, including those pertinent to phase
separation. Dissociation constants (K_d) were determined by fitting the FP data to a log(agonist) vs.
response model and are reported as mean ±SEM: WT (98 ± 1 nM), R252W (51 ± 2 nM), Q400R (145 ±
26 nM), D466N (49 ± 6 nM), S87L (107 ± 24 nM), S218L (92 ± 10 nM), F256L (69 ± 2 nM), R266A (63

± 5 nM), and T424R (242 ± 13 nM). Several variants, including R252W, D466N, F256L, and R266A, demonstrated increased binding affinity relative to WT, while others, such as T424R and Q400R, exhibited markedly reduced binding.

(*e*). Live-cell FRAP images of HeLa cells overexpressing EGFP–MORC2 full length (FL), E236G, and T424R mutants. Scaler bar: 1 μm.

(*f*). FRAP analysis of EGFP–MORC2 condensates in the nucleus. Data represent mean ± SEM, with n ≥ 32 replicates.

(*g*) A Mechanistic Model of MORC2-Mediated Chromatin Condensation. MORC2 functions as a multivalent DNA-binding machine. The C-terminal CC3-IBD and IDR serve as the structural engine for phase separation, while DNA acts as a scaffold that lowers the phase boundary. The N-terminal ATPase domain undergoes conformational cycling (Open/Closed), providing the dynamic fluidity necessary for the assembly to remain within a functional 'viscoelastic window'. In this window, MORC2 achieves the local density required for DNA compaction and gene silencing

Supplemental Figures

Fig. S1 Full-length MORC2 is a dimeric protein, with dimerization mediated by its C-terminal CC3 domain.

(*a*) *Negative-stain electron microscopy revealed that MORC2FL particles purified from HEK293F cells appeared poorly resolved and exhibited considerable morphological heterogeneity. And a zoomed-in particle was to comparing morphology from negative stain EM with AlphaFold predicted structure. Scaler bar: 100 nm and 10 nm.*

(*b*) *Structural prediction of MORC2FL using AlphaFold3 under a two-copy modeling condition revealed the presence of an extended disordered region at the C-terminus and suggested a potential dimerization interface within this region.*

(*c*) *Summary of theoretical and measured molecular weight of MORC2 constructs, including MORC2FL, MORC2ΔCC3-IBD, CC3-IBD, CC3, and CC3 mutants. Data are derived from recombinant protein purification and static light-scattering (SLS) analysis.*

(*d*) *AlphaFold3-based modeling of MORC2 with the N-terminal domain removed revealed structural features of the C-terminus, including extensive disorder and a putative dimerization interface.*

(*e*) *Molecular model of the CC3 dimer highlights hydrophobic core residues (I908, L911, L915, F921, F922, F951, Y954). Key intermolecular distances include I908-I908' (3.4 Å), L911-L915' (3.7 Å), F922-F922' (3.9 Å) and L958-L958' (4.0 Å), supporting dimer stability.*

(*f*) *Molecular model of the CC3 dimer formation interface show the additional hydrophobic interactions.*

(*g*) *Point mutations in key residues (L911, L915, F922) affect dimerization. Additional mutants (I908Q, F951Q, Y921A, Y954A, L958Q) show no significant change in molecular weight compared to WT, confirming their minimal impact on dimer formation. SLS analysis shows molecular weight distributions, with WT CC3-IBD forming dimers (~ 60 kDa) as indicated by red curves, while mutants exhibit higher molecular weights indicative of aggregation. Constructs were tagged with a Trx tag (14 kDa) for solubility.*

Fig. S2 In vitro and in vivo evidence that MORC2 undergoes phase separation.

(*a*) *Concentration gradients of EGFP-MORC2FL proteins (0.9, 1.8, 3.6, and 7.2 μM) were analyzed for in vitro phase separation. The proteins were examined in a buffer containing 150 mM NaCl, and phase separation was characterized by fluorescence microscopy with 488 nm excitation for EGFP. Scaler bar: 2 μm.*

(*b*) *SDS-PAGE analysis of EGFP-MORC2FL in sedimentation-based assays. Protein distribution between supernatant (S) and pellet (P) fractions is shown for increasing concentrations (0.9, 1.8, 3.6, and 7.2 μM).*

(*c*) *Time-lapse imaging of EGFP-MORC2FL phase separation over 120 s highlights dynamic droplet fusion. Scaler bar: 2 μm.*

(*d*) *Schematic of CRISPR-Cas9 strategy for generating MORC2 knockout (KO) HeLa cells. sgRNA targeting sequences and protospacer adjacent motifs (PAM) are highlighted. Genomic sequencing confirms insertion-deletion mutations introduced at target sites in MORC2 allelic genes.*

(*e*) *Immunostaining images of WT and MORC2 KO HeLa cells validate the specificity of the custom-made MORC2 antibody. Scaler bar: 5 μm.*

(*f*) *The fusion of condensates of EGFP-MORC2 over expression in HeLa cell.*

1207

1208 *Fig. S3 Genomic loci of MORC2a in chimeric mice and its subcellular localization during mitosis.*

1209 (*a*) *Generation of endogenous EGFP-MORC2 chimeric mice. Genomic sequencing confirms*
1210 *insertion-EGFP mutations introduced at target sites in MORC2 allelic genes.*

1211 (*b*) *Live-cell imaging revealed that EGFP-MORC2FL exhibits dynamic subcellular localization*
1212 *throughout mitosis, without forming detectable condensates. Scaler bar: 10 μm.*

1213

Fig. S4 The C-terminal domain (CTD) of MORC2 mediates nuclear condensate formation.

(*a*) *EGFP-tagged MORC2 CTD forms nuclear condensates in HeLa cells, whereas the NTD does not. Scaler*
bar: 10 μm.

(*b*) *Truncation analysis of EGFP-tagged CTD reveals that both the CC3-IBD and IDR-NLS regions are*
important for condensate formation. Scaler bar: 10 μm.

(*c*) *EGFP-tagged CTD constructs lacking the IDR-NLS region but fused to an exogenous N-terminal nuclear*
localization signal ("PKKKRKV") successfully localize to the nucleus but fail to form condensates,
indicating that the IDR is essential for phase separation. Scaler bar: 10 μm.

(*d*) *Targeted expression of IDR-NLS and CC3-IBD fragments in the nucleus failed to induce condensate*
formation, indicating that these domains alone are not sufficient to drive phase separation. Scaler bar: 10
μm.

(*e*) *The statistical data for droplet area in (Fig. 3a) demonstrate that the deletion of IDR or CC3 is*
imperative for condensate formation. The data are expressed as the mean ± standard error of the mean
*(SEM). n ≥ 15 fields; one-way ANOVA with Tukey's post hoc test. ****P < 0.0001.*

1228

Fig. S5 The biophysical behavior of CTD and CTDΔCW in vitro.

(*a*) *SDS-PAGE analysis of Cy3-labeled CTD showing expected size and degradative behavior.*

(*b*) *Fluorescence microscopy images of Cy3-labeledCTD at increasing protein concentrations*
(1.25 μM, 2.5 μM, 5 μM, and 10 μM) in standard phase separation buffer (50 mM Tris-Cl, pH
8.0, 150 mM NaCl). Despite increasing concentration, CW-CTD formed only small, dense puncta that
did not exhibit robust size expansion or morphological changes. Scaler bar: 10 μm.

(*c*) *Representative bright-field and Cy3 fluorescence microscopy images of labeled CTDΔCW protein*
droplets, showing intact condensate morphology under standard phase separation conditions. Scaler
bar: 5 μm.

(*d*) *SDS-PAGE analysis comparing Cy3-labeled and unlabeled CTDΔCW, indicating no degradation or*
mobility shift.

(*e*) *Static light scattering (SLS) profiles of Cy3-labeled and unlabeled CTDΔCW proteins reveal similar*
molecular weight distributions and hydrodynamic behavior.

(*f*) *Time-lapse imaging of Cy3-labeled CTDΔCW over 180 s reveals dynamic droplet behavior in vitro.*
Scaler bar: 2 μm.

1244

1245 *Fig. S6 In vitro assays to investigate isolated fragment fails to induce phase separation.*

1246 (*a*) *Sedimentation assay showing domain containing 1-900 (5μM) and 1-900 (5μM) is not induced*
1247 *phase separation in 100 mM NaCl buffer.*

1248 (*b*) *Representative images showing that Cy3-labeled MORC2 1-900 (5μM) could not undergo phase*
1249 *separation.*

(*c*) In vitro assays to investigate the contribution of IDR-IBD interactions to phase separation.

(*d*) Sedimentation assay showing domain containing 1-536 (10 μM) and 537-1032 (10 μM) is not induced phase separation in 100 mM NaCl buffer.

(*e*) Representative images showing that Cy3-labeled MORC2 1-536 (10μM) could not undergo phase separation in including 50 nM 601 DNA.

Fig. S7 NMR titration about IDR-IBD. *The corresponding CSP plots in the process of NMR titration related to* Fig. 4

Fig. S8 Phase diagram of MORC2-DNA.

DNA-dependent phase behavior of the protein in vitro. Phase diagram showing the formation of liquid droplets and amorphous structures as a function of DNA and protein concentrations. Each circle represents an individual experimental condition. Black regions indicate regimes favoring liquid droplet formation, whereas the darker shaded region at high DNA and protein concentrations denotes the formation of amorphous assemblies. At low protein concentration (0.5 μM) and low DNA levels, no phase separation was observed in shaded regions.

Fig. S9 IBD-deficient mutant (ΔIBD) in HEK293T cells

(a) Representative single-cell images of EGFP-MORC2-expressing HEK293T cells. Scaler bar: 10 μm.

(b) Quantification of the percentage of cells showing visible nuclear condensates in each construct. Data are presented as mean ± SEM; $n \geq 11$ replicates. Statistical significance was assessed using unpaired Student's t-test. ****$p < 0.0001$.

Fig. S10 Condensation and dynamics-defective MORC2 mutants impair transcriptional regulation

(**a**) GO-based molecular function terms enriched among all differentially expressed genes upon EGFP-MORC2FL overexpression.

(**b**) Venn diagram illustrating the overlap between differentially expressed genes identified in this study and previously characterized targets regulated in MORC2 knockout HeLa cells[12]. Fold-change (log□ fold change) analysis of the 102 overlapping genes showing significant expression changes upon MORC2FL overexpression, compared with their corresponding fold changes following overexpression of the three MORC2 mutants. Dashed lines connect the same gene across the four transfection conditions.

Fig. S11 Compared to the wild-type protein, N-terminally truncated MORC2-CTD forms nuclear puncta in HeLa cells with significantly reduced internal dynamics, as revealed by FRAP assays.

Fig. S12 Representative SDS-PAGE gels for all purified protein constructs.

(**a**) MORC2 fragments used for conformational analysis by FPLC-MALS (related to Fig. 1).

(**b**) MORC2 CC3 domain fragments used for dimerization analysis by FPLC-MALS (related to Fig. S1).

(**c**) MORC2 fragments expressed in E. coli for EMSA-based DNA-binding assays (related to Fig. 5).

(**d**) MORC2 fragments expressed in HEK293F cells for ATPase activity measurements with or without 601 DNA (related to Fig. 5).

1292　　*(e) Disease-associated MORC2 mutants expressed in HEK293F cells for ATPase and DNA-binding*

1293　　*assays, while E236G mutant exhibited poor biochemical properties, precluding further analysis in vitro*

1294　　*(related to Fig. 7).*

1295　　*(f) Agarose gel analysis for MORC2 constructs to confirm the absence of nucleic acids. The pET32*

1296　　*vector as the positive control, the protein preparation for analysis is 0.05 mg.*

a HEK 293F system

b Spin down assay — supernatant (S) / pellet (P)

c EGFP-FL-hMORC2: 1000 mM NaCl; 150 mM NaCl

d EGFP-FL-hMORC2 — Droplet area (μm)² vs Concentration (μM): 0.9, 1.8, 3.6, 7.2

e Time-lapse FRAP imaging of EGFP-FL-hMORC2 droplets — Pre-bleaching, 0 s, 60 s, 90 s, 120 s

f Normalized intensity vs Time (s): Unbleaching, Bleaching

g endogenous MORC2 (antibody detected) — MORC2, DAPI, Merge (Hela)

h EGFP-MORC2 over-expression — EGFP-MORC2, H2A-mCherry, Merge (Hela)

i EGFP-MORC2 over-expression — Pre-bleaching, 0.2 s, 1.6 s, 3.0 s, 4.4 s, 5.8 s, 7.2 s

j Normalized intensity vs Time (s): Unbleaching, Bleaching

k endogenous GFP-MORC2 chimeric mice — EGFP-MORC2, DAPI, EGFP-MORC2/DAPI, NeuN, Merge (Brain, Spinal cord)

l Pre-bleaching, Bleaching, Post-bleaching — Normalized intensity vs Time (s): Unbleaching, Bleaching

a Negative-staining electron microscopy

MORC2FL

10 nm

100 nm

b Structure of MORC2FL prediction (AF3, 2 copies)

C (1032)   C' (1032)

90°

c

Protein	Theroretical MW (kDa)	Measured MW (kDa)	Dimerization
MORC2FL	119.8	239.0 ± 1.0	Dimer
MORC2ΔCC3-IBD	102.5	103.6 ± 0.5	Monomer
CC3-IBD	15.3	29.2 ± 0.2	Dimer
CC3	12.1	23.8 ± 0.4	Dimer
CTDΔCW	56.0	110.3 ± 0.2	Dimer
TCD-CC3-IBD	33.2	63.1 ± 0.5	Dimer
Trx (tag)-CC3-IBD	29.3	61.7 ± 0.2	Dimer
Trx (tag)-truncated CC3 (901-975)	22.7	45.7 ± 0.4	Dimer

d Structure of CTD prediction

N (604)   N' (604)

C' (1032)   C (1032)

e

N   N (904)

C (1000)   C

3.4 Å   Layer 0
I908'   I908

L911'   3.7 Å   *Layer 1*
L911

L915'   L915   *Layer 2*

Y951'   F922' F922   F951   *Layer 3*
Y921   Y921'
3.9 Å   Layer 4
Y954'   Y954

L958'   L958   Layer 5
4.0 Å

f

I908'   I908

F951
Y921'

F922'
Y954

L958'   L958

g

901 □□ 1032 (WT)   901 □□ 1032 (F922Q)
901 □□ 1032 (L915Q)   901 □□ 1032 (L911Q)

113.2 ± 0.2 kDa
79.8 ± 0.4 kDa
61.7 ± 0.2 kDa

I908Q   Dimer

F951Q   Dimer

Y921A   Dimer

L958Q   Dimer

F954Q   Dimer

a

0.9 μM   1.8 μM   3.6 μM   7.2 μM

EGFP-MORC2FL

2 μm

b

Mr (kDa)   0.9 μM   1.8 μM   3.6 μM   7.2 μM
            S   P   S   P   S   P   S   P

180 –
130 –

c

EGFP-MORC2FL

0 s   10 s   30 s   60 s   90 s   120 s

2 μm

d

humman *MORC2* genome loci   exon 4   exon 5

18965   Target   PAM   19011

TCGGCCAAGCGA**ACACCTGAGTCTACTCAGAT****TGG**GCAGTACGGGAA

Wild Type   AAGGCTCAGACCTTTGACT // AGCGA**ACACCTGAGTCTACTCAGAT****TGG**GCAGT

Mutant #1   AAGGCTCAGACCT- - - - - - // -GGGCAGT

Mutant #2   AAGGCTCAGACCTTTGACT // AGCGA**ACACCTGAGTCTACTCAG**A**T****TGG**GCAGT
                                                                    TACGGGA
                                                                    Insert

e

MORC2 Knockout Validation

MORC2   DAPI

HeLa

5 μm

f

0.00 s   0.63 s   1.26 s   1.89s   2.52 s   3.15 s   3.78 s   4.41 s   5.04 s

5.67 s   6.3 s   6.93 s   7.56 s   8.19 s   8.82 s   9.45 s   10.08 s   10.71 s

1 μm

EGFP-MORC2 over-expression

a

mouse *MORC2a* genome loci

exon 1 (306-1316)   exon 2 (9541-9594)   41346 bp

1224   1316

CAAA GACTGAAGACTCATTGCTGTCA TGG CCTTCACCAATTACAGCAGTCTCAACCGAGCTCAACTAACCTTTGAATATCTCCACACAAATTC

Target   PAM

Met Ala Phe Thr Asn Tyr Ser Ser Leu Asn Arg Ala Gln Leu Thr Phe Glu Tyr Leu His Thr Asn

EGFP-Linker sequence insertion

Donor

850 bp   741 bp   850 bp
Left homology arm   EGFP-Linker   Right homology arm
(397-1246)   (1249-2098)

397   1246 Met Val Ser Lys Gly Glu Glu Leu Phe Thr Gly

CGAGCTGCAGGCAACGCCCGAT   CTCATTGCTGATGGTGAGCAAGGGCGAGGAGCTGTTCACCGGG

Sequencing

left homology arm (397-1246)   partical EGFP coding sequence

b

| EGFP-MORC2FL | H2A-mCherry | Merge |

Prorphase

Telophase

10 μm

| EGFP-MORC2FL | H2A-mCherry | Merge |

Metarphase

a CTD

Mr (kDa)
72
55

b 150 mM NaCl

Cy3-CTD

1.25 μM 2.5 μM 5 μM 20 μM

10 μm

c Cy3-CTDΔCW

10 μM

DIC

Cy3

5 μm

d CTDΔCW Cy3-CTDΔCW

Mr (kDa)
72
55

e Static light-scattering (SLS) analysis

CTDΔCW
Cy3-CTDΔCW

118.2 ± 1.7 kDa

117.6 ± 0.7 kDa

dRI

Molecular Weight (kDa)

volume (mL)

f Cy3-CTDΔCW

0 s 9 s 63 s 108 s 180 s

2 μm

a

	EGFP	Bright-field	Merge
MORC2FL			
MORC2ΔIBD			

10 μm

b

HEK293T cells

EGFP-MORC2FL
EGFP-MORC2ΔIBD

Percent of Hela cells with condensates (%)

****

a

Top 20 GO Molecular Functions

ubiquitin−like protein ligase binding
cadherin binding
ubiquitin protein ligase binding
transcription coactivator activity
DNA−binding transcription factor binding
ribonucleoprotein complex binding
chromatin DNA binding
ATP hydrolysis activity
actin binding
double−stranded RNA binding
RNA polymerase II−specific DNA−binding transcription factor binding
ribosome binding
extracellular matrix structural constituent
ATP−dependent activity, acting on DNA
kinase regulator activity
promoter−specific chromatin binding
translation factor activity
integrin binding
DNA−binding transcription repressor activity
protein sequestering activity

$-\log10(p.adjust)$

Gene Count
- 20
- 40
- 60

Rich Factor
1.0
0.8
0.6
0.4
0.2
0.0

b

EGFP overexpression vs EGFP-MORC2FL overexpression

Genes previously characterized as being dysregulated by MORC2FL overexpression

1572 — 102 — 360

70 downregulated genes trends

Log2 Fold Change

FL   ΔCC3   ΔIDRa   +KS

32 upregulated genes trends

Log2 Fold Change

FL   ΔCC3   ΔIDRa   +KS

a Purified in HEK293F cell system for Fig. 1d and 1f

b Purified in HEK293F cell system for Fig. S1g

c Purified in *E. coli* cell system for Fig. 5a-f

d Purified in HEK293F cell system for Fig. 5j

e Purified in HEK293F cell system for Fig. 7c and 7d