A switch in kappa opioid receptor signaling from inhibitory to excitatory induced by stress in a subset of cortically-projecting dopamine neurons

Elyssa B. Margolis[1][*]

[1]UCSF Weill Institute for Neurosciences, Neuroscience Graduate Program, Department of Neurology, University of California, San Francisco, CA, United States

*Corresponding Author

Email: Elyssa.Margolis@ucsf.edu

1 Abstract

The kappa opioid receptor (KOR) has shown potential as a therapeutic target for several
neuropsychiatric disorders including major depressive disorder, pain, and substance use
disorder. *In vivo* signaling of G protein coupled receptors like the KOR is generally thought to
change in magnitude but not sign in such behavior states. Here we investigated KOR
modulation of ventral tegmental area (VTA) neurons following an acute, behaviorally aversive
manipulation. We found this switches KOR signaling from inhibitory to excitatory in a subset of
VTA dopamine neurons. Brief corticotrophin releasing factor (CRF) exposure *ex vivo* rapidly
induces a similar switch in KOR signaling, specifically in dorsal medial prefrontal cortex
(dmPFC) projecting neurons, but not nucleus accumbens or basolateral amygdala projecting
neurons. These KOR mediated excitations depend on G protein activation, but where
somatodendritic VTA KORs activate a K^+ conductance to hyperpolarize dopamine neurons in
control conditions, depolarizations require HCN channel function. One behavioral impact of this
change is a loss of the aversiveness of intra-VTA KOR activation, providing direct evidence that
rapid changes in GPCR signaling pathways can be triggered by activity at other GPCRs and
significantly alter behavioral responses driven by neuromodulators.

17 Introduction

Activation of the KOR system mediates immediate processing of and long-lasting behavioral
changes in response to aversive stimuli. Pharmacological blockade or genetic inactivation of
KORs can decrease the acute aversiveness of stressors, including pain, and promote resilience
to the development of depressive-like behaviors including decreased food reward seeking,
decreased social interactions, and increased immobility in forced swim and social defeat
paradigms[1–4]. Therefore, KOR antagonism has been proposed as a therapeutic approach to
neuropsychiatric disorders with symptom profiles that include aversive emotional and anhedonic
states, such as anxiety and depression, and to mitigate stress induced relapse to drug use[5].

The opioid receptor family of GPCRs commonly signals via inhibitory $G\alpha_{i/o}$ activation[6]. Agonists
at somatodendritic KORs hyperpolarize neurons in a variety of brain regions including the dorsal
raphe, nucleus raphe magnus, and ventral pallidum[7–9]. In the VTA, KOR activation inhibits a
subset of dopamine neurons through G protein-gated inwardly rectifying K^+ channel (GIRK)
activation while non-dopamine neurons show no response[10]. Behaviorally, KOR activation in the
VTA is aversive[11], and intra-VTA blockade of KORs prevents the aversiveness of systemic KOR
agonist administration[12]. Intra-VTA KOR blockade also prevents stress induced reinstatement of
cocaine seeking[13]. Paradoxically, aversive stress blunts some physiological KOR responses,
such as the KOR coupled GIRK conductance in serotonergic dorsal raphe neurons[8].
Understanding KOR function in relevant behavior states will improve our ability to treat the
target neuropsychiatric disorders.

Here we investigated postsynaptic KOR agonist responses in VTA neurons after an acute
aversive stressor in order to improve our understanding of what functionality KOR antagonism
might have in a therapeutically relevant state. Rather than a change in response magnitude,
acute aversive stress caused a change in the sign of the physiological response to KOR
activation in a subset of VTA neurons. Importantly, this change was specific to VTA dopamine
neurons projecting to the dmPFC, a site where dopamine function contributes to executive
function and reward processing[14–17].

44 Results

We performed *ex vivo* whole cell recordings in rat VTA neurons after a single session of
aversive foot shock stress, making slices 30-60 min after the stressor. All statistics are
described in Table 1. In addition to VTA neurons that responded to the KOR selective agonist
U69,593 (1 μM) with a hyperpolarization or no change in membrane potential, similar to prior
observations in untreated animals[10,18,19], we unexpectedly observed depolarizations in a distinct
subset of neurons from both male and female rats (Fig. 1a-d). This includes neurons confirmed
as dopaminergic with immunocytochemistry (Fig. 1b, Supplementary Fig. 1a; recording
locations in Supplementary Fig. 1b). Depolarizations were still observed 3 (4/18 neurons), but
not 5 (0/7 neurons), days after the foot shock stress session (Fig 1d). Action potential (AP)
threshold was more depolarized in neurons from foot shock stressed rats compared to shams,
but did not alter other basal properties of VTA neurons including baseline input resistance (Ri),
AP duration, AP peak, h current magnitude, or spontaneous firing rate (Supplementary Fig. 2).
In the recordings from stressed animals, initial Ri was greater in neurons that subsequently
showed a depolarization in response to U69,593 compared to neurons that subsequently
showed a hyperpolarization (Supplementary Fig. 3). No other differences were detected in
baseline physiological properties across VTA neurons grouped by their U69,593 responses
(Supplementary Fig. 3). One possibility is that the aversive stressor decreases GIRK function in
a subset of neurons, unmasking the observed depolarizing effect of U69,593. However, when
we blocked GIRKs in slices from control, untreated animals with $BaCl_2$ (100 μM), the distribution
of responses to U69,593 was clearly different from those from stressed animals with no
depolarization > 0.8 mV (Fig. 1d,e). Dopamine D2 receptor activation hyperpolarizes a subset of
VTA neurons through GIRK channel activation[20]. If acute, aversive stress decreases GIRK
function in VTA neurons rather than change KOR signaling, we would expect to observe a shift
in responses to the D2R agonist quinpirole (1 μM) as well. The means and distributions of
quinpirole responses were not different between VTA neurons from sham and foot shocked rats
(Fig. 1f).

The corticotrophin releasing factor (CRF) system is upstream of the KOR system in encoding
the aversiveness of stressful experiences[21], and blocking the KORs can prevent CRF-induced
anxiety-like behaviors[22,23]. As foot shock stress drives CRF release in the VTA[24], we tested the
hypothesis that incubating VTA slices from control animals with no behavioral manipulations in
CRF would produce similar changes in KOR signaling to the *in vivo* aversive stressor exposure.
In fact, in VTA slices incubated in 200 nM CRF for 5 min, a subset of neurons responded to
subsequent U69,593 application with a depolarization (Fig. 2a). Next we tested whether
neurons excited by KOR activation after CRF were the same neurons inhibited by KOR
activation under control conditions or a previously KOR-insensitive population. All neurons that
showed a depolarization in response to KOR activation after CRF showed a hyperpolarization to
KOR activation before CRF (Fig. 2b,c). In fact, for neurons that showed a depolarization after
CRF, there was a clear inverse correlation (n = 5 neurons; coeff. = -0.9) between the
magnitudes of KOR responses before and after CRF application (Fig. 2c).

We previously reported that VTA dopamine neurons that project to the mPFC and basolateral
amygdala (BLA), but not the nucleus accumbens (NAc), are inhibited by somatodendritic KOR
activation in rat[18,19]. To test whether the signaling switch happens in a projection selective
manner, we recorded from retrogradely labeled VTA neurons that project to dmPFC (including
prelimbic and anterior cingulate cortices), ventral mPFC (infralimbic cortex), medial NAc, and
BLA blind to injection site. Most distinctly, dmPFC-projecting VTA neurons showed the strongest
depolarizations after CRF exposure while none of the vmPFC-projecting neurons depolarized
(Fig. 3). NAc- and BLA-projecting showed no and minimal depolarizations, respectively (Fig. 3).
Thus, the switch in KOR signaling from inhibitory to excitatory in a specific subpopulation of
dmPFC-projecting VTA neurons.

Next we probed the signaling mechanism underlying the KOR mediated excitations triggered by
CRF exposure. To test whether the KOR excitations are due to indirect effects involving local
connectivity in the slice we measured KOR agonist responses while blocking AMPARs (DNQX,
10 μM), GABA$_A$Rs (gabazine 20 μM), and action potential activity (500 nM TTX). This did not
prevent U69,593 induced depolarizations (Fig. 4a). To block G protein signaling in the recorded
neuron we included GDP-β-s in the intracellular solution (500 μM), and broke in to whole cell
configuration only after CRF application was completed. This G protein block eliminated both
depolarizations and hyperpolarizations induced by U69,593 (Fig. 4a). Together these data show
that the KOR induced depolarizations are not circuit dependent, and because the G protein
blockade was limited to the recorded neuron the KORs in question appear to be expressed in
the recorded neuron. GPCR activation could cause depolarization by closing GIRKs[25], however
blocking GIRKs with BaCl$_2$ (100 μM) did not prevent U69,593 induced depolarizations (Fig. 4a).
We previously reported that mu and delta opioid receptors (MORs and DORs) on VTA neurons
can drive depolarizations that require Ca^{2+} channel conductances[26,27], however in the present
study, blocking Ca^{2+} channels with bath application of CdCl$_2$ (100 μM) did not prevent U69,593
induced depolarizations (Fig. 4a). A Na$^+$ leak current contributes to the maintenance of a
depolarized potential and spontaneous AP firing in VTA neurons[28,29] that can be inhibited by
G$\alpha_{i/o}$ protein coupled receptors[30]. Blocking this conductance did not prevent U69,593 induced
depolarizations (Fig. 4a).

I_h is a hyperpolarization activated, non-selective cation conductance that we found is prominent
in mPFC-projecting VTA neurons[19]. In nodose ganglion neurons MOR activation inhibits I_h[31],
and in the nucleus raphe magnus (NRM) KOR activation augments I_h[32]. The I_h blocker ZD7288
prevented U69,593 induced depolarizations (Fig. 4a). Given that the reversal potential for I_h is
depolarized compared to typical membrane potentials, the simplest possibility is that KOR
activation augments I_h conductance. Consistent with this, we found that U69,593 increased I_h
amplitude in a subset of neurons recorded in slices pretreated with CRF, and this response that
was not observed in control slices (Fig. 4b). CRF pretreatment did not impact other I_h properties
measured during U69,593 application (Supplementary Fig. 4).

Bruchas et al. reported that P38 MAPK contributes to some KOR mediated behavioral changes
following aversive stressors *in vivo*[33]. Here we tested for a contribution of p38 to this KOR
switch or KOR signaling. Incubating slices from control rats with the p38 blocker SB203580 prior
to CRF application prevented subsequent U69,593 induced depolarizations (Fig. 4c). However,
when slices were exposed to CRF prior to incubation in SB203580, U69,593 induced
depolarizations persisted (Fig. 4a). We conclude that CRF receptor activation alters the
signaling of a subset of KORs in the VTA through a p38 dependent signaling pathway, but
KORs are not signaling through p38 directly to depolarize neurons.

Finally, we tested for a behavioral consequence of this change in KOR signaling. Decreases in
dopamine neuron activity canonically encode punishment and aversion, and therefore intra-VTA
KOR agonist induced aversion is widely thought to be due to inhibition of dopamine neurons. In
the context of this model, a switch in KOR signaling in dopamine neurons from inhibitory to
excitatory would be predicted to diminish conditioned place aversion (CPA) to intra-VTA KOR
agonist microinjection. Since we found the switch in signaling sign to last between 3 and 5 days,
we designed a paradigm where the animals received foot shock on days 1 and 6 in order to
maintain stress-induced changes to KOR signaling. Pairings of either intra-VTA U69,593 (0.2
mM, 0.5 mL/side) or vehicle once per day were performed on non-stress days, for a total of four
pairings per side (Fig. 5a), in order to minimize the formation of a direct association between the
foot shock stress and the place preference manipulation. Indeed, foot shock stress pretreatment

eliminated the development of CPA to intra-VTA U69,593 (Fig. 5b, placements in
Supplementary Fig. 5).

Discussion

Together, these data show that a single stressful event can dramatically change the signaling
sign of a modulatory GPCR system. The change occurs in a specific neural circuit and can
persist for days. This stress induced switch at KORs can also be induced by the stress-
associated CRF neuropeptide-GPCR system in the same brain region, and while the signaling
is still G protein mediated, It appears that the key conductance involved is I_h, a switch from
GIRK coupling.

The contributions of VTA dopamine neurons to the processing of aversive experiences
encompass motivational and learning components and are circuit-specific[34]. *In vivo*
electrophysiology studies show that while many midbrain putative and opto-tagged dopamine
neurons decrease their firing when a predicted reward is omitted (i.e., an aversive outcome)[34,35],
some putative and directly identified dopamine neurons increase their firing in response to an
aversive sensory stimulus[36,37]. We previously demonstrated that in control rats, KOR activation
inhibits VTA dopamine neurons that project to the mPFC and BLA, but not the NAc[18,19]. With
microdialysis, we confirmed that *in vivo* VTA KOR activation decreases dopamine release in
mPFC but not in the NAc[19]. Here we show a switch from inhibitory to excitatory signaling in a
subpopulation of dmPFC-projecting VTA neurons after CRF exposure. This excitatory KOR
effect may contribute to the reported stress induced activation of dmPFC-projecting VTA
dopamine neurons[35,36]. The loss of intra-VTA KOR agonist induced CPA after the stress
induced switch in KOR signaling also implicates the dopaminergic projection to dmPFC in value
signaling. Consistent with this, MOR activation in the VTA causes an increase in dopamine in
the anterior cingulate cortex (ACC, included in the dmPFC in this study), and lesioning
dopamine terminals there decreases the magnitude of intra-VTA MOR agonist induced
conditioned place preference[15]. Also, genetic deletion of KOR from TH neurons prevents stress
induced augmentation of cocaine CPP[37]. The transfer function of dopamine activity in the mPFC
has been extensively studied, but a synthesized understanding of its contributions to mPFC
processing remains somewhat elusive[38–40].

There are at least a dozen possible sources of CRF input to the VTA including the bed nucleus
of the stria terminalis, dorsal raphe, parabrachial nucleus, lateral dorsal tegmentum, and local
VTA neurons[41,42] that might be activated by aversive stressors. A subset of VTA neurons
express CRF receptor 1 (CRF-R1) and the CRF binding protein, but not CRF receptor 2[43,44].
Blocking CRF-R1 in the VTA decreases foot shock stress induced reinstatement of cocaine
seeking[35]. Further, intra-VTA KOR blockade[13] and intra-dmPFC dopamine antagonists also
mitigate stress induced reinstatement of cocaine seeking[35,45], a provocative parallel to the
neuromodulatory circuit identified here.

We found that blocking I_h prevented KOR agonist induced depolarizations. Even though I_h is
generated by hyperpolarization activated, cyclic nucleotide-gated (HCN) channels[46,47] and thus
opioid receptors might interact with them though cAMP modulation, in the NRM KOR activation
in tissue from control rats augments I_h through a cAMP-independent signaling pathway that
involves intracellular release of Ca^{2+} [32]. In these neurons, cAMP activation causes a positive
shift in the voltage dependence of I_h, but does not change the maximum I_h amplitude. In the
VTA, the adenylyl cyclase activator forskolin causes a depolarizing shift in the I_h $V_{1/2}$[48]. The
isoform HCN3, which is expressed in the vast majority of VTA dopamine neurons and in many
more VTA neurons than isoforms 1, 2 or 4[48], is relatively insensitive to cAMP[46,49]. Here, KOR
activation after CRF did not alter I_h $V_{1/2}$, consistent with a cAMP-independent effect.

Consistent with our observations here, an increase in HCN channel function in the VTA has
been implicated in behavioral changes induced by aversive stressors. For instance, after social
defeat stress in mice, I_h magnitude is larger in VTA neurons of susceptible mice compared to
those in control mice[50]. Blocking I_h in the VTA[50,51] and systemically[52] reverse the social
avoidance induced by social defeat. Our data suggest that *in vivo* activity at KOR in the VTA
after stress increases I_h function in dmPFC-projecting dopamine neurons. Thus decreasing I_h
function in the VTA may be one common therapeutic target, be it through direct HCN blockade
or by KOR antagonism.

What are the *in vivo* consequences of this switch in KOR signaling? At the circuit level, this
change in KOR signaling means that the change in VTA dmPFC-projecting dopamine neuron
firing will have an opposite sign when an endogenous KOR ligand is released into the VTA. That
is, the peptidergic inputs to the VTA may have the same behaviorally related firing patterns as
before an aversive stressor, but the impact on dmPFC dopamine release will be the inverse. In
addition to contributing to value encoding, dopamine function in the dmPFC also contributes to
executive function in several ways[16,17]. For instance, dopamine lesions in in the ACC impairs
attentional set shifting[41] and dopamine receptor antagonists delivered to the prelimbic cortex
degrade decision making[42]. These and other studies therefore suggest that ongoing or
increases in dopamine release in the dmPFC contribute to saliency processing, and support
working memory[43], which are particularly advantageous following an aversive stressor.
Activating mPFC-projecting VTA dopamine neurons reverses chronic stress induced immobility
and loss of sucrose preference[44], and an increase in dopamine signaling in dmPFC also
relieves ongoing pain[45]. Provocatively, stress induced analgesia is also blocked by systemic
administration of the KOR selective antagonist norBNI[46], which in the context of our results
would prevent an increase in dopamine release in the dmPFC, although it is not yet known if
dmPFC-projecting VTA dopamine neurons contribute to this phenomenon. Together, these
studies raise the possibility that this switch in VTA KOR function is adaptive. An optimal
concentration of dopamine in the mPFC facilitates learning, and learning about acutely aversive
experiences, and how to avoid them, is critical to survival. Here we discovered a convergence of
two neuropeptide systems, one changing the sign of the physiological response of the other in a
specific neural circuit.

Methods

Animals.
All experiments were performed in accordance with the guidelines of the National Institutes of
Health Guide for the Care and Use of Laboratory Animals and the Institutional Animal Care and
Use Committees (IACUC) at the University of California San Francisco. Male and female
Sprague Dawley rats were obtained from Inotiv or Charles River Laboratories. Rats were
allowed access to food and water ad libitum and maintained on a reverse 12 h:12 h light/dark
cycle. Rats were group housed except those with cannula implants.
226
Foot shock stress.
Operant chambers equipped with a foot shock grid were used for administering aversive
stressors (physical boxes manufactured by Med. Associates with custom updated electronics
controlled by microcontroller (Arduino)). Foot shock or sham sessions were 10 min, delivering
10, 0.8 mA intensity, 0.5 s duration shocks. Intervals between shocks were randomized (10 –
110 sec) in order to avoid predictability. Foot shock stress moderately elevated blood CORT,
samples taken directly after foot shock session (Supplementary Fig. 6).
234
235 **Electrophysiology.**

Rats were deeply anesthetized with isoflurane, decapitated, and brains were quickly removed and cooled with ice-cold artificial cerebrospinal fluid (aCSF) consisting of (in mM): 119 NaCl, 2.5 KCl, 1.0 NaH_2PO_4, 26.2 $NaHCO_3$, 11 glucose, 1.3 $MgSO_4$, 2.5 $CaCl_2$, saturated with 95% O_2-5% CO_2, with a measured osmolarity 310–320 mOsm/L. Horizontal sections (150 μm) containing the VTA were cut with a Leica VT 1000 S or a Campden 7000smz-2 vibratome. Tissue containing the DiI injection sites was drop fixed in 4% formaldehyde to confirm targeting, with recordings in DiI labeled neurons made blind to injection target. Slices were incubated in oxygenated aCSF at 33 °C and allowed to recover for at least one hour. A single slice was placed in the recording chamber and continuously superfused with oxygenated aCSF (2 mL/min). Neurons were visualized with an upright microscope (Zeiss AxioExaminer.D1) equipped with infrared-differential interference contrast, Dodt optics, and fluorescent illumination. Whole-cell recordings were made at 34 °C using borosilicate glass microelectrodes (3–5 MΩ) filled with K-gluconate internal solution containing (in mM): 123 K-gluconate, 10 HEPES, 8 NaCl, 0.2 EGTA, 2 MgATP, 0.3 Na_3GTP, and 0.1% biocytin (pH 7.2 adjusted with KOH; 275 mOsm/L). Liquid junction potentials were not corrected. Input resistance was monitored throughout voltage clamp experiments with a −4 mV step every 30 s.

Signals were recorded using an Axopatch 1D (Molecular Devices, San Jose, CA) or IPA (Sutter Instruments, Novato, CA). Signals were filtered at 5 kHz and collected at 20 kHz using IGOR Pro (Wavemetrics) and custom data acquisition routines or collected at 10 kHz using SutterPatch software (Sutter Instruments).

Baseline hyperpolarization-activated cation currents (I_h) were recorded in each neuron by voltage clamping cells at -60 and stepping to -40, -50, -70, -80, -90, -100, -110, and -120 mV. Recordings were then made in current clamp mode (I = 0 pA). Changes in I_h properties were measured in voltage clamp, alternating between two step series (Supplementary Fig. 4) with one series commencing each 2.17 min.

After recordings, slices were drop fixed in 4% PFA for at least 2 h at 4 °C and processed for tyrosine hydroxylase (TH) immunocytochemistry and biocytin labeling. Recording were made from throughout the VTA.

Immunocytochemistry.
Slices were preblocked for 2 h at room temperature in PBS with 0.3% Tween20 and 5% normal goat serum, then incubated at 4°C with a rabbit anti-TH polyclonal antibody (1:400; EMD Millipore, RRID: AB_390204). Slices were then washed thoroughly in PBS with 0.3% Tween20 before being agitated overnight at 4°C with Cy5 anti-rabbit secondary antibody (1:1000; Jackson ImmunoResearch, RRID: AB_2534032) and FITC streptavidin (1:200; Jackson ImmunoResearch, AB_2337236). Tween20 was excluded in slices with DiI. Sections were rinsed and mounted on slides using VECTASHIELD® Antifade Mounting Media (Vector Laboratories) and visualized with an Axioskop FS2 Plus (Zeiss) with an Axiocam MRm (Zeiss) running Neurolucida (MBF Biosciences).

CORT ELISA.
Trunk blood was collected at time of slice preparation for electrophysiology animals. Samples were centrifuged, then serum was frozen and stored at -80°C. ELISA kits were used as directed (DetectX Corticosterone Enzyme Immunoassay Kit (product # K014-H1); Arbor Assays).

Stereotaxic surgeries.
Tracer injections. Rats weighing 275–300 g were anesthetized with 3–5% isoflurane (Henry Schein) via inhalation and secured in a stereotaxic frame. Bilateral craniotomies were created with a dental drill above the injection site.

Tracer injections. Bilateral injections of DiI (7% in ethanol; Biotium, Hayward, CA) were made in
the dmPFC (2.0 mm AP, ± 0.6 mm ML, -2.8 mm V), vmPFC (2.0 mm AP, ± 0.6 mm ML, -4.8
mm V), NAc (1.2 mm AP, ± 0.8 mm ML, -7.4 mm V), or amygdala (-3.3 mm AP, ± 5.0 mm ML, -
8.4 mm V) using a Nanoject II (Drummond Scientific, Broomall, PA). 50 – 300 nL was injected in
each hemisphere over ~5 min. The injector tip was left in place for at least 2 minutes after
injection to decrease backflow and spread to tissue dorsal to the injection site. Recordings were
made 7-8 days later.
294
295 *Cannulation.* For microinjections into the VTA, bilateral guide cannulae were implanted 1 mm
296 above the VTA (−5.8 mm AP, ±0.5 mm ML,−7.2 mm DV). A dummy stylet was inserted to
297 maintain patency of the cannulae. Cannulae were anchored with flat point screws and dental
298 cement.
299
300 *Analgesia and recovery*. Rats were treated with subcutaneous Carprofen (5 mg/kg, Zoetis) and
301 topical 2% lidocaine (Phoenix Pharmaceutical, Inc.) during the surgery for pain control. After
302 surgery, animals had access to liquid Tylenol (~1:40) in their drinking water for 3–5 days or
303 were administered Meloxicam (s.c. 2 mg/kg, Pivetal) once per day for two days. All DiI and
304 cannulae placements were histologically verified postmortem.
305
Conditioned place preference.
Rats were allowed to recover for 1–2 weeks prior to behavioral testing. First, all rats were tested
for bias in the conditioning boxes. The conditioning boxes (physical boxes purchased from Med.
Associates, Georgia, VT, USA with custom updated electronics (Arduino; IR beam breaks
(Adafruit)) and software have three divisions: two conditioning chambers (25 cm × 21 cm × 21
cm) with distinct visual (horizontal vs. vertical stripes) and textural (thick vs. thin mesh flooring)
cues, separated by a smaller gray neutral chamber (12 cm × 21 cm × 21 cm). Animals were
allowed up to three opportunities to show neutrality across the chambers during 20-min baseline
sessions. Rats that displayed a consistent baseline preference (>65% of time spent in one
chamber) were excluded from the study. Rats were pseudorandomly assigned to receive
U69,593 in one of the conditioning chambers, and assignments were counter-balanced for each
cohort.
The paradigm commenced with a foot shock or sham treatment as described above, completed
in a different room from the conditioning chambers. Starting the next day, conditioning pairings
(20 min) occurred once daily for 4 consecutive days alternating U69,593 (0.2 mM) and vehicle
(0.5 mL/side). This 5 day series (including foot shock/sham) was repeated a second time for a
total of 4 pairings for U69,593 and vehicle, each. Finally, on test day, rats were drug free and
had access to all three chambers for 20 min. Difference score was defined as (Time
spent in U69,593 paired chamber)−(Time spent in vehicle paired chamber).
325
326 *Microinjections*. Rats were lightly restrained in a cloth wrap for intracranial microinjections. A
327 bilateral 33 G microinjector (PlasticsOne) that extended 1 mm ventrally beyond the guide
328 cannula was inserted to deliver drug. Hamilton syringes were driven by a dual syringe pump to
329 deliver the injection volume over 2 min. Injectors were left in place for 1 min after injection.
330
Brain removal and histochemistry. To check cannulae placements, rats were deeply
anesthetized with an intraperitoneal injection of Euthasol (0.1 mg/kg, Virbac Animal Health) after
0.3 µL of Chicago Sky Blue (in 2% PBS) was injected through the cannulae to mark injection
locations. After becoming unresponsive to noxious stimuli, the rats were transcardially perfused
with 400 mL of saline, followed by 400 mL of 4% paraformaldehyde (PFA) in 0.1 M phosphate
buffer. The brains were extracted and immersion-fixed in PFA for 2 h at room temperature (RT),
washed two times with PBS to remove excess PFA, and stored in 1X PBS at 4°C.

Perfused brains and drop fixed DiI containing tissue were sectioned (50 µm) using a vibratome
(Leica VT 1000S). Alternating slices were Nissl stained. Sections were mounted on slides using
VECTASHIELD® mounting medium (Vector Laboratories). Microinjection locations for rats
included in the behavior study are summarized in Supplementary Fig. 5.
342
Data analysis.
The magnitude of U69,593 induced changes in V_m were quantified by comparing the mean
values over the last 4 min of U69,593 application to the 4 minutes of baseline just prior to
starting the drug application. Responses were tested for significance by statistically comparing
30 sec bins during these two time periods (unpaired Student's t-test). I_h magnitude was
measured as the difference between the initial capacitive response to a voltage step from −60 to
−120 mV and the final current during the same step. Reported baseline Ri and firing rate are the
average measurements over the first 2 min of I_{clamp} recording. For AP duration measurements,
at least 8 waveforms were averaged together and duration was calculated from threshold on the
ascending phase to recrossing threshold on the falling phase. Threshold was identified as the
first datapoint where the derivative of the waveform exceeds 10 V/s. For evaluation of KOR
agonist induced changes in I_h properties, see measurements described in Supplementary Fig. 4.
355
356 Population statistics were performed in R. Where datasets failed tests of normality or
357 homogeneity of variance, non-parametric permutation analyses (https://github.com/eb-margolis-
358 neuroscience-lab/R-general-Margolislab/tree/main) were used to test for statistical significance.
359 For behavior data, mixed measures two way ANOVA was used with Bonferroni correction for
360 multiple comparisons.
361

Acknowledgements
I thank Joseph R. Driscoll, Thomas J. Cirino, Peter Fong, Aashka Popat, Ben J. Ngu, Benjamin
Snyder, Kasra Mansorian, and Madelyn Moulton for technical assistance. This work was
supported by National Institutes of Health grants R01DA030529, R01DA042025 and
R01DA008863, and funds from the State of California for medical research on alcohol and
substance use.

368
Figure Captions
Figure 1. Following a single foot shock session, a subset of VTA neurons depolarize in
response to KOR activation. (a) Rats experienced a single intermittent foot shock session (10
min) prior to *ex vivo* whole cell recordings. (b) Example recording from a VTA neuron (biocytin
fill: red) that was confirmed as dopaminergic with TH immunocytochemistry (green), scale bar
10 µm. In whole cell current clamp configuration (I = 0 pA) bath application of the KOR agonist
U69,592 (1 µM) caused a depolarization. Inset: I_h measured in voltage clamp, scale bars 200 pA
and 200 ms. (c) Mean time course of responses in neurons that were significantly depolarized
by U69,593. (d) Summarized data, each circle is a neuron, from same day sham shock, same
day shock, 3 days after shock and 5 days after shock treated rats. Circles and triangles indicate
VTA neurons from male and female rats, respectively. Permutation analysis comparing means
of U69,593 responses in sham vs shock, p = 0.02 (Table 1) Shading indicates within cell
statistics (inset) (e) Summarized U69,593 responses in VTA neurons from naïve rats, and from
naïve rats where U69,593 responses were measured in $BaCl_2$ (100 µM). Statistical comparison
performed was permutation analysis comparing variances (SDs; $ indicates p<0.05), p = 0.04.
(f) Summarized responses to the dopamine D2 receptor agonist quinpirole (1 µM) in VTA
neurons from naïve, sham, and foot shocked rats. One way ANOVA sham vs shock, p = 0.9.
386

Figure 2. *ex vivo* exposure to CRF can switch KOR signaling from inhibitory to excitatory.
(a) Summarized data, each circle is a neuron. Slices were incubated for 5 min with CRF (200
nM), CRF was washed out for at least 5 min, then U69,593 (1 μM) was bath applied for 5 min.
Statistical comparison to U69,593 responses in VTA neurons from naïve rats (Fig. 1e left),
Permutation analysis of means, p = 0.00004. Shading indicates within cell statistics. (b) Left,
U69,593 (1 μM) was applied once, washed out, then reapplied (paired t-test p = 1.0). Right,
U69,593 (1 μM) was applied once, washed out, CRF was washed on (5 min, 200 nM), washed
out, then U69,593 was reapplied (paired t-test p = 0.08). (c) Data from (b) graphed as
magnitude of response to first U69,593 application to second application.
396
397 **Figure 3. CRF switches KOR signaling in a subpopulation of dmPFC-projecting VTA**
398 **neurons.** U69,593 responses were recorded in retrogradely labeled neurons after slices were
399 incubated in CRF (5 min, 200 nM). Kruskal-Wallis rank sum test p = 0.006, pairwise
400 comparisons listed in Table 1. Shading indicates within cell statistics.
401
Figure 4. KOR mediated depolarizations require I_h. (a) Signaling mechanism(s) that
contribute to U69,593 (1 μM) induced depolarizations were investigated after exposing slices *ex*
vivo to CRF (5 min, 200 nM). TTX (500 nM), DNQX (10 μM), and gabazine (20 μM) blocked
synaptic transmission; intracellular GDP-β-s (500 μM) blocked intracellular G protein signaling;
$BaCl_2$ (100 μM) blocked K^+ channels; $CdCl_2$ (100 μM) blocked Ca^{2+} channels; replacement of
Na^+ with NMDG (151 mM, replacing NaCl) in aCSF blocked Na^+ leak current; ZD7288 (10 μM)
blocked HCN channels (I_h); SB203580 (10 μM) blocked p38/MAPK signaling. (b) I_h magnitude
was measured during U69,593 (1 μM) application in voltage clamp, stepping from -60 mV to -
120 mV, in control slices or slices pretreated with CRF (5 min, 200 nM). (c) p38/MAPK was
blocked with SB203580 application during both CRF and U69,593 application. Shading
indicates within cell statistics.
413
Figure 5. Foot shock stress eliminates intra-VTA KOR mediated aversion. (a) Experiment
timeline where rats were trained in a conditioned place preference paradigm, to intra-VTA
U69,593 (0.2 mM, 0.5 mL/side) or vehicle. Rats were pretreated with sham handling or foot
shock stress on non-pairing days. (b) Baseline vs post test for place conditioning. Two way
mixed measures ANOVA) Pairwise comparisons: Sham, baseline vs post test: 0.0004 Shock,
baseline vs post test: 0.6 (Bonferroni corrected).

Supplementary Figure 1. (a) Summary of responses to U69,593 in neurons identified at TH(+)
with immunocytochemistry. These are a subset of the data in Fig. 1d. (b) Locations of VTA
neurons recorded from same day foot shock stress rats. Color indicates change in Vm in
response to U69,593.

Supplementary Figure 2. Physiological properties recorded in VTA neurons from foot shock
(same day) or sham rats. Quantifications of data collected during the first 150 s immediately
after whole cell access is achieved. (a) Ri, (b) AP duration, (c) AP threshold, (d) AP peak, (e) I_h
magnitude (f) spontaneous firing rate. **p < 0.01

Supplementary Figure 3. Foot shock rat data from Supplementary Figure 2, sorted by the type
of response to U69,593. (a) Ri, (b) AP duration, (c) AP threshold, (d) AP peak, (e) I_h magnitude
*p < 0.05

Supplementary Figure 4. I_h measurements. (a) top, schematic of V step commands from V=-
40 mV to quantify $V_{1/2}$ and activation τ. Bottom, example response. (b) top, schematic of V step
commands from V = -40 mV to quantify reversal V. Bottom, example response from the same

neuron as in (a). (c) Example single exponential fit to data in (a). (d) Schematic of experiment
design. Step series shown in (a) and (b) were applied to each neuron for a "baseline"
measurement and again during U69,593 application. (e) Difference between $V_{1/2}$ measured at
baseline and in U69,593. Grey is no pretreatment, Orange is CRF pretreatment. (f) Difference
between $V_{reverse}$ measured at baseline and in U69,593. Grey is no pretreatment, Orange is CRF
pretreatment. (g) I_h activation τ at baseline vs in U69,593. Grey is no pretreatment, Orange is
CRF pretreatment.

Supplementary Figure 5. Microinjection sites for intra-VTA place conditioning experiments in
Figure 5.

Supplementary Figure 6. CORT measured (ELISA) from trunk blood samples collected from
same day foot shock and sham animals during brain slice preparation. * p = 0.05

References

1. Varlinskaya, E. I., Spear, L. P. & Diaz, M. R. Stress alters social behavior and sensitivity to

pharmacological activation of kappa opioid receptors in an age-specific manner in

Sprague Dawley rats. *Neurobiol. Stress* **9**, 124–132 (2018).

2. McLaughlin, J. P., Li, S., Valdez, J., Chavkin, T. A. & Chavkin, C. Social defeat stress-

induced behavioral responses are mediated by the endogenous kappa opioid system.

Neuropsychopharmacology **31**, 1241–8 (2006).

3. Massaly, N. *et al.* Pain-Induced Negative Affect Is Mediated via Recruitment of The

Nucleus Accumbens Kappa Opioid System. *Neuron* (2019)

doi:10.1016/j.neuron.2019.02.029.

4. Groblewski, P. A., Zietz, C., Willuhn, I., Phillips, P. E. M. & Chavkin, C. Repeated stress

exposure causes strain-dependent shifts in the behavioral economics of cocaine in rats.

Addict. Biol. **20**, 297–301 (2015).

5. Carlezon Jr., W. A. & Krystal, A. D. Kappa-Opioid Antagonists for Psychiatric Disorders:

From Bench to Clinical Trials. *Depress. Anxiety* **33**, 895–906 (2016).

6. Reeves, K. C., Shah, N., Muñoz, B. & Atwood, B. K. Opioid Receptor-Mediated Regulation

of Neurotransmission in the Brain. *Front. Mol. Neurosci.* **15**, 919773 (2022).

7.   Ji, M. *et al.* Dynorphin promotes stress-induced depressive behaviors by inhibiting

ventral pallidal neurons in rats. *Acta Physiol.* **236**, (2022).

8.   Lemos, J. C. *et al.* Repeated stress dysregulates κ-opioid receptor signaling in the dorsal

raphe through a p38α MAPK-dependent mechanism. *J. Neurosci. Off. J. Soc. Neurosci.* **32**,

12325–12336 (2012).

9.   Pan, Z. Z., Tershner, S. A. & Fields, H. L. Cellular mechanism for anti-analgesic action of

agonists of the kappa-opioid receptor. *Nature* **389**, 382–385 (1997).

10. Margolis, E. B., Hjelmstad, G. O., Bonci, A. & Fields, H. L. Kappa-opioid agonists directly

inhibit midbrain dopaminergic neurons. *J Neurosci* **23**, 9981–6 (2003).

11. Bals-Kubik, R., Ableitner, A., Herz, A. & Shippenberg, T. S. Neuroanatomical sites

mediating the motivational effects of opioids as mapped by the conditioned place

preference paradigm in rats. *J Pharmacol Exp Ther* **264**, 489–95 (1993).

12. Ehrich, J. M. *et al.* Kappa Opioid Receptor-Induced Aversion Requires p38 MAPK

Activation in VTA Dopamine Neurons. *J Neurosci* **35**, 12917–31 (2015).

13. Graziane, N. M., Polter, A. M., Briand, L. A., Pierce, R. C. & Kauer, J. A. Kappa opioid

receptors regulate stress-induced cocaine seeking and synaptic plasticity. *Neuron* **77**,

942–954 (2013).

14. Vijayraghavan, S., Major, A. J. & Everling, S. Neuromodulation of Prefrontal Cortex in

Non-Human Primates by Dopaminergic Receptors during Rule-Guided Flexible

Behavior and Cognitive Control. *Front. Neural Circuits* **11**, 91 (2017).

15. Narita, M. *et al.* Implication of dopaminergic projection from the ventral tegmental area

to the anterior cingulate cortex in μ-opioid-induced place preference. *Addict. Biol.* **15**,

434–447 (2010).

16. Tichelaar, J. G., Sayalı, C., Helmich, R. C. & Cools, R. Impulse control disorder in

Parkinson's disease is associated with abnormal frontal value signalling. *Brain J. Neurol.*

146, 3676–3689 (2023).

17. Cools, R., Froböse, M., Aarts, E. & Hofmans, L. Dopamine and the motivation of cognitive

control. *Handb. Clin. Neurol.* **163**, 123–143 (2019).

18. Margolis, E. B., Mitchell, J. M., Ishikawa, J., Hjelmstad, G. O. & Fields, H. L. Midbrain

dopamine neurons: projection target determines action potential duration and

dopamine D(2) receptor inhibition. *J Neurosci* **28**, 8908–13 (2008).

19. Margolis, E. B. *et al.* Kappa opioids selectively control dopaminergic neurons projecting

to the prefrontal cortex. *Proc Natl Acad Sci U A* **103**, 2938–42 (2006).

20. Seutin, V., Johnson, S. W. & North, R. A. Effect of dopamine and baclofen on N-methyl-D-

aspartate-induced burst firing in rat ventral tegmental neurons. *Neuroscience* **58**, 201–

206 (1994).

21. Land, B. B. *et al.* The dysphoric component of stress is encoded by activation of the

dynorphin kappa-opioid system. *J Neurosci* **28**, 407–14 (2008).

22. Bruchas, M. R., Land, B. B., Lemos, J. C. & Chavkin, C. CRF1-R activation of the

dynorphin/kappa opioid system in the mouse basolateral amygdala mediates anxiety-

like behavior. *PloS One* **4**, e8528 (2009).

23. Van't Veer, A., Yano, J. M., Carroll, F. I., Cohen, B. M. & Carlezon, W. A. Corticotropin-

releasing factor (CRF)-induced disruption of attention in rats is blocked by the κ-opioid

receptor antagonist JDTic. *Neuropsychopharmacol. Off. Publ. Am. Coll.*

Neuropsychopharmacol. **37**, 2809–2816 (2012).

24. Wang, B. *et al.* Cocaine experience establishes control of midbrain glutamate and

dopamine by corticotropin-releasing factor: a role in stress-induced relapse to drug

seeking. *J Neurosci* **25**, 5389–96 (2005).

25. Xia, Y.-F., Margolis, E. B. & Hjelmstad, G. O. Substance P inhibits GABAB receptor

signalling in the ventral tegmental area. *J. Physiol.* **588**, 1541–1549 (2010).

26. Margolis, E. B., Hjelmstad, G. O., Fujita, W. & Fields, H. L. Direct bidirectional µ-opioid

control of midbrain dopamine neurons. *J Neurosci* **34**, 14707–16 (2014).

27. Margolis, E. B., Fujita, W., Devi, L. A. & Fields, H. L. Two delta opioid receptor subtypes

are functional in single ventral tegmental area neurons, and can interact with the mu

opioid receptor. *Neuropharmacology* **123**, 420–432 (2017).

28. Cobb-Lewis, D. E., Sansalone, L. & Khaliq, Z. M. Contributions of the Sodium Leak

Channel NALCN to Pacemaking of Medial Ventral Tegmental Area and Substantia Nigra

Dopaminergic Neurons. *J. Neurosci. Off. J. Soc. Neurosci.* **43**, 6841–6853 (2023).

29. Khaliq, Z. M. & Bean, B. P. Pacemaking in dopaminergic ventral tegmental area neurons:

depolarizing drive from background and voltage-dependent sodium conductances. *J.*

Neurosci. Off. J. Soc. Neurosci. **30**, 7401–7413 (2010).

30. Philippart, F. & Khaliq, Z. M. Gi/o protein-coupled receptors in dopamine neurons

inhibit the sodium leak channel NALCN. *eLife* **7**, e40984 (2018).

31. Ingram, S. L. & Williams, J. T. Opioid inhibition of Ih via adenylyl cyclase. *Neuron* **13**,

179–186 (1994).

32. Pan, Z. Z. κ-Opioid receptor-mediated enhancement of the hyperpolarization-activated

current (Ih) through mobilization of intracellular calcium in rat nucleus raphe magnus.

J. Physiol. **548**, 765–775 (2003).

33. Bruchas, M. R. *et al.* Stress-induced p38 mitogen-activated protein kinase activation

mediates kappa-opioid-dependent dysphoria. *J Neurosci* **27**, 11614–23 (2007).

34. Morales, M. & Margolis, E. B. Ventral tegmental area: cellular heterogeneity,

connectivity and behaviour. *Nat Rev Neurosci* (2017) doi:10.1038/nrn.2016.165.

35. Vranjkovic, O. *et al.* Enhanced CRFR1-Dependent Regulation of a Ventral Tegmental

Area to Prelimbic Cortex Projection Establishes Susceptibility to Stress-Induced

Cocaine Seeking. *J. Neurosci. Off. J. Soc. Neurosci.* **38**, 10657–10671 (2018).

36. Deutch, A. Y. *et al.* Stress selectively increases fos protein in dopamine neurons

innervating the prefrontal cortex. *Cereb. Cortex N. Y. N 1991* **1**, 273–292 (1991).

37. Abraham, A. D., Casello, S. M., Land, B. B. & Chavkin, C. Optogenetic stimulation of

dynorphinergic neurons within the dorsal raphe activate kappa opioid receptors in the

ventral tegmental area and ablation of dorsal raphe prodynorphin or kappa receptors

in dopamine neurons blocks stress potentiation of cocaine reward. *Addict. Neurosci.* **1**,

100005 (2022).

38. Sheynikhovich, D., Otani, S., Bai, J. & Arleo, A. Long-term memory, synaptic plasticity and

dopamine in rodent medial prefrontal cortex: Role in executive functions. *Front. Behav.*

Neurosci. **16**, 1068271 (2023).

39. Durstewitz, D. & Seamans, J. K. The dual-state theory of prefrontal cortex dopamine

function with relevance to catechol-o-methyltransferase genotypes and schizophrenia.

Biol. Psychiatry **64**, 739–749 (2008).

40. Robinson, S. E. & Sohal, V. S. Dopamine D2 Receptors Modulate Pyramidal Neurons in

Mouse Medial Prefrontal Cortex through a Stimulatory G-Protein Pathway. *J. Neurosci.*

Off. J. Soc. Neurosci. **37**, 10063–10073 (2017).

41. Soden, M. E. *et al.* Distinct Encoding of Reward and Aversion by Peptidergic BNST

Inputs to the VTA. *Front. Neural Circuits* **16**, 918839 (2022).

42. Grieder, T. E. *et al.* VTA CRF neurons mediate the aversive effects of nicotine

withdrawal and promote intake escalation. *Nat. Neurosci.* **17**, 1751–1758 (2014).

43. Van Pett, K. *et al.* Distribution of mRNAs encoding CRF receptors in brain and pituitary

of rat and mouse. *J. Comp. Neurol.* **428**, 191–212 (2000).

44. Wang, H.-L. & Morales, M. Corticotropin-releasing factor binding protein within the

ventral tegmental area is expressed in a subset of dopaminergic neurons. *J. Comp.*

Neurol. **509**, 302–318 (2008).

45. McFarland, K. & Kalivas, P. W. The circuitry mediating cocaine-induced reinstatement of

drug-seeking behavior. *J. Neurosci. Off. J. Soc. Neurosci.* **21**, 8655–8663 (2001).

46. Combe, C. L. & Gasparini, S. Ih from synapses to networks: HCN channel functions and

modulation in neurons. *Prog. Biophys. Mol. Biol.* **166**, 119–132 (2021).

47. Monteggia, L. M., Eisch, A. J., Tang, M. D., Kaczmarek, L. K. & Nestler, E. J. Cloning and

localization of the hyperpolarization-activated cyclic nucleotide-gated channel family in

rat brain. *Mol. Brain Res.* **81**, 129–139 (2000).

48. Mu, L. *et al.* cAMP-mediated upregulation of HCN channels in VTA dopamine neurons

promotes cocaine reinforcement. *Mol. Psychiatry* **28**, 3930–3942 (2023).

49. Mistrík, P. *et al.* The Murine HCN3 Gene Encodes a Hyperpolarization-activated Cation

Channel with Slow Kinetics and Unique Response to Cyclic Nucleotides. *J. Biol. Chem.*

280, 27056–27061 (2005).

50. Cao, J.-L. *et al.* Mesolimbic dopamine neurons in the brain reward circuit mediate

susceptibility to social defeat and antidepressant action. *J. Neurosci. Off. J. Soc. Neurosci.*

30, 16453–16458 (2010).

51. Cai, M. *et al.* HCN channel inhibitor induces ketamine-like rapid and sustained

antidepressant effects in chronic social defeat stress model. *Neurobiol. Stress* **26**,

100565 (2023).

52. Teichman, E. M. *et al.* Design and validation of novel brain-penetrant HCN channel

inhibitors to ameliorate social stress-induced susceptible phenotype. *Mol. Psychiatry*

(2025) doi:10.1038/s41380-025-02972-8.

Table 1 – Assumption testing and statistical comparisons for summary data

Experiment	Figure	# Neurons; # Outliers; # Extreme outliers	Shapiro-Wilk test of Normality Statistic; p val	Test for Homogeneity of Variances Statistic; p val (Test used)	Parametric test Statistic; p val (Test used)	Non-parametric test Statistic and/or p val (Test used)
U69,593 response (mV), acute stress vs sham	1d	67; 3; 0	0.9; 0.00008	7.5; 0.008 (Levene's)	n/a	0.02 (Permutation, mean)
U69,593 response in BaCl$_2$ vs acute stress	1d,e	49; 4; 0	0.9; 0.0007	3.0; 0.09 (Levene's)	n/a	0.04 (Permutation, SD)
Quinpirole response (mV), acute stress vs sham	1f	27; 2; 1	1.0; 0.6	1.4; 0.2 (Levene's)	0.02; 0.9 (one way ANOVA)	n/a
U69.593 response naïve vs after CRF	1e, 2a	78; 2; 0	1.0; 0.02	0.9; 0.4 (Levene's)	n/a	0.00004
U69,593, first application vs second application	2b, left	6; 0; 0	0.9; 0.6	0.02; 0.9	0.0003; 1.0 (paired t-test)	
U69,593 responses pre- vs post- CRF	2b, right	10; 0; 0	1.0; 0.7	2.0; 0.2	-2.0; 0.08 (paired t-test)	
U69,593, first application vs second application	2c	6; 0; 0			5.1; 0.007 (Pearson's product moment correlation, coeff = 0.9)	
U69,593 responses pre- vs post- CRF, in neurons with a post-CRF U69,593 induced depolarization	Subset in 2c	5			-3.7; 0.03 (Pearson's product moment correlation, coeff = -0.9)	
U69,593 responses differ by projection target	3	48; 6; 3	0.8; 0.0000001	1.4; 0.3	n/a	12.4; 0.006 (Kruskal-Wallis rank sum test) Pairwise comparisons (p adjustment method BH): dmPFC-BLA: 0.05 dmPFC-NAc: 0.05 dmPFC-vmPFC: 0.03 BLA-NAc: 0.4 BLA-vmPFC: 0.4 NAc-vmPFC: 0.2
U69,593 after CRF vs after CRF with GDP-β-s	4a	34; 1; 0	0.9; 0.007	4.4; 0.05	n/a	0.01 (Permutation, SD)
U69,593 after CRF vs after CRF with ZD7288	4a	39; 2; 1	0.9; 0.06	2.5; 0.1	0.7; 0.4 (one way ANOVA)	
U69,593 after CRF vs after CRF with TTX, DNQX, gabazine	4a	47; 4; 0	0.9; 0.005	1.2; 0.3	n/a	0.3 (Permutation, mean)
U69,593 after CRF vs after CRF with BaCl$_2$	4a	38; 1; 0	0.9; 0.01	2.5; 0.1	n/a	0.5 (Permutation, mean)
U69,593 after CRF vs after CRF with CdCl$_2$	4a	33; 1; 0	0.9; 0.007	0.01; 0.9	n/a	0.6 (Permutation, mean)
U69,593 after CRF vs after CRF with NMDG	4a	31; 1; 0	0.9; 0.01	1.1; 0.3	n/a	0.2 (Permutation, mean)
U69,593 after CRF vs after CRF with SB203580 (only during U69,593)	4a	36; 1; 0	0.9; 0.00008	7.5; 0.008	n/a	0.4 (Permutation, mean)
I_h magnitude after U69,593, with vs without CRF pretreatment	4b	18; 2; 0	0.9; 0.02	1.7; 0.2	n/a	0.3 (Permutation, mean)
U69,593 after CRF vs after CRF with SB203580 (applied	4a,c	34; 2; 0	1.0; 0.2	0.0007; 1.0	3.8; 0.06	

during CRF and U69,593)						
Intra-VTA U69,593 place conditioning, sham- vs shock-treated rats	5b	20 rats; 0; 0	Base-sham: 1.0; 0.8 Base-shock: 0.9; 0.6 Sham-test: 0.9; 0.5 Shock-test: 0.9; 0.4	Sham: 0.9; 0.4 Shock: 3.5; 0.08	Pretreatment-U69,593 interaction: 9.0; 0.008 (Two way mixed measures ANOVA) Pairwise comparisons: Sham, Base vs test: 0.0004 Shock, Base vs test: 0.6 (Bonferroni corrected)	
Baseline Ri in neurons from stress vs sham animals	Supp 2	139; 0; 0	1.0; 0.01	0.5; 0.5		0.2 (Permutation, mean)
Spontaneous AP duration in neurons from stress vs sham animals	Supp 2	81; 1; 0	0.9; 0.00002	0.9; 0.3		0.8 (Permutation, mean)
Spontaneous AP threshold in neurons from stress vs sham animals	Supp 2	81; 0; 0	1.0; 0.8	0.9; 0.3	9.5; 0.003 (one way ANOVA)	
Spontaneous AP peak in neurons from stress vs sham animals	Supp 2	81; 1; 0	1.0; 0.09	1.5; 0.2	0.2; 0.7 (one way ANOVA)	
I_h magnitude in neurons from stress vs sham animals	Supp 2	129; 17; 10	0.6; <0.000001	0.5; 0.5		0.4 (Permutation, mean)
Baseline spontaneous firing rate in neurons from stress vs sham animals	Supp 2	60; 5; 3	0.8; <0.000001	0.4; 0.5		0.3 (Permutation, mean)
Proportion of spontaneously active neurons from stress vs sham animals						0.9 (two tailed Fisher exact test)
Baseline Ri in neurons from stress animals grouped by U69,593 response type	Supp 3	46; 0; 0	1.0; 0.4	0.2; 0.8	3.2; 0.04 (one way ANOVA) Pairwise comparisons: Inhib-excit: 0.04 Inhib-no resp: 0.2 Excit-no resp: 0.9	
AP duration in neurons from stress animals grouped by U69,593 response type	Supp 3	27; 0; 0	0.9; 0.1	4.4; 0.02		3.1; 0.2 (Kruskal-Wallis rank sum test)
AP threshold in neurons from stress animals grouped by U69,593 response type	Supp 3	27; 0; 0	1.0; 0.4	1.5; 0.2	0.3; 0.7 (one way ANOVA)	
AP peak in neurons from stress animals grouped by U69,593 response type	Supp 3	27; 1; 0	0.9; 0.1	0.6; 0.5	0.8 (one way ANOVA)	

| CORT from recording animals, foot shock vs sham | Supp 6 | 16 rats; 1; 0 | 0.9; 0.2 | 0.09; 0.8 | 0.05 (one way ANOVA) | |

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5