Title: Complement C3 recognition by C3 convertases

Authors: Changhao Jia[1,3]†, Xiaoke Yang[2]†, Ming-hui Zhao[1]*, Ying Tan[1]*, Junyu Xiao[2,4]*

Affiliations:

[1]Renal Division, Department of Medicine, Peking University First Hospital; Institute of Nephrology, Peking University; Key Laboratory of Renal Disease, Ministry of Health of China; Key Laboratory of CKD Prevention and Treatment, Ministry of Education of China; Research Units of Diagnosis and Treatment of Immune-Mediated Kidney Diseases, Chinese Academy of Medical Sciences; Beijing, P.R. China.

[2]State Key Laboratory of Gene Function and Modulation Research, School of Life Sciences, Peking-Tsinghua Center for Life Sciences, Peking University, Beijing, P.R. China.

[3]Academy for Advanced Interdisciplinary Studies, Center for Biomed-X Research, Peking University, Beijing, P.R. China.

[5]Changping Laboratory, Beijing, P.R. China.

†These authors contributed equally to this work.

*Correspondence:

Ming-hui Zhao (mhzhao@bjmu.edu.cn), Ying Tan (tanying@bjmu.edu.cn), Junyu Xiao (junyuxiao@pku.edu.cn)

Abstract

The complement system plays a fundamental role in human immunity, and its dysregulation is implicated in numerous diseases. Activation of the complement occurs through three main pathways: classical, lectin, and alternative; which converge at the central component, C3. The classical and lectin pathways utilize the C4b2a convertase to cleave C3 and initiate complement activation, while the alternative pathway employs the C3bBb convertase, which is further stabilized by properdin. The molecular mechanisms governing C3 recognition by these convertase complexes remain incompletely understood. Here, we first present a 3.1 Å cryo-electron microscopy (cryo-EM) structure of the C4b2a–C3 complex, alongside the 2.9 Å and 3.1 Å structures of the C4b2 zymogen in the loading and activation states. These structural snapshots elucidate the structural basis for C3 engagement by C4b2a, and illustrate sequential conformational changes during the classical/lectin pathway convertase maturation. Furthermore, we determine a 2.6 Å cryo-EM structure of the C3bBb–properdin–C3 complex, which uncovers unique substrate-binding features of C3bBb and sheds light on how properdin stabilizes the alternative pathway convertase. These results offer comprehensive mechanistic insights into complement activation.

Main Text

The complement system is a vital component of innate immunity, comprising a complex network of over 50 soluble proteins and cell surface receptors that play essential roles in pathogen clearance, immune modulation, and the maintenance of homeostasis (*1, 2*). Dysregulation of the complement system has been associated with various human disorders, including renal, rheumatic, neurological, and cardiovascular diseases, etc. (*3-6*). More recently, complement dysregulation has also been implicated in SARS-CoV-2-related pathogenesis, particularly with respect to thrombosis (*7-8*).

The complement system operates through three distinct but interconnected activation pathways: the classical pathway, the lectin pathway, and the alternative pathway. Each of these pathways converges at the central component, C3, which is cleaved by the C3 convertases to produce the fragments C3a and C3b. C3a functions as an anaphylatoxin, triggering inflammatory responses through its interaction with the G protein-coupled receptor C3aR, while C3b acts as an opsonin that binds to pathogens, enhancing their recognition and clearance by phagocytes. Additionally, C3b serves as a critical subunit for the alternative pathway C3 convertase, promoting the amplification of the complement response.

C3 convertases are key enzymatic complexes within the complement system that play a crucial role in activating the complement cascade (Fig. 1A). There are two main forms of C3 convertase: C4b2a (here the convention where 'a' denotes the enzymatically active fragment of C2 is used) in the classical and lectin pathways, and C3bBb in the alternative pathway (*9, 10*). The C4b2a convertase is formed through the binding of C4b, generated by the serine protease C1s in the classical pathway or MASP-2 in the lectin pathway, to C2, which is further cleaved by C1s or MASP-2, resulting in the active C4b2a form. In contrast, the C3bBb convertase arises from the binding of C3b to factor B (FB), which is activated by factor D (FD). The resulting C3bBb convertase is stabilized by properdin (P) (*11*). Alternatively, a small fraction of C3 can undergo spontaneous hydrolysis, known as "C3 tick-over", yielding C3(H2O) that adopts a C3b-like conformation (*12*), and can thereby lead to the formation of the C3(H2O)Bb convertase. Recent studies have shown that granzyme K (GZMK) can also activate the complement cascade by cleaving C4 and C2, which leads to the formation of the C4b2a convertase (*13*), or by directly cleaving and activating C3 (*14*).

Both C4b2a and C3bBb are inherently labile, dissociating within minutes after formation (*15, 16*), posing significant challenges for their study. Previous breakthroughs, in particular the determination of the C3bB zymogen structure in complex with factor D (*17*) and the C3bBb– SCIN (Staphylococcal complement inhibitor) complex (*18*), have provided foundational

insights into the alternative pathway convertase assembly. The general location of the C3 substrate binding site on C3bBb was inferred from the crystallographic dimer interface formed between the two C3b molecules in the C3bBb–SCIN structure. However, mechanistic details underlying C3 recognition by the C3bBb convertase remain unclear. Similarly, the assembly and substrate engagement mechanisms of the classical/lectin pathway C4b2a convertase are poorly understood, limiting therapeutic targeting of these complexes.

In this study, we employ cryo-electron microscopy (cryo-EM) to resolve the molecular architecture of these convertases in functional states. We report a 3.1 Å structure of the C4b2a–C3 complex, elucidating the structural basis for C3 recognition by the classical/lectin pathway convertase. We also resolve the 2.9 Å and 3.1 Å structures of the C4b2 zymogen (standardized notation for the C4bC2 complex) in the loading and activation states, and thereby reveal sequential conformational transitions during the classical/lectin pathway convertase formation and maturation. Furthermore, we capture a 2.6 Å structure of the C3bBbP–C3 complex, which unveils distinct features governing substrate engagement by C3bBb. Detailed analysis also delineates how properdin stabilizes the C3b–Bb interaction, thereby positively regulating the alternative pathway convertase. These high-resolution snapshots address long-standing questions in complement biology, and provide a foundation for advancing the development of C3-targeted therapeutics.

Cryo-EM structure of the C4b2a–C3 complex

To capture a stable C3 convertase for cryo-EM analysis, we first isolated C3 and C4 from human plasma as previously described (*19*) (Fig. 1B, C). C4b was then generated by incubating C4 with C1s. Additionally, we produced a catalytically inactive mutant of C2, containing the S679A mutation, using HEK293F cells. Following this, C4b, C2, and C3 were mixed in an equal molar ratio, with Ni^{2+} included to enhance the C4b–C2 interaction (*20*). Afterwards, this sample was briefly treated with C1s and then rapidly plunged in liquid ethane for cryo-EM analysis. The C1s treatment resulted in partial cleavage of C2, leading to the formation of the C4b2a convertase, which readily engages C3 and enables us to trap the classical pathway C3 convertase-substrate complex and visualize the structure at a resolution of 3.1 Å (Fig. 2A; Fig. S1; Table S1). Additionally, we captured two functional states of the C4b2 pro-convertase containing intact C2: a loading state at 2.9 Å and an activation state at 3.1 Å.

In the C4b2a–C3 complex, C4b adopts an active conformation typical of C3b and C4b, resembling a "puppeteer" structure (*21*) (Fig. 2B). This includes a head (C345C domain), a neck (anchor), shoulders (MG7–8), a body (MG1–6, LNK), and a downward-extending arm

(CUB domain), with a "puppet" (TED domain) dropped to the level of MG1–MG4–MG5.

Compared to the structure of C4b alone (*22*), the C345C domain exhibits a slight rotation to

accommodate C2a (Fig. S2A). C2a binds to the C345C domain of C4b via its vWA domain,

while its serine protease (SP) domain swings toward C3. The substrate C3 adopts a

characteristic inactive conformation, with the TED domain securely tucked beneath the CUB

and MG8 domains, effectively concealing the Cys1010–Gln1013 thioester warhead (*23*) (Fig.

S2B). The ANA (anaphylatoxin) or C3a domain, consisting of four helices, points towards the

SP domain of C2a, facilitating the docking of the α'-NT segment, which contains the Arg748–

Ser749 cleavage site, into the catalytic center of C2a.

Substrate recognition mechanism of C4b2a

The structure of the C4b2a–C3 complex illustrates the specific recognition mechanism

employed by the classical/lectin pathway C3 convertase. Both C4b and C2a are involved in C3

recognition, contributing to the stringent substrate specificity. The interface between C4b and

C3 buries 2140 $Å^2$ from each molecule and involves multiple MG domains from both proteins

(Fig. 2C). Specifically, C4b-MG4 interacts with C3-MG5, facilitated by residues $Pro383_{C4b}$,

$Pro434_{C4b}$, and the $Leu514_{C3}$–$Pro518_{C3}$ strand (Fig. 2D). Between C4b-MG5 and C3-MG5,

$Pro478_{C4b}$ and $Asp479_{C4b}$ form interactions with $Leu461_{C3}$ and $Thr463_{C3}$, while $Arg481_{C4b}$ and

$Arg494_{C4b}$ engage with $Glu469_{C3}$ and $Asp513_{C3}$, respectively. Additional interactions occur

between C4b-MG5 and C3-MG4, where $Arg524_{C4b}$ is surrounded by $Asn412_{C3}$–$Pro415_{C3}$, and

$Thr525_{C4b}$–$Leu526_{C4b}$ pack against $Met368_{C3}$–$Pro369_{C3}$. The C4b-MG6 domain, which

consists of two split subdomains (MG6α and MG6β), interacts with three domains in C3 (Fig.

S2C): $Asp576_{C4b}$ forms a salt bridge with $Lys264_{C3}$ in C3-MG3; $Glu585_{C4b}$ bonds with

$Arg855_{C3}$ in C3-MG7; and $Arg828_{C4b}$ interacts with $Asp572_{C3}$ in C3-MG6. Additionally, C4b-

MG7 and C3-MG7 align closely, with $Arg831_{C4b}$, $Glu832_{C4b}$, and $Leu859_{C4b}$ accommodating

$Phe920_{C3}$. Overall, two dozen hydrogen bonds and salt bridge contacts are formed between

C4b and C3, ensuring their specific interaction.

The interaction between C2a and C3 buries 1360 $Å^2$ from each molecule. The α'-NT

segment (residues 743–756), which contains the scissile loop of C3, exhibits clear densities

and is positioned within the active site of C2a-SP (Fig. 2E; Fig. S2D). $Arg510_{C2a}$ contacts the

C-terminal end of the α4 helix in the ANA/C3a domain, facilitating the positioning of $Leu744_{C3}$

(P5 site). Residues $Glu556_{C2a}$, $Tyr558_{C2a}$, and $Lys676_{C2a}$ orient $Gly745_{C3}$–$Leu746_{C3}$–$Ala747_{C3}$

(P4–P2 residues) through hydrogen bond interactions with their main-chain groups, and

$Met648_{C2a}$ further packs on $Leu746_{C3}$. $Gly700_{C2a}$–$Asn703_{C2a}$, located at the beginning of the

long loop 2, accommodate Arg748$_{C3}$ (P1). Additionally, Lys676$_{C2a}$ interacts with the main-chain of Ser749$_{C3}$ (P1'), while Lys487$_{C2a}$ forms a salt bridge with Glu753$_{C3}$ (P5'). Notably, in the absence of C3 binding, Lys676$_{C2a}$ is drawn to Ser698$_{C2a}$, causing its main-chain carbonyl group to form a hydrogen bond with Ser679$_{C2a}$, which distorts the oxyanion hole (*24*). Upon C3 binding, Lys676$_{C2a}$ interacts with Ala747$_{C3}$ and Ser749$_{C3}$, leading to the establishment of a functional oxyanion hole (Fig. S2E). In addition to engaging the scissile loop of C3, C2a also anchors to the C3-MG3 domain (Fig. 2C). A pocket formed by Lys485$_{C2a}$, Leu517$_{C2a}$, Arg519$_{C2a}$, and Glu532$_{C2a}$ accommodates Arg290$_{C3}$, while Trp529$_{C2a}$ packs with His333$_{C3}$ (Fig. S2F). As a result of these interactions with both C2a and C4b, C3-MG3, along with the upper tip of C3-MG4, undergoes a ~25° rotation within the MG1–6 ring (Fig. S2B). Collectively, these interactions ensure the precise positioning of the Arg748$_{C3}$–Ser749$_{C3}$ peptide bond, priming it for cleavage by the Asp561$_{C2a}$–His507$_{C2a}$–Ser679(Ala)$_{C2a}$ catalytic triad.

Formation of the classical/lectin pathway C3 convertase

We also captured the structure of the C4b2 pro-convertase at the loading and activation states. Together with the C4b2a structure in complex with the C3 substrate, these structures allow us to visualize the formation of the C3 convertase at discrete steps (Fig. 3A).

The loading state of the C4b2 structure exhibits similarities to the factor B complexed with cobra venom factor (CVF) (*25*), a homolog of C3b (Fig. S3A). In this state, the C2-vWF region, namely the CCP1–3 domains or the C2b pro-segment, and the C2-vWA domain interact with C4b. In particular, C2-CCP1 and C2-vWA together engage the C4b-C345C domain. Notably, akin to the interaction observed between CVF and FB, the carboxyl group of the terminal Val1744$_{C4b}$ coordinates with the Ni^{2+} ion in the metal ion-dependent adhesion site (MIDAS) of C2-vWA, completing the metal ion coordination (Fig. 3B). Additionally, the CCP2–3 domains contact C4b-MG2, C4b-MG6, C4b-MG7, C4b-CUB, and the C4b-α'NT region (Fig. S3B). While the C2-SP domain exhibits flexibility in this state and cannot be clearly modeled despite the overall high resolution of the map, its general positioning suggests close proximity to the vWA domain (Fig. S3C). This arrangement would obstruct the access to the scissile site within the CCP3-vWA linker by the C1s protease, also reminiscent of the structural arrangement in the CVF-B complex.

The activation state of the C4b2 structure closely resembles the C3bB structure (*17*) (Fig. S3D), and provides a complete visualization of C2 (Fig. 3A). In this state, C2-SP adopts a stable conformation and is anchored to the MG2 and CUB domains of C4b. Additionally, the

vWA-SP linker in C2 is more ordered, and is positioned between C2-vWA and C2-SP (Fig. S3E). This change pulls C2-vWA closer to C2-SP, which in turn induces a rotation of the C4b-C345C domain. As a result, the interaction between C2-CCP1 and C4b-C345C is reduced (Fig. 3C; Fig. S3F). This displacement likely contributes to the release of C2b pro-segment following C2 cleavage. Models of C1s or MASP2 can be generated based on Factor D, utilizing the C3bBD complex structure as a template (Fig. 3D). In the resulting models, the scissile loop would be readily accommodated in the active sites of these proteases, facilitating C2 cleavage and C4b2 convertase maturation.

In the C4b2a–C3 complex, C2a adopts a conformation closely resembling that of the C2a alone structure (24) (Fig. S3G), with the C2a-SP drawn towards the substrate C3 as described above. Consequently, a tectonic reorientation of C2a is observed when compared to the C2a region in the activation state of C4b2 (Fig. 3E, Movie 1). The C2a-vWA domain shifts into the position previously occupied by CCP1–3, while the C2a-SP undergoes a ~118° swing. Thus, the CCP1-3 pro-segment must be released for the activated C2a to effectively dock onto C3. The large swing of the C2a-SP is further facilitated by a ~63° rotation of the α7 helix in the C2a-vWA domain, accompanied by a complete reorientation of the vWA-SP linker (Fig. 3E, Movie 1). The C2a-vWA domain remains attached to the C4b-C345C domain. Notably, in the C3-bound state, the carboxyl group of $Val1744_{C4b}$ is positioned further from the Ni^{2+} ion in the MIDAS motif of C2a-vWA, as evidenced by the density map (Fig. 3B). This suggests a loosened interaction between C4b and C2a, preparing C2a for release following C3 cleavage. It is likely that the smaller ionic radius of Mg^{2+} would further expedite this dissociation process *in vivo*. As such, engagement of C2a to C3 would prompt the concurrent release of C2a from C4b, explaining the labile nature of the C4b2a convertase.

Cryo-EM structure of the C3bBbP–C3 complex

To investigate C3 recognition by the AP C3 convertase, we generated C3b from C3 through limited trypsin processing (26). A catalytically inactive factor B containing the S699A mutation was prepared using HEK293 cells. Additionally, we produced a two-chained monomeric properdin by inserting a Tobacco Etch Virus (TEV) protease cleavage site between its TSR3 and TSR4 domains (Fig. 1B), and then processing the purified properdin oligomer with TEV, as previously described (27). Using a similar approach to the generation of the C4b2a–C3 complex, we mixed purified C3b, factor B, properdin, and C3 in equal molar ratios with Ni^{2+}. The mixture was then briefly treated with factor D, converting factor B into the active fragment Bb. Subsequently, cryo-EM analysis was conducted, resulting in the structure determination of

the C3bBbP–C3 complex at an atomic resolution of 2.6 Å (Fig. 4A, B; Fig. S4; Table S1).

In this complex, C3b and Bb individually closely resemble their respective structures observed in the C3bBb–SCIN (Staphylococcal complement inhibitor) complex (*18*) (Fig. S5A). It is apparent that Bb can now properly latch onto the substrate C3 without the steric hindrance imposed by SCIN. The C3 molecule adopts a conformation nearly identical to that seen in the C4b2a–C3 complex (Fig. S5B). Similar to the C4b2a–C3 complex, both C3b and Bb engage the C3 molecule. C3b forms a quasi-dimer with C3, involving the MG3–7 domains of both proteins. This C3b–C3 quasi-dimer shares similarities with the C3b–C3b homodimer observed in the C3bBb–SCIN crystal structure, especially in the lower region of the MG rings (Fig. S5C). However, the upper half of the MG ring including the MG2–3–6 domains in the C3 molecule, as well as the MG7 domain, undergoes rotations and leans further toward C3b in the C3b–C3 interaction. Compared to the C4b–C3 interface in the C4b2a–C3 structure, the C3b–C3 interaction involves fewer hydrogen bonds and salt bridge interactions (15 vs. 24), and the interface area is slightly smaller (2040 Å^2 vs. 2140 Å^2), suggesting that C4b forms a more optimized interaction with C3 than C3b.

In contrast, the interaction between Bb and C3 exhibits unique features and appears enhanced compared to the C2a–C3 interaction (Fig. 4C). For instance, beyond targeting the α'-NT segment and MG3 domain of C3, as seen with C2a, Bb also contacts C3-MG8 via several polar residues uniquely present in its loop 2. The β-hairpin comprising the first two β-strands in the Bb-SP domain are longer than the corresponding region in C2a-SP and effectively packs onto Asn750$_{C3}$–Asp752$_{C3}$ (P2'–P4' residues) by inducing the formation of a short strand (Fig. 4D, Fig. S5D). The tip of this β-hairpin also contacts C3-MG3. A hydrophobic pocket formed by Pro570$_{Bb}$ and Ala663$_{Bb}$–Tyr666$_{Bb}$ accommodates Leu744$_{C3}$ and Leu746$_{C3}$ (Fig. 4E). Gly720$_{Bb}$–Asp723$_{Bb}$ at the beginning of loop 2 holds Arg748$_{C3}$ in a similar manner as Gly700$_{C2a}$–Asn703$_{C2a}$, with the positive charge of Arg748$_{C3}$ better neutralized by Asp723$_{Bb}$. The Asp576$_{Bb}$–His526$_{Bb}$–Ser699(Ala)$_{Bb}$ triad adopts nearly identical positions to Asp561$_{C2a}$–His507$_{C2a}$–Ser679(Ala)$_{C2a}$, ready to attack the Arg748$_{C3}$–Ser749$_{C3}$ peptide bond.

Stabilization of C3bBb by properdin

Properdin is the only known positive regulator of the complement pathway. Previous studies have demonstrated that properdin primarily interacts with the C345C domain of C3b through two critical loops: the TSR5-stirrup (or thumb) and TSR6-stirrup (or index finger) (*28, 29*). Focused refinement in this region allows us to pinpoint how these two loops insert into the C3b–Bb interface (Fig. 5A). Notably, Arg330$_P$ in the TSR5-stirrup loop forms a salt bridge

with both $Asp1534_{C3b}$ and the main-chain oxygen of $Leu1533_{C3b}$, effectively positioning itself between $Leu1533_{C3b}$–$Asp1534_{C3b}$ on one side and $Pro1662_{C3b}$ on the other. This arrangement exerts pressure on $Pro1662_{C3b}$ and facilitates the engagement of the terminal $Asn1663_{C3b}$ with the metal ion (Ni^{2+} in this case) in the MIDAS motif of the Bb-vWA domain (Fig. 5B). Additionally, $Arg329_P$ in the TSR5-stirrup forms a hydrogen bond with the main-chain oxygen of $Leu349_{Bb}$, while stacking on $Phe1659_{C3b}$ as previously described (Fig. 5C) (*29*). $Ile340_P$ and $Pro341_P$ engage $Leu349_{Bb}$ through hydrophobic interactions. In the TSR6-stirrup, $Ser419_P$ and $Met420_P$ interact with $Tyr317_{Bb}$ and $Lys350_{Bb}$ via their main-chain oxygens. Collectively, these interactions from properdin serve to glue Bb onto C3b, thereby stabilizing the alternative pathway C3 convertase.

Discussion

   Targeting the complement C3 convertase represents a promising therapeutic strategy for various diseases, particularly those linked to complement dysregulation, such as certain kidney disorders. Pegcetacoplan, a member of the compstatin family of cyclic peptide C3 inhibitors, has been approved for treating paroxysmal nocturnal hemoglobinuria (PNH) and geographic atrophy (*30, 31*). Structural studies have shown that the compstatin molecules bind at the MG4–MG5 interface of C3 (*32, 33*). Consistent with previous analyses, compstatin molecules like Cp40, when engaged with C3, hinder the binding of C3 to either C4b2a or C3bBb, effectively blocking C3 activation across all complement pathways (Fig. 6A; Fig. S6). Notably, Cp40 can also bind to C3b, inhibiting the interaction between C3bBb and its C3 substrate, providing an additional layer of inhibitory effect. This dual mechanism makes compstatins ideally suited for broad-spectrum complement inhibition (Fig. 6A). Iptacopan, a factor B inhibitor, is another therapeutic agent used for treating PNH. Structural comparisons of the Bb-SP–Iptacopan complex (*34*) with our C3bBbP–C3 structure reveal that Iptacopan directly obstructs the entry of the α'-NT segment into the active site (Fig. 6B).

   Additionally, ongoing research is directed towards developing antibodies or nanobodies that specifically target C3 or the C3 convertases, examining these molecules in the context of our structural models to confirm their proposed mechanisms. For instance, both the S77 antibody and the hC3Nb1 nanobody bind to the C3b-MG7 domain (*35, 36*), disrupting the interaction between the C3bBb convertase and C3 (Fig. S6C). Furthermore, C4b-targeting nanobodies, such as NbE3, inhibit the C4b–C2a and C4b2a–C3 interactions, respectively (Fig. S6D) (*37*).

   In summary, we have delineated the molecular mechanisms by which C3 is recognized by

273   both C4b2a and C3bBb. Our results highlight distinct substrate-binding characteristics of the

274   classical/lectin and alternative convertases and elucidate the unique role of properdin in the

275   alternative pathway. These findings provide a fundamental understanding of these essential

276   molecular machineries and will guide the design and optimization of future therapeutic

277   molecules aimed at modulating complement activation.

References and Notes

Hajishengallis, G., Reis, E. S., Mastellos, D. C., Ricklin, D. & Lambris, J. D. Novel mechanisms and functions of complement. *Nat Immunol* **18**, 1288-1298, doi:10.1038/ni.3858 (2017).

Mastellos, D. C., Hajishengallis, G. & Lambris, J. D. A guide to complement biology, pathology and therapeutic opportunity. *Nat Rev Immunol* **24**, 118-141, doi:10.1038/s41577-023-00926-1 (2024).

Petr, V. & Thurman, J. M. The role of complement in kidney disease. *Nat Rev Nephrol* **19**, 771-787, doi:10.1038/s41581-023-00766-1 (2023).

Holers, V. M. Complement therapeutics are coming of age in rheumatology. *Nat Rev Rheumatol* **19**, 470-485, doi:10.1038/s41584-023-00981-x (2023).

Dalakas, M. C., Alexopoulos, H. & Spaeth, P. J. Complement in neurological disorders and emerging complement-targeted therapeutics. *Nat Rev Neurol* **16**, 601-617, doi:10.1038/s41582-020-0400-0 (2020).

Maffia, P., Mauro, C., Case, A. & Kemper, C. Canonical and non-canonical roles of complement in atherosclerosis. *Nat Rev Cardiol* **21**, 743-761, doi:10.1038/s41569-024-01016-y (2024).

Afzali, B., Noris, M., Lambrecht, B. N. & Kemper, C. The state of complement in COVID-19. *Nat Rev Immunol* **22**, 77-84, doi:10.1038/s41577-021-00665-1 (2022).

Cervia-Hasler, C. *et al.* Persistent complement dysregulation with signs of thromboinflammation in active Long Covid. *Science* **383**, eadg7942, doi:10.1126/science.adg7942 (2024).

Ricklin, D., Reis, E. S., Mastellos, D. C., Gros, P. & Lambris, J. D. Complement component C3 - The "Swiss Army Knife" of innate immunity and host defense. *Immunol Rev* **274**, 33-58, doi:10.1111/imr.12500 (2016).

Geisbrecht, B. V., Lambris, J. D. & Gros, P. Complement component C3: A structural perspective and potential therapeutic implications. *Semin Immunol* **59**, 101627, doi:10.1016/j.smim.2022.101627 (2022).

Pedersen, D. V., Lorentzen, J. & Andersen, G. R. Structural studies offer a framework for understanding the role of properdin in the alternative pathway and beyond. *Immunol Rev* **313**, 46-59, doi:10.1111/imr.13129 (2023).

Gadeberg, T. A. F. *et al.* Cryo-EM analysis of complement C3 reveals a reversible major opening of the macroglobulin ring. *Nat Struct Mol Biol*, doi:10.1038/s41594-024-01467-4 (2025).

Donado, C. A. *et al.* Granzyme K activates the entire complement cascade. *Nature*, doi:10.1038/s41586-025-08713-9 (2025).

Lan, F. *et al.* GZMK-expressing CD8(+) T cells promote recurrent airway inflammatory

315   diseases. *Nature* **638**, 490-498, doi:10.1038/s41586-024-08395-9 (2025).

316   15   Kerr, M. A. The human complement system: assembly of the classical pathway C3 convertase.
317   *Biochem J* **189**, 173-181, doi:10.1042/bj1890173 (1980).

318   16   Pangburn, M. K. & Muller-Eberhard, H. J. The C3 convertase of the alternative pathway of
319   human complement. Enzymic properties of the bimolecular proteinase. *Biochem J* **235**, 723-
320   730, doi:10.1042/bj2350723 (1986).

321   17   Forneris, F. *et al.* Structures of C3b in complex with factors B and D give insight into
322   complement convertase formation. *Science* **330**, 1816-1820, doi:10.1126/science.1195821
323   (2010).

324   18   Rooijakkers, S. H. *et al.* Structural and functional implications of the alternative complement
325   pathway C3 convertase stabilized by a staphylococcal inhibitor. *Nat Immunol* **10**, 721-727,
326   doi:10.1038/ni.1756 (2009).

327   19   Gadjeva, M. The complement system. Overview. *Methods Mol Biol* **1100**, 1-9,
328   doi:10.1007/978-1-62703-724-2_1 (2014).

329   20   Fishelson, Z. & Muller-Eberhard, H. J. C3 convertase of human complement: enhanced
330   formation and stability of the enzyme generated with nickel instead of magnesium. *J Immunol*
331   **129**, 2603-2607 (1982).

332   21   Janssen, B. J., Christodoulidou, A., McCarthy, A., Lambris, J. D. & Gros, P. Structure of C3b
333   reveals conformational changes that underlie complement activity. *Nature* **444**, 213-216,
334   doi:10.1038/nature05172 (2006).

335   22   Mortensen, S. *et al.* Structural Basis for the Function of Complement Component C4 within
336   the Classical and Lectin Pathways of Complement. *J Immunol* **194**, 5488-5496,
337   doi:10.4049/jimmunol.1500087 (2015).

338   23   Janssen, B. J. *et al.* Structures of complement component C3 provide insights into the function
339   and evolution of immunity. *Nature* **437**, 505-511, doi:10.1038/nature04005 (2005).

340   24   Milder, F. J. *et al.* Structure of complement component C2A: implications for convertase
341   formation and substrate binding. *Structure* **14**, 1587-1597, doi:10.1016/j.str.2006.08.008
342   (2006).

343   25   Janssen, B. J. *et al.* Insights into complement convertase formation based on the structure of
344   the factor B-cobra venom factor complex. *EMBO J* **28**, 2469-2478,
345   doi:10.1038/emboj.2009.184 (2009).

346   26   Wu, J. *et al.* Structure of complement fragment C3b-factor H and implications for host
347   protection by complement regulators. *Nat Immunol* **10**, 728-733, doi:10.1038/ni.1755 (2009).

348   27   Pedersen, D. V. *et al.* Functional and structural insight into properdin control of complement
349   alternative pathway amplification. *EMBO J* **36**, 1084-1099, doi:10.15252/embj.201696173
350   (2017).

28   Pedersen, D. V. *et al.* Structural Basis for Properdin Oligomerization and Convertase

Stimulation in the Human Complement System. *Front Immunol* **10**, 2007,
doi:10.3389/fimmu.2019.02007 (2019).

29   van den Bos, R. M., Pearce, N. M., Granneman, J., Brondijk, T. H. C. & Gros, P. Insights Into
Enhanced Complement Activation by Structures of Properdin and Its Complex With the C-
Terminal Domain of C3b. *Front Immunol* **10**, 2097, doi:10.3389/fimmu.2019.02097 (2019).

30   Lamers, C., Mastellos, D. C., Ricklin, D. & Lambris, J. D. Compstatins: the dawn of clinical
C3-targeted complement inhibition. *Trends Pharmacol Sci* **43**, 629-640,
doi:10.1016/j.tips.2022.01.004 (2022).

31   Mastellos, D. C. & Lambris, J. D. Pegcetacoplan (Syfovre(TauMu)): targeting complement C3
in geographic atrophy. *Trends Pharmacol Sci* **45**, 1175-1176, doi:10.1016/j.tips.2024.10.001
(2024).

32   Janssen, B. J., Halff, E. F., Lambris, J. D. & Gros, P. Structure of compstatin in complex with
complement component C3c reveals a new mechanism of complement inhibition. *J Biol Chem*
282, 29241-29247, doi:10.1074/jbc.M704587200 (2007).

33   Lamers, C. *et al.* Insight into mode-of-action and structural determinants of the compstatin
family of clinical complement inhibitors. *Nat Commun* **13**, 5519, doi:10.1038/s41467-022-
33003-7 (2022).

34   Schubart, A. *et al.* Small-molecule factor B inhibitor for the treatment of complement-mediated
diseases. *Proc Natl Acad Sci U S A* **116**, 7926-7931, doi:10.1073/pnas.1820892116 (2019).

35   Katschke, K. J., Jr. *et al.* Structural and functional analysis of a C3b-specific antibody that
selectively inhibits the alternative pathway of complement. *J Biol Chem* **284**, 10473-10479,
doi:10.1074/jbc.M809106200 (2009).

36   Jensen, R. K. *et al.* A potent complement factor C3-specific nanobody inhibiting multiple
functions in the alternative pathway of human and murine complement. *J Biol Chem* **293**, 6269-
6281, doi:10.1074/jbc.RA117.001179 (2018).

37   De la, O. B. K. I. *et al.* Multifaceted Activities of Seven Nanobodies against Complement C4b.
J Immunol **208**, 2207-2219, doi:10.4049/jimmunol.2100647 (2022).

38   Punjani, A., Rubinstein, J. L., Fleet, D. J. & Brubaker, M. A. cryoSPARC: algorithms for rapid
unsupervised cryo-EM structure determination. *Nat Methods* **14**, 290-296,
doi:10.1038/nmeth.4169 (2017).

39   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583-
589, doi:10.1038/s41586-021-03819-2 (2021).

40   Krishnan, V., Xu, Y., Macon, K., Volanakis, J. E. & Narayana, S. V. The crystal structure of C2a,
the catalytic fragment of classical pathway C3 and C5 convertase of human complement. *J Mol*
Biol **367**, 224-233, doi:10.1016/j.jmb.2006.12.039 (2007).

41   Pettersen, E. F. *et al.* UCSF Chimera--a visualization system for exploratory research and
analysis. *J Comput Chem* **25**, 1605-1612, doi:10.1002/jcc.20084 (2004).

42 Emsley, P., Lohkamp, B., Scott, W. G. & Cowtan, K. Features and development of Coot. *Acta*
Crystallogr D Biol Crystallogr **66**, 486-501, doi:10.1107/S0907444910007493 (2010).
43 Adams, P. D. *et al.* PHENIX: a comprehensive Python-based system for macromolecular
structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221,
doi:10.1107/S0907444909052925 (2010).
44 Pettersen, E. F. *et al.* UCSF ChimeraX: Structure visualization for researchers, educators, and
developers. *Protein Sci* **30**, 70-82, doi:10.1002/pro.3943 (2021).
396

Acknowledgments:

We are grateful to the Cryo-EM and the high-performance computing platforms of Peking

University and Changping Laboratory for support with data collection and computation. We

also thank the National Center for Protein Sciences at Peking University for assistance with the

AKTA facility.

Funding:

The National Natural Science Foundation of China (32325018)

National Key R&D Program of China (2024YFA1306203)

The Qidong-SLS Innovation Fund, and the Frontier Innovation Fund of Peking University

Chengdu Academy for Advanced Interdisciplinary Biotechnologies to J.X.

Capital's Funds for Health Improvement and Research (2024-2-4076) to Y.T.

Author contributions:

C.J. and X.Y. made equal contributions to this work.

Protein purification and complex assembling: C.J., X.Y.

Cryo-EM sample preparation, data acquisition, image processing, structure determination:

X.Y., C.J.

Funding acquisition: J.X., Y.T., M.H.Z.

Project administration: J.X., Y.T., M.H.Z.

Supervision: J.X., Y.T., M.H.Z.

Writing – original draft: C.J., X.Y.

Writing – review & editing: J.X., Y.T., M.H.Z.

Competing interests:

The authors declare no competing interests.

Data and materials availability:

Cryo-EM density maps have been deposited in the Electron Microscopy Data Bank with

accession codes EMD-63893 (C4b2 in loading state), EMD-63892 (C4b2 in activation state),

EMD-63894 (C4b2a–C3 complex) and EMD-63895 (C3bBbP–C3 complex). Structural

coordinates have been deposited in the Protein Data Bank (PDB) with the accession codes

9U60, 9U5Z, 9U61 and 9U62. The other structural coordinates involved in this study are

available from PDB (2ODP, 2WIN, 6S0B, 6EHG, 3G6J, 6RAV, 1ELV, 5JPM, 2XWB and

3HS0).

Fig. 1: Overview of the complement C3 convertases.

(A) A simplified diagram of the complement cascade.

(B) Schematic representations of the domain organization of C3 convertase components, including C4, C2, C3, factor B (FB), and properdin. MG: Macroglobulin domain; LNK: Linker domain; α'-NT: C3 α' chain N-terminal; CUB: Complement C1r-C1s, UEGF, BMP1; TED: Thioester domain; vWF: von Willebrand factor; vWA: von Willebrand A domain; SP: Serine protease domain. The yellow connecting lines indicate interchain disulfide bonds, while the triangular arrows denote cleavage sites on C4 and C3.

(C) SDS-PAGE analysis of purified proteins, including C4, C4b, C2, C3, C3b, FB, properdin, and factor D (FD). Asterisks (*) indicate contaminant proteins.

Fig. 2: Cryo-EM structure of the classical/lectin pathway C3 convertase.

(A) Two views of the cryo-EM reconstruction of the C4b2a–C3 complex. Complement proteins C4b, C3, and C2b are depicted in blue, orange, and green, respectively, while the C3a(ANA) domain is highlighted in red.

(B) A ribbon diagram of the C4b2a–C3 structure accompanied by a schematic representation.

(C) The interaction between C4b and C3 involves multiple MG domains from both proteins.

(D) Interaction details between C3-MG4, C3-MG5 and C4b-MG5, C4b-MG4.

(E) Interaction between the C3 scissile loop and the C2a-SP domain. C3a(ANA) and α'-NT are shown in red and orange, respectively. The cleavage site is indicated by a black arrowhead. The oxanion hole in C2a-SP is highlighted in blue, and the catalytic triad Ser679(A), His507, Asp561 are highlighted in yellow. The hydroxyl group of Ser679 is hypothetical and is represented in a lighter color.

Fig. 3: Formation of the classical/lectin pathway C3 convertase.

(A) The C3 convertase at three distinct steps: loading, activation, and C3-engaged. C4b, C2b (CCP1–3), and C2a (vWA–SP) are depicted in purple, salmon and green, respectively. In the loading state, the C2a-SP domain exhibits flexibility and cannot be clearly modeled, and is represented by a dashed circle.

(B) The positions of C4b-V1744 relative to the Ni^{2+} ion in C2-vWA. Ser262, Ser264, Thr337 and Asp376 in the MIDAS motif of C2-vWA are highlighted in orange. The density maps are shown at the same contour level.

(C) Structural comparison between the loading and activation states of C4b2. The loading state is presented in gray, while the activation state uses the same color scheme as in panel a.

(D) Structural models of C1s or MASP-2 bound to C2. These models are generated by aligning

466   the activation state of the C4b2 structure with that of C3bBD (PDB: 2XWB) and then

467   aligning the structures of C1s (PDB: 1ELV) and MASP-2 (PDB: 5JPM) to Factor D.

468   **(E)** Structural comparison between the activation and C3-engaged states. The C3-engaged state

469   is shown in gray, while the activation state is presented in color.

Fig. 4: Cryo-EM structure of the alternative pathway C3 convertase.

(A) Two views of the C3bBbP–C3 complex cryo-EM reconstruction. C3b, Bb, and C3 are depicted in khaki, light blue, and orange, respectively; while properdin is shown in aquamarine. C3a is highlighted in red.

(B) Ribbon rendering of the C3bBbP–C3 structure.

(C) Interaction between Bb-SP and C3, with Bb-SP, C3a, α'-NT, C3-MG3, C3-MG8 shown in light blue, red, orange, cyan, and dark green, respectively. The β-hairpin in Bb-SP is highlighted in pink.

(D) The β-hairpin in Bb-SP is longer than that of C2a-SP, inducing the formation of a short strand in C3.

(E) Interaction between the C3 scissile loop and Bb-SP.

Fig. 5: Stabilization of C3bBb by properdin.

(A) The TSR5-stirrup and TSR6-stirrup loops of properdin promote the interaction between C3b and Bb. Bb-vWA, C3b-C345C, and properdin are shown in light blue, khaki and aquamarine, respectively. The density map of the structural interface is shown below.

(B) Arg330$_P$ facilitates the interaction between C3b and the metal ion in the MIDAS motif of Bb-vWA.

(C) Additional interactions between properdin and C3bBb.

Fig. 6: Insights into therapeutic drugs targeting the C3 convertase.

(A) Mechanism of CP40 inhibition on the engagement of C3 with C4b2a and C3bBb. The C4b2a–C3 inhibition model is generated by superimposing C3b-Cp40 (PDB: 7BAG) onto the C3 structures of the C4b2a–C3 complex (left panel). The C3bBbP–C3 inhibition model is created by superimposing C3b-Cp40 onto both C3b and C3 (right panel). The binding of CP40 would create steric clashes that hinder C3 binding to either C4b2a or C3bBb.

(B) Mechanism of Iptacopan inhibition on the engagement of C3 with C3bBbP. The inhibition model is generated by superimposing Bb-Iptacopan (PDB: 6RAV) onto the structure of Bb-C3a-α'-NT, which is separated from the C3bBbP–C3 complex. Iptacopan is represented in space-filling format.

Fig. S1: Cryo-EM 3D reconstruction of the C4b2a–C3 complex, C4b2(loading) and C4b2(activation).

(A) Flowchart of cryo-EM data processing for C4b2a–C3, C4b2 (loading) and C4b2(activation). The density maps of C4b2a–C3 and C4b2 were generated from the same data set.

(B)-(D) Angular particle distribution heat map and Gold-standard Fourier shell correlation

508   (GSFSC) curves of C4b2a–C3, C4b2 (loading) and C4b2(activation), respectively.

509   **(E)-(G)** Resolution estimations for the final maps of C4b2a–C3, C4b2 (activation) and C4b2

510   (loading), respectively.

Fig. S2: Structure analysis of C4b2a–C3 complex.

(A) C4b structure in the C4b2a–C3 complex compared to C4b alone (PDB: 5jtw).

(B) C3 structure in the C4b2a–C3 complex compared to C3 alone (PDB: 2A73).

(C) C4b-MG6 and C4-MG7 interact with C3-MG3, C3-MG6, and C3-MG7.

(D) Density map of C3-ANA and α'-NT. Black arrow indicates the cleavage site between R748 and S749.

(E) A functional oxyanion hole is established in C2a-SP upon C3 binding, when compared to the C2a alone structure (PDB: 2i6q).

(F) Interaction between C2a-SP and C4b-MG3.

Fig. S3: Activation of classical pathway C3 convertase.

(A) Structural comparison of CVF-B (PDB: 3hs0) and the loading state of C4b2. CVF-B is shown in yellow, and C4b2 is shown in purple, green and salmon.

(B) Interaction between the vWF domain of C2 and C4b in the loading state.

(C) A low resolution density map of the C4b2 loading state reveals the general location of the C2a-SP.

(D) Structural comparison of C3bB (PDB: 3xwb) and the activation state of C4b2. C3bB is shown in grey, and the C4b2 activation state structure is shown in purple, green and salmon.

(E) The linker between C2a-vWA and C2a-SP domain undergo conformational changes from the loading state to activation state, and subsequently to C3-bound state.

(F) The docking of C2-SP onto C4b results in reduced interaction between C2a-vWF and C4b-C345C, facilitating the detachment of C2a-vWF.

(G) Structure comparison of C2a in the C3-bound state and the C2a alone structure (PDB: 2i6q).

Fig. S4: Cryo-EM 3D reconstruction of the C3bBbP–C3 complex.

(A) Flowchart of cryo-EM data processing for the C3bBbP–C3 complex.

(B)-(D) Resolution estimations for the final maps of the C3bBbP–C3 complex and the focused map of C3bBbP–C3 complex, respectively.

(E)-(G) Angular particle distribution heat map and Gold-standard Fourier shell correlation (GSFSC) curves of the C3bBbP–C3 complex and the focused map of C3bBbP–C3 complex, respectively.

Fig. S5: Cryo-EM structure analysis of the C3bBbP–C3 complex.

(A) Structural superimposition of the C3bBbP–C3 and C3bBb–SCIN (PDB: 2win). C3bBb–SCIN is shown in pink. C3 in the C3bBbP–C3 complex is shown using a surface representation. The presence of SCIN would prevent the binding of C3 to C3bBb due to steric hindrance.

(B) Structural superimposition of C3bBbP–C3 and C4b2a–C3.

(C) Structural comparison between the C4b-C3 heterodimer, C3b-C3 quasi-dimer and C3b-C3b homodimer. Red boxes indicate MG-rings within C3 and C3b.

(D) Interactions between the Bb-SP hairpin structure and C3 α'-NT.

Fig. S6: C3 inhibitory molecules.

(A) Chemical structure of the cyclic peptide CP40.

(B) Chemical structure of the Iptacopan (Fabhalta).

(C) Structure of hC3bN1-C3b and S77-C3b. hC3bN1 is a C3b-MG7 binding nanobody, whereas S77 is a C3b-MG7 binding antibody. Both hC3bN1 and S77 would inhibit C3bBb by obstructing the binding of the substrate C3 molecule.

(D) Structure of NbE3-C4b. NbE3 would block the binding C3 to C4b and thereby inhibit the C4b2a convertase.

562 **Table S1. Cryo-EM data collection, refinement and validation statistics.**

	C4b2_loading	C4b2_activation	C4b2a_C3	C3bBbP_C3
Data collection and processing				
Magnification		130,000		
Voltage (kV)		300		
Electron exposure (e–/Å^2)		60		
Defocus range (μm)		−1.1 to −1.5		
Pixel size (Å)		0.95		
Symmetry imposed		C1		
Initial particle images (no.)	2,093,466	2,093,466	453,264	848,419
Final particle images (no.)	109,606	36,0922	109,455	389,342
Map resolution (Å) FSC threshold: 0.143	2.91	3.06	3.09	2.57
Refinement				
Initial model used (PDB code)	Alphafold2	Alphafold2	Alphafold2 2ODP	Alphafold2 2WIN, 6S0B
Model resolution (Å) FSC threshold: 0.143	2.91	3.06	3.09	2.57
Map sharpening B factor (Å^2)	62.0	44.0	59.6	83.9
Model composition				
Non-hydrogen atoms	14,719	16,447	29,000	29,466
Protein residues	1,899	2,116	3,674	3,709
Ligands	1	1	1	1
B factors (Å^2)				
Protein	81.19	122.56	92.51	458.80
Ligand	106.13	130.72	100.96	127.37
R.m.s. deviations				
Bond lengths (Å)	0.004	0.007	0.005	0.004
Bond angles (°)	0.813	1.141	0.837	0.829
Validation				
MolProbity score	1.62	1.76	1.41	1.69
Clashscore	2.87	4.74	3.15	3.25
Poor rotamers (%)	0.56	0.28	0.41	3.30
Ramachandran plot				
Favored (%)	95.67	91.05	95.65	96.92
Allowed (%)	4.33	8.95	4.33	3.08
Disallowed (%)	0.00	0.00	0.03	0.00