Profiling of *Burkholderia pseudomallei* variants derived from

Queensland's clinical isolates

Pauline M.L. Coulon[1], Kay Ramsay[3], Aven Lee[2], Edita Ritmejeryte[2], Miranda E. Pitt[1], Joyce

To[1], Daniel G. Mediati[1], Ian Gassiep[3], Sarah Reed[2], Patrick N. A. Harris[3] and Garry S.A.

Myers[1]

Affiliations:

[1]Australian Institute for Microbiology and Infection, Faculty of Science, University of

Technology Sydney, Ultimo, NSW, Australia

[2]Mass Spectrometry Facility, University of Queensland, Centre for Clinical Research,

Queensland, Australia

[3]The University of Queensland Centre for Clinical Research (UQCCR), Faculty of Medicine,

The University of Queensland, Queensland, Australia

Corresponding author:

Pauline M.L. Coulon, Australian Institute for Microbiology and Infection, Faculty of Science,

University of Technology Sydney, NSW, Australia, +61390353555,

pauline.coulon@uts.edu.au

Abstract

Burkholderia pseudomallei (*Bp*), an environmental bacterium and opportunistic pathogen endemic to tropical regions, is highly adaptive and thrives in diverse environments, from soil to human hosts. Bacterial adaptation is critical for survival, virulence modulation, and persistence during infection and can manifest as colony morphotype variation (CMV). While Bp adaptation has been well studied, CMV remains poorly understood. Here, we characterized five clinical Bp isolates exhibiting heterogeneous populations with rough and smooth colony morphologies. We used phenotypic assays, whole-genome sequencing, and proteomics to investigate the molecular pathways impacted by CMV - by comparing smooth and rough morphotypes. While phenotypic differences in protease activity, haemolysis, mucoid, iron uptake and antibiotic sensitivity —including to antimicrobial agents commonly used to treat infections—were rare, these traits alone could not distinguish morphotypes or group of isolates. Genomic comparisons revealed either no differences or limited isolate-specific mutations, which does not explain the overall difference in phenotypes. In contrast, proteomic analysis uncovered consistent shifts in protein abundance related to virulence, including quorum sensing, DNA methylation, and secretion systems. Rough variants showed higher expression of EPS-associated proteins, the BpsI3/R3 quorum sensing system, and the global regulator ScmR, whereas smooth variants upregulated type III/VI secretion and siderophore biosynthesis pathways. These findings suggest that CMV is driven by phase variation and regulatory mechanisms rather than punctual genomic modifications. Our study underscores the limitations of phenotype or genome-based classification alone in the context of CMV and highlights the value of integrated multi-omics approaches to uncover CMV-associated biomarkers, with potential applications in diagnostics and the development of targeted therapies against persistent and drug-resistant Bp infections.

Keywords: colony morphotype variation, epigenetics, quorum sensing, omics

Importance

Burkholderia pseudomallei (Bp), the causative agent of melioidosis, is endemic to Australia, Asia, Africa, and the Americas. It predominantly affects Indigenous populations and individuals with diabetes, chronic lung or kidney disease, or alcoholism. *Bp* is known for its exceptional genomic and phenotypic plasticity, enabling rapid adaptation to diverse environments. This adaptability is reflected by colony morphotype variation (CMV), including reversible phase variation between smooth and rough colonies. In this study, we report rough and smooth colonies from clinical samples and emphasize the importance of characterizing CMV through multi-omics approaches rather than relying solely on genomics and phenotypic traits. By integrating genomics, phenotypic and proteomic data, we identified that a limited number of mutations-including one in a putative regulatory element-likely drive major molecular changes between morphotypes. These affect the expression of quorum sensing systems, the transcriptional regulator ScmR, DNA methyltransferase, and virulence-associated genes.

Introduction

Burkholderia are highly adaptable Gram-negative environmental saprophyte bacteria found in diverse ecological niches ranging from soil and plant roots to human hosts. Their remarkable phenotypic plasticity enables them to persist in various ecological niches, often facilitated by transmission through contaminated water, soil, and products (1-7). Among them, *Burkholderia pseudomallei* (*Bp*) poses a significant public health threat as the causative agent of melioidosis, a severe and often fatal disease in animals and humans (1, 8). While *Bp* is endemic to tropical and sub-tropical regions of Asia, Africa, Central and South America

and Australia (1), it has been recognized since 2023 as naturally established in the

environment of the United States (9).

Although most healthy individuals exposed to *Bp* do not develop melioidosis,

vulnerable population with underlying conditions such as cystic fibrosis (in rare occurrence),

diabetes, chronic kidney disease, chronic lung disease, alcoholism or with a compromised

immune system often -are the most at risks. In these individuals, *Bp* infections can cause

organ abscesses or severe pneumonia, often leading to fatal sepsis with a mortality rate

reaching up to 52% (1, 10-17). The incidence of *Bp* infections is strongly correlated with

heavy rains and flooding, and cases are expected to increase as climate change intensifies

extreme weather events (18). In mid-2025, Queensland (Australia), reports it worst year of

melioidosis in three decades with over 230 cases and 31 deaths (Queensland Health).

Additionally, there is a growing concern that melioidosis cases could (1) spread into Central

Brisbane and New South Wales (Australia), where the disease is less recognised by

healthcare providers and (2) be transmitted directly by pets, livestock or wild animals through

contact with infected wounds or via shared environment–even if no zoonotic transmission

have been confirmed to date (8).

Burkholderia's rapid adaptation to environmental changes and stressors is often associated

with modifications in outer membrane proteins, antibiotic resistance, and virulence factors

(19-23). This adaptability is reflected by colony morphotype variation (CMV), a process by

which bacteria switch between distinct colony morphologies. CMV is now used as an

umbrella term encompassing reversible changes (phase variation without genomic plasticity),

irreversible genomic plasticity (e.g., chromosome loss, (24)), antigenic variation, and

phenotypic switching (25, 26). Although CMV plays a critical role in persistence during

infection, it remains underexplored. While a few studies have shown that CMV can result in

mutations in global regulators, two-component systems, genome reduction and duplication,

bacteriophage cluster integration and DNA methylation, as reviewed by Coulon and

colleagues (26).

Among *Burkholderia* species, *Bp* has been the most extensively studied in terms of CMV.

Chantratita and colleagues identified up to seven dynamic and reversible colony

morphologies in *Bp* (19). Under laboratory conditions, these variants emerge in response to

stress conditions and revert to the parental morphotype once the stress is removed (19, 23,

27). Notably, CMVs also occur during infection: in 10% out of over 450 clinical *Bp* isolates,

a mixed mucoid and non-mucoid CMV population was observed on blood agar. The mucoid

phenotype was attributed to a difference in O-antigen Lipopolysaccharides (LPS; [OPS])

production, while no mutations or altered expression of the *wbiA* O-antigen acetylase was

linked (28). Further, the emergence of small-colony variants (SCVs) from a wrinkled *Bp*

isolate during long-term infections in murine was attributed to upregulation of the LPS

biosynthesis cluster genes (29, 30). Using experimental evolution in the same model, Gierok

and colleagues have demonstrated the emergence of two reversible yellow colony variants

(YA and YB) from the environmental *Bp* K96243. These variants showed upregulation of the

σ-54 dependent regulator *yelR,* leading to resistance to hypoxic stress and enhanced

colonization and persistence in the murine stomach (31).

In this study, we characterised five additional *Bp* isolates undergoing CMV using phenotypic,

genomic, and proteomic profiling to uncover the similarities and dissimilarities in molecular

pathways impacted by CMV.

Ethical Approval

This study received ethical approval from the Royal Brisbane and Women's Hospital Ethics

Committee (LNR/2020/QRBW/65573), with site-specific authority obtained from the

Townsville Hospital and Health Service and approval under the Queensland Public Health

Act.

Material and Method

Colony morphology screening

In brief, 292 Bp isolates were collected from Townsville between 1999 and 2022 (32).

Among those classified as 'mixed population' (Etipola and colleagues, unpublished), five

isolates were selected for this study. Among the isolates listed in **Table S1**, each was streaked

on Luria Broth (LB) agar plates containing 0.01% Congo red (CRLA) from -80°C stock and

incubated at 37°C for three overnights – to confirm the mixed population phenotype.

Phenotypic assays

Production of virulence factors. Individual colonies (four biological replicates) were picked

and stabbed into LB agar plate containing 1.5% skimmed milk to assess proteolytic activity

(33); 0.5 g/L yeast extract supplemented with 4g/L D-mannitol (YEM) and agar to assess

mucoid (34); a modified recipe of Chrome Azurol S agar (using 10X chelated LB instead of

minimal media 9 supplemented with chelated casamino acid) to assess siderophore

production (35) and ensure the growth of clinical isolates; and Columbia horse blood agar

plate to assess hemolysis activity (36). Plates were incubated at 37°C for 3 days. Inhibition

circles were measured and confirmed by statistical analyses (see specific text in the legend of

Table 1).

Antimicrobial sensitivity assay. Individual colonies were resuspended in 0.9% NaCl solution

to achieve a turbidity of 0.5-0.63 McFarland. Each bacterial suspension was then spread on

Muller Hinton II agar plates to assess susceptibility to meropenem using gradient MIC test

strips (ETEST, range0.002-32 µg/mL; Biomérieux), and disk diffusion for ceftazidime

(10µg) and sulfamethoxazole-trimethoprim (25µg) (Oxoid). Antimicrobial susceptibility was

determined according to EUCAST guidelines (v15). For CMVs from the same isolate

showing considerable visual changes in antibiotic susceptibility, three biological replicates

were conducted on independent days, and the average inhibition zone diameters were used

for susceptibility classification.

Statistical analyses. All datasets were first tested for normality using the Shapiro–Wilk test.

Based on the results, comparisons of virulence factor production between rough and smooth

CMVs were performed using either a Wilcoxon rank-sum test or a t-test, with a Bonferroni

adjusted pvalue.

Genomic analysis

gDNA extraction. Genomic DNA was extracted from a mixed of subsamples of three

macrocolonies grown for three nights at 37 °C on CRLA using either gDNA spin or high

molecular weight extraction kits (NEB). Short-read sequencing (Illumina) was conducted on

all CMVs. Briefly, libraries were prepared using 1 ng of gDNA via the Nextera XT kit and

sequenced using the NovaSeq XPlus platform (37). On select isolates, long-read sequencing

(Oxford Nanopore Technologies) was implemented. Libraries were prepared using 400 ng

gDNA with the SQK-NBD114.24 kit and sequenced using R10.4.1 flow cells on the GridION

platform. Reads were base-called using Dorado 7.6.7.

Genome assembly. Nanopore and Illumina reads were quality-filtered and trimmed using

Nanofilt (v.2.8.0; (38)) and fastp (v0.24.1; (39)) respectively. Nanopore reads were

assembled using Autocycler (v0.4.0; (40)). Assemblies were polished using Polypolish

(v0.6.0; (41)) and then POLCA from the pypolca toolkit (v3.0.1; (42)) with Illumina reads.

Final assemblies were annotated using Bakta (v1.11; (43)). For the second CMV from each

isolate, Illumina reads were assembled into contigs using Unicycler (short read option,

v.0.5.1; (44)).

Genomic modification analyses. To identify SNPs and indels, Illumina reads from the second

CMV were mapped onto the corresponding first CMV assembly using Snippy (v4.6.0,

haploid mode; - Galaxy implementation) (45). As a control, Illumina reads from the first

CMV were also mapped onto their own assemblies, confirming the absence of variation.

Prediction of regulatory targets in non-coding regions. At least 50-nt of the upstream and

downstream sequences of non-coding genomic variants (including SNPs, insertions or

deletions) were extracted and used as input into IntaRNA (46) to predict regulatory targets in

Bp. RNA-RNA interactions were filtered for those occurring within 50-nt upstream and 25-nt

downstream of mRNA start codons (AUG), at least 7 nucleotides of base-pairing, and

containing no G-U base-pairs at the 5' or 3' seed ends.

Prediction of secondary structure in non-coding regions. The non-coding genomic variant

regions (50-nt upstream and downstream, as *above*) were extracted and used as input into the

RNAfold software package to calculate the minimum free energy secondary structure

formation of sequences(47). The analyses were performed using the -p, -d2, and --noLP

default parameters.

Identification of genes encoding for DNA methyltransferase and their linked motifs. Pod5

files obtained from Nanopore sequencing were merged, using pod5 merge function and

analyzed   for   DNA   methylation   motifs   using   the   Dorado   with   both

dna_r10.4.1_e8.2_400bps_hac@v4.3.0_6mA@v2,

dna_r10.4.1_e8.2_400bps_hac@v4.3.0_5mC_5hmC@v1   models

(https://github.com/nanoporetech/dorado). To identify the location and number of detected

motifs, as well as the genes encoding DNA methyltransferases, the call_methylation and

annotate_rm functions from MicrobeMod were used. These tools were provided with the

corresponding genome assembly ".fasta" and annotation ".gff3" files for each isolate (**Table**

S3-4).

Proteomic analysis

Protein extraction and peptide preparation. After three days on CRLA incubated at 37°C,

bacterial colonies were resuspended in lysing buffer (1M Tris-HCL containing 1% SDS) and

were boiled at 95°C and cool down 10 min at 4°C, three time. Bacterial lysates were then

sonicated in an ultrasonic bath YJ5120-1 (Labtex) for 3 minutes, followed shaking at 1,000

rpm at 95°C for 15 minutes on a Thermomixer Compact (Eppendorf). The mixture was

allowed to cool down to room temperature and 30 µL was mixed with 270 µL of Buffer I (6

M guanidine chloride, 50 mM Tris pH 8.0, 10 mM DTT) and incubated at 30°C for 40 min.

Following addition of 9 µL of 0.5 M iodoacetamide, the sample was incubated at 30°C for 50

min. The reduced and alkylated protein sample was transferred to a 10 kDa cut-off Amicon

Ultra device (Merck) which was placed into a collection tube and centrifuged at $14,000 \times g$

for 30 min. The flow-through from the column was removed and 300 µL of 100 mM

ammonium bicarbonate was added to the column followed by centrifugation at $14,000 \times g$ for

30 min. To digest proteins in the column, 130 µl of 100 mM ammonium bicarbonate

containing 2 µg Trypsin was added. The column was incubated at 37°C for 24 h and then

centrifuged in a new collection tube at $14,000 \times g$ for 30 min. Fifty µL of water with 0.1%

formic acid was then added to the spin column followed by centrifugation at $14,000 \times g$ for

30 min to elute the remaining digested peptides. Enzymes and salts were removed using C_{18}

ZipTips (Merck) following the manufacturer's instructions and desalted peptides were eluted

with 80% acetonitrile/0.1% formic acid into microcentrifuge tubes. Digested peptide samples

were dried down using a vacuum concentrator (Concentrator plus, Eppendorf) and

reconstituted with 50 μL water with 0.1% formic acid prior to LC-MS/MS analysis.

LC-MS/MS analysis. The samples were chromatographically separated and resulting

proteomics data was acquired using an Acquity M-class micro-LC system (Waters, USA)

coupled with ZenoTOF 7600 LC-MS/MS system using the OptiFlow Turbo V ion source.

(AB Sciex). Two μL of sample were loaded onto Micro TRAP C18 column (0.3×10 mm;

Phenomenex) and washed for 10 min at 10 μL/min with 97% solvent A (water with 0.1%

[v/v] formic acid) and 3% solvent B (99.9% [v/v] acetonitrile, 0.1% [v/v] formic acid).

Liquid chromatography was performed at 5 μL/min using a 2.7 μm Peptide C18 160Å; ($0.3 \times$

150 mm; Bioshell) column with the column oven temperature maintained at 40°C. The

gradient started at 3% solvent B which was increased to 5% by 0.5 min, followed by an

increase to 35% solvent B by 38.9 min and then another increase to 80% solvent B by 39

min. The column was washed with 80% solvent B for 3.9 min before equilibrating with 3%

solvent B from 43 to 45 min. Data was acquired using Zeno SWATH DIA, using a 65

variable width SWATH windows. The TOF MS parameters were as follows: precursor mass

range was 400 – 1500 m/z; declustering potential (DP) was set to 80 V and accumulation

time was set at 0.25 s. The TOF MS/MS parameters were as follows: fragmentation mass

range was set to 100 – 1500 m/z; with an accumulation time of 0.025s; fragmentation mode

was set to CID with Zeno pulsing selected. The collision energy for the MS/MS acquisition

was automatically adjusted according to the m/z and charge of the peptide.

Data analyses. Raw label free quantification (LFQ) LC-MS/MS data were converted to

.mzML files, using MS convert, to allow the data search using Fragpipe v22 (48). Protein

identification was conducted against the respective annotated proteomes using the

DIA_SpecLib_Quant workflow, which builds a spectral library with MSFragger-DIA and

performs quantification with DIA-NN. The output "diann-output.pg_matrix.tsv" was filtered

to remove contaminants and retain proteins with at least two non-missing values for at least

one group. Data was normalized using the variance stabilization normalization (49). Proteins

with two or more missing values in one group were categorized as "detected/not detected."

For proteins with a single missing value in one group, missing data were imputed using the

mean of the other replicates in that group. Differential protein expression between groups was

assessed using Welch's t-test, with Benjamini–Hochberg correction applied to control the

false discovery rate (adjusted $p < 0.05$) (50, 51). Analyses were performed on three biological

replicates (Table S5-9).

Proteomic representation. All differences in protein abundances observed among the

Townsville *Bp* isolates in this study were mapped back to the reference strain *Bp* K96243,

widely regarded as the model organism for Bp research and the basis for most virulence

factor and pathway annotations. To identify homologous proteins between the Townsville Bp

isolates and Bp K96243, a megablastp was set up with only one output match for each gene

in query using the following parameters: -evalue 10 -outfmt 6 -num_threads 3 -

max_target_seqs 1 -max_hsps 1 (Table S10; (52). Clustering of orthologous groups (COGs)

for annotated proteins were determined using EggNOG tool v5 (Table S11; (53)). Heatmap

representation, made using 'pheatmap' package in R (54), was based on protein with at least

two isolates showing differences with either three not detected by LC-MS/MS or with

maximum two not differential abundance between CMV.

Multi-omics integration

Principal component analysis. Data were transformed using binary variable to be able to

create a principal component analysis using 'stats' R package function 'prcomp' (55). PCA

was conducted separately for each omics layer using, respectively 16 phenotypes, 19 genomic

top 30 proteomic variables. For integrative analysis, all genomic and phenotypic variables,

along with the top 30 most variable proteomic features, were combined to explore multi-

omics clustering patterns across isolates and colony morphotypes.

Data availability

Genomes from this study are available in GenBank under the Bioproject: PRJNA1295092

(56, 57). Proteomic raw data deposited to the ProteomeXchange Consortium via the PRIDE

partner repository with the dataset identifier PXD065768 (58).

Results

Colony morphotype variation does not translate into distinct phenotypic clusters

Among over 200 clinical Bp isolates from their Townsville collection (32), Etipola and

colleagues (unpublished) classified them into wrinkled, mucoid, or mixed colony

morphotypes when grown on Ashdown selective agar. From the group classified as having

"mixed" morphologies, we selected seven wrinkled clinical isolates and their corresponding

mucoid variants for further genomic, proteomic, and phenotypic investigation (**Table S1**).

Since CMV can occur spontaneously under laboratory conditions – such as removing

antibiotic pressure (19, 23, 27) – we streaked each isolate onto CRLA to confirm its capacity

to generate variants.

Isolates TSV267 and TSV274 produced only one type of colony morphology on CRLA.

Therefore, the five remaining isolates (TSV82, 140, 268, 270 and 287) were selected for

further characterization. To confirm morphology differences, parental and variants were also

grown on Ashdown agar (**Fig. 1**). Interestingly, TSV82, TSV268, and TSV287 smooth

variants were consistently classified as type SI on CRLA and type II (TSV82 and TSV287

smooth) and type I (TSV268 smooth) on Ashdown agar, whereas their respective rough CMV

forms were further differentiated. On CRLA, TSV82 and TSV268 rough variants were

classified as RI, and TSV287 rough as SI. On Ashdown, these rough variants were more

distinct, with TSV82 rough classified as type VIII, TSV268 rough as type II, and TSV287

rough as type VII. In contrast, TSV140 and TSV270 smooth variants were classified as SI on

CRLA and type II and III on Ashdown, respectively, while their rough forms were more

divergent: TSV140 rough was RII on CRLA and type V on Ashdown, and TSV270 rough

was RIII on CRLA and type V on Ashdown (19, 59). With extended incubation, some

morphotypes shifted (**Figure 1B**). TSV270 rough on CRLA developed a lighter inner colony,

while TSV268 smooth lost its wrinkled texture and became fully smooth.

309
Since CMV often reflects underlying adaptive phenotypes, we next assessed virulence-

associated traits, including proteolytic and haemolytic activity, mucoid, and antibiotic

susceptibility (**Table 1**). All ten rough and smooth CMVs were sensitive to meropenem

(MEM), ceftazidime (CAZ). However, while all ten rough CMVs were sensitive to TMP-

SMX, the smooth variant of TSV140 was resistant, whereas the smooth variant of TSV268

remained classified as 'Susceptible, Increased Exposure' but was more sensitive than its rough

counterpart.

As expected, smooth variants were generally more mucoid on YEM agar than their respective

rough parental forms. Additionally, rough variants of TSV82, TSV270, and TSV287

displayed haemolytic activity, in contrast to their smooth counterparts. TSV82 and TSV270

rough forms also produced significantly more protease and siderophore than the smooth

forms, while TSV287 rough only displayed a higher production of siderophore compared to

its smooth variant.

324　Table 1 1 Phenotypic profiling of seven Bp clinical isolates and their respective mucoid
325　variants

Isolate	Colony morphology	Antibiotic Susceptibility test			Mucoidy	Hemolysis	Proteolytic Activity (cm^2)	Siderophore Production (cm^2)
		MEM	CAZ	TMP-SMX				
TSV 82	rough	S	I	I	-	beta	1.81 ± 0.51 *	7.473 ± 1.239 *
	smooth	S	I	I	++	gamma	0.89 ± 0.09	5.097 ± 0.911
TSV 140	rough	S	I	I	+	gamma	0.73 ± 0.10	4.113 ± 0.728
	smooth	S	I	R⍰	++	gamma	0.71 ± 0.00	4.634 ± 1.237
TSV 268	rough	S	I	I	-	gamma	0.00 ± 0.00	2.058 ± 0.548
	smooth	S	I	I⍰	+	gamma	0.00 ± 0.00	2.307 ± 0.288
TSV 270	rough	S	I	I	+	beta	1.80 ± 0.23 **	13.888 ± 2.865 **
	smooth	S	I	I	++	gamma	0.64 ± 0.07	3.753 ± 1.46
TSV 287	rough	S	I	I	-	beta	0.75 ± 0.10	7.707 ± 0.999 *
	smooth	S	I	I	+	gamma	0.92 ± 0.13	5.6 ± 0.916

EUCAST: S = Susceptible; I= Susceptible increased exposure; R= Resistant; Meropenem (MEM): S≤2; R>2; Ceftazidime (CAZ; disk

10µg): S≥50mm; R<18mm; Trimethoprim-Sulphamethadoxone (TMP-SMX; disk 1.25-23.75µg): S≥50mm; R<17mm;

"⍰" represent a TMP-SMX susceptibility statistically different between rough and smooth CMV of TSV140 and TSV268 using a ttest;

Mucoidy: - = non-mucoid; + = mucoid; ++ = more mucoid;

Hemolysis: beta = complete lysis of red blood cell; gamma = no lysis;

Proteolytic activity: average ± stand deviation; comparison TSV 270 rough vs smooth by ttest and TSV 82 rough vs smooth by wilcoxon

test; Siderophore production: average ± stand deviation; comparison by ttest.

Pvalues: "*" represents p value<0.05; "**" represents 0.01<p value<0.05.

To further explore CMV, we assessed the formation of sectors (bacterial growth on the edges

of the main colony), reflecting in variant emergence, after extended incubation on each

medium (**Fig. 1B**). CRLA induced smooth variants in the initial rough forms of TSV268 and

TSV270, while blood agar promoted smooth variants in the rough forms of TSV82, TSV268,

and TSV287 but also in the smooth form of TSV82. Ashdown agar triggered morphotype

changes only in TSV82 and TSV287. Finally, TSV82 smooth variants emerged from its

initial rough form on LB agar supplement with skimmed milk. TSV140 showed no variant

emergence across any media.

Taken together, these results demonstrate that Bp colony morphology varies across some

media but does not reliably correlate with virulence-related phenotypes.

Genomic analyses reveal distinct mutations including in noncoding region with

potential predicted regulatory elements

After confirming that both rough and smooth colony morphotype from each isolate shared the

same sequence type (ST; **Table S1**), we investigated whether differences in colony

morphology were due to underlying genomic variations. We first assembled the genome of

TSV82 rough, TSV 140, TSV268, TSV270 and TSV270 smooth colonies and mapped

Illumina reads from TSV82 smooth, TSV140, TSV268, TSV270 and TSV270 rough on their

respective counterparts assembled genomes using Snippy (**Table 2**).

Table 2: Genomic modifications between rough and smooth CMV

Comparison	Average of illumina coverage	Chromosome	Position	Type	Reference	Alternative	Evidence	Frame Type	Effect	Locustag	Gene	Product	Bp K9623 locustag
TSV82 Smooth vs Rough	Contig 1 50.5X Contig 2 41.4X	1	87085	snp	G	A	A:39 G:3	CDS	synonymous c.3825G>A p.Thr1275Thr	RS_00350	mbaJ	siderophore related no-ribosomal peptide synthase	BPSL1778
		1	87236	snp	T	C	C:42 T:2	CDS	synonymous c.3976T>C p.Leu1326Leu				
		1	87274	snp	C	A	A:40 C:1	CDS	missense c.4014C>A p.His1338Gln				
		1	87454	snp	C	T	T:50 C:1	CDS	synonymous c.4194C>T p.Ile1398Ile				
		1	88446	snp	C	T	T:16 C:1	CDS	missense c.5186C>T p.Ala1729Val				
		1	88575	snp	A	C	C:21 A:2	CDS	missense c.5315A>C p.Asp1772Ala				
		1	88933	snp	C	T	T:16 C:1	CDS	synonymous c.5673C>T p.Arg1891Arg				
		1	89173	snp	G	A	A:24 G:2	CDS	synonymous_ c.5913G>A p.Glu1971Glu				
		1	89512	snp	C	A	A:20 C:1	CDS	synonymous c.6252C>A p.Ala2084Ala				
		1	153859	snp	T	A	A:28 T:2	CDS	missense c.167A>T p.His56Leu	RS_00580		AMP-binding domain protein	BPSL1734
		1	991823	snp	A	T	T:13 A:1	CDS	missense c.217A>T p.Thr73Ser	RS_04110		Oxidoreductase, FAD-binding family protein	BPSL2405
		1	2552774	del	CG	C	C:54 CG:4	intergenic region				Putative non coding region with regulatory element in 5' RS_11330	BPSL0367
		1	2942569	snp	G	A	A:10 G:0	CDS	synonymous c.636G>A p.Ala212Ala	RS_12950		Phosphoribosyl transferase	BPSL0718
		1	2942620	snp	C	G	G:10 C:0	CDS	synonymous c.687C>G p.Ala229Ala				
		1	2942657	snp	G	A	A:15 G:0						
		2	262512	snp	T	C	C:27 T:2	CDS	missense c.628A>G p.Lys210Glu	RS_18625		DNA-binding protein	BPSS0830
		2	486060	snp	G	A	A:28 G:1	CDS	missense c.601G>A p.Glu201Lys	RS_19540		SDR family NAD(P)-dependent oxidoreductase	BPSS1007
		2	537160	complex	GGCCGG	AATCGA	AATCGA:10 GGCCGG:0	CDS	missense c.869_874delCCGGCCinsTCGATT p.ProGlyPro290LeuAspSer	RS_19610		Helix-turn-helix domain protein putative AraC family members	BPSS1020
		2	538080	snp	G	T	T:16 G:1	intergenic region				Putative non coding region with regulatory elementin 5' RS_19610	
		2	547475	snp	C	A	A:16 C:1	CDS	synonymous c.285G>T p.Ala95Ala	RS_19655		Branched-chain amino acid ABC transporter permease	BPSS1030
		2	685277	snp	G	A	A:29 G:1	CDS	missense c.194C>T p.Ala65Val	RS_20305	hyfB	hydrogenase 4 subunit B	BPSS1147
		2	737147	snp	A	G	G:30 A:2						
		2	746638	snp	A	G	G:30 A:0	CDS	synonymous c.783T>C p.Ser261Ser	RS_20510		Luciferase oxidoreductase, group 1 family protein	BPSS1190
		2	988639	snp	A	G	G:55 A:4	CDS	missense c.164A>G p.Gln55Arg	RS_21370		Peptidase, S9A (Prolyl	BPSS1346

										oligopeptidase) family	
	2	1313262	snp	C	T	T:12 C:0	CDS	synonymous c.207C>T p.Ile69Ile	RS_22655		PPC domain-containing protein	BPSS1588
	2	1791476	snp	C	T	T:22 C:1	CDS	synonymous c.2052C>T p.Ala684Ala	RS_24525	rbbA	ribosome-associated ATPase/putative transporter RbbA	BPSS1938
	2	2200644	snp	G	T	T:20 G:1	intergenic region				Putative non coding region with regulatory elementin 5' RS_26200	BPSS2260
	2	3013633	snp	G	C	C:69 G:0	CDS	synonymous c.30C>G p.Ala10Ala	RS_29435	potI	ABC-type spermidine/putrescine transport system, permease component II	BPSS0464
	2	3015293	snp	A	G	G:38 A:2	CDS	synonymous c.492T>C p.Arg164Arg	RS_29445	potG	Spermidine/putrescine import ATP-binding protein	BPSS0466
	2	3015662	snp	A	G	G:37 A:1	CDS	synonymous. 123T>C p.Ala41Ala				
	2	3016583	snp	G	C	C:53 G:4	CDS	synonymous c.513C>G p.Ala171Ala	RS_29450	potF	Putrescine-binding periplasmic protein	BPSS0467
	2	3016945	snp	G	A	A:47 G:1	CDS	missense c.151C>T p.Pro51Ser				
	2	3017125	snp	G	A	A:24 G:0	promotor					
TSV140 Rough vs Smooth	Contig 1 67.8X Contig 2 69.8X Contig 3 90X	2	2170684	del	AGCTTCG	A	A:10 AGCTTCG:0	CDS	missense c.53_58delAGCTTCG p.PheGly103Phe	RS_30990		serine/arginine repetitive matrix 1 CDS	BPSS1650
TSV268 Rough vs Smooth	Contig 1 55.7X Contig 2 56.7X	-										
TSV270_Rough vs Smooth	Contig 1 74X Contig 2 76.3X	-										
TSV287 Rough vs Smooth	Contig 1 64.6X Contig 2 55.4X	1	774690	del	GACGAATCAGT	G	G:32 GACGAATCAGT:0	intergenic region		RS_03535		Putative non coding region with regulatory elementbetween 3' ispA (RS_03530) and 5' trnP tRNA (RS_03540)	BPSL3006-BPSL3007

Firstly, no genomic variation was detected between the rough and smooth forms of TSV268

and TSV270, suggesting that their CMV is likely driven by phase variation. In contrast,

TSV82 smooth differed from its rough counterpart by 33 mutations including four in

intergenic regions, 16 synonymous and eleven missenses. To confirm the accuracy of these

mutations, we mapped contigs from the TSV82 smooth Illumina assembly onto the TSV82

rough reference to ensure they were not artifacts associated with contig ends. However, the

Illumina coverage was 50.5X for contig 1 and 41.4X for contig 2, which could account for

the observed mutations (**Table 2**).

We then assessed the predicted impact of intergenic and missense mutations based on amino

acid substitution significance and genomic context (e.g., within open reading frames or

promoter regions). MbaJ, part of the siderophore biosynthesis cluster (which could explain

the observed difference in siderophore production for this isolate; **Table 1 and Fig. S1A**);

PotF, a periplasmic binding protein of the PotFGHI system involved in putrescine import in

E. coli (**Fig. S1B**); TSV82_RS_18625/BPSS0830, a DNA-binding protein; a NAD(P)-

dependent oxidoreductase (**Fig. S1C**); and TSV82_RS_19610/BPSS0120, a helix-turn-helix

domain-containing protein and putative member of the AraC family of transcriptional

regulators (**Fig. S1D**). To evaluate the potential structural consequences of these mutations,

we used AlphaFold 4 to predict and align the tertiary structures of wild-type and mutated

proteins. Mutations were either located in regions with low prediction confidence or did not

significantly alter predicted secondary or tertiary structures (**Fig. S1**). Further experimental

validation will be necessary to assess their functional impact. Finally, we identified mutations

between smooth and rough colonies within predicted ncRNAs located in the 5′ untranslated

regions (UTRs) of TSV82_RS_19610 (a G-to-T substitution), TSV82_RS_11330 (a deletion

of "CG"), and TSV82_RS_26200 (a G-to-T substitution).

For TSV140, TSV140 rough and smooth variants differed by a deletion of one set of

"GCTTCG" repeat sequence within TSV140_RS_30990 encoding for a putative

serine/arginine repetitive matrix 1 (**Table 2**). This deletion induces a frame shift resulting in a

putative truncated protein. In TSV287, the rough variant lacked one "ACGAATCAGT"

repeat between the 3' UTR of *ispA* (TSV287_RS_03530) and 5' UTR *trnP* tRNA

(TSV287_RS_03535) (**Table 2**).

Proteomic profiles of Bp colony morphotypes reflect adaptive changes in virulence,

quorum sensing, and DNA methylation

In this study, we assessed phenotypes associated with key virulence factors known as

essential for *Bp* pathogenicity, including protease, siderophore and EPS production, along

with susceptibility to antibiotics commonly used for melioidosis treatment. However, *Bp* is

also known to modulate its production of lipopolysaccharide (LPS) and capsular

polysaccharide (CPS) to escape host immunity, resistance to antibiotic and phage attack,

employ multiple secretion systems to invade within macrophages (to replicate) and then

escape (60-70). To gain deeper insights on the molecular pathways affected by CMV, we

performed proteomic analyses comparing smooth and rough colony morphologies when

grown on CRLA plates for 3 days (**Fig. 2; Tables S4-10**). Differential abundance was

observed in proteins involved in a wide range of biological functions including amino acid

and carbohydrate metabolism and transport, cell cycle control and division, cell

wall/membrane/envelope biogenesis, defense mechanisms, energy production and

conversion, inorganic ion transport and metabolism, lipid metabolism, nucleotide

metabolism, post-translational modification and chaperone function, secondary structure,

signal transduction, transcription, and translation. These proteins were consistently either

undetected or significantly lower or higher in abundance in at least three smooth colonies

compared to their rough counterparts. Notably, proteins associated with cell motility and

intracellular trafficking, and secretion were typically undetected or present in lower

abundance in rough colonies relative to smooth colonies.

More specifically, proteins involved in the CAC^{6}AG DNA methylation motif (**Table S2-3**),

antimicrobial resistance (PenA, AmrR), bactobolin biosynthesis (BtaN, L, H, J, Q), haem

uptake (BhuS, T, V), CPS biosynthesis (WbcN), chemotaxis (CheR, CheD), flagellar

biosynthesis (FliN, G, FlgG, H, K FlhA, FliD, H, I, J, O, S, P), Macrophage infectivity

potentiators (Mip2), Malleilactone and malleicyprol biosynthesis (MalB, I, M), Malleobactin

synthesis (FmtA, MbaI, F, A), Pyochelin biosynthesis (FptA, PchA, B, C, D, G, H, I,) T3SS-

3 secretion system (BprQ), T6SS-1 secretion system (TssD-5 aka Hcp1), two-component

signal transduction system (BprR), Type II-OPS biosynthesis (Wzt) were not detected or in

lower abundance in rough colonies relative to smooth colonies. In contrast, proteins involved

in bacterial cellulose synthase (BcsA), EPS cepacian cluster I and II (BceA, C, D, E, F, G, H,

N, M, R, T), quorum sensing (BpsI3/R3 and ScmR), T3SS-3 cascade regulator BspR, T6SS-6

lps (TagM-1, lipoprotein), Type III O–PS polysaccharide (e.g.: RbfbF, G, H) Type IV pilus

biosynthesis (PilS), were not detected or in lower abundance in smooth colony when

compared to rough colonies (**Fig. 2; Tables S4-10**).

Altogether these results indicate that differential regulation of processes such as quorum

sensing and DNA methylation leads to distinct production patterns of numerous proteins

involved in virulence, which in turn are reflected in the observed CMV phenotypes. While

these findings highlight that CMV reflects underlying adaptive changes linked to virulence

modulation, further studies are required to determine whether ScmR, quorum sensing, and

DNA methylation directly regulate CMV and associated modulation of virulence factor

production.

Mutations in non-coding regulatory elements may contribute to differential protein abundance, but are unlikely to drive CMV

Given the identification of mutations in non-coding intergenic regions that may contain regulatory elements such as small (s)RNA (with their predicted structure in **Fig. S2**), we investigated whether these regulatory elements could account for the observed differential protein expression (**Table S11**). Firstly, none of the potential predicted ncRNA had been previously reported (71, 72). Target prediction linked these predicted ncRNAs to genes involved in Type III (*bsaV, bsaK*) and Type IV (*vgrG, bimA*) secretion systems, antimicrobial resistance (*penA*), capsular polysaccharide biosynthesis (*wcbK*), chemotaxis (*cheY*), and secondary metabolite biosynthesis (e.g., *pchD/E* for pyochelin, *malF/M* for malleilactone, *mbaE* for malleobactin, *bceQ* for EPS cepacian, and *bhuU* for Burkholderia haem uptake). However, most of these targets did not exhibit significant differences in protein abundance between rough and smooth colonies in our proteomic analyses. Thus, while the non-coding regions merit further investigation for potential regulatory elements, they do not appear to be the primary drivers of colony morphotype variation.

Deeper -omics profiling shows potential to better characterize *Bp* isolate

Identification based solely on phenotype is unreliable due to the inherent variation in *Burkholderia* colony morphotypes (19, 59), which is further complicated by CMV occurring within individual isolates as an adaptive response. Our combined phenotypic and genomic analyses revealed that these genomic modifications do not explain alone the observed changes (**Fig. 1, Table 2**). This was confirmed by principal component analysis (PCA), where phenotypic and genomic data failed to cluster similar morphotypes or isolates (**Fig. 3A**). In contrast, proteomic profiling allowed more consistent discrimination between rough and smooth CMV variants and even between distinct smooth variants (**Fig. 3A**).

However, proteomics alone lacks biological context without accompanying phenotypic data,

particularly for virulence traits, pathogenicity, and responses to antimicrobial treatment or

stress, which may also be shaped by spontaneous mutations. Therefore, integrating all

datasets provides the most biologically relevant insights (**Fig. 3B**). While smooth isolates are

not further distinct by integrating phenotypic and genomic data, further rearrangements are

observed between rough isolates.

Discussion

Bp has been previously reported to exhibit a heterogeneous population, with both mucoid and

non-mucoid colonies, and to undergo CMV depending on the presence of environmental

pressures. While genomic profiling is often used to study *Bp* adaptation during infection and

phenotypic assays are used to characterize CMV, integrated studies examining the

relationship between genomic changes, phenotypic shifts and proteomic alterations remain

rare (26, 27, 73). In this study, we employed a multi-omics approach to investigate

phenotypic, and proteomic changes associated with CMV, along with underlying genomic

modifications. Our aim was to reveal both conserved and isolate-specific features associated

with the switch between rough and smooth morphotypes – which might transition from an

acute-like to a chronic-adapted state during infection (66).

In several isolates, phase variation appeared to drive the switch between rough and smooth

colony morphologies in the absence of detectable genomic variation. In contrast, other

isolates exhibited limited but notable genomic modifications. In TSV82, SNPs led to

missense mutations in a protein involved in siderophore biosynthesis, along with changes in

both the promoter region and ORF of potF, a gene encoding a potential protective antigen for

melioidosis (74). However, it is important to note that the genomic variation observed in

TSV82 may be partially attributed to lower Illumina read coverage (below 50X).

Additionally, across TSV82 and TSV287 isolates, either a missing repeat sequence or a SNP

was detected within a putative novel mRNA regulatory element (71, 72).

Predicted targets of the putative regulatory element suggest involvement in macrophage

escape, giant multinucleated cell formation - possibly through predicted interacting with

genes involvement in type III and IV secretion systems - and the regulation of several

biosynthetic gene clusters linked to virulence and antimicrobial resistance - including those

responsible for malleilactone, malleicyprol, malleobactin, and pyochelin biosynthesis. While

some key regulators, such as BspR—known to regulate the T3SS-3 and influence T6SS-5—

showed differential abundance in the proteomics analysis, other secretion system components

remained unchanged. This suggests that further studies are needed to clarify the regulatory

changes underlying CMV that influence both secretion systems, which are critical for

macrophage invasion and intracellular survival during infection (60-69).

Nonetheless, our data show extensive changes in pathways related to flagella, antimicrobial

molecules, siderophore biosynthesis, CPS, LPS, and OMPs involved in antimicrobial

resistance. These findings support previous studies describing the emergence of SCVs in Bp

through phase variation, as well as the role of genomic modifications in adaptation during

chronic infection (17, 26, 75, 76). The limited number of point mutations observed suggests

that while they may contribute to phenotypic changes, they are unlikely to be the primary

drivers of CMV. If genomic modifications were the main cause, we would expect more

extensive alterations across regulatory genes and isolates.

Instead, the broad phenotypic and molecular changes associated with CMV are more likely

driven by global regulatory systems—such as ScmR, quorum sensing, and epigenetic

mechanisms like DNA methylation. These pathways were affected at the protein level in our

study. This aligns with previous work demonstrating that ScmR regulates virulence factor

production and quorum sensing in *B. thailandensis*, a surrogate model of *Bp*, and that both

quorum sensing and DNA methylation influence phase variation in *B. ambifaria*, a member

of the *Burkholderia cepacia* complex (77-79).

This study aimed to serve as a preliminary investigation, ultimately demonstrating that

integrating of multi-omics data —phenotypic, genomic, and proteomic— provides valuable

insights into the biological impact reflected by *Bp* adaptation through CMV. This approach

allowed us to differentiate isolates by morphotype and to uncover mechanisms underlying

bacterial adaptation. However, limitations remain, particularly in proteomic analyses, where

LC-MS/MS detection thresholds may restrict peptide identification. Future work will involve

expanding this approach to a broader collection of isolates from environmental, animal, and

human sources, spanning acute and chronic infections. We aim to generate a comprehensive

database that integrates multi-layered omics data, including epigenomics, transcriptomics and

metabolomics, with virulence phenotypes (e.g., macrophage invasion and escape), antibiotic

responses, and clinical outcomes. This will improve our ability to discover the various

mechanisms underlying bacterial adaptation – visible through CMV - correlate molecular

features with disease severity and persistence, and to identify stable biomarkers for diagnostic

and therapeutic targets.

Acknowledgment

The authors thank Julien L. Breton-Robin for helping set command lines tools using

singularity. This work was supported by the funding AIMI and UTS Faculty of Science seed

fundings hold by Dr Pauline M.L. Coulon, the strategic research accelerator funding (SRA

2726229) hold by Prof Garry Myers and EL2 Investigator grant from the NHMRC

(APP2033851) hold by Dr Patrick N.A. Harris. Part of this work was carried out during the

award of a Chancellor's Research Fellowship to Daniel G. Mediati. Royal Australasian

College of Physicians Queensland Regional Committee Research Development Grant hold by

Ian Gassiep allowed the collection and storage of *Bp* isolates. Experimental work was done at

UQCCR which would not have been possible without the ISME scholar funding awarded to

Dr Pauline M.L. Coulon.

P.M.L.C: Concept and visualisation, Funding, Experimental work; K.R: Experimental work

and reviewing D.G.M: Experimental work: data analyses and predictions, and reviewing;

M.E.P and J.T: Experimental work: DNA libraries and sequencing and reviewing; A.L, E.R

and S.R: Experimental work: sample preparation and LCMS runs for the proteomic

experiment, and reviewing; I.G & P.N.A.H: Reviewing and Funding; G.S.A.M: Reviewing

and Funding.

References

1. Meumann EM, Limmathurotsakul D, Dunachie SJ, Wiersinga WJ, Currie BJ. 2024. *Burkholderia pseudomallei* and melioidosis. Nat Rev Microbiol 22:155-169.

2. Kunakom S, Eustaquio AS. 2019. *Burkholderia* as a Source of Natural Products. J Nat Prod 82:2018-2037.

3. Torbeck L, Raccasi D, Guilfoyle DE, Friedman RL, Hussong D. 2011. *Burkholderia cepacia*: This Decision Is Overdue. PDA J Pharm Sci Technol 65:535-43.

4. Cundell T. 2019. Excluding *Burkholderia cepacia* complex from aqueous, non-sterile drug products. American Pharmaceutical Review 22.

5. Tavares M, Kozak M, Balola A, Coutinho CP, Godinho CP, Hassan AA, Cooper VS, Sá-Correia I. 2020. Adaptation and Survival of Burkholderia cepacia and B. contaminans During Long-Term Incubation in Saline Solutions Containing Benzalkonium Chloride. Frontiers in Bioengineering and Biotechnology 8:630.

6. Gee JE, Bower WA, Kunkel A, Petras J, Gettings J, Bye M, Firestone M, Elrod MG, Liu L, Blaney DD, Zaldivar A, Raybern C, Ahmed FS, Honza H, Stonecipher S, O'Sullivan BJ, Lynfield R, Hunter M, Brennan S, Pavlick J, Gabel J, Drenzek C, Geller R, Lee C, Ritter JM, Zaki SR, Gulvik CA, Wilson WW, Beshearse E, Currie BJ, Webb JR, Weiner ZP, Negrón ME, Hoffmaster AR. 2022. Multistate Outbreak of Melioidosis Associated with Imported Aromatherapy Spray. New England Journal of Medicine 386:861-868.

7. McRobb E, Sarovich DS, Price EP, Kaestli M, Mayo M, Keim P, Currie BJ. 2015. Tracing melioidosis back to the source: using whole-genome sequencing to investigate an outbreak originating from a contaminated domestic water supply. J Clin Microbiol 53:1144-8.

8. Sprague LD, Neubauer H. 2004. Melioidosis in Animals: A Review on Epizootiology, Diagnosis and Clinical Presentation. Journal of Veterinary Medicine, Series B 51:305-320.

9. Petras JK, Elrod MG, Ty MC, Dawson P, O'Laughlin K, Gee JE, Hanson J, Boutwell C, Ainsworth G, Beesley CA, Saile E, Tiller R, Gulvik CA, Ware D, Sokol T, Balsamo G, Taylor K, Salzer JS, Bower WA, Weiner ZP, Negron ME, Hoffmaster AR, Byers P. 2023. Locally Acquired Melioidosis Linked to Environment - Mississippi, 2020-2023. N Engl J Med 389:2355-2362.

10. Limmathurotsakul D, Golding N, Dance DA, Messina JP, Pigott DM, Moyes CL, Rolim DB, Bertherat E, Day NP, Peacock SJ, Hay SI. 2016. Predicted global distribution of *Burkholderia pseudomallei* and burden of melioidosis. Nat Microbiol 1.

11. Chantratita N, Phunpang R, Yarasai A, Dulsuk A, Yimthin T, Onofrey LA, Coston TD, Thiansukhon E, Chaisuksant S, Tanwisaid K, Chuananont S, Morakot C, Sangsa N, Chayangsu S, Silakun W, Buasi N, Chetchotisakd P, Day NPJ, Lertmemongkolchai G, West TE. 2023. Characteristics and one year outcomes of melioidosis patients in Northeastern Thailand: a prospective, multicenter cohort study. The Lancet Regional Health - Southeast Asia 9.

12. Currie BJ, Mayo M, Ward LM, Kaestli M, Meumann EM, Webb JR, Woerle C, Baird RW, Price RN, Marshall CS, Ralph AP, Spencer E, Davies J, Huffam SE, Janson S, Lynar S, Markey P, Krause VL, Anstey NM. 2021. The Darwin Prospective Melioidosis Study: a 30-year prospective, observational investigation. Lancet Infect Dis 21:1737-1746.

13.   Currie BJ, Fisher DA, Howard DM, Burrow JN. 2000. Neurological melioidosis. Acta
Trop 74:145-51.

14.   Chadwick DR, Ang B, Sitoh YY, Lee CC. 2002. Cerebral melioidosis in Singapore: a
review of five cases. Trans R Soc Trop Med Hyg 96:72-6.

15.   Schwartzman G, Reddy SA, Berg SH, Currie BJ, Saavedra AP. 2023. Cutaneous
melioidosis: An updated review and primer for the dermatologist. J Am Acad
Dermatol 89:1201-1208.

16.   Currie BJ. 2015. Melioidosis: evolving concepts in epidemiology, pathogenesis, and
treatment. Semin Respir Crit Care Med 36:111-25.

17.   Price EP, Viberg LT, Kidd TJ, Bell SC, Currie BJ, Sarovich DS. 2018.
Transcriptomic analysis of longitudinal Burkholderia pseudomallei infecting the
cystic fibrosis lung. Microbial Genomics 4:465 14.

18.   Currie BJ, Jacups SP. 2003. Intensity of rainfall and severity of melioidosis, Australia.
Emerg Infect Dis 9:1538-42.

19.   Chantratita N, Wuthiekanun V, Boonbumrung K, Tiyawisutsri R, Vesaratchavest M,
Limmathurotsakul D, Chierakul W, Wongratanacheewin S, Pukritiyakamee S, White
NJ, Day NPJ, Peacock SJ. 2007. Biological Relevance of Colony Morphology and
Phenotypic Switching by Burkholderia pseudomallei. Journal of Bacteriology
189:807--817.

20.   Coulon PML, Groleau M-C, Hachani A, Padula MP, Stinear TP, Déziel E. 2024.
Quorum sensing and DNA methylation play active roles in clinical
Burkholderia phase variation. bioRxiv
doi:10.1101/2024.09.30.615881:2024.09.30.615881.

21.   Vial L, Groleau M-C, Lamarche MG, Filion G, Castonguay-Vanier J, Dekimpe V,
Daigle F, Charette SJ, Déziel E. 2009. Phase variation has a role in Burkholderia
ambifaria niche adaptation. The ISME Journal 4:49.

22.   Mariappan V, Barathan M, Zulpa AK, Vadivelu J, Vellasamy KM. 2023. Small
colony variants of *Burkholderia pseudomallei*: alteration of the virulence factors.
Journal of Taibah University for Science 17:2244657.

23.   Häußler S, Rohde M, Steinmetz I. 1999. Highly resistant Burkholderia pseudomallei
small colony variants isolated in vitro and in experimental melioidosis. Medical
Microbiology and Immunology 188:91--97.

24.   Agnoli K, Schwager S, Uehlinger S, Vergunst A, Viteri DF, Nguyen DT, Sokol PA,
Carlier A, Eberl L. 2011. Exposing the third chromosome of Burkholderia cepacia
complex strains as a virulence plasmid. Molecular Microbiology 83:362--378.

25.   Wisniewski-Dyé F, Vial L. 2008. Phase and antigenic variation mediated by genome
modifications. Antonie van Leeuwenhoek 94:493--515.

26.   Coulon PML, Agnoli K, Myers GSA. 2025. Colony morphotype variation in
Burkholderia: implications for success of applications and therapeutics. Journal of
Bacteriology doi:10.1128/jb.00521-24.

27.   Chantratita N, Tandhavanant S, Wikraiphat C, Trunck LA, Rholl DA, Thanwisai A,
Saiprom N, Limmathurotsakul D, Korbsrisate S, Day NPJ, Schweizer HP, Peacock
SJ. 2012. Proteomic analysis of colony morphology variants of Burkholderia
pseudomallei defines a role for the arginine deiminase system in bacterial survival.
Journal of Proteomics 75:1031--1042.

28.   Wikraiphat C, Saiprom N, Tandhavanant S, Heiss C, Azadi P, Wongsuvan G,
Tuanyok A, Holden MTG, Burtnick MN, Brett PJ, Peacock SJ, Chantratita N. 2015.
Colony Morphology Variation of Burkholderia pseudomallei Is Associated with
Antigenic Variation and O-Polysaccharide Modification. Infection and Immunity
83:2127--2138.

29.   Bernhards RC, Cote CK, Amemiya K, Waag DM, Klimko CP, Worsham PL, Welkos
SL. 2017. Characterization of in vitro phenotypes of Burkholderia pseudomallei and
Burkholderia mallei strains potentially associated with persistent infection in mice.
Archives of Microbiology 199:277--301.

30.   Al-Maleki AR, Vellasamy KM, Mariappan V, Venkatraman G, Tee TS, Vadivelu J.
2019. Transcriptome analysis of Burkholderia pseudomallei SCV reveals an
association with virulence, stress resistance and intracellular persistence. Genomics
112:501--512.

31.   Austin CR, Goodyear AW, Bartek IL, Stewart A, Sutherland MD, Silva EB, Zweifel
A, Vitko NP, Tuanyok A, Highnam G, Mittelman D, Keim P, Schweizer HP,
Vázquez-Torres A, Dow SWC, Voskuil MI. 2015. A Burkholderia pseudomallei
Colony Variant Necessary for Gastric Colonization. mBio 6:e02462--14.

32.   Gassiep I, Chatfield MD, Permana B, Burnard D, Bauer MJ, Cuddihy T, Forde BM,
Mayer-Coverdale J, Norton RE, Harris PNA. 2024. The Genomic Epidemiology of
Clinical Burkholderia pseudomallei Isolates in North Queensland, Australia.
Pathogens 13:584.

33.   Chapalain A, Vial L, Laprade N, Dekimpe V, Perreault J, Déziel E. 2013.
Identification of quorum sensing□controlled genes in Burkholderia ambifaria.
MicrobiologyOpen 2:226 242.

34.   Zlosnik JE, Hird TJ, Fraenkel MC, Moreira LM, Henry DA, Speert DP. 2008.
Differential mucoid exopolysaccharide production by members of the *Burkholderia*
cepacia complex. J Clin Microbiol 46:1470-3.

35.   Lynne AM, Haarmann D, Louden BC. 2011. Use of Blue Agar CAS Assay for
Siderophore Detection. Journal of Microbiology \& Biology Education 12:51 53.

36.   Buxton R. 2005. Blood agar plates and hemolysis protocols. American Society for
Microbiology 15:1-9.

37.   Gaio D, Anantanawat K, To J, Liu M, Monahan L, Darling AE. 2022. Hackflex: low-
cost, high-throughput, Illumina Nextera Flex library construction. Microb Genom 8.

38.   De Coster W, D'Hert S, Schultz DT, Cruts M, Van Broeckhoven C. 2018. NanoPack:
visualizing and processing long-read sequencing data. Bioinformatics 34:2666-2669.

39.   Chen S. 2023. Ultrafast one-pass FASTQ data preprocessing, quality control, and
deduplication using fastp. iMeta 2:e107.

40.   Wick RR, Howden BP, Stinear TP. 2025. Autocycler: long-read consensus assembly
for bacterial genomes. bioRxiv doi:10.1101/2025.05.12.653612:2025.05.12.653612.

41.   Bouras G, Judd LM, Edwards RA, Vreugde S, Stinear TP, Wick RR. 2024. How low
can you go? Short-read polishing of Oxford Nanopore bacterial genome assemblies.
Microbial Genomics 10.

42.   Zimin AV, Salzberg SL. 2020. The genome polishing tool POLCA makes fast and
accurate corrections in genome assemblies. PLOS Computational Biology
16:e1007981.

43.   Schwengers O, Jelonek L, Dieckmann MA, Beyvers S, Blom J, Goesmann A. 2021.
Bakta: rapid and standardized annotation of bacterial genomes via alignment-free
sequence identification. Microb Genom 7.

44.   Wick RR, Judd LM, Gorrie CL, Holt KE. 2017. Unicycler: Resolving bacterial
genome assemblies from short and long sequencing reads. PLoS Comput Biol
13:e1005595.

45.   Seemann T. 2015. Snippy: rapid haploid variant calling and core SNP phylogeny.
GitHub Available at: github com/tseemann/snippy.

46. Wright PR, Georg J, Mann M, Sorescu DA, Richter AS, Lott S, Kleinkauf R, Hess
WR, Backofen R. 2014. CopraRNA and IntaRNA: predicting small RNA targets,
networks and interaction domains. Nucleic Acids Res 42:W119-23.
47. Mathews DH, Disney MD, Childs JL, Schroeder SJ, Zuker M, Turner DH. 2004.
Incorporating chemical modification constraints into a dynamic programming
algorithm for prediction of RNA secondary structure. Proc Natl Acad Sci U S A
101:7287-92.
48. Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI. 2017.
MSFragger: ultrafast and comprehensive peptide identification in mass spectrometry-
based proteomics. Nature Methods 14:513-520.
49. Huber W, von Heydebreck A, Sultmann H, Poustka A, Vingron M. 2002. Variance
stabilization applied to microarray data calibration and to the quantification of
differential expression. Bioinformatics 18 Suppl 1:S96-104.
50. Coombs KM, Berard A, Xu W, Krokhin O, Meng X, Cortens JP, Kobasa D, Wilkins
J, Brown EG. 2010. Quantitative proteomic analyses of influenza virus-infected
cultured human lung cells. Journal of Virology 84:10888--10906.
51. Kammers K, Cole RN, Tiengwe C, Ruczinski I. 2015. Detecting significant changes
in protein abundance. EuPA Open Proteomics 7:11--19.
52. Johnson M, Zaretskaya I, Raytselis Y, Merezhuk Y, McGinnis S, Madden TL. 2008.
NCBI BLAST: a better web interface. Nucleic Acids Research 36:W5 W9.
53. Huerta-Cepas J, Szklarczyk D, Heller D, Hernández-Plaza A, Forslund SK, Cook H,
Mende DR, Letunic I, Rattei T, Jensen Lars J, von Mering C, Bork P. 2019. eggNOG
5.0: a hierarchical, functionally and phylogenetically annotated orthology resource
based on 5090 organisms and 2502 viruses. Nucleic Acids Research 47:D309-D314.
54. Raivo K. 2025. pheatmap: Pretty Heatmaps. .
55. Team RDC. 2018. R: A language and environment for statistical computing.
56. Sayers EW, Bolton EE, Brister JR, Canese K, Chan J, Comeau DC, Connor R, Funk
K, Kelly C, Kim S, Madej T, Marchler-Bauer A, Lanczycki C, Lathrop S, Lu Z,
Thibaud-Nissen F, Murphy T, Phan L, Skripchenko Y, Tse T, Wang J, Williams R,
Trawick BW, Pruitt KD, Sherry ST. 2022. Database resources of the national center
for biotechnology information. Nucleic Acids Research 50:D20-D26.
57. Sayers EW, Cavanaugh M, Clark K, Ostell J, Pruitt KD, Karsch-Mizrachi I. 2019.
GenBank. Nucleic Acids Research 47:D94 D99.
58. Perez-Riverol Y, Bai J, Bandla C, Garcia-Seisdedos D, Hewapathirana S,
Kamatchinathan S, Kundu DJ, Prakash A, Frericks-Zipper A, Eisenacher M, Walzer
M, Wang S, Brazma A, Vizcaino JA. 2022. The PRIDE database resources in 2022: a
hub for mass spectrometry-based proteomics evidences. Nucleic Acids Res 50:D543-
D552.
59. Gierok P, Kohler C, Steinmetz I, Lalk M. 2016. Burkholderia pseudomallei Colony
Morphotypes Show a Synchronized Metabolic Pattern after Acute Infection. PLOS
Neglected Tropical Diseases 10:e0004483.
60. Broek CWV, Stevens JM. 2017. Type III Secretion in the Melioidosis Pathogen
Burkholderia pseudomallei. Frontiers in Cellular and Infection Microbiology 7:255.
61. Chen Y, Wong J, Sun GW, Liu Y, Tan G-YG, Gan Y-H. 2011. Regulation of Type VI
Secretion System during Burkholderia pseudomallei Infection. Infection and
Immunity 79:3064--3073.
62. Chieng S, Mohamed R, Nathan S. 2015. Transcriptome analysis of Burkholderia
pseudomallei T6SS identifies Hcp1 as a potential serodiagnostic marker. Microbial
Pathogenesis 79:47--56.

63. Gong L, Cullinane M, Treerat P, Ramm G, Prescott M, Adler B, Boyce JD, Devenish RJ. 2011. The Burkholderia pseudomallei Type III Secretion System and BopA Are Required for Evasion of LC3-Associated Phagocytosis. PLoS ONE 6:e17852.

64. Lennings J, West TE, Schwarz S. 2019. The Burkholderia Type VI Secretion System 5: Composition, Regulation and Role in Virulence. Frontiers in Microbiology 9:3339.

65. Majerczyk C, Schneider E, Greenberg EP. 2016. Quorum sensing control of Type VI secretion factors restricts the proliferation of quorum-sensing mutants. eLife 5:e14712.

66. Sun GW, Chen Y, Liu Y, Tan GYG, Ong C, Tan P, Gan YH. 2010. Identification of a regulatory cascade controlling Type III Secretion System 3 gene expression in Burkholderia pseudomallei. Molecular Microbiology 76:677--689.

67. Toesca IJ, French CT, Miller JF. 2014. The Type VI Secretion System Spike Protein VgrG5 Mediates Membrane Fusion during Intercellular Spread by Pseudomallei Group Burkholderia Species. Infection and Immunity 82:1436--1444.

68. Treerat P, Alwis P, D'Cruze T, Cullinane M, Vadivelu J, Devenish RJ, Prescott M, Adler B, Boyce JD. 2015. The Burkholderia pseudomallei Proteins BapA and BapC Are Secreted TTSS3 Effectors and BapB Levels Modulate Expression of BopE. PLoS ONE 10:e0143916.

69. Warawa J, Woods DE. 2005. Type III secretion system cluster 3 is required for maximal virulence of Burkholderia pseudomallei in a hamster infection model. FEMS Microbiology Letters 242:101--108.

70. Kespichayawattana W, Rattanachetkul S, Wanun T, Utaisincharoen P, Sirisinha S. 2000. Burkholderia pseudomallei induces cell fusion and actin-associated membrane protrusion: a possible mechanism for cell-to-cell spreading. Infection and immunity 68:5377--84.

71. Khoo JS, Chai SF, Mohamed R, Nathan S, Firdaus-Raih M. 2012. Computational discovery and RT-PCR validation of novel *Burkholderia* conserved and *Burkholderia pseudomallei* unique sRNAs. BMC Genomics 13 Suppl 7:S13.

72. Mohd-Padil H, Damiri N, Sulaiman S, Chai SF, Nathan S, Firdaus-Raih M. 2017. Identification of sRNA mediated responses to nutrient depletion in *Burkholderia pseudomallei*. Sci Rep 7:17173.

73. Al-Maleki AR, Mariappan V, Vellasamy KM, Tay ST, Vadivelu J. 2015. Altered Proteome of Burkholderia pseudomallei Colony Variants Induced by Exposure to Human Lung Epithelial Cells. Plos One 10:e0127398.

74. Harland DN, Chu K, Haque A, Nelson M, Walker NJ, Sarkar-Tyson M, Atkins TP, Moore B, Brown KA, Bancroft G, Titball RW, Atkins HS. 2007. Identification of a LolC homologue in *Burkholderia pseudomallei*, a novel protective antigen for melioidosis. Infect Immun 75:4173-80.

75. Price EP, Sarovich DS, Mayo M, Tuanyok A, Drees KP, Kaestli M, Beckstrom-Sternberg SM, Babic-Sternberg JS, Kidd TJ, Bell SC, Keim P, Pearson T, Currie BJ. 2013. Within-host evolution of *Burkholderia pseudomallei* over a twelve-year chronic carriage infection. mBio 4.

76. Vipond J, Kane J, Hatch G, McCorrison J, Nierman WC, Losada L. 2013. Sequence Determination of Burkholderia pseudomallei Strain NCTC 13392 Colony Morphology Variants. Genome Announcements 1:e00925--13.

77. Coulon PML, Groleau M-C, Hachani A, Padula MP, Stinear T, P., Déziel E. 2025. Quorum sensing and DNA methylation play active roles in clinical Burkholderia phase variation. Journal of Bacteriology 207:e00531-24.

78. Le Guillouzer S, Groleau M-C, Mauffrey F, Déziel E. 2020. ScmR, a Global
Regulator of Gene Expression, Quorum Sensing, pH Homeostasis, and Virulence in
Burkholderia thailandensis. Journal of Bacteriology 202.
79. Mao D, Bushin LB, Moon K, Wu Y, Seyedsayamdost MR. 2017. Discovery of scmR
as a global regulator of secondary metabolism and virulence in Burkholderia
thailandensis E264. Proceedings of the National Academy of Sciences of the United
States of America 114:E2920 E2928.
795

Figures

Figures

Figure 1 Characterization of Bp isolates by colony morphology and phenotypes. A) Bp

isolates grown on CRLA or Ashdown agar for 3 days, Roman number represent the type of

colony morphology. B) Variant emergence among Bp isolates when grown on various agar

for 8 days; arrows targets variants emerging from parental isolates. Note: all rough variants

retained a macroscopically wrinkled or dry texture when observed by eye, across media.

Figure 2 Proteomic profile of Bp isolates by comparing smooth to rough colonies.

Heatmap showing protein abundance difference, for each Bp isolates for common proteins

between isolates.

Figure 3 Cluster of Bp isolates, and their variants based on three -omics layers. A)

Individual PCA analysis based on either phenotype (16), genomic (19) or proteomic (1429)

variables. B) Integrative PCA analysis using on all genomic and phenotypes variable and top

30 proteins differentiating isolates.

A

3 days

	0.01% Congo red LB Agar		Ashdown Agar	
	Rough	Smooth	Rough	Smooth
TSV82	RI	SI	VIII	II
TSV140	RII	SI	V	II
TSV268	RI	SI	II	I
TSV270	RIII	SI	V	III
TSV287	RI	SI	VII	II

B

8 days

	0.01% Congo red LB Agar		Ashdown Agar		Columbian Horse Blood agar		1.5% skimmed milk LB agar		Yeast Extract Mannitol Agar	
	Rough	Smooth	Rough	Smooth	Rough	Smooth	Rough	Smooth	Rough	Smooth
TSV82										
TSV140										
TSV268										
TSV270										
TSV287										