Sex differences in crossover interference in house mice

Andrew P Morgan*

*Department of Medicine, University of North Carolina, Chapel Hill, NC 27514, USA

Corresponding author:
Andrew P Morgan (`andrew.parker.morgan@gmail.com`)
Department of Medicine
University of North Carolina
101 Manning Drive #7085
Chapel Hill NC 27599-7085

Keywords: meiosis; crossover interference

Abstract

Meiotic recombination ensures the fidelity of chromosome segregation in most organisms with sexual repro-duction. The distribution of crossovers along chromosomes is governed in part by interference, which prevents multiple crossovers from occurring in close proximity, though not all crossovers are subject to interference. Neither the factors that control strength of interference, nor the extent to which they vary within and between species, are well understood. Here we confirm that that crossover interference is stronger in male than in female meiosis in house mice (*Mus musculus*), provide the first estimate of the proportion of non-interfering crossovers in female mice, and show that this proportion is lower than in males. Interference is stronger on shorter chro-mosomes in both sexes, but the frequency of interference escape is similar across the range of chromosome size. Together with evidence that interference varies across strains and subspecies, our results provide a foundation for studying the evolution and sexual dimorphism in this important feature of meiosis in mice.

Introduction

Meiotic recombination is essential for faithful chromosome segregation in most sexually-reproducing organisms (Hassold and Hunt 2001). Both the number and the spatial distribution of crossovers are tightly regulated, but vary within and between species. At fine scale, the position of crossovers is determined in part by the location of double-strand breaks, which in turn are associated with specific sequence motifs and/or epigenetic marks which differ by taxa (reviewed in Paigen and Petkov (2010)). At the chromosome scale, crossovers tend to be spaced more evenly than expected by chance. This observation dates back to the first linkage maps inferred from *Drosophila* (Sturtevant 1913, 1915) and is known as crossover interference.

The mechanisms and evolutionary significance of crossover interference remain incompletely understood (Hunter 2015). Muller (1916) first speculated that even spacing of crossovers might promote orderly chromosome segregation. There is now ample evidence that aberrant placement of crossovers with respect to the centromere, telomere or to each other contributes to nondisjunction (Hassold *et al.* 1991; Koehler *et al.* 1996; Hassold and Hunt 2001). Others (Kleckner *et al.* 2004; Wang *et al.* 2015) have proposed that interference arises as a consequence of processes that ensure at least one crossover per bivalent (Jones 1984) and maintain the total number of crossovers per meiosis within some optimum range (Martini *et al.* 2006; Cole *et al.* 2012). Further, a subset of crossovers are also processed via a distinct pathway that is not subject to interference (Santos *et al.* 2003; Higgins *et al.* 2004; Guillon *et al.* 2005; Holloway *et al.* 2008). Like other features of the recombination landscape (Morelli and Cohen 2005; Sardell and Kirkpatrick 2020)), both the magnitude of interference and the frequency of interference escape vary between sexes in several mammal species (Otto and Payseur 2019). Comparative studies are hindered by the relative paucity of empirical estimates of interference parameters in both sexes. Recombination tends to be much easier to study in the male than in the female germline – spermatocytes are usually more readily obtained than oocytes, and breeding many progeny from one male is often less resource-intensive than breeding the same number of progeny from one female. Our knowledge of meiosis in mammals is thus biased towards the male germline.

Here we use a large panel of 8-way intercross pedigrees from the Collaborative Cross (CC) (Churchill *et al.* 2004) to provide sex-specific estimates of the strength of crossover interference and the proportion of non-interfering crossovers in both sexes of house mice (*Mus musculus*). The CC is an excellent vehicle for studying recombination (Liu *et al.* 2014) because it comprises hundreds of informative meioses through both sexes in a randomized genetic background with balanced contributions from 8 founder strains representing all the major subspecies of mice (*M. m. domesticus*, *M. m. musculus* and *M. m. castaneus*). We confirm that interference is stronger in males than females and show for the first time that the proportion of non-interfering crossovers is also greater in males than females. Longer chromosomes show stronger interference than shorter chromosomes in both sexes, but no difference in proportion of non-interfering crossovers.

Figure 1: **Estimating crossover interference genome-wide.** (**A**) Pedigree of a representative intercross family. Founder haplotypes are color-coded, and each family is uniquely defined by the order of founder strains in the first generation. Crossovers identified in the G_2:F_1 offspring can be assigned to exactly one meiosis. (**B**) Sex-specific genetic maps, smoothed over 1 Mb windows. (**C,D**) Posterior distribution of global sex-specific estimates of ν and p. Solid dots show posterior mean; bars show $25\% - 75\%$ile (thick) and $2.5\% - 97.5\%$ (thin) quantile intervals.

Results and discussion

We obtained fully phased autosomal haplotypes from one male and one female offspring from each of 237 intercross families with the structure shown in **Figure 1A**. Briefly, the 8 founder strains (G_0 generation, not shown) were randomly intercrossed to create the G_1 hybrids, which were again intercrossed to create the G_2 hybrids. The offspring of the G_2 pairs are denoted the G_2:F_1 and their genomes are a readout of the meioses in the G_1 and G_2 animals. Pedigree constraints allow each haplotype junction to be assigned to a crossover event in exactly one of 8 meioses, so the 474 offspring provide at least partial information for the outcome of 1 896 meioses (see Liu *et al.* (2014)). (Crossovers in the G_1 meioses are only observed if they are transmitted to the G_2:F_1.) We limit our attention to the 948 fully observed meioses in the G_2 generation, which resulted in a total of 12 191 crossovers (6 537 in female and 5 654 in male meioses). Of 18 012 transmitted chromosomes, 7 457 (41.4%) were non-recombinant, 8 950 (49.7%) had exactly one crossover and 1 605 (8.9%) had two or more crossovers. An average of 13.8 crossovers were transmitted per meiosis in females and 11.9 in males. On chromosomes with more than crossover, mean distance between adjacent crossovers was 44.8 cM (SD 14.2 cM) in females and 44.4 cM (SD 13.0 cM) in maless, on the sex-specific genetic maps estimated from this population (**Figure 1B**).

Sex-specific interference parameters were estimated under the "gammma-sprinkling" model of crossover

interference (Housworth and Stahl 2003) using an extension of a Bayesian procedure we have described previously (Morgan and Payseur 2024) (**Figure 1C,D**). Interference was 1.6 times stronger in males ($\nu_{\mathrm{male}} = 18.6$ [95% HPDI $16.3 - 21.1$]) than in females ($\nu_{\mathrm{female}} = 11.2$ [95% HPDI $10.2 - 12.3$]). The proportion of crossovers that escape interference was also 2.9 times greater in males ($p_{\mathrm{male}} = 0.027$ [95% HPDI $0.019 - 0.036$]) than in females ($p_{\mathrm{female}} = 0.0094$ [95% HPDI $0.0054 - 0.016$]).

Many prior studies have shown that crossover interference is heterogeneous across chromosomes. We extended our model to allow chromosome-specific values of ν and p but were not able to obtain stable estimates owing to limited sample size. Results from a reduced model without interference escape were more stable and showed strong negative correlation between ν and chromosome length in both sexes (Spearman's $\rho = -0.39$, $p = 0.017$; **Figure S1**). We reasoned that we could at least partially capture this signal by aggregating over chromosomes of similar size, and therefore divided the them into three groups, each comprising about one-third of the cumulative genetic map. Results are shown in **Figure 2**. Interference was clearly stronger on shorter relative to longer chromosomes – 1.5 times (95% HPDI $1.1 - 1.9$) stronger in females and 2.4 times (95% HPDI $1.6 - 3.7$) stronger in males. The proportion of non-interfering crossovers was similar across the range of chromosome size (just 0.96 times greater in females and 1.0 times greater in males, not significantly different from unity). Sex differences in interference parameters were preserved across the range of chromosome size, though the difference in strength of interference between short and long chromosomes was greater in males. Considering female meioses only, interference on the X chromosome was similar in magnitude to that on autosomes of similar size (**Figure S2**).

Previous studies in mice based on linkage mapping of chromosome 1 (Petkov *et al.* 2007) and cytological analyses (de Boer *et al.* 2006; Peterson and Payseur 2021) have shown stronger interference in males than females. We confirm those findings. Our estimate of $\nu_{\mathrm{female}} = 11.2$ was nearly identical to the only previous estimate from linkage data (11.3) in (*Mus musculus* $\times$ *Mus spretus*) F_1 females (Broman *et al.* 2002). As with cattle (Wang *et al.* 2016), pigs (Brekke *et al.* 2025), domestic dogs (Campbell *et al.* 2016) and humans (Campbell *et al.* 2015), we observed an inverse relationship between (genetic) chromosome size and chromosome-level interference in both sexes. This results in a negative correlation between the genome-wide strength of interference and the average genetic length of a chromosome arm across species (Spearman's $\rho = -0.66$, $p = 0.022$; **Figure S3**), though there is no such relationship with the proportion of non-interfering crossovers (Spearman's $\rho = 0.28$, $p = 0.38$). The proportion of non-interfering crossovers in mice was about half as large as in these other species (**Table 1**). At least in this small group of species, there is no consistent pattern as to whether interference is stronger in males or females; in the sex with the longer or shorter genetic map; or in the sex with more or fewer non-interfering crossovers.

Why does crossover interference differ between sexes? One possibility is that the physical process(es) that

species	sex	cM	N	FN	ν	p	reference
mouse	M	1221	19	19	18.60	0.0268	this study
	F	1355			11.20	0.009 37	
dog	M	1816	38	38	14.05	0.055	Campbell *et al.* (2016)
	F	2162			30.64	0.035	
cow (Holstein)	M	2519	29	58	8.70	0.0834	Wang *et al.* (2016)
	F	2372			9.96	0.1002	
cow (Jersey)	M	2369	29	58	6.70	0.0220	Wang *et al.* (2016)
	F	2229			7.55	0.0289	
pig	M	1915	18	32	8.54	0.049	Brekke *et al.* (2025)
	F	2664			6.45	0.043	
human	M	2708	23	46	8.93	0.067	Campbell *et al.* (2015)
	F	4355			7.19	0.078	

Table 1: Sex-specific interference parameters for available mammal species. cM, length of the autosomal recombination map. N, haploid chromosome number. FN, number of haploid chromosome arms ("fundamental number", (Matthey 1945)).

constitute interference, whatever they are, scale with the length of the synaptonemal complex (SC). The SC is longer in females in humans (Tease and Hultén 2004) and mice (de Boer *et al.* 2006), and crossover counts are positively correlated with SC length within and across individuals in many species (Lynn *et al.* 2002; de Boer *et al.* 2006; Dumont and Payseur 2011; Wang *et al.* 2019a,b). Petkov *et al.* (2007) showed that the coefficient of coincidence, a measure of interference between pairs of loci, depends on distance between loci on the SC (in µm) rather than in genomic coordinates (base pairs) in mice. This predicts that interference will be stronger in the sex with shorter SC relative to genome size and shorter genetic map. Humans, mice and pigs fit this pattern; dogs and cows do not. The relationship between SC length and recombination rate is very well-conserved while the direction of sex differences in interference is not. Factors beyond SC length must therefore be at play and it seems likely that some of these are lineage-specific. Since crossovers tend to be suppressed near centromeres (Hunter 2015) but the number of crossovers per meiosis is proportional to the number of chromosome arms (Pardo-Manuel de Villena and Sapienza 2001), lineage-specific features of chromosome organization and karyotype may contribute. So too may differences in life history: longer generation time means longer duration of meiotic arrest, and may favor weaker interference in females for optimal positioning of crossovers for secure attachment to the spindle (Kong *et al.* 2004; Campbell *et al.* 2015). It is tempting to speculate that sex differences in crossover interference are related to the unique susceptibility of the female germline to meiotic drive – to the extent that weaker interference alters the probability of a crossover between the centromere and a driving locus, it could oppose the spread of the driver if it acts at meiosis I, or favor spread if it acts at meiosis II (Brandvain and Coop 2012). The direction and magnitude of effect would depend on the genetic position, mechanism of

action, and fitness effect of the driving allele. Finally, sex differences in interference may arise indirectly from the balance between sex-specific (and possibly sexually-antagonistic) selective pressures on recombination rate (Mank 2009; Dumont 2017) and mechanisms that ensure crossover homeostasis.

What explains the divergent relationship between ν and p on chromosome size? The biophysical constraints on the maximum number of crossovers and minimum spacing of crossovers compatible with successful chromosome segregation, may be such that chromosomes below some threshold size can accommodate a one and only one crossover. That corresponds to complete interference ($\nu \to \infty$), and indeed, small chromosomes tend to have larger values of ν across species. If the non-interfering pathway is not be subject to the same constraints, we predict that p would be uncorrelated with chromosome size.

Some caveats do apply to our results. First, by using haplotypes transmitted in live offspring as the substrate for analysis, we limit our study to viable gametes only. To the extent that interference is systematically different in gametes which are eliminated by meiotic checkpoints or cannot produce a viable zygote, our findings may be biased. Second, some caution is required when extending conclusions about interference and interference escape as statistical phenomena to the underlying molecular processes. In particular, though we infer the proportion of crossovers that escape interference in aggregate (p), we have no way of designating specific crossovers as non-interfering. The gamma-sprinkling model generally has better fit to empirical data than simpler models (Otto and Payseur 2019), but the relationship between p and the proportion of crossovers molecularly defined as non-interfering still needs to be validated.

Materials and methods

This work re-analyzes published crossover data from the mouse Collaborative Cross (CC) project (Churchill *et al.* 2004). Briefly, the CC is a panel of recombinant inbred mouse lines (RILs), each of which was initiated with an 8-way intercross followed by inbreeding. One male and one female of the last outbreeding generation in a subset of 237 lines were genotyped using a high-density SNP array and used to characterize the genome of the incipient RILs. Mouse breeding, DNA extraction, genotyping, haplotype inference, and construction of the recombination map are described in detail in Liu *et al.* (2014). Coordinates of haplotype segments were defined on the sex-specific genetic map estimated from this cross.

Crossover interference was analyzed under the "gamma-sprinkling" (or Housworth-Stahl) model. Some proportion $p \in (0,1)$ of crossovers are modeled as non-interfering, while the remaining $1 - p$ are subject to interference under the gamma model (Broman and Weber 2000) in which the strength of interference is represented by the single unitless parameter $\nu > 0$. (Note that our parameterization assumes that p is strictly positive, though it can be arbitrarily small. This differs from some other implementations.) We allow both ν and p to vary across groups of meioses (indexed by $i = 1, \ldots m$) and groups of chromosomes (indexed by

Figure 2: **Interchromosomal variation in crossover interference.** (**A**) Strength of interference estimated for chromosomes binned by size ("long", chromosomes 1-5; "medium", chromosomes 6-12; "short", chromosomes 13-19). (**B,C**) Proportion of non-interfering crossovers. (**C**) Sex contrasts for ν and p, expressed as ratios. (**D,E**) Sex-specific contrasts between ν and p for "short" and "long" chromosomes.

$j = 1, \dots n$), using an extension of the hiearchical Bayesian approach described in (Morgan and Payseur 2024).

Let group-specific parameters ν_{ij}, p_{ij} be defined as follows:

$$\log(\nu_{ij}) = \beta_0 + \beta_i + \beta_j \tag{1}$$

$$\mathrm{logit}(p_{ij}) = \alpha_0 + \alpha_i + \alpha_j \tag{2}$$

where β_0 is a global intercept term (to apply some shrinkage) and the β_i and β_j are subject- and chromosome-specific effects. The prior distribution for $\beta_0 \sim \mathcal{N}\left(0, \sigma_\nu^2\right)$ while $\boldsymbol{\beta} \sim \mathrm{MVN}\left(\mathbf{0}, \sigma_\nu^2 \mathbf{I}_m\right)$. That is, group-specific β_i are assumed uncorrelated. The α terms are defined similarly. The likelihood of a set of transmitted chromosome segments, given the parameter vector $\boldsymbol{\theta} = (\boldsymbol{\beta}, \boldsymbol{\alpha})$, was calculated with the C routine R_stahl_loglik from the R package xoi v0.72 (Broman 2023). Inference was done by a simple Metropolis-coupled Markov chain Monte Carlo (MCMC) sampler, with proposal distribution tuned to achieve a rejection rate of around 0.4. The sampler was run for $25\,000$ iterations, of which the first $5\,000$ were discarded as burn-in. Quantities of interest were calculated directly from the posterior distributions, with summary statistics calculated using R package coda v0.19-4.1.

Data availability

Processed data underlying key analyses in this manuscript will be available on Figshare:

`doi:10.6084/m9.figshare.29876591`

Analysis code is available on Github:

`https://github.com/andrewparkermorgan/mouse_crossover_interference_2`

Acknowledgements

We thank Bret Payseur for helpful discussions.

Funding

Support was provided by the following grants from the National Institutes of Health: P50 GM076468 (Gary A Churchill) and F30 MH103925 (Andrew Morgan).

Conflicts of interest

None of the authors have any competing financial interests in the work described in this manuscript.

References

Brandvain, Y. and G. Coop, 2012 Scrambling eggs: meiotic drive and the evolution of female recombination rates. Genetics **190**: 709–723.

Brekke, C., A. B. Gjuvsland, P. Berg, and S. E. Johnston, 2025 Independent genetic basis of meiotic crossover positioning and interference in domestic pigs. Sci. Rep. **15**: 9260.

Broman, K. W., 2023 *xoi: Tools for Analyzing Crossover Interference*.

Broman, K. W., L. B. Rowe, G. A. Churchill, and K. Paigen, 2002 Crossover interference in the mouse. Genetics **160**: 1123–1131.

Broman, K. W. and J. L. Weber, 2000 Characterization of human crossover interference. Am. J. Hum. Genet. **66**: 1911–1926.

Campbell, C. L., C. Bhérer, B. E. Morrow, A. R. Boyko, and A. Auton, 2016 A pedigree-based map of recombination in the domestic dog genome. G3 .

Campbell, C. L., N. A. Furlotte, N. Eriksson, D. Hinds, and A. Auton, 2015 Escape from crossover interference increases with maternal age. Nat. Commun. **6**: 6260.

Churchill, G. A., D. C. Airey, H. Allayee, J. M. Angel, A. D. Attie, *et al.*, 2004 The collaborative cross, a community resource for the genetic analysis of complex traits. Nat. Genet. **36**: 1133–1137.

Cole, F., L. Kauppi, J. Lange, I. Roig, R. Wang, *et al.*, 2012 Homeostatic control of recombination is implemented progressively in mouse meiosis. Nat. Cell Biol. **14**: 424–430.

de Boer, E., P. Stam, A. J. J. Dietrich, A. Pastink, and C. Heyting, 2006 Two levels of interference in mouse meiotic recombination. Proc. Natl. Acad. Sci. U. S. A. **103**: 9607–9612.

Dumont, B. L., 2017 X-chromosome control of genome-scale recombination rates in house mice. Genetics **205**: 1649–1656.

Dumont, B. L. and B. A. Payseur, 2011 Evolution of the genomic recombination rate in murid rodents. Genetics **187**: 643–657.

Guillon, H., F. Baudat, C. Grey, R. M. Liskay, and B. de Massy, 2005 Crossover and noncrossover pathways in mouse meiosis. Mol. Cell **20**: 563–573.

Hassold, T. and P. Hunt, 2001 To err (meiotically) is human: the genesis of human aneuploidy. Nat. Rev. Genet. **2**: 280–291.

Hassold, T. J., S. L. Sherman, D. Pettay, D. C. Page, and P. A. Jacobs, 1991 XY chromosome nondisjunction in man is associated with diminished recombination in the pseudoautosomal region. Am. J. Hum. Genet. **49**: 253–260.

Higgins, J. D., S. J. Armstrong, F. C. H. Franklin, and G. H. Jones, 2004 The arabidopsis MutS homolog AtMSH4 functions at an early step in recombination: evidence for two classes of recombination in arabidopsis. Genes Dev. **18**: 2557–2570.

Holloway, J. K., J. Booth, W. Edelmann, C. H. McGowan, and P. E. Cohen, 2008 MUS81 generates a subset of MLH1-MLH3-independent crossovers in mammalian meiosis. PLoS Genet. **4**: e1000186.

Housworth, E. A. and F. W. Stahl, 2003 Crossover interference in humans. Am. J. Hum. Genet. **73**: 188–197.

Hunter, N., 2015 Meiotic recombination: The essence of heredity. Cold Spring Harb. Perspect. Biol. **7**: a016618.

Jones, G. H., 1984 The control of chiasma distribution. Symp. Soc. Exp. Biol. **38**: 293–320.

Kleckner, N., D. Zickler, G. H. Jones, J. Dekker, R. Padmore, *et al.*, 2004 A mechanical basis for chromosome function. Proc. Natl. Acad. Sci. U. S. A. **101**: 12592–12597.

Koehler, K. E., R. S. Hawley, S. Sherman, and T. Hassold, 1996 Recombination and nondisjunction in humans and flies. Hum. Mol. Genet. **5 Spec No**: 1495–1504.

Kong, A., J. Barnard, D. F. Gudbjartsson, G. Thorleifsson, G. Jonsdottir, *et al.*, 2004 Recombination rate and reproductive success in humans. Nat. Genet. **36**: 1203–1206.

Liu, E. Y., A. P. Morgan, E. J. Chesler, W. Wang, G. A. Churchill, *et al.*, 2014 High-resolution sex-specific linkage maps of the mouse reveal polarized distribution of crossovers in male germline. Genetics **197**: 91–106.

Lynn, A., K. E. Koehler, L. Judis, E. R. Chan, J. P. Cherry, *et al.*, 2002 Covariation of synaptonemal complex length and mammalian meiotic exchange rates. Science **296**: 2222–2225.

Mank, J. E., 2009 The evolution of heterochiasmy: the role of sexual selection and sperm competition in determining sex-specific recombination rates in eutherian mammals. Genet. Res. (Camb.) **91**: 355–363.

Martini, E., R. L. Diaz, N. Hunter, and S. Keeney, 2006 Crossover homeostasis in yeast meiosis. Cell **126**: 285–295.

Matthey, R., 1945 L'évolution de la formule chromosomiale chez les vertébrés. Experientia **1**: 78–86.

Morelli, M. A. and P. E. Cohen, 2005 Not all germ cells are created equal: aspects of sexual dimorphism in mammalian meiosis. J Reprod Fertil **130**: 761–781.

Morgan, A. P. and B. A. Payseur, 2024 Genetic background affects the strength of crossover interference in house mice. Genetics **228**.

Muller, H. J., 1916 The mechanism of crossing-over. Am. Nat. **50**: 193–221.

Otto, S. P. and B. A. Payseur, 2019 Crossover interference: Shedding light on the evolution of recombination. Annu. Rev. Genet. **53**: 19–44.

Paigen, K. and P. Petkov, 2010 Mammalian recombination hot spots: properties, control and evolution. Nat. Rev. Genet. **11**: 221–233.

Pardo-Manuel de Villena, F. and C. Sapienza, 2001 Recombination is proportional to the number of chromosome arms in mammals. Mamm. Genome **12**: 318–322.

Peterson, A. L. and B. A. Payseur, 2021 Sex-specific variation in the genome-wide recombination rate. Genetics **217**: 1–11.

Petkov, P. M., K. W. Broman, J. P. Szatkiewicz, and K. Paigen, 2007 Crossover interference underlies sex differences in recombination rates. Trends Genet. **23**: 539–542.

Santos, T. d. L., N. Hunter, C. Lee, B. Larkin, J. Loidl, *et al.*, 2003 The Mus81-Mms4 endonuclease acts independently of double-holliday junction resolution to promote a distinct subset of crossovers during meiosis in budding yeast. Genetics **164**: 81–94.

Sardell, J. M. and M. Kirkpatrick, 2020 Sex differences in the recombination landscape. Am. Nat. **195**: 361–379.

Sturtevant, A. H., 1913 The linear arrangement of six sex-linked factors in drosophila, as shown by their mode of association. J. Exp. Zool. **14**: 43–59.

Sturtevant, A. H., 1915 The behavior of the chromosomes as studied through linkage. Z. Indukt. Abstamm. Vererbungsl. **13**: 234–287.

Tease, C. and M. A. Hultén, 2004 Inter-sex variation in synaptonemal complex lengths largely determine the different recombination rates in male and female germ cells. Cytogenet. Genome Res. **107**: 208–215.

Wang, R. J., B. L. Dumont, P. Jing, and B. A. Payseur, 2019a A first genetic portrait of synaptonemal complex variation. PLoS Genet. **15**: e1008337.

Wang, S., C. Veller, F. Sun, A. Ruiz-Herrera, Y. Shang, *et al.*, 2019b Per-nucleus crossover covariation and implications for evolution. Cell **177**: 326–338.e16.

Wang, S., D. Zickler, N. Kleckner, and L. Zhang, 2015 Meiotic crossover patterns: obligatory crossover, interference and homeostasis in a single process. Cell Cycle **14**: 305–314.

Wang, Z., B. Shen, J. Jiang, J. Li, and L. Ma, 2016 Effect of sex, age and genetics on crossover interference in cattle. Sci. Rep. **6**: 37698.

Supplementary material

File S1. Table of 21 993 crossovers (COs) reported in (Liu *et al.* 2014). Columns are as follows.

column	contents
funnel	family ID (referred to as "funnel" due to shape of pedigree)
order	order of founder strains at the G_0 generation; this is unique for each family
meiosis	code identifying one of 8 meioses in the pedigree, with respect to the G_2:F_1 individual: MGM, maternal grandmother (G_1); MGF, maternal grandfather (G_1); PGM, paternal grandmother (G_1); PGF, paternal grandfather (G_1); Mf, mother of female G_2:F_1 individual (G_2); Pf, father of female G_2:F_1 individual (G_2); Mm, father of male G_2:F_1 individual (G_2); Pm, father of male G_2:F_1 individual (G_2).
generation	pedigree generation in which this CO occurred (G_1 or G_2)
sex	sex of the parent in which this CO occurred (M or F)
chr	chromosome on which this CO occured
start	physical position of proximal (left-most) marker defining this CO (bp)
end	physical position of distal (right-most) marker defining this CO (bp)
from	founder haplotype proximal to CO
to	founder haplotype distal to CO
pos	physical midpoint of CO interval (bp)
cM	genetic position (in cM) of this CO, on the sex-specific map

Additional notes

Physical positions are reported on the mm9/GRCm37 assembly, as this was used during the original analysis, without liftover to the current mm39 assembly. Only genetic positions are actually used in this manuscript. Founder strains are reported as 1-letter codes: A = A/J; B = C57BL/6J; C = 129S1/SvImJ; D = NOD/ShiLtJ; E = NZO/HlLtJ; F = CAST/EiJ; G = PWK/PhJ; H = WSB/EiJ.

Figure S1: **Chromosome-specific interference parameters under the simple gamma model without interference escape.** (**A**) Chromosome-specific values of ν with 95% HPDIs plotted against genetic chromosome length. (**B**) Comparison of male versus female estimates of ν for each chromosome. Dotted reference line passes through the intercept and has slope 1.

Figure S2: **Chromosome-specific interference parameters under the simple gamma model without interference escape, in females only.** Chromosome-specific values of ν with 95% HPDIs plotted against genetic chromosome length, for labelled autosomes (grey) and the X chromosome (black).

Figure S3: **Interference vs length of genetic map.** Sex-specific values of ν (**A**) and p (**B**) plotted against the average genetic length of a chromosome arm in 5 mammal species.