RESCUE: recovery of unattributed expression patterns in spatial transcriptomics

Young Joo Lee[1], Seokjin Yeo[2], Alex W. Schrader[3], Ian M. Traniello[4], Amy Cash Ahmed[5],

Gene E. Robinson[5,6,7], Hee-Sun Han[2,3,5,6*,] Sihai Dave Zhao[1,5*]

[1]Department of Statistics, University of Illinois Urbana-Champaign

[2]Department of Bioengineering, University of Illinois Urbana-Champaign

[3]Department of Chemistry, University of Illinois Urbana-Champaign

[4]Lewis-Sigler Institute for Integrative Genomics, Princeton University

[5]Carl R. Woese Institute of Genomic Biology, University of Illinois Urbana-Champaign

[6]Neuroscience Program, University of Illinois Urbana-Champaign

[7]Department of Entomology, University of Illinois Urbana-Champaign

* Corresponding authors: sdzhao@illinois.edu (S.D.Z), hshan@illinois.edu (H.-S.H)

ABSTRACT

Spatial transcriptomics (ST) enables gene expression profiling while preserving the

spatial architecture of intact tissue. Analyzing ST data often proceeds by first extracting

cell-level information, typically through cell segmentation or cell-type deconvolution.

However, a critical oversight has been that a substantial portion of molecular expression

is systematically lost or unannotated by these methods. This lost expression can arise

from diverse and biologically important sources like fragile or underrepresented cell types,

subcellular structures like neurites, and extracellular expression. These omissions can

result in biased analyses and incorrect or incomplete biological interpretations. We

describe a new computational method, RESCUE, that can recover the unattributed spatial

expression patterns missed by existing ST analysis methods and enable robust inference

even when reference is incomplete. We validate RESCUE using MERFISH data from the

honey bee brain and apply it to multiple ST datasets to demonstrate how it can reveal

novel insights into complex tissue biology.

INTRODUCTION

Profiling cellular expression within its spatial context has become possible with the recent

development of spatial transcriptomics (ST) technologies, which measures gene

expression in intact tissue sections[1,2]. Analyzing ST data typically proceeds by extracting

cell-level information from the spatial expression data, as cells are natural units of

analysis. With high-resolution platforms such as MERFISH[3], 10x Visium HD[4], or 10x

Xenium[5], this is usually done using cell segmentation[6–8]. With low-resolution platforms

like 10x Visium[9] and Slide-seq[10,11], where each spatial location can contain multiple cells,

it is common to deconvolve the observed expression into separate contributions from

different cell types, based on reference single-cell/nuclei RNA-seq (sc/snRNA-seq)

data[12–16].

However, a major problem with these existing ST analysis methods is that they

can fail to capture important expression patterns. First, they may omit entire cells or cell

types. Cell segmentation algorithms for high-resolution data have difficulty with cells with

complex morphologies, such as fibroblasts, or cells lacking nuclei, like erythrocytes[17,18].

Deconvolution methods for low-resolution data cannot account for cell types that are

absent from the reference sc/snRNA-seq data because they are too fragile to survive

tissue dissociation[19–21]. Second, existing ST approaches can overlook expression

patterns unique to cellular substructures. Segmentation methods simply drop transcripts

located in substructures that are difficult to segment, such as the axons and dendrites of

neurons. These fragile structures are also typically destroyed during tissue dissociation

and therefore are not represented in reference single-cell data used in deconvolution.

Indeed, it has been estimated that over 40% of the total transcriptome of an adult mouse

brain may be missed by sc/snRNA-seq[22]. Similarly, reference snRNA-seq data, by

definition, ignores cytoplasmic expression. Finally, existing methods do not capture

extracellular tissue expression patterns, which cannot be studied by segmentation or

deconvolution.

These difficult-to-capture expression patterns can be critically important. First, they

can bias analyses that do not take them into account. An example with dramatic

implications is that deconvolution estimates of cell-type proportions can be invalid if the

reference is missing any cell types[23], as we illustrate later in this work. Second, there may

be important biological processes that are unique to the overlooked expression patterns.

For example, important missed cell types include adipocytes[5], which play crucial roles in

metabolic disease and cancer[24], and neutrophils[25], which are core components of the

immune system. Important missed substructural expression includes axonal or dendritic

transcripts that are central to synaptic development and plasticity[26–30] and diseases such

as amyotrophic lateral sclerosis[31,32], as well as cytoplasmic transcripts that regulate key

processes such as metabolism, signaling, and stress responses[33–35] and are involved in

Alzheimer's disease[36,37]. Important missed extracellular expression patterns include

extracellular RNAs that mediate intercellular communication in cancer[38–40].

We describe a new computational method, RESCUE, that can recover the spatial

expression patterns missed by single-cell inference on ST data. Our approach is based

on making a fundamental correction to the class of statistical models currently used for

integrating ST data and reference data, such as sc/snRNA or cell segmentation results.

Briefly, our method partitions observed gene expression into one component explainable

by the reference data, which we refer to as "canonical" expression, and another

component that is not, which we refer to as "unattributed" expression. Our purified

canonical ST expression component can be used to correct existing cell-type

deconvolution methods. Our recovered unattributed ST expression component gives us

the new ability to specifically target biology that is distinct from our reference data, as

discussed above. The unattributed expression recovery step in RESCUE is analogous to

negative selection in experimental biology[41–44], where we identify and remove canonical

expression. We demonstrate that RESCUE accurately recovers unattributed expression,

enables robust deconvolution of ST data despite incomplete reference data, and reveals

biologically meaningful patterns overlooked by existing methods. While we focus here on

ST data, we note that RESCUE can also be directly applied to non-spatial modalities,

such as bulk RNA-seq data.

RESULTS

Overview of RESCUE

RESCUE is predicated on a new model that we propose for integrating ST and reference

single-cell data, and can be applied to diverse ST platforms, including MERFISH and

Visium, across different organisms. Most existing models[12–16] posit that ST expression at

each location is composed of a linear combination of expression patterns present in the

reference, where the coefficients of this linear combination are interpreted as the

proportions of each reference cell type in that location. However, as described in the

introduction, there are numerous sources of ST expression that are left out of this model;

we will refer to these as unattributed expression patterns, because they are not attributed

to canonical expression patterns reflected in the reference.

To address this issue, we propose a corrected model (**Fig. 1**) that allows the ST

profile to have both canonical and unattributed expression components. While it appears

minor, this change has important implications. Under this model, results from the statistics

literatures[45–47] imply that under certain conditions it is possible to simultaneously recover

unattributed expression and estimate the linear combination of reference patterns that

constitute canonical expression. The conditions are that only relatively few genes show

non-zero, or "sparse" expression, in the unattributed component, which is reasonable

given the nature of unattributed expression and the read depth/detection efficiency of ST

technologies (**Methods**).

Leveraging this insight, we developed RESCUE, an algorithm that simultaneously

deconvolves ST data while rescuing spatial unattributed expression. RESCUE solves a

regularized robust multivariate regression problem and can be applied to both low- and

high-resolution ST data. Unlike existing deconvolution methods[12–16], it is robust to

incomplete references, such as snRNA-seq data which lack cytoplasmic expression

profiles. We emphasize that the interpretation of the unattributed component depends on

both the biological context of the tissue and the choice of the reference dataset

(**Methods**). As a result, the unattributed component may comprise multiple biological

sources, including expression from cell types missing in the reference, substructural

expression, and extracellular RNA. RESCUE is designed to recover these signals jointly

rather than to assign them to predefined categories. The rescued unattributed expression

can then be further decomposed using standard unsupervised approaches, such as

clustering and non-negative matrix factorization (NMF), to identify distinct expression

patterns within the unattributed component. In addition, differentially expressed gene

(DEG) and spatially variable gene (SVG) analyses can be applied to characterize the

biological and spatial features of the recovered unattributed signals. Careful consideration

of biological structures and tissue organization is therefore essential when interpreting

the unattributed expression. We illustrate this in a broad range of examples below.

Simulations

To evaluate the performance of RESCUE, we conducted simulation studies on synthetic

data designed to mimic real ST measurements with varying degrees of unattributed

expression. We simulated ST data using real scRNA-seq data from the honey bee brain[48]

by introducing synthetic unattributed expression of a small number of genes (**Methods**).

Unlike other tissues where cell bodies and non-cell-body structures are spatially

intermingled, honey bee brains provide a unique benefit for simulating unattributed

expression due to the existence of the neuropil region, where axons and dendrites are

spatially segregated from cell bodies. Molecular expression in these regions is largely

absent in single-cell sequencing data due to the structural fragility of neurites. During the

tissue dissociation step, a necessary step in single-cell sequencing, neurites are mostly

sheared off and lost. In contrast, ST measurements performed on intact tissue sections

preserve these structures and therefore capture molecular expression within neuropils.

To guide the choice of sparsity levels used in our simulations, we examined

empirical expression patterns in the neuropil region of our in-house MERFISH data. We

found that neuropil regions exhibit highly sparse gene expression, with an average

sparsity level of 0.92 across spatial locations (**Fig. 2a**). Here, sparsity refers to the

structure of the unattributed component (i.e., that only a subset of genes is present at a

given spatial location), rather than to low expression levels (**Methods**). Based on this

empirical distribution, we simulated unattributed expression across a range of sparsity

levels by varying the proportion of zero entries (ranging from 0.8 to 0.95). In addition, we

varied the proportion of unattributed expression (ranging from 0 to 0.8) to cover different

levels of the overall magnitude of unattributed expression. Importantly, these parameter

ranges fall within those observed in real biological samples and are not artificially chosen

to favor RESCUE.

We first compared the deconvolution performance of RESCUE with existing

methods. **Fig. 2b** summarizes the accuracy of cell-type proportion estimates across

methods, measured using the average Lin's concordance correlation coefficient[49] (CCC),

root mean squared error (RMSE), and Jensen-Shannon divergence (JSD) (**Methods**). As

expected, the performance of standard deconvolution methods deteriorates sharply as

the level of unattributed expression increases. In contrast, RESCUE maintains high

accuracy across all conditions, demonstrating robustness to both high unattributed

content and reduced sparsity. These results indicate that failing to explicitly model

unattributed expression can lead to systematic biases in cell-type proportion estimates.

162   We also evaluated the accuracy of unattributed expression recovery (**Fig. 2c**)

using the relative Frobenius error between the true and estimated unattributed expression

matrices (**Methods**). Overall, RESCUE achieves the lowest recovery error even under

high unattributed expression and low sparsity levels, whereas other methods deviate

substantially from the ground truth, especially when unattributed signals become strong.

These results indicate that explicitly modeling unattributed expression enables accurate

recovery of hidden transcriptional signals across a wide range of sparsity and signal-

strength regimes.

170   To assess robustness and generalizability, we expanded the simulation study by

varying noise levels via library size variability across spots and varying the effective grid

resolution via changes in the average total UMI count per spot (**Supplementary Fig. 1**).

As expected, overall performance tends to decrease with increasing library size variability

and increase with higher average total UMIs. Across these regimes, RESCUE remains

among the top-performing methods, demonstrating robustness across a range of noise

levels and grid sizes.

RESCUE accurately recovers the unattributed transcriptome in real tissues

Using a custom MERFISH platform[50,51], we generated single molecule-resolution ST data

from coronal sections of a honey bee brain, targeting 130 genes relevant to aggression

(**Supplementary Table S1**)[52]. Transcripts in axons and dendrites, which are collectively

termed as neurites, have been difficult to profile due to segmentation challenges. Cell

segmentation methods, such as Cellpose[6] and Baysor[7], can only identify the boundaries

of neuronal cell bodies, also called somata. Although individual axons and dendrites can

be stained in 2D cell cultures[53], in native tissues they extend across multiple imaging

planes and therefore cannot be reliably segmented as individual cellular units. As a result,

neurite-associated expression has traditionally been ignored or incorrectly merged into

nearby cytosolic expression.

We first experimentally validated that RESCUE can accurately recover gene

expression in the neuropil region of the honey bee brain (**Fig. 3a**). This validation is

possible because the physical separation of neurites and somata in the brain makes it

simple for existing algorithms to separate somatic from unattributed expression. Cell

soma were segmented using Cellpose[6], revealing that only 73.5% of transcripts were

localized within cells, with the remaining 26.5% comprising the unattributed expression.

Using honey bee brain scRNA-seq data[48] as a reference, we then applied RESCUE and

competing methods to our MERFISH data after dividing the brain into grids of size

$20 \times 20 \, \mu m^2$ (**Methods**). For each grid, we estimated the proportion of canonical

expression relative to total expression and obtained a ground-truth estimate by identifying

transcripts localized within cell bodies via cell segmentation. RESCUE's estimates closely

matched the ground truth (**Fig. 3b**) and were more accurate than those from other

methods, as quantified by Lin's CCC and RMSE (**Methods**). Notably, RESCUE relies

only on reference scRNA-seq data and does not use cell segmentation information.

To further clarify the source of discrepancies among methods, we evaluated

canonical and unattributed expression separately in anatomically well-defined regions.

Specifically, we examined grids corresponding to pure cell bodies and compared the

average canonical expression of each gene against the ground truth derived from

segmentation (**Fig. 3c**). RESCUE closely matched the true somatic expression levels,

whereas competing methods systematically underestimated canonical expression. This

behavior reflects the fact that existing deconvolution approaches are designed to estimate

cell-type proportions rather than expression levels, and under the subtraction-based

framework (**Methods**), expression that is not well explained by the reference profiles is

left as residual rather than attributed to the canonical component. Conversely, in grids

containing no identifiable cell bodies, RESCUE accurately recovered unattributed

expression in neuropil-only regions (**Fig. 3d**), while competing methods were significantly

less accurate. Together, these results demonstrate that RESCUE effectively partitions

the full transcriptome into canonical and unattributed components in real tissue, capturing

valuable expression signals that would otherwise be lost using conventional approaches.

Importantly, transcripts outside segmented somata reflect genuine biological

signals rather than technical artifacts: in our MERFISH sample (**Fig. 3a**), neuropil-

associated transcripts form structured spatial patterns that are clearly segregated from

DAPI-stained nuclei (**Supplementary Fig. 2**). Moreover, the MERFISH protocol includes

gel embedding, expansion, and clearing steps that reduce autofluorescence and

suppress nonspecific signal[54,55]. These signals, which would have been ignored by

standard cell segmentation approaches, highlight the importance of RESCUE in capturing

overlooked transcriptomic information.

Finally, to investigate whether the unattributed expression recovered by RESCUE

reflects structured biological organization rather than unstructured noise, we performed

clustering analysis of the unattributed expression profiles across spatial locations.

Clustering revealed anatomically distinct transcriptional niches within neuropil regions

(**Fig. 3e**), indicating that the recovered unattributed component exhibits coherent spatial

patterns. Specifically, distinct clusters localize to known neuropil structures, including the

mushroom body calyces, antennal lobes, optic lobes, and regions proximal to the central

complex. In contrast, spatial locations corresponding to somatic regions collapsed into a

single cluster, consistent with the fact that canonical expression is explicitly removed, and

only minimal unattributed signal remains in these regions. Examination of representative

genes within the neuropil-associated clusters showed spatially structured expression

patterns that are largely absent from the canonical component (**Fig. 3f**). These findings

further support that the unattributed expression recovered by RESCUE captures

biologically meaningful and spatially organized signals, revealing hidden tissue structure

that is not accessible through conventional analyses.

RESCUE enables accurate deconvolution of ST data with incomplete reference

Here, we illustrate how RESCUE's estimated canonical expression component can be

used to achieve accurate deconvolution of ST data when key cell types are missing from

the reference, a challenge for all existing methods that has been noted in a recent study[23].

We analyzed paired single-cell FFPE-seq (scFFPE-seq) and 10x Visium ST data from

human breast cancer tissue[5]. In this ST dataset, adipocytes are manually annotated in

approximately 10% of spatial spots (**Fig. 4a**). However, adipocytes are absent from the

scFFPE-seq reference, likely due to their high lipid content interfering with fixation and

processing. These cells are prone to floating, rupturing, or sticking to plastic surfaces

during tissue dissociation, which hinders their recovery in single-cell sequencing

protocols[5,17]. As a result, reference-based methods that assume the observed expression

is fully explained by the available reference cell types tend to misattribute adipocyte-

derived transcripts to other cell types.

In adipocyte-rich regions, existing deconvolution methods fail to account for the

missing adipocyte transcriptome and instead misattribute the signal to other reference

cell types, such as stromal and endothelial cells (**Fig. 4b**). This systematic misassignment

likely arises from transcriptional similarity, as shown by the strong correlation between

adipocyte and stromal/endothelia profiles in pseudo-bulk measurements from Visium and

scFFPE-seq (**Fig. 4c**). In contrast, RESCUE explicitly models unattributed expression

and thus avoids forcing unassigned signal into existing reference cell types, which results

in more accurate and biologically plausible estimates. As shown in **Fig. 4d-e**, RESCUE's

estimated proportions closely align with the scFFPE-seq composition, while other

methods exhibit significant distortions. Together, these results demonstrate that RESCUE

improves deconvolution accuracy in settings with incomplete references by preventing

the systematic misassignment of unattributed transcriptomes to incorrect cell types.

To further interpret the biological content of the unattributed expression recovered

by RESCUE, we applied NMF to the unattributed component. This analysis revealed

multiple latent factors capturing structured expression patterns that are not explained by

the canonical component (**Fig. 4f**). Notably, the NMF Factor 1 (Unattributed) captures the

spatial distribution of adipocyte-rich regions, effectively recovering a missing cell type

absent from the reference. Genes with the highest loadings in this factor include well-

established adipocyte markers such as *FABP4*, *CFD*, and *G0S2*[5,56] (**Fig. 4g**). In addition,

274 the NMF Factor 2 (Unattributed) is enriched for genes such as *DCN*, *COL14A1*, and *C1R*,

275 which have been implicated in extracellular matrix (ECM) domains[57–59]. Together, these

276 findings indicate that the unattributed component captured by RESCUE is biologically

277 meaningful, thereby supporting that explicitly modeling it leads to improved deconvolution

278 in the presence of incomplete references.

279  Finally, to assess whether these deconvolution improvements generalize beyond

280 the human breast cancer example, we benchmarked RESCUE using additional datasets,

281 including human dorsolateral prefrontal cortex (DLPFC)[60] and cichlid fish telencephalon[61]

282 Visium samples. In both datasets, reference profiles were derived from snRNA-seq, which

283 inherently underrepresent cytoplasmic and process-localized RNA, leading standard

284 deconvolution methods to misattribute unexplained ST signals. Across these datasets,

285 RESCUE consistently outperformed competing methods (**Supplementary Fig. 3-4**),

286 indicating that the benefits of explicitly modeling unattributed expression extend across

287 tissues, species, and reference modalities.

289 **RESCUE reveals unattributed biological processes in diverse tissues**

290 In many neuroscience experiments, cell-level information is often characterized using

291 snRNA-seq[62,63], which can leave a substantial fraction of biologically relevant expression

292 unattributed. Here, we illustrate how RESCUE can uncover unique biological processes

293 within the unattributed expression of the human DLPFC and cichlid fish telencephalon,

294 using snRNA-seq as a reference.

Human DLPFC. Maynard et al.[60] profiled the DLPFC of neurotypical adult donors using

paired snRNA-seq and 10x Visium ST. We applied RESCUE to eight samples in which

all six cortical layers and the white matter (WM) were annotated, to recover unattributed

expression patterns not represented in the snRNA-seq reference (**Fig 5a**). As shown in

Fig. 5b, RESCUE reveals that Layer 1 and the WM contain substantially higher levels of

unattributed expression compared to other cortical layers. This is consistent with the fact

that Layer 1 has sparser cell densities and contains neuropil-rich regions[57], and that WM

contains a substantial amount of axons and myelin[64], whose transcriptomic content is

mostly cytoplasmic and thus is absent from the reference snRNA-seq data.

RESCUE enables us to study biology specific to the unattributed transcriptome of

Layer 1. We first identified DEGs by comparing Layer 1 to all other layers using limma[65],

based on either the raw observed ST expression or the unattributed expression recovered

by RESCUE. These two analyses gave substantially different results (**Fig. 5c**). For

example, the top five genes most enriched in Layer 1 based on unattributed expression

had either fold-changes in the opposite direction (*MT-ND1*, *MT-ND2*, and *MT-CO1*) or

negligible fold-changes (*MT3* and *CST3*) in the differential expression analysis using raw

data. We also identified genes such as *CLU*, *GFAP*, and *MALAT1* that exhibit clear spatial

enrichment within the unattributed component of Layer 1 (**Fig. 5d**). These genes are

associated with local translation and neurite outgrowth, and their enrichment outside

nucleus is consistent with known roles of astrocytic processes and non-somatic cellular

structures in Layer 1 of the cortex[66,67].

To explore the underlying biological processes, we performed gene set enrichment

analysis separately for DEGs derived from the unattributed expression and from the raw

ST data (**Fig. 5e**). Notably, gene ontology (GO) terms enriched among the unattributed

DEGs include mitochondrial processes: "oxidative phosphorylation", "energy derivation

by oxidation of organic compounds", and "generation of precursor metabolites and

energy". In contrast, no mitochondria-related pathways reached statistical significance in

the analysis based on observed ST expression; only the broad term "response to stress"

passed the FDR < 0.05 threshold. These results align with the elevated metabolic

demands of a neurite-enriched Layer 1, which is densely populated with synapses and

reliant on local energy production to support axonal and dendritic activity[68,69]. Collectively,

these findings demonstrate that RESCUE can uncover previously obscured yet

functionally important gene expression programs in Layer 1 of the human DLPFC.

Cichlid Fish Telencephalon. Johnson et al.[61] collected paired snRNA-seq and Visium

ST data from the dorsal telencephalon of individual cichlid fishes exhibiting distinct social

behaviors. As shown in **Fig. 6a**, the dorsal telencephalon is characterized by prominent

radial glial (RG) processes that originate in the ventricular zone and extend long

projections toward superficial pallial regions[70,71]. These RG projections span nearly the

entire depth of the pallium and may play a role in shaping spatial gene expression profiles

across distinct layers and compartments, but they do not survive tissue dissociation and

thus their transcriptomes are not captured by snRNA-seq.

We applied RESCUE to this dataset to recover unattributed transcriptomic signals

using the snRNA-seq data as a reference. To assess spatial heterogeneity, we performed

SVG analysis using SPARK-X[72] on both the observed ST expression and the unattributed

expression. As shown in **Fig. 6b**, a substantial number of SVGs are uniquely identified in

the unattributed transcriptome. In **Fig. 6c**, we show the expression maps of those SVGs

that were undetectable in the observed ST expression yet exhibited clear spatial structure

once nuclear signals were removed. Among these, *app*, *cntn1*, *dner*, and *eno1* mark

distinct transcriptional programs that may operate along RG projections. These genes

highlight transcriptional signals that are likely to reflect localized molecular activities along

RG scaffolds: *app* encodes a transmembrane protein that promotes synapse formation

and plasticity[73,74], *cntn1* (Contactin-1) is a GPI-anchored adhesion molecule essential for

axon guidance and neurite outgrowth[75], *dner* mediates neuron-glia interactions that are

critical for astrocyte differentiation[76], and *eno1* (α-enolase) promotes neuritogenesis and

contributes to structural remodeling in the developing nervous system[77]. Indeed, Johnson

et al.[61] found that bower-building behavior in cichlid fishes may be related to neuronal

rebalancing driven by a subpopulation of RG cells and their processes, highlighting the

importance of studying transcriptional processes in RG projections, which RESCUE

enables.

To further characterize the structure of the unattributed transcriptome, we applied

NMF to the unattributed expression component (**Fig. 6d**). Several unattributed factors are

enriched for genes associated with synaptic and process-localized functions, including

slc1a2, *calm2*, and mitochondrial ATP synthase (**Fig. 6e**), consistent with transcriptional

activity in extended RG scaffolds and neuronal processes. In addition, one factor shows

strong enrichment for ribosomal protein genes, suggesting localized protein synthesis

within these structures[78]. Notably, these structured transcriptional programs emerge only

after the canonical expression is removed, indicating that the unattributed expression

recovered by RESCUE captures biologically meaningful signals associated with radial

glial and cytoplasmic processes that are not represented in snRNA-seq reference data.

DISCUSSION

RESCUE is the first method designed to acknowledge and recover the unattributed transcriptome, which encompasses gene expression patterns not captured by cell segmentation or deconvolution approaches in ST data analysis. These unattributed signals, though typically overlooked, can carry substantial functional and biological significance. While the field has developed numerous transcriptomic profiling methods, such as scRNA-seq for isolating gene expression at the single-cell level, Ribo-seq for ribosome-associated transcripts, and CLIP-seq for binding-protein interactions, each of these technologies inevitably captures only a fraction of the total transcriptome. RESCUE provides a systematic framework to quantify and analyze what is missing: the molecular signals outside the scope of reference datasets.

By formulating the problem as a robust sparse recovery framework, RESCUE decomposes observed ST data into a reference-aligned canonical component and a biologically meaningful unattributed component. RESCUE focuses on characterizing what is not explained by reference profiles and introduces a broader conceptual innovation: computational negative selection. Rather than focusing solely on what to include (as in positive selection), we define a reference component to subtract out and focus on what remains. In this work, we used scRNA-seq and snRNA-seq data as reference profiles and focused primarily on brain tissue, but the concept is general and can be applied to a wide range of other settings, for example, muscle tissue, where multinucleated cells make segmentation challenging and single-cell reference data difficult to interpret. The choice of reference therefore requires careful design and quality control.

One striking finding emerging from our brain tissue analyses is the substantial

amount of expression (26.5% of total expression) found in the unattributed transcriptome

of the honey bee brain MERFISH data. While localization of mRNAs to neurites is a well-

studied phenomenon[26–30], our results, along with the estimates for the adult mouse brain

from Ament and Poulpoulos[22], additionally suggest that these local expression patterns

may be different from those found in cell soma. RESCUE enables a reexamination of

existing datasets to more carefully study this apparently divergent expression. Despite

these promising biological insights revealed by RESCUE, reliance on computational

inference alone may have potential limitations, underscoring the value of experimental

validation when feasible. Orthogonal approaches, such as *in situ* hybridization, could

further strengthen and refine these findings; however, such validation is often impractical

in tissues where cell bodies and non-cell-body structures are spatially intermingled and

difficult to segment. These challenges highlight the importance of discovery-oriented

computational frameworks like RESCUE, which are designed to uncover biologically

meaningful patterns and guide interpretation in settings where experimental confirmation

is difficult.

There are several potential directions for future work. First, in cases where no

suitable reference is available, extensions of RESCUE to reference-free or unsupervised

frameworks may be possible. For example, methods based on robust principal

component analysis (PCA)[79,80] could be adapted to extract unattributed structure without

predefined reference matrices. Second, applying RESCUE to high-resolution ST data

requires creating a predefined grid, and further research is required to develop a

systematic method to determine the optimal grid size. Third, while the gene-level sparsity

assumption underlying RESCUE is biologically plausible in many settings, there may exist

scenarios in which the unattributed component is not sparse but instead exhibits

alternative structure, such as low-rank or low-rank plus sparse residuals. Extending

RESCUE to accommodate such residual regimes represents an important direction for

future research. Despite these remaining challenges, we have already shown that

RESCUE can still provide meaningful analyses of unattributed expression.

Finally, RESCUE reminds us that the biology we wish to study and the

measurements we are able to make are not always one and the same. For instance, we

typically accept single-cell reference data as truly representing the expression of the

entire single cell, but in many cases it does not, as we have illustrated above. Any

measurement technology has inherent biases. As a result, recognizing, acknowledging

and properly interpreting what is missed, in other words, the "unattributed expression", is

important for a comprehensive understanding of a system.

METHODS

RESCUE model

We assume that our ST data profiles G genes in J grids across the tissue section. In low-

resolution ST technologies, these grids are predefined as multicellular spots, whereas in

higher-resolution ST technologies, these can be constructed at any desired size, which

allows inference at different biological scales. To introduce the statistical model behind

RESCUE, we begin by reviewing existing linear deconvolution methods[12–16]. These

methods decompose the ST expression matrix $Y \in \mathbb{R}_{\geq 0}^{G \times J}$ of G genes in J grids as

433
$$Y = X\alpha + E, \tag{1}$$

where $X \in \mathbb{R}_{\geq 0}^{G \times K}$ is a matrix of K cell-type expression reference profiles, $\alpha \in \mathbb{R}_{\geq 0}^{K \times J}$ is a

matrix of inferred cell-type proportions within each grid, and $E \in \mathbb{R}^{G \times J}$ is assumed to be a

mean-zero noise matrix. The issue is that (1) implies that expectation of ST expression

is $\mathbb{E}(Y) = X\alpha$, which ignores unattributed transcriptome.

Instead, we further decompose the error E, motivated by the robust regression

framework[45–47]. Specifically, we consider the multivariate mean-shift model:

440
$$Y = X\alpha + R + \mathcal{E}, \tag{2}$$

where $X\alpha$ represents the expected expression values of each gene in each grid due to

the presence of cells whose types are represented in the reference sc/snRNA-seq data,

$R \in \mathbb{R}_{\geq 0}^{G \times J}$ is an unattributed term that allows for outliers that cannot be captured by the

reference data, and $\mathcal{E} \in \mathbb{R}^{G \times J}$ is a noise matrix that allows for residual variability between

the ST sample and the reference data.

Following the mean-shift model in the robust regression framework, we assume

that the unattributed component R is sparse. Specifically, the sparsity assumption reflects

the expectation that only a subset of genes exhibits unattributed expression within each

grid. This assumption is required for identifiability of the decomposition and is biologically

plausible, as we expect relatively few genes to exhibit unattributed expression within each

grid. Consistent with this assumption, we observe empirically that unattributed expression

in neuropil regions of the honey bee brain is highly sparse, with 92% of genes showing

no unattributed expression on average. Moreover, even in extracellular environments,

ECM proteins typically involve highly sparse, restricted gene sets, with only 1.3–1.5% of

genes showing nonzero expression, rather than transcriptome-wide activation[81].

Optimization problem

RESCUE aims to simultaneously recover the sparse non-negative matrix R and non-

negative cell-type proportions α in our model (2). Specifically, we consider the following

optimization problem:

$$\min_{\alpha,R\geq0}\|W \odot R\|_1 \ \text{ subject to } \ \|Y - R - X\alpha\|_F^2 \leq \lambda,$$

where W is a weight matrix, $\odot$ denotes the Hadamard product, $\|\cdot\|_F$ denotes the

Frobenius norm, and $\lambda > 0$ is a tuning parameter. We set each entry of the weight matrix

as $W_{gj} = 1/\sqrt{Var(R_{gj})}$ to account for the fact that different genes have different variances

across different locations. These weights will be estimated through an iterative variance

estimation procedure. The ℓ_1 objective function enforces sparsity of our estimated R,

while the constraint ensures that the residual variability $\mathcal{E}$ is not too large, in other words,

that the ST data do not deviate too much from the reference sc/snRNA-seq data.

In practice, we solve the Lagrangian form of this problem:

$$\min_{\alpha,R\geq0} f(\alpha, R) = \|W \odot R\|_1 + \frac{\mu}{2}\|Y - R - X\alpha\|_F^2, \tag{3}$$

where $\mu > 0$ is a tuning parameter. We apply an efficient alternating minimization to solve

two convex sub-problems in each iteration:

$$\hat{\alpha} = \underset{\alpha \geq 0}{\text{argmin}} \|Y - R - X\alpha\|_F^2, \tag{4}$$

$$\hat{R} = \underset{R \geq 0}{\text{argmin}} \|W \odot R\|_1 + \frac{\mu}{2}\|Y - R - X\alpha\|_F^2 = S_+\left(Y - X\alpha, \frac{W}{\mu}\right), \tag{5}$$

where $S_+(x, \tau) = \max(x - \tau, 0)$ denotes the entry-wise non-negative soft-thresholding

function.

Estimation of adaptive weights

Kong et al.[47] proposed adaptive lasso-type weights to reduce masking and swamping

effects in the presence of multiple outliers. Our weighting scheme plays a similar role, but

differs in that the weights are learned iteratively through variance estimation of the

unattributed component. In ST deconvolution models, the observed count $\tilde{Y}_{gj}$ is often

modeled under a Poisson assumption[13,16]:

$$\tilde{Y}_{gj}|\lambda_{gj} \sim \text{Poisson}(N_j\lambda_{gj})$$

485 where $N_j = \sum_g \tilde{Y}_{gj}$ represents the total transcript count, or number of UMIs for grid j, and

486 λ_{gj} denotes the expected expression rate of gene g at grid j. Rather than directly

487 modeling $\tilde{Y}_{gj}$, we assume that it can be decomposed as $\tilde{Y}_{gj} = \tilde{L}_{gj} + \tilde{R}_{gj}$, where

$$\tilde{L}_{gj}|\lambda_{gj}^L \sim \text{Poisson}(N_j^L\lambda_{gj}^L) \quad \text{and} \quad \tilde{R}_{gj}|\lambda_{gj}^R \sim \text{Poisson}(N_j^R\lambda_{gj}^R).$$

Here, $N_j^L = \sum_g \tilde{L}_{gj}$ and $N_j^R = \sum_g \tilde{R}_{gj}$ represent the total transcript counts of the canonical

and unattributed expression for grid j, respectively, and $(\lambda_{gj}^L, \lambda_{gj}^R)$ denote the expected

expression rates of these components. We can write $N_j^L = (1 - \pi_j)N_j$ and $N_j^R = \pi_j N_j$,

where $\pi_j \in [0, 1]$ denotes the proportion of unattributed expression for grid j.

When using counts per million (CPM) normalization, i.e., $Y_{gj} = \tilde{Y}_{gj}/N_j \times 10^6$, we

can estimate the variance of unattributed component as

$$\text{Var}(R_{gj}) = \text{Var}\left(\frac{\tilde{R}_{gj}}{N_j} \times 10^6\right) = \left(\frac{10^6}{N_j}\right)^2 \times \text{Var}(\tilde{R}_{gj}) \approx \frac{10^6}{N_j} \times \hat{R}_{gj}, \qquad (6)$$

496   where the approximation follows from $\widehat{\text{Var}}(\tilde{R}_{gj}) = \widehat{N_j^R \lambda_{gj}^R}$ and $\tilde{R}_{gj} = R_{gj}/10^6 \times N_j$. Here,

497   the term $\widehat{N_j^R \lambda_{gj}^R}$ can be iteratively updated using the plug-in estimator $\tilde{R}_{gj} \approx N_j^R \lambda_{gj}^R$. To

498   avoid division by zero, we set $W_{gj} = 1/\max\left(\sqrt{\frac{10^6}{N_j} \times R_{gj}}, \delta\right)$, where δ is a small positive

499   constant. We outline the iterative algorithm for RESCUE below (**Algorithm 1**).

Algorithm 1 Iterative algorithm for RESCUE

Input: Normalized expression $Y \in \mathbb{R}_{\geq 0}^{G \times J}$, observed UMI counts $N \in \mathbb{N}^J$, reference profile
    $X \in \mathbb{R}_{\geq 0}^{G \times K}$, $\mu > 0$, $\delta > 0$, and $\epsilon > 0$.

Output: Cell-type proportion $\alpha \in \mathbb{R}_{\geq 0}^{K \times J}$ and unattributed expression $R \in \mathbb{R}_{\geq 0}^{G \times J}$.

  1: Initialize $R^{(0)} = 0$, $W_{gj}^{(0)} = 1$, and $t = 1$.
  2: **repeat**
  3:    $\alpha^{(t)} = \text{NNLS}(X, Y - R^{(t-1)})$ from (4).
  4:    $R^{(t)} = S_+(Y - X\alpha^{(t)}, W/\mu)$ from (5).
  5:    **for** $g = 1$ to G and $j = 1$ to J **do**
  6:       $W_{gj}^{(t)} = 1/\max\left(\sqrt{\frac{10^6}{N_j} \times R_{gj}^{(t)}}, \delta\right)$ from (6).
  7:    **end for**
  8:    $t = t + 1$.
  9: **until** $|f(\alpha^{(t)}, R^{(t)}) - f(\alpha^{(t-1)}, R^{(t-1)})|/f(\alpha^{(t)}, R^{(t)}) < \epsilon$, where f is defined in (3).
Return: $\alpha^{(t)}$ and $R^{(t)}$.

Tuning parameter selection

RESCUE aims to recover unattributed expression by balancing the sparsity of R, which

reflects the number of rejected outliers, against the level of residual variability $\mathcal{E}$. Following

standard practices in robust regression and robust PCA[45,79], we set $\mu = GJ/C\|Y\|_1$ and

focus on selecting the optimal value of the tuning parameter C. However, choosing the

optimal tuning parameter is challenging in the presence of outliers[80,82].

To select the optimal tuning parameter, we adopt a data-driven strategy recently

proposed for related deconvolution problems in bulk RNA-seq anaysis[83]. The key idea is

to generate simulated data that closely resemble the observed ST data and use this for

evaluation. Specifically, we use a surrogate ST dataset with "ground truth" proportions of

canonical and unattributed expression. To construct this surrogate ST dataset, we first

apply NNLS to the observed ST data to obtain initial estimates for α and R:

514
$$\hat{R}_{NNLS} = (Y - X\hat{\alpha}_{NNLS})_+, \quad \text{where} \quad \hat{\alpha}_{NNLS} = \underset{\alpha \geq 0}{\operatorname{argmin}} \|Y - X\alpha\|_F^2, \tag{7}$$

where $(x)_+ = \max(x, 0)$ denotes the zero-thresholding function. We then simulate a

surrogate ST data $\tilde{Y}_{\text{sim}}$ from a Poisson distribution with rescaled expected expression

rates, preserving the overall signal-to-noise characteristics observed in the real data.

RESCUE is applied to $\tilde{Y}_{\text{sim}}$ across a range of μ values, and the optimal C is selected as

the value that minimizes the recovery error of the canonical and unattributed components

in the surrogate dataset. We summarize the tuning procedure below (**Algorithm 2**).

Algorithm 2 Tuning parameter selection via surrogate match

Input: Normalized expression $Y \in \mathbb{R}_{\geq 0}^{G \times J}$, observed UMI counts $N \in \mathbb{N}^J$, reference profile
    $X \in \mathbb{R}_{\geq 0}^{G \times K}$, and $S > 0$.
Output: Optimal tuning parameter μ.
  1: **for** $s = 1$ to S **do**
  2:    Obtain $\hat{L}_{\text{NNLS}} = X\hat{\alpha}_{\text{NNLS}}$ and $\hat{R}_{\text{NNLS}}$ from (7).
  3:    Simulate ST data surrogate $\tilde{Y}_{\text{sim}} \sim \text{Pois}(N_j \lambda_{gj})$ with $\lambda_{gj} = (\hat{L}_{\text{NNLS}} + \hat{R}_{\text{NNLS}})/10^6$.
  4:    Apply Algorithm 1 with normalized Y_{sim} to obtain $\hat{L} = X\hat{\alpha}$ and $\hat{R}$.
  5:    Evaluate $\eta_C^{(s)} = \max\left(\|\hat{L}_{\text{NNLS}} - \hat{L}\|_F / \|\hat{L}_{\text{NNLS}}\|_F, \|\hat{R}_{\text{NNLS}} - \hat{R}\|_F / \|\hat{R}_{\text{NNLS}}\|_F\right)$.
  6: **end for**
Return: $\mu = GJ/C\|Y\|_1$, where $C = \operatorname{argmin}_C\left(\frac{1}{S}\sum_{s=1}^S \eta_C^{(s)}\right)$.

Platform effect normalization

To mitigate cross-platform effects between ST data and sc/snRNA-seq reference data, we apply platform effect normalization. This step is essential because differences in experimental platforms can introduce systematic shifts in gene expression levels that confound downstream analysis. Without proper alignment of scale, these differences may lead to biased estimation of cell-type contributions or unattributed signals. We adopt the normalization strategy from RCTD[13], which rescales each gene in the reference data to match the overall expression level of the corresponding gene in the ST data. Specifically, it first computes pseudo-bulk measurements for both ST and reference datasets and then apply a gene-wise log-scale correction to harmonize them.

Data normalization and reference matrix

To account for technical variability and the effect of library size, we adopt CPM normalization but fit RESCUE in the linear space. We avoid log-transformation for two main reasons. First, log-transformed input tends to underestimate true signals in gene expression deconvolution[84]. Second, log-transformation requires adding pseudo-counts, which can introduce significant artificial distortion between zeros and non-zeros[85,86]. Instead, we work in the linear scale, employing an iteratively reweighted algorithm (**Algorithm 1**) to stabilize variance adaptively. This approach allows the model to adjust for varying data scales and minimizes the influence of outliers without artificially altering data distributions. The normalized reference matrix is then used to construct average cell-type expression.

Reference dependence of unattributed expression

RESCUE is a computational negative selection method which relies on the choice of reference, and interpretation of the recovered unattributed expression therefore requires consideration of both biological context and reference completeness. To further assess the impact of reference selection within the same tissue, we extended the honey bee brain analysis using snRNA-seq[87] and segmented MERFISH cells as alternative reference profiles (**Supplementary Fig. 5**). Using segmented MERFISH cells produced unattributed expression patterns highly consistent with those obtained using scRNA-seq, as both references capture cellular transcripts beyond the nucleus and therefore explain similar fractions of the ST signal. In contrast, the snRNA-seq reference yielded higher levels of unattributed expression on average, reflecting its enrichment for nuclear RNA and underrepresentation of cytoplasmic and neuritic transcripts. These results indicate that variation in the recovered unattributed component reflects biologically meaningful differences in reference coverage rather than technical artifacts, and that interpretation of unattributed expression should be guided by the biological properties of both the tissue and the reference dataset.

Choice of grid resolution for honey bee brain MERFISH data

For high-resolution ST platforms such as MERFISH, RESCUE operates on spatially aggregated expression profiles defined over user-specified grids. The choice of grid size reflects a trade-off between spatial resolution and sparsity: smaller grids preserve fine spatial structure but increase sparsity, whereas larger grids reduce sparsity at the cost of spatial detail. As a practical guideline, we recommend selecting a grid size such that the

average total transcript count per grid is comparable to the typical total transcript count

per cell. In the honey bee MERFISH analysis, we used a grid size of $20 \times 20 \; \mu m^2$, which

approximates cell-scale resolution. To assess sensitivity to grid resolution, we varied grid

size across a range of spatial scales and found that RESCUE produces consistent results

across a broad range of grid resolutions (**Supplementary Fig. 6**).

Subtraction-based estimation of unattributed expression for existing methods

Most existing ST deconvolution methods are designed to estimate cell-type proportions,

rather than to recover expression levels directly. Because these methods do not explicitly

model unattributed expression, we adopted a subtraction-based framework, following a

strategy used in related deconvolution settings for bulk RNA-seq data[23]. Specifically, we

first estimated cell-type proportions using each deconvolution method and then recovered

the unattributed expression by subtracting the estimated cellular contributions from the

observed expression, followed by zero-thresholding to avoid negative values. Under this

framework, the unattributed expression is defined as the portion of the observed ST

expression that cannot be explained by the reference-based cellular expression implied

by each method's estimated cell-type proportions.

Data generating process for simulation studies

For a total of $J = 2{,}500$ grids, we first generated the total UMI count for grid j from a

negative binomial distribution $N_j \sim \mathrm{NB}(\mathrm{mean} = M, \mathrm{size} = 1/\sigma)$. We generated the raw

count for each gene in each grid as $\tilde{Y}_{gj} = \tilde{L}_{gj} + \tilde{R}_{gj}$, where $\tilde{R}_{gj}$ denotes the unattributed

expression and $\tilde{L}_{gj}$ represents the canonical expression that could have been captured

by sc/snRNA-seq reference data.

To generate $\tilde{L}_{gj}$, we used the real scRNA-seq data from honey bee brains[48] as the

source of canonical expression profiles. We selected the top $G = 500$ number of highly

variable genes and constructed the reference matrix X using $K = 20$ number of cell types

annotated via Louvain clustering in Seurat[88]. Latent mixing proportions were drawn as

$\beta_{kj} \sim \text{Beta}(1,3)$ and normalized as $\alpha_{kj} = \beta_{kj}/\sum_k \beta_{kj}$, with at most 10 cell types per spot.

Defining $\lambda_{gj}^L = (X\alpha)_{gj}$, we generated the cell-type-driven canonical expression as

599
$$\tilde{L}_{gj} \sim \text{Multinomial}\left((1 - \pi_j)N_j, P_{gj} = \frac{\lambda_{gj}^L}{\sum_g \lambda_{gj}^L}\right).$$

600   To generate sparse unattributed expression, we sampled

601
$$\tilde{R}_{gj} \sim \text{Multinomial}\left(\pi_j N_j, Q_{gj} = \frac{\lambda_{gj}^R}{\sum_g \lambda_{gj}^R}\right),$$

where $\lambda_{gj}^R \sim \text{Gamma}(1, 1/3)$ with probability $(1 - s)$ and $\lambda_{gj}^R = 0$ with probability s. Here,

we introduced a sparsity parameter s, which makes $(s \times 100)$ percentage of intensity

parameter λ_{gj}^R zero-inflated. Under this setting, we can generate $\tilde{L}_{gj}$ and $\tilde{R}_{gj}$ that satisfy

$\sum_g(\tilde{L}_{gj} + \tilde{R}_{gj}) = N_j$. In our simulations, we fixed $M = 10,000$ and $\sigma = 0.5$, while varying

$\pi_j \in \{0, 0.2, 0.4, 0.6, 0.8\}$ and $s \in \{0.8, 0.85, 0.9, 0.95\}$ to control the proportion of canonical

expression and the sparsity level of unattributed transcriptome, respectively (**Fig. 2**). We

then examined the robustness of RESCUE under additional simulation settings by varying

the average total UMI count per spot (M), as well as varying noise levels by adjusting

library size variability across spots (σ) (**Supplementary Fig. 1**).

Evaluation metrics for cell-type deconvolution in simulation studies

To evaluate the accuracy of cell-type deconvolution, we considered three complementary metrics: Lin's CCC, RMSE, and JSD. We first measured Lin's CCC between the true and estimated proportion of the kth cell type, defined as

$$\text{CCC}(\alpha_k, \hat{\alpha}_k) = \frac{2\text{Cov}(\alpha_k, \hat{\alpha}_k)}{\sigma_{\alpha_k}^2 + \sigma_{\hat{\alpha}_k}^2 + \left(\mu_{\alpha_k} - \mu_{\hat{\alpha}_k}\right)^2},$$

where μ_{α_k} and $\mu_{\hat{\alpha}_k}$ are the means of the two variables, and $\sigma_{\alpha_k}^2$ and $\sigma_{\hat{\alpha}_k}^2$ denote the corresponding variances. We used Lin's CCC because it captures shifts in both location and scale, making it robust to high leverage points[83]. The concordance increases as the CCC value gets closer to 1. We report the average CCC across all cell types.

To quantify estimation error at the spatial location level, we computed the RMSE between the true and estimated cell-type proportion vectors. For spatial location j, RMSE is defined as

$$\text{RMSE}_j = \sqrt{\frac{1}{K}\sum_{k=1}^{K}\left(\alpha_{kj} - \hat{\alpha}_{kj}\right)^2},$$

where K is the number of cell types, and α_{kj} and $\hat{\alpha}_{kj}$ denote the true and estimated proportions of cell type k at location j, respectively. We report the average RMSE across all spatial locations, with smaller values indicating better performance.

Finally, we computed the JSD between the true and estimated cell-type proportion distributions at each spatial location. For spatial location j, JSD is defined as

$$\text{JSD}_j = \frac{1}{2}\text{KL}(\alpha_j \| m_j) + \frac{1}{2}\text{KL}(\hat{\alpha}_j \| m_j), \qquad m_j = \frac{1}{2}(\alpha_j + \hat{\alpha}_j),$$

631   where $\mathrm{KL}(\cdot \parallel \cdot)$ denotes the Kullback-Leibler divergence. We report the average JSD

632   across all spatial locations, with smaller values indicating better performance.

634   **Evaluation metrics for unattributed expression recovery in simulation studies**

635   To evaluate the accuracy of unattributed expression recovery, we measured the relative

636   Frobenius error, defined as

637
$$(\text{Relative Frobenius Error}) = \frac{\|R - \hat{R}\|_F}{\|R\|_F},$$

638   where R and $\hat{R}$ denote the true and estimated unattributed expression matrices,

639   respectively.

641   **Evaluation metrics for honey bee brain MERFISH data**

642   We measured Lin's CCC and the RMSE between the segmentation-based (true) and

643   estimated unattributed expression proportions, defined as follows:

644
$$\mathrm{CCC}(\rho, \hat{\rho}) = \frac{2\mathrm{Cov}(\rho, \hat{\rho})}{\sigma_\rho^2 + \sigma_{\hat{\rho}}^2 + (\mu_\rho - \mu_{\hat{\rho}})^2} \quad \text{and} \quad \mathrm{RMSE}(\rho, \hat{\rho}) = \sqrt{\frac{1}{J}\sum_{j=1}^{J}(\rho_j - \hat{\rho}_j)^2},$$

where $\rho_j = \sum_g \tilde{R}_{gj} / \sum_g \tilde{Y}_{gj}$ denotes the true unattributed expression proportion in grid j,

and $\hat{\rho}_j$ denotes its estimated value. Here, μ_ρ and $\mu_{\hat{\rho}}$ are the means of two variables, and

σ_ρ^2 and $\sigma_{\hat{\rho}}^2$ denote the corresponding variances.

Benchmarking computational runtime

We evaluated the computational runtime of each method across ST datasets of varying

sizes (**Supplementary Fig. 7**). Across all benchmarks, RESCUE completed within

practical time limits for contemporary ST datasets, demonstrating scalability to datasets

with hundreds to thousands of spatial locations and genes. All analyses were performed

using CPU-only computation on a standard laptop (Apple M3 Pro with 36GB of memory),

without GPUs or high-performance computing clusters.

DATA AVAILABILITY

Human breast cancer dataset[5]: 10x Visium and scFFPE-seq data are available via GEO

accession GSE243280. Human DLPFC dataset[60]: 10x Visium and snRNA-seq data are

available at https://github.com/LieberInstitute/spatialLIBD. Cichlid fish brain dataset[61]:

10x Visium and snRNA-seq data are available via GEO accession GSE217619. Honey

bee brain MERFISH data is available upon reasonable request. Honey bee brain scRNA-

seq data[48] is available at https://doi.org/10.6084/m9.figshare.16832518.

CODE AVAILABILITY

RESCUE is available as an R package at https://github.com/brunoyjlee/RESCUE.

ACKNOWLEDGEMENTS

We thank Dr. Tina Barbasch on valuable discussions on the analysis of fish brain data.

Funding: This work was supported by the National Institutes of Health (R21HG013180 to

Y.J.L., S.Y., H.-S.H. and S.D.Z., R35GM147420 to A.W.S and H.-S.H. and R01AT013189

to S.D.Z.).

FIGURES

Figure 1. Schematic overview of RESCUE. RESCUE is a novel computational negative

selection method designed to recover the spatial expression patterns missed by standard

single-cell inference on ST data. It takes as input ST data to be decomposed, along with

a reference matrix constructed from sc/snRNA-seq data from the same tissue, and

performs robust sparse recovery via a regularized multivariate regression framework.

RESCUE partitions observed gene expression into a reference-explainable "canonical"

component and a residual "unattributed" component. As a result, the purified canonical

expression enables more accurate deconvolution, while the recovered unattributed

expression enables downstream analyses such as clustering, differentially expressed

gene (DEG), and spatially variable gene (SVG) analyses. Created with BioRender.com.

Figure 2. Simulation results. a. Schematic of the simulation framework. ST data are

generated as the sum of reference-driven canonical expression and synthetic unattributed

expression. The distribution of sparsity levels observed in neuropil regions of the honey

bee brain motivates the simulated sparsity regimes. Representative examples of synthetic

unattributed expression under low- and high-sparsity/expression levels are shown. **b.**

Cell-type deconvolution performance across methods, evaluated using average Lin's

concordance correlation coefficient (CCC; top), root mean squared error (RMSE; middle),

and Jensen-Shannon divergence (JSD; bottom) as a function of the sparsity/proportion

levels of the unattributed expression. **c.** Unattributed expression recovery performance

measured by relative Frobenius error under the same simulation settings.

Figure 3. Application to the honey bee brain MERFISH data. a. MERFISH profiling of a coronal honey bee brain section. Each dot represents an individual transcript, with genes shown in distinct colors. A substantial fraction of transcripts localize to neurites, rather than to cell bodies. **b.** Comparison of segmentation-based ground-truth ("Cell Mask") and estimated somatic proportions. Cell mask ratio is computed as the proportion of MERFISH spots in each grid lying within a cell. Lin's concordance correlation coefficient (CCC) and root mean squared error (RMSE) between ground-truth and estimated proportions are reported. **c.** Comparison of estimated versus ground-truth canonical gene expression levels in grids corresponding to pure cell body regions. **d.** Comparison of estimated versus ground-truth unattributed gene expression levels in neuropil-only grids lacking segmented somata. For **c** and **d,** each point represents a gene averaged across grids; grey dashed lines indicate the identity line, and blue lines denote linear regression fits. Lin's CCC and RMSE between ground-truth and estimated average expression are reported. **e.** Clustering of the unattributed expression profiles inferred by RESCUE reveals distinct transcriptional niches within neuropil regions, while somatic regions collapse into a single cluster. **f.** Spatial expression patterns of representative genes, shown separately for canonical and unattributed components. Expression values are min-max normalized separately for each panel.

Figure 4. Application to the human breast cancer Visium data. a. Manual annotation of Visium ST data from human breast cancer tissue. **b.** Estimated cell-type proportions in adipocyte regions across methods. Red box highlights that existing deconvolution methods misattribute adipocyte-derived signal to other reference cell types due to the absence of adipocytes in the reference, whereas RESCUE mitigates such misassignment by modeling unattributed component. **c.** Heatmap of correlations between pseud-bulk Visium ST and scFFPE-seq reference cell types. Red box highlights high correlations between adipocytes and other cell types including stromal and endothelial cells. **d.** Comparison of cell-type deconvolution accuracy across methods using Lin's concordance correlation coefficient (CCC), root mean squared error (RMSE), and Jensen-Shannon divergence (JSD). **e.** Comparison of estimated versus reference cell-type proportions across methods. Each point represents a cell type, with grey dashed lines indicating the identity line. **f.** Non-negative matrix factorization (NMF) of the canonical and unattributed expression component inferred by RESCUE. Weight values are min-max normalized separately for each panel. **g.** Spatial expression patterns of adipocyte marker genes, shown separately for canonical and unattributed components. Expression values are min-max normalized separately for each panel.

Figure 5. Application to the human DLPFC Visium data. a. Anatomical annotations of

Visium ST samples from the DLPFC dataset, with cortical layers (Layer 1-6) and white

matter (WM) labeled. **b.** Proportions of unattributed expression estimated by RESCUE

across cortical layers and WM. Layer 1 and WM exhibit substantially higher unattributed

expression compared to other layers. **c.** Comparison of log-fold changes for differentially

expressed gene (DEG) analysis comparing Layer 1 to all other layers using limma on raw

observed ST expression (x-axis) versus unattributed expression recovered by RESCUE

(y-axis). Blue dashed lines indicate zero-fold change. Genes highlighted in red are top

Layer 1 markers, detected from unattributed expression. **d.** Spatial expression patterns

of Layer 1 marker genes derived from unattributed expression, shown separately for

canonical and unattributed components. Expression values are min-max normalized

separately for each panel. **e.** Gene ontology (GO) terms enriched among Layer 1-

enriched DEGs based on unattributed (left) and raw (right) expression.

Figure 6. Application to the cichlid fish telencephalon Visium data. a. Schematic

illustration of radial glial (RG) anatomy[61] in the dorsal telencephalon, highlighting RG cell

bodies in the ventricular zone and long processes extending toward superficial pallial

regions. **b.** Comparison of adjusted p-values for spatially variable gene (SVG) detection

using SPARK-X on raw observed ST expression (x-axis) versus unattributed expression

recovered by RESCUE (y-axis). Blue dashed lines indicate the significance threshold at

adjusted p-value = 0.05. Genes labeled in red are SVGs uniquely detected from the

unattributed expression. Adjusted p-values are obtained using SPARK-X's default option:

Benjamini-Yekutieli procedure. **c.** Spatial expression patterns of unattributed-only SVGs,

shown separately for canonical and unattributed components. Expression values are min-

max normalized separately for each panel. **d.** Non-negative matrix factorization (NMF) of

the canonical and unattributed expression component inferred by RESCUE. Weight

values are min-max normalized separately for each panel. **e.** Heatmaps of genes with

high loadings in canonical (left) and unattributed (right) factors.

REFERENCES

1. Rao, A., Barkley, D., França, G. S. & Yanai, I. Exploring tissue architecture using spatial transcriptomics. *Nature* **596**, 211–220 (2021).

2. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat Methods* **19**, 534–546 (2022).

3. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* https://doi.org/10.1126/science.aaa6090 (2015) doi:10.1126/science.aaa6090.

4. Nagendran, M. *et al*. 1457 Visium HD enables spatially resolved, single-cell scale resolution mapping of FFPE human breast cancer tissue. *J Immunother Cancer* **11**, (2023).

5. Janesick, A. *et al*. High resolution mapping of the tumor microenvironment using integrated single-cell, spatial and in situ analysis. *Nat Commun* **14**, 8353 (2023).

6. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular segmentation. *Nat Methods* **18**, 100–106 (2021).

7. Petukhov, V. *et al*. Cell segmentation in imaging-based spatial transcriptomics. *Nat Biotechnol* **40**, 345–354 (2022).

8. Polański, K. *et al*. Bin2cell reconstructs cells from high resolution Visium HD data. *Bioinformatics* **40**, btae546 (2024).

9. Ståhl, P. L. *et al*. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78–82 (2016).

10. Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide

expression at high spatial resolution. *Science* **363**, 1463–1467 (2019).

11. Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular resolution

with Slide-seqV2. *Nat Biotechnol* **39**, 313–319 (2021).

12. Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF

regression to deconvolute spatial transcriptomics spots with single-cell

transcriptomes. *Nucleic Acids Res* **49**, e50 (2021).

13. Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial

transcriptomics. *Nat Biotechnol* **40**, 517–526 (2022).

14. Ma, Y. & Zhou, X. Spatially informed cell-type deconvolution for spatial

transcriptomics. *Nat Biotechnol* **40**, 1349–1359 (2022).

15. Zhou, Z., Zhong, Y., Zhang, Z. & Ren, X. Spatial transcriptomics deconvolution at single-

cell resolution using Redeconve. *Nat Commun* **14**, 7930 (2023).

16. Liu, Z., Wu, D., Zhai, W. & Ma, L. SONAR enables cell type deconvolution with spatially

weighted Poisson-Gamma model for spatial transcriptomics. *Nat Commun* **14**, 4727

(2023).

17. Si, Y. *et al.* FICTURE: scalable segmentation-free analysis of submicron-resolution

spatial transcriptomics. *Nat Methods* **21**, 1843–1854 (2024).

18. Mitchel, J., Gao, T., Cole, E., Petukhov, V. & Kharchenko, P. V. Impact of Segmentation

Errors in Analysis of Spatial Transcriptomics Data. 2025.01.02.631135 Preprint at

https://doi.org/10.1101/2025.01.02.631135 (2025).

19. Nguyen, H., Nguyen, H., Tran, D., Draghici, S. & Nguyen, T. Fourteen years of cellular

deconvolution: methodology, applications, technical evaluation and outstanding

challenges. *Nucleic Acids Res* **52**, 4761–4783 (2024).

20. Garmire, L. X. *et al.* Challenges and perspectives in computational deconvolution of

genomics data. *Nat Methods* **21**, 391–400 (2024).

21. Gaspard-Boulinc, L. C., Gortana, L., Walter, T., Barillot, E. & Cavalli, F. M. G. Cell-type

deconvolution methods for spatial transcriptomics. *Nat Rev Genet* 1–19 (2025)

doi:10.1038/s41576-025-00845-y.

22. Ament, S. A. & Poulopoulos, A. The brain's dark transcriptome: Sequencing RNA in

distal compartments of neurons and glia. *Current Opinion in Neurobiology* **81**, 102725

(2023).

23. Ivich, A. *et al.* Missing cell types in single-cell references impact deconvolution of bulk

data but are detectable. *Genome Biology* **26**, 86 (2025).

24. Han, J. *et al.* Single-cell sequencing unveils key contributions of immune cell

populations in cancer-associated adipose wasting. *Cell Discov* **8**, 122 (2022).

25. Fetit, R. *et al.* Characterizing Neutrophil Subtypes in Cancer Using scRNA Sequencing

Demonstrates the Importance of IL1β/CXCR2 Axis in Generation of Metastasis-specific

Neutrophils. *Cancer Research Communications* **4**, 588–606 (2024).

26. Sutton, M. A. *et al.* Miniature Neurotransmission Stabilizes Synaptic Function via Tonic

Suppression of Local Dendritic Protein Synthesis. *Cell* **125**, 785–799 (2006).

27. Cajigas, I. J. *et al.* The Local Transcriptome in the Synaptic Neuropil Revealed by Deep

Sequencing and High-Resolution Imaging. *Neuron* **74**, 453–466 (2012).

28. Hafner, A.-S., Donlin-Asp, P. G., Leitch, B., Herzog, E. & Schuman, E. M. Local protein

synthesis is a ubiquitous feature of neuronal pre- and postsynaptic compartments.

Science **364**, eaau3644 (2019).

29. Glock, C. *et al.* The translatome of neuronal cell bodies, dendrites, and axons.

Proceedings of the National Academy of Sciences **118**, e2113929118 (2021).

30. von Kügelgen, N. & Chekulaeva, M. Conservation of a core neurite transcriptome

across neuronal types and species. *WIREs RNA* **11**, e1590 (2020).

31. Lee, K.-W., Kim, H.-J., Sung, J.-J., Park, K.-S. & Kim, M. Defective neurite outgrowth in

aphidicolin/cAMP-induced motor neurons expressing mutant Cu/Zn superoxide

dismutase. *International Journal of Developmental Neuroscience* **20**, 521–526 (2002).

32. Markmiller, S. *et al.* Persistent mRNA localization defects and cell death in ALS neurons

caused by transient cellular stress. *Cell Reports* **36**, 109685 (2021).

33. Buxbaum, A. R., Haimovich, G. & Singer, R. H. In the right place at the right time:

visualizing and understanding mRNA localization. *Nat Rev Mol Cell Biol* **16**, 95–109

(2015).

34. Gandalovičová, A., Vomastek, T., Rosel, D. & Brábek, J. Cell polarity signaling in the

plasticity of cancer cell invasiveness. *Oncotarget* **7**, 25022–25049 (2016).

35. Backlund, M. *et al.* Plasticity of nuclear and cytoplasmic stress responses of RNA-

binding proteins. *Nucleic Acids Res* **48**, 4725–4740 (2020).

36. Apicco, D. J. *et al.* Reducing the RNA binding protein TIA1 protects against tau-

mediated neurodegeneration in vivo. *Nat Neurosci* **21**, 72–80 (2018).

37. Wolozin, B. & Ivanov, P. Stress granules and neurodegeneration. *Nat Rev Neurosci* **20**,

649–666 (2019).

38. Ridder, K. *et al.* Extracellular vesicle-mediated transfer of functional RNA in the tumor

microenvironment. *OncoImmunology* **4**, e1008371 (2015).

39. Li, I. & Nabet, B. Y. Exosomes in the tumor microenvironment as mediators of cancer

therapy resistance. *Molecular Cancer* **18**, 32 (2019).

40. O'Brien, K., Breyne, K., Ughetto, S., Laurent, L. C. & Breakefield, X. O. RNA delivery by

extracellular vesicles in mammalian cells and its applications. *Nat Rev Mol Cell Biol*

21, 585–606 (2020).

41. Landry, F. J. & Findlay, S. R. Purification of human basophils by negative selection.

Journal of Immunological Methods **63**, 329–336 (1983).

42. Hansel, T. T. *et al.* Purification of human blood eosinophils by negative selection using

immunomagnetic beads. *Journal of Immunological Methods* **122**, 97–103 (1989).

43. Hansel, T. T. *et al.* An improved immunomagnetic procedure for the isolation of highly

purified human blood eosinophils. *Journal of Immunological Methods* **145**, 105–110

(1991).

44. Munoz, N. M. & Leff, A. R. Highly purified selective isolation of eosinophils from human

peripheral blood by negative immunomagnetic selection. *Nat Protoc* **1**, 2613–2620

(2006).

45. She, Y. & Owen, A. B. Outlier Detection Using Nonconvex Penalized Regression. *Journal*

of the American Statistical Association **106**, 626–639 (2011).

46. Lee, Y., MacEachern, S. N. & Jung, Y. Regularization of Case-Specific Parameters for

Robustness and Efficiency. *Statistical Science* **27**, 350–372 (2012).

47. Kong, D., Bondell, H. D. & Wu, Y. Fully Efficient Robust Estimation, Outlier Detection

and Variable Selection Via Penalized Regression. *Statistica Sinica* **28,** 1031–1052

(2018).

48. Traniello, I. M. *et al.* Single-cell dissection of aggression in honeybee colonies. *Nat Ecol*

Evol **7**, 1232–1244 (2023).

49. Lin, L. I.-K. A Concordance Correlation Coefficient to Evaluate Reproducibility.

Biometrics **45**, 255–268 (1989).

50. Kumar, A. *et al.* Intracellular spatial transcriptomic analysis toolkit (InSTAnT). *Nat*

Commun **15**, 7794 (2024).

51. Kim, Y. *et al.* SpaceExpress enables differential spatial transcriptomics with natural

coordinate systems. 2024.12.19.628720 Preprint at

https://doi.org/10.1101/2024.12.19.628720 (2025).

52. Yeo, S., Schrader, A. W., Lee, J., Asadian, M. & Han, H.-S. Spot-Based Global

Registration for Subpixel Stitching of Single-Molecule Resolution Images for Tissue-

Scale Spatial Transcriptomics. *Anal. Chem.* **96**, 6517–6522 (2024).

53. Wang, G. *et al.* Spatial organization of the transcriptome in individual neurons.

2020.12.07.414060 Preprint at https://doi.org/10.1101/2020.12.07.414060 (2020).

54. Moffitt, J. R. *et al.* High-performance multiplexed fluorescence in situ hybridization in

culture and tissue with matrix imprinting and clearing. *Proceedings of the National*

Academy of Sciences **113**, 14456–14461 (2016).

55. Wang, G., Moffitt, J. R. & Zhuang, X. Multiplexed imaging of high-density libraries of

RNAs with MERFISH and expansion microscopy. *Sci Rep* **8**, 4847 (2018).

56. Khaledian, B., Thibes, L. & Shimono, Y. Adipocyte regulation of cancer stem cells.

Cancer Sci **114**, 4134–4144 (2023).

57. Hu, X. *et al*. Decorin-mediated suppression of tumorigenesis, invasion, and metastasis

in inflammatory breast cancer. *Commun Biol* **4**, 72 (2021).

58. Cai, R. *et al*. Primary breast tumor induced extracellular matrix remodeling in

premetastatic lungs. *Sci Rep* **13**, 18566 (2023).

59. de Freitas Oliveira-Tore, C. *et al*. Non-canonical extracellular complement pathways

and the complosome paradigm in cancer: a scoping review. *Front. Immunol.* **16**,

(2025).

60. Maynard, K. R. *et al*. Transcriptome-scale spatial gene expression in the human

dorsolateral prefrontal cortex. *Nat Neurosci* **24**, 425–436 (2021).

61. Johnson, Z. V. *et al*. Cellular profiling of a recently-evolved social behavior in cichlid

fishes. *Nat Commun* **14**, 4891 (2023).

62. Lake, B. B. *et al*. A comparative strategy for single-nucleus and single-cell

transcriptomes confirms accuracy in predicted cell-type expression from nuclear RNA.

Sci Rep **7**, 6031 (2017).

63. Habib, N. *et al*. Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nat*

Methods **14**, 955–958 (2017).

64. DeFelipe, J. *et al*. Cortical white matter: beyond the pale remarks, main conclusions

and discussion. *Front. Neuroanat.* **4**, 1692 (2010).

65. Ritchie, M. E. *et al.* limma powers differential expression analyses for RNA-sequencing

and microarray studies. *Nucleic Acids Res* **43**, e47 (2015).

66. Sakers, K. *et al.* Astrocytes locally translate transcripts in their peripheral processes.

Proceedings of the National Academy of Sciences **114,** E3830–E3838 (2017).

67. Jiang, T. *et al.* The lncRNA MALAT1/miR-30/Spastin Axis Regulates Hippocampal

Neurite Outgrowth. *Front. Cell. Neurosci.* **14,** (2020).

68. Overly, C. C., Rieff, H. I. & Hollenbeck, P. J. Organelle motility and metabolism in axons

vs dendrites of cultured hippocampal neurons. *J Cell Sci* **109**, 971–980 (1996).

69. Harris, J. J., Jolivet, R. & Attwell, D. Synaptic Energy Use and Supply. *Neuron* **75**, 762–

777 (2012).

70. Than-Trong, E. & Bally-Cuif, L. Radial glia and neural progenitors in the adult zebrafish

central nervous system. *Glia* **63**, 1406–1428 (2015).

71. Miranda-Negrón, Y. & García-Arrarás, J. E. Radial glia and radial glia-like cells: Their role

in neurogenesis and regeneration. *Front. Neurosci.* **16,** (2022).

72. Zhu, J., Sun, S. & Zhou, X. SPARK-X: non-parametric modeling enables scalable and

robust detection of spatial expression patterns for large spatial transcriptomic studies.

Genome Biology **22**, 184 (2021).

73. Abramsson, A. *et al.* The zebrafish amyloid precursor protein-b is required for motor

neuron guidance and synapse formation. *Developmental Biology* **381**, 377–388 (2013).

74. Newman, M., Ebrahimie, E. & Lardelli, M. Using the zebrafish model for Alzheimer's

disease research. *Front. Genet.* **5,** (2014).

75. Haenisch, C. *et al.* The neuronal growth and regeneration associated Cntn1

(F3/F11/Contactin) gene is duplicated in fish: expression during development and

retinal axon regeneration. *Molecular and Cellular Neuroscience* **28**, 361–374 (2005).

76. Eiraku, M. *et al.* DNER acts as a neuron-specific Notch ligand during Bergmann glial

development. *Nat Neurosci* **8**, 873–880 (2005).

77. Díaz-Ramos, À., Roig-Borrellas, A., García-Melero, A. & López-Alemany, R. α-Enolase, a

Multifunctional Protein: Its Role on Pathophysiological Situations. *BioMed Research*

International **2012**, 156795 (2012).

78. Buxbaum, A. R., Haimovich, G. & Singer, R. H. In the right place at the right time:

visualizing and understanding mRNA localization. *Nat Rev Mol Cell Biol* **16**, 95–109

(2015).

79. Zhou, Z., Li, X., Wright, J., Candès, E. & Ma, Y. Stable Principal Component Pursuit. in

2010 IEEE International Symposium on Information Theory 1518–1522 (2010).

doi:10.1109/ISIT.2010.5513535.

80. Mateos, G. & Giannakis, G. B. Robust PCA as Bilinear Decomposition With Outlier-

Sparsity Regularization. *Trans. Sig. Proc.* **60**, 5176–5190 (2012).

81. Manabe, R. *et al.* Transcriptome-based systematic identification of extracellular matrix

proteins. *Proceedings of the National Academy of Sciences* **105**, 12849–12854 (2008).

82. Giannakis, G. B., Mateos, G., Farahmand, S., Kekatos, V. & Zhu, H. USPACOR:

Universal sparsity-controlling outlier rejection. in *2011 IEEE International Conference*

on Acoustics, Speech and Signal Processing (ICASSP) 1952–1955 (2011).

doi:10.1109/ICASSP.2011.5946891.

83. Huang, P., Cai, M., Lu, X., McKennan, C. & Wang, J. Accurate estimation of rare cell-

type fractions from tissue omics data via hierarchical deconvolution. *The Annals of*

Applied Statistics **18**, 1178–1194 (2024).

84. Zhong, Y. & Liu, Z. Gene expression deconvolution in linear space. *Nat Methods* **9**, 8–9

(2012).

85. Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical

variability in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562–578 (2018).

86. Townes, F. W., Hicks, S. C., Aryee, M. J. & Irizarry, R. A. Feature selection and

dimension reduction for single-cell RNA-Seq based on a multinomial model. *Genome*

Biology **20**, 295 (2019).

87. Mu, X. *et al.* Single-nucleus and spatial transcriptomics identify brain landscape of

gene regulatory networks associated with behavioral maturation in honeybees. *Nat*

Commun **16**, 3343 (2025).

88. Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat*

Biotechnol **36**, 411–420 (2018).

977