**RESCUE: recovery of idiosyncratic expression patterns in spatial transcriptomics**

Young Joo Lee[1], Seokjin Yeo[2], Alex W. Schrader[3], Ian M. Traniello[4], Amy Cash Ahmed[5],

Gene E. Robinson[5,6,7], Hee-Sun Han[2,3,5,6*,] Sihai Dave Zhao[1,5*]

[1]Department of Statistics, University of Illinois Urbana-Champaign

[2]Department of Bioengineering, University of Illinois Urbana-Champaign

[3]Department of Chemistry, University of Illinois Urbana-Champaign

[4]Lewis-Sigler Institute for Integrative Genomics, Princeton University

[5]Carl R. Woese Institute of Genomic Biology, University of Illinois Urbana-Champaign

[6]Neuroscience Program, University of Illinois Urbana-Champaign

[7]Department of Entomology, University of Illinois Urbana-Champaign

* Corresponding authors: sdzhao@illinois.edu (S.D.Z), hshan@illinois.edu (H.-S.H)

**ABSTRACT**

Spatial transcriptomics (ST) enables gene expression profiling while preserving the

spatial architecture of intact tissue. Analyzing ST data often proceeds by first extracting

cell-level information, typically through cell segmentation or cell-type deconvolution.

However, they can fail to capture important expression patterns, for example from fragile

or underrepresented cell types, subcellular structures like neurites, and extracellular

expression. These omissions can result in biased analyses and incorrect or incomplete

biological interpretations. We describe a new computational method, RESCUE, that can

recover the idiosyncratic spatial expression patterns missed by existing ST analysis

methods and enable robust inference even when reference data are incomplete. We

validate RESCUE using MERFISH data from the honey bee brain and apply it to multiple

ST datasets to demonstrate how it can reveal novel insights into complex tissue biology.

**INTRODUCTION**

Profiling cellular expression within its spatial context has become possible with the recent

development of spatial transcriptomics (ST) technologies, which measures gene

expression in intact tissue sections[1,2]. Analyzing ST data typically proceeds by extracting

cell-level information from the spatial expression data, as cells are natural units of

analysis. With high-resolution platforms such as MERFISH[3], 10x Visium HD[4], or 10x

Xenium[5], this is usually done using cell segmentation[6–8]. With low-resolution platforms

like 10x Visium[9] and Slide-seq[10,11], where each spatial location can contain multiple cells,

it is common to deconvolve the observed expression into separate contributions from

different cell types, based on reference single-cell/nuclei RNA-seq (sc/snRNA-seq)

data[12–16].

However, a major problem with these existing ST analysis methods is that they

can fail to capture important expression patterns. First, they may omit entire cells or cell

types. Cell segmentation algorithms for high-resolution data have difficulty with cells with

complex morphologies, such as fibroblasts, or cells lacking nuclei, like erythrocytes[17,18].

Deconvolution methods for low-resolution data cannot account for cell types that are

absent from the reference sc/snRNA-seq data because they are too fragile to survive

tissue dissociation[19–21]. Second, existing ST approaches can overlook expression

patterns unique to cellular substructures. Segmentation methods simply drop transcripts

located in substructures that are difficult to segment, such as the axons and dendrites of

neurons. These fragile structures are also typically destroyed during tissue dissociation

and therefore are not represented in reference single-cell data used in deconvolution.

Indeed, it has been estimated that over 40% of the total transcriptome of an adult mouse

brain may be missed by sc/snRNA-seq[22]. Similarly, reference snRNA-seq data, by

definition, ignores cytoplasmic expression. Finally, existing methods do not capture

extracellular tissue expression patterns, which cannot be studied by segmentation or

deconvolution.

These difficult-to-capture expression patterns can be critically important. First, they

can bias analyses that do not take them into account. An example with dramatic

implications is that deconvolution estimates of cell-type proportions can be invalid if the

reference is missing any cell types[23], as we illustrate later in this work. Second, there may

be important biological processes that are unique to the overlooked expression patterns.

For example, important missed cell types include adipocytes[5], which play crucial roles in

metabolic disease and cancer[24], and neutrophils[25], which are core components of the

immune system. Important missed substructural expression includes axonal or dendritic

transcripts that are central to synaptic development and plasticity[26–30] and diseases such

as amyotrophic lateral sclerosis[31,32], as well as cytoplasmic transcripts that regulate key

processes such as metabolism, signaling, and stress responses[33–35] and are involved in

Alzheimer's disease[36,37]. Important missed extracellular expression patterns include

extracellular RNAs that mediate intercellular communication in cancer[38–40].

We describe a new computational method, RESCUE, that can recover the spatial

expression patterns missed by single-cell inference on ST data. Our approach is based

on making a fundamental correction to the class of statistical models currently used for

integrating ST data and reference data, such as sc/snRNA or cell segmentation results.

Briefly, our method partitions observed gene expression into one component explainable

by the reference data, which we refer to as "canonical" expression, and another

component that is not, which we refer to as "idiosyncratic" expression. Our purified

canonical ST expression component can be used to correct existing cell-type

deconvolution methods. Our recovered idiosyncratic ST expression component gives us

the new ability to specifically target biology that is distinct from our reference data, as

discussed above. The idiosyncratic expression recovery step in RESCUE is analogous

to negative selection in experimental biology[41–44], where we identify and remove

canonical expression. We demonstrate that RESCUE accurately recovers idiosyncratic

expression, enables robust deconvolution of ST data despite incomplete reference data,

and reveals biologically meaningful patterns overlooked by existing methods. While we

focus here on ST data, we note that RESCUE can also be directly applied to non-spatial

modalities, such as bulk RNA-seq data.

**RESULTS**

**Overview of RESCUE**

RESCUE is predicated on a new model that we propose for integrating ST and reference

single-cell data, and can be applied to diverse ST platforms, including MERFISH and

Visium, across different organisms. Most existing models[12–16] posit that ST expression at

each location is composed of a linear combination of expression patterns present in the

reference, where the coefficients of this linear combination are interpreted as the

proportions of each reference cell type in that location. However, as described in the

introduction, there are numerous sources of ST expression that are left out of this model;

we will refer to these as idiosyncratic expression patterns, because they are different from

the canonical expression patterns reflected in the reference.

To address this issue, we propose a corrected model (**Fig. 1**) that allows the ST

profile to have both canonical and idiosyncratic expression components. While it appears

minor, this change has important implications. Under this model, results from the statistics

literatures[45–47] imply that under certain conditions it is possible to simultaneously recover

idiosyncratic expression and estimate the linear combination of reference patterns that

constitute canonical expression. The conditions are that only relatively few genes show

non-zero, or "sparse" expression, in the idiosyncratic component, which is reasonable

given the nature of idiosyncratic expression and the read depth/detection efficiency of ST

technologies.

Leveraging this insight, we developed RESCUE, an algorithm that simultaneously

deconvolves ST data while rescuing spatial idiosyncratic expression. RESCUE solves a

regularized robust multivariate regression problem and can be applied to both low- and

high-resolution ST data. Unlike existing deconvolution methods[12–16], it is robust to

incomplete references, such as snRNA-seq data which lack cytoplasmic expression

profiles. The rescued idiosyncratic expression can be analyzed using standard

approaches, such as clustering, differentially expressed gene (DEG) analysis, and

spatially varying gene (SVG) analysis, to discover new biology hidden from existing ST

analyses. We illustrate this in a broad range of examples below. We note that the

interpretation of idiosyncratic expression depends on both the tissue context and the

choice of reference dataset.

**Simulations**

To evaluate the performance of RESCUE, we conducted a simulation study on synthetic

data that mimics real experimental ST data with varying degrees of idiosyncratic

expression. As no existing methods are specifically designed to recover idiosyncratic

expression, we adapted four widely used deconvolution approaches: non-negative least

squares (NNLS), weighted NNLS (WNNLS), RCTD[13], and CARD[14], to recover the

idiosyncratic component. For each method, we first estimated cell-type proportions using

its standard deconvolution procedure and then recovered the idiosyncratic expression by

subtracting the estimated cellular contributions from the observed ST expression.

We simulated ST data using real scRNA-seq data from the honey bee brain[48] by

introducing synthetic idiosyncratic expression of a small number of genes (Methods).

Honey bee brains provide a unique benefit for simulating idiosyncratic expression due to

the existence of the neuropil region, where axons and dendrites are spatially segregated

from cell bodies. The molecular expression in these regions is largely absent in single-

cell sequencing data due to the structural fragility of neurites. During the tissue

dissociation step, a necessary step in single-cell sequencing, neurites are mostly sheared

off and lost. In contrast, ST measurements performed on intact tissue sections preserve

these structures, thus the molecular expression in neuropils as well. Our in-house

collected MERFISH data shows that neuropil regions have less than 10% of genes

expressed on average, confirming a highly sparse idiosyncratic expression component.

We simulated both highly sparse and moderately sparse idiosyncratic expression by

varying the idiosyncratic sparsity level (e.g., 0.95 for "highly sparse" and 0.80 for

"moderately sparse") and idiosyncratic expression proportion (ranging from 0 to 0.8).

We first compared the deconvolution performance with existing methods. **Fig. 2A**

shows the accuracy of cell-type proportion estimates across methods, measured by

average Lin's concordance correlation coefficient[49] (CCC) (Methods). Expectedly, the

performance of standard deconvolution methods deteriorates sharply as idiosyncratic

expression level increases. On the other hand, RESCUE maintains high accuracy across

all conditions, demonstrating robustness to both high idiosyncratic content and reduced

sparsity. Our findings imply that idiosyncratic expression, if not properly accounted for,

can introduce systematic biases in cell-type proportion estimates.

We also evaluated the quality of idiosyncratic transcriptome recovery (**Fig. 2B**),

using relative Frobenius error between the true and estimated idiosyncratic expression

matrices (Methods). Overall, RESCUE achieves the lowest recovery error, even under

high idiosyncratic expression and low sparsity, while other methods deviate substantially

from the ground truth, especially when idiosyncratic signals are dense. Our findings show

that idiosyncratic expression, if not properly accounted for, can obscure cell-type

deconvolution and other downstream analyses, which we will illustrate below. This has

major implications for deconvolution with incomplete reference data and analyses of

complex tissue structures with significant idiosyncratic transcription. RESCUE accurately

recovers the hidden transcriptome and cell-type composition with minimal distortion.

**RESCUE accurately recovers the idiosyncratic transcriptome in real tissues**

Using a custom MERFISH platform[50], we generated single molecule-resolution ST data

from coronal sections of a honey bee brain, targeting 130 genes relevant to aggression

(**Supplementary Table S1**)[51]. Transcripts in axons and dendrites, which are collectively

termed as neurites, have been difficult to profile due to segmentation challenges. Cell

segmentation methods, such as Cellpose[6] and Baysor[7], can only identify the boundaries

of neuronal bodies, also called somata. Individual axons and dendrites could be stained

in 2D cell cultures[52], but in native tissues they likely will not lie in one imaging plane. Thus,

traditionally, neurite expression has been either ignored or incorrectly merged into nearby

cytosolic expression.

We first experimentally validated that RESCUE can accurately recover gene

expression in the neuropil region of the honey bee brain. The honey bee brain serves as

an ideal test case due to its spatially segregated somata and neuropil regions, allowing

precise profiling of neurite expression without interference from somatic expression. Cell

soma were segmented using Cellpose[6]. Remarkably, only 73.5% of transcripts were

localized within cells, indicating that the idiosyncratic transcriptome accounts for 26.5%

of the observed ST expression. These signals, which would have been ignored by

standard cell segmentation approaches, highlight the importance of RESCUE in capturing

overlooked transcriptomic information.

Using honey bee brain scRNA-seq data[48] as a reference, we applied each method

to our MERFISH data after dividing the brain into grids of size $20 \times 20 \ \mu m^2$. We first used

RESCUE and the other methods to estimate the proportion of canonical expression

relative to the total expression in each of our grids. Leveraging the spatial segregation of

the honey bee brain, we then obtained a ground truth proportion of transcripts in each

grid that are localized inside cell bodies via cell segmentation. RESCUE's estimates

closely matched those from cell segmentation (**Fig. 3A**) and were more accurate than

estimates from other methods based on two metrics: Lin's CCC and RMSE between the

segmentation-based and estimated proportions (**Fig. 3B**, Methods). Note that, in this

example, RESCUE solely relies on reference scRNA-seq data and does not rely on cell

segmentation.

Finally, we evaluated how well each method recovered the gene expression

specifically within the somatic regions (**Fig. 3C**). We calculated the Lin's CCC between

the true and estimated gene expression within the grids located in somata. Again,

RESCUE significantly outperformed other methods, accurately recovering canonical as

well as idiosyncratic expression. Collectively, these results validate that RESCUE is

highly effective in partitioning the full transcriptome into its somatic and neuropil

components, capturing valuable expression signals that would otherwise be lost using

conventional approaches.

**RESCUE enables accurate deconvolution of ST data with incomplete reference**

Here, we illustrate how RESCUE's estimated canonical expression component can be

used to achieve accurate deconvolution of ST data when key cell types are missing from

the reference, a challenge for all existing methods that has been noted in a recent study[23].

We analyzed paired single-cell FFPE-seq (scFFPE-seq) and 10x Visium ST data from

human breast cancer tissue[5]. In this ST dataset, adipocytes are manually annotated in

approximately 10% of spatial spots (**Fig. 4A**). However, adipocytes are absent from the

scFFPE-seq reference, likely due to their high lipid content interfering with fixation and

processing. These cells are prone to floating, rupturing, or sticking to plastic surfaces

during tissue dissociation, which hinders their recovery in single-cell sequencing

protocols[5,17]. As a result, reference-based methods that assume the observed expression

is fully explained by the available reference cell types tend to misattribute adipocyte-

derived transcripts to other cell types.

In adipocyte-rich regions, all four existing methods (NNLS, WNNLS, RCTD, and

CARD) fail to account for the missing adipocyte transcriptome and instead misattribute

the signal to other reference cell types, such as stromal and endothelial cells (**Fig. 4C**).

This systematic misassignment likely arises from transcriptional similarity, as shown by

the strong correlation between adipocyte and stromal/endothelia profiles in pseudo-bulk

measurements from Visium and scFFPE-seq (**Fig. 4B**). On the other hand, RESCUE

accommodates idiosyncratic expression and avoids forcing unassigned signal into

existing reference cell types, which results in more accurate and biologically plausible

estimates. As shown in **Fig. 4D**, RESCUE's estimated proportions closely align with the

scFFPE-seq composition, while other methods exhibit significant distortions. Together,

these results demonstrate that RESCUE improves deconvolution accuracy in settings

with incomplete references by preventing the systematic misassignment of idiosyncratic

transcriptomes to incorrect cell types.

**RESCUE reveals idiosyncratic biological processes in diverse tissues**

In many neuroscience experiments, cell-level information is often characterized using snRNA-seq[53,54], which does not capture cytoplasmic or neurite expression. Here, we illustrate how RESCUE can uncover unique biological processes within the idiosyncratic expression of the human dorsolateral prefrontal cortex (DLPFC) and cichlid fish telencephalon, using snRNA-seq as a reference.

**Human DLPFC.** Maynard et al.[55] profiled the DLPFC of neurotypical adult donors using paired snRNA-seq and 10x Visium ST. We applied RESCUE to eight samples in which all six cortical layers and the white matter (WM) were annotated, to recover idiosyncratic expression patterns not represented in the snRNA-seq reference (**Fig 5A**). As shown in **Fig. 5B**, RESCUE reveals that Layer 1 and the WM contain substantially higher levels of idiosyncratic expression compared to other cortical layers. This is consistent with the fact that Layer 1 has sparser cell densities and contains neuropil-rich regions[55], and that WM contains a substantial amount of axons and myelin[56], whose transcriptomic content is mostly cytoplasmic and thus is absent from the reference snRNA-seq data.

RESCUE enables us to study biology specific to the idiosyncratic transcriptome of Layer 1. We first identified DEGs by comparing Layer 1 to all other layers using either the raw observed ST expression or the idiosyncratic expression recovered by RESCUE. These two analyses gave substantially different results (**Fig. 5C**). For example, the top five genes most enriched in Layer 1 based on idiosyncratic expression had either fold-

changes in the opposite direction (*MT-ND1*, *MT-ND2*, and *MT-CO1*) or negligible fold-

changes (*MT3* and *CST3*) in the differential expression analysis using raw data.

To explore the underlying biological processes, we performed gene ontology (GO)

enrichment analysis separately for DEGs derived from the idiosyncratic expression and

from the raw ST data (**Fig. 5D**). Idiosyncratic GO terms include mitochondrial processes:

"oxidative phosphorylation", "energy derivation by oxidation of organic compounds", and

"generation of precursor metabolites and energy". In contrast, no mitochondria-related

pathways reached statistical significance in the analysis based on observed ST

expression; only the broad term "response to stress" passed the FDR < 0.05 threshold.

These results align with the elevated metabolic demands of a neurite-enriched region like

Layer 1, which is densely populated with synapses and reliant on local energy production

to support axonal and dendritic activity[57,58]. Collectively, these findings demonstrate that

isolating idiosyncratic expression can uncover previously obscured yet functionally

important gene expression programs in Layer 1 of the human DLPFC.

**Cichlid Fish Telencephalon.** Johnson et al.[59] collected paired snRNA-seq and Visium

ST data from the dorsal telencephalon of individual cichlid fishes exhibiting distinct social

behaviors. As shown in **Fig. 6A**, anatomical annotations delineate multiple subdivisions

of the dorsal pallium, structured around radial glial (RG) zones that extend long

projections from the ventricular zone into superficial pallial regions[60,61]. These RG

projections span nearly the entire depth of the pallium and may play a role in shaping

spatial gene expression profiles across distinct layers and compartments, but they do not

survive tissue dissociation and thus their transcriptomes are not captured by snRNA-seq.

We applied RESCUE to this dataset to recover idiosyncratic transcriptomic signals

using the snRNA-seq data as a reference. To assess spatial structure in the rescued

transcriptome, we performed SVG detection using SPARK-X[62] on both the observed ST

expression and the idiosyncratic expression. As shown in **Fig. 6B**, a substantial number

of SVGs are uniquely identified in the idiosyncratic transcriptome. In **Fig. 6C**, we show

the idiosyncratic expression maps of those SVGs that were undetectable in the observed

ST expression yet exhibited clear spatial structure once nuclear signals were removed.

Among these, *app*, *cntn1*, *dner*, and *eno1* mark distinct transcriptional programs

that may operate along RG projections. These genes highlight transcriptional signals that

are likely to reflect localized molecular activities along RG scaffolds: *app* encodes a

transmembrane protein that promotes synapse formation and plasticity[63,64], *cntn1*

(Contactin-1) is a GPI-anchored adhesion molecule essential for axon guidance and

neurite outgrowth[65], *dner* mediates neuron-glia interactions that are critical for astrocyte

differentiation[66], and *eno1* ($\alpha$-enolase) promotes neuritogenesis and contributes to

structural remodeling in the developing nervous system[67]. Indeed, Johnson et al.[59] found

that bower-building behavior in cichlid fishes may be related to neuronal rebalancing

driven by a subpopulation of RG cells and their processes, highlighting the importance of

studying transcriptional processes in RG projections, which RESCUE enables.

**DISCUSSION**

RESCUE is the first method designed to acknowledge and recover the idiosyncratic

transcriptome, which encompasses gene expression patterns not captured by cell

segmentation or deconvolution approaches in ST data analysis. These idiosyncratic

signals, though typically overlooked, can carry substantial functional and biological

significance. While the field has developed numerous transcriptomic profiling methods,

such as scRNA-seq for isolating gene expression at the single-cell level, Ribo-seq for

ribosome-associated transcripts, and CLIP-seq for binding-protein interactions, each of

these technologies inevitably captures only a fraction of the total transcriptome. RESCUE

provides a systematic framework to quantify and analyze what is missing: the molecular

signals outside the scope of reference datasets.

By formulating the problem as a robust sparse recovery framework, RESCUE

decomposes observed ST data into a reference-aligned canonical component and a

biologically meaningful idiosyncratic component. RESCUE focuses on characterizing

what is not explained by reference profiles and introduces a broader conceptual

innovation: computational negative selection. Rather than focusing solely on what to

include (as in positive selection), we define a reference component to subtract out and

focus on what remains. In this work, we used scRNA-seq and snRNA-seq data as

reference profiles and focused primarily on brain tissue, but the concept is general and

can be applied to a wide range of other settings, for example, muscle tissue, where

multinucleated cells make segmentation challenging and single-cell reference data

difficult to interpret. The choice of reference therefore requires careful design and quality

control.

One striking finding emerging from our brain tissue analyses is the substantial

amount of expression (26.5% of total expression) found in the idiosyncratic transcriptome

of the honey bee brain MERFISH data. While localization of mRNAs to neurites is a well-

studied phenomenon[26–30], our results, along with the estimates for the adult mouse brain

from Ament and Poulpoulos[22], additionally suggest that these local expression patterns

may be different from those found in cell soma. RESCUE enables a reexamination of

existing datasets to more carefully study this apparently divergent expression.

There are several potential directions for future work. First, in cases where no

suitable reference is available, extensions of RESCUE to reference-free or unsupervised

frameworks may be possible. For example, methods based on robust principal

component analysis[68,69] could be adapted to extract idiosyncratic structure without

predefined reference matrices. Second, applying RESCUE to high-resolution ST data

requires creating a predefined grid, and further research is required to develop a

systematic method to determine the optimal grid size. Despite these remaining

challenges, we have already shown that RESCUE can still provide meaningful analyses

of spatial idiosyncratic expression.

Finally, RESCUE reminds us that the biology we wish to study and the

measurements we are able to make are not always one and the same. For instance, we

typically accept single-cell reference data as truly representing the expression of the

entire single cell, but in many cases it does not, as we have illustrated above. Any

measurement technology has inherent biases. As a result, recognizing, acknowledging

and properly interpreting what is missed, in other words, the "idiosyncratic expression", is

important for a comprehensive understanding of a system.

## METHODS

### RESCUE model

We assume that our ST data profiles $G$ genes in $J$ grids across the tissue section. In low-resolution ST technologies, these grids are predefined as multicellular spots, whereas in higher-resolution ST technologies, these can be constructed at any desired size, which allows inference at different biological scales. To introduce the statistical model behind RESCUE, we begin by reviewing existing linear deconvolution methods[12–16]. These methods decompose the ST expression matrix $Y \in \mathbb{R}_{\geq 0}^{G \times J}$ of $G$ genes in $J$ grids as

$$Y = X\alpha + E, \tag{1}$$

where $X \in \mathbb{R}_{\geq 0}^{G \times K}$ is a matrix of $K$ cell-type expression reference profiles, $\alpha \in \mathbb{R}_{\geq 0}^{K \times J}$ is a matrix of inferred cell-type proportions within each grid, and $E \in \mathbb{R}^{G \times J}$ is assumed to be a mean-zero noise matrix. The issue is that (1) implies that expectation of ST expression is $\mathbb{E}(Y) = X\alpha$, which ignores idiosyncratic transcriptome.

Instead, we further decompose the error $E$, motivated by the robust regression framework[45–47]. Specifically, we consider the multivariate mean-shift model

$$Y = X\alpha + R + \mathcal{E}, \tag{2}$$

where $X\alpha$ represents the expected expression values of each gene in each grid due to the presence of cells whose types are represented in the reference sc/snRNA-seq data, $R \in \mathbb{R}_{\geq 0}^{G \times J}$ is an idiosyncratic term that allows for outliers that cannot be captured by the reference data, and $\mathcal{E} \in \mathbb{R}^{G \times J}$ is a noise matrix that allows for residual variability between the ST sample and the reference data. Our key assumption is that $R$ is sparse. This is necessary for identifiability and is also reasonable because we expect relatively few

genes to exhibit idiosyncratic transcriptome within each grid. We also assume that both

$\alpha$ and $R$ are non-negative, for biological interpretation.

**Estimation algorithm**

RESCUE aims to simultaneously recover the sparse non-negative matrix $R$ and non-

negative cell-type proportions $\alpha$ in our model (2). Specifically, we consider the following

optimization problem:

$$\min_{\alpha, R \geq 0} \|W \odot R\|_1 \text{ subject to } \|Y - R - X\alpha\|_F^2 \leq \lambda,$$

where $W$ is a weight matrix, $\odot$ denotes the Hadamard product, $\|\cdot\|_F$ denotes the

Frobenius norm, and $\lambda > 0$ is a tuning parameter. We set each entry of the weight matrix

as $W_{gj} = 1/\sqrt{Var(R_{gj})}$ to account for the fact that different genes have different variances

across different locations. The $\ell_1$ objective function enforces sparsity of our estimated $R$,

while the constraint ensures that the residual variability $\mathcal{E}$ is not too large, in other words,

that the ST data do not deviate too much from the reference sc/snRNA-seq data.

In practice, we solve the Lagrangian form of this problem:

$$\min_{\alpha, R \geq 0} f(\alpha, R) = \|W \odot R\|_1 + \frac{\mu}{2}\|Y - R - X\alpha\|_F^2, \tag{3}$$

where $\mu > 0$ is a tuning parameter. We apply an efficient alternating minimization to solve

two convex sub-problems in each iteration. Specifically, our proposed algorithm is a

multivariate extension of the soft-IPOD[45], incorporating non-negative constraints and the

weight matrix:

$$\hat{\alpha} = \operatorname*{argmin}_{\alpha \geq 0}\|Y - R - X\alpha\|_F^2, \tag{4}$$

$$\hat{R} = \operatorname*{argmin}_{R \geq 0}\|W \odot R\|_1 + + \frac{\mu}{2}\|Y - R - X\alpha\|_F^2 = S_+\left(Y - X\alpha, \frac{W}{\mu}\right), \tag{5}$$

where $S_+(x, \tau) = \max(x - \tau, 0)$ denotes the entry-wise non-negative soft-thresholding

function. In the presence of multiple outliers, it is important to address masking and

swamping effects. Our proposed weight plays a similar role as the adaptive weight used

in Kong et al.[47], but it can be iteratively estimated within the algorithm, not requiring a

robust initial estimator.

To estimate $W_{gj}$, we iteratively estimate the variance of $R_{gj}$. In ST deconvolution

models, it is common to model the observed gene expression count $\tilde{Y}_{gj}$ under a Poisson

assumption[13,16]:

$$\tilde{Y}_{gj} | \lambda_{gj} \sim \text{Poisson}(N_j \lambda_{gj})$$

where $N_j = \sum_g \tilde{Y}_{gj}$ represents the total transcript count, or number of UMIs for grid $j$, and

$\lambda_{gj}$ denotes the expected expression rate of gene $g$ at grid $j$. Rather than directly

modeling $\tilde{Y}_{gj}$, we assume that it can be decomposed as $\tilde{Y}_{gj} = \tilde{L}_{gj} + \tilde{R}_{gj}$, where

$$\tilde{L}_{gj} | \lambda_{gj}^L \sim \text{Poisson}(N_j^L \lambda_{gj}^L) \quad \text{and} \quad \tilde{R}_{gj} | \lambda_{gj}^R \sim \text{Poisson}(N_j^R \lambda_{gj}^R).$$

Here, $N_j^L = \sum_g \tilde{L}_{gj}$ and $N_j^R = \sum_g \tilde{R}_{gj}$ represent the total transcript counts of the canonical

and idiosyncratic expression for grid $j$, respectively, and $(\lambda_{gj}^L, \lambda_{gj}^R)$ denote the expected

expression rates of these components. We can write $N_j^L = \pi_j N_j$ and $N_j^R = (1 - \pi_j) N_j$,

where $\pi_j \in [0, 1]$ denotes the proportion of canonical expression for grid $j$.

When using counts per million (CPM) normalization, i.e., $Y_{gj} = \tilde{Y}_{gj} / N_j \times 10^6$, we

can estimate the variance of idiosyncratic component as

$$\text{Var}(R_{gj}) = \text{Var}\left(\frac{\tilde{R}_{gj}}{N_j} \times 10^6\right) = \left(\frac{10^6}{N_j}\right)^2 \times \text{Var}(\tilde{R}_{gj}) \approx \frac{10^6}{N_j} \times \hat{R}_{gj}, \qquad (6)$$

where the approximation follows from $\widehat{\mathrm{Var}}(\tilde{R}_{gj}) = N_j^R \lambda_{gj}^R$ and $\tilde{R}_{gj} = R_{gj}/10^6 \times N_j$. Here,

the term $N_j^R \lambda_{gj}^R$ can be iteratively updated using the plug-in estimator $\tilde{R}_{gj} \approx N_j^R \lambda_{gj}^R$. To

avoid division by zero, we set $W_{gj} = 1/\max\left(\sqrt{\frac{10^6}{N_j}} \times R_{gj}, \delta\right)$, where $\delta$ is a small positive

constant.

**Tuning parameter selection**

RESCUE aims to recover idiosyncratic transcriptome by balancing the sparsity of $R$,

representing the number of outliers rejected, against the level of residual variability $\mathcal{E}$.

Following the approaches in robust regression and robust PCA[45,68], we set $\mu = GJ/C\|Y\|_1$

and instead focus on finding the optimal $C$. However, selecting the optimal tuning

parameter can be challenging in the presence of outliers[69,70].

To address this, we adopt a surrogate matching approach[71]. The idea is to choose

the optimal tuning parameter $C$ using a surrogate ST dataset with "ground truth" canonical

and idiosyncratic transcriptomes. To construct this surrogate ST dataset, we apply NNLS

to the observed ST data to obtain rough estimates for $\alpha$ and $R$:

$$\hat{R}_{NNLS} = (Y - X\hat{\alpha}_{NNLS})_+, \quad \text{where} \quad \hat{\alpha}_{NNLS} = \underset{\alpha \geq 0}{\mathrm{argmin}}\|Y - X\alpha\|_F^2, \tag{7}$$

where $(x)_+ = \max(x, 0)$ denotes the zero-thresholding function. We then simulate a

surrogate ST data $\tilde{Y}_{\mathrm{sim}}$ from a Poisson distribution with a rescaled expected expression

rate, preserving the overall signal-to-noise characteristics observed in the real data.

RESCUE is applied to $\tilde{Y}_{\mathrm{sim}}$ across a range of $\mu$ values, and the optimal $C$ is selected as

the value that minimizes the recovery error of the canonical and idiosyncratic components

in the surrogate dataset.

**Platform effect normalization**

To mitigate cross-platform effects between ST data and sc/snRNA-seq reference data, we apply platform effect normalization. This step is essential because differences in experimental platforms can introduce systematic shifts in gene expression levels that confound downstream analysis. Without proper alignment of scale, these differences may lead to biased estimation of cell-type contributions or idiosyncratic signals. We adopt the normalization strategy from RCTD[13], which rescales each gene in the reference data to match the overall expression level of the corresponding gene in the ST data. Specifically, we first compute pseudo-bulk measurements for both ST and reference datasets and then apply a gene-wise log-scale correction to harmonize them.

**Data normalization and reference matrix**

To account for technical variability and the effect of library size, we adopt CPM normalization but fit RESCUE in the linear space. We avoid log-transformation for two main reasons. First, log-transformed input tends to underestimate true signals in gene expression deconvolution[72]. Second, log-transformation requires adding pseudo-counts, which can introduce significant artificial distortion between zeros and non-zeros[73,74]. Instead, we work in the linear scale, employing an iteratively reweighted algorithm to stabilize variance adaptively. This approach allows the model to adjust for varying data scales and minimizes the influence of outliers without artificially altering data distributions. The normalized reference matrix is then used to construct average cell-type expression.

**446**    **Data generating process for simulation studies**

**447**    For a total of $J = 2,500$ grids, we first generated the total UMI count for grid $j$ from a

**448**    negative binomial distribution $N_j \sim \text{NB}(\text{mean} = 10,000, \text{size} = 2)$. We generated the raw

**449**    count for each gene in each grid as $\tilde{Y}_{gj} = \tilde{L}_{gj} + \tilde{R}_{gj}$, where $\tilde{R}_{gj}$ denotes the idiosyncratic

**450**    expression and $\tilde{L}_{gj}$ represents the canonical expression that could have been captured

**451**    by sc/snRNA-seq reference data.

**452**    To generate $\tilde{L}_{gj}$, we used the real scRNA-seq data from honey bee brains[48] as the

**453**    source of canonical expression profiles. We selected the top $G = 500$ number of highly

**454**    variable genes and constructed the reference matrix $X$ using $K = 20$ number of cell types

**455**    annotated via Louvain clustering in Seurat[75]. Latent mixing proportions were drawn as

**456**    $\beta_{kj} \sim \text{Beta}(1,3)$ and normalized as $\alpha_{kj} = \beta_{kj}/\sum_k \beta_{kj}$, with at most 10 cell types per spot.

**457**    Defining $\lambda_{gj}^L = (X\alpha)_{gj}$, we generated the cell-type-driven canonical expression as

**458**
$$\tilde{L}_{gj} \sim \text{Multinomial}\left(\pi_j N_j, P_{gj} = \frac{\lambda_{gj}^L}{\sum_g \lambda_{gj}^L}\right).$$

**459**    To generate sparse idiosyncratic expression, we sampled

**460**
$$\tilde{R}_{gj} \sim \text{Multinomial}\left((1 - \pi_j)N_j, Q_{gj} = \frac{\lambda_{gj}^R}{\sum_g \lambda_{gj}^R}\right),$$

**461**    where $\lambda_{gj}^R \sim \text{Gamma}(1, 1/3)$ with probability $(1 - s)$ and $\lambda_{gj}^R = 0$ with probability $s$. Here,

**462**    we introduced a sparsity parameter $s$, which makes $(s \times 100)$ percentage of intensity

**463**    parameter $\lambda_{gj}^R$ zero-inflated. Under this setting, we generated $\tilde{L}_{gj}$ and $\tilde{R}_{gj}$ to satisfy and

**464**    $\sum_g(\tilde{L}_{gj} + \tilde{R}_{gj}) = N_j$. In our simulations, we considered $\pi_j \in \{0, 0.2, 0.4, 0.6, 0.8\}$ and $s \in$

**465**    $\{0.8, 0.85, 0.9, 0.95\}$ to control the proportion of canonical expression and the sparsity level

**466**    of idiosyncratic transcriptome, respectively.

**Evaluation metrics for simulation studies**

469 To evaluate the robust estimation of cell-type proportions, we measured Lin's CCC

470 between the true and estimated proportion of the $k$th cell type, defined as

$$471 \quad \text{CCC}(\alpha_k, \hat{\alpha}_k) = \frac{2\text{Cov}(\alpha_k, \hat{\alpha}_k)}{\sigma_{\alpha_k}^2 + \sigma_{\hat{\alpha}_k}^2 + \left(\mu_{\alpha_k} - \mu_{\hat{\alpha}_k}\right)^2},$$

where $\mu_{\alpha_k}$ and $\mu_{\hat{\alpha}_k}$ are the means of the two variables, and $\sigma_{\alpha_k}^2$ and $\sigma_{\hat{\alpha}_k}^2$ denote the

corresponding variances. We used Lin's CCC because it captures shifts in both location

and scale, making it robust to high leverage points[71]. The concordance increases as the

CCC value gets closer to 1. We took the average of Lin's CCC across all cell types to

compare the cell-type deconvolution performance.

To evaluate the accuracy of idiosyncratic transcriptome recovery, we measured

relative Frobenius error, defined as

$$479 \quad (\text{Relative Frobenius Error}) = \frac{\|R - \hat{R}\|_F}{\|R\|_F},$$

480 where $R$ and $\hat{R}$ denote the true and estimated idiosyncratic transcriptomes.

**Evaluation metrics for honey bee brain MERFISH data**

483 We measured Lin's CCC and the root mean squared error (RMSE) between the

484 segmentation-based (true) and estimated idiosyncratic proportions, defined as follows:

$$485 \quad \text{CCC}(\rho, \hat{\rho}) = \frac{2\text{Cov}(\rho, \hat{\rho})}{\sigma_{\rho}^2 + \sigma_{\hat{\rho}}^2 + \left(\mu_{\rho} - \mu_{\hat{\rho}}\right)^2} \quad \text{and} \quad \text{RMSE}(\rho, \hat{\rho}) = \sqrt{\frac{1}{J}\sum_{j=1}^{J}\left(\rho_j - \hat{\rho}_j\right)^2},$$

where $\rho_j = \sum_g \tilde{R}_{gj} / \sum_g \tilde{Y}_{gj}$ denotes the true idiosyncratic proportion in grid $j$, and $\hat{\rho}_j$

denotes its estimated value. Here, $\mu_\rho$ and $\mu_{\hat{\rho}}$ are the means of two variables, and $\sigma_\rho^2$ and

$\sigma_{\hat{\rho}}^2$ denote the corresponding variances.

**DATA AVAILABILITY**

Human breast cancer dataset[5]: 10x Visium and scFFPE-seq data are available via GEO

accession GSE243280. Human DLPFC dataset[55]: 10x Visium and snRNA-seq data are

available at https://github.com/LieberInstitute/spatialLIBD. Cichlid fish brain dataset[59]:

10x Visium and snRNA-seq data are available via GEO accession GSE217619. Honey

bee brain scRNA-seq data[48] is available at https://doi.org/10.6084/m9.figshare.16832518.

496
**CODE AVAILABILITY**

RESCUE is available as an R package at https://github.com/brunoyjlee/RESCUE.

**ACKNOWLEDGEMENTS**

We thank Dr. Tina Barbasch on valuable discussions on the analysis of fish brain data.

Funding: This work was supported by the National Institutes of Health (R21HG013180 to

Y.J.L., S.Y., H.-S.H. and S.D.Z., R35GM147420 to A.W.S and H.-S.H. and R01AT013189

to S.D.Z.).

## REFERENCES

1. Rao, A., Barkley, D., França, G. S. & Yanai, I. Exploring tissue architecture using spatial transcriptomics. *Nature* **596**, 211–220 (2021).

2. Moses, L. & Pachter, L. Museum of spatial transcriptomics. *Nat. Methods* **19**, 534–546 (2022).

3. Chen, K. H., Boettiger, A. N., Moffitt, J. R., Wang, S. & Zhuang, X. Spatially resolved, highly multiplexed RNA profiling in single cells. *Science* (2015) doi:10.1126/science.aaa6090.

4. Nagendran, M. *et al*. 1457 Visium HD enables spatially resolved, single-cell scale resolution mapping of FFPE human breast cancer tissue. *J. Immunother. Cancer* **11**, (2023).

5. Janesick, A. *et al*. High resolution mapping of the tumor microenvironment using integrated single-cell, spatial and in situ analysis. *Nat. Commun.* **14**, 8353 (2023).

6. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular segmentation. *Nat. Methods* **18**, 100–106 (2021).

7. Petukhov, V. *et al*. Cell segmentation in imaging-based spatial transcriptomics. *Nat. Biotechnol.* **40**, 345–354 (2022).

8. Polański, K. *et al*. Bin2cell reconstructs cells from high resolution Visium HD data. *Bioinformatics* **40**, btae546 (2024).

9. Ståhl, P. L. *et al*. Visualization and analysis of gene expression in tissue sections by spatial transcriptomics. *Science* **353**, 78–82 (2016).

10.    Rodriques, S. G. *et al.* Slide-seq: A scalable technology for measuring genome-wide

expression at high spatial resolution. *Science* **363**, 1463–1467 (2019).

11.    Stickels, R. R. *et al.* Highly sensitive spatial transcriptomics at near-cellular

resolution with Slide-seqV2. *Nat. Biotechnol.* **39**, 313–319 (2021).

12.    Elosua-Bayes, M., Nieto, P., Mereu, E., Gut, I. & Heyn, H. SPOTlight: seeded NMF

regression to deconvolute spatial transcriptomics spots with single-cell transcriptomes.

*Nucleic Acids Res.* **49**, e50 (2021).

13.    Cable, D. M. *et al.* Robust decomposition of cell type mixtures in spatial

transcriptomics. *Nat. Biotechnol.* **40**, 517–526 (2022).

14.    Ma, Y. & Zhou, X. Spatially informed cell-type deconvolution for spatial

transcriptomics. *Nat. Biotechnol.* **40**, 1349–1359 (2022).

15.    Zhou, Z., Zhong, Y., Zhang, Z. & Ren, X. Spatial transcriptomics deconvolution at

single-cell resolution using Redeconve. *Nat. Commun.* **14**, 7930 (2023).

16.    Liu, Z., Wu, D., Zhai, W. & Ma, L. SONAR enables cell type deconvolution with

spatially weighted Poisson-Gamma model for spatial transcriptomics. *Nat. Commun.*

**14**, 4727 (2023).

17.    Si, Y. *et al.* FICTURE: scalable segmentation-free analysis of submicron-resolution

spatial transcriptomics. *Nat. Methods* **21**, 1843–1854 (2024).

18.    Mitchel, J., Gao, T., Cole, E., Petukhov, V. & Kharchenko, P. V. Impact of

Segmentation Errors in Analysis of Spatial Transcriptomics Data. 2025.01.02.631135

Preprint at https://doi.org/10.1101/2025.01.02.631135 (2025).

19.    Nguyen, H., Nguyen, H., Tran, D., Draghici, S. & Nguyen, T. Fourteen years of

cellular deconvolution: methodology, applications, technical evaluation and

outstanding challenges. *Nucleic Acids Res*. **52**, 4761–4783 (2024).

20.    Garmire, L. X. *et al*. Challenges and perspectives in computational deconvolution of

genomics data. *Nat. Methods* **21**, 391–400 (2024).

21.    Gaspard-Boulinc, L. C., Gortana, L., Walter, T., Barillot, E. & Cavalli, F. M. G. Cell-

type deconvolution methods for spatial transcriptomics. *Nat. Rev. Genet.* 1–19 (2025)

doi:10.1038/s41576-025-00845-y.

22.    Ament, S. A. & Poulopoulos, A. The brain's dark transcriptome: Sequencing RNA in

distal compartments of neurons and glia. *Curr. Opin. Neurobiol*. **81**, 102725 (2023).

23.    Ivich, A. *et al*. Missing cell types in single-cell references impact deconvolution of

bulk data but are detectable. *Genome Biol*. **26**, 86 (2025).

24.    Han, J. *et al*. Single-cell sequencing unveils key contributions of immune cell

populations in cancer-associated adipose wasting. *Cell Discov*. **8**, 122 (2022).

25.    Fetit, R. *et al*. Characterizing Neutrophil Subtypes in Cancer Using scRNA

Sequencing Demonstrates the Importance of IL1β/CXCR2 Axis in Generation of

Metastasis-specific Neutrophils. *Cancer Res. Commun*. **4**, 588–606 (2024).

26.    Sutton, M. A. *et al*. Miniature Neurotransmission Stabilizes Synaptic Function via

Tonic Suppression of Local Dendritic Protein Synthesis. *Cell* **125**, 785–799 (2006).

27.    Cajigas, I. J. *et al*. The Local Transcriptome in the Synaptic Neuropil Revealed by

Deep Sequencing and High-Resolution Imaging. *Neuron* **74**, 453–466 (2012).

28.    Hafner, A.-S., Donlin-Asp, P. G., Leitch, B., Herzog, E. & Schuman, E. M. Local

protein synthesis is a ubiquitous feature of neuronal pre- and postsynaptic

compartments. *Science* **364**, eaau3644 (2019).

29.    Glock, C. *et al*. The translatome of neuronal cell bodies, dendrites, and axons. *Proc.*

*Natl. Acad. Sci.* **118**, e2113929118 (2021).

30.    von Kügelgen, N. & Chekulaeva, M. Conservation of a core neurite transcriptome

across neuronal types and species. *WIREs RNA* **11**, e1590 (2020).

31.    Lee, K.-W., Kim, H.-J., Sung, J.-J., Park, K.-S. & Kim, M. Defective neurite outgrowth

in aphidicolin/cAMP-induced motor neurons expressing mutant Cu/Zn superoxide

dismutase. *Int. J. Dev. Neurosci.* **20**, 521–526 (2002).

32.    Markmiller, S. *et al*. Persistent mRNA localization defects and cell death in ALS

neurons caused by transient cellular stress. *Cell Rep.* **36**, 109685 (2021).

33.    Buxbaum, A. R., Haimovich, G. & Singer, R. H. In the right place at the right time:

visualizing and understanding mRNA localization. *Nat. Rev. Mol. Cell Biol.* **16**, 95–109

(2015).

34.    Gandalovičová, A., Vomastek, T., Rosel, D. & Brábek, J. Cell polarity signaling in the

plasticity of cancer cell invasiveness. *Oncotarget* **7**, 25022–25049 (2016).

35.    Backlund, M. *et al*. Plasticity of nuclear and cytoplasmic stress responses of RNA-

binding proteins. *Nucleic Acids Res.* **48**, 4725–4740 (2020).

36.    Apicco, D. J. *et al*. Reducing the RNA binding protein TIA1 protects against tau-

mediated neurodegeneration in vivo. *Nat. Neurosci.* **21**, 72–80 (2018).

37.    Wolozin, B. & Ivanov, P. Stress granules and neurodegeneration. *Nat. Rev.*

*Neurosci.* **20**, 649–666 (2019).

38.    Ridder, K. *et al.* Extracellular vesicle-mediated transfer of functional RNA in the

tumor microenvironment. *OncoImmunology* **4**, e1008371 (2015).

39.    Li, I. & Nabet, B. Y. Exosomes in the tumor microenvironment as mediators of

cancer therapy resistance. *Mol. Cancer* **18**, 32 (2019).

40.    O'Brien, K., Breyne, K., Ughetto, S., Laurent, L. C. & Breakefield, X. O. RNA delivery

by extracellular vesicles in mammalian cells and its applications. *Nat. Rev. Mol. Cell*

*Biol.* **21**, 585–606 (2020).

41.    Landry, F. J. & Findlay, S. R. Purification of human basophils by negative selection. *J.*

*Immunol. Methods* **63**, 329–336 (1983).

42.    Hansel, T. T. *et al.* Purification of human blood eosinophils by negative selection

using immunomagnetic beads. *J. Immunol. Methods* **122**, 97–103 (1989).

43.    Hansel, T. T. *et al.* An improved immunomagnetic procedure for the isolation of

highly purified human blood eosinophils. *J. Immunol. Methods* **145**, 105–110 (1991).

44.    Munoz, N. M. & Leff, A. R. Highly purified selective isolation of eosinophils from

human peripheral blood by negative immunomagnetic selection. *Nat. Protoc.* **1**, 2613–

2620 (2006).

45.    She, Y. & Owen, A. B. Outlier Detection Using Nonconvex Penalized Regression. *J.*

*Am. Stat. Assoc.* **106**, 626–639 (2011).

46.    Lee, Y., MacEachern, S. N. & Jung, Y. Regularization of Case-Specific Parameters for

Robustness and Efficiency. *Stat. Sci.* **27**, 350–372 (2012).

47.    Kong, D., Bondell, H. D. & Wu, Y. Fully Efficient Robust Estimation, Outlier Detection

and Variable Selection Via Penalized Regression. *Stat. Sin.* **28**, 1031–1052 (2018).

48.    Traniello, I. M. *et al.* Single-cell dissection of aggression in honeybee colonies. *Nat.*

*Ecol. Evol.* **7**, 1232–1244 (2023).

49.    Lin, L. I.-K. A Concordance Correlation Coefficient to Evaluate Reproducibility.

*Biometrics* **45**, 255–268 (1989).

50.    Kumar, A. *et al.* Intracellular spatial transcriptomic analysis toolkit (InSTAnT). *Nat.*

*Commun.* **15**, 7794 (2024).

51.    Yeo, S., Schrader, A. W., Lee, J., Asadian, M. & Han, H.-S. Spot-Based Global

Registration for Subpixel Stitching of Single-Molecule Resolution Images for Tissue-

Scale Spatial Transcriptomics. *Anal. Chem.* **96**, 6517–6522 (2024).

52.    Wang, G. *et al.* Spatial organization of the transcriptome in individual neurons.

2020.12.07.414060 Preprint at https://doi.org/10.1101/2020.12.07.414060 (2020).

53.    Lake, B. B. *et al.* A comparative strategy for single-nucleus and single-cell

transcriptomes confirms accuracy in predicted cell-type expression from nuclear RNA.

*Sci. Rep.* **7**, 6031 (2017).

54.    Habib, N. *et al.* Massively parallel single-nucleus RNA-seq with DroNc-seq. *Nat.*

*Methods* **14**, 955–958 (2017).

55.    Maynard, K. R. *et al.* Transcriptome-scale spatial gene expression in the human

dorsolateral prefrontal cortex. *Nat. Neurosci.* **24**, 425–436 (2021).

56.    DeFelipe, J. *et al.* Cortical white matter: beyond the pale remarks, main conclusions

and discussion. *Front. Neuroanat.* **4**, 1692 (2010).

57.    Overly, C. C., Rieff, H. I. & Hollenbeck, P. J. Organelle motility and metabolism in

axons vs dendrites of cultured hippocampal neurons. *J. Cell Sci.* **109**, 971–980 (1996).

58.    Harris, J. J., Jolivet, R. & Attwell, D. Synaptic Energy Use and Supply. *Neuron* **75**,

762–777 (2012).

59.    Johnson, Z. V. *et al.* Cellular profiling of a recently-evolved social behavior in cichlid

fishes. *Nat. Commun.* **14**, 4891 (2023).

60.    Than-Trong, E. & Bally-Cuif, L. Radial glia and neural progenitors in the adult

zebrafish central nervous system. *Glia* **63**, 1406–1428 (2015).

61.    Miranda-Negrón, Y. & García-Ararás, J. E. Radial glia and radial glia-like cells: Their

role in neurogenesis and regeneration. *Front. Neurosci.* **16**, (2022).

62.    Zhu, J., Sun, S. & Zhou, X. SPARK-X: non-parametric modeling enables scalable and

robust detection of spatial expression patterns for large spatial transcriptomic studies.

*Genome Biol.* **22**, 184 (2021).

63.    Abramsson, A. *et al.* The zebrafish amyloid precursor protein-b is required for motor

neuron guidance and synapse formation. *Dev. Biol.* **381**, 377–388 (2013).

64.    Newman, M., Ebrahimie, E. & Lardelli, M. Using the zebrafish model for Alzheimer's

disease research. *Front. Genet.* **5**, (2014).

65.    Haenisch, C. *et al.* The neuronal growth and regeneration associated Cntn1

(F3/F11/Contactin) gene is duplicated in fish: expression during development and retinal

axon regeneration. *Mol. Cell. Neurosci.* **28**, 361–374 (2005).

66.    Eiraku, M. *et al.* DNER acts as a neuron-specific Notch ligand during Bergmann glial

development. *Nat. Neurosci.* **8**, 873–880 (2005).

67.    Díaz-Ramos, À., Roig-Borrellas, A., García-Melero, A. & López-Alemany, R. α-

Enolase, a Multifunctional Protein: Its Role on Pathophysiological Situations. *BioMed*

*Res. Int.* **2012**, 156795 (2012).

68.    Zhou, Z., Li, X., Wright, J., Candès, E. & Ma, Y. Stable Principal Component Pursuit.

in *2010 IEEE International Symposium on Information Theory* 1518–1522 (2010).

doi:10.1109/ISIT.2010.5513535.

69.    Mateos, G. & Giannakis, G. B. Robust PCA as Bilinear Decomposition With Outlier-

Sparsity Regularization. *Trans Sig Proc* **60**, 5176–5190 (2012).

70.    Giannakis, G. B., Mateos, G., Farahmand, S., Kekatos, V. & Zhu, H. USPACOR:

Universal sparsity-controlling outlier rejection. in *2011 IEEE International Conference on*

*Acoustics, Speech and Signal Processing (ICASSP)* 1952–1955 (2011).

doi:10.1109/ICASSP.2011.5946891.

71.    Huang, P., Cai, M., Lu, X., McKennan, C. & Wang, J. Accurate estimation of rare cell-

type fractions from tissue omics data via hierarchical deconvolution. *Ann. Appl. Stat.* **18**,

1178–1194 (2024).

72.    Zhong, Y. & Liu, Z. Gene expression deconvolution in linear space. *Nat. Methods* **9**,

8–9 (2012).

73.    Hicks, S. C., Townes, F. W., Teng, M. & Irizarry, R. A. Missing data and technical

variability in single-cell RNA-sequencing experiments. *Biostatistics* **19**, 562–578 (2018).

74.    Townes, F. W., Hicks, S. C., Aryee, M. J. & Irizarry, R. A. Feature selection and

dimension reduction for single-cell RNA-Seq based on a multinomial model. *Genome*

*Biol.* **20**, 295 (2019).

75.    Butler, A., Hoffman, P., Smibert, P., Papalexi, E. & Satija, R. Integrating single-cell

transcriptomic data across different conditions, technologies, and species. *Nat.*

*Biotechnol*. **36**, 411–420 (2018).

**Figure Legends**

**Figure 1. Schematic overview of RESCUE.** Overview of RESCUE. RESCUE is a

computational negative selection method designed to recover the idiosyncratic

transcriptome. It takes as input ST data to be decomposed, along with a reference matrix

constructed from sc/snRNA-seq data from the same tissue. With these two data inputs,

RESCUE performs robust sparse recovery of the idiosyncratic transcriptome through a

regularized multivariate regression problem. To account for gene variability across

different locations, RESCUE iteratively updates a weight matrix within its alternating

minimization algorithm. RESCUE partitions the full transcriptome profile into the canonical

and idiosyncratic expression, which can be further used for downstream analysis, such

as DEG and SVG analyses. Created with BioRender.com.

**Figure 2. Simulation results of deconvolution and idiosyncratic transcriptome**

**recovery. A.** Comparisons of cell-type deconvolution performance using the average

Lin's CCC across cell types. **B.** Comparisons of idiosyncratic transcriptome recovery

performance using the relative Frobenius error. The proportion of idiosyncratic expression

ranges from 0% to 80% and the idiosyncratic sparsity level ranges from 80% to 95%.

**Figure 3. Application to the honey bee brain MERFISH data. A.** Comparison of

segmentation-based ("Cell Mask") and estimated somatic proportions. Cell mask ratio is

computed as the proportion of MERFISH spots in each grid lying within a cell. **B.** Bar plots

of Lin's CCC and RMSE between segmentation-based and estimated idiosyncratic

proportions. For each grid, idiosyncratic proportions are calculated as 1 – (proportion of

grid's somatic expression). **C.** Comparison of segmented and estimated transcript counts

for each grid in cell body regions where the cell mask ratio is 1. Grey dashed lines indicate

$y = x$ and blue solid lines represent the smoothed line from estimates.

**Figure 4. Application to the human breast cancer Visium data. A.** Manual annotation

of Visium ST data from human breast cancer tissue. **B.** Heatmap of gene expression

correlations between pseudo-bulk Visium and scFFPE-seq cell types. Red box highlights

high similarity between adipocytes and other cell types including stromal cells. **C.**

Estimated cell-type proportions in adipocyte regions across methods. Existing methods

overestimate stromal and endothelial populations due to the absence of adipocytes in the

reference. RESCUE avoids such misattributions by modeling idiosyncratic expression. **D.**

Comparison of estimated versus reference cell-type proportions across all methods.

**Figure 5. Application to the human DLPFC Visium data. A.** Anatomical annotations of

10x Visium samples from the DLPFC dataset. **B.** Idiosyncratic expression proportions in

each layer estimated by RESCUE. RESCUE estimates higher idiosyncratic proportions

in Layer 1 and WM, consistent with their enrichment in neuropil and myelinated axons,

respectively. **C.** Differential expression analysis comparing Layer 1 to other layers using

observed ST expression ("raw") versus idiosyncratic expression. Each point represents a

gene, with log-fold changes from raw expression (x-axis) versus idiosyncratic expression

(y-axis). Blue dashed lines indicate the thresholds at log-fold changes = 0. Red-labeled

genes are top markers of Layer 1, detected from the idiosyncratic expression. **D.** Gene

ontology (GO) enrichment analysis results of Layer 1-enriched differentially expressed

genes (DEGs) based on idiosyncratic (left) and raw (right) expression.

**Figure 6. Application to the cichlid fish telencephalon Visium data. A.** Schematic of

radial glial (RG) anatomy (left; Johnson et al.[59]) and regional annotation of Visium ST data

from the dorsal telencephalon of cichlid fish (right). **B.** Comparison of adjusted p-values

for SVG detection using raw (x-axis) versus idiosyncratic (y-axis) expression. Blue

dashed lines indicate the significance threshold at adjusted p-values = 0.05. Red-labeled

genes are examples of SVGs detected from idiosyncratic expression but not detectable

from raw data. **C.** Spatial expression maps of idiosyncratic expression for unique SVGs.

**Figure 1.**

**Figure 2.**

**Figure 3.**

**Figure 4.**

**Figure 5.**

A

B

C

D

**Figure 6.**