Preventing unintentional bat captures in canopy traps used for insect surveys

A. Sallé[1✉], L. Arthur[2], G. Parmain[3], D. Williams[4], A. Chrétien[5], E. Le Souchu[1], C. Moliard[3], A.
Roques[6], C. Bouget[3]

[1] Laboratoire Physiologie, Ecologie et Environnement (P2E), Université d'Orléans, UR 1207,
USC-INRAE 1328, 1 rue de Chartres, Orléans 45067, France

[2] NGO « Chauve-qui-peut », L'observatoire, 22 rue Ranchot, 18000 Bourges

[3] UR EFNO, INRAE, Domaine des Barres, Nogent-sur-Vernisson, France

[4] Forest Research, Centre of Forest Protection, Alice Holt Lodge, Farnham, UK

[5] Muséum d'Histoire Naturelle de Bourges, Parc Saint Paul, 18000 Bourges, France

[6] URZF UR 0633, INRAE, Orléans, France

ORCID number

A. Sallé: 0000-0003-3876-9272

G. Parmain: 0000-0001-6275-2898

E. Le Souchu: 0000-0002-8186-0349

A. Roques: 0000-0002-3734-3918

C. Bouget: 0000-0002-5206-7560

✉ Corresponding author: Aurélien Sallé, aurelien.salle@univ-orleans.fr

Acknowledgements

This work was also supported by Région Centre-Val de Loire Project no. 2018-00124136
(CANOPEE) coordinated by A. Sallé. We thank O. Denux (INRAE) and X. Pineau (University of
Orléans) for their technical assistance. The DREAL provided legal authorizations for the
incidental capture of bats, according to article R411-7 of the French Environmental Code (#
41-2021-04-26-00004, 36-2021-04-07-00008, DDT-2021/001).

Author contributions

Aurélien Sallé: conceptualization; methodology; investigation; validation; formal analysis;
funding acquisition; writing – original draft; writing – review and editing; project
administration. Laurent Arthur: investigation; writing – review and editing. Guillem Parmain:
investigation; data curation; writing – review and editing. Dave Williams: methodology;
writing – review and editing. Amélie Chrétien: investigation; writing – review and editing.
Elodie Le Souchu: investigation; writing – review and editing. Carl Moliard: methodology.
Alain Roques: conceptualization; methodology; investigation; writing – review and editing.
Christophe Bouget: conceptualization; methodology; investigation; writing – review and
editing.

Funding information

This work was also supported by Région Centre-Val de Loire Project no. 2018-00124136
(CANOPEE) coordinated by A. Sallé.

Conflict of interest statement

The authors declare no conflicts of interest.

Abstract

There is growing interest in deploying insect traps in forest canopies for ecological research,
invasive species surveillance, and biodiversity monitoring. However, forest canopies also
harbor a diverse and abundant bat community. In this study, we investigated the impact of
flight interception traps and multi-funnel traps deployed in oak canopies on incidental bat
captures, and evaluated whether trap modifications could reduce this bycatch. We also
examined how these modifications influenced the species richness, abundance, and mean
body size of several beetle taxa, including Buprestidae, Cerambycidae, Cetoniinae, and
Scolytinae. Both trap types captured bats. Flight interception traps collected seven species -
Myotis bechsteinii, *Myotis daubentonii*, *Myotis mystacinus*, *Nyctalus noctula*, *Nyctalus
leisleri*, *Plecotus auritus*, and *Pipistrellus pipistrellus* - with an average of 0.38 to 0.45 bats per
trap. Multi-funnel traps primarily captured males of *P. pipistrellus*, with an average of 0.24 to
0.33 bats per trap. Adding a 13-mm mesh above the collection container significantly
reduced bat bycatch to 0.02 to 0.05 bats per trap. This modification had no significant effect
on insect species richness or mean body size, although it did reduce beetle abundance in
Cerambycidae in black multi-funnel traps, and Scolytinae in flight interception traps, likely
due to more frequent clogging by twigs and leaves. We recommend modifying flight
interception and multi-funnel traps deployed in forest canopies to prevent unintentional
harm to protected bat species.

Keywords

Lindgren trap, multi-funnel, flight interception trap, ecological trap, bycatch

Introduction

Insects are routinely sampled in forests for purposes related to biodiversity monitoring, invasive species surveillance and pest management. A variety of survey techniques - often involving traps - have been developed for the detection, monitoring, or inventory of either broad insect communities or specific target species (Brockerhoff et al., 2023). However, these traps are rarely taxon-specific and frequently capture a wide range of non-target organisms. Most reported bycatch consists of invertebrates (e.g., Zumr and Stary, 1991; Rojas-Nossa et al., 2018; Hafsi et al., 2020; Sukovata et al., 2022), and in some cases, it provides an opportunity to detect rare or previously unrecorded native and exotic species (Vincent et al., 2020; Le Souchu et al., 2024a; Mas et al., 2023). However, certain trap designs can also unintentionally capture and occasionally kill vertebrates. For example, conventional pitfall traps, commonly used to sample ground-dwelling invertebrates, may also cause mortality in small mammals and amphibians (Pearce et al., 2005). Avoiding bycatch is important not only for ethical considerations related to the killing of beneficial and/or patrimonial species, but also for regulatory compliance, as some non-target species may be legally protected. Moreover, the incidental capture of non-target organisms may compromise the quality and representativeness of insect samples. Decomposing bodies of vertebrates accelerate the decomposition of insect samples and may attract additional insect species (Pearce et al., 2005), while collecting a large volume of non-target organisms may also reduce trap efficiency for targeted species (Spears et al., 2021). This would in turn bias the sample and increase the time necessary to sort the samples (Pearce et al., 2005). To avoid these issues, while maintaining the effectiveness of the trap, trap designs often need to be adapted (e.g. Pearce et al., 2005; Spears et al., 2021).

The canopy of forests provides a diverse range of specific trophic resources, microhabitat and microclimatic conditions for insects (Sallé et al., 2021). As a result, it harbors diverse and unique, yet largely understudied, arthropod communities (Floren and Schmidl, 2008; Bouget et al., 2011; Ulyshen, 2011; Urban-Mead et al., 2021). In this regard, temperate forest canopies may serve as reservoirs for rare and conservation-important species (Plewa et al., 2017; Le Souchu et al., 2024a). The canopy may also provide critical resources to fulfill the life cycle of forest insect species associated with other forest strata and may consequently provide an optimal location for trapping certain species (Ulyshen, 2011; Sallé et al., 2021). Setting traps in the canopy may consequently improve the detection and monitoring of both patrimonial insect species and major forests pests (Plewa et al., 2017; Rassati et al., 2019; Sallé et al., 2020).

Many species of bats also rely on the forest canopy for roosting and foraging (Müller et al., 2013; Law et al., 2016; Erasmy et al., 2021). This is particularly true for the canopy of old-growth oak forests which harbor a large amount and diversity of microhabitats, including loose bark, cracks and cavities that are used by bats for roosting and nesting (Regnery et al., 2013). Forest insects are an important food source, and bats may play a role in the control of forest pests (Beilke and O'Keefe, 2023; Augusto et al., 2024). Bat populations in Europe are

thought to have experienced significant declines in the last century, and some species have
undergone significant range contractions (Browning et al., 2020). In France, eight of the 36
bat species are classified as threatened, and eight are considered near threatened (Arthur
and Lemaire, 2021). In Europe, 25 percent of the 40 bat species are threatened (Temple and
Terry, 2009), and all bat species consequently benefit from legal protection across Europe
(Council Directive 92/43/EEC, 1992). Nevertheless, several studies have reported incidental
captures of bats in traps intended for insect sampling (McLeod et al., 2021; Holzinger et al.,
2023; Ancillotto et al., 2024).

This study reports the incidental capture of bats by three types of insect trapping devices
deployed in the canopy of oak stands in France, designed primarily for sampling canopy-
dwelling insect fauna. Given the growing interest in canopy trapping for ecological research,
invasive insect surveillance, and monitoring programs, we propose a simple modification to
these devices to minimize bat bycatch while maintaining trapping efficacy for the targeted
insect groups. We first document the species composition and abundance of bats captured,
then evaluate the effectiveness of the modified trap design in reducing these unintended
captures. Lastly, we assess the impact of change in trapping design on the abundance,
species richness, and mean body size of the targeted insect taxa.

Material and methods

The study was conducted in six oak-dominated forests located in France (Table 1). The two
dominant tree species in these forests were *Quercus robur* L. and *Quercus petraea* (Matt)
Liebl. In each forest we selected 3 to 9 stands. The main objective of the study was to
characterise the community of insects dwelling in oak canopies. For this, three kinds of traps
were used: black and green multi-funnel traps (BMF and GMF respectively, Chemtica
Internacional, San Jose, Costa Rica), with 12 fluon-coated funnels (figure 1), and flight
interception traps with transparent panels (FI, Ecole d'ingénieurs de Purpan) (figure 1). Traps
were suspended among the lower branches in the canopy (i.e. approximately 10 - 15 m
above the ground). The collectors were filled with a solution of 50% (v/v) monopropylene
glycol and water plus a drop of detergent. Setting GMF traps in the canopy layer is
considered as one of the most efficient sampling approaches to collect *Agrilus* species
(Rassati et al., 2019; Sallé et al., 2020), and can be used either to monitor native species or
detect exotic ones (Santoiemma et al., 2024; Le Souchu et al., 2024b), but also to monitor
and detect a wide array of canopy-dwelling insects (Sallé et al., 2020 ; Vincent et al., 2020 ;
Le Souchu et al., 2024a). BMF traps were baited with volatiles (a blend of longhorn
pheromones, ethanol and alpha-pinene, see Roques et al., 2023). These traps have been
developed for early detection of exotic longhorn beetles (Roques et al., 2023). FI traps are
routinely used to sample saproxylic beetles (Bouget et al., 2009). The traps were installed
during three consecutive years, from 2019 to 2021, approximately from April to September,

and were emptied every month. The number of stands, and traps per stand, varied among
forests and years (table 1).

In 2019, all traps were installed as supplied by the manufacturers. In 2020 and 2021, to
prevent further captures of bats, as they are protected species, all traps were modified
accordingly, and no unmodified (control) traps were used in parallel. In 2020, GMF and FI
traps were equipped with wooden sticks (5 mm diameter), set as cross, in the last funnel
over the collection cup (figure 1). This modification is similar to the one proposed by
Holzinger et al. (2023) to prevent bats from being captured by FI traps. BMF traps were not
used in 2020. Since this modification proved ineffective (see Results), we implemented and
evaluated an alternative modification in 2021. A 13-mm wire mesh was fixed onto a small
funnel inserted inside the last funnel of the traps, over the collection cup of all trap types
(figure 1), as proposed by McLeod et al. (2021). Preliminary assays conducted at the
Museum of Bourges using live common pipistrelles from an animal shelter indicated that
larger mesh sizes (2 x 2 cm and 2 x 4 cm) were easily crossed by the bats and, consequently,
were too wide. The mesh was connected to the rim of the last funnel to allow bats to exit
the trap. We obtained a legal authorization from regional authorities (Direction Régionale de
l'Environnement de l'Aménagement et du Logement) for the incidental capture of bats for
scientific purposes according to article R411-7 of the French Environmental Code, pending
traps were adapted to avoid these captures.

Figure 1: Flight interception trap (FI, top left) and black multi-funnel trap (BMF, top right). Modified
funnels with wooden sticks (bottom left) and a 13-mm metallic mesh (bottom right) placed just
above the collection cup.

Table 1. Sampling sites and years, and number of traps used. Lat.: latitude, Long.: longitude, GMF:
green multi-funnel traps, BMF: black multi-funnel traps, FI: flight interception traps. See text for
details. WS-: traps modified with wooden sticks, MM-: traps modified with a metallic mesh.

Location	Lat.	Long.	Years of monitoring	Number of stands	Type of traps used (n)
State forest of Boulogne	47.57	1.53	2021	6	MM-GMF (12) MM-BMF (6) MM-FI (12)
Forest of Chantilly	49.17	2.51	2021	3	MM-GMF (6) MM-BMF (3) MM-FI (6)
State forest of Châteauroux	46.74	1.72	2021	6	MM-GMF (12) MM-BMF (6) MM-FI (12)
State forest of l'Abbaye	46.23	3.37	2019	3	GMF (6) BMF (3) FI (6)
			2021	3	MM-GMF (6) MM-BMF (3) MM-FI (6)
State forest of Orléans	47.87	2.41	2019	9	GMF (18) BMF (9) FI (18)
			2020	6	WS-GMF (12) WS-FI (12)
			2021	6	MM-GMF (12) MM-BMF (6) MM-FI (12)
State forest of Vierzon	47.46	2.19	2019	9	GMF (18) BMF (9) FI (18)
			2020	6	WS-GMF (12) WS-FI (12)
			2021	6	MM-GMF (12) MM-BMF (6) MM-FI (12)

Bats and insects identification

We counted the bats caught in all types of traps during the survey. Bats caught in GMF and
FI traps in 2019 and 2020 were kept for identification by the Museum of Natural History of
Bourges (France). Bats collected were also sexed. Bats collected in BMF traps were counted
but not identified. Saproxylic beetles collected were identified at the species level for most
families.

Statistical analyses

All analyses were performed in R, version 4.2.1 (R Core Team 2023). To assess whether mean
bat catches per trap varied among years, and after the addition of the wire mesh, we
performed pairwise Wilcoxon tests, with a Holm correction for multiple comparisons.

In addition, we have tested whether the change in trap design affected the abundance,
species richness and average size of beetles collected. We focussed these analyses on key
taxonomical groups of beetles. For this, we selected several families or subfamilies of
saproxylic beetles which are either significant forest pests and/or patrimonial species,
exhibit various body sizes, and were well sampled by our different trap designs.
Consequently we selected four groups: (i) longhorn beetles (Cerambycidae) which have
average to large body sizes (13.5 ± 0.5 mm body length, calculated from the database
mentioned below), include both pest and patrimonial species, and are specifically targeted
by BMF traps, (ii) jewel beetles (Buprestidae) which have average body sizes (9.0 ± 0.5 mm),
include pest species, and are specifically targeted by GMF traps, (iii) flower chafers

(Cetoniinae) which have large body sizes (17.4 ± 1.2 mm), include patrimonial species and
are well sampled by FI traps and (iv) bark beetles (Scolytinae), which have small body sizes
(2.5 ± 0.1 mm), include pest species, and are also well sampled by FI traps. We assessed
whether the mean abundance and species richness per trap varied before (2019) and after
(2021) the addition of the wire mesh with Kruskal-Wallis tests. Bark beetles were not
identified in BMF traps in 2021 and were consequently discarded from the analyses for this
trap type.

Finally, to assess the effect of wire mesh on the average body size of collected beetles, we
first assigned an average body size to each of the sampled taxa, within the four above
mentioned taxonomic groups, using the database provided by Bouget and coll. (2019). For
each group, we then calculated the average body size per trap in 2019 and 2021. For each
trap type, we restricted the analyses to the most frequently sampled taxonomic groups (i.e.
Buprestidae for GMF traps, Cerambycidae for BMF traps and Cetoniinae and Scolytinae for FI
traps). To assess whether the average body sizes varied significantly, we performed Kruskal-
Wallis tests. For all tests, the significance level was set at $\alpha = 0.05$.

Results

Bat captures

In 2019, 33 bats were captured: 5 in the 21 BMF traps, 11 in the 42 GMF traps and 17 in the
42 FI traps. In 2020, 17 bats were caught: 8 in the 24 GMF traps, and 9 in the 24 FI traps. One
FI trap collected 6 bats during a single survey. In 2021, 5 bats were captured: 1 in the 30 BMF
traps, 3 in one of the 60 GMF traps and 1 in FI traps (table S1). While the average number of
bats did not differ between 2019 and 2020 in both FI and GMF traps, it significantly
decreased in 2021, in all trap types (figure 2).

In 2019 and 2020, FI traps collected six species of bats, mostly Bechstein's bats (37%) and
common pipistrelles (30%), while GMF traps collected only two species, mostly common
pipistrelles (95%) (table 2). Overall, the captured specimens were predominantly male (83%).
Approximately half of the captures occurred in July, with the remainder mostly distributed
between May and June.

Table 2: number of individuals (males and/or females) of bat species captured in flight interception traps (FI)
and green multi-funnel traps (GMF) installed in oak canopies in 2019 and 2020. nd: not determined

Trap type	*Myotis bechsteinii* (Kuhl)	*Myotis daubentonii* (Kuhl)	*Myotis mystacinus* (Kuhl)	*Nyctalus noctula* (Schreber)	*Nyctalus leisleri* (Kuhl)	*Plecotus auritus* (L.)	*Pipistrellus pipistrellus* (Schreber)
FI	10 (4m, 6f)	2 (m)	1 (nd)	2 (1m, 1 nd)	3 (m)	1 (m)	8 (m)
GMF			1 (m)				18 (m)

Figure 2: mean number of bats (± standard error) collected in black multi-funnel traps (BMF), green multi-funnel traps (GMF), and flight interception traps (FI) in 2019, in 2019 (with a standard collection cup), 2020 (with a wooden cross above the collection cup) and 2021 (with wire mesh above the collection cup)

Insect surveys

The mean abundance of beetles per trap decreased in 2021 for Cerambycidae in BMF traps and for Scolytinae in FI traps, but did not significantly vary for other taxonomical groups (figure 3). The mean species richness per trap of the four taxonomic groups considered did not significantly vary between 2019 and 2021 (figure 3).

Figure 3: mean abundance (± standard error) (left) and mean species richness (± standard error) (right) per trap of Buprestidae, Cerambycidae, Cetoniinae and Scolytinae collected in green multi-funnel traps, black multi-funnel traps, and flight interception traps in 2019, i.e. with a standard collection cup, and in 2021, with a wire mesh above the collection cup. Symbols: ˙ indicates a p.value < 0.1, * indicates a p.value < 0.05. Note that abundance scales are different among families and trap types.

The average body size per trap slightly increased in 2021 for the Cerambycidae in BMF traps (figure 4). It did not significantly vary for the other taxonomic groups considered, although it tended to decrease for the Cetoniinae (figure 4).

249 Figure 4: mean body size (± standard error) per trap of species of Buprestidae, Cerambycidae, Scolytinae and
250 Cetoniinae collected in green multi-funnel traps (green bars: Buprestidae), black multi-funnel traps (grey bars:
251 Cerambycidae), and flight interception traps (white and blue bars: Cetoniinae and Scolytinae) in 2019, i.e. with
252 a standard collection cup, and in 2021, with a modified collection cup. Symbols: · indicates a p.value < 0.1, *
253 indicates a p.value < 0.05.

Discussion

All trapping devices set in oak canopies incidentally captured bats. However, while FI traps
collected a relatively diverse range of species, MF traps almost exclusively captured males of
common pipistrelles. This may seem surprising given that bats use echolocation to detect
prey and avoid obstacles (Jones and Holderied, 2007). Nevertheless, several studies have
reported that bats can collide with large stationary objects (Orbach and Fenton, 2010; Greif
et al., 2017; Tanalgo et al., 2025). Smooth vertical surfaces may function as ecological traps,
as bats can perceive tilted smooth vertical plates as open flyways (Greif et al., 2017). This is a
plausible explanation for the bycatch observed in FI traps (Holzinger et al., 2023), but it does
not account for the captures in MF traps. An alternative explanation is that bats are
attracted to insects resting on the traps, or to the sounds produced by insects trapped within
the collectors, and fall into the traps while trying to catch them. A third possibility, specific to
MF traps, is that pipistrelles mistake these devices for potential roosting sites, resembling
loose bark or bark crevices, which they commonly use for summer roosting (Bat Tree Habitat
Key, 2018; Holzinger et al., 2023).

The bat species captured with both trap types are commonly found in French oak forests
(Tillon et al., 2016; L. Arthur, pers. comm.). Two additional species, namely *Myotis alcathoe*
Helversen & Heller and *Barbastella barbastellus* (Schreber), are also known to occur in the
region's forests but were not captured during our survey. FI traps deployed in an Austrian
oak forest also caught *M. bechsteinii* and *N. noctula* as well as *M. alcathoe, Myotis brandtii*
(Eversmann) and *Pipistrellus pygmaeus* (Leach) (Holzinger et al., 2023). In our study, most of

the pipistrelle captures occurred during the maternity and juvenile-rearing period (May to
July). During this time, colonies are primarily composed of females, which typically settle in
anthropogenic sites such as towns and villages (Mathews et al., 2023). Solitary males, on the
other hand, tend to forage in distinct areas, including forests (L. Arthur, pers. obs.), which
may explain why they were overwhelmingly predominant in our captures as well as in those
reported in Austria (Holzinger et al., 2023). We observed several times two or more bats in
the same collector, with a maximum of six female Bechstein's bats caught together. This
explains the high standard errors for mean bat catch per trap (figure 2). It is possible that
trapped individuals emitted distress calls that attracted conspecifics, especially if a roosting
site was in the vicinity (Russ et al., 1998).

Few previous studies have reported bats being captured in FI or MF traps (McLeod et al.,
2021; Holzinger et al., 2023). MF traps were originally developed to collect bark beetles
(Lindgren, 1983) and have since been widely used across the globe for sampling forest
insects. Similarly, FI traps are commonly employed for forest insect monitoring (Bouget et
al., 2009). However, these traps are typically placed at ground level, where they might be
less likely to interfere with bat flight paths. Nevertheless, some evidence exists that Western
long-eared bats (*Myotis evotis* (H. Allen)) have been unintentionally captured in MF traps set
in harvested Douglas fir stands (McLeod et al., 2021). Since the introduction of the emerald
ash borer (*Agrilus planipennis*, Fairmaire) in North America, GMF traps have been
extensively deployed for the detection and monitoring of this invasive pest (Crook et al.,
2014; Santoiemma et al., 2024), and it is recommended to place these traps within the tree
canopy (Crook et al., 2014), as in our sampling design. This maximizes detection of *A.*
planipennis, as newly emerged adults feed and mate in the foliage (Crook et al., 2014). The
common pipistrelle is not present in North America but is widely distributed across the
Western Palearctic (Mathews et al., 2023). This could explain why, despite the widespread
use of MF traps in that region, there have not been reports of large-scale bat captures. The
emerald ash borer has also been introduced into Russia and is currently spreading westward
through Europe, posing a serious threat to native ash populations (Orlova-Bienkowskaja,
2014; Orlova-Bienkowskaja et al., 2020). In Europe, deploying GMF traps in ash canopies is
recommended for the early detection of this pest, allowing for timely implementation of
management measures (European Food Safety Authority et al., 2020; Evans et al., 2020;
Santoiemma et al., 2024). However, if not properly adapted, these traps could inadvertently
cause significant bat mortality, including among species classified as vulnerable in France
(e.g., *N. noctula*) or Europe (e.g., *M. bechsteinii*).

The addition of wooden sticks did not prevent bats from falling into the collectors,
suggesting that the trap modification proposed by Holzinger et al. (2023) may not be
effective in mitigating bat bycatch. However, installing a 13-mm metallic mesh significantly
reduced the number of bat captures. Since we did not operate standard and modified traps

simultaneously, out of concern for further bat mortality, we cannot entirely rule out the
possibility that the decrease in bat captures observed in 2021 was due to interannual
fluctuations in bat populations. However, this explanation appears unlikely, as it would imply
a simultaneous population decline across all bat species. Furthermore, as these modified
traps have since been used for insect sampling in forest canopies in both France and the
United Kingdom, and bat captures have been rarely observed, this supports the conclusion
that adding a mesh is an effective measure to reduce incidental bat captures. The mesh did
not completely prevent bat captures, and a few individuals still fell into the collectors in
2021. Close examination of these traps suggests that the gap between the mesh and the
upper funnel may have been too narrow, preventing bats from escaping and forcing them to
pass through the mesh into the collector. Particular care should be taken in positioning the
mesh to avoid such issues. Using a finer mesh may further reduce bat captures, though this
could also affect trap efficiency for insects. In addition, it may be preferable to use a square
mesh rather than a hexagonal one, as bats' forelimbs might get caught in the latter (A.
Chrétien, pers. obs.).

Adding a mesh could have reduced both the quantity and diversity of insects captured,
particularly larger specimens that may have escaped before falling into the collector (Pearce
et al., 2005). Nonetheless, we did not observe any significant reduction in mean species
richness associated with mesh installation. Similarly, no significant decrease in average
beetle size was found, even among larger taxa such as Cerambycidae and Cetoniinae. Only a
decreasing size trend was observed in Cetoniinae. Nonetheless, we did observe a significant
decrease in abundance between 2019 and 2021 in some trap types for Cerambycidae and
Scolytinae, with a similar trend for Buprestidae. As with bat captures, because standard and
modified traps were not used in parallel, we cannot conclusively attribute these reductions
to the mesh, especially given the known interannual variability in forest insect populations
(e.g., Garnas et al., 2023; Le Souchu et al., 2024b). It is possible though that reduced
abundance was due to mesh clogging by leaves, twigs, or other tree debris, which may have
blocked insects from reaching the collector, hence more frequent maintenance and
monitoring would likely mitigate this issue. BMF traps are commonly used for the
biosurveillance of wood-boring insects at points of entry such as ports and airports (Roques
et al., 2023). In this context, maximizing the number of individuals captured is crucial, as
exotic species are typically detected in very low numbers. Adding a mesh to MF traps may
reduce their capture efficiency and, consequently, the likelihood of detecting quarantine or
other targeted exotic species.

In conclusion, adding a 13-mm mesh to the funnel just above the collector proved to be a
simple and effective modification that reduced bat bycatch without significantly affecting
beetle diversity, although it may reduce beetle abundance. We recommend the systematic

modification of traps deployed within the tree canopy, at least within Europe, and urge trap
manufacturers to consider this issue in future trap designs.

References

Ancillotto, L., Cosentino, F., Festa, F. & Mori, E. (2024). Adhesive traps for suppressing pest
insects represent a serious threat to bats across Europe. *European Journal of Wildlife*
Research 70, 119.

Arthur, L. & Lemaire, M. (2021). Les chauves-souris de France Belgique, Luxembourg et
Suisse, 3rd ed. Muséum National d'Histoire Naturelle, Paris ; Biotope, Mèze, 592 p.

Augusto, A. M., Raposeira, H., Horta, P., Mata, V. A., Aizpurua, O., Alberdi, A., Jones, G.,
Razgour, O., Santos, S.A.P., Russo, D. & Rebelo, H. (2024). Bat diversity boosts ecosystem
services: Evidence from pine processionary moth predation. *Science of the Total*
Environment, *912*, 169387.

Bat Tree Habitat Key (2018). Bat Roosts in Trees: A Guide to Identification and Assessment
for Tree-Care and Ecology Professionals. Pelagic Publishing Ltd, 264 p.

Beilke, E. A., & O'Keefe, J. M. (2023). Bats reduce insect density and defoliation in temperate
forests: An exclusion experiment. *Ecology*, *104*, e3903.

Bouget, C., Brustel, H., Noblecourt, T., & Zagatti, P. (2019). *Les Coléoptères saproxyliques de*
France: Catalogue écologique illustré. Muséum National d'Histoire Naturelle, Paris, 744 p.

Bouget, C., Brustel, H., Brin, A., & Valladares, L. (2009). Evaluation of window flight traps for
effectiveness at monitoring dead wood-associated beetles: the effect of ethanol lure under
contrasting environmental conditions. *Agricultural and Forest Entomology*, *11*, 143-152.

Bouget, C., Brin, A., & Brustel, H. (2011). Exploring the "last biotic frontier": are temperate
forest canopies special for saproxylic beetles? *Forest Ecology and Management*, *261*, 211-
220.

Brockerhoff, E. G., Corley, J. C., Jactel, H., Miller, D. R., Rabaglia, R. J., & Sweeney, J. (2023)
Monitoring and surveillance of forest insects. In *Forest Entomology and Pathology, volume 1:*
Entomology, Allison J.D., Paine, T.D., Slippers, B., & Wingfield, M. J. (eds). 669-705.

Browning, E., Barlow, K. E., Burns, F., Hawkins, C., & Boughey, K. (2021). Drivers of European
bat population change: a review reveals evidence gaps. *Mammal Review*, *51*, 353-368.

Council Directive 92/43/EEC (1992) of 21 May 1992 on the conservation of natural habitats
and of wild fauna and flora. *Official Journal of the European Union*, *206*, 50.

Crook, D. J., Francese, J. A., Rietz, M. L., Lance, D. R., Hull-Sanders, H. M., Mastro, V. C., ... &
Ryall, K. L. (2014). Improving detection tools for emerald ash borer (Coleoptera:
Buprestidae): comparison of multifunnel traps, prism traps, and lure types at varying
population densities. *Journal of Economic Entomology*, *107*, 1496-1501.

Erasmy, M., Leuschner, C., Balkenhol, N., & Dietz, M. (2021). Three-dimensional stratification
pattern in an old-growth lowland forest: How does height in canopy and season influence
temperate bat activity? *Ecology and Evolution*, *11*, 17273-17288.

European Food Safety Authority (EFSA), Schans, J., Schrader, G., Delbianco, A., Graziosi, I., &
Vos, S. (2020). Pest survey card on *Agrilus planipennis*. *EFSA Supporting Publications*, *17*,
1945E.

Evans, H. F., Williams, D., Hoch, G., Loomans, A., & Marzano, M. (2020). Developing a
European Toolbox to manage potential invasion by emerald ash borer (*Agrilus planipennis*)
and bronze birch borer (*Agrilus anxius*), important pests of ash and birch. *Forestry: An
International Journal of Forest Research*, *93*, 187-196.

Floren, A., & Schmidl, J. (2008). Canopy arthropod research in Europe: basic and applied
studies from the high frontier. Bioform, Nuremberg, 576 p.

Garnas, J. R., Ayres, M. P., & Lombardero, M. J. (2023). Forest insect population dynamics. In
Forest Entomology and Pathology, volume 1: Entomology, Allison J.D., Paine, T.D., Slippers,
B., & Wingfield, M. J. (eds). pp. 115-140.

Greif, S., Zsebők, S., Schmieder, D., & Siemers, B. M. (2017). Acoustic mirrors as sensory
traps for bats. *Science*, *357*(6355), 1045-1047.

Hafsi, A., Abbes, K., Harbi, A., & Chermiti, B. (2020). Field efficacy of commercial food
attractants for Ceratitis capitata (Diptera: Tephritidae) mass trapping and their impacts on
non-target organisms in peach orchards. *Crop Protection*, *128*, 104989.

Holzinger, W.E., Huemer, S., Komposch, B. (2023). Window traps are a potential threat for
bats. *Hystrix Italian Journal of Mammalogy*, 34, 71-72

Jones, G., & Holderied, M. W. (2007). Bat echolocation calls: adaptation and convergent
evolution. *Proceedings of the Royal Society B: Biological Sciences, 274*(1612), 905-912.

Law, B., Park, K. J., & Lacki, M. J. (2016). Insectivorous bats and silviculture: balancing timber
production and bat conservation. Bats in the Anthropocene: conservation of bats in a
changing world, 105-150.

Le Souchu, E., Cours, J., Cochenille, T., Bouget, C., Bankhead-Dronnet, S., Braet, Y., ... & Sallé,
A. (2024a). Responses of the hyper-diverse community of canopy-dwelling Hymenoptera to
oak decline. *Insect conservation and diversity*, *17*, 430-448.

Le Souchu, E., Bouget, C., & Sallé, A. (2024b). Environmental drivers of local and temporal
variations in the community of oak-associated borers (Coleoptera: Buprestidae). *European*
Journal of Forest Research, *143*, 603-616.

Lindgren, B. S. (1983). A multiple funnel trap for scolytid beetles (Coleoptera). *The Canadian*
Entomologist, *115*(3), 299-302.

Mathews, F., Anderson, M., Coomber, F., Finch, D., Le Marquand, C., O'Malley, K., & Wright,
P. (2023). Common Pipistrelle *Pipistrellus pipistrellus* (Schreber, 1774). In *Chiroptera* (pp.
555-589). Springer International Publishing.

McLeod, E., Quamme, D., Lausen, C., Swan, F. (2021). The Incidental Capture of Bats in
Douglas-Fir Beetle Funnel Traps. NACFOR's biology intern poster.
https://www.nakuspcommunityforest.com/the-incidental-capture-of-bats-in-douglas-fir-
beetle-funnel-traps/

Müller, J., Brandl, R., Buchner, J., Pretzsch, H., Seifert, S., Strätz, C., ... & Fenton, B. (2013).
From ground to above canopy—Bat activity in mature forests is driven by vegetation density
and height. *Forest Ecology and Management*, *306*, 179-184.

Orbach, D. N., & Fenton, B. (2010). Vision impairs the abilities of bats to avoid colliding with
stationary obstacles. *PLoS One*, *5*(11), e13912.

Orlova-Bienkowskaja, M. J. (2014). Ashes in Europe are in danger: the invasive range of
Agrilus planipennis in European Russia is expanding. *Biological Invasions*, *16*, 1345-1349.

Orlova-Bienkowskaja, M. J., Drogvalenko, A. N., Zabaluev, I. A., Sazhnev, A. S., Peregudova, E.
Y., Mazurov, S. G., ... & Bieńkowski, A. O. (2020). Current range of *Agrilus planipennis*
Fairmaire, an alien pest of ash trees, in European Russia and Ukraine. *Annals of Forest*
Science, *77*, 1-14.

Pearce, J. L., Schuurman, D., Barber, K. N., Larrivée, M., Venier, L. A., McKee, J., & McKenney,
D. (2005). Pitfall trap designs to maximize invertebrate captures and minimize captures of
nontarget vertebrates. *The Canadian Entomologist, 137*, 233-250.

Plewa, R., Jaworski, T., Hilszczański, J., & Horák, J. (2017). Investigating the biodiversity of the
forest strata: The importance of vertical stratification to the activity and development of
saproxylic beetles in managed temperate deciduous forests. *Forest Ecology and*
Management, 402, 186-193.

Rassati, D., Marini, L., Marchioro, M., Rapuzzi, P., Magnani, G., Poloni, R., ... & Sweeney, J.
(2019). Developing trapping protocols for wood-boring beetles associated with broadleaf
trees. *Journal of Pest Science, 92*, 267-279.

Regnery, B., Couvet, D., Kubarek, L., Julien, J. F., & Kerbiriou, C. (2013). Tree microhabitats as
indicators of bird and bat communities in Mediterranean forests. *Ecological Indicators*, *34*,
221-230.

Rojas-Nossa, S. V., Novoa, N., Serrano, A., & Calviño-Cancela, M. (2018). Performance of
baited traps used as control tools for the invasive hornet *Vespa velutina* and their impact on
non-target insects. *Apidologie*, *49*, 872-885.

Roques, A., Ren, L., Rassati, D., Shi, J., Akulov, E., Audsley, N., ... & Millar, J. G. (2023).
Worldwide tests of generic attractants, a promising tool for early detection of non-native
cerambycid species. *NeoBiota*, *84*, 169-209.

Russ, J. M., Racey, P. A., & Jones, G. (1998). Intraspecific responses to distress calls of the
pipistrelle bat, *Pipistrellus pipistrellus*. *Animal Behaviour, 55*, 705-713.

Sallé, A., Parmain, G., Nusillard, B., Pineau, X., Brousse, R., Fontaine-Guenel, T., ... & Bouget,
C. (2020). Forest decline differentially affects trophic guilds of canopy-dwelling beetles.
Annals of Forest Science, 77, 1-21.

Sallé, A., Cours, J., Le Souchu, E., Lopez-Vaamonde, C., Pincebourde, S., & Bouget, C. (2021).
Climate change alters temperate forest canopies and indirectly reshapes arthropod
communities. *Frontiers in Forests and Global Change, 4*, 710854.

Santoiemma, G., Williams, D., Booth, E. G., Cavaletto, G., Connell, J., Curletti, G., ... & Rassati,
D. (2024). Efficacy of trapping protocols for *Agrilus* jewel beetles: a multi-country
assessment. *Journal of Pest Science*, *97*, 1795-1810.

Spears, L. R., Christman, M. E., Koch, J. B., Looney, C., & Ramirez, R. A. (2021). A review of
bee captures in pest monitoring traps and future directions for research and collaboration.
Journal of Integrated Pest Management, *12*, 49.

Sztencel-Jabłonka, A., & Bogdanowicz, W. (2012). Population genetics study of common
(*Pipistrellus pipistrellus*) and soprano (Pipistrellus pygmaeus) pipistrelle bats from central
Europe suggests interspecific hybridization. *Canadian Journal of Zoology*, *90*, 1251-1260.

Sukovata, L., Dziuk, A., Plewa, R., & Jaworski, T. (2022). The effect of trap color on catches of
Monochamus galloprovincialis and three most numerous non-target insect species. *Insects*,
13, 220.

Tanalgo, K. C., Cruz, K. C. D., & Russo, D. (2025). Susceptibility of bats to ecological and
evolutionary traps. *Biological Conservation*, *305*, 111110.

Temple, H. J., & Terry, A. (2009). European mammals: Red List status, trends, and
conservation priorities. *Folia Zoologica*, *58*(3), 248.

Tillon, L., Bouget, C., Paillet, Y., & Aulagnier, S. (2016). How does deadwood structure
temperate forest bat assemblages?. *European Journal of Forest Research*, *135*(3), 433-449.

Ulyshen, M. D. (2011). Arthropod vertical stratification in temperate deciduous forests:
implications for conservation-oriented management. *Forest Ecology and Management*,
261(9), 1479-1489.

Urban-Mead, K. R., Muñiz, P., Gillung, J., Espinoza, A., Fordyce, R., van Dyke, M., ... &
Danforth, B. N. (2021). Bees in the trees: Diverse spring fauna in temperate forest edge
canopies. *Forest Ecology and Management*, *482*, 118903.

Vincent, A., Tillier, P., Vincent-Barbaroux, C., Bouget, C., & Sallé, A. (2020). Influence of forest
decline on the abundance and diversity of Raphidioptera and Mecoptera species dwelling in
oak canopies. *European Journal of Entomology*, *117*.

Zumr, V., & Starý, P. (1991). Effects of baited pitfall traps (*Hylobius abietis* L.) on non-target
forest insects. *Journal of Applied Entomology*, *112*, 525-530.

499