**Title**

Deciphering Intercellular Communication in the Cerebellar External Granule Layer: Insights

into Non-Classical Connections in Neural Development

**Authors**

Malalaniaina Rakotobe[1,2*], Shiyu Liu[1,3], Garima Virmani[1], Gabriel Kaddour[4], Nathaly Dongo

Mendoza[1], Frédéric Doussau[5], Jean Livet[4], Laurence Cathala[2], Chiara Zurzolo[1*]

**Affiliation**

[1]Membrane Traffic and Pathogenesis, Institut Pasteur, Université Paris Cité, CNRS UMR

3691, Paris, France.

[2]Sorbonne Université, Institut du Cerveau - Paris Brain Institute - ICM, Inserm, CNRS,

APHP, Hôpital de la Pitié Salpêtrière, Paris, France

[3]Structure et Dynamique des Systèmes Vivants, Université Paris-Saclay, 91190 Gif-sur-

Yvette, France

[4]Sorbonne Université, INSERM, CNRS, Institut de la Vision, Paris 75012, France

[5]Institut des Neurosciences Cellulaires et Intégrative, Université de Strasbourg, CNRS UPR

3212, Strasbourg, France

Corresponding authors: chiara.zurzolo@pasteur.fr (Chiara Zurzolo),

malalaniainarakotobe@gmail.com (Malalaniaina Rakotobe)

<u>**Abstract**</u>

Intercellular communication is essential for brain development. While classical modes—

paracrine, juxtacrine, and synaptic signaling—are well characterized, emerging evidence

suggests that membranous bridges, such as tunneling nanotubes (TNTs), forming de novo

between cells and mainly described *in vitro*, may also contribute. Yet their presence and

function in vivo remain unclear, partly due to the difficulty of distinguishing them from other

intercellular connections (ICs). Building on connectomic observations in fixed tissue

reporting ICs in the external granule layer (EGL) of the developing cerebellum, we examined

their nature in postnatal day 7 (P7) mice. Using immunofluorescence, sparse genetic labelling,

[1] The present address of M.R is at NBX Biosciences, 34000 Montpellier, France; G.V is at Instytut Biologii Doświadczalnej im. M. Nenckiego PAN, 02-093 Warszawa, Poland; N.D.M is at The EKFZ for *Digital* Health, 01307 Dresden, Germany; L.C is at University of Colorado Anschutz Medical Campus, 80045 CO, USA

and live imaging, we distinguished division-independent ICs from cytokinetic bridges (CBs)

and intercellular bridges (IBs). CBs were detected in the EGL, whereas IBs were not observed

in this region. In addition to CBs, we identified membranous protrusions which appeared to

link clonally and non-clonally related cells. The presence of these ICs reinforces previous

connectomic evidence in fixed tissue and raises the possibility that they may participate in

intercellular communication during cerebellar development. These findings warrant further

investigation into a potentially underexplored mode of communication in the developing

brain.

**Keywords**

Intercellular communication, intercellular connections (ICs), tunneling nanotubes (TNTs),

cytokinetic bridges (CBs), intercellular bridges (IBs), cerebellum development.

**Introduction**

Intercellular communication plays a crucial role in brain development, ensuring the precise

regulation and coordination of processes including neurogenesis, cell migration,

differentiation and the establishment of functional neuronal networks. This communication

occurs through adhesion molecules, the secretion of extracellular vesicles and soluble

molecules (Bahram Sangani et al., 2021; Sahay et al., 2020), as well as through synaptic

interactions via electrical and chemical synapses (Pereda, 2014). Disruptions in these

pathways can lead to neurodevelopmental disorders, highlighting their significance in brain

maturation.

Beyond classical mechanisms, recent studies have identified additional structures involved in

intercellular communication, including cytonemes, intercellular bridges (IBs), and tunneling

nanotubes (TNT) (Korenkova et al., 2020). Cytonemes are thin, F-actin-based, close-ended

protrusions (1–700 µm in length) that mediate signal transduction between cells via

membrane-tethered receptors and ligands, without establishing cytoplasmic continuity (Hall et

al., 2024). IBs (0.4–350 µm in length) arise from incomplete cytokinesis and allow

cytoplasmic exchange between connected cells (Haglund et al., 2011; Rujano et al., 2022).

Unlike cytokinetic bridges (CBs), which are abscised at the end of cytokinesis, IBs persist

longer, facilitating sustained communication between connected cells. TNTs, which can

extend up to 200 μm, are F-actin based, open-ended membranous structures, formed *de novo*

between two distinct cells. They facilitate the transfer of cargo such as organelles, proteins,

and nucleic acid. The open-ended nature of TNTs can be assessed using high-resolution

electron microscopy (EM) or live imaging which demonstrates uninterrupted transfer of large

cargo such as organelles (Sáenz-de-Santa-María et al., 2023). However, these techniques are

difficult to apply to tissue samples. Identifying TNTs in *vivo* is particularly challenging due to

the lack of specific molecular markers, the fragility of these structures, and their sensitivity to

fixatives like paraformaldehyde, which can compromise structural preservation.

Consequently, such structures are typically referred to as "TNT-like" in tissue until their

ability to transfer cellular cargos between connected cells is confirmed (Palese et al., 2025).

Due to these hurdles, TNTs have been extensively studied *in vitro*, but their *in vivo*

characterization remains limited, with only recent evidence confirming functional TNTs in the

zebrafish embryo (Korenkova et al., 2024).

In the developing cerebellum, the external granule layer (EGL) constitutes a dynamic region

where granule cells (GCs), the most numerous neurons of the brain, are generated. The EGL

is subdivided into an outer proliferative zone (oEGL) and an inner differentiating zone

(iEGL), as originally defined by Altman & Bayer (Altman & Bayer, 1997). In rodents,

granule cell progenitors (GCPs) proliferate in the oEGL for approximately three weeks before

differentiating into postmitotic GCs, which migrate through the iEGL and molecular layer

(Curran & O'Connor) to reach the inner granular layer (IGL) (Consalez et al., 2020).

Intercellular communication within the EGL is essential for coordinating interactions not only

between GCs but also with other neuronal populations (Kim et al., 2023). Combining serial-

section scanning electron microscopy (ssSEM) with computational connectomics in fixed

tissue, we recently identified open-ended intercellular connections (ICs) linking immature

GCs in the mouse EGL. By analyzing cellular parameters such as shape, Golgi apparatus

number, and cilium presence, our data suggested that these structures could represent either

stable IBs or TNTs. However, their precise nature and function could not be precisely

determined since the ssSEM  approach did not allow the use of specific markers to distinguish

IBs from other types of cellular connections (Cordero Cervantes et al., 2023).

In this study, we examined the presence and characteristics of ICs in the EGL of postnatal day
7 (P7) mice. Using immunofluorescence, live imaging, and sparse genetic labelling, we
differentiated cytokinetic bridges (CBs) and intercellular bridges (IBs) from other connections
that appeared to span fully between cells. While CBs were identified in the iEGL, our data
indicate that they do not account for all previously described ICs (Cordero Cervantes et al.,
2023), suggesting that other types of connections—such as TNT-like structures—may also
occur during cerebellar development.

## **Materials and methods**

### **Animals**

All mice used were generated and maintained in the animal facility of the Institut Pasteur,
Paris. The care and handling of the animals prior or during the experimental procedures
followed European Union rules and were approved by the French Animal Protection
Committee and the local Ethics Committees. Animals were maintained on 12h light/dark
cycles with food and water provided *ad libitum*. C57BL/6 mice at P7 were used for the
quantification of CBs and mitotic cells in the iEGL. Rj:SWISS mice were used for the
electroporation of the *pCAG-Dendra2-E2A-EGFP-Cep55* plasmid.

### **Transgenic mouse lines**

We used a *CAG-Cytbow* reporter line bearing a *Cytbow* MAGIC Marker transgene (Loulier et
al., 2014) that stochastically expresses mTurquoise2, eYFP or tdTomato under the broad *CAG*
promoter in a Cre-dependent manner (this line will be described elsewhere). To generate
experimental animals, we crossed homozygous *CAG-Cytbow*$^{+/+}$ males with *Nestin-CreER*$^{T2+/-}$
females (Lagace  et al., 2007) to obtain *Nestin-CreER*$^{T2+/-}$; *CAG-Cytbow*$^{+/-}$ offspring. To
increase the probability of combinatorial fluorescent protein expression of in a Cre-dependent
manner, these *Nestin-CreER*$^{T2+/-}$; *CAG-Cytbow*$^{+/-}$ females were backcrossed with *CAG-*
*Cytbow*$^{+/+}$ males to generate *Nestin-CreER*$^{T2+/-}$; *CAG-Cytbow*$^{+/+}$experimental animals.

### **Tamoxifen injection**

To sparsely label neural cells, *Nestin-CreER$^{T2+/-}$; CAG-Cytbow$^{+/+}$* newborn mice received

subcutaneous injection of 25 µL of tamoxifen (234-118-0, Merck) diluted in corn oil once per

day at P0, P1, P2 and P3 at a concentration of 5 mg/mL.

**Plasmid**

119  Coding sequence of *pCAG-Dendra2-E2A-EGFP-Cep55* construct was cloned from *pDEST-*

120  *Dendra2-E2A-EGFP-Cep55* construct (used in Korenkova et al., 2024) into the *pCAG-GFP*

121  plasmid from Addgene (Plasmid #11150).

**Postnatal electroporation**

Vetergesic, at a concentration of 0.05 mg/kg, was injected 30 minutes prior to surgery for

analgesia. Anesthesia was induced with 4% isoflurane for 10 minutes and then maintained at

2% throughout the procedure. Before making the incision, laocaine at a concentration of 2.5

mg/kg was injected at the incision site to alleviate pain. To prevent desiccation, the eyes were

protected with Ocry-gel. Following the craniotomy, the *pCAG-Dendra2-E2A-EGFP-Cep55*

plasmid was injected at the surface of the cerebellum using a glass micropipette. The plasmid

was delivered at 4–5 sites for a total injection volume of 1 µl at a concentration of 5.9 µg/µl.

Triple electrodes coated with conductive gel were used for electroporation. The electrode

connected to the anode was placed on a micromanipulator and lowered into contact with the

craniotomy under optical control, while the two cathode electrodes were positioned on either

side of the animal's head. Electroporation (BTX electroporator) was performed with the

following parameters: 5 pulses of 50 ms at 1 Hz, at 80V. Immediately after electroporation,

the mouse was placed on a heating pad and stimulated to restart respiration in case of

cardiorespiratory arrest. The incision was sealed with Vetbond surgical glue, and a

subcutaneous injection of 20-30 µl of 0.9% NaCl was administered gently to rehydrate the

mouse.

**Live imaging of acute cerebellar slices**

This protocol was adapted from a Star Protocol by Qu. X et al. (Qu et al., 2021). Briefly, pups

were decapitated and the brain dissected out in oxygenated, ice-cold complete HBSS (cHBSS)

buffer. The brain was then transferred to the slicing chamber of a vibratome (Leica Biosystem

VT 1000S), which contained cHBSS to keep the brain submerged, while oxygenation is

maintained in the chamber. Cerebellar slices (250 µm thick) were prepared using the

vibratome at a speed of 0.08–0.1 mm/s and collected in a "nest beaker" containing complete

cHBSS medium, bubbled at 33°C. Cerebellar slices expressing fluorescent reporters were

selected and transferred to a µ-Slide 8 Well (ibidi, Cat. No: 80807) with 1 mL of warm, fresh,

sterile medium. The ibidi well were placed in a temperature-controlled chamber attached to

the microscope, where both temperature (37°C) and CO□ flow (5%) were regulated and

maintained to promote cell viability and ensure optical stability during live imaging.

| Component of the complete HBSS | Stock concentration | Final concentration | Volume |
|---|---|---|---|
| HBSS (Thermo Fisher, 14065056) | 10x | 1x | 50 mL |
| HEPES (pH 7.4) (Thermo Fisher, 15630080) | 1 M | 2.5 mM | 1.25 mL |
| D-Glucose (Sigma-Aldrich, G8270) | 1 M | 30 mM | 15 mL |
| CaCl2 (Sigma-Aldrich, C5670) | 100 mM | 1 mM | 5 mL |
| MgSO4 (Sigma-Aldrich, M2643) | 100mM | 1mM | 5 mL |
| NaHCO3 (Sigma-Aldrich, 144-55-8) | 1 M | 4 mM | 2 mL |
| ddH2O | | | 421.75 mL |
| Total | | | 500 mL |
| Sterile filter with a 0.2 µm filter. Store at 4°C. | | | |

## Immunohistochemistry

Brains were dissected out and fixed overnight in 4% paraformaldehyde (PFA) at 4°C, then

rinsed in PBS and cryoprotected in 30% sucrose in PBS at 4°C overnight before embedding

and freezing in OCT (Cryomatrix™, Thermo) at −80°C. The cerebellum was sliced in sagittal

sections of 40 µm using a Leica CM 3050S cryostat. Sections were then mounted on

SuperFrost+ slides (Fisher Scientific). Slices underwent an antigen unmasking step: they were

incubated in unmasking solution (citrate buffer, pH 6) for 15 minutes in the 2100 Antigen

Retriever. They were then cooled in PBS for 10 minutes before being permeabilized for 10

minutes in PBS containing 0.5% Triton. Next, slices were incubated for 1 hour in blocking

solution (PBS with 5% fetal calf serum (FCS) and 2% bovine serum albumin (BSA)). To

quantify CBs and mitotic cells, two rabbit-derived primary antibodies were used sequentially

(the rabbit anti-Aurora B kinase antibody and the rabbit anti-Ki67 antibody). For those

samples a two-step IHC was run. First, slices were immuno-labelled with the rabbit anti-

Aurora B primary antibody before running a second IHC with the other antibodies including

the rabbit anti-Ki67 antibody. The first Rabbit antibody was included in the blocking solution

at room temperature for 2 hours. After PBS washes, the slices were incubated overnight at

4°C with a Fab fragment secondary antibody in blocking solution. Then the second step of the

IHC procedure was the same for all samples. Primary antibodies were incubated overnight at

4°C in blocking solution. The next day, secondary antibodies were incubated for 2 hours at

room temperature in blocking solution. Finally, sections were incubated with DAPI (1 µg/ml)

in PBS for 30 minutes and mounted using Fluoromount-Gold. Primary antibodies used were:

rabbit anti-Aurora B kinase (Abcam, ab2254, 1:500), rabbit anti-Ki67 (Abcam, ab15580,

1:200), goat anti-TAG1 (rndsystems, AF4439, 1:200), mouse anti-citron kinase (BD

Biosciences, 611376, 1:200), goat anti-anillin (Abcam, ab5910, 1:50), rabbit anti-anillin

(Bethyl laborories, A301-406A, 1:500), chicken anti-GFP (Aves Lab, GFP-1010, 1:1000),

goat anti-tdTomato (Sicgen, AB8181, 1:200), rat anti-PH3 (Abcam, ab10543, 1:200).

Secondary antibodies used (Jackson ImmunoResearch, 1:500) were: Fab Fragment anti-rabbit

Alexa Fluor™647, anti-rabbit Alexa Fluor™594, anti-rabbit Alexa Fluor™647, anti-goat

Alexa Fluor™488, anti-mouse Alexa Fluor™488, anti-goat Alexa Fluor™647, anti-chicken

488, anti-goat Alexa Fluor™594, anti-rat Alexa Fluor™488.

**Microscopy and image analysis**

*Imaging*

Images of cerebellar sections were acquired on a Zeiss LSM900 confocal microscope, except

for a subset of samples electroporated with the *pCAG-Dendra2-E2A-EGFP-Cep55* plasmid

which were imaged on a Nikon Ti2E Spinning disk microscope. In addition, a Leica SP8

confocal microscope was used to image the 4 fluorescent proteins (eBFP2, mTurquoise2,

eYFP, tdTomato) expressed by *Nestin-CreERT2+/-; Cytbow+/+* mice.

The Zeiss LSM900 was equipped with a 63x/1.4 NA oil-immersion objective (**Fig. 1, 2, 4,**

**S4E**) or a 10x/0.45 NA DRY objective (**Fig. 3B, S3 and S4D**), as indicated on the figure

legends. Images were acquired at 16-bit depth, with a pixel size of 0.1 µm and a z-step size of

0.35 µm. The following fluorochromes and detection settings were used: DAPI (excited at
405 nm, emission detected at 415-480 nm), Alexa Fluor 488 (488 nm, 500-550 nm), Alexa
Fluor 594 (561 nm, 580-620 nm), and Alexa Fluor 647 (633 nm, 640-700 nm).

The Nikon Ti2E inverted microscope was equipped with a Yokogawa CSU-W1 spinning disk
unit and a Photometrics Prime 95B back-illuminated sCMOS camera. A 60x/1.4 NA oil-
immersion objective was used (**Fig. 3C, 3F**). Images were collected in 12-bit mode, with a
pixel size of 0.11 µm and a z-step of 0.3 µm. Standard bandpass filters optimized for each
fluorophore were used including 447/60 for DAPI and 525/50 for AF488.

The Leica SP8 confocal microscope was used with either a 20x/0.75 NA water immersion
objective (**Fig. S4B**) or a 63x/1.4 oil-immersion oil objective (**Fig. S4C**), as indicated in the
figure legends. Images were collected in an 8-bit mode for low-magnification tile scans (**Fig.
S4B**) and in 16-bit mode for high resolution acquisition (**Fig. S4C**). For the 20x objective, the
pixel size was 1.14 µm with a z-step of 3.8 µm. For the 63x objective, images were single
tiles with a pixel size of 0.09 µm and a z-step of 0.35 µm. A 405 nm laser and white laser
(470-670) were used to excite the following fluorophores: eBFP2 (excited at 405 nm,
emission detected at 415-465 nm), mTurquoise2 (excited at 470 nm, emission detected at 480-
530 nm), mEYP (excited at 514 nm, emission detected at 524-574 nm), tdTomato (excited at
554 nm, emission detected at 564-604 nm).

All image quantifications were performed on unprocessed datasets. Brightness and contrast
adjustments applied to figures were linear and applied across entire images, solely to enhance
visualization of thin TNT-like structures.

*Quantification of mitotic cells*

Connected nuclei were assigned to one of three Ki67 pattern classes based on the nuclear
distribution of the Ki67 signal: (i) Ki67$^{mitotic}$, showing a dense, compact/perichromosomal
pattern characteristic of mitotic cells; (ii) Ki67$^{interphase}$, displaying a punctate or dotted nuclear
pattern typical of interphase; and (iii) Ki67$^{G0}$, lacking detectable nuclear Ki67 signal. These
categories follow published descriptions of Ki67 dynamics across the cell cycle (Sobecki et
al., 2016; Sun & Kaufman, 2018). Scoring was performed manually on z-stacks using DAPI
as a nuclear reference. For each image, only mitotic cells connected by a CB were quantified.

Quantification of CBs and mitotic cells in the iEGL were normalized to the total number of

nuclei within the EGL to allow comparison with our previous study (Cordero Cervantes et al.,

2023). Nuclei were segmented using a multi-step computational pipeline combining deep

learning and classical segmentation. First, a U-Net convolutional neural network was trained

to segment nuclei in 3D confocal stacks (**Fig. S2A**). The resulting binary masks were then

used to remove background signals.

Next, nuclear detection was refined in 2D using Cellpose (Stringer et al., 2021) which proved

more reliable than the previously used watershed algorithm (**Table 1**). To estimate the number

of nuclei in 3D, Cellpose-based 2D counts were corrected using a scaling factor (F$_i$) (F$_i$ =

0.092) determined by comparing automatic 2D counts to manual 3D annotations on

representative stacks. The correction factor Fi  for each region was calculated as below and

applied uniformly to all images:

$$Fi \ = \ \frac{Manual\ 3D\ count}{Cellpose\ 2D\ count} \ \ ....\ (Eq\ 1)$$

| Sample | Region | Cellpose 2D count | Manually count | | Relative error (%) | Correction Factor (Fi) (according to Eq. 1) |
|---|---|---|---|---|---|---|
| | | | 2D | 3D | | |
| **Sample 1** | Region n°1 | 547 | 503 | 56 | 8.75% | 0.11 |
| **Sample 2** | Region n°2 | 735 | 692 | 62 | 6.21% | 0.09 |
| **Sample 3** | Region n°3 | 695 | 668 | 62 | 4.04% | 0.09 |
| | | | | Average | 6.33 % | 0.092 |

*Table 1. Estimation of a 2D-to-3D correction factor for Cellpose-based nucleus counting. Correction factors*
*were derived by comparing Cellpose 2D counts to manual annotations of the same slice (Manual 2D count) and*
*to the full Z-stack (Manual 3D Count). Relative Error quantifies the discrepancy between Cellpose 2D and*
*manual 2D counts. The average Correction Factor (F$_i$)=0,092 was then used to extrapolate total 3D counts*
*from Cellpose 2D count across the full dataset.*

*Nuclei segmentation*

Nuclei segmentation with the U-Net CNN (VGG-16 encoder) involved four sequential
steps: data preparation, learning, testing, and inference. The DAPI channel was used as input;
slices were rescaled (0-255), equalized, and manually annotated in VAST (Berger et al., 2018)
to generate ground-truth masks.

Training data were split into overlapping 1x320x320-pixels tiles (Z, X, Y), with a
stride of 1x270x270 pixels. Pixel dimensions were calibrated to 0.35 µm/pixel in X and Y
axes, and the Z-spacing between slices was 3.5 µm. All image tiles were rescaled before
training and split randomly into training (80%) and test (20%) sets.

Augmentations included random rotations, flips, ±10% translations, and ±20% contrast
adjustments. The U-Net model was trained for 40 epochs using the Adam optimizer (initial
learning rate: $1 \times 10^{\square\square}$, batch size 2) and selected based on the best intersection-over-union
(IoU) score.

During inference, the trained model was applied to the full 3D volumes using a sliding-
window approach with the same tile size and overlap used during training. For each tile,
probability maps were thresholded at 0.5 to produce binary masks, stitched, and post-
processed by hole filling and border-object removal to refine the final segmentation. The
resulting masks were used to isolate nuclei and eliminate background signals.

*EGL and iEGL classification*

The iEGL was delineated using a CNN-based classifier (was the same as described in
our previous paper (Cordero Cervantes et al., 2023)) trained on three-channel inputs (DAPI,
Ki67, TAG1; **Fig. S2B**). Input volumes were split into non-overlapping patches of 100×100
pixels (X, Y) (0.35 µm/pixel). The network comprised three convolutional blocks with
increasing filter sizes (16, 32, 64), each followed by max-pooling, and a dense layer of 128
units with dropout (rate 0.2). Input images were resized to 100x100 pixels, and pixel values
were normalized to the [0, 1] range. The training dataset consisted of randomly chosen and
manually labeled iEGL and Background tiles. A total of 5488 random tiles were used for
training and 1372 for validation. To enhance model generalization, data augmentation was
applied during training, including random horizontal flips, ±10% rotations, and ±10% zoom.

For iEGL class after filtering: 2658 tiles (Training: 2126, Test: 532)

For Background class after filtering: 8404 tiles (Training: 6723, Test: 1681)

The final model achieved F1-scores of 0.99897541 for iEGL and 0.99833055 for background

on the test data. During inference, patches with probability > 0.5 for iEGL and < 0.6 for

background were classified as iEGL, and binary masks were reconstructed into full 2D maps.

The resulting classified patches were reconstructed into binary masks delineating the iEGL.

To generate the full EGL mask, the post-processed iEGL mask was expanded by padding,

with the padding direction (left, right, up, down, right-up, or left-down) manually selected

depending on the anatomical orientation of the EGL in the image (**Fig. S2C**).

The number of nuclei in the iEGL was determined by segmenting nuclei in 2D and counting

those that overlapped with the iEGL mask. To estimate the corresponding 3D nuclear count,

the 2D count was multiplied by the previously derived scaling factor $F\square=0.092$ (**Fig. S2D**).

The same approach was used to quantify the nuclei in the EGL (**Fig. S2D'**).

*Image analysis and quantification*

Analyses in **Fig. 1** and **Fig. 2** were performed on sections from the medial vermis (lobule

VIII), consistent with our previous study (Cordero Cervantes et al., 2023). Analyses in **Fig. 3**

and **Fig. 4** were conducted across the entire cerebellum, although labelled cells were

predominantly located in lobule IX and X. 2D images were processed with Fiji (ImageJ). 3D

renderings of acquired z-stacks were generated using Imaris® software (version 10, Oxford

Instruments). TNT-like connections were identified based on morphological criteria: a

membranous protrusion was counted as a TNT-like connection if it appeared continuous, did

not express CBs or IBs markers, and connected two labelled cells. The total number of

labelled cells was counted manually using the Spots function in Imaris. The length of every

TNT-like connection was measured using the Measurement points tool.

All quantifications were performed on confocal images acquired from multiple cerebellar

sections per mouse. Counts of labeled structures were normalized per image to the total

number of DAPI$\square$ nuclei within the relevant region (EGL or iEGL), defined using CNN- and

Cellpose-based segmentation for CBs and mitotic cells, or by manual counting of the total

number of labeled cells for TNT-like connections. For **Fig.1** and **Fig.2,** per-image percentages

were averaged across all images from each mouse to obtain a mean value per animal. For **Fig.**

**4,** the percentage of TNT-like connections was calculated by summing all connections and

labeled cells across images for each mouse, then averaging these values across all mice.

Percentages within categories (Ki67 classes, Aurora B☐ and/or anillin☐ bridges) were

calculated per image and averaged per mouse as described above. N refers to the number of m

ice analyzed, and data are reported as mean ± SEM across biological replicates.

Statistical analysis was performed using Microsoft Excel and GraphPad Prism. To compare

relative numbers of CBs and mitotic cells Student's T-test, as the variance was equal and the

data were normally distributed. Illustrations of the quantifications are displayed as violin plot,

dot plot, bar, and pie chart.  Each experiment was performed in at least 3 biological replicates.

**Results**

<u>Cytokinetic bridges (CBs) are present throughout the EGL but less frequent in the iEGL</u>

To assess the presence of mitotic cells and CBs in the EGL in P7 mice, we labeled mitotic

cells with an anti-Ki67 antibody and CBs with an anti-Aurora B kinase antibody (**Fig. 1A,**

**S1A**). Expression of the Transient Axonal Glycoprotein TAG-1, together with Ki67, was used

to distinguish the iEGL from the oEGL through a CNN-based approach that automatically

generated a mask of the iEGL region. In the oEGL, mitotic cells were found in high density

and displayed a distinct and robust Ki67 expression pattern (**Fig. 1A**) (Sobecki et al., 2016;

Sun & Kaufman, 2018) consistent with previous reports identifying this layer as the primary

proliferative zone where cell division is most frequent (Legué et al., 2016; Nakashima et al.,

2015)—and consequently CB formation. The iEGL contrasted with a sparse distribution of

mitotic cells (**Fig. 1A, B, white arrowheads**), indicating that only a small number of GCs

continue to divide within this layer. This was corroborated by anti-Aurora B kinase labelling,

which presence is indicative of CB at the end of mitosis (**Fig. 1A, 1C** yellow arrowhead) with

a progressive decrease in CB number toward the ML (**Fig. 1C0**) (N = 4 mice; 55 images, 615

CBs quantified across the EGL). As expected, their proportion in the iEGL was also lower

compared to the oEGL (**Fig. 1C**). These findings align with previous studies characterizing

the EGL as a dynamic proliferative zone (Cordero Cervantes et al., 2023; Hanzel et al., 2019).

Moreover, our quantification of mitotic cells is consistent with our previous finding, despite

different methodological approaches (Cordero Cervantes et al., 2023) (**Fig. S1A**).

To determine whether the previously identified intercellular connections (ICs) (Cordero

Cervantes et al., 2023) could be fully accounted for by CBs, we replicated our previous

analysis by quantifying mitotic cells (prophase, prometaphase, metaphase, anaphase, and telophase) and Aurora B kinase–positive CBs in the iEGL using immunofluorescence, and compared these values with those previously obtained using an ssSEM-based approach (**Fig. 1D, S1A, S1B**). All quantifications were normalized to the total number of nuclei within the EGL, obtained from CNN- and Cellpose-based segmentation of the DAPI channel (see **Fig. S2** and Methods). Immunolabeling marked similar percentages of mitotic cells (0.69%) and CBs (0.66%) in the iEGL, consistent with ongoing mitotic activity in this layer and suggesting that the labeled structures likely represent transient post-mitotic intermediates. However, the frequency of CBs obtained with this approach was substantially lower than the overall frequency of ICs (3.76%) previously reported in our earlier study using ssSEM (Cordero Cervantes et al., 2023) (see **Fig. S1A** for a comparison), which suggests that additional types of connection, such as IBs or TNT-like connections, may also be present in the iEGL.

<u>The EGL exhibited no detectable intercellular bridges (IBs)</u>

To assess whether IBs are present in the iEGL at P7, we performed quantitative immunofluorescence to evaluate the proliferative state of GCs connected by Aurora B kinase-positive bridges. Cell-cycle classification was based on Ki67 nuclear distribution patterns rather than absolute intensity (see Methods), enabling discrimination among 3 stages (**Fig. 2A**) : mitosis ($Ki67^{mitotic}$), interphase ($Ki67^{interphase}$), and quiescence ($Ki67^{G0}$) (Sobecki et al., 2016; Sun & Kaufman, 2018). Connected GCs were nearly equally distributed between the "$Ki67^{mitotic}$" (51%) and "$Ki67^{interphase}$" (49%) classes (**Fig. 2B**). In most cases, both connected cells exhibited comparable Ki67 patterns, consistent with their shared mitotic origin. Occasionally, asymmetric Ki67 labeling was observed within a pair, likely reflecting early post-mitotic divergence or initiation of cell-cycle exit in one daughter cell. However, Aurora B kinase-positive CBs were never observed between pairs of "$Ki67^{G0}$" GCs (**Fig. 2A, yellow arrowhead, and 2B**)**.** These findings indicate that CBs can persist transiently following mitotic exit, but do not remain in quiescent cells, consistent with the rapid down-regulation of Ki67 expression within 1-2 days after entering the G0 phase (Miller et al., 2018; Sobecki et al., 2016).

Aurora B kinase has been extensively used as a marker for CBs, however its expression in IBs remains uncharacterized. To further differentiate stable IBs from CBs, we labelled  midbody-containing bridges using an anti-citron kinase antibody, which marks midbodies in CBs but

not mature IBs (Price et al., 2023) and an anti-anillin antibody, previously reported to label

IBs in the Drosophila larval brain (Haglund et al., 2011), (**Fig. 2C, S1B**). We found that

81.33% of the bridges were labelled with both Aurora B kinase and anillin (**Fig. 2c$_1$ and 2D**),

2.78% were only labelled with Aurora B kinase (**Fig. 2c$_2$ and 2D**), and 15.89% were only

labelled with anillin (**Fig. 2c$_3$ and 2D**) (N = 4 mice, 66 images). The absence of anillin or

Aurora B kinase in those two groups may reflect differences in the timing or levels of protein

expression, potentially making them undetectable at certain stages of cytokinesis (Halcrow et

al., 2022; Renshaw et al., 2014). Importantly, all 3 groups express the midbody protein citron

kinase suggesting that all those ICs are CBs (**Fig. 2c$_1$-c$_3$ and 2D**). While we cannot fully

exclude the presence of IBs due to the lack of well-characterized markers in somatic cells, our

data do not support the presence of IBs in the EGL. Together, these findings indicate that CBs

alone cannot account for all the ICs previously identified in the EGL (Cordero Cervantes et

al., 2023). Moreover, differences in their distribution and molecular profiles relative to those

earlier observations suggest the existence of additional forms of intercellular connectivity.

This prompted us to investigate whether other forms of connections could be detected in the

developing cerebellum.

<u>Putative TNT-like connections are present in the developing cerebellum</u>

Having established that the ICs observed in the EGL cannot all be accounted for by CBs or

IBs, we next attempted to better characterize these structures (**Fig. S1B**). We developed an

approach to visualize fine structures between sparsely labelled GCs. P5 pups were

electroporated to transfer a *pCAG-Dendra2-E2A-GFP-Cep55* plasmid in GCs progenitors

(**Fig. S3**), leading to cytoplasmic expression of a fusion protein comprising Dendra2, GFP and

the midbody marker CEP55. Pups were sacrificed 2 days after electroporation (Korenkova et

al., 2024).

Cerebellar slices were immunostained with anti-GFP to enhance the endogenous cytoplasmic

GFP and anti-anillin to visualize CBs (**Fig. 3A-3C**). We used anti-anillin rather than anti-

Aurora B kinase, as it identifies a greater number of CBs (**Fig. 2D**). Anti-citron kinase and

anti-Cep55 antibodies were not targeted because their punctate staining made specific

labelling difficult to distinguish from background. In several cases, we observed pairs of cells

that appeared to be linked by a putative bridging structure or a cellular protrusion, compatible

with the presence of an IC, although we can not exclude the possibility that the labeled GC

processes were merely in close contact with neighboring cell. Some of these potential
connections were marked by Anillin, identifying them as CBs. However, among the putative
GC connections visualized by GFP staining, 77% did not express anillin (**Fig. 3D**; N = 3
mice, 83 images, 36 unlabeled connections and 11 CBs), suggesting that the majority of them
are unlikely to be CBs and therefore may correspond to putative TNT-like structures. This
proportion closely matches that reported in gastrula zebrafish embryo, where TNT-like
connections, which were also compared to CBs, account for 81% of ICs (Korenkova et al.,
2024). However, in contrast to zebrafish, where most TNT-like connections exceeded 5 µm in
length, allowing length-based criteria to distinguish CBs/IBs from TNT-like connections, we
observed only a modest size difference. Although some putative TNT-like connections
extended beyond 5 µm, their mean length was 5.38 µm, compared to 2.92 µm for CBs (**Fig.
3E**), indicating that connection type could not be reliably distinguished by length alone. To
directly observe their formation, we performed time-lapse imaging on acute brain slices,
capturing a putative TNT-like connection developing via a filopodia-driven mechanism,
similar to previously described *in vitro* (Korenkova et al., 2020) (**Fig. 3F-3F', S3B**). We
observed protrusions extending from each cell (white arrowheads), with an apparent
connecting bridge between the two cells fully forming within 20 minutes and persisting for at
least 10 minutes. This temporal persistence suggests that they are not filopodia, which are
much more dynamic structures that protrude and retract within seconds. However, in the
absence of functional evidence for cargo exchange, we cannot rule out the possibility that
some of the observed ICs correspond to close-ended cell processes or to cytonemes
(Korenkova et al., 2020), which are commonly observed during development in *Drosophila*
and zebrafish. To assess their open-ended nature and functionality, we attempted to
photoconvert the cytoplasmic protein Dendra2 in a single cell and monitor its transfer to the
connected cell. However, due to the close apposition of the two cells and light refraction
caused by tissue thickness, photoconversion resulted in Dendra2 activation in both cells,
precluding definitive interpretation.

Characteristics of putative TNT-like connections in the EGL from brain slice

To complement the experimental description of those connections and determine whether they
occur between clonally or non-clonally related GC, we used a Brainbow-derived strategy.
*Nestin-Cre$^{ERT2+/-}$* mice in which tamoxifen induces CRE expression only in neural progenitors

were crossed with *Cytbow*[+/+] mice (Loulier et al., 2014) in which mTurquoise, eYFP and

tdTomato are stochastically expressed following CRE-mediated recombination. Tamoxifen

was administered from P0 to P3 (**Fig. 4A**; see Methods) and resulted in stochastic expression

of one of these three fluorescent proteins in GCs, or combinations of two due to the

homozygous *Cytbow*[+/+] genotype (**Fig. S4A**). Since these fluorescent reporters are genetically

encoded and stably inherited through cell division, they enabled inference of putative clonal

relationships among GCs based on color. Although, in principle, neighboring non-clonally

related cells could express the same color, sparse labeling minimized this likelihood.

Accordingly, cells sharing the same color were considered putatively clonally related, whereas

cells labeled with different colors were classified as non-clonally related, as they cannot

originate from the same cell division (**Fig. 4A**). Pups were sacrificed at P7. Similar to the

*pCAG-Dendra2-E2A-EGFP-Cep55* plasmid electroporation approach, this strategy enabled

sparse labelling of GCs and the identification of putative connections in the densely populated

EGL, particularly in lobule X (**Fig. S4B**). In this context, connections between non-clonally

related cells (typically of different colors), which cannot result from cell division, may

potentially correspond to de novo TNT-like connections. However, given the microscopy

limitations, we cannot rule out the possibility that some of these structures in fixed samples

are filopodia-like extensions fully spanning two cells. Endogenous fluorescence signals were

amplified to optimize optical visualization of these thin structures (**Fig. S4C-E**). The

tdTomato signal was enhanced using an anti-tdTomato antibody, while eYFP was amplified

using an anti-GFP antibody which also detected mTurquoise2, and to a lesser extent eBFP2

(**Fig. S4C-D**). Importantly, because anti-GFP amplifies both mTurquoise2 and eYFP, these

two reporters were not separable within the amplification channel. Nonetheless, given the

*Cytbow* design and the sparse labelling, the probability that a mTurquoise2[+] GC is physically

connected to a neighboring eYFP[+] GC remained limited, minimizing the likelihood of

overlooking connections between non-clonally related GCs detected by the anti-GFP

antibody. In addition, cerebellar slices were also labelled with an anti-anillin antibody (**Fig.**

**4A**) to detect CBs between labelled GCs (**Fig. S4E**). We identified both clonally related (87%;

**Fig. 4B, 4D, 4H, S4F, S4D**) and non-clonally related (13%; **Fig. 4C, 4D**) GCs connected via

putative TNT-like connections (N = 3 mice, 200 images, 27 pairs of putatively clonal-GCs

and 4 pairs of non-clonally related GCs). Overall, 8.89% of labelled GCs were connected via

putative TNT-like connections (**Fig. 4E**; number of TNT-like connections normalized to the

total number of labelled cells).

Among connected cells, 79% possessed a single TNT-like connection (53 out of 67 GCs; **Fig.**
**4B, F, H, Fig. S4F**) while 21% had multiple connections (14 out of 67 GCs; **Fig. 4C, F,**
**S4d$_1$**), a feature previously observed *in vitro* (Sáenz-de-Santa-María et al., 2023). Notably,
these connections originated from either soma or neurites, comparable to TNTs described *in*
*vitro* (Tardivel et al., 2016; Vargas et al., 2019). Of the 41 putative TNT-like connections
identified, 54% connected GC somas (22/41; **Fig. 4b$_1$-b$_2$, 4H, S4d$_1$, S4F**) 41% connected a
soma to a neurite (17/41; **Fig. 4c, 4F purple arrowhead**), and 5% connected two neurites
(2/41 connections; **Fig. 4F orange arrowhead**). Additionally, based on the DAPI pattern, we
occasionally observed TNT-like connections connecting dividing (identified by compact
chromatin) and non-dividing cells (which displayed more diffuse staining and darker
nucleolar regions) (**Fig. 4b$_1$**), a phenomenon previously reported in developing zebrafish
(Korenkova et al., 2024). To further assess the mitotic state of those connected GCs,
cerebellar slices were post-stained with anti-PH3 (mitotic marker Phospho-histone H3) to
more reliably identify mitotic cells compared to using DAPI staining alone. Additional
staining with anti-anillin was used to exclude CBs, and anti-tdTomato was applied to improve
the visualization of TNT-like connections (**Fig. 4H;** N = 3 mice, 86 images). We only found
putative TNT-like connections between non-dividing cells, suggesting that connections
between dividing and non-dividing cells were very rare (**Fig. 4I**). Indeed, although these
structures were distributed throughout the EGL, their location varied; some were in the oEGL
close to the pia (**Fig. 4b$_1$**), others located in the middle of the EGL (**Fig. 4b$_2$, S4F**), but the
majority were in the iEGL near the ML border, where GCs are mostly postmitotic (**Fig. 4C,**
**4H, S4d$_1$**).

**Conclusion**

This study identifies ICs in the EGL of the developing mouse cerebellum, focusing on CBs,
IBs, and other types of connections with TNT-like features. We provide evidence for the
presence of CBs in the iEGL, whereas IBs do not appear to be present in this region. In
addition, we observed diverse morphologies of potential ICs throughout the EGL—thin or
thick projections emerging from somata or neurites—which resemble previously reported
TNT-like structures *in vitro* and *in vivo* (Chang et al., 2025; Vargas et al., 2019). Notably, our
observations suggest that multiple ICs could arise between the same pair of cells. Although

GCs predominate in the EGL, our labelling strategy does not exclude the involvement of other

cell types in forming these connections (Cadilhac et al., 2021; Palese et al., 2025).

Since all ICs previously identified by ssSEM were open at both ends (Cordero Cervantes et

al., 2023), we speculate that some of the connections described here may also be open-ended.

However, we cannot exclude the possibility that certain structures represent filopodia-like

processes crossing over neighbouring cells or resemble DNTs—closed-end connections—

recently reported in the developing mouse cortex (Chang et al., 2025). The lower number of

ICs observed here compared to TNT-like structures in zebrafish embryos (35%) (Korenkova

et al., 2024) may reflect differences in tissue type, species, developmental timing, or

methodology. Fixation and immunostaining procedures are known to disrupt fragile TNTs

(Cordero Cervantes & Zurzolo, 2021; Delage et al., 2016; Matejka et al., 2024), and the use of

cytoplasmic rather than membrane-targeted reporters may have prevented full visualization of

very thin ICs, potentially leading to an underestimation of their frequency.

Overall, our findings are consistent with previous connectomic analyses of the developing

mouse cerebellum (Cordero Cervantes et al., 2023), revealing unusual connections between

neural cells and excluding a cytokinetic origin for these structures. Our observations are

compatible with the presence of TNT-like connections in the EGL that could contribute to

cerebellar development by enabling targeted communication before synapse formation

(Zurzolo, 2021), coordinating granule-cell migration (Teddy & Kulesa, 2004), or influencing

cerebellar organization and synaptogenesis (Kim et al., 2023). Future studies integrating

functional analysis with high-resolution live imaging will be essential to determine the nature,

dynamics, and developmental roles of these structures in cerebellar circuit assembly.

**Limitations of the Study**

Although we identified potential ICs based on 3D morphology, this study could not assess

their functional properties. Additional validation is required to determine whether these

structures correspond to true TNTs capable of molecular exchange or instead represent

cytonemes or filopodia-like protrusions in close proximity to neighboring cells. Functional

approaches, including live imaging to track their dynamics and cargo transfer, as well as

genetic or pharmacological disruption of their formation, will be critical to establish their role

in cerebellar circuit assembly. Our observations, together with our previous ssSEM-based

connectomic study (Cordero Cervantes et al., 2023), show that some of these ICs lack

cytokinetic and intercellular-bridge markers (Halcrow et al., 2022), arise between non-

dividing cells, and appear as thin processes linking distant granule cells, features compatible

with TNTs. However, current imaging resolution limits prevent definitive assessment of open-

ended continuity, and we cannot exclude the possibility that some structures correspond to

transient or incidental protrusions. Functional assays used to study TNT-like structures in

other systems, including zebrafish embryos (Korenkova et al., 2024) and interpericyte TNTs

(Alarcón-Martínez et al., 2020), have not yet been applied in the developing cerebellum. The

lack of specific TNT markers in neural tissue also limits discrimination from other IC types.

In addition, the developmental timing of TNT-like connection formation remains unclear;

approaches combining pulse–chase labelling with midbody or TNT-associated markers may

help define when these structures emerge relative to granule-cell maturation.

Addressing these limitations will be essential to determine whether TNT-mediated

communication contributes to cerebellar development and whether its dysregulation may be

relevant to neurodevelopmental disorders or cerebellar malformations.

**<u>Acknowledgments</u>**

The authors would like to acknowledge the help of the following Institut Pasteur facilities:

HistoPathology Core Facility, Photonic Bioimaging platform and the Image Analysis Hub.

The authors would like to thank all members of the Zurzolo group for fruitful discussions,

with special thanks to Thea Chrysostomou for helping to implement postnatal electroporation

at Institut Pasteur and Olga Korenkova for her technical and scientific feedback.

This work was supported by a dedicated grant called Big Brain Theory Program co-funded by

Institut Pasteur and Paris Brain Institute to CZ and LC, the Inception program (Investissement

d'Avenir grant) ANR-16-CONV-0005, and Equipe FRM - EQU202103012630 ANR

AAPG2023 UnProSec 2024-2027, FRM "MND202310017892" 2023-2027 to CZ, Conseil

régional d'Ile-de-France, DIM ELICIT and "Financement de Fin de thèse" de la FRM to GK.

**<u>Author contributions</u>**

Conceptualization: MR, CZ and LC. Methodology: MR, SL, GK, CZ, LC, FD and JL.

Investigation: MR, GK, GV, NDM. Formal analysis: MR, GV. Writing—original draft: MR.

Writing—review and editing: MR, GV, SL, CZ, LC, GK, JL, FD. Funding acquisition: CZ,

LC

## **<u>Declaration of interests</u>**

551    The authors declare no competing interests.

## References

Altman, J., & Bayer, S. A. (1997). *Development of the cerebellar system : in relation to its*
*evolution, structure, and functions*. CRC Press.
Bahram Sangani, N., Gomes, A. R., Curfs, L. M. G., & Reutelingsperger, C. P. (2021). The
role of Extracellular Vesicles during CNS development. *Prog Neurobiol*, *205*, 102124.
https://doi.org/10.1016/j.pneurobio.2021.102124
Berger, D. R., Seung, H. S., & Lichtman, J. W. (2018). VAST (Volume Annotation and
Segmentation Tool): Efficient Manual and Semi-Automatic Labeling of Large 3D
Image Stacks. *Front Neural Circuits*, *12*, 88. https://doi.org/10.3389/fncir.2018.00088
Cadilhac, C., Bachy, I., Forget, A., Hodson, D. J., Jahannault-Talignani, C., Furley, A.
J.,…Ango, F. (2021). Excitatory granule neuron precursors orchestrate laminar
localization and differentiation of cerebellar inhibitory interneuron subtypes. *Cell Rep*,
*34*(13), 108904. https://doi.org/10.1016/j.celrep.2021.108904
Chang, M., Krüssel, S., Parajuli, L. K., Kim, J., Lee, D., Merodio, A.,…Kwon, H. B. (2025).
Intercellular communication in the brain through a dendritic nanotubular network.
*Science*, *390*(6768), eadr7403. https://doi.org/10.1126/science.adr7403
Consalez, G. G., Goldowitz, D., Casoni, F., & Hawkes, R. (2020). Origins, Development, and
Compartmentation of the Granule Cells of the Cerebellum. *Front Neural Circuits*, *14*,
611841. https://doi.org/10.3389/fncir.2020.611841
Cordero Cervantes, D., Khare, H., Wilson, A. M., Mendoza, N. D., Coulon-Mahdi, O.,
Lichtman, J. W., & Zurzolo, C. (2023). 3D reconstruction of the cerebellar germinal
layer reveals tunneling connections between developing granule cells. *Sci Adv*, *9*(14),
eadf3471. https://doi.org/10.1126/sciadv.adf3471
Cordero Cervantes, D., & Zurzolo, C. (2021). Peering into tunneling nanotubes-The path
forward. *EMBO J*, *40*(8), e105789. https://doi.org/10.15252/embj.2020105789
Curran, B., & O'Connor, J. J. (2001). The pro-inflammatory cytokine interleukin-18 impairs
long-term potentiation and NMDA receptor-mediated transmission in the rat
hippocampus in vitro. *Neuroscience*, *108*(1), 83-90. https://doi.org/10.1016/s0306-
4522(01)00405-5
Delage, E., Cervantes, D. C., Penard, E., Schmitt, C., Syan, S., Disanza, A.,…Zurzolo, C.
(2016). Differential identity of Filopodia and Tunneling Nanotubes revealed by the
opposite functions of actin regulatory complexes. *Sci Rep*, *6*, 39632.
https://doi.org/10.1038/srep39632
Haglund, K., Nezis, I. P., & Stenmark, H. (2011). Structure and functions of stable intercellular
bridges formed by incomplete cytokinesis during development. *Commun Integr Biol*,
*4*(1), 1-9. https://doi.org/10.4161/cib.4.1.13550
Halcrow, E. F. J., Mazza, R., Diversi, A., Enright, A., & D'Avino, P. P. (2022). Midbody
Proteins Display Distinct Dynamics during Cytokinesis. *Cells*, *11*(21).
https://doi.org/10.3390/cells11213337
Hall, E. T., Dillard, M. E., Cleverdon, E. R., Zhang, Y., Daly, C. A., Ansari, S. S.,…Ogden, S.
K. (2024). Cytoneme signaling provides essential contributions to mammalian tissue
patterning. *Cell*, *187*(2), 276-293.e223. https://doi.org/10.1016/j.cell.2023.12.003
Hanzel, M., Rook, V., & Wingate, R. J. T. (2019). Mitotic granule cell precursors undergo
highly dynamic morphological transitions throughout the external germinal layer of the
chick cerebellum. *Sci Rep*, *9*(1), 15218. https://doi.org/10.1038/s41598-019-51532-y
Kim, M., Jun, S., Park, H., Tanaka-Yamamoto, K., & Yamamoto, Y. (2023). Regulation of
cerebellar network development by granule cells and their molecules. *Front Mol*
*Neurosci*, *16*, 1236015. https://doi.org/10.3389/fnmol.2023.1236015
Korenkova, O., Liu, S., Prlesi, I., Pepe, A., Albadri, S., Del Bene, F., & Zurzolo, C. (2024).
Tunneling nanotubes enable intercellular transfer in zebrafish embryos. *Dev Cell*.
https://doi.org/10.1016/j.devcel.2024.10.016

Korenkova, O., Pepe, A., & Zurzolo, C. (2020). Fine intercellular connections in development: TNTs, cytonemes, or intercellular bridges? *Cell Stress*, *4*(2), 30-43. https://doi.org/10.15698/cst2020.02.212

Lagace , D. C., Whitman, M. C., Noonan, M. A., Ables, J. L., DeCarolis, N. A., Arguello, A. A.,…Eisch, A. J. (2007). Dynamic Contribution of Nestin-Expressing Stem Cells to Adult Neurogenesis. *Journal of Neuroscience*, *27*(46), 12623-12629. https://doi.org/10.1523/JNEUROSCI.3812-07.2007

Legué, E., Gottshall, J. L., Jaumouillé, E., Roselló-Díez, A., Shi, W., Barraza, L. H.,…Joyner, A. L. (2016). Differential timing of granule cell production during cerebellum development underlies generation of the foliation pattern. *Neural Dev*, *11*(1), 17. https://doi.org/10.1186/s13064-016-0072-z

Loulier, K., Barry, R., Mahou, P., Le Franc, Y., Supatto, W., Matho, K. S.,…Livet, J. (2014). Multiplex cell and lineage tracking with combinatorial labels. *Neuron*, *81*(3), 505-520. https://doi.org/10.1016/j.neuron.2013.12.016

Matejka, N., Amarlou, A., Neubauer, J., Rudigkeit, S., & Reindl, J. (2024). High-Resolution Microscopic Characterization of Tunneling Nanotubes in Living U87 MG and LN229 Glioblastoma Cells. *Cells*, *13*(5). https://doi.org/10.3390/cells13050464

Miller, I., Min, M., Yang, C., Tian, C., Gookin, S., Carter, D., & Spencer, S. L. (2018). Ki67 is a Graded Rather than a Binary Marker of Proliferation versus Quiescence. *Cell Rep*, *24*(5), 1105-1112.e1105. https://doi.org/10.1016/j.celrep.2018.06.110

Nakashima, K., Umeshima, H., & Kengaku, M. (2015). Cerebellar granule cells are predominantly generated by terminal symmetric divisions of granule cell precursors. *Dev Dyn*, *244*(6), 748-758. https://doi.org/10.1002/dvdy.24276

Palese, F., Rakotobe, M., & Zurzolo, C. (2025). Transforming the concept of connectivity: unveiling tunneling nanotube biology, and their roles in brain development and neurodegeneration. *Physiol Rev*. https://doi.org/10.1152/physrev.00023.2024

Pereda, A. E. (2014). Electrical synapses and their functional interactions with chemical synapses. *Nat Rev Neurosci*, *15*(4), 250-263. https://doi.org/10.1038/nrn3708

Price, K. L., Tharakan, D. M., & Cooley, L. (2023). Evolutionarily conserved midbody remodeling precedes ring canal formation during gametogenesis. *Dev Cell*, *58*(6), 474-488.e475. https://doi.org/10.1016/j.devcel.2023.02.008

Qu, X., Kumar, A., & Bartolini, F. (2021). Live imaging of microtubule dynamics at excitatory presynaptic boutons in primary hippocampal neurons and acute hippocampal slices. *STAR Protoc*, *2*(1), 100342. https://doi.org/10.1016/j.xpro.2021.100342

Renshaw, M. J., Liu, J., Lavoie, B. D., & Wilde, A. (2014). Anillin-dependent organization of septin filaments promotes intercellular bridge elongation and Chmp4B targeting to the abscission site. *Open Biol*, *4*(1), 130190. https://doi.org/10.1098/rsob.130190

Rujano, M. A., Briand, D., Đelić, B., Marc, J., & Spéder, P. (2022). An interplay between cellular growth and atypical fusion defines morphogenesis of a modular glial niche in Drosophila. *Nat Commun*, *13*(1), 4999. https://doi.org/10.1038/s41467-022-32685-3

Sáenz-de-Santa-María, I., Henderson, J. M., Pepe, A., & Zurzolo, C. (2023). Identification and Characterization of Tunneling Nanotubes for Intercellular Trafficking. *Curr Protoc*, *3*(11), e939. https://doi.org/10.1002/cpz1.939

Sahay, A., Kale, A., & Joshi, S. (2020). Role of neurotrophins in pregnancy and offspring brain development. *Neuropeptides*, *83*, 102075. https://doi.org/10.1016/j.npep.2020.102075

Sobecki, M., Mrouj, K., Camasses, A., Parisis, N., Nicolas, E., Llères, D.,…Fisher, D. (2016). The cell proliferation antigen Ki-67 organises heterochromatin. *Elife*, *5*, e13722. https://doi.org/10.7554/eLife.13722

Stringer, C., Wang, T., Michaelos, M., & Pachitariu, M. (2021). Cellpose: a generalist algorithm for cellular segmentation. *Nat Methods*, *18*(1), 100-106. https://doi.org/10.1038/s41592-020-01018-x

Sun, X., & Kaufman, P. D. (2018). Ki-67: more than a proliferation marker. *Chromosoma*, *127*(2), 175-186. https://doi.org/10.1007/s00412-018-0659-8

Tardivel, M., Bégard, S., Bousset, L., Dujardin, S., Coens, A., Melki, R.,…Colin, M. (2016).
Tunneling nanotube (TNT)-mediated neuron-to neuron transfer of pathological Tau
protein assemblies. *Acta Neuropathol Commun*, *4*(1), 117.
https://doi.org/10.1186/s40478-016-0386-4
Teddy, J. M., & Kulesa, P. M. (2004). In vivo evidence for short- and long-range cell
communication in cranial neural crest cells. *Development*, *131*(24), 6141-6151.
https://doi.org/10.1242/dev.01534
Vargas, J. Y., Loria, F., Wu, Y. J., Córdova, G., Nonaka, T., Bellow, S.,…Zurzolo, C. (2019).
The Wnt/Ca [2+] pathway is involved in interneuronal communication mediated by
tunneling nanotubes. *EMBO J*, *38*(23), e101230.
https://doi.org/10.15252/embj.2018101230
Zurzolo, C. (2021). Tunneling nanotubes: Reshaping connectivity. *Curr Opin Cell Biol*, *71*,
139-147. https://doi.org/10.1016/j.ceb.2021.03.003

**Fig. 1. Assessment of the presence of CBs in the iEGL**

(A) Sagittal sections of P7 WT cerebellum, taken from the medial vermis (lobule VIII), labelled with anti-TAG-1 (green), anti-Ki67 (red), anti-Aurora B kinase (grey) and DAPI staining (blue). TAG-1 and Ki67 expression were used to distinguish the iEGL from the oEGL using a CNN-based classifier that automatically generated a mask for the iEGL; the resulting boundary is indicated by the dotted line. Solid lines delineate the EGL, with the pia located above and the ML below. Yellow arrowhead points at a CB and white arrowheads at mitotic cells in the oEGL. (B) Higher magnification views of the regions shown in (A) highlighting (a) a mitotic cell (white arrowheads) and (b) a CB (yellow arrowhead) within the iEGL. (C) Distribution of CBs across the EGL, represented as the relative distance from the ML to the pia (0 = ML, 1= pia). The oEGL-iEGL transition corresponds approximately to the mid-point of this axis. (D) Comparison of the percentage of mitotic cells (0.69%) and CBs (0.66%) in the iEGL. The data are presented as mean ± SEM; p-value= 0.7 analyzed using a Student T-test. (C-D) N = 4 mice, 55 images were analyzed in total. All images are maximum-intensity projections of six z-slices acquired at 63x magnification.

**Fig. 2. Assessment of the nature of CBs in the iEGL**

(A) Sagittal sections of WT P7 cerebellum, taken from the medial vermis (lobule VIII), labelled with anti-TAG-1 (green), anti-Ki67 (red), anti-Aurora B kinase (grey) and DAPI staining (blue). The expression of TAG-1 was used to identify the iEGL using a CNN-based method that creates a mask on the iEGL area. The white arrowheads show Ki67+ cells that are connected by a CB. The yellow arrowheads show Ki67- cells, which were never observed to be connected by a CB. (B) Distribution of CBs connecting cells in the different proliferative states in the iEGL. The proliferative state is determined based on the pattern and expression level of Ki67. "Ki67$^{mitotic}$" (51%) represents cells that are undergoing mitosis, "Ki67$^{interphase}$" (49%) represents cells in earlier cell cycle phases or those that have recently exited proliferation, "Ki67$^{G0}$" (0%) represents cells in late G0. Per-image percentages were averaged across all images from each mouse to obtain a mean value per animal. N = 4 mice; 55 images were analyzed in total. (C) Labelling with anti-citron kinase (green), anti-anillin (red), anti-Aurora B kinase (grey) and DAPI staining (blue). (c1-c3) Higher magnification of the areas indicated in (C), showing CBs labelled with both anillin and Aurora B kinase (c1), only Aurora B kinase (c2) or only anillin (c3). All CBs expressed citron kinase. (D) Proportion of CBs in the categories shown in c1-c3: 81.33% were Aurora B kinase$^+$/Anillin$^+$/citron kinase$^+$, 2.78% were Aurora B kinase$^+$/Anillin$^-$/citron kinase$^+$, and 15.89% labelled Aurora B kinase$^-$/Anillin$^+$/citron kinase$^+$. N = 4 mice, 66 images were analyzed in total. Pannel (A) corresponds to a z-slice, (C) and (c1-c3) are maximum projections of 6 z-slices (z-step=0.35 μm). All images were acquired at 63x magnification.

**Fig. 3. Identification of potential GC connections compatible with TNT-like bridges**

(A) Schematic representation of the strategy used to sparsely label cells and visualize ICs: The pCAG-Dendra2-E2A-GFP-Cep55 plasmid is electroporated in the EGL of P5 pups. The pups are sacrificed at P7 and cerebellar slices were stained with anti-anillin to distinguish CBs from putative connections non-derived from cells division (ie. putative TNT-like connections). (B) Representative image showing the expression of the plasmid in the EGL. (C) Sagittal cerebellar slices labeled with anti-GFP (green), anti-anillin (red), and DAPI, showing a CB (anillin⁺ connection) and a TNT-like connection (anillin- connection). The white arrowheads point at the connected cells. Dotted lines delineate the connections. (D) Graph representing the proportion of CBs and TNT-like connections over the total number of ICs identified. (E) Length of the connections identified in (D) showing an average TNT-like connections and IB's length of 5.38 µm and 2.92 µm, respectively. (D-E) N = 3 mice, 36 TNT-like connections and 11 CBs identified, 83 images were analyzed in total. Although cells were labelled, no connections were found in the EGL of one mouse, likely due to the rarity of these connections. In (E) data are presented as mean ± SEM; p-value= 0.09 analyzed using a Welch T-test. (F) Example of TNT-like connection forming between two cells, via the filopodia driven mechanism in a sagittal cerebellar slice. A zoomed-in view of the area where the connection forms (indicated by the dotted purple square) is shown in the bottom right corner of each time point. The two protrusions establish connection approximately 20 min after the start of the recording. (F') Plot profile of the connection shown in (F) at the different time points, showing a continuous connection (grey value > 0) between 20-30 min. Panel (B) shows a z-maximal projection at x10 magnification. (C,F) images correspond to 3D views at 60x magnification. Solid lines delineate the EGL.

**A**

NestinCre$^{ERT2+/-}$ ; Cytbow$^{+/+}$
**Tamoxifen injection P0-P3**

EGL

**Sacrifice P7**

Staining

Anillin+   CB
Anillin-   **TNT-like** connection
Anillin-   **TNT-like** connection

**B** — DAPI / tdTomato / Anillin

Pia
EGL
b₂  b₁
ML
10 µm

b₁ — Pia / EGL / 5 µm
b₂ — EGL

**C** — DAPI / GFP / tdTomato

EGL
ML
5 µm

**D** — Types of connected cells

13%
87%

Same color
Different color

**E**

% of connected cells in the EGL

8,89

Mean percentage of connected cells

**F** — Number of connections per connected cells

21%
79%

1 connection
>1 connection

**G** — Types of connections

5%
41%
54%

Soma-soma
Soma-neurite
Neurite-neurite

**H** — DAPI / PH3 / tdTomato / Anillin

EGL
5 µm

**I**

| | | |
|---|---|---|
| | **Total number of connections** | 10 |
| **TdTomato+ cells** | % of connections between a pair of PH3- cells | 100 |
| | % of connections between a PH3+ and a PH3- cells | 0 |
| | % of connections between a pair of PH3+ cells | 0 |

**J** — Distribution of connected cells (anillin- connections) throughout the EGL

% of connected cells

**Localisation of connected cells**

| | |
|---|---|
| 11.76 | Both close to the pia |
| 8.82 | One close to the pia, the other in the mid-EGL |
| 14.71 | Both in the mid-EGL |
| 5,88 | One in the mid-EGL, the other close to the ML |
| 52.94 | Both close to the ML |
| 5.88 | Once close to ML, the other in the ML |

**Fig. 4. Assessment of the features of potential connections in the EGL**

(A) Schematic representation of the strategy used to sparsely label cells and visualize ICs: *NestinCre$^{ERT2+/-}$ ; Cytbow$^{+/+}$* mice were injected with tamoxifen between P0 and P3 to induce the mosaic expression of mTurquoise2, eYFP, tdTomato or a combination of two of these reporters in neurons. Mice were sacrificed at P7 and were stained with DAPI, anti-anillin (grey) to distinguish CBs from putative TNT-like connections, anti-GFP (green) and anti-tdTomato (red) to amplify reporter signals and differentiate clonally vs. non-clonally related cells. (B) Representative image of TNT-like connections (anillin-) between tdTomato+ cells in the EGL. (b$_1$- b$_2$) Represent a higher magnification of the areas indicated in (B) showing TNT-like connections close to the pia (b$_1$) and in the middle of the EGL (b$_2$). The asterisk marks a dividing cell, as identified by the DAPI pattern. (C) Representative image showing a TNT-like connection between a tdTomato$^+$ cell and a tdTomato$^+$GFP$^+$ cell near the ML border. The blue arrowhead points at a connection formed between the soma and neurite. The orange arrowhead points at a connection between the neurites. (D) Graph presenting the percentage of TNT-like connection between same-color vs. different-color cells. (E) Graph representing the relative percentage of connected cells over the total labelled population. The dotted line indicates the mean. SEM is ± 0.061. (F) Graph presenting the percentage of cells forming one (53 out of 67 GCs) vs. multiple (>1) TNT-like connections (14 out of 67 GCs). (G) Graph presenting the percentage of TNT-like connections categorized by location: soma-soma (22/41 connections), neurite-neurite (17/41 connections), soma-neurite (2/41 connections). (H) Labelling of EGL sagittal sections with anti-PH3 (in green), anti-Tdtomato (in red), anti-anillin (in grey) and DAPI staining (in blue). (J) Spatial distribution of connected (I) Quantification representing the percentage of TNT-like connections (anillin$^-$) between non-dividing cells (PH3$^-$), dividing cells (PH3$^+$), one dividing (PH3$^+$) and one non-dividing cell (PH3$^-$) cells across the EGL. (b$_1$-b$_2$, C, F, I) The white arrowheads indicate connected cells. The dashed lines delineate TNT-like connections. (D-E, G-H, K) N = 3 mice, 200 images. (I-J) N = 3 mice, 86 images. All images correspond to 3D views and were taken at 63x magnification. Solid lines delineate the EGL. Dotted lines delineate the connections.