Legacy of warming and microbial treatments shape root exudates and

rhizosphere fungal communities of a tropical tree

Joel Masanga[1*], Parker Bartz[2], Gabriela Hernandes Villani[2], Iana F. Grullón-Penkova[3], Benedicte

Bachelot[2], Jesse R. Lasky[1,4]

[1]Department of Biology, Pennsylvania State University, University Park, PA, USA.

[2]Department of Plant Biology Ecology and Evolution, Oklahoma State University, Stillwater, OK,

USA.

[3]USDA Forest Service International Institute of Tropical Forestry Río Piedras, PR, USA.

[4]PAC Herbarium, Pennsylvania State University, University Park, PA, USA.

*Corresponding   author.

Summary

- Plant metabolites play a pivotal role in shaping rhizosphere microbial communities, yet how plant and microbial functions respond to environmental change remains poorly understood. We combined ecometabolomics and fungal community profiling to investigate how multiple abiotic and biotic treatments influence root exudate chemistry and fungal community assemblage in a tropical tree.

- *Guarea guidonia* seedlings were grown in sterilized soil primed with inoculum from long-term experimentally warmed or ambient plots in Puerto Rico. We manipulated soil microbes, soil moisture, and plant density and tested metabolite-fungi-plant trait associations.

- Metabolite diversity was significantly influenced by soil microbial legacy and moisture, while antimicrobial treatments altered metabolite composition without affecting diversity. Metabolite clusters exhibited distinct treatment-specific patterns. Fungal alpha diversity increased under low moisture, community composition shifted with antimicrobial treatments, and certain families showed strong treatment-specific responses. Although fungal diversity was not correlated with metabolite diversity, fungal community structure was significantly associated with metabolite composition. We also found global and pathway-specific correlations between fungal and metabolite distances, and weak but significant associations between metabolite/fungal composition and seedling traits.

- These results underscore how environmental conditions shape belowground interactions, highlighting metabolite–fungal associations as potential early indicators of plant response to disturbance.

Keywords: Abiotic stress; Ecometabolomics; Experimental warming; Plant–microbe interactions; Root exudates; Soil microbial legacy; Tropical forest ecology.

Introduction

Climate change has intensified disturbance events in the tropics (Emanuel, 2005; Walsh & Ryan,

2000), with major consequences for forest structure, composition, and ecosystem function

(Andrewin et al., 2015; Webster et al., 2005; Lugo, 2008). Ecosystem recovery from related

disturbances likely hinges on microbial feedbacks, with mutualists enhancing plant resilience

and pathogens compounding stress effects (Brooker et al., 2021; Wright et al., 2017). Past

evidence from experimental warming and drought studies in tropical forests suggest that such

disturbances may for instance reduce pathogen pressure while favoring beneficial microbes

(Bachelot et al., 2020), although the biochemical mechanisms mediating these shifts remain

unclear. Research on ecological disturbance has largely centered on temperate systems,

aboveground traits and single stressors, which not only leaves belowground dynamics

underexplored (Melillo et al., 2017; Schindlbacher et al., 2011) but also overlooks the

potentially non-additive and complex effects of interacting stressors (Eaton et al., 2020;

Robertson & Platt, 2001; Vargas et al., 2010). The Stress Gradient Hypothesis (Bertness &

Callaway, 1994) posits that increasing stress favors facilitative over competitive interactions. In

this context, Caribbean forests, which are chronically disturbed by storms and warming, offer a

valuable system to study how multiple stressors shape belowground ecological dynamics.

Plant-microbe interactions are central to nutrient cycling, seedling establishment, and plant–

soil feedbacks (Van Der Heijden et al., 2008; Wagg et al., 2014), yet this key interaction is rarely

considered in studies that explore how ongoing changes in climate will shape the future of

plant communities. In addition, much of our current understanding of plant-microbe

interactions under changing environments comes from crop systems and single-stressor

experiments, leaving important gaps in how plants and microbes jointly respond to complex

interacting disturbances under natural environments (Trivedi et al., 2022). Plant–microbe–soil

interactions heavily rely on root exudates, which contain a repertoire of primary and secondary

metabolites that influence rhizosphere microbial composition (Hu et al., 2018) and nutrient

cycling (Majumdar et al., 2023; Müller & Junker, 2022; Pang et al., 2021). Further, root

exudation and chemistry have been shown to dynamically respond to abiotic and biotic stimuli

(Gidman et al., 2003; Gidman et al., 2005; Jones et al., 2011; Lake et al., 2009; Widarto et al.,

2006). The interaction between plants, their chemical profiles and rhizosphere microbiome can

generate plant–soil feedbacks that influence performance and community dynamics of future

communities (Bever, 1994; Gfeller et al., 2023; Hannula et al., 2021; Van der Putten et al.,

2013). This might be by altering the soil biotic environment, either positively through

recruitment of mutualists such as arbuscular mycorrhizal fungi (AMF) (Bennett et al., 2012;

Berendsen et al., 2012; Tamburini et al., 2020), or negatively via pathogen accumulation and

resource depletion (Maron et al., 2016; Thakur et al., 2021; Yang et al., 2015). While these

interactions have been widely explored in agroecosystems and invasion ecology (Engelkes et al.,

2008; Majumdar et al., 2023; van der Putten et al., 2016), their role in mediating tropical forest

responses to environmental stress remains underexplored.

Here, we investigated how altered biotic and abiotic conditions influence root exudate

chemistry, rhizosphere fungal communities, and seedling traits in the tropical tree species *G.*

guidonia. We hypothesized that: (H1) Variation in biotic (microbial alteration) and abiotic

(warming, soil moisture and plant density) conditions leads to significant shifts in root exudate

metabolic profiles; (H2) These conditions influence rhizosphere fungal community diversity,

composition, and abundance; (H3) Differences in root exudate chemistry are associated with

corresponding changes in fungal community structure; and (H4) Variation in seedling

performance is linked to treatment-driven changes in both root exudate profiles and fungal

community composition (Fig. 1a). To test these hypotheses, we employed untargeted

metabolomics and amplicon-based fungal community profiling to detect metabolites in root

exudates and identify rhizosphere fungal taxa, respectively. We grew *G. guidonia* seedlings in

sterilized soil primed with inoculum from either experimentally warmed (+4◻) or ambient plots

at the Tropical Response to Altered Climate Experiment (TRACE) in Puerto Rico, a long-term

field warming experiment (Kimball et al., 2018) that has spanned multiple climatic events

including major droughts and hurricanes (Alonso-Rodriquez et al., 2022). This design enabled us

to assess how microbial communities that resulted from exposure to long-term warming (as a

proxy for climate-driven plant-soil-feedback) affect plant metabolite exudation and rhizosphere

assembly. We manipulated soil and foliar microbes (via fungicides and heat sterilization), soil

moisture (reflecting drought projections described by Bhardwaj et al., 2018), and plant density

to test for context-dependent responses. We then evaluated associations between metabolite

composition, fungal communities, and seedling traits, with a view that such links reflect early-

stage feedbacks relevant to tropical forest resilience and recovery under climate shifts (Jiang et

al., 2024; Noecker et al., 2019).

Material and Methods

Experimental treatments, plant material and data collection

This study was conducted under shade house conditions, between January and July 2023 at the

USDA Forest Service Sabana Field Research Station in the Luquillo Experimental Forest,

northeastern Puerto Rico (18°18𝔼N, 65°50𝔼W). Forest soil was collected from the neighboring

forest, a wet subtropical forest with clay-rich, acidic inceptisols. The soil was sieved, sterilized at

70𝔼°C for 48𝔼hours, and 250𝔼g filled into 4-liter plastic pots. Half of the pots (n = 120) were

primed with 5𝔼g of soil inoculum collected from either experimentally warmed or ambient

control plots at the nearby TRACE field warming experiment. Experimental warming plots have

been maintained at +4𝔼°C above ambient temperature since 2016, with a one-year pause in

warming following Hurricanes Irma and Maria in September 2017 (Reed et al., 2020). These

inocula, intended to transfer microbial legacies, are hereafter referred to as "warmed soil" and

"ambient soil," respectively. The pots were further split into four equal batches in which the

inoculum was subjected to one of the following antimicrobial treatments; soil fungicide

(treated with Banrot™ (Prabhakaran & Dann, 2022); foliar fungicide (treated with Abound

Flowable™ (Starkey et al., 2013); heat treatment (oven-sterilized at 70𝔼 for 48 hours), and

control (no fungicide/heat sterilization treatment). These are collectively referred to as

antimicrobial treatments (Fig. 1b). Pots were randomly arranged on raised benchtops, inside a

shade house constructed using a gardening tarp that guarantees low light (60% shade) and

allows air circulation but restricts insect entry.

We studied the shade tolerant tree *G. guidonia*, a dominant late-successional broadleaved

evergreen species common in tropical understories (Pennington and Clarkson 2013). The

seedling stage was targeted due to its documented sensitivity to soil microbial variation and

environmental stress (Elger et al., 2009). To ensure uniformity across plant material, we grew

apparently phenotypically uniform seedlings, germinated from surface sterilized (in 5% NaOCl)

seeds collected from mother plants in close proximity. Seedlings were transplanted at low (1

plant/pot), moderate (3 plants/pot), and high (5 plants/pot) densities to test for plant–plant

interactions. Water availability was manipulated with water-stress pots receiving 60 mL

distilled water twice a week and well-watered controls receiving 120 mL at the same frequency

(Fig. 1b). After three months of growth, root exudates were collected by exposing a single

lateral root (still attached to the plant) in each pot, gently cleaning it with ethanol-soaked

cotton balls, and submerging it in 5 mL deionized water for 48 hours under darkness provided

by foil-wrap. Samples were stored at –80 °C until metabolomic profiling. Plants were then

phenotyped for aboveground traits (leaf surface area, specific leaf area, and leaf number), prior

to harvesting. Roots were scanned using WinRHIZO™ (Regent Instruments Inc., Canada) to

assess total root length, root surface area, root diameter, root volume, and number of root tips.

Subsamples were flash-frozen at –80 °C for rhizosphere fungal community sequencing.

Fig. 1 Root exudate metabolomes of *G. guidonia* are shaped by warming-induced microbial legacy and antimicrobial treatments. (a) Metabolite diversity (Functional Hill numbers) across antimicrobial treatments (x-axis), colored based on microbial legacy conditions (ambient vs. warmed soil). Boxplot whiskers represent 25th and 75th percentiles, while dots indicate biological replicates. (b) NMDS ordination of root exudate metabolite composition based on Bray–Curtis distances of normalized peak intensities. (c) Linear discriminant analysis (LDA) highlighting separation of metabolomic profiles across treatments. (d) Volcano plots of differentially abundant metabolites. Dotted lines indicate raw P-value threshold (P=0.05). Points are colored by FDR-adjusted significance: red = enriched, blue = depleted, grey = not significant.

Liquid Chromatography-Mass Spectrometry

Root exudates were centrifuged (300 rpm, 20 min) and concentrated using solid-phase
extraction (SPE) prior to LC–MS/MS analysis. Samples were passed through C18 columns
(HyperSep SPE, Thermo Fisher, USA) preconditioned with acidified methanol (1% formic acid)
and HPLC-grade water under vacuum (Resprep manifold, Restek). Metabolites were eluted in
3 mL methanol, concentrated in a centrifugal vacuum concentrator (SpeedVac), and
reconstituted in 100 µL of 60:40 methanol:water containing 0.1 µM chlorpropamide (internal
standard). LC–MS/MS was performed at the Huck Metabolomics Core Facility, Pennsylvania
State University, on a Thermo Vanquish Horizon system (Thermo Fisher Scientific, Waltham,
MA, USA) equipped with a Waters BEH C18 column (2.1 × 150 mm, 1.7 µm) at 55 °C. The
mobile phase consisted of 0.1% formic acid in water (solvent A) and 0.1% formic acid in
acetonitrile (solvent B), with the following gradient: 10% B (0–2 min), ramped to 40% B
(12 min), then to 100% B (15 min), held until 21 min, returned to 10% B at 22 min, and re-
equilibrated until 30 min. The flow rate was 0.3 mL/min. The eluate was introduced into a
Thermo Orbitrap Exploris 120 mass spectrometer using heated electrospray ionization. Full scan
data (m/z 100–1000) were acquired at 120,000 resolution, followed by ddMS2 scans at 15,000
resolution using normalized collision energies of 15, 30, and 45 V. Capillary voltages were
2500 V (negative mode) and 3500 V (positive mode). Sheath gas was set to 40, auxiliary gas to
6, and sweep gas to 0. The ion transfer tube and vaporizer were set to 325 °C and 350 °C,
respectively. Data was obtained in both positive and negative modes, and analyzed using
MetaboAnalyst 6.0 (Pang et al., 2024).

Fungal community characterization

To assess fungal community structure, we sampled roots from pots with surviving seedlings
following exudate collection at the end of the 3 months growth period. Root samples were
lysed using a TissueLyser II (Qiagen, Germantown, MD, USA) with ceramic beads, and total DNA
extracted using the PowerPlant Pro Kit (Qiagen), following the manufacturer's protocol with
minor modifications. PCR amplification was performed targeting the fungal internal transcribed

spacer 2 (ITS2) region using primers ITS4_Nextera and JL0015.8SR_Nextera. PCR cycling

conditions were: 95 °C for 3 min (initial denaturation), followed by 30 cycles of 98 °C for 20 s,

65.7 °C for 15 s, and 72 °C for 45 s, with a final extension at 72 °C for 5 min (Gohl et al.,

2016). Sequencing was performed on the NextSeq P1 XLEAP platform at the University of

Minnesota Genomic Center (UMGC). Initial quality control and demultiplexing were also

conducted by UMGC. Sequence processing and operational taxonomic units (OTU) clustering

were performed on the Pete supercomputer at Oklahoma State University using a modified

Mothur pipeline (Schloss et al., 2009). After detecting and removing chimeras, this pipeline

generates operational taxonomic units (OTUs) by grouping sequences at 97% identity (Gweon

et al., 2015) using the UNITE v9 database (Abarenkov et al., 2024). Taxonomic classification and

ecological guild assignment were completed using FUNGuild (Nguyen et al., 2016), enabling

annotation of fungal OTUs by functional role.

Statistical analyses

Treatment effects on metabolite diversity and composition

We analyzed exudate metabolite diversity using the 'chemodiv' R package (Petrén et al., 2023),

which calculates Functional Hill numbers integrating compound richness, relative abundance,

and structural dissimilarity (Chiu & Chao, 2014). Compounds were categorized using the

function *NPCTable*, and pairwise dissimilarities computed with *compDis*. Diversity metrics were

compared across antimicrobial treatments, microbial legacy, plant density, and soil moisture

using two-tailed t-tests or one-way ANOVA. To assess treatment effects on metabolite

composition, we computed Bray–Curtis dissimilarity matrices and conducted PERMANOVA

using *adonis2* function in 'vegan'. We used Linear Discriminant Analysis (LDA), implemented in

the 'LDA' package to identify metabolites that best discriminated among treatment groups.

Significant contributors ($p < 0.05$) were ranked based on LDA scores. Differential abundance of

individual metabolites was evaluated using the 'limma' package, relative to control (untreated

inoculum) as the reference group. Metabolites were considered significantly modulated if they

had a false discovery rate (FDR) < 0.05. To explore non-linear treatment effects, we employed

the package 'cluster' (Freire-Zapata et al., 2024) to identify metabolite clusters based on co-

occurrence patterns across samples. Only metabolites detected in ≥50% of samples were

included. Pairwise dissimilarities were computed based on Manhattan distances using the

function *daisy*, while clustering was performed using Partitioning Around Medoids (PAM) using

the function *pam*. The optimal number of clusters (k) was determined by silhouette analysis.

Consensus values, reflecting sample-level contributions to each cluster, were calculated for use

in downstream correlations with fungal diversity and seedling traits.

Treatment effects on fungal diversity, composition and differential abundance

To characterize fungal diversity across experimental treatments, we calculated *alpha* diversity

indices from OTU tables after rarefaction using the package 'phyloseq' (McMurdie & Holmes,

2013). These metrics were compared across treatments using t-tests or ANOVA. Fungal *beta*

diversity was estimated via Bray–Curtis dissimilarity and tested using PERMANOVA. Differential

abundance analysis was conducted using the package 'ANCOM-BC' (Lin & Peddada, 2020),

which accounts for compositional bias and sample-level heterogeneity. Taxa were compared

relative to controls, and significance was determined at FDR<0.05 based on the Benjamini–

Hochberg correction.

Metabolite-fungal community associations

We examined metabolite–fungal associations using multiple complementary approaches. (i)

Mantel tests, conducted in 'vegan' with 999 permutations, were performed between Bray–

Curtis dissimilarity matrices of metabolites and fungal communities for all samples that had

both datasets (n = 144). This included all detected compounds (including unknowns, n=612) and

pathway subsets, namely alkaloids (n=150), amino acids and peptides (n=68), carbohydrates

(n=28), fatty acids (n=91), polyketides (n=23), shikimates and phenylpropanoids (n=110), and

terpenoids (n=79). (ii) Partial Spearman's rank correlations, implemented in the 'ppcor' v.1.1

package (Kim, 2015) were used to test associations between individual metabolite features and

fungal Shannon diversity and OTU abundance. All models were adjusted for microbial legacy,

antimicrobial treatment, plant density, and soil moisture. Results were corrected for multiple

comparisons using FDR. (iii) Redundancy Analysis (RDA) was conducted using the first 10

metabolite principal components (PCs) to quantify explained variance in fungal community

composition. Contributions of individual PCs to RDA axes were assessed via biplot loadings,

while outlier fungal taxa associated with metabolite gradients were identified based on

Mahalanobis distances from the RDA centroid. These taxa were interpreted as drivers of

community-level metabolomic associations. (iv) Spearman's rank tests, implemented in the

'Hmisc' package, were used to correlate metabolite cluster consensus values with fungal

Shannon diversity, and results corrected for multiple comparisons using FDR.

Metabolite- and fungi-seedling trait associations

To test whether variation in metabolite or fungal composition was linked to seedling traits, we

first constructed a Euclidean distance matrix from six morphological traits (root surface area,

root diameter, root volume, dry root mass, leaf surface area, and specific leaf area). This matrix

was correlated with metabolite and fungal community dissimilarities using Mantel tests. To

detect coordinated divergence, we extracted sample pairs in the upper-right quadrant (top

25%) of both trait vs metabolite and trait vs fungal Mantel test results. Treatment combinations

within these high-divergence pairs were summarized in contingency tables and visualized as

heatmaps to identify dominant treatment contrasts driving divergence. Finally, to detect

metabolite–trait associations beyond global metabolomic distance metrics, we correlated

metabolite cluster consensus values (from cluster analysis) with individual plant traits using

Spearman's rank tests, followed by FDR correction.

Results

Warming-induced microbial legacy and heat sterilization of soil inoculum

influence exudate metabolomes

We hypothesized that biotic (microbial alteration) and abiotic (warming, soil moisture and plant

density) conditions and/or their combinations influence root exudate metabolite profiles in *G*.

guidonia (H1). To test this hypothesis, we conducted untargeted metabolomics on root
exudates from 162 individuals and identified 612 putative metabolites (471 in positive and 141
in negative ion modes). NPClassifier assigned the compounds to seven major pathways:
alkaloids (150), amino acids and peptides (68), carbohydrates (28), fatty acids (91), polyketides
(23), shikimates and phenylpropanoids (110), and terpenoids (79). Notably, 63 features could
not be classified. Next, we quantified metabolite diversity using Functional Hill numbers and
compared diversity across treatments. Significantly higher metabolite diversity was observed
under ambient than warmed soil inoculum (t = 3.71, p = 0.0003), and under high than low soil
moisture (t = 2.12, p = 0.036) (Fig. 2a; Fig. S1). In contrast, neither antimicrobial treatments nor
plant density significantly affected metabolite diversity (both p > 0.05). Likewise, antimicrobial
treatments did not alter the overall distribution of metabolic pathways (Fig. S1). Metabolite
composition was significantly influenced by microbial legacy and antimicrobial treatments (p <
0.05 for both), but not by soil moisture or plant density (p > 0.05) (Fig. 2b). Together, these
results suggest that warming legacy effects on microbes and soil moisture level primarily
influence metabolite richness. In contrast, microbial alteration and plant density have a
stronger effect on metabolite composition, shifting the relative abundances of specific
compounds rather than the total number present.

To further understand how treatments shaped metabolite composition, we performed linear
discriminant analysis (LDA), and observed clear separation based on microbial legacy (Fig. 2c).
The top 20 metabolites contributing to each LDA axis are shown in Fig. S1, while the full list of
differentially discriminant features is provided in **Dataset S1**. To identify specific compounds
driving these patterns, we applied the 'limma' package in R for differential enrichment and
depletion of metabolites under microbial legacy and antimicrobial treatments, two factors that
significantly influenced composition (Fig. 2b). We identified 255 metabolites that differed
significantly between warmed and ambient inoculum sources (126 enriched and 129 depleted;
FDR < 0.05). Additionally, 183 and 148 metabolites were differentially abundant in soil fungicide
and foliar fungicide treatments, respectively, compared to untreated controls (FDR < 0.05). The

heat-treated condition yielded 39 metabolites that were nominally significant (P < 0.05) but did

not pass FDR correction (FDR > 0.05) (Fig. 12; Fig. S2).

Fig. 2 Diversity and composition of rhizosphere fungal communities associated with *G.*

guidonia. (a) Shannon diversity of fungal communities across antimicrobial treatments (x-axis)

and moisture. H and L denote high and low moisture levels, respectively. (b) Relative

abundance of dominant fungal families across antimicrobial treatments. (c) Principal coordinate

analysis (PCoA) based on Bray–Curtis dissimilarities illustrating shifts in fungal community

composition in response to treatments. (d) Differential abundance of fungal taxa based on

ANCOM-BC analysis. Enrichment and depletion patterns are shown relative to the untreated

control group.

Metabolite clusters reveal distinct treatment-specific responses

We identified three metabolite clusters based on co-occurring features. The clusters consisted

of 88 (Cluster 1), 141 (Cluster 2), and 53 (Cluster 3) metabolites, with all seven annotated

metabolite pathways represented across clusters (Fig. S2). Heatmaps revealed clear treatment-

specific shifts in metabolite abundance relative to control conditions across the three clusters

(Fig. S2). To further evaluate treatment effects, we fitted linear models with interaction terms

and found that metabolites in Cluster 1 were significantly influenced by all experimental

treatments ($p < 0.05$) except low plant density ($p > 0.05$). Cluster 2 metabolites responded

significantly to microbial legacy ($p = 0.004$), antimicrobial treatments ($p < 0.001$), and moisture

levels ($p < 0.001$). Cluster 3 metabolites showed a similar treatment response to Cluster 1.

Linear model results are summarized in **Dataset S2-S4**. By grouping metabolites based on co-

occurrence, we uncovered nuanced environmental effects that were potentially obscured in

traditional single-metabolite models.

Soil moisture and antimicrobial treatments influence rhizosphere fungal diversity and community composition

Amplicon sequencing and rarefaction identified 359 fungal OTUs in the *G. guidonia* rhizosphere,

spanning 8 phyla, 18 classes, 59 orders, 130 families, 144 genera, and 200 species. Fungal

Shannon diversity was significantly higher under low moisture relative to well-watered

conditions ($t = 2.33$, $p = 0.022$; Fig. 3a). In contrast, no significant differences in Shannon or

other *alpha* diversity metrics were observed across microbial legacy, antimicrobial, or plant

density treatments ($p > 0.05$; Fig. S3–S4). At the family level, antimicrobial treatments induced

compositional shifts, with distinct variation in fungal taxa across treatments (Fig. 3b). To assess

fungal community responses to experimental treatments, we applied principal coordinate

analysis (PCoA) on Bray–Curtis dissimilarity matrices followed by PERMANOVA. Fungal

community composition differed significantly across antimicrobial treatments ($p = 0.008$), but

not across soil inoculum source, plant density, or moisture when considered individually ($p >$

0.05 for all). However, a significant interaction effect ($p = 0.005$; Fig. 3c) emerged when all

treatment variables were considered jointly, suggesting that fungal assemblages are shaped by

the combined influence of combined environmental alteration.

Next, we used ANCOM-BC to examine family-level shifts in fungal differential abundance across

antimicrobial treatments, controlling for microbial legacy, plant density, and soil moisture. Taxa

from control pots (no treatment on soil inoculum) served as the baseline for differential

abundance analysis. We found that the soil fungicide treatment resulted in elevated abundance

of Hydnodontaceae and Stachybotryaceae, while heat sterilization caused significant

enrichment of Polyporaceae and Glomerellaceae. The foliar fungicide treatment increased the

abundance of Bionectriaceae and Junewangiaceae. Notably, Hydnodontaceae was enriched

under soil fungicide but depleted in heat and foliar fungicide treatments, whereas

Glomerellaceae showed an opposite trend, enriched under heat treatment but depleted by

fungicides. These patterns (Fig. 3d) support our earlier findings that antimicrobial treatments

distinctly shape rhizosphere fungal assemblages.

Fig. 3 Associations between fungal communities and root exudate metabolites. (a–h) Mantel

test results showing correlations between fungal Bray–Curtis dissimilarity with (a) all

metabolites, (b–h) individual metabolite classes (x-axis). (i) Partial Spearman's rank correlations

between fungal Shannon diversity and individual metabolites, adjusted for soil inoculum,

antimicrobial treatments, soil moisture, and plant density. Red-positive associations; Blue:

negative associations. Dotted line indicates p = 0.05 threshold. Top 3 positively and negatively

associated metabolites are labeled. *-6,10a-dihydroxy-4-(hydroxymethyl)-4,7,11b-trimethyl-

1,2,3,4a,5,6,6a,7,11,11a-decahydronaphtho[2,1-f][1]benzofuran-9-one.

Metabolite composition is associated with fungal community structure

To assess the relationship between rhizosphere metabolite composition and fungal community

structure, we used redundancy analysis (RDA), with the first 10 metabolite principal

components (PCs) as explanatory variables. These PCs captured most of the variation in

metabolite profiles while reducing dimensionality. The RDA model explained a modest

proportion of fungal community variation (R^2 = 0.0614; Adjusted R^2 = –0.00008; Fig. S5a), with

PC2 and PC10 contributing most to RDA1, and PC1 and PC4 to RDA2. Most PCs were

characterized by predominantly positive metabolite loadings, suggesting that metabolite

patterns captured by these axes were broadly associated with shifts in fungal composition (Fig.

S5b). While PCs reflect underlying metabolite structure, RDA scores reflect taxon responses to

those multivariate gradients. References to PC contributions to RDA axes illustrate the strength

of association between metabolite-derived variation and community composition, rather than a

direct mapping of metabolite loadings onto fungal taxa. Outlier fungal taxa, identified via

Mahalanobis distance, included 64 OTUs. These taxa were predominantly from phylum

Ascomycota (84%), with Nectriaceae making up over half of the total (Fig. S5c). To complement

this multivariate approach, we used Mantel tests to assess the strength of correlation between

fungal community distances and metabolite dissimilarity. We found a significant positive

correlation between global metabolite composition (all 612 features) and fungal community

structure (r = 0.006, p < 0.001; Fig. 4a). Importantly, all seven metabolite classes showed

statistically significant associations with fungal distances (p <0.05 for all comparisons; Fig. 4b–

h). These results indicate a strong covariation between rhizosphere fungal communities and

exuded metabolite profiles, consistent with potential reciprocal influences between plant

chemistry and microbial community composition. To explore whether metabolite profiles were

associated with fungal diversity rather than just composition, we used partial Spearman's rank

correlation models adjusted for soil inoculum source, antimicrobial treatments, soil moisture,

and plant density. We identified several positive and negative metabolite associations with

fungal Shannon diversity (Fig. 4i). However, none remained significant after multiple testing

(FDR >0.05). We also checked for additional metabolite vs fungal association patterns, focusing

on consensus metabolite cluster values, and observed that Clusters 2 and 3 exhibited significant

positive correlations with fungal Shannon diversity (p < 0.05), whereas Cluster 1 had a weak

negative association (Table S1).

Fig. 4 Associations between seedling trait divergence and metabolite and fungal community

dissimilarity. (a–b) Mantel test results showing pairwise correlations between seedling trait

distances with (a) root exudate metabolite dissimilarity and (b) fungal community dissimilarity.

(c–d) Heatmaps summarizing the frequency of treatment pairings contributing to high-
divergence sample pairs extracted from the upper 25% quantile (≥75th percentile) for both trait
vs metabolite or fungal distances. Warmer colors represent greater co-occurrence of treatment
combinations among highly divergent samples, highlighting treatment-driven shifts in
associations.

Association between seedling traits with exudate metabolites and rhizosphere
fungal communities

All studied seedling traits, with the exception of dry root mass, did not significantly vary with
experimental treatments (Fig. S6-7 and Table S2). To test whether belowground processes were
linked to trait expression (H4), we correlated seedling traits with exudate metabolite and fungal
community data. Seedling traits showed no significant association with metabolite diversity
(Functional Hill index). In contrast, trait dissimilarity was significantly correlated with metabolite
composition (Mantel R^2 = 0.018, p < 0.001) (Fig. 5a), suggesting that compositional shifts mirror
morphological variation more strongly than diversity metrics alone. With regards to the
metabolite clusters, leaf surface area exhibited the strongest association with consensus cluster
values, negatively correlating with Cluster 1 and positively with Clusters 2 and 3 (FDR-adjusted
p<0.005), suggesting possible trade-offs in carbon dynamics. Associations with dry root mass
and specific leaf area were weaker and less consistent (Table S1). Fungal community
composition was significantly correlated with trait distance (Mantel R^2 = 0.001, p = 0.0031; Fig.
5b), while no association was observed between fungal Shannon diversity and seedling traits (p
>0.05).

To further assess treatment-driven divergence, we compared high-dissimilarity sample pairs
from Mantel plots, focusing on points in the ≥75th percentile in both axes across
metabolite/trait and fungal/trait associations. Trait–metabolite divergence was strongest in
comparisons involving heat-treated inoculum (particularly when compared to control and foliar
fungicide treatments), highlighting a strong legacy effect of microbial depletion or abiotic stress
on exudation and plant development. By contrast, soil fungicide treatments showed little trait–

metabolite divergence, suggesting that AMF-specific disruption had limited impact. Trait–fungal

divergence was more broadly distributed, with mild increases in pairs involving foliar fungicide

and heat treatments (Fig. 5c–d). Overall, these results suggest that root exudate metabolites

are more tightly linked to seedling trait variation than fungal communities, with treatment

history potentially playing a key role in shaping these interactions.

Fig. 5 Associations between seedling trait divergence and metabolite and fungal community

dissimilarity. (a–b) Mantel test results showing pairwise correlations between seedling trait

distances with (a) root exudate metabolite dissimilarity and (b) fungal community dissimilarity.

(c–d) Heatmaps summarizing the frequency of treatment pairings contributing to high-

divergence sample pairs extracted from the upper 25% quantile (≥75[th] percentile) for both trait

vs metabolite or fungal distances. Warmer colors represent greater co-occurrence of treatment

combinations among highly divergent samples, highlighting treatment-driven shifts in

associations.

Discussion

Tropical forests face simultaneous environmental disturbances (Bennett et al., 2021; Lugo,
2008), whose impacts have broad implications for global ecosystem function. While
aboveground biodiversity responses are more frequently studied (Malhi et al., 2014),
belowground processes- particularly those linking root exudation, microbial assembly, and soil
biotic legacies- remain underexplored. Root exudates are key drivers of rhizosphere
microbiome composition and can influence plant resilience to biotic and abiotic stresses (Badri
& Vivanco, 2009; Canarini et al., 2019), yet how climate-induced shifts in microbial communities
feed back into these interactions remains poorly understood.

Here, we examined how warming-induced soil microbial legacy influences metabolite
composition in root exudates and associated belowground interactions in a tropical tree
species. Using inoculum from experimentally warmed and ambient plots, combined with
manipulations of fungal communities, plant density, and soil moisture, we found that exudate
metabolite–fungi relationships are shaped by specific treatments rather than strong and broad
global correlations. Metabolite diversity was significantly influenced by warming-induced
microbial legacy and high moisture, whereas composition shifted with microbial legacy and
antimicrobial treatments. Variations in metabolite profiles coincided with shifts in fungal
community structure, particularly among families such as Hydnodontaceae, Glomerellaceae,
and Nectriaceae. In addition, metabolite classes were significantly correlated with fungal
composition. While fungal communities had modest relationships with seedling traits,
metabolite profiles exhibited weak but significant associations with trait variation, particularly
leaf surface area. Together, our results point to a context-dependent relationship between root
exudate metabolites and fungal communities, structured by both abiotic stress and microbial
legacy. These findings underscore the complexity of belowground interactions in tropical
systems and highlight root exudates as early biochemical signals in plant responses to
environmental change.

444 Biotic and abiotic treatments alter plant metabolic profiles

Environmental change is a powerful driver of belowground ecological dynamics, shaping both
soil microbial communities and the diverse repertoire of plant metabolites released in root
exudates in response to abiotic and biotic stressors (Bardgett & Caruso, 2020; Hu et al., 2018;
Ma et al., 2022; Pang et al., 2021; Trivedi et al., 2022; Van der Putten et al., 2013; Yang et al.,
2025). Our results show that *G. guidonia* responds to environmental alterations through
changes in metabolite exudation and rhizosphere fungal community structure. Specifically,
microbial legacy emerged as a key determinant of metabolite composition, supporting the idea
that long-term changes in microbial communities can influence the synthesis and secretion of a
wide range of plant metabolites, with potential feedback effects on the rhizosphere
microbiome (Cao et al., 2020; Wagg et al., 2014). It is important to note that some of the
exuded compounds may originate from microbial metabolism or transformation of plant-
derived compounds, especially under fluctuating environmental conditions. These shifts likely
reflect plasticity in plant chemical investment, potentially mediated by altered nutrient cycling
or microbial signaling that regulate secondary metabolism (Badri & Vivanco, 2009; Canarini et
al., 2019). Microbe-altering treatments further modified exudate composition, possibly through
feedbacks that reconfigure carbon allocation to defense or nutrient acquisition. Abiotic factors,
such as warming and drought, have been shown to influence exudate profiles by modulating
enzyme activity and metabolite transport (Yang et al., 2025). Collectively, these findings point
to the integrative nature of plant metabolic responses to environmental and microbial contexts.

464 Environmental conditions influence fungal diversity, composition, and abundance

Fungal taxa exhibit strong ecological differentiation, often shaped by subtle biotic and abiotic
variations that reflect their functional roles and life history traits (Bahram & Netherway, 2022;
Bui et al., 2020). These patterns intensify under environmental disturbance, where altered soil
chemistry and microbial interactions act as filters, favoring competitively dominant or stress-
tolerant taxa (Delgado-Baquerizo et al., 2020). In this study, higher soil moisture was associated
with greater fungal diversity, likely due to improved conditions for microbial activity and plant–

microbe interactions. This aligns with previous findings showing that water limitation can
induce persistent shifts in forest microbial communities (Jaeger et al., 2023). Differential
abundance analyses further revealed treatment-driven restructuring of fungal assemblages,
consistent with taxon-specific ecological strategies. For instance, Hydnodontaceae was
enriched in soil fungicide treatments (likely reflecting reduced AMF), but depleted under foliar
fungicide and heat sterilization. This suggests potential dependency on complex microbial
networks involving both AMF and pathogens. In contrast, Glomerellaceae increased in heat-
treated soils, possibly exploiting reduced microbial competition, but declined under fungicide
applications, indicating sensitivity to broader microbial disruptions. These shifts reflect
differential responses to altered resource and interaction landscapes. Nectriaceae, a
predominantly pathogenic family (Lombard et al., 2015), also responded strongly to biotic and
abiotic treatments, with elevated prevalence under altered conditions. This suggests that
community restructuring may create ecological opportunities for certain taxa, potentially with
implications for plant health and rhizosphere dynamics (Osborne et al., 2018; Taylor et al.,
2014). Taken together, these results highlight fine-scale functional differentiation within fungal
communities, shaped by variation in stress tolerance, resource use, and microbe–microbe
interactions.

Fungal community structure and metabolite composition are tightly linked

Root exudates shape rhizosphere microbial assembly through the release of primary and
secondary metabolites that recruit or deter specific taxa (Leach et al., 2017; Nobori et al.,
2018). In this study, *G. guidonia* metabolites were significantly associated with rhizosphere
fungal community structure, reinforcing the role of exudates as key mediators of belowground
plant–microbe interactions (Broeckling et al., 2008; Zhalnina et al., 2018). Mantel tests revealed
significant associations between fungal communities and all major metabolite classes, namely
alkaloids, carbohydrates, fatty acids, polyketides, shikimates and phenylpropanoids, amino
acids, and terpenoids. This suggests that diverse compound groups contribute to shaping fungal
assemblages, supporting the idea that chemically diverse exudates serve as ecological filters
that help select for fungi that metabolize or tolerate particular compounds (Clocchiatti et al.,

2021; Pang et al., 2021). Positive correlations between metabolite and fungal diversity also

suggest that richer chemical environments may support higher microbial diversity, likely

through expanded resource niches or signaling pathways (Zhalnina et al., 2018). However,

redundancy analysis showed that aggregate metabolomic profiles only explained a small

proportion of fungal community variation, implying that specific metabolite–taxon associations

rather than global chemical shifts could be the primary ecological drivers. While fungal

composition was sensitive to microbe-altering treatments (e.g., fungicides), it was not strongly

linked to overall metabolite diversity. This implies an asymmetric relationship where

metabolites could be playing a dominant role in structuring early fungal communities although

fungal feedbacks may be limited at this seedling stage. Fine-scale pairwise analyses revealed

significant associations between individual metabolites and specific fungal OTUs, underscoring

the importance of targeted interactions that may be masked in community-level metrics. Taken

together, these results point to a close link between rhizosphere fungal communities and

metabolite composition. However, given that soil inoculum (microbial legacy) also influenced

metabolite profiles, the directionality of these associations is likely context-dependent and may

reflect complex feedbacks between plant exudation and microbial community dynamics.

Metabolite and fungal variation is linked to seedling traits

Although overall treatment effects on seedling traits were modest, we observed significant

associations between metabolite clusters, metabolite principal components, and plant

morphological traits. Fungal community structure was also significantly correlated with trait

variation. These findings suggest that belowground chemical and microbial variation may

influence early plant phenotypes, albeit subtly and in a context-dependent manner. Notably,

leaf surface area and specific leaf area (SLA) exhibited clear relationships with metabolite

composition. Specifically, SLA was positively associated with Cluster 1 metabolites, which was

dominated by secondary metabolites, and negatively associated with Clusters 2 and 3 that were

primarily enriched in primary metabolites (Fig. S2b). This pattern may reflect shifts in carbon

allocation strategy, where seedlings with thinner leaves potentially prioritize the production of

secondary metabolites such as phenolics and terpenoids, which have known roles in defense

and microbial signaling (Badri & Vivanco, 2009). In contrast, seedlings associated with Clusters 2

and 3 may favor primary metabolism that supports nutrient uptake, microbial facilitation, or

root biomass production. These divergent associations suggest possible trade-offs between

growth and defense, consistent with broader patterns in plant functional trait ecology (Freschet

et al., 2015). Associations with dry root mass followed a similar but less consistent trend,

suggesting a potential but weaker role for belowground allocation in shaping exudate profiles.

The stronger correlation between metabolite composition and trait dissimilarity, compared to

that between metabolite richness and traits, highlights the ecological relevance of qualitative

shifts in exudate chemistry. This indicates that variation in the type, rather than the number, of

metabolites more strongly influences plant morphological outcomes.

Fungal community composition also correlated with trait dissimilarity, albeit with lower

explanatory power, suggesting a more diffuse or indirect role that is potentially mediated by

altered microbial signaling or nutrient dynamics. Such effects may reflect indirect or delayed

induction of pathways through which rhizosphere composition shapes trait expression,

consistent with prior evidence that plant–microbe interactions often intensify over time or

under prolonged stress (Ke et al., 2021; Ravanbakhsh et al., 2019). For instance, trait–

metabolite divergence was strongest in comparisons involving heat-treated inoculum,

indicating a legacy effect of microbial depletion or abiotic stress. Conversely, soil fungicide

treatments caused relatively little divergence, suggesting that selective AMF suppression alone

had limited influence on trait expression or exudate patterns within the experimental window.

Trait–fungal divergence was more broadly distributed but less pronounced, further supporting

a primary role for exudate composition in driving morphological outcomes. Overall, our results

suggest that seedling trait variation is more closely linked to root exudate composition than

rhizosphere fungal diversity, and that treatment history plays a key role in structuring these

interactions. Compositional shifts in the exudate profile, particularly those involving secondary

metabolites, may represent an underrecognized axis of adaptive plasticity during early plant

development. Future work across ontogeny and in field conditions will be essential to resolve

the ecological and mechanistic significance of these belowground–aboveground trait linkages.

Conclusion

This study reveals how environmental conditions reshape plant–soil–microbe interactions by altering both root exudate chemistry and fungal community composition in a tropical tree species. We show that treatment effects, particularly warming-induced microbial legacy and soil moisture, modulate root exudate metabolite profiles and that microbial manipulations significantly shift metabolite composition and fungal structure. These shifts suggest that root exudation is governed by a dynamic interplay between abiotic conditions and microbial context, with metabolites in turn acting as ecological filters that shape fungal assemblages. While our data reveal significant associations between metabolite composition and fungal community structure, links to seedling traits were weak at this developmental stage suggesting that belowground feedbacks influencing plant morphology may be subtle during early growth. It is possible that the association could strengthen over time or under cumulative environmental stress. The responses of distinct fungal families, such as Hydnodontaceae and Nectriaceae, further support the idea of functional filtering driven by environmental and microbial cues. Taken together, our results underscore the need to consider metabolite-mediated pathways in understanding plant resilience to climate change. They highlight the importance of multifactorial approaches that integrate abiotic stress, microbial dynamics, and metabolic plasticity. Future work should examine how these interactions evolve over time, how different microbial guilds contribute, and how specific metabolite pathways influence plant fitness and ecosystem recovery under ongoing global change.

Acknowledgements

This work was funded by NSF grant DEB 2120085, Department of Energy (DE-SC0012000, DE-SC-0011806, 89243018S-SC-000014, 89243018S-SC-000017, DE-SC-0018942, DE-SC0022095, 89243021S-SC-000076) and NSF (DEB-1754713). We would like to acknowledge the Pennsylvania State University's Huck Metabolomics Core Facility (RRID:SCR_023864) for use of the Vanquish Horizon UHPLC System and Ashley Shay for helpful discussions on sample

581 preparation. We also thank Shelby McMahan, William Mejía, Alberto Ibáñez, Deyaneira Iglesias,

582 and Laura Rubio for assistance with shade house experiments.

Competing interests

584 None declared.

Author contributions

586 **JRL**, and **BB** conceptualized and designed the study and acquired funding. **JM**, **PB** and **GHV**

587 performed shade house experiments, with input from **IFG**. **JM** processed samples for

588 metabolomics analysis. **BB** oversaw sample processing for fungal DNA sequencing. **JM** analyzed

589 the data with input from **JRL** and **BB**. **JM** wrote the manuscript. All authors provided feedback

590 on the manuscript.

Data availability

592 All data supporting the findings of this study are included in the manuscript and supplementary

593 information files. Code used for data analysis is found at

594 https://github.com/Jmasanga/Guarea_guidonia_analysis

References

Abarenkov, K., Nilsson, R. H., Larsson, K. H., Taylor, A. F. S., May, T. W., Frøslev, T. G., Pawlowska,

J., Lindahl, B., Põldmaa, K., Truong, C., Vu, D., Hosoya, T. et al. (2024). The UNITE database for

molecular identification and taxonomic communication of fungi and other eukaryotes:

sequences, taxa and classifications reconsidered. Nucleic Acids Research, 52(D1), D791–D797.

Alonso-Rodríguez, A. M., Wood, T. E., Torres-Díaz, J., Cavaleri, M. A., Reed, S. C., & Bachelot, B.

(2022). Understory plant communities show resistance to drought, hurricanes, and

experimental warming in a wet tropical forest. Frontiers in Forests and Global Change, 5,

733967.

Andrewin, A. N., Rodriguez-Llanes, J. M., & Guha-Sapir, D. (2015). Determinants of the lethality of

climate-related disasters in the Caribbean Community (CARICOM): a cross-country analysis.

Scientific Reports 2015 5:1, 5(1), 1–10.

Bachelot, B., Alonso-Rodríguez, A. M., Aldrich-Wolfe, L., Cavaleri, M. A., Reed, S. C., & Wood, T. E.
(2020). Altered climate leads to positive density-dependent feedbacks in a tropical wet forest.
Global Change Biology, 26(6), 3417–3428.

Badri, D. V., & Vivanco, J. M. (2009). Regulation and function of root exudates. Plant, Cell and
Environment, 32(6), 666–681.

Bahram, M., & Netherway, T. (2022). Fungi as mediators linking organisms and ecosystems. FEMS
Microbiology Reviews, 46(2), 1–16.

Bardgett, R. D., & Caruso, T. (2020). Soil microbial community responses to climate extremes:
resistance, resilience and transitions to alternative states. Philosophical Transactions of the
Royal Society B, 375(1794).

Bennett, A. C., Dargie, G. C., Cuni-Sanchez, A., Mukendi, J. T., Hubau, W., Mukinzi, J. M., Phillips,
O. L., Malhi, Y., Sullivan, M. J. P., Cooper, D. L. M., et al. (2021). Resistance of African tropical
forests to an extreme climate anomaly. Proceedings of the National Academy of Sciences of
the United States of America, 118(21).

Bennett, A. J., Bending, G. D., Chandler, D., Hilton, S., & Mills, P. (2012). Meeting the demand for
crop production: The challenge of yield decline in crops grown in short rotations. Biological
Reviews, 87(1), 52–71.

Berendsen, R. L., Pieterse, C. M. J., & Bakker, P. A. H. M. (2012). The rhizosphere microbiome and
plant health. Trends in Plant Science, 17(8), 478–486.

Bertness, M. D., & Callaway, R. (1994). Positive interactions in communities. Trends in Ecology
and Evolution, 9(5), 191–193. https://doi.org/10.1016/0169-5347(94)90088-4

Bever, J. D. (1994). Feedback between plants and their soil communities in an old field
community. Ecology, 75(7), 1965–1977.

Bhardwaj, A., Misra, V., Mishra, A., Wootten, A., Boyles, R., Bowden, J. H., & Terando, A. J. (2018).
Downscaling future climate change projections over Puerto Rico using a non-hydrostatic
atmospheric model. Climatic Change, 147(1–2), 133–147.

Broeckling, C. D., Broz, A. K., Bergelson, J., Manter, D. K., & Vivanco, J. M. (2008). Root exudates
regulate soil fungal community composition and diversity. Applied and Environmental
Microbiology, 74(3), 738–744.

Brooker, R. W., George, T. S., Homulle, Z., Karley, A. J., Newton, A. C., Pakeman, R. J., & Schöb, C.

(2021). Facilitation and biodiversity–ecosystem function relationships in crop production

systems and their role in sustainable farming. Journal of Ecology, 109(5), 2054–2067.

Bui, A., Orr, D., Lepori-Bui, M., Konicek, K., Young, H. S., & Moeller, H. V. (2020). Soil fungal

community composition and functional similarity shift across distinct climatic conditions. FEMS

Microbiology Ecology, 96(12), 193.

Canarini, A., Kaiser, C., Merchant, A., Richter, A., & Wanek, W. (2019). Root exudation of primary

metabolites: Mechanisms and their roles in plant responses to environmental stimuli. Frontiers

in Plant Science, 10, 422679.

Cao, J., Lin, T. C., Yang, Z., Zheng, Y., Xie, L., Xiong, D., & Yang, Y. (2020). Warming exerts a

stronger effect than nitrogen addition on the soil arbuscular mycorrhizal fungal community in

a young subtropical Cunninghamia lanceolata plantation. Geoderma, 367, 114273.

Chiu C-H., & Chao A. (2014). Distance-based functional diversity measures and their

decomposition: a framework based on hill numbers. PLoS ONE 9: e100014.

Clocchiatti, A., Hannula, S. E., van den Berg, M., Hundscheid, M. P. J., & de Boer, W. (2021).

Evaluation of Phenolic Root Exudates as Stimulants of Saptrophic Fungi in the Rhizosphere.

Frontiers in Microbiology, 12, 644046.

Delgado-Baquerizo, M., Reich, P. B., Trivedi, C., Eldridge, D. J., Abades, S., Alfaro, F. D., Bastida, F.,

Berhe, A. A., Cutler, N. A., Gallardo, A., García-Velázquez, L., Hart, S. C., et al. (2020). Multiple

elements of soil biodiversity drive ecosystem functions across biomes. Nature Ecology &

Evolution 2020 4:2, 4(2), 210–220.

Eaton, W. D., McGee, K. M., Alderfer, K., Jimenez, A. R., & Hajibabaei, M. (2020). Increase in

abundance and decrease in richness of soil microbes following Hurricane Otto in three primary

forest types in the Northern Zone of Costa Rica. PLOS ONE, 15(7), e0231187.

Elger, A., Lemoine, D. G., Fenner, M., & Hanley, M. E. (2009). Plant ontogeny and chemical

defence: Older seedlings are better defended. Oikos, 118(5), 767–773.

Emanuel, K. (2005). Increasing destructiveness of tropical cyclones over the past 30 years. Nature,

436(7051), 686–688.

Engelkes, T., Morriën, E., Verhoeven, K. J. F., Bezemer, T. M., Biere, A., Harvey, J. A., McIntyre, L.
M., Tamis, W. L. M., & Van Der Putten, W. H. (2008). Successful range-expanding plants
experience less above-ground and below-ground enemy impact. Nature, 456(7224), 946–948.
Freire-Zapata, V., Holland-Moritz, H., Cronin, D. R., Aroney, S., Smith, D. A., Wilson, R. M.,
Ernakovich, J. G., Woodcroft, B. J., Bagby, S. C., Mondav, R., et al. (2024). Microbiome–
metabolite linkages drive greenhouse gas dynamics over a permafrost thaw gradient. Nature
Microbiology 2024 9:11, 9(11), 2892–2908.
Freschet, G. T., Swart, E. M., & Cornelissen, J. H. C. (2015). Integrated plant phenotypic responses
to contrasting above- and below-ground resources: Key roles of specific leaf area and root
mass fraction. New Phytologist, 206(4), 1247–1260.
Gweon, H. S., Oliver, A., Taylor, J., Booth, T., Gibbs, M., Read, D. S., Griffiths, R. I., & Schonrogge,
K. (2015). PIPITS: An automated pipeline for analyses of fungal internal transcribed spacer
sequences from the Illumina sequencing platform. Methods in Ecology and Evolution, 6(8),
973–980.
Gfeller, V., Waelchli, J., Pfister, S., Deslandes-Hérold, G., Mascher, F., Glauser, G., Aeby, Y.,
Mestrot, A., Robert, C. A. M., Schlaeppi, K., & Erb, M. (2023). Plant secondary metabolite-
dependent plant-soil feedbacks can improve crop yield in the field. ELife, 12, e84988.
Gidman, E. A., Goodacre, R., Emmett, B., Wilson, D. B., Carroll, J. A., Caporn, S. J. M., Cresswell, N.,
& Gwynn-Jones, D. (2005). Metabolic fingerprinting for bio-indication of nitrogen responses in
Calluna vulgaris heath communities. Metabolomics, 1(3), 279–285.
Gidman, E., Goodacre, R., Emmett, B., Smith, A. R., & Gwynn-Jones, D. (2003). Investigating
plant–plant interference by metabolic fingerprinting. Phytochemistry, 63(6), 705–710.
Hannula, S. E., Heinen, R., Huberty, M., Steinauer, K., De Long, J. R., Jongen, R., & Bezemer, T. M.
(2021). Persistence of plant-mediated microbial soil legacy effects in soil and inside roots.
Nature Communications, 12(1), 1–13.
Hu, L., Robert, C. A. M., Cadot, S., Zhang, X., Ye, M., Li, B., Manzo, D., Chervet, N., Steinger, T., Van
Der Heijden, M. G. A., Schlaeppi, K., & Erb, M. (2018). Root exudate metabolites drive plant-
soil feedbacks on growth and defense by shaping the rhizosphere microbiota. Nature
Communications 2018 9:1, 9(1), 1–13.

Jaeger, A. C. H., Hartmann, M., Six, J., & Solly, E. F. (2023). Contrasting sensitivity of soil bacterial
and fungal community composition to one year of water limitation in Scots pine mesocosms.
FEMS Microbiology Ecology, 99(6), 1–17.

Jiang, P., Wang, Y., Zhang, Y., Fei, J., Rong, X., Peng, J., Yin, L., & Luo, G. (2024). Intercropping
enhances maize growth and nutrient uptake by driving the link between rhizosphere
metabolites and microbiomes. New Phytologist, 243(4), 1506–1521.

Jones, O. A. H., Maguire, M. L., Griffin, J. L., Jung, Y. H., Shibato, J., Rakwal, R., Agrawal, G. K., &
Jwa, N. S. (2011). Using metabolic profiling to assess plant-pathogen interactions: An example
using rice (Oryza sativa) and the blast pathogen Magnaporthe grisea. European Journal of
Plant Pathology, 129(4), 539–554.

Ke, P. J., Zee, P. C., & Fukami, T. (2021). Dynamic plant–soil microbe interactions: the neglected
effect of soil conditioning time. New Phytologist, 231(4), 1546–1558.

Kim, S. (2015). ppcor: An R Package for a Fast Calculation to Semi-partial Correlation Coefficients.
Communications for Statistical Applications and Methods, 22(6), 665.

Kimball, B. A., Alonso-Rodríguez, A. M., Cavaleri, M. A., Reed, S. C., González, G., & Wood, T. E.
(2018). Infrared heater system for warming tropical forest understory plants and soils. Ecology
and Evolution, 8(4), 1932–1944.

Lake, J. A., Field, K. J., Davey, M. P., Beerling, D. J., & Lomax, B. H. (2009). Metabolomic and
physiological responses reveal multi-phasic acclimation of Arabidopsis thaliana to chronic UV
radiation. Plant, Cell and Environment, 32(10), 1377–1389.

Leach, J. E., Triplett, L. R., Argueso, C. T., & Trivedi, P. (2017). Communication in the Phytobiome.
Cell, 169(4), 587–596.

Lin, H., & Peddada, S. D. (2024). Multigroup analysis of compositions of microbiomes with
covariate adjustments and repeated measures. Nature Methods, 21(1), 83–91.

Lombard, L., van der Merwe, N. A., Groenewald, J. Z., & Crous, P. W. (2015). Generic concepts in
Nectriaceae. Studies in Mycology, 80, 189–245.

Lugo, A. E. (2008). Visible and invisible effects of hurricanes on forest ecosystems: An
international review. Austral Ecology, 33(4), 368–398.

Ma, W., Tang, S., Dengzeng, Z., Zhang, D., Zhang, T., & Ma, X. (2022). Root exudates contribute to
belowground ecosystem hotspots: A review. Frontiers in Microbiology, 13, 937940.

Majumdar, S., Kaur, H., Rinella, M. J., Kundu, A., Vadassery, J., Erbilgin, N., Callaway, R. M.,
Cadotte, M. W., & Inderjit. (2023). Synergistic effects of canopy chemistry and autogenic soil
biota on a global invader. Journal of Ecology, 111(7), 1497–1513.

Malhi, Y., Gardner, T. A., Goldsmith, G. R., Silman, M. R., & Zelazowski, P. (2014). Tropical forests
in the anthropocene. Annual Review of Environment and Resources, 39(Volume 39, 2014),
125–159.

Maron, J. L., Smith, A. L., Ortega, Y. K., Pearson, D. E., & Callaway, R. M. (2016). Negative plant-
soil feedbacks increase with plant abundance, and are unchanged by competition. Ecology,
97(8), 2055–2063.

McMurdie, P. J., & Holmes, S. (2013). phyloseq: An R Package for Reproducible Interactive
Analysis and Graphics of Microbiome Census Data. PLOS ONE, 8(4), e61217.

Melillo, J. M., Frey, S. D., DeAngelis, K. M., Werner, W. J., Bernard, M. J., Bowles, F. P., Pold, G.,
Knorr, M. A., & Grandy, A. S. (2017). Long-term pattern and magnitude of soil carbon feedback
to the climate system in a warming world. Science, 358(6359), 101–105.

Müller, C., & Junker, R. R. (2022). Chemical phenotype as important and dynamic niche dimension
of plants. New Phytologist, 234(4), 1168–1174.

Nguyen, N. H., Song, Z., Bates, S. T., Branco, S., Tedersoo, L., Menke, J., Schilling, J. S., & Kennedy,
P. G. (2016). FUNGuild: An open annotation tool for parsing fungal community datasets by
ecological guild. Fungal Ecology, 20, 241–248.

Nobori, T., Mine, A., & Tsuda, K. (2018). Molecular networks in plant–pathogen holobiont. FEBS
Letters, 592(12), 1937–1953.

Noecker, C., Chiu, H.-C., McNally, C. P., & Borenstein, E. (2019). Defining and Evaluating Microbial
Contributions to Metabolite Variation in Microbiome-Metabolome Association Studies.
MSystems, 4(6).

Osborne, O. G., De-Kayne, R., Bidartondo, M. I., Hutton, I., Baker, W. J., Turnbull, C. G. N., &
Savolainen, V. (2018). Arbuscular mycorrhizal fungi promote coexistence and niche divergence
of sympatric palm species on a remote oceanic island. New Phytologist, 217(3), 1254–1266.

Pang, Z., Chen, J., Wang, T., Gao, C., Li, Z., Guo, L., Xu, J., & Cheng, Y. (2021). Linking Plant
Secondary Metabolites and Plant Microbiomes: A Review. Frontiers in Plant Science, 12.

Pang, Z., Lu, Y., Zhou, G., Hui, F., Xu, L., Viau, C., Spigelman, A. F., Macdonald, P. E., Wishart, D. S.,
Li, S., & Xia, J. (2024). MetaboAnalyst 6.0: towards a unified platform for metabolomics data
processing, analysis and interpretation. Nucleic Acids Research, 52(W1), W398–W406.

Pennington, T. D., & Clarkson, J. J. (2013). A REVISION OF GUAREA (MELIACEAE). Edinburgh
Journal of Botany, 70(2), 179–362.

Petrén, H., Köllner, T. G., & Junker, R. R. (2023). Quantifying chemodiversity considering
biochemical and structural properties of compounds with the R package chemodiv. New
Phytologist, 237(6), 2478–2492.

Prabhakaran, A. D., & Dann, E. K. (2022). Evaluation of Fungicide Soil Drench Treatments to
Manage Black Root Rot Disease of Avocado. Plant Disease, 106(8), 2026–2030.

Ravanbakhsh, M., Kowalchuk, G. A., & Jousset, A. (2019). Root-associated microorganisms
reprogram plant life history along the growth–stress resistance tradeoff. ISME Journal, 13(12),
3093–3101.

Reed, S. C., Reibold, R., Cavaleri, M. A., Alonso-Rodríguez, A. M., Berberich, M. E., & Wood, T. E.
(2020). Soil biogeochemical responses of a tropical forest to warming and hurricane
disturbance. Advances in Ecological Research, 62, 225–252.

Robertson, K. M., & Platt, W. J. (2001). Effects of multiple disturbances (fire and hurricane) on
epiphyte community dynamics in a subtropical forest, Florida, U.S.A. Biotropica, 33(4), 573–
582.

Schindlbacher, A., Rodler, A., Kuffner, M., Kitzler, B., Sessitsch, A., & Zechmeister-Boltenstern, S.
(2011). Experimental warming effects on the microbial community of a temperate mountain
forest soil. Soil Biology and Biochemistry, 43(7), 1417–1425.

Tamburini, G., Bommarco, R., Wanger, T. C., Kremen, C., van der Heijden, M. G. A., Liebman, M.,
& Hallin, S. (2020). Agricultural diversification promotes multiple ecosystem services without
compromising yield. Science Advances, 6(45).

Taylor, D. L., Hollingsworth, T. N., McFarland, J. W., Lennon, N. J., Nusbaum, C., & Ruess, R. W.
(2014). A first comprehensive census of fungi in soil reveals both hyperdiversity and fine-scale
niche partitioning. Ecological Monographs, 84(1), 3–20.
Thakur, M. P., van der Putten, W. H., Wilschut, R. A., Veen, G. F. (Ciska), Kardol, P., van Ruijven, J.,
Allan, E., Roscher, C., van Kleunen, M., & Bezemer, T. M. (2021). Plant–Soil Feedbacks and
Temporal Dynamics of Plant Diversity–Productivity Relationships. Trends in Ecology &
Evolution, 36(7), 651–661.
Trivedi, P., Batista, B. D., Bazany, K. E., & Singh, B. K. (2022). Plant–microbiome interactions under
a changing world: responses, consequences and perspectives. New Phytologist, 234(6), 1951–
1959.
Van Der Heijden, M. G. A., Bardgett, R. D., & Van Straalen, N. M. (2008). The unseen majority: soil
microbes as drivers of plant diversity and productivity in terrestrial ecosystems. Ecology
Letters, 11(3), 296–310.
Van der Putten, W. H., Bardgett, R. D., Bever, J. D., Bezemer, T. M., Casper, B. B., Fukami, T.,
Kardol, P., Klironomos, J. N., Kulmatiski, A., Schweitzer, J. A., et al. (2013). Plant-soil feedbacks:
The past, the present and future challenges. Journal of Ecology, 101(2), 265–276.
van der Putten, W. H., Bradford, M. A., Pernilla Brinkman, E., van de Voorde, T. F. J., & Veen, G. F.
(2016). Where, when and how plant–soil feedback matters in a changing world. Functional
Ecology, 30(7), 1109–1121.
Vargas, R., Hasselquist, N., Allen, E. B., & Allen, M. F. (2010). Effects of a hurricane disturbance on
aboveground forest structure, arbuscular mycorrhizae and belowground carbon in a restored
tropical forest. Ecosystems, 13(1), 118–128.
Wagg, C., Bender, S. F., Widmer, F., & Van Der Heijden, M. G. A. (2014). Soil biodiversity and soil
community composition determine ecosystem multifunctionality. Proceedings of the National
Academy of Sciences of the United States of America, 111(14), 5266–5270.
Walsh, K. J. E., & Ryan, B. F. (2000). Tropical Cyclone Intensity Increase near Australia as a Result
of Climate Change. Journal of Climate, 13(16), 3029–3036.

Webster, P. J., Holland, G. J., Curry, J. A., & Chang, H. R. (2005). Atmospheric science: Changes in
tropical cyclone number, duration, and intensity in a warming environment. Science,
309(5742), 1844–1846.

Widarto, H. T., Van Der Meijden, E., Lefeber, A. W. M., Erkelens, C., Kim, H. K., Choi, Y. H., &
Verpoorte, R. (2006). Metabolomic differentiation of Brassica rapa following herbivory by
different insect instars using two-dimensional nuclear magnetic resonance spectroscopy.
Journal of Chemical Ecology, 32(11), 2417–2428.

Wright, A. J., Wardle, D. A., Callaway, R., & Gaxiola, A. (2017). The Overlooked Role of Facilitation
in Biodiversity Experiments. Trends in Ecology & Evolution, 32(5), 383–390.

Yang, C. X., Chen, S. J., Hong, X. Y., Wang, L. Z., Wu, H. M., Tang, Y. Y., Gao, Y. Y., & Hao, G. F.
(2025). Plant exudates-driven microbiome recruitment and assembly facilitates plant health
management. FEMS Microbiology Reviews, 49, 8.

Yang, L., Maron, J. L., & Callaway, R. M. (2015). Inhibitory effects of soil biota are ameliorated by
high plant diversity. Oecologia, 179(2), 519–525.

Zhalnina, K., Louie, K. B., Hao, Z., Mansoori, N., Da Rocha, U. N., Shi, S., Cho, H., Karaoz, U., Loqué,
D., Bowen, B. P., et al. (2018). Dynamic root exudate chemistry and microbial substrate
preferences drive patterns in rhizosphere microbial community assembly. Nature Microbiology
2018 3:4, 3(4), 470–480.

822