Many facets of self-directed behaviours in macaques

Debottam Bhattacharjee[1,2,3]*, Eva S.J. van Dijk[1], Roos E.M. de Vries[1], Charlotte E. Kluiver[1],
Mia Valomy[4], Adam N. Zeeman[5,6], and Jorg J.M. Massen[1,7]

Affiliations:
[1] Animal Behaviour & Cognition, Department of Biology, Utrecht University; 3584 CH Utrecht,
The Netherlands.
[2] Department of Infectious Diseases and Public Health, Jockey Club College of Veterinary
Medicine and Life Sciences, City University of Hong Kong; Kowloon, Hong Kong SAR.
[3] Centre for Animal Health and Welfare, Jockey Club College of Veterinary Medicine and Life
Sciences, City University of Hong Kong; Kowloon, Hong Kong SAR.
[4] Faculté de Médecine de Créteil, École Nationale Vétérinaire d'Alfort; 94700 Maisons-Alfort,
France.
[5] Institute of Biodiversity and Ecosystem Dynamics, Department of Evolutionary and
Population Biology, University of Amsterdam; 1098 XH Amsterdam, The Netherlands.
[6] ARTIS Amsterdam Royal Zoo; 1018 CZ Amsterdam, The Netherlands.
[7] Royal Rotterdam Zoological & Botanical Gardens; 3041 JG Rotterdam, The Netherlands.

*Corresponding author: Debottam Bhattacharjee (bhattacharjee.debottam@gmail.com)

Abstract

Emotional states like anxiety are adaptive, assisting individuals during social uncertainty and threats. While humans can self-report anxiety, assessing comparable states in animals requires objective measures. Self-directed behaviours (SDBs), including self-scratch, autogroom, and yawn, are indicators of anxiety, particularly in nonhuman primates. Yet it is unclear whether they reflect a common underlying process or distinct responses shaped by demography, social interactions and status, and the broader social dynamics. We studied 13 macaque groups across six species along a *despotic–egalitarian* social style gradient to investigate the factors associated with variation in SDBs. Our findings revealed partial correlational consistency among the SDBs, where only self-scratch and autogroom positively correlated. Further, we found that males yawned more than females, and individuals receiving less aggression exhibited longer autogroom bouts. Dominance rank showed behaviour-specific effects, with weak evidence for an association with self-scratch, longer autogrooming in subordinates, and a non-linear pattern in yawning, which was lowest at intermediate ranks. Alpha status was linked to more frequent yawning than others. And finally, social style showed no consistent effects on SDBs. Our research demonstrates that SDBs in macaques reflect differential responses related to social uncertainty rather than one anxiety dimension, providing insights into primate emotion regulation.

Keywords

Affective states, Despotism, Emotional arousal, *Macaca*, Social anxiety, Social complexity, Social unpredictability, Stress

1. Introduction

Individual emotional states are mediated by physiological and behavioural changes across species, offering benefits that range from welfare to survival [1,2]. In social contexts, a common emotional state is anxiety – an individual's response to aversive situations or threats [3,4]. While adult humans can self-report their experience of anxiety, assessing this psychosocial or affective state in animals requires objective methodological approaches. Self-directed behaviours (SDBs), defined by behaviours directed toward the self, such as self-scratch, autogroom, and yawn [5], are powerful and widely used proxies to capture anxiety-related emotional states in animals [6]. Although these behaviours can serve varying functions, from environmental to hygienic, their social functions have received considerable attention [6–10]. By testing the discriminant validity of SDBs, previous research has found their positive correlations with physiological measures of anxiety and broadly 'stress'. For instance, post-conflict SDBs positively correlated with maximum heart rate in greylag geese (*Anser anser*) [11]. In European starlings (*Sturnus vulgaris*), acute crowding led to elevated SDBs, heart rates, and corticosterone concentrations [12]. SDBs increased when long-tailed macaques (*Macaca fascicularis*) were given anxiety-inducing drugs [13,14]. However, a lack of positive correlation between SDBs and those physiological measures was also reported (see [15–17]). Nonetheless, research on various social contexts, such as during allogrooming and conflict, has found systematic fluctuations in SDBs [18–21], establishing their mediating roles in anxiety. Thus, the consensus is that SDBs are displacement behaviours that do not necessarily reflect chronic stress (but see [22]) and are instead associated with transient or short-term emotional states with *coping* benefits [5,6,11,17,23–27], for example, helping individuals overcome challenges related to stressful situations (e.g., during intragroup conflicts). Yet, considerable between-individual variation in the expression of SDBs is prevalent, and comprehensive evidence about their underlying proximate mechanisms is still poorly understood.

Demographic factors such as age and sex, as well as social factors like aggression, are known to be associated with SDBs and their variations. A decline in physical conditions with ageing can be associated with higher negative affective states [28]. Consequently, older individuals may use SDBs as a functional means of coping, or they may increase reliance on frequent hygienic activities [8,29]. By contrast, general life experience with ageing may lead to better emotional regulation [30], where individuals become more certain about their relationships with group members, potentially leading to a decrease in SDBs. Sex-level differences have also been reported, with males often exhibiting more SDBs than females [31]. For instance, yawning in males can lead to the exposure of large canines, which carry social information on dominance and promote group stability. Inter-sex competition, including aggression [32], can also potentially lead to variations in SDBs. Thus, sex-level differences can be explained by anatomical (*sensu* sexual dimorphism), physiological (such as concentration of androgens), as well as social (such as competition and aggression) factors [32–34]. However, sex-based variations in SDBs are not conclusive, as comparable rates between females and males have also been reported (see [35,36]). Moreover, aggression can influence SDBs, irrespective of sex differences [8,37–39], with a positive association between receiving frequent aggression and exhibiting higher rates of SDBs. Furthermore, many studies have analysed SDBs as a single composite index (e.g., see [17,29,40,41]), based on the assumption that the elemental behaviours (i.e., self-scratch, autogroom, and yawn) covary, or simply due to limited sampling effort, resulting in insufficient data on each

behaviour separately. Overall, while there is still a critical knowledge gap regarding how age,
sex, and aggression are associated with specific elemental anxiety-related behaviours, the
existing inconclusive evidence also points to the need to investigate other social factors.

In stable social hierarchies, social status or dominance rank may influence SDBs. In
particular, the broad mechanisms – (un-)certainty in the outcomes of within-group social
interactions and rank-related energetic constraints – can be grouped under two non-mutually
exclusive hypotheses. The uncertainty hypothesis suggests that subordinate group members
experience greater unpredictability in the outcomes of social interactions and therefore
express more frequent SDBs, as these behaviours help them cope with tense situations or
avoid conflict [38,41–45]. Particularly, such 'reactive' coping benefits may come from
heightened 'flight' responses, helping subordinates to avoid higher-ranking group members
and potential conflicts, or to seek support from closely affiliated group members [6,38,46–
48]. By contrast, the energetic constraint hypothesis posits that different ranks within a stable
hierarchy are associated with unique costs and benefits, and that stress-related responses
covary with rank [49]. Accordingly, the top rank or the 'alpha status' is linked to elevated
energetic demands (*sensu* 'executive stress syndrome') [49], from physiological costs of
mating and functional needs for vigilance (e.g., against outgroup threats, but also intragroup
competitors/challengers) to group maintenance. Empirical findings across primates reflect
both mechanisms, leading to inconclusive patterns: in some systems, high-ranking
individuals are the most stressed, particularly when dominance is maintained through
frequent physical aggression; in others, lower-ranking individuals show more stress when
hierarchies rely primarily on intimidation; and in more egalitarian societies, rank-related
stress differences may be weak or absent [50]. These contrasts highlight that the
relationship between rank and stress may depend on the species' social style, the stability of
dominance relationships, and the broader dynamics of within-group interactions [50,51].
However, comparative research investigating SDBs across phylogenetically close species
with varying social styles is limited.

The genus *Macaca* represents ~22 species of group-living primates, with similar social
organizations yet strikingly different social tolerance levels or social styles [52–54], providing
an excellent opportunity to investigate SDBs and their proximate mechanisms comparatively.
The broader social styles (interchangeably referred to as tolerance grades) in macaques are
shaped by phylogenetically co-varying structural traits [52–55], primarily reflected in how
individuals tolerate one another within a group. These co-varying traits include hierarchical
steepness of dominance, coalitionary tendencies, and nepotistic and kin biases. While
robust comparative empirical evidence is still scarce, some studies indicate that these traits
indeed covary, and macaque species or societies can be categorized into four distinct
tolerance grades along a despotic–egalitarian gradient [56–58]. Despotic societies are
primarily characterized by steeper dominance hierarchies and more frequent aggressive
interactions than egalitarian ones [56,59]. From grade one (or the despotic end) to four (or
the egalitarian end), social tolerance is considered to increase (but see [60]). Accordingly,
primatologists use these grades to distinguish the different macaque species and to test
various ecological, social, and cognitive hypotheses. As varying social styles can influence
individuals at the physiological [61], behavioural [62], and cognitive [63] levels, SDBs can
also be potentially shaped by them, and by studying SDBs at the grade level, we can
account for the phylogenetic inertia. Furthermore, rank-related uncertainties are higher in
more egalitarian than in despotic societies, which are often described as "complex" with

regard to within-group communication and interaction [64,65]. Yet, whether SDBs generally
vary across social styles remains a rarely investigated topic. Therefore, a comparative
assessment of SDBs across macaque societies representing the four social styles can
provide critical insights into their functions and, broadly, evolution.

Here we comparatively investigated three different SDBs (self-scratch, autogroom, and
yawn) across six macaque species: Japanese (*Macaca fuscata*), rhesus (*M. mulatta*), long-
tailed (*M. fascicularis*), lion-tailed (*M. silenus*), barbary (*M. sylvanus*), and crested (*M. nigra*)
macaques. These species represented all four social styles (Grade 1: *M. fuscata* and *M.*
mulatta; Grade 2: *M. fascicularis*; Grade 3: *M. silenus* and *M. sylvanus*; Grade 4: *M. nigra*, cf.
[52–54]). Using a comprehensive dataset including 113 individuals (>4 years of age), we
investigated (i) whether the three SDBs covary or exhibit correlational consistency, and (ii)
the associations of SDBs with age, sex, aggression, dominance rank, and social styles or
grades. Although we used SDB to denote the three behaviours, we analysed them
separately and at the behaviour level.

We made a two-tailed prediction for age: older individuals would exhibit either higher (due to
the general decline of physical conditions leading to increased hygienic requirements) or
lower (due to better emotion regulation and more certainty of social relationships) amounts
of SDBs. Aligning with the dimorphism hypothesis, we expected that SDBs, particularly yawn
behaviour, would differ between females and males, with males exhibiting higher SDBs than
females [33,34]. We expected a relationship between aggression and SDBs, specifically with
individuals receiving more frequent aggression exhibiting higher amounts of SDBs. Due to
the non-mutually exclusive nature of the uncertainty and energetic constraint hypotheses, we
expected complex non-linear relationships between ranks and SDBs (cf. [49]). On one hand,
we predicted that lower-ranking individuals would display higher amounts of SDBs than their
higher-ranking group members due to uncertainty in the outcomes of social interactions. On
the other hand, if higher ranks are associated with unique costs, such as elevated energy
demands for mating and vigilance, and/or exhibition of dominance (cf. [49]), then the highest
ranks or the alpha status would be associated with higher amounts of SDBs than the rest of
the group members. Finally, for social styles, we had two alternative predictions. If
despotism intensifies social stress through frequent aggressive interactions and intimidation
(cf. [52,59]), then individuals in more despotic societies are expected to exhibit higher rates
of SDBs than those in more egalitarian societies. Alternatively, if egalitarian societies are
characterized by more fluid and less predictable (i.e., more complex) social relationships (cf.
[41,63]), individuals in these societies would experience higher social uncertainty than those
in despotic societies, subsequently leading to increased SDBs.

2. Materials and Methods

2.1. Study subjects

We observed 13 captive macaque groups across six species—*Macaca fuscata*, *M. mulatta*,
M. fascicularis, *M. silenus*, *M. sylvanus*, and *M. nigra*—in their existing social environments.
See **Table S1** for details on the subjects, and housing and husbandry protocols. Only
individuals ≥ 4 years – considered sexually mature [66,67] – at the time of testing (n = 113)

were included in the study and analyses (see **Table S1**). All groups had unrestricted access
to fresh drinking water. No alterations were made to feeding protocols for this research. The
animals were well habituated to the presence of the observers and exhibited no signs of
stress during data collection.

2.2. Behavioural data collection using continuous focal sampling

Behavioural observations were carried out using a continuous focal sampling method
between late 2020 and early 2023. We conducted these observations from 0800 to 1830
hours. Each focal observation lasted 20 minutes without periods when the individual was out
of sight. The order in which focal animals were observed was randomized. From the 113
individuals (≥ 4 years, range: 4 – 29 years; female = 92, male = 21), we obtained a total of
31,480 minutes of behavioural data (**Table S1**). The mean (± SD) observation minutes per
individual was 278.58 ± 75.23 (range = 160 – 460 min). The number of observation minutes
differed across groups, largely due to the varying group sizes. Focal data were used to
assess SDBs, aggressive behaviours, and dominance ranks.

2.3. Self-directed behaviours

We investigated three widely recognised SDBs – self-scratch, autogroom, and yawn
[5,6,13,16,23,41,68]. Self-scratch was defined as the act of an individual using their finger,
hand, or foot to rake across their own skin. Autogroom was defined by an individual
grooming themselves by closely inspecting the skin, hands, nails, toes, or other body parts.
Finally, we employed the widely accepted definition of a yawn (cf. [68]) – an involuntary and
powerful gaping of the jaw accompanied by deep inspiration, followed by a temporary period
of peak muscle contraction and passive closure of the jaw with shorter expiration. We could
not differentiate between self-directed and directed display yawns, as they are challenging to
disentangle in social contexts [69]. We calculated the rates of self-scratch and yawn
(frequency per minute) separately to control for the varying observation minutes across
groups and species. For autogroom, we quantified the duration and controlled for the
observation variation by calculating seconds per minute. A new autogroom bout was
considered if it was resumed after an interval of at least 5 seconds to account for data
independence.

2.4. Aggression initiated and received

207 We extracted data on five behavioural variables (*lunge, open-mouth threat, physical
208 attack/hit*, *bite,* and *chase*, see [70] for definitions) that represented both contact- and non-
209 contact aggression. For each individual, we calculated the rates (frequency per minute) of
210 aggression initiated (toward others) and aggression received (from others). By calculating
211 rates, we controlled for the varying observation minutes across groups and species.

2.5. Dominance rank and social style

Based on five variables that represent unprovoked submissive behaviours (*avoid*, *be
displaced*, *silent-bared teeth*, *flee*, and *social presence*, see [70] for definitions), we
assessed individual dominance (or social) ranks as well as the steepness of the within-group
dominance hierarchies using a Bayesian Elo-rating method. For all groups, except the *M.
fuscata*, we used previously published data to obtain these values [60]. Individual dominance

ranks were obtained as ordinal ranks at the group level. For *M. fuscata*, the limited number
of dyadic unprovoked submissive interactions could not be used to calculate the ordinal
ranks reliably. Therefore, we used previously published results on the matrilines to which
those individuals belong [71]. Since the sampled individuals were adult females only in the
M. fuscata group (cf. **Table S1**), matrilineal rank inheritance can be assumed [72], and these
alternative measures can be useful for determining the ordinal ranks of the individuals.
Nonetheless, we took caution while performing statistical analyses using this group in some
cases. Due to the relatively long-term data collection (spanning ~8 months per group),
hierarchies were assumed to be stable, and no rank reversals were observed during the
study.

The steepness of hierarchies followed the assumption of the co-variation framework (cf. [52–
54]) that more despotic societies show higher steepness than more egalitarian ones [60].
Previously published data on our study groups revealed a moderate to strong correlation
between macaque tolerance grades (or social styles) and hierarchy steepness (with all
individuals: r = -0.33; with only adult females: r = -0.61) [60]. The fulfillment of this
assumption, based on empirical data, validated our assignment of social styles to the
species we investigated.

2.6. Data coding

Video-recorded data were coded using BORIS (version 7.13.6) or in a frame-by-frame
manner using PotPlayer (version 240618). Due to data coding by multiple trained coders (by
D.B., E.S.J.vD., R.E.M.dV., C.E.K., M.V., and A.N.Z., with assistance from 14 more coders),
we comprehensively checked for inter-rater reliability. The agreement check was performed
after all the coders became fully trained to code behavioural data. Overall, the agreement
was high (Intraclass correlation coefficient or ICC(3,k) = 0.88, including SDBs, dominance
rank, and aggression). Particularly, for the SDBs, the ICC(3,k) value was 0.94, suggesting
excellent agreement among all coders.

2.7. Statistical models and packages

All statistical analyses were conducted in R (version 4.1.1) [73]. We used a Bayesian
modeling approach using the "*brms*" package [74]. SDBs (response variables) were
modelled using multivariate Gaussian models with identity links, which allowed them to be
estimated jointly with a residual correlation structure. Three models were constructed to test
the predictions of the study. In all models, we included a nested random effect of species
and group (i.e., group within species) to account for the hierarchical sampling structure.

Model 1 was used to investigate the association between the three SDBs, specifically
whether they exhibited strong correlational consistency. We used the default weakly
informative priors implemented in *brms*: flat priors for intercepts, half-student-t (3, 0, 10)
priors on the residual standard deviations, and an LKJ(1) prior for the residual correlation
matrix.

In *Model 2*, we tested the effects of age, sex, aggression initiated and received, dominance
rank, and tolerance grade or social style on the expression of SDBs. For dominance rank,
we included a penalized cubic regression spline (*k=3*) to capture potential non-linear
patterns. An interaction term of rank and grade was used. In the absence of a strong effect,

we investigated the individual effects of rank and grade. Although the SDBs were corrected
for varying sampling durations, we further checked the effect of observation time.
Additionally, we used group size in the model to investigate any potential effects. All three
response variables shared the same predictor structure and random-effects structure. We
used weakly informative priors for all model parameters. Fixed-effect coefficients were
assigned normal (0, 1) priors, and intercepts were given normal priors centred near the
empirical means of each response. Residual standard deviations and the standard
deviations of all random intercepts were assigned exponential priors with the rate parameter
value of 1. The residual correlation matrix among the three SDBs was given an LKJ(2) prior,
favoring weakly correlated structures while remaining only mildly informative.

Finally, in *Model 3*, we evaluated the effects of alpha status (alpha vs. non-alpha) on the
expression of SDBs, as alpha status may incur an elevated energetic cost compared to
intermediate or more subordinate group members [49]. Unlike in other models, here we
excluded the data on *M. fuscata* for two reasons: males were not included in the sample,
and dominance rank relationships were assessed through matrilineal ranks. Accordingly, we
had 12 alpha and 75 non-alpha individuals. Like previous models, we used weakly
informative priors for all model parameters. The priors used in this model fully resembled the
priors of Model 2.

For sufficient convergence, all Bayesian models were subjected to at least 5500 iterations
with 1000 warmup sessions. We reported median estimate coefficients (*Est*), 95% credible
interval (*crl*), containing 95% of the posterior probability density function, and probability of
direction (*pd*) that indicated certainty of an effect. We established model convergence by
following Bayesian statistics guidelines [75]. Furthermore, we checked trace and
autocorrelation plots, Gelman-Rubin convergence estimations, and density histograms of
posterior distributions. The posterior distributions were sampled using 10,000 iterations and
2000 warmups. We used the "*DHARMa.helpers*" package [76] and investigated model fits by
examining the residual distributions (i.e., deviation, dispersion, and outliers). Square-root
transformation of yawn was performed for improving model fits of 2 and 3. We applied the
Leave-one-out (LOO) cross validation method for model selection based on the expected log
pointwise predictive density (ELPD). Only best-fitted models are presented in the results.

3. Results

3.1. Self-directed behaviours in macaques show only partial correlational consistency

We found only partial correlational consistency among the three SDBs (**Figure 1, Table S2**).
Our model, which controlled for species- and group-level structure, revealed a moderate
positive residual association between self-scratch and autogroom (*Est* = 0.21, 95% *crl* [0.02
– 0.38], *pd* = 0.98). By contrast, the residual correlations between self-scratch and yawn (*Est*
= 0.08, 95% *crl* [-0.12 – 0.27], *pd* = 0.77) and between autogroom and yawn (*Est* = 0.02,
95% *crl* [-0.17 – 0.21], *pd* = 0.57) were weak. Taken together, these results suggest that
while some self-directed behaviours may share partially overlapping sources of variation,
their associations are generally weak. Thus, the three SDBs do not appear to be driven by a
single common underlying factor, supporting our decision to model them as separate
response variables rather than combining them into a single composite index.

Figure 1. Self-directed behaviours partially correlate with each other. Posterior estimates of residual correlations among self-directed behaviours. Solid black points show the posterior median correlation coefficient for each behavioural pair, while horizontal bars represent the 95% *crl*. The violins show data distribution. The vertical red dashed line indicates a parameter estimate of zero, i.e., the overlap of the *crl* with this line suggests no strong correlations. The yellow color highlights a moderately strong correlation.

3.2. Factors associated with self-directed behaviours in macaques

The LOO model comparison identified the multivariate model, which included sex, aggression received, dominance rank, tolerance grade, observation time, and group size, as the best-fitting model for predicting self-scratch, autogroom, and yawn. The dropped variables had no influence on the expression of SDBs. See **Table S3** for model comparison and selection using the LOO method. We found that the key predictors had varying associations with the three self-directed behaviours (**Table S3**).

3.2.1. Associations with sex and aggression

Sex had no effect on self-scratch (*Est* = -0.04, 95% *crl* [-0.08 − 0.01], *pd* = 0.95) and autogroom (*Est* = -0.77, 95% *crl* [-1.82 − 0.30], *pd* = 0.92), but it strongly predicted yawning,

with males showing more frequent yawning than females (*Est* = -0.08, 95% *crl* [-0.10 − -
0.05], *pd* = 1.00, **Figure 2**). Aggression received was not associated with self-scratch (*Est* =
0.00, 95% *crl* [-0.03 − 0.03], *pd* = 0.52) and yawn (*Est* = -0.01, 95% *crl* [-0.02 − 0.01], *pd* =
0.83), but was negatively related to autogroom, suggesting that individuals who received
lower aggression exhibited prolonged autogroom bouts (*Est* = -0.84, 95% *crl* [-1.61 − 0.09],
pd = 0.98).

Figure 2. Males exhibit more frequent yawn than females. Square-root transformed yawn rates
between male and female macaques. Boxes represent interquartile ranges, and whiskers represent
the upper and lower limits of the data. The horizontal bars within the boxes represent the median
values. Solid black dots show the distribution of data. Macaque photo credit: Monkey Reality project
from Animal Behaviour & Cognition group, Utrecht University.

3.2.2. Associations with dominance rank

We found evidence of both linear and non-linear relationships between dominance rank and
the SDBs (**Figure 3**). While our analysis on rank showed limited evidence for an effect on
self-scratch (*Est* = 0.03, 95% *crl* [-0.00 – 0.06], *pd* = 0.97, **Figure 3a**), it was strongly
associated with autogroom (*Est* = 1.14, 95% *crl* [0.35 – 1.94], *pd* = 0.99, **Figure 3b**) and
yawn (*Est* = 0.02, 95% *crl* [0.01 – 0.04], *pd* = 0.99, **Figure 3c**) behaviours. These results
suggest that low-ranking individuals engaged in longer autogrooming bouts than their high-
ranking counterparts. By contrast, the non-linear relationship between rank and yawning
behaviour suggests that yawn rates were lowest at intermediate ranks and increased toward
both high- and low-ranking individuals.

Figure 3. Relationships between dominance rank and self-directed behaviours in macaques.
(a) relationship between self-scratch and ordinal rank, (b) relationship between autogroom and ordinal
rank, (c) relationship between square-root transformed yawn and ordinal rank. Solid black points
show raw data points and overlaid in blue are the posterior predictions. Dashed blue line indicates no
strong relationship, whereas the solid lines suggest strong relationship between rank and self-directed
behaviours.

3.2.3. Effects of social styles

The mean (±SD) rates of self-scratch were 0.18 ± 0.08, 0.36 ± 0.25, 0.46 ± 0.21, and 0.41 ±
0.22 for Grades 1, 2, 3, and 4, respectively (**Figure 4a**), whereas the mean durations of
autogroom bouts were 3.14 ± 2.98, 8.61 ± 5.70, 5.12 ± 3.34, and 3.13 ± 2.52 (**Figure 4b**).
Likewise, the mean rates of yawn were 0.01 ± 0.02, 0.02 ± 0.04, 0.03 ± 0.04, and 0.03 ±
0.04 for the four tolerance grades (**Figure 4c**). Despite a visual pattern of the most despotic
societies exhibiting lower SDBs, tolerance grade showed no consistent associations, with all
estimates highly uncertain and 95% *crI* values spanning zero (**Table S3**). However, we found
substantial species- and group-level variation in autogrooming (species SD = 2.17; group SD
= 2.12), moderate variation in self-scratch, and minimal variation in yawning (**Table S3**). No
strong effects of observation time and group size were found (**Table S3**).

Figure 4. Self-directed behaviours across macaque social styles or tolerance grades. (a) rate of
self-scratch behaviour, **(b)** duration of autogroom, and **(c)** rate of yawn behaviour. Boxes represent
interquartile ranges, and whiskers represent the upper and lower limits of the data. The horizontal
bars within the boxes represent median values. Solid black dots show the distribution of data.

3.3. Yawn under pressure? Alpha status is linked to increased rates of yawn

The mean (±SD) rates of self-scratch were 0.326 ± 0.154 for alpha individuals and 0.362 ±
0.244 for non-alpha individuals. Similarly, the mean (±SD) durations of autogrooming bouts
were 5.105 ± 3.628 for alphas and 4.984 ± 4.533 for non-alphas, while yawning rates were
0.068 ± 0.054 and 0.022 ± 0.032 for alpha and non-alpha individuals, respectively. Alpha
individuals did not differ from non-alpha individuals in autogrooming (*Est* = 0.49, 95% *crl* [-
1.65 – 2.57], *pd* = 0.63, **Table S4**) or self-scratch (*Est* = -0.06, 95% *crl* [-0.15 – 0.02], *pd* =
0.92, **Table S4**). By contrast, yawning rates were higher in alpha individuals compared to
non-alpha (*Est* = 0.12, 95% *crl* [0.07 – 0.17], *pd* = 1.00, **Figure 5**, **Table S4**).

Figure 5. Rate of yawn between alpha and non-alpha individuals. Boxes represent interquartile
ranges, and whiskers represent the upper and lower limits of the data. The horizontal bars within the
boxes represent the median values. Solid black dots show the distribution of square-root transformed
data.

Discussion

SDBs are considered displacement behaviours that mediate anxiety-like emotional states
and serve functions, like assisting individuals to avoid potential conflicts or seek support from
closely affiliated group members through visual means [6,13,22,41]. Yet conclusive evidence
on the proximate drivers and the associated demographic and social factors underlying
variations in SDBs is lacking. Moreover, existing research predominantly includes specific
social contexts, like affiliative interactions and conflicts, and often uses social variables like
proximity to neighbors (to assess dominance and indirect social influence of group members
on others). However, complex emotional states like anxiety are shaped by the continuous
and multifaceted processes of social interactions, including broader social dynamics. We
conducted behavioural observations on macaques from six different species that varied
along a despotic–egalitarian gradient, and investigated three widely recognised SDBs: self-
scratch, autogroom, and yawn. Our results suggest that these behaviours share only a
limited correlational structure and are influenced differently by demographic and social
factors. While we found little to no evidence to support the idea that SDBs are influenced by
the broader social dynamics of a species (cf. [50,51]), dominance rank-related effects were
complex and more pronounced. These results indicate that SDBs do not form a simple
unified anxiety construct and instead may stem from a complex set of functions like social
uncertainty, signaling, and hygiene.

Partially overlapping social functions of self-scratch, autogroom, and yawn

The functions of SDBs can range from social and environmental to hygienic [6–10]. A study
comparing the explanatory power of these three hypotheses found stronger support for the
parasitic or hygienic model over the other two (cf. [8]). However, they also found support for
the prediction that social status (dominance rank) and the uncertainty (presence of nearby
group members) of social interactions predict SDBs. In the current study, which investigates
the social and emotional aspects of SDBs, we found that not all three behaviours covaried
positively, indicating their functional heterogeneity. This is further corroborated by the fact
that each of these behaviours was differentially influenced by some of the demographic and
social factors we looked at (see later). Consequently, our findings suggest the need for
reevaluation of several previous studies that used composite SDB measures as proxies for
anxiety, and we strongly advocate for treating the elemental measures separately. Besides,
future research should carefully evaluate the context-specific expression of these
behaviours.

Demographic factors are weakly associated with self-directed behaviours

We found no evidence of associations between age and the SDBs. In humans, greater
social contact has been found to alleviate anxiety associated with ageing [77]. Thus, the
comparable expression of SDBs across the age range of our study animals (4–29 years)
potentially indicates that social living conditions may facilitate better emotional regulation
[78]. Nonetheless, the lack of association between age and SDBs corroborated the findings
of some previous research [79,80]. Unlike age, sex was strongly associated with yawning,
but not with self-scratch and autogroom. Males displayed higher rates of yawning than
females. A sex-specific bias in yawning can be explained by sexual dimorphism (cf. [33,34]).
Since males are more dominant than females in macaque societies, males may exert
dominance and signal threats by yawning more, as they possess larger canines than

females. Therefore, instead of anxiety, frequent yawning in males may serve as a form of
social signaling [69].

Social factors underlying self-directed behaviours

Contrary to our prediction, we found that individuals who received more frequent aggression
from group members engaged in autogrooming for shorter durations. Additionally, there was
no association between aggression and self-scratch or yawn. These findings contradict the
results of previous research [8,37–39,81]. An inverse relationship between aggression
received and autogroom may indicate the presence of alternate coping strategies mediating
aggression-related anxiety, like social buffering [47,48]. Furthermore, the non-significant
relationships of self-scratch and yawn may indicate that direct social interactions in the form
of aggression may not be a prerequisite to elicit anxiety-related emotional states (but see
[37–39]), or that these behaviours primarily serve other functions as discussed before.

Social status or dominance rank exhibited complex effects on autogroom and yawn
behaviours, consistent with both uncertainty and energetic-constraint hypotheses. The effect
of rank on self-scratch was weak but showed an inverse U-shaped non-linear trend. The
relatively higher self-scratch rates among intermediate-ranking individuals could potentially
be due to the high probabilities of rank mobility, whereas the highest and lowest ends of the
spectrum remain somewhat reserved [82]. By contrast, the prolonged autogrooming bouts in
subordinate individuals clearly indicates an influence of social uncertainty (but see [81]).
Finally, a U-shaped relationship pattern emerged between rank and yawn, which could be
attributed to both rank-related uncertainty (i.e., for lower ranks) and the energetic costs (i.e.,
for higher ranks) of social ranks (cf. [38,41,44]). The elevated rates of yawning among
higher-ranking individuals indicate the heightened physiological demands of mating,
vigilance, and group maintenance. This was even further strongly supported by the finding of
increased yawning among alpha individuals in comparison to the non-alpha (cf. [49]). In
particular, frequent yawning by alpha individuals can convey social signals of dominance to
other group members [83,84], which can help maintain the stability of social groups.
Alternatively, although not mutually exclusive, yawns are known to function as a brain
cooling mechanism and may become more frequent in alpha individuals due to their high
loads of vigilance [85]. While elevated rates of yawning in higher-ranking individuals can
reflect on certain energy demands, it might be indicative of different emotional costs in lower-
ranking individuals. Thus, it is possible that the lower-ranking individuals are probably
'certain that they are uncertain' in predicting the outcomes of social interactions (also see
[86]), subsequently leading to elevated rates of yawning. These results may also suggest the
probable mechanisms of vigilance and social tension at play for subordinate individuals.
Nonetheless, the functions of yawning are multifaceted, and more interdisciplinary research
is required to assess its role regarding social dynamics. In summary, the rank-related effects
and their directions were not consistent for self-scratch, autogroom, and yawn, further
stressing the differential mechanisms governing these behaviours in macaques.

Our results revealed no strong social style level variations in self-scratch, autogroom, and
yawn. However, individuals belonging to the most despotic social style (i.e., Grade 1)
showed a tendency to exhibit lower levels of these behaviours than others (cf. Figure 4).
Nonetheless, these patterns are contradictory to the prediction that despotism inflicts
pressure on individuals in the form of power asymmetries and frequent aggressive

interactions, and that individuals are highly susceptible to negative emotions like anxiety
[52,54,59,87]. In line with our findings, a comparative study between despotic long-tailed and
egalitarian Tonkean macaques (*Macaca tonkeana*) showed that long-term cortisol
concentrations (used as an indicator of stress) were higher among individuals of Tonkean
than those of long-tailed macaques [61]. While harsh leadership (such as in despotic
societies) can induce stress among humans [88], macaques belonging to the despotic social
styles can rely on alternative mechanisms to alleviate and cope with anxiety, such as
forming and maintaining selectively strong bonds [89–91] as a social buffer [92] (*sensu*
social buffering [47]) and engaging in discriminate cooperative and prosocial interactions
[60,93–97]. By contrast, in egalitarian societies, such inter-individual bonds are not
necessarily predictable, strong, and stable [59–61,98], potentially dampening the positive
effects of tolerance [48]. Moreover, a recent study using the same groups (except the
Japanese macaques) as the current, experimentally showed that co-feeding tolerance levels
were higher in more despotic societies (cf. [60]). Therefore, the strong selective bonds in
despotic societies might have been translated into higher tolerance, which in turn resulted in
less frequent exhibition of SDBs.

The alternative prediction, i.e., higher social uncertainty in more egalitarian than despotic
societies, was partially supported, at least based on the same trend in the most despotic
society. Due to the predictable nature of social relationships in despotic societies, individuals
are certain about the possible outcomes of social interactions, subsequently exhibiting lower
levels of SDBs. By contrast, higher uncertainty in less despotic or more egalitarian societies
can lead to individuals showing greater levels of SDBs. Notably, social uncertainty is
considered a critical measure of social complexity [64,65]. Thus, in other words, we can
assume that social complexity may play a role in driving anxiety-like behaviours, which
should be tested in future studies.

Ethics

We received ethical approval from the Animal Experiments Committee and the Animal
Welfare Organisation at the Biomedical Primate Research Centre (BPRC) (approval no.
019E). At each participating zoo, internal ethical committees further approved and carefully
monitored the research procedures. Observations were conducted from a considerable
distance, without deliberately influencing the behaviour of the animals. The entire study was
strictly non-invasive in accordance with European Directive 2010/63, and we followed the
guidelines set forth by the American Society of Primatologists regarding the appropriate
treatment and inclusion of animals.

Data accessibility

511 Upon publication, data and code will be made publicly available.

Declaration of AI use

513 We have not used AI-assisted technologies in creating this article.

Authors' contributions

D.B.: conceptualization, methodology, investigation, formal analysis, writing – original draft,
writing – review and editing, visualization, supervision, project administration, funding
acquisition; E.S.J.vD.: investigation; R.E.M.dV.: investigation; C.E.K.: investigation; M.V.:
investigation; A.N.Z.: investigation; J.J.M.M.: conceptualization, methodology, writing –
original draft, writing – review and editing, supervision, project administration, funding
acquisition. All authors gave final approval for publication and agreed to be held accountable
for the work performed therein.

Conflict of interest declaration

523 We declare we have no competing interests.

Funding

525 The study received funding from the European Union's Horizon 2020 – Marie Skłodowska-
526 Curie Actions research and innovation program (H2020-MSCA-IF-2019-893016 to D.B.).

Acknowledgements

We thank William Allen, Elena Belli, Paula Escriche Chova, Jolanda de Jong, Aníta
Guðjónsdóttir, Elja Jeunink, Penny Kuijer, Esmee Middelburg, Veera Schroderus, Sjoerd
Sijbrandij, Paola Meems, Amber Kozanli, Yesper Bos, and Mary Maximiadi for their
assistance with data collection and organization. We thank the staff members of the
Affenberg Zoobetriebsgesellschaft mbH (Austria), Biomedical Primate Research Centre (the
Netherlands), Diergaarde Blijdorp (the Netherlands), Gaia Zoo (the Netherlands), Apenheul
Primate Park (the Netherlands), Planckendael Zoo (Belgium), and Artis Zoo (the
Netherlands) for their support during our research. We thank Peter Gaubatz, Svenja
Gaubatz, Jan Langermans, Annet Louise Louwerse, Jos Hartog, Linda Bruins-van
Sonsbeek, Emile Prins, Lisette van den Berg, Marjolein Osieck, Lena Pflüger, Karline

Janmaat, and Elisabeth Sterck for facilitating this research at respective zoos and
institutions. We also thank George Hodgson and Alan McElligott for providing valuable
comments on an earlier version of the manuscript.

References

1.   Kremer L, Klein Holkenborg SEJ, Reimert I, Bolhuis JE, Webb LE. 2020 The nuts and bolts of animal emotion. *Neurosci Biobehav Rev* **113**, 273–286. (doi:10.1016/j.neubiorev.2020.01.028)

2.   Nesse RM, Ellsworth PC. 2009 Evolution, emotions, and emotional disorders. *Am Psychol* **64**, 129–139. (doi:10.1037/a0013503)

3.   Bateson M, Brilot B, Nettle D. 2011 Anxiety: An Evolutionary Approach. *Can J Psychiatry* **56**, 707–715. (doi:10.1177/070674371105601202)

4.   Brosnan SF, Tone EB, Williams L. 2017 The evolution of social anxiety. In T. K. Shackelford & V. Zeigler-Hill (Eds.), *The evolution of psychopathology,* Springer Nature publishing. 93–116. (doi:10.1007/978-3-319-60576-0_4)

5.   Soler M. 2018 Self-directed behavior. In *Encyclopedia of Animal Cognition and Behavior*, pp. 1–5. Cham: Springer international publishing. (doi:10.1007/978-3-319-47829-6_1522-1)

6.   Maestripieri D, Schino G, Aureli F, Troisi A. 1992 A modest proposal: displacement activities as an indicator of emotions in primates. *Anim Behav* **44**, 967–979. (doi:10.1016/S0003-3472(05)80592-5)

7.   Hart BL. 1990 Behavioral adaptations to pathogens and parasites: Five strategies. *Neurosci Biobehav Rev* **14**, 273–294. (doi:10.1016/S0149-7634(05)80038-7)

8.   Duboscq J, Romano V, Sueur C, MacIntosh AJJ. 2016 Scratch that itch: revisiting links between self-directed behaviour and parasitological, social and environmental factors in a free-ranging primate. *R Soc Open Sci* **3**, 160571. (doi:10.1098/rsos.160571)

9.   Mooring MS, Blumstein DT, Stoner CJ. 2004 The evolution of parasite-defence grooming in ungulates. *Biol J Linn Soc* **81**, 17–37. (doi:10.1111/j.1095-8312.2004.00273.x)

10.   Troisi A, Schino G. 1987 Environmental and Social Influences on Autogrooming Behaviour in a Captive Group of Java Monkeys. *Behaviour* **100**, 292–302.

11.   Wascher CAF, Fraser ON, Kotrschal K. 2010 Heart Rate during Conflicts Predicts Post-Conflict Stress-Related Behavior in Greylag Geese. *PLoS One* **5**, e15751. (doi:10.1371/journal.pone.0015751)

12.   Nephew BC, Romero LM. 2003 Behavioral, physiological, and endocrine responses of starlings to acute increases in density. *Horm Behav* **44**, 222–232. (doi:10.1016/j.yhbeh.2003.06.002)

13.   Schino G, Perretta G, Taglioni AM, Monaco V, Troisi A. 1996 Primate displacement activities as an ethopharmacological model of anxiety. *Anxiety* **2**, 186–191. (doi:10.1002/(SICI)1522-7154(1996)2:4<186::AID-ANXI5>3.0.CO;2-M)

14.   Schino G, Troisi A, Perretta G, Monaco V. 1991 Measuring anxiety in nonhuman primates: Effect of lorazepam on macaque scratching. *Pharmacol Biochem Behav* **38**, 889–891. (doi:10.1016/0091-3057(91)90258-4)

15.   Ulyan MJ, Burrows AE, Buzzell CA, Raghanti MA, Marcinkiewicz JL, Phillips KA. 2006
The effects of predictable and unpredictable feeding schedules on the behavior and
physiology of captive brown capuchins (Cebus apella). *Appl Anim Behav Sci* **101**,
154–160. (doi:10.1016/j.applanim.2006.01.010)

16.   Ellis JJ, MacLarnon AM, Heistermann M, Semple S. 2011 The social correlates of self-
directed behaviour and faecal glucocorticoid levels among adult male olive baboons (
Papio hamadryas anubis) in Gashaka-Gumti National Park, Nigeria. *Afr Zool* **46**,
302–308. (doi:10.1080/15627020.2011.11407503)

17.   Higham JP, MacLarnon AM, Heistermann M, Ross C, Semple S. 2009 Rates of self-
directed behaviour and faecal glucocorticoid levels are not correlated in female wild
olive baboons (*Papio hamadryas anubis*). *Stress* **12**, 526–532.
(doi:10.3109/10253890902756565)

18.   Fraser ON, Stahl D, Aureli F. 2008 Stress reduction through consolation in
chimpanzees. *Proc Natl Acad Sci USA* **105**, 8557–8562.
(doi:10.1073/pnas.0804141105)

19.   Manson JH, Perry S. 2000 Correlates of Self-directed behaviour in wild white-faced
capuchins. *Ethology* **106**, 301–317. (doi:10.1046/j.1439-0310.2000.00527.x)

20.   Castles DL, Whiten A, Aureli F. 1999 Social anxiety, relationships and self-directed
behaviour among wild female olive baboons. *Anim Behav* **58**, 1207–1215.
(doi:10.1006/anbe.1999.1250)

21.   Wright CI, Majolo B, Schino G, Ventura R, Romero T. 2024 Social tension after
grooming in wild Japanese macaques *(Macaca fuscata yakui)* is sex specific and
sensitive to social relationships. *Am J Primatol* **86**. (doi:10.1002/ajp.23664)

22.   Troisi A. 2002 Displacement activities as a behavioral measure of stress in nonhuman
primates and human subjects. *Stress* **5**, 47–54. (doi:10.1080/102538902900012378)

23.   Laméris DW, Verspeek J, Salas M, Staes N, Torfs JRR, Eens M, Stevens JMG. 2022
Evaluating self-directed behaviours and their association with emotional arousal
across two cognitive tasks in bonobos (Pan paniscus). *Animals* **12**, 3002.
(doi:10.3390/ani12213002)

24.   Leeds A, Lukas K. 2018 Experimentally evaluating the function of self-directed
behaviour in two adult mandrills (Mandrillus sphinx). *Anim Welf* **27**, 81–86.
(doi:10.7120/09627286.27.1.081)

25.   van Dijk ESJ, Bhattacharjee D, Belli E, Massen JJM. 2023 Hand preference predicts
behavioral responses to threats in Barbary macaques. *Am J Primatol* **85**.
(doi:10.1002/ajp.23499)

26.   Gustison ML, MacLarnon A, Wiper S, Semple S. 2012 An experimental study of
behavioural coping strategies in free-ranging female Barbary macaques (*Macaca*
sylvanus). *Stress* **15**, 608–617. (doi:10.3109/10253890.2012.668589)

27.   Briefer EF, Oxley JA, McElligott AG. 2015 Autonomic nervous system reactivity in a
free-ranging mammal: effects of dominance rank and personality. *Anim Behav* **110**,
121–132. (doi:10.1016/j.anbehav.2015.09.022)

28.   Steptoe A, Deaton A, Stone AA. 2015 Subjective wellbeing, health, and ageing. *The*
Lancet **385**, 640–648. (doi:10.1016/S0140-6736(13)61489-0)

29. Reamer L, Tooze Z, Coulson C, Semple S. 2010 Correlates of self-directed and
stereotypic behaviours in captive red-capped mangabeys (Cercocebus torquatus
torquatus). *Appl Anim Behav Sci* **124**, 68–74. (doi:10.1016/j.applanim.2010.01.012)

30. Charles ST, Carstensen LL. 2010 Social and Emotional Aging. *Annu Rev Psychol* **61**,
383–409. (doi:10.1146/annurev.psych.093008.100448)

31. Troisi A, Aureli F, Schino G, Rinaldi F, de Angelis N. 1990 The Influence of Age, Sex,
and Rank on Yawning Behavior in Two Species of Macaques (*Macaca fascicularis*
and *M. fuscata*). *Ethology* **86**, 303–310. (doi:10.1111/j.1439-0310.1990.tb00437.x)

32. Bernstein IS, Ehardt CL. 1985 Age-sex differences in the expression of agonistic
behavior in rhesus monkey (Macaca mulatta) groups. *J Comp Psychol* **99**, 115–132.
(doi:10.1037/0735-7036.99.2.115)

33. Thorén S, Lindenfors P, Kappeler PM. 2006 Phylogenetic analyses of dimorphism in
primates: Evidence for stronger selection on canine size than on body size. *Am J*
Phys Anthropol **130**, 50–59. (doi:10.1002/ajpa.20321)

34. Zannella A, Stanyon R, Palagi E. 2017 Yawning and social styles: Different functions
in tolerant and despotic macaques (Macaca tonkeana and Macaca fuscata). *J Comp*
Psychol **131**, 179–188. (doi:10.1037/com0000062)

35. Sclafani V, Norscia I, Antonacci D, Palagi E. 2012 Scratching around mating: factors
affecting anxiety in wild Lemur catta. *Primates* **53**, 247–254. (doi:10.1007/s10329-
012-0294-6)

36. Peignot P, Jankowski F, Anderson JR. 2004 On self-scratching in captive mandrills
(Mandrillus sphinx). *Folia Primatol* **75**, 160–164. (doi:10.1159/000078307)

37. Cooper MA, Aureli F, Singh M. 2007 Sex Differences in reconciliation and post-conflict
anxiety in bonnet macaques. *Ethology* **113**, 26–38. (doi:10.1111/j.1439-
0310.2006.01287.x)

38. Kaburu SSK, MacLarnon A, Majolo B, Qarro M, Semple S. 2012 Dominance rank and
self-scratching among wild female Barbary macaques (*Macaca sylvanus*). *Afr Zool*
47, 74–79. (doi:10.1080/15627020.2012.11407525)

39. Schino G, Rosati L, Geminiani S, Aureli F. 2007 Post-conflict anxiety in Japanese
macaques (*Macaca fuscata*): aggressor's and victim's perspectives. *Ethology* **113**,
1081–1088. (doi:10.1111/j.1439-0310.2007.01407.x)

40. Pomerantz O, Terkel J. 2009 Effects of positive reinforcement training techniques on
the psychological welfare of zoo-housed chimpanzees (*Pan troglodytes*). *Am J*
Primatol **71**, 687–695. (doi:10.1002/ajp.20703)

41. Castles DL, Whiten A, Aureli F. 1999 Social anxiety, relationships and self-directed
behaviour among wild female olive baboons. *Anim Behav* **58**, 1207–1215.
(doi:10.1006/anbe.1999.1250)

42. Radford AN. 2012 Post-allogrooming reductions in self-directed behaviour are
affected by role and status in the green woodhoopoe. *Biol Lett* **8**, 24–27.
(doi:10.1098/rsbl.2011.0559)

43. Molesti S, Majolo B. 2013 Grooming increases self-directed behaviour in wild Barbary
macaques, Macaca sylvanus. *Anim Behav* **86**, 169–175.
(doi:10.1016/j.anbehav.2013.05.008)

44. Abbott DH *et al.* 2003 Are subordinates always stressed? a comparative analysis of
rank differences in cortisol levels among primates. *Horm Behav* **43**, 67–82.
(doi:10.1016/S0018-506X(02)00037-5)

45. Zhang Q, Wang X, Zhu Y, Garber PA, Xia D, Li J. 2025 Self-directed behaviors prior to
approaching conspecifics in wild Tibetan macaques (*Macaca thibetana*). *Am J
Primatol* **87**. (doi:10.1002/ajp.70051)

46. Bhattacharjee D, Guðjónsdóttir AR, Chova PE, Middelburg E, Jäckels J, de Groot NG,
Wallner B, Massen JJM, Pflüger LS. 2024 Behavioral, physiological, and genetic
drivers of coping in a non-human primate. *iScience* **27**, 108890.
(doi:10.1016/j.isci.2024.108890)

47. Cohen S, Wills TA. 1985 Stress, social support, and the buffering hypothesis. *Psychol
Bull* **98**, 310–357. (doi:10.1037/0033-2909.98.2.310)

48. Hennessy MB, Kaiser S, Sachser N. 2009 Social buffering of the stress response:
Diversity, mechanisms, and functions. *Front Neuroendocrinol* **30**, 470–482.
(doi:10.1016/j.yfrne.2009.06.001)

49. Gesquiere LR, Learn NH, Simao MCM, Onyango PO, Alberts SC, Altmann J. 2011
Life at the top: rank and stress in wild male baboons. *Science (1979)* **333**, 357–360.
(doi:10.1126/science.1207120)

50. Sapolsky RM. 2005 The influence of social hierarchy on primate health. *Science
(1979)* **308**, 648–652. (doi:10.1126/science.1106477)

51. Creel S. 2001 Social dominance and stress hormones. *Trends Ecol Evol* **16**, 491–497.
(doi:10.1016/S0169-5347(01)02227-3)

52. Thierry B. 2007 Unity in diversity: Lessons from macaque societies. *Evol Anthropol*
16, 224–238. (doi:10.1002/evan.20147)

53. Thierry B. 2000 Covariation of conflict management patterns across macaque
species. *Natural Conflict Resolution* (doi:10.1006/anbe.2001.1656)

54. Thierry B. 2022 Where do we stand with the covariation framework in primate
societies? *Am J Biol Anthropol* **178**, 5–25. (doi:10.1002/ajpa.24441)

55. Shultz S, Opie C, Atkinson QD. 2011 Stepwise evolution of stable sociality in
primates. *Nature* **479**, 219–222. (doi:10.1038/nature10601)

56. Balasubramaniam KN *et al.* 2012 Hierarchical steepness, counter-aggression, and
macaque social style scale. *Am J Primatol* **74**, 915–925. (doi:10.1002/ajp.22044)

57. Balasubramaniam KN *et al.* 2018 The influence of phylogeny, social style, and
sociodemographic factors on macaque social network structure. *Am J Primatol* **80**.
(doi:10.1002/ajp.22727)

58. Sueur C, Petit O, De Marco A, Jacobs AT, Watanabe K, Thierry B. 2011 A comparative
network analysis of social style in macaques. *Anim Behav* **82**, 845–852.
(doi:10.1016/j.anbehav.2011.07.020)

59. Duboscq J, Micheletta J, Agil M, Hodges K, Thierry B, Engelhardt A. 2013 Social
Tolerance in Wild Female Crested Macaques (*Macaca nigra)* in Tangkoko-Batuangus
Nature Reserve, Sulawesi, Indonesia. *Am J Primatol* **75**, 361–375.
(doi:10.1002/ajp.22114)

60. Bhattacharjee D *et al.* 2025 Despotism promotes cooperation through enhanced
interdependencies in non-human primate societies. *biorxiv*
(doi:10.1101/2025.03.27.645840)

61. Sadoughi B, Lacroix L, Berbesque C, Meunier H, Lehmann J. 2021 Effects of social
tolerance on stress: hair cortisol concentrations in the tolerant Tonkean macaques (
Macaca tonkeana) and the despotic long-tailed macaques (*Macaca fascicularis*).
Stress **24**, 1033–1041. (doi:10.1080/10253890.2021.1998443)

62. Ciani F, Dall'Olio S, Stanyon R, Palagi E. 2012 Social tolerance and adult play in
macaque societies: a comparison with different human cultures. *Anim Behav* **84**,
1313–1322. (doi:10.1016/j.anbehav.2012.09.002)

63. Loyant L, Waller BM, Micheletta J, Meunier H, Ballesta S, Joly M. 2023 Tolerant
macaque species are less impulsive and reactive. *Anim Cogn* (doi:10.1007/s10071-
023-01789-8)

64. Rincon A V, Waller BM, Duboscq J, Mielke A, Pérez C, Clark PR, Micheletta J. 2023
Higher social tolerance is associated with more complex facial behavior in macaques.
elife **12**. (doi:10.7554/eLife.87008.3)

65. Rebout N *et al.* 2022 Tolerant and intolerant macaques differ in the context specificity
of their calls and how they 'comment' on the interactions of others. *Behav Ecol
Sociobiol* **76**, 67. (doi:10.1007/s00265-022-03177-7)

66. Bercovitch FB, Berard JD. 1993 Life history costs and consequences of rapid
reproductive maturation in female rhesus macaques. *Behav Ecol Sociobiol* **32**, 103–
109. (doi:10.1007/BF00164042)

67. Kinnally EL, Lyons LA, Abel K, Mendoza S, Capitanio JP. 2008 Effects of early
experience and genotype on serotonin transporter regulation in infant rhesus
macaques. *Genes Brain Behav* **7**, 481–486. (doi:10.1111/j.1601-183X.2007.00383.x)

68. Barbizet J. 1958 Yawning. *J Neurol Neurosurg Psychiatry* **21**, 203–209.
(doi:10.1136/jnnp.21.3.203)

69. Moyaho A, Flores Urbina A, Monjaraz Guzmán E, Walusinski O. 2017 Yawning: a cue
and a signal. *Heliyon* **3**, e00437. (doi:10.1016/j.heliyon.2017.e00437)

70. Kluiver CE, de Jong JA, Massen JJM, Bhattacharjee D. 2022 Personality as a
Predictor of Time-Activity Budget in Lion-Tailed Macaques (Macaca silenus). *Animals*
12, 1495. (doi:10.3390/ani12121495)

71. Pflüger LS, Pink KE, Wallner B, Radler C, Dorner M, Huffman MA. 2021 Twenty-three-
year demographic history of the Affenberg Japanese macaques (Macaca fuscata), a
translocated semi-free-ranging group in southern Austria. *Primates* **62**, 761–776.
(doi:10.1007/s10329-021-00928-4)

72. Chapais B. 1988 Experimental matrilineal inheritance of rank in female Japanese
macaques. *Anim Behav* **36**, 1025–1037. (doi:10.1016/S0003-3472(88)80062-9)

747	73.	R Development Core Team. 2019 R Core Team (2020). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/. *R Foundation for Statistical Computing.* **2**.
748		
749		

73. R Development Core Team. 2019 R Core Team (2020). R: A language and
environment for statistical computing. R Foundation for Statistical Computing, Vienna,
Austria. URL https://www.R-project.org/. *R Foundation for Statistical Computing.* **2**.

74. Bürkner P-C. 2017 brms: An R Package for Bayesian Multilevel Models Using Stan. *J*
Stat Softw **80**. (doi:10.18637/jss.v080.i01)

75. Depaoli S, van de Schoot R. 2017 Improving transparency and replication in Bayesian
statistics: The WAMBS-Checklist. *Psychol Methods* **22**, 240–261.
(doi:10.1037/met0000065)

76. Rodríguez-Sánchez F. 2024 DHARMa.helpers: Helper functions to check models not
(yet) directly supported by DHARMa.

77. Brunton RJ, Scott G. 2015 Do we fear ageing? A multidimensional approach to ageing
anxiety. *Educ Gerontol* **41**, 786–799. (doi:10.1080/03601277.2015.1050870)

78. Charles ST, Carstensen LL. 2010 Social and emotional aging. *Annu Rev Psychol* **61**,
383–409. (doi:10.1146/annurev.psych.093008.100448)

79. Polizzi di Sorrentino E, Schino G, Tiddi B, Aureli F. 2012 Scratching as a window into
the emotional responses of wild tufted capuchin monkeys. *Ethology* **118**, 1072–1084.
(doi:10.1111/eth.12008)

80. Dell'Anna F, Schino G, Aureli F. 2022 Anxiety in Geoffroy's spider monkeys (*Ateles*
geoffroyi): can scratching be used as an indicator? *Am J Primatol* **84**.
(doi:10.1002/ajp.23373)

81. Daniel JR, dos Santos AJ, Vicente L. 2008 Correlates of Self-directed Behaviors in
Captive Cercopithecus aethiops. *Int J Primatol* **29**, 1219–1226. (doi:10.1007/s10764-
008-9300-7)

82. Blersch R, Beisner BA, Vandeleest JJ, McCowan B. 2024 The dynamics of
dominance in a 'despotic' society. *Anim Behav* **218**, 55–64.
(doi:10.1016/j.anbehav.2024.09.009)

83. Zannella A, Stanyon R, Maglieri V, Palagi E. 2021 Not all yawns tell the same story:
The case of Tonkean macaques. *Am J Primatol* **83**. (doi:10.1002/ajp.23263)

84. Leone A, Ferrari PF, Palagi E. 2014 Different yawns, different functions? Testing social
hypotheses on spontaneous yawning in Theropithecus gelada. *Sci Rep* **4**, 4010.
(doi:10.1038/srep04010)

85. Massen JJM, Dusch K, Eldakar OT, Gallup AC. 2014 A thermal window for yawning in
humans: Yawning as a brain cooling mechanism. *Physiol Behav* **130**, 145–148.
(doi:10.1016/j.physbeh.2014.03.032)

86. Sinha A. 1998 Knowledge acquired and decisions made: triadic interactions during
allogrooming in wild bonnet macaques, Macaca radiata. *Philos Trans R Soc Lond B*
Biol Sci **353**, 619–631. (doi:10.1098/rstb.1998.0230)

87. Adams MJ *et al.* 2015 Personality structure and social style in macaques. *J Pers Soc*
Psychol **109**, 338–353. (doi:10.1037/pspp0000041)

88. Diebig M, Bormann KC, Rowold J. 2016 A double-edged sword: Relationship between
full-range leadership behaviors and followers' hair cortisol level. *Leadersh Q* **27**, 684–
696. (doi:10.1016/j.leaqua.2016.04.001)

89.   Schino G. 2007 Grooming and agonistic support: a meta-analysis of primate
reciprocal altruism. *Behav Ecol* **18**, 115–120. (doi:10.1093/beheco/arl045)

90.   Ventura R, Majolo B, Koyama NF, Hardie S, Schino G. 2006 Reciprocation and
interchange in wild Japanese macaques: grooming, cofeeding, and agonistic support.
Am J Primatol **68**, 1138–1149. (doi:10.1002/ajp.20314)

91.   Aere S, Pedruzzi L, Facondini G, Böye M, Palagi E, Lemasson A. 2025 The curious
case of rhesus macaques: despotism does not prevent third-party postconflict
affiliation. *Anim Behav* **219**, 123022. (doi:10.1016/j.anbehav.2024.10.036)

92.   Young C, Majolo B, Heistermann M, Schülke O, Ostner J. 2014 Responses to social
and environmental stress are attenuated by strong male bonds in wild macaques.
Proc Natl Acad Sci USA **111**, 18195–18200. (doi:10.1073/pnas.1411450111)

93.   Bhattacharjee D, Cousin E, Pflüger LS, Massen JJM. 2023 Prosociality in a despotic
society. *iScience* **26**, 106587. (doi:10.1016/j.isci.2023.106587)

94.   Bhattacharjee D, Waasdorp S, Middelburg E, Sterck EHM, Massen JJM. 2024
Personality heterophily and friendship as drivers for successful cooperation. *Proc R*
Soc B **291**. (doi:10.1098/rspb.2023.2730)

95.   Sterck EHM, Olesen CU, Massen JJM. 2015 No costly prosociality among related
long-tailed macaques (Macaca fascicularis). *J Comp Psychol* **129**, 275–282.
(doi:10.1037/a0039180)

96.   Sigmundson R, Stribos MS, Hammer R, Herzele J, Pflüger LS, Massen JJM. 2021
Exploring the cognitive capacities of Japanese macaques in a cooperation game.
Animals **11**, 1497. (doi:10.3390/ani11061497)

97.   de Laat EJAM, Waasdorp S, Roth TS, Massen JJM, Sterck EHM. 2025 Despotic long-
tailed macaques benefit others in a group service paradigm. *Behaviour* , 1–29.
(doi:10.1163/1568539X-bja10338)

98.   Duboscq J, Neumann C, Agil M, Perwitasari-Farajallah D, Thierry B, Engelhardt A.
2017 Degrees of freedom in social bonds of crested macaque females. *Anim Behav*
123, 411–426. (doi:10.1016/j.anbehav.2016.11.010)

827 **Electronic supplementary information to**

828 # Many facets of self-directed behaviours in macaques

830 **Table S1. Summary of study species and groups with sampling and demographic**
831 **information.**

Species	Group ID	Focal sampled	Age (years)	Sex
Macaca fuscata (Japanese)	Affenberg	N = 26; 280 min/focal	4 – 29 (median age = 10)	Female = 26; Male = 0
Macaca mulatta (Rhesus)	R3G2	N = 12; 160 min/focal	4 – 14 (median age = 5)	Female = 11; Male = 1
	R3G7	N = 10; 210 min/focal	4 – 25 (median age = 13)	Female = 9; Male = 1
Macaca fascicularis (Long-tailed)	J1G4	N = 8; 320 min/focal	4 – 9 (median age = 4)	Female = 7; Male = 1
	J1G7	N = 10; 320 min/focal	4 – 14 (median age = 12.5)	Female = 9; Male = 1
	NWR	N = 4; 320 min/focal	4 – 6 (median age = 4)	Female = 0; Male = 4
Macaca silenus (Lion-tailed)	Blijdorp	N = 3; 240 min/focal	6 – 18 (median age = 7)	Female = 2; Male = 1
	Apenheul	N = 6; 340 min/focal	7 – 27 (median age = 18.5)	Female = 5; Male = 1
Macaca sylvanus (Barbary)	Gaia	N = 14; 240 min/focal	4 – 20 (median age = 10.5)	Female = 8; Male = 6
	Apenheul	N = 9; 420 min/focal	4 – 20 (median age = 7)	Female = 8; Male = 1
Macaca nigra (Crested)	Blijdorp	N = 3; 460 min/focal	5 – 21 (median age = 13)	Female = 2; Male = 1
	Artis	N = 3; 320 min/focal	9 – 21 (median age = 15)	Female = 3; Male = 0
	Planckendael	N = 5; 180 min/focal	4 – 15 (median age = 6)	Female = 2; Male = 3

The *M. fuscata* group was located at Affenberg Landskron Park in Villach, Austria, within a
large 40,000 m² enclosure characterized by native mixed woodlands typical of Southern
Austria. This group comprised approximately 170 individuals. Two *M. mulatta* groups (R3G2
and R3G7), with 30 and 25 members respectively, were housed at the BPRC in Rijswijk, the
Netherlands. Both groups lived in facilities featuring identical enclosure sizes: 74 m² indoors
and 250 m² outdoors. At the same facility (BPRC), we also studied three *M. fascicularis*
groups: J1G4 (15 individuals), J1G7 (18 individuals), and NWR (4 individuals). J1G4 and
J1G7 had access to the same enclosure sizes (49 m² indoor and 183 m² outdoor), while the
smaller NWR group was housed in smaller enclosures measuring 3.55 m² indoors and 3.88
m² outdoors. The two groups of *M. Silenus* were housed at Blijdorp Zoo and Apenheul

Primate Park in the Netherlands. The Blijdorp group comprised 3 individuals and had 100 m²
indoor and 106 m² outdoor spaces, whereas the Apenheul group included 8 individuals, with
access to an 80 m² indoor enclosure and a significantly larger 768 m² outdoor area. Two *M.*
sylvanus groups were included: one at Gaia Zoo in the Netherlands and the other at
Apenheul Primate Park. The Gaia Zoo group consisted of 14 individuals and had access to a
large 3522 m² outdoor area and a 108 m² indoor space. The Apenheul group, with 13
individuals, lived exclusively in a 3829 m² outdoor enclosure featuring natural elements such
as vegetation, rocks, and a creek. Finally, the three *M. nigra* groups were housed at Blijdorp
Zoo, Artis Zoo (both in the Netherlands), and Planckendael Zoo in Belgium. Group sizes
were 5, 4, and 6 individuals, respectively. The Blijdorp group was kept in a 70 m² indoor and
a 160 m² outdoor enclosure. At Artis Zoo, the enclosures measured 65 m² indoors and 761
m² outdoors, while the Planckendael group had access only to an 82 m² indoor enclosure.

In all cases where both indoor and outdoor spaces were provided, macaques could freely
move between them via connecting tunnels. An exception was the Apenheul *M. sylvanus*
group, where only the outdoor space was available during observations. Group sizes include
all members, including infants under one year of age.

Although husbandry practices varied slightly between groups—reflecting differences in
species-specific needs and institutional protocols—all followed the standards set by the
European Association of Zoos and Aquaria for animal housing and care. Enclosures were
equipped with various enrichment items, including climbing structures, ropes, tree trunks,
wooden elements, and slides. With the exception of the *M. fuscata* and Apenheul *M.*
sylvanus groups, which were kept exclusively in outdoor habitats, indoor housing areas were
climate-controlled and featured concrete floors layered with sawdust bedding. Feeding
schedules differed across groups to meet specific dietary requirements but remained
consistent with their usual routines during our observations. Standard diets included
commercial monkey pellets, a variety of fruits and vegetables, and seed mixes such as
sunflower and corn.

Table S2. Summary of Bayesian multivariate hierarchical model (*Model 1*)
investigating the correlational consistency among self-scratch, autogroom, and yawn
behaviours.

Fixed (population-level) effects

Response	Parameter	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Self-scratch	Intercept	0.35	0.10	0.15	0.54
Autogroom	Intercept	4.42	1.09	2.20	6.55
Yawn	Intercept	0.03	0.01	0.00	0.06

877 **Residual correlation**

Paired behaviours	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Self-scratch – Autogroom	0.21	0.10	0.02	0.38
Self-scratch – Yawn	0.08	0.10	–0.12	0.27
Autogroom – Yawn	0.02	0.10	–0.17	0.21

879 **Table S3. Summary of Bayesian multivariate hierarchical model (*Model 2*)**
880 **investigating the effects of sex, aggression received, dominance rank, and tolerance**
881 **grade or social style on self-scratch, autogroom, and yawn behaviours.**

883 **Self-scratch**

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	0.31	0.36	–0.66	0.92
Grade 1	0.04	0.80	–1.12	2.25
Grade 2	0.01	0.75	–1.71	1.50
Grade 3	–0.04	0.62	–1.56	1.03
Aggression received	0.00	0.02	–0.03	0.03
Sex (male)	–0.04	0.02	–0.08	0.01
Mean observation	0.02	0.08	–0.15	0.18
Group size	–0.24	0.61	–2.05	0.54

885 **Autogroom**

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	3.46	1.31	0.64	5.73
Grade 1	–4.08	3.08	–10.72	1.75
Grade 2	3.17	2.74	–2.76	8.65
Grade 3	0.70	2.32	–3.82	5.65
Aggression received	–0.84	0.39	–1.61	–0.09
Sex (male)	–0.77	0.53	–1.82	0.30
Mean observation	–0.29	0.83	–1.89	1.42
Group size	0.82	2.22	–3.44	5.67

887 **Yawn**

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	0.13	0.23	–0.47	0.53
Grade 1	0.08	0.44	–0.63	1.19
Grade 2	–0.03	0.46	–1.07	0.89
Grade 3	–0.01	0.37	–0.83	0.67
Aggression received	–0.01	0.01	–0.02	0.01
Sex (male)	–0.08	0.01	–0.10	–0.05
Mean observation	–0.02	0.03	–0.07	0.03
Group size	–0.13	0.25	–0.83	0.19

888
889

Smooth terms (dominance rank)

Response	Smooth term	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Self-scratch	s(ord_rank)	0.03	0.02	–0.00	0.06
Autogroom	s(ord_rank)	1.14	0.40	0.35	1.94
Yawn (sqrt)	s(ord_rank)	0.02	0.01	0.01	0.04

Note: Best-fitted model presented (Model3). Model selection was done using LOO method.
Result of LOO comparison of models –

```
         elpd_diff    se_diff
Model5   0.0          0.0
Model4   -0.3         0.5
Model3   -0.6         1.1
Model2   -3.0         2.2
Model1   -3.9         2.8
```

Variables included in Models:

Model1: s(ord_rank, k = 3) + grade + agginitiate + aggreceive + age + sex + meanobs + groupsize + (1 | species / group)

Model2: s(ord_rank, k = 3) + grade + agginitiate + aggreceive + sex + meanobs + groupsize + (1 | species / group)

Model3: s(ord_rank, k = 3) + grade + aggreceive + sex + meanobs + groupsize + (1 | species / group)

Model4: s(ord_rank, k = 3) + grade + aggreceive + sex + groupsize + (1 | species / group)

Model5: s(ord_rank, k = 3) + grade + aggreceive + sex + (1 | species / group)

[ord_rank = dominance rank; agginitiate = aggression initiate; aggreceive = aggression receive; meanobs = observation minutes]

Table S4. Summary of Bayesian multivariate hierarchical model (*Model 3*) investigating the alpha vs non-alpha individuals on self-scratch, autogroom, and yawn behaviours.

Self-scratch

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	0.38	0.13	0.13	0.62
Alpha vs non-alpha	−0.06	0.05	−0.15	0.02

Autogroom

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	4.40	1.26	1.81	6.89
Alpha vs non-alpha	0.49	1.07	−1.65	2.57

Yawn

Predictor	Estimate	SE	95% CrI (Lower)	95% CrI (Upper)
Intercept	0.12	0.04	0.05	0.19
Alpha vs non-alpha	0.12	0.03	0.07	0.17