Scaling and foraging behavior drive the evolution of humeral shape in hummingbirds

Juan Camilo Ríos-Orjuela[1,2,*], Carlos Daniel Cadena[1], Alejandro Rico-Guevara[3, 4], Ilias

Berberi[5], Lauren Miner[5] & Roslyn Dakin[5]

[1]Laboratorio de Biología Evolutiva de Vertebrados, Departamento de Ciencias Biológicas,

Universidad de los Andes, Bogotá, Colombia

[2]Grupo de Morfología y Ecología Evolutiva, Instituto de Ciencias Naturales, Universidad

Nacional de Colombia, Sede Bogotá, Colombia

[3]Department of Biology, University of Washington, Seattle, WA, U.S.A.

[4]Burke Museum of Natural History and Culture, University of Washington, Seattle, WA,

U.S.A.

[5]Department of Biology, Carleton University, Ottawa, Ontario, Canada

* Corresponding author (jcriosor@gmail.com)

Orcid: JCR: 0000-0001-6976-9131, DC: 0000-0003-4530-2478, ARG: 0000-0003-4067-

5312, IB: 0000-0003-3051-8145, LM: 0009-0008-1991-7472, RD: 0000-0002-3140-3975.

Author contributions: JCR: Conceptualization, Methodology, Software, Formal analysis,

Investigation, Data curation, Writing – Original draft, Writing – Review & Editing,

Visualization. **RD:** Conceptualization, Validation, Investigation, Resources, Writing –

Original draft, Writing – Review & Editing, Supervision, Funding acquisition. **CDC:**

Conceptualization, Writing – Review & Editing, Supervision, Funding acquisition. **IB:**

Investigation, Data curation, Writing – Review & Editing. **LM:** Investigation, Data curation.

ARG: Conceptualization, Investigation, Writing – Review & Editing, Supervision.

Funding: This research was supported by an NSERC Discovery Grant (Canada), the Faculty

of Sciences at Universidad de los Andes (Bogotá, Colombia), the Walt Halperin Endowed

Professorship and the Washington Research Foundation as Distinguished Investigator to A.R.-

G., and the Emerging Leaders in the Americas Program (Global Affairs Canada).

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgments: We thank Steven Cardiff and Nick Mason for their support during

fieldwork and specimen access, as well as the Louisiana State University Museum of Natural

Science for facilitating access to their ornithological collections.

Abstract

Understanding how locomotion-related skeletal elements evolve under biomechanical and ecological constraints is central to animal evolutionary biology. In hummingbirds (Trochilidae), the humerus plays a key role in force transmission during hovering and flapping flight, yet the drivers of its shape evolution have not been examined. We combined geometric morphometrics with phylogenetic comparative analyses to examine humeral shape variation, evolutionary rates, and phenotypic integration in male hummingbirds from 78 species. Our analyses identified humerus allometry as the dominant predictor of shape, revealing a pattern in which larger humeri show broader proximal epiphyses, increased shaft robustness, and reduced curvature. In addition to this scaling pattern, we find that male humerus shape differs subtly among hummingbird species that differ in the use of aggression during nectar foraging. Evolutionary rates of humeral shape were heterogeneous and decoupled from ecological predictors. We also find phenotypic integration between the proximal and distal regions of the humerus, indicating coordinated evolution. Together, these results show that humeral evolution in hummingbirds is governed primarily by biomechanical scaling and internal integration, with foraging ecology introducing secondary, size-dependent modifications. This work highlights the importance of considering scaling and internal integration when interpreting morphological evolution in locomotor systems across vertebrates.

Keywords: Adaptive evolution, Allometry, Biomechanics, Bird flight, Geometric morphometrics, Phenotypic modularity and integration.

Introduction

Among skeletal elements, the avian humerus plays a central role in flight mechanics and ecological adaptation, connecting the axial skeleton to the forelimb and anchoring the primary muscles driving the wingbeat [1]. Variation in humeral morphology is expected to influence wingbeat kinematics and, indirectly, the magnitude and direction of aerodynamic forces, as well as flight control and maneuverability [2]. Despite this biomechanical importance, comparative studies of avian flight have largely focused on external features such as wing aspect ratio or wing loading [3], which capture overall aerodynamic performance, but fail to reflect underlying skeletal variation and musculoskeletal function.

In hummingbirds (family Trochilidae), the humerus is short and robust, reflecting the extreme mechanical demands of their specialized flight mode, characterized by sustained hovering, high wingbeat frequencies owing to their small body size, and rapid wing rotation throughout the stroke [2]. Unlike the long, slender humeri typical of birds adapted for gliding or forward-powered flight [4], the hummingbird humerus functions as a rigid lever transmitting large forces from enlarged flight muscles that drive lift production and rapid wing rotation [5]. This configuration enables precise control during hovering and agile maneuvers but also imposes high mechanical loads [2]. As a result, compared to other birds, the hummingbird humerus exhibits increased cortical thickness, a relatively straight shaft, and pronounced muscle attachment sites at the deltopectoral crest (e.g., Fig. 1A; [6]). This morphology reflects a biomechanical trade-off between minimizing inertial resistance during rapid wing movements and maintaining the structural stiffness required to transmit large aerodynamic forces. Such opposing demands are expected to strongly constrain humeral shape, making it a key element for understanding the evolution of flight-related morphology in hummingbirds. Despite the functional centrality of the humerus, and the exceptional diversity of hummingbird family

(>350 species), comparative analyses of its evolution across this group have not been

examined.

From a biomechanical perspective, the evolution of load-bearing skeletal elements is

constrained by scaling relationships between size, force production, and structural resistance.

Under geometric similarity, body mass scales with linear dimensions cubed, whereas

resistance to bending or axial loads scales with cross-sectional area and thus with linear

dimensions squared [7]. Maintaining constant mechanical stress therefore requires load-

bearing bones to scale at greater than isometry. This expectation applies broadly to locomotor

elements, including the humerus, which experiences substantial bending and torsional loads

during the wingbeat. Even in the absence of ecological differentiation, fundamental

biomechanical principles predict a disproportionate increase in humeral robustness with body

size.

The above general biomechanical expectations are further modified by the exceptional

allometric scaling of the wing in hummingbirds, a group where species vary ~10-fold in body

mass. Unlike most birds, larger hummingbird species exhibit disproportionately large wings

relative to body mass, and unusually low wing loading [8–10], a departure from isometry that

is central to sustained hovering flight. As a result, aerodynamic and inertial forces acting on

the wing do not increase proportionally with body mass alone. Because these forces are

transmitted proximally through the wing skeleton, the humerus must accommodate both

increases in body size and the disproportionate scaling of wing dimensions. Consequently,

humeral morphology reflects an interaction between general biomechanical constraints and

clade-specific wing allometry, making hummingbirds an informative system for studying size-

related skeletal evolution.

Hummingbirds also exhibit species differences in foraging behavior that are linked to the use

of different flight modes. In some species, males primarily defend floral territories, engaging

in aggressive chases, rapid accelerations, and tight turns, whereas other species adopt

traplining strategies involving predictable routes and sustained flight among dispersed

resources [9,11]. These differences in the use of aggressive behavior while foraging are

expected to impose different biomechanical demands on the flight apparatus. Territorial

hummingbirds may benefit from shorter, more robust humeri that enhance force production

and maneuverability [2], whereas traplining behavior may favor longer, more slender humeri

associated with sustained flight efficiency [12,13]. However, foraging strategies are coarse

ecological proxies and may be confounded with body size, habitat structure, and sex-specific

behaviors. In many hummingbird species, males and females differ in morphology and flight

performance, often as a result of sexual selection for enhanced agility in males [14–17]. Here,

we focus primarily on humerus evolution in males, but we present measurements from a

smaller sample of females in the Supplementary Material.

Beyond foraging strategy, additional species differences related to the flight environment and

larger-scale movements may also shape humeral morphology. Habitat density modulates

maneuvering demands, with closed environments requiring frequent turning and fine-scale

control, whereas open habitats are expected to favor fast flight and straighter flight paths

[18,19]. Migratory behavior represents another axis influencing morphological evolution,

from resident species to those undertaking seasonal migrations that require sustained long-

distance flight [20]. Broad environmental gradients such as elevation, latitude, and geographic

range size may further shape humeral morphology. High-elevation environments impose

hypoxia and reduced air density, increasing the mechanical and metabolic demands of flight

and driving divergence in wing shape, body size, and metabolic traits in hummingbirds [21–

23]. Latitude and geographic range size similarly capture variation in climatic heterogeneity,

which may influence dispersal and exposure to diverse flight conditions [24,25].

Given this background, we expect humeral morphology in hummingbirds to reflect a complex

interplay among biomechanical scaling, ecological demands, and evolutionary history.

However, the extent to which ecological variables predict humeral shape independently of

body size, and whether they are associated with differences in evolutionary rates, remains

underexplored [26,27]. In this context, evaluating evolutionary rates provides a

complementary axis for testing ecological effects beyond static shape differences, as

accelerated shape evolution may reflect ecological opportunity or functional innovation,

whereas low rates may indicate stabilizing selection or strong biomechanical constraints

[28,29].

The hummingbird humerus, with distinct proximal and distal functions, also provides an

opportunity to investigate phenotypic integration. Proximally, the humerus anchors the

pectoralis and supracoracoideus muscles that power the downstroke and upstroke, whereas

distally it articulates with the radius and ulna, transmitting forces that shape wing motion and

orientation [1,30]. Understanding whether these regions evolve in a coordinated manner offers

insights into developmental constraints and adaptive flexibility [31–33]. Locomotor traits

rarely evolve independently [34], instead exhibiting integration or modularity driven by

shared developmental pathways and functional covariation [35,36]. While most studies

examine integration across multiple skeletal elements, analyzing integration within a single

bone can reveal how functionally distinct regions remain evolutionarily coordinated despite

localized biomechanical specialization.

Here, we investigate how biomechanical scaling, species ecology, environmental gradients,

and evolutionary processes jointly shape humeral morphology in male hummingbirds. First,

we tested whether humeral shape exhibits strong allometric scaling with body size. In this

clade, wing length increases more than expected from isometric scaling with body mass

(approximately mass^0.5), resulting in disproportionately long wings in larger species [8–10].

Because the humerus transmits aerodynamic and inertial forces from the wing to the body, we

predict that humeral robustness exhibits positive allometry (i.e., scales greater than isometry)

with body size. We further assessed whether allometric variation in humeral shape is

associated with differences in evolutionary rates, recognizing that size-related constraints may

promote directional change or impose stabilizing selection.

Second, we examined whether ecological context modulates humeral morphology and

evolutionary rates beyond size and allometry. We hypothesize that differences in foraging

behavior will impose distinct biomechanical demands: we expect territorial species to show

shorter, more robust humeri consistent with enhanced force transmission during high-power

flight, whereas traplining species may exhibit more elongated humeri associated with

sustained flight efficiency [9,37]. We further predicted complementary effects of habitat

structure and migratory strategy, with closed habitats favoring greater maneuverability and

open habitats favoring straighter flight trajectories; we expect migratory species to exhibit

morphologies linked to sustained directional flight [12,18,38]. In contrast, when examining

evolutionary rates, we did not assume the ecological predictors would have a single

directional effect, as selection may promote innovation or impose stabilizing constraints

depending on ecological variability and biomechanical trade-offs [39].

Third, we assessed whether broad environmental gradients shape humeral morphology and

evolutionary rates. High elevation imposes hypoxia and reduced air density, increasing

aerodynamic and metabolic demands. We therefore predicted that elevation may be associated

with shifts in humeral morphology, whereas its effects on evolutionary rates could arise either

through increased selective pressure promoting faster change, or through environmental

constraints limiting phenotypic divergence. Latitude and species' geographic range size are

broad descriptors of environmental variation, and we accounted for these in our analyses,

although we did not have a priori directional expectations for these two predictors.

Lastly, we hypothesized strong internal phenotypic integration in the hummingbird humerus,

with coordinated evolution between proximal and distal regions due to their shared role in

force transmission and wing mechanics. This is expected because modifications in one region

(e.g., altering muscle attachment sites) would require compensatory changes to maintain flight

performance. However, bones may also evolve modularly when regions face distinct

functional demands [34,40,41]. Whether such modularity also occurs within the hummingbird

humerus remains an open question, with implications for how evolutionary change may be

distributed across the structure. By integrating ecological predictors and morphometric

analyses within a phylogenetic framework, we provide a comprehensive assessment of the

drivers of humeral morphological evolution.

Methods

Specimens and image acquisition

We imaged humeral morphology of skeletal specimens from 80 hummingbird species (~22%

of extant Trochilidae with 7 of 9 clades represented in our sample [42]) at the Louisiana State

University Museum of Natural Science (LSUMNS, see Table S1 for specimen details).

Sampling was uneven due to limited skeletal material: ~87% of species (70/80) were

represented by a single male specimen. We focused our main analyses on male specimens to

minimize variation associated with sexual dimorphism and to ensure sampling consistency, as

male skeletal material is more widely available [43]. We provide a preliminary

characterization of a smaller sample of females in the supplement.

The right humerus was photographed in cranial view with a scale bar (0.1 mm divisions)

using a Canon EOS M50 (6000 × 4000 px) with a 28 mm macro lens. Images were acquired

with orthogonal orientation and fixed camera distance using a copy stand to ensure cross-

species comparability [44].

We digitized 25 homologous anatomical landmarks per humerus, primarily along the outline

(Fig. 1A–B), using TPSDig2 v2.32 [45]. Landmarks were selected for anatomical consistency

and biomechanical relevance; semilandmarks were not used (see Table S2 for details). We

screened each photograph for humerus orientation, planar alignment, and landmark visibility

and excluded specimens that failed to meet these criteria. Two species (*Calypte anna* and

Goethalsia bella) were excluded because the specimens (one individual each) did not meet

landmark quality criteria. After screening, 78 species remained for downstream analyses.

To assess the potential influence of geometric uncertainty on the morphometric dataset, we

considered several non-statistical sources of measurement error. Image resolution introduces

≤1–2 pixels of landmark uncertainty (~0.1% of humeral length), intra-observer digitization

error is typically 0.5–1.5%, inter-observer error is negligible (single observer), and local

anatomical ambiguity may add ~1–3% variability. Combined uncertainty (~2–4% of inter-

landmark distances) is well below interspecific shape differences and unlikely to affect major

axes of variation or phylogenetic patterns.

Morphometric data preparation

We conducted all analyses in R 4.4.2 [46]. We aligned landmark configurations using

Generalized Procrustes Analysis (GPA) to remove effects of position, orientation, and scale

[47], using gpagen() in *geomorph* [48]. For species represented by more than one specimen,

we used the species-average humerus shape in all subsequent analyses. We used principal

component analysis (PCA) of the Procrustes-aligned coordinates to visualize overall shape

variation and morphospace structure (Fig. 1C). We did not use PCA scores for hypothesis

testing.

Species traits

For each species, we compiled ecological, geographic, and morphological predictors. Species

body mass (g) from Tobias et al., (2022) [49] was included as a proxy for biomechanical

scaling, given its relationship with aerodynamic force production. We compiled estimates of

species geographic range size (km²) and range centroid latitude (absolute value) from Tobias

et al., (2022) [49], and species maximum elevation (m) from Quintero and Jetz (2018) [50].

Body mass and range size values were log-transformed prior to analysis to improve linearity

and reduce heteroscedasticity. As a measure of humerus size, we used landmark centroid size,

a matrix-derived measure based on Procrustes analysis [51].

Each species' primary foraging strategy was classified into three categories—territorial,

trapliner, and opportunist—using Feinsinger's framework [52] and Rombaut et al., (2022)

[53]. According to this framework, we classify species as territorial if the males primarily

defend spatially stable floral patches and exclude competitors through aggressive interactions,

whereas species are classified as trapliners if the males primarily visit dispersed flowers and

generally do not defend feeding territories. We classify species as opportunists if the males

that exhibit flexible or context-dependent foraging behavior (i.e., species that are described as

using territorial or non-territorial foraging, depending on the location or conditions). For each

species, we manually reviewed foraging strategies assigned in Rombaut et al., (2022) [53] and

the corresponding species account in *Birds of the World* [54] to ensure consistency, and to

classify species that were not included in Rombaut et al., (2022) [53] (see Table S3 for

details).

Habitat density was obtained from Tobias et al., (2022) [49], with each species categorized as

using dense, semi-open, or open habitats. Species migratory behavior was classified into four

categories: (i) resident, (ii) altitudinal migrant, (iii) short-distance latitudinal migrant, and (iv)

long-distance latitudinal migrant, based on species accounts in *Birds of the World* [54],

BirdLife International [55], and Barçante (2017) [56]. We considered a species to be

migratory (ii - iv) only when explicit evidence of documented movement was available.

Phylogenetic signal

We assessed phylogenetic signal in humeral shape using Blomberg's multivariate K statistic

[33,57], using physignal() in *geomorph*, based on 1,000 permutations. All analyses used an

ultrametric phylogeny from McGuire et al., (2014) [58] pruned to the final species set.

Phylogenetic modeling of humeral shape variation

To evaluate how ecological and morphological predictors influence humeral shape, we used a

phylogenetic generalized least squares (PGLS) framework using procD.pgls() in *geomorph*,

with Pagel's λ estimated by maximum likelihood in the final model [59].

We first evaluated a set of *a priori* candidate ecological models (Table S4) and then fitted a

final multivariate model including humerus centroid size, body mass, foraging strategy,

migratory status, habitat density, geographic range size, range centroid latitude (absolute

value), and maximum elevation. Categorical predictors were coded using sum-to-zero

contrasts so that tests reflect overall deviations of each factor from the grand mean rather than

comparisons to a single reference level. We verified that collinearity among predictors was

low (Pearson's $|r| < 0.36$; all adjusted GVIF < 1.5, Table S5). Statistical significance was

assessed using 1,000 permutations of residuals under a phylogenetic correlation structure.

Because sequential (Type I) sums of squares are sensitive to predictor order (Table S6), we

based our inferences on Type III sums of squares, providing order-invariant tests of each

predictor's partial effect while controlling for all others.

We evaluated the sensitivity of PGLS analyses to assumptions about phylogenetic signal

(Pagel's λ) by refitting the final additive model across a range of fixed λ values (0–1) (see

supplement Table S7 for details). We performed additional sensitivity analyses to consider

interactions between species size and foraging behavior using the same inferential framework

(Table S8).

274 Discriminant analysis of humeral shape by foraging strategy

To test whether humeral shape differs among foraging strategies, we conducted a Canonical

Variates Analysis (CVA) using size-corrected Procrustes coordinates. Allometric effects were

removed by regressing the aligned shape coordinates on log centroid size and using the

residuals as inputs [51]. CVA was implemented with CVA() in *Morpho* [60], with foraging

strategy as the grouping factor, and group separation was evaluated using 1,000 permutation

tests. Phylogenetic structure in canonical space was visualized by projecting species scores

onto the phylogeny using *phytools* [61].

282 Evolutionary rates of humeral shape

We tested whether humeral shape has evolved at different rates in hummingbirds with

different ecological traits using compare.evol.rates() in geomorph with 1,000 permutations.

We conducted these rate comparisons for the following categorical predictors: foraging

strategy, habitat density, and migratory behavior, by evaluating whether group-specific mean

rates deviated from expectations given the phylogeny and the distribution of shape change

estimates across species.

For continuous predictors, we tested associations between evolutionary rates and each

predictor using search.trend()in *RRphylo* [62], which fits phylogenetically informed

regressions between branch-specific rate estimates and predictor values. We applied standard

RRphylo diagnostics to identify potential outliers using the built-in functions

RRphylo::outliers() and *RRphylo*::search.trend(), summarized in Table S9.

We additionally modeled species-specific evolutionary rates as a univariate response using

phylogenetic generalized least squares (PGLS) fitted with *nlme*::gls() under a Brownian-

motion correlation structure derived from the same ultrametric phylogeny. Each predictor was

evaluated in a separate univariate model.

Phenotypic modularity and integration

To evaluate phenotypic integration within the humerus, we partitioned the Procrustes-aligned

landmark data into two functional modules defined based on avian flight anatomy; we refer to

this as the *a priori* modularity hypothesis. According to this hypothesis, the proximal module

includes the humeral head, tubercles, and associated articular surfaces involved in shoulder

rotation and load transfer during both hovering and flapping. The distal–shaft module

comprises the diaphysis, trochlea, capitulum, and distal muscle insertion sites, which

primarily mediate force transmission, wingbeat mechanics, and resistance to bending loads

during forward flight [30,63]. This subdivision reflects the well-established biomechanical

differentiation between the articulation region of the humerus and its load-bearing shaft. We

quantified evolutionary covariation between the *a priori* modules using phylogenetic partial

least squares implemented with phylo.integration() in *geomorph*, assessing significance of

modularity with 1,000 phylogenetically informed permutations. We did not perform

alternative modularity searches (e.g., EMMLi-based optimization) because our objective was

to test an anatomically motivated prediction of integration rather than explore the full modular

landscape of the humerus.

 Results

 Phylogenetic signal

Humeral shape exhibited a significant phylogenetic signal (K = 0.309, p = 0.001, Fig. 1D),

indicating that closely related hummingbird species tend to share similar humeral

morphologies. The geometric configuration used to quantify humeral shape revealed

consistent variation in landmark positions across species, highlighting major axes of

morphological differentiation (Fig. 1A–C).

Predictors of humeral shape variation

The multivariate PGLS analysis identified humerus centroid size as the strongest predictor of

humeral shape variation ($R^2 = 0.098$, Z = 3.26, p = 0.001; Table 1; Fig. 2). Foraging strategy

also showed a significant association with humeral shape ($R^2 = 0.034$, Z = 1.98, p = 0.025).

Several additional predictors exhibited marginal effects: body mass ($R^2 = 0.017$, p = 0.070),

migratory behavior ($R^2 = 0.045$, p = 0.058), maximum elevation ($R^2 = 0.015$, p = 0.095), and

absolute latitude ($R^2 = 0.014$, p = 0.112) each explained small proportions of shape variation,

but did not reach statistical significance. We did not detect effects of habitat density ($R^2 =$

0.018, p = 0.394), or geographic range size ($R^2 = 0.010$, p = 0.369) on humeral shape.

Across all predictors, effect sizes were modest, and the final model explained a limited

fraction of total multivariate variation in humeral shape (ca. 25%), with size-related effects

accounting for the largest share. Sensitivity analyses across values of Pagel's λ showed that

the effects of humerus size and foraging strategy were consistent (Table S7). The effects of

humerus size and foraging strategy on humerus shape were also robust to the inclusion of an

interaction between these two variables (Table S8).

Morphospace differentiation by foraging strategy

PCA revealed substantial overlap in humeral morphospace among foraging strategies (Fig.

3A). The morphospaces of territorial and traplining species were largely contained within the

broader morphospace occupied by opportunistic species, as illustrated by convex hulls.

Projection of phylogenetic relationships onto the morphospace indicated some local clustering

of related taxa, consistent with the detected phylogenetic signal, but no strong global

structuring by clade.

Canonical Variates Analysis further characterized differences among foraging strategies (Fig.

3C). The first canonical axis primarily separated traplining species from territorial and

opportunistic taxa, whereas the second axis distinguished territorial from opportunistic

species. However, cross-validated classification accuracy was low (42.3%), only slightly

exceeding random expectations, and all strategies showed extensive overlap in canonical

space. Traplining species exhibited higher correct classification (47.6%) than territorial and

opportunistic species (36–43.7%). Consistent with these patterns, a permutational multivariate

test on size-corrected shape data detected small but significant differences in mean humeral

shape among foraging strategies ($R^2 = 0.060$, p = 0.001).

Evolutionary rates of humeral shape

Estimates of evolutionary rates derived from *RRphylo* revealed a strongly right-skewed

distribution, with most species exhibiting low rates of humeral shape evolution (median =

0.0063; interquartile range = 0.0030–0.0120, Fig. 1D). Only a small number of branches

showed elevated rates, indicating isolated episodes of accelerated morphological change

(Table S9). Phylogenetic signal in evolutionary rates was weak ($\lambda \approx 0.03$) and not

significantly different from zero (LR = 0.040, p = 0.840), indicating that closely related

species did not evolve at more similar rates than expected under phylogenetic independence.

Group-wise comparisons detected no significant differences in evolutionary rates among

foraging strategies ($p = 0.138$), habitat density categories ($p = 0.503$), or migratory classes (p

$= 0.953$; Table 2). Similarly, univariate PGLS models showed no significant associations

between evolutionary rates and continuous predictors, including body mass, humerus centroid

size, geographic range size, elevation, or latitude (all $p > 0.13$; Table 2). These results indicate

that variation in humeral evolutionary tempo is not systematically associated with these

ecological or morphological predictors.

Phenotypic modularity and integration

Phylogenetic partial least-squares analysis revealed strong integration between the proximal

and distal regions of the humerus (r-PLS = 0.869, Z = 4.014, $p = 0.001$; Fig. 5). This indicates

that morphological changes in one region are closely mirrored by changes in the other,

consistent with coordinated evolution across the bone. The magnitude and significance of

integration were robust to phylogenetic correction and did not depend on ecological or size-

related predictors.

376   **Discussion**

Allometric scaling as the dominant driver of humeral shape evolution

Among all predictors evaluated, humerus centroid size showed the strongest and most

consistent association with humeral shape, whereas body mass exerted only a marginal effect.

This allometric signal indicates that increasing humeral size is associated with broader

proximal epiphyses, increased mediolateral shaft robustness, and reduced curvature. Similar

size-related reinforcement of proximal limb bones has been documented across volant birds

and other flying vertebrates, where increases in cross-sectional dimensions enhance resistance

to bending and torsional stresses [12,30,63]. These trends are consistent with general scaling

principles linking skeletal strength to geometry rather than body mass per se [7,63].

Biologically, this pattern indicates that selection acting on flight performance in larger

hummingbirds favors humeral morphologies that enhance joint stability and load-bearing

capacity. Expanded articular surfaces increase muscle attachment area and force transmission

at the shoulder, particularly for the pectoralis and supracoracoideus, whereas straighter and

more robust diaphyses reduce bending moments generated during flapping and hovering

[1,10,30]. Comparable relationships between bone robustness, joint morphology, and force

transmission have been reported in avian forelimb mechanics [64,65]. This interpretation is

especially informative given the unusual scaling of the hummingbird wing: distal

components, particularly the manus and primary feathers, scale disproportionately with body

mass, producing relatively long wings and low wing loading [8–10]. As a result, aerodynamic

and inertial forces generated distally increase nonlinearly with size and must be transmitted

proximally through the humerus. This contrast between elongation of distal wing elements

and structural reinforcement of the humerus is consistent with previous work showing that

elongation of distal elements contributes to increased aerodynamic efficiency [12,18,66].

Overall, these results show that allometric scaling imposes a dominant and predictable

constraint on humeral evolution in hummingbirds, shaping mechanically robust

configurations that stabilize force transmission during hovering and flapping flight across

ecological contexts.

Ecological effects as secondary modifiers of humeral shape

Although allometric scaling explains most of the variation in male humerus shape, our results

suggest that foraging and competitive behavior act as a secondary modifier. Foraging strategy

showed a statistically significant but modest association with humeral shape, with differences

among species that use territorial, traplining, and opportunistic foraging behavior. These

differences are modest, such that foraging categories overlap within a shared morphospace

adapted to hovering flight, rather than discrete or mutually exclusive morphotypes. Principal

component and canonical variate analyses also revealed that opportunistic hummingbird

species span most of the variation occupied by territorial and traplining taxa. Although

multivariate tests detected statistically significant differences in mean shape after accounting

for size, these differences were small relative to overall variation. Similarly modest

differentiation has been reported for other avian skeletal traits [67,68].

Despite their modest magnitude, associations between humerus shape and foraging behaviour

were consistent with our predictions. Traplining species tend to exhibit relatively longer, more

slender humeri, consistent with sustained forward flight over extended circuits, where

elongation of the proximal wing segment may contribute to increased effective wingspan and

reduced wingbeat frequency [12,13,66,69]. Territorial species tend to show shorter, thicker,

more curved humeri; these features are expected to reduce shoulder moment of inertia,

facilitating rapid angular acceleration and tight maneuvering during aggressive chases and

vertical ascents [9,11,37,38].

Humeral elongation is not the sole route to increased wingspan or aerodynamic efficiency:

similar outcomes can arise from elongation of distal wing elements, often with lower inertial

costs [12,18]. Accordingly, we do not assume a one-to-one correspondence between humeral

length and wing length; rather, humeral geometry likely interacts with distal skeletal and

feather elements to shape overall wing planform and flight performance, making humeral

elongation a plausible but not exclusive pathway contributing to strategy-associated

differences. Our analyses also suggest that foraging effects may be size-dependent, rather than

uniform. We found evidence for an interaction between foraging strategy and humerus

centroid size (Table S8). These results suggest that foraging strategy may bias humeral shape

evolution conditional on size, instead of acting as an independent driver of morphology.

Other ecological and geographic variables, including migratory behavior, habitat density,

latitude, and elevation, showed marginal or no detectable associations with humeral shape,

consistent with reports for other avian morphological traits once size and phylogeny are

accounted for [25,70]. Habitat density and migratory categories are proxies for flight-space

structure and movement ecology. Their lack of detectable associations with humerus shape

suggests that they may primarily act on distal wing/feather components, or their effects may

be overshadowed by size-related load-transmission constraints in the humerus.

The humerus—by virtue of its mechanically integrated, load-bearing role—may be buffered

against many ecological pressures, with selection acting primarily within limits imposed by

allometric scaling and structural integration [30,37]. Future studies using three-dimensional

humeral models may reveal additional axes of shape variation—particularly related to

articular surfaces and torsional geometry—not captured by the present two-dimensional

approach, potentially uncovering stronger or complementary associations with ecological and

functional predictors.

Phylogenetic structure as background context

Across all analyses, humeral shape exhibited significant phylogenetic signal, indicating

resemblance among related species. This pattern is typical of complex morphological traits

and reflects shared ancestry, developmental pathways, and conserved biomechanical solutions

[33,57]. Importantly, this phylogenetic structure provides the historical backdrop against

which both shape variation and evolutionary rates are interpreted. By accounting for shared

ancestry, size-dependent scaling and ecological effects can be evaluated as deviations from

inherited morphological similarity, rather than artifacts of phylogenetic relatedness [39].

Evolutionary rates: heterogeneity without ecological predictors

In contrast to patterns of shape variation, evolutionary rates of male humeral morphology showed no consistent association with ecological predictors. Rates were highly heterogeneous across the phylogeny, with most branches exhibiting low values and only a few showing elevated rates. Rate estimates also lacked detectable phylogenetic signal, and we did not detect any associations between evolutionary rates and any other species traits. Together, these patterns suggest that variation in evolutionary tempo arises idiosyncratically rather than being structured by shared ancestry [28,39]. Similar decoupling between morphological disparity and evolutionary tempo has been reported in other macroevolutionary studies, where strong constraints on form coexist with heterogeneous, lineage-specific rates of change [27,29].

These results caution against adaptive narratives that link particular ecological strategies to accelerated skeletal evolution without explicit support from evolutionary rate analyses. Although maximum elevation showed weak explanatory power in the multivariate models, individual lineages exhibited elevated evolutionary rates, including *Metallura phoebe*, a high-elevation species identified as a rate outlier (Table S9). Such cases suggest that localized increases in evolutionary rate may arise from recent, lineage-specific processes—such as shifts in functional integration, developmental constraints, or population-level dynamics—rather than reflecting a general effect of elevation on humeral evolution. Importantly, the presence of these outliers does not contradict the overall absence of a strong elevational signal, but instead highlights the distinction between global macroevolutionary trends and idiosyncratic evolutionary trajectories at the species level [29].

Phenotypic integration as a key constraint on humeral evolution

We find significant phenotypic integration between proximal and distal regions of the

hummingbird humerus. This finding indicates coordinated evolution, consistent with the

shared role of the proximal and distal regions in force transmission during flight [1,30]. Such

integration is expected in highly loaded locomotor structures, where changes in one region

require compensatory adjustments elsewhere to preserve functional integrity [36,71].

This integration also provides an explanation for several observed patterns in our

phylogenetic analysis. It helps explain why ecological effects on humeral shape are modest, as

selection cannot act independently on isolated regions of the bone, and it indicates that

evolutionary change in the humerus is channeled along constrained trajectories defined by

internal structural coherence. Similar roles of integration in shaping evolutionary constraint

have been documented across vertebrate skeletal systems [72,73]. Together, these results

show that phenotypic integration simultaneously enables coordinated functional refinement

and limits morphological disparity. Even when ecological pressures promote divergence in

humeral shape, internal structural coherence remains evolutionarily conserved.

Conclusion

Taken together, our results show that humeral evolution in hummingbirds is governed

primarily by biomechanical scaling and internal structural integration, maintaining a skeletal

configuration optimized for sustained hovering flight, with ecological factors introducing

secondary, constrained modifications to shape. While behaviors such as foraging strategy

leave detectable signatures on humeral morphology, these effects are modest relative to the

dominant influence of size-dependent mechanics. Moreover, evolutionary rates of humeral

change appear largely decoupled from ecological predictors, highlighting the importance of

constraint and historical contingency in shaping macroevolutionary patterns. This integrative

perspective underscores that, even within a lineage renowned for flight specialization, the

evolution of skeletal form reflects a balance between universal biomechanical principles and

the selective nuances of ecological diversification.

507   **Data availability statement**

508 Data and code for all analyses are available on Figshare at:

509 https://figshare.com/s/2c012d7ba82c9eeda5ed

References

1. Tobalske BW, Biewener AA. 2008 Contractile properties of the pigeon supracoracoideus during different modes of flight. *Journal of Experimental Biology* **211**, 170–179. (doi:10.1242/jeb.007476)

2. Agrawal S, Tobalske BW, Anwar Z, Luo H, Hedrick TL, Cheng B. 2022 Musculoskeletal wing-actuation model of hummingbirds predicts diverse effects of primary flight muscles in hovering flight. *Proceedings of the Royal Society B: Biological Sciences* **289**. (doi:10.1098/rspb.2022.2076)

3. Hieronymus TL. 2015 Qualitative skeletal correlates of wing shape in extant birds (Aves: Neoaves). *BMC Evol. Biol.* **15**, 30. (doi:10.1186/s12862-015-0303-7)

4. Dial KP. 1992 Avian Forelimb Muscles and Nonsteady Flight: Can Birds Fly without Using the Muscles in Their Wings? *Auk* **109**, 874–885. (doi:10.2307/4088162)

5. Warrick DR, Tobalske BW, Powers DR. 2005 Aerodynamics of the hovering hummingbird. *Nature* **435**, 1094–1097. (doi:10.1038/nature03647)

6. Serrano FJ, Costa-Pérez M, Navalón G, Martín-Serra A. 2020 Morphological Disparity of the Humerus in Modern Birds. *Diversity (Basel).* **12**, 173. (doi:10.3390/d12050173)

7. Vogel S. 2003 *Comparative Biomechanics: Life's Physical World*. 1st edn. Princeton, NJ.: Princeton University Press.

8. Greenewalt CH. 1962 Dimensional Relationships for Flying Animals. *Smithsonian Miscellaneous Collections* **144**, 1–46.

9. Altshuler DL, Stiles FG, Dudley R. 2004 Of Hummingbirds and Helicopters: Hovering Costs, Competitive Ability, and Foraging Strategies. *Am. Nat.* **163**, 16–25. (doi:10.1086/380511)

10. Skandalis DA *et al.* 2017 The biomechanical origin of extreme wing allometry in hummingbirds. *Nat. Commun.* **8**, 1047. (doi:10.1038/s41467-017-01223-x)

11. Sargent AJ, Groom DJE, Rico-Guevara A. 2021 Locomotion and Energetics of Divergent Foraging Strategies in Hummingbirds: A Review. *Integr. Comp. Biol.* **61**, 736–748. (doi:10.1093/icb/icab124)

12. Nudds RL, Dyke GJ, Rayner JM V. 2007 Avian brachial index and wing kinematics: putting movement back into bones. *J. Zool.* **272**, 218–226. (doi:10.1111/j.1469-7998.2006.00261.x)

13. Sievwright H, Macleod N. 2012 Eigensurface analysis, ecology, and modelling of morphological adaptation in the falconiform humerus (Falconiformes: Aves). *Zool. J. Linn. Soc.* **165**, 390–419. (doi:10.1111/j.1096-3642.2012.00818.x)

14. Wilcox SC, Clark CJ. 2022 Sexual selection for flight performance in hummingbirds. *Behavioral Ecology* **xx**, 2–14. (doi:10.1093/beheco/arac075)

15.   Temeles EJ, Miller JS, Rifkin JL. 2010 Evolution of sexual dimorphism in bill size and
shape of hermit hummingbirds (Phaethornithinae): a role for ecological causation.
Philosophical Transactions of the Royal Society B: Biological Sciences **365**, 1053–
1063. (doi:10.1098/rstb.2009.0284)

16.   Garzón-Agudelo F, Mansfield L, Epperly K, Rico-Guevara A. 2025 Sharper, straighter,
stiffer, stronger: sexually dimorphic bill shape enhances male stabbing performance in
the green hermit hummingbird (Phaethornis guy). *Journal of Experimental Biology*
228. (doi:10.1242/jeb.250769)

17.   Stiles FG, Altshuler DL, Dudley R. 2005 Wing Morphology and Flight Behavior of
Some North American Hummingbird Species. *Auk* **122**, 872–886.
(doi:10.1093/auk/122.3.872)

18.   Norberg UM. 1990 *Vertebrate Flight*. Berlin, Heidelberg: Springer Berlin Heidelberg.
(doi:10.1007/978-3-642-83848-4)

19.   Dudley R. 2002 Mechanisms and Implications of Animal Flight Maneuverability.
Integr. Comp. Biol. **42**, 135–140. (doi:10.1093/icb/42.1.135)

20.   Alerstam T, Hedenström A, Åkesson S. 2003 Long☐distance migration: evolution and
determinants. *Oikos* **103**, 247–260. (doi:10.1034/j.1600-0706.2003.12559.x)

21.   Altshuler DL. 2006 The physiology and biomechanics of avian flight at high altitude.
Integr. Comp. Biol. **46**, 62–71. (doi:10.1093/icb/icj008)

22.   Stiles FG. 2008 Ecomorphology and phylogeny of Hummingbirds: Divergence and
Convergence in Adaptations to High Elevations. *Ornitol. Neotrop.* **19**, 511–519.

23.   Altshuler DL, Dudley R, McGuire JA. 2004 Resolution of a paradox: Hummingbird
flight at high elevation does not come without a cost. *Proceedings of the National*
Academy of Sciences **101**, 17731–17736. (doi:10.1073/pnas.0405260101)

24.   Yang J, Yang C, Lin H, Lees AC, Tobias JA. 2024 Elevational constraints on flight
efficiency shape global gradients in avian wing morphology.
(doi:10.1101/2024.07.12.603304)

25.   Claramunt S, Derryberry EP, Remsen J V., Brumfield RT. 2012 High dispersal ability
inhibits speciation in a continental radiation of passerine birds. *Proceedings of the*
Royal Society B: Biological Sciences **279**, 1567–1574. (doi:10.1098/rspb.2011.1922)

26.   Chira AM, Cooney CR, Bright JA, Capp EJR, Hughes EC, Moody CJA, Nouri LO,
Varley ZK, Thomas GH. 2018 Correlates of rate heterogeneity in avian
ecomorphological traits. *Ecol. Lett.* **21**, 1505–1514. (doi:10.1111/ele.13131)

27.   Teplitsky C *et al.* 2014 Assessing Multivariate Constraints to Evolution across Ten
Long-Term Avian Studies. *PLoS One* **9**, e90444. (doi:10.1371/journal.pone.0090444)

28.   Harmon LJ *et al.* 2010 Early bursts of body size and shape evolution are rare in
comparative data. *Evolution (N Y).* , no-no. (doi:10.1111/j.1558-5646.2010.01025.x)

29. Cooney CR, Bright JA, Capp EJR, Chira AM, Hughes EC, Moody CJA, Nouri LO,
Varley ZK, Thomas GH. 2017 Mega-evolutionary dynamics of the adaptive radiation
of birds. *Nature* **542**, 344–347. (doi:10.1038/nature21074)

30. Poore SO, Ashcroft A, Sánchez-Haiman A, Goslow GE. 1997 The Contractile
Properties of the M. Supracoracoideus in the Pigeon and Starling: a Case for Long-
Axis Rotation of the Humerus. *Journal of Experimental Biology* **200**, 2987–3002.
(doi:10.1242/jeb.200.23.2987)

31. Hallgrímsson B, Katz DC, Aponte JD, Larson JR, Devine J, Gonzalez PN, Young NM,
Roseman CC, Marcucio RS. 2019 Integration and the Developmental Genetics of
Allometry. *Integr. Comp. Biol.* **59**, 1369–1381. (doi:10.1093/icb/icz105)

32. Felice RN, Watanabe A, Cuff AR, Noirault E, Pol D, Witmer LM, Norell MA,
O'Connor PM, Goswami A. 2019 Evolutionary Integration and Modularity in the
Archosaur Cranium. *Integr. Comp. Biol.* **59**, 371–382. (doi:10.1093/icb/icz052)

33. Adams DC, Felice RN. 2014 Assessing Trait Covariation and Morphological
Integration on Phylogenies Using Evolutionary Covariance Matrices. *PLoS One* **9**,
e94335. (doi:10.1371/journal.pone.0094335)

34. Orkney A, Bjarnason A, Tronrud BC, Benson RBJ. 2021 Patterns of skeletal integration
in birds reveal that adaptation of element shapes enables coordinated evolution
between anatomical modules. *Nat. Ecol. Evol.* **5**, 1250–1258. (doi:10.1038/s41559-
021-01509-w)

35. Hallgrímsson B, Jamniczky H, Young NM, Rolian C, Parsons TE, Boughner JC,
Marcucio RS. 2009 Deciphering the Palimpsest: Studying the Relationship Between
Morphological Integration and Phenotypic Covariation. *Evol. Biol.* **36**, 355–376.
(doi:10.1007/s11692-009-9076-5)

36. Klingenberg CP. 2008 Morphological Integration and Developmental Modularity.
Annu. Rev. Ecol. Evol. Syst. **39**, 115–132.
(doi:10.1146/annurev.ecolsys.37.091305.110054)

37. Tobalske BW, Biewener AA, Warrick DR, Hedrick TL, Powers DR. 2010 Effects of
flight speed upon muscle activity in hummingbirds. *Journal of Experimental Biology*
213, 2515–2523. (doi:10.1242/jeb.043844)

38. Dakin R, Segre PS, Straw AD, Altshuler DL. 2018 Morphology, muscle capacity, skill,
and maneuvering ability in hummingbirds. *Science (1979).* **359**, 653–657.
(doi:10.1126/science.aao7104)

39. Revell LJ, Harmon LJ, Collar DC. 2008 Phylogenetic Signal, Evolutionary Process,
and Rate. *Syst. Biol.* **57**, 591–601. (doi:10.1080/10635150802302427)

40. Felice RN, Goswami A. 2018 Developmental origins of mosaic evolution in the avian
cranium. *Proceedings of the National Academy of Sciences* **115**, 555–560.
(doi:10.1073/pnas.1716437115)

41. Navalón G, Marugán-Lobón J, Bright JA, Cooney CR, Rayfield EJ. 2020 The
consequences of craniofacial integration for the adaptive radiations of Darwin's finches
and Hawaiian honeycreepers. *Nat. Ecol. Evol.* **4**, 270–278. (doi:10.1038/s41559-019-
1092-y)

42. Gill F, Donsker D, Rasmussen P. 2023 IOC World Bird List 13.1. *IOC World Bird List
Datasets*. (doi:10.14344/IOC.ML.13.1)

43. Cardini A, Elton S. 2007 Sample size and sampling error in geometric morphometric
studies of size and shape. *Zoomorphology* **126**, 121–134. (doi:10.1007/s00435-007-
0036-2)

44. Zelditch ML, Swiderski DL, Sheets HD, Fink WL. 2012 *Geometric Morphometrics for
Biologists*. 2nd edn. Elsevier. (doi:10.1016/B978-0-12-778460-1.X5000-5)

45. Rohlf FJ. 2015 The tps series of software. *Hystrix* **26**, 1–4. (doi:10.4404/hystrix-26.1-
11264)

46. R Core Team. 2025 R: A language and environment for statistical computing.

47. Rohlf FJ, Slice D. 1990 Extensions of the Procrustes Method for the Optimal
Superimposition of Landmarks. *Syst. Zool.* **39**, 40. (doi:10.2307/2992207)

48. Adams D, Collyer M, Kaliontzopoulou A, Baken E. 2012 geomorph: Geometric
Morphometric Analyses of 2D and 3D Landmark Data. *CRAN: Contributed Packages*.
(doi:10.32614/CRAN.package.geomorph)

49. Tobias JA *et al.* 2022 AVONET: morphological, ecological and geographical data for
all birds. *Ecol. Lett.* **25**, 581–597. (doi:10.1111/ele.13898)

50. Quintero I, Jetz W. 2018 Global elevational diversity and diversification of birds.
Nature **555**, 246–250. (doi:10.1038/nature25794)

51. Klingenberg CP. 2016 Size, shape, and form: concepts of allometry in geometric
morphometrics. *Dev. Genes Evol.* **226**, 113–137. (doi:10.1007/s00427-016-0539-2)

52. Feinsinger P. 1976 Organization of a Tropical Guild of Nectarivorous Birds. *Ecol.
Monogr.* **46**, 257–291. (doi:10.2307/1942255)

53. Rombaut LMK, Capp EJR, Hughes EC, Varley ZK, Beckerman AP, Cooper N, Thomas
GH. 2022 The evolution of the traplining pollinator role in hummingbirds:
specialization is not an evolutionary dead end. *Proceedings of the Royal Society B:
Biological Sciences* **289**. (doi:10.1098/rspb.2021.2484)

54. Billerman SM, Keeney BK, Kirwan GM, Medrano F, Sly ND, Schmidt MG. 2025
Birds of the World. *Cornell Laboratory of Ornithology, Ithaca, NY, USA.
https://birdsoftheworld.org/bow/* (doi:https://doi.org/10.2173/bow)

55. BirdLife International. 2024 IUCN Red List species accounts and distribution maps.
https://datazone.birdlife.org.

56.   Barçante L, M. Vale M, S. Alves MA. 2017 Altitudinal migration by birds: a review of
658   the literature and a comprehensive list of species. *J. Field Ornithol.* **88**, 321–335.
659   (doi:10.1111/jofo.12234)

660   57.   Blomberg SP, Garland T, Ives AR. 2003 Testing for phylogenetic signal in comparative
661   data: behavioral traits are more labile. *Evolution (N Y).* **57**, 717–745.
662   (doi:10.1111/j.0014-3820.2003.tb00285.x)

663   58.   McGuire JA, Witt CC, Remsen JV, Corl A, Rabosky DL, Altshuler DL, Dudley R. 2014
664   Molecular Phylogenetics and the Diversification of Hummingbirds. *Current Biology*
665   **24**, 910–916. (doi:10.1016/j.cub.2014.03.016)

666   59.   Collyer ML, Baken EK, Adams DC. 2022 A standardized effect size for evaluating and
667   comparing the strength of phylogenetic signal. *Methods Ecol. Evol.* **13**, 367–382.
668   (doi:10.1111/2041-210X.13749)

669   60.   Schlager S. 2017 Morpho and Rvcg – Shape Analysis in R. In *Statistical Shape and
670   Deformation Analysis*, pp. 217–256. Elsevier. (doi:10.1016/B978-0-12-810493-
671   4.00011-0)

672   61.   Revell LJ. 2012 Phytools: an R package for phylogenetic comparative biology (and
673   other things). *Methods Ecol. Evol.* **3**, 217–223. (doi:10.1111/j.2041-
674   210X.2011.00169.x)

675   62.   Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, Di
676   Febbraro M, Raia P. 2018 A new method for testing evolutionary rate variation and
677   shifts in phenotypic evolution. *Methods Ecol. Evol.* **9**, 974–983. (doi:10.1111/2041-
678   210X.12954)

679   63.   Biewener AA. 1982 Bone Strength in Small Mammals and Bipedal Birds: Do Safety
680   Factors Change With Body Size? *Journal of Experimental Biology* **98**, 289–301.
681   (doi:10.1242/jeb.98.1.289)

682   64.   Tobalske BW, Altshuler DL, Powers DR. 2004 Take-off mechanics in hummingbirds
683   (Trochilidae). *Journal of Experimental Biology* **207**, 1345–1352.
684   (doi:10.1242/jeb.00889)

685   65.   Biewener AA, Dial KP. 1995 In vivo strain in the humerus of pigeons (*Columba livia*)
686   during flight. *J. Morphol.* **225**, 61–75. (doi:10.1002/jmor.1052250106)

687   66.   Feinsinger P, Chaplint SB. 1975 On the relationship between wing disc loading and
688   foraging strategy in hummingbirds. *Amer. Natur.* **109**.

689   67.   Bright JA, Marugán-Lobón J, Cobb SN, Rayfield EJ. 2016 The shapes of bird beaks
690   are highly controlled by nondietary factors. *Proceedings of the National Academy of
691   Sciences* **113**, 5352–5357. (doi:10.1073/pnas.1602683113)

68.   Shatkovska O V., Ghazali M. 2020 Integration of skeletal traits in some passerines:
impact (or the lack thereof) of body mass, phylogeny, diet and habitat. *J. Anat.* **236**,
274–287. (doi:10.1111/joa.13095)

69.   Zusi RL. 2013 Introduction to the Skeleton of Hummingbirds (Aves: Apodiformes,
Trochilidae) in Functional and Phylogenetic Contexts. *Ornithol. Monogr.* **77**, 1–94.
(doi:10.1525/om.2013.77.1.1)

70.   Drury JP, Clavel J, Tobias JA, Rolland J, Sheard C, Morlon H. 2024 Limited ecological
opportunity influences the tempo of morphological evolution in birds. *Current Biology*
34, 661-669.e4. (doi:10.1016/j.cub.2023.12.055)

71.   Goswami A, Randau M, Polly PD, Weisbecker V, Bennett CV, Hautier L, Sánchez-
Villagra MR. 2016 Do Developmental Constraints and High Integration Limit the
Evolution of the Marsupial Oral Apparatus? *Integr. Comp. Biol.* **56**, 404–415.
(doi:10.1093/icb/icw039)

72.   Randau M, Goswami A. 2017 Unravelling intravertebral integration, modularity and
disparity in Felidae (Mammalia). *Evol. Dev.* **19**, 85–95. (doi:10.1111/ede.12218)

73.   Fabre AC *et al.* 2020 Metamorphosis shapes cranial diversity and rate of evolution in
salamanders. *Nat. Ecol. Evol.* **4**, 1129–1140. (doi:10.1038/s41559-020-1225-3)

Tables

Table 1. Humerus size is the primary predictor of humerus shape variation, with

secondary effects of foraging strategy and migration. Results of multivariate PGLS models

testing ecological and morphological predictors of male humerus shape across hummingbird

species. For each predictor, degrees of freedom (Df), partial R^2, F- and Z-statistics, and p-

values are shown. Phylogenetic signal was strong, with maximum-likelihood $\lambda \approx 1$.

Significance codes: *** $p \leq 0.001$, * $p \leq 0.05$, . $p \leq 0.1$.

Predictor	Df	R^2	F	Z	p-value	Significance
Humerus size	1	0.0982	9.3602	3.2617	0.001	***
Body mass	1	0.0165	1.5747	1.5033	0.07	.
Foraging strategy	2	0.0344	1.641	1.9796	0.025	*
Migration	3	0.0447	1.4188	1.6304	0.058	.
Habitat	2	0.0182	0.8662	0.227	0.394	
Range	1	0.0097	0.9285	0.3686	0.369	
Latitude	1	0.014	1.3299	1.1993	0.112	
Elevation	1	0.0148	1.408	1.3298	0.095	.
Estimated lambda (ML)	~ 1					

Table 2. Evolutionary rates of humerus shape do not differ significantly among

ecological or morphological predictors. Group-wise rate variance analyses testing for

differences in evolutionary rates of male humerus shape among categorical ecological

predictors, and univariate PGLS models evaluating associations between evolutionary rates

and continuous predictors. No predictor exhibited a statistically significant effect on humerus

shape evolutionary rates.

Group-wise rate variance analyses for categorical predictors			
Predictor	Observed Rate Ratio	Effect size	p-value
Foraging strategy	2.5141	1.135	0.138
Habitat	2.0279	-0.0196	0.503
Migration	1.6409	-1.6145	0.953
Univariate PGLS values			
Predictor	Estimate	Std. Error	Lambda (ML)	p-value
Body mass	0.0078	0.0092	0.0579	0.3996
Humerus size	-1.1709	1.5639	0.0088	0.4563
Range	-0.0026	0.0017	0.0327	0.1349
Elevation	0.000002	0.000003	0.0276	0.5455
Latitude	-0.0001	0.0003	0.0217	0.8738

 Figures

Figure 1. Hummingbird humeral shape shows strong phylogenetic structure and

evolutionary rate heterogeneity across major clades. (a) Right humerus illustrating the 25

anatomical landmarks used to quantify humerus shape, averaged across male specimens and

scaled to mean size. (b) Procrustes-aligned landmark dispersion across specimens, illustrating

overall humeral shape variation within the hummingbird family. (c) Thin-plate spline

deformation grids for PC1 and PC2, showing the major axes of humerus shape variation

captured by our morphometric analysis. (d) Phylogenetic distribution of humerus shape

evolutionary rates across the 78 hummingbird species analyzed; branch colors indicate

estimated evolutionary rates, and tip symbols denote foraging strategy. Insets show the null

distribution of phylogenetic signal with the observed value indicated by an arrow, and the

distribution of log-transformed mean evolutionary rates.

Figure 2. Humerus shape variation depends on humerus size and foraging strategy in

hummingbirds. Multivariate relationships between male humerus shape (PC1) and

ecological and morphological predictors across 78 hummingbird species. Boxplots show

group-level differences for categorical predictors. Scatterplots for continuous predictors show

the estimated regression fit. Humerus shape varies with humerus size and foraging strategy,

whereas other predictors show weak or non-significant associations. Silhouettes illustrate

shape change along the PC1 axis. Body mass was analyzed as the natural logarithm of mass in

grams, and geographic range size as the natural logarithm of area in km². Humerus size is

derived from landmark coordinates, in relative (coordinate) units.

Figure 3. Hummingbird foraging strategies differ in humeral morphospace. Foraging

strategies differ in their occupation of humeral morphospace, with territorial, traplining, and

opportunistic species forming clusters in both unsupervised (PCA) and supervised (CVA)

shape spaces. Each point represents a species (n = 78), colored by foraging strategy. (a) PCA

of Procrustes-aligned humeral shapes, with PC1 and PC2 explaining 17.97% and 16.39% of

variation, respectively. Shaded convex hulls denote group morphospaces, and thin black lines

represent phylogenetic relationships projected into shape space. (b) Mean humeral shape

variation across specimens visualized as a thin-plate spline deformation of the consensus

configuration; red points denote landmarks. (c) Canonical Variate Analysis maximizing group

separation based on foraging strategy. Dashed ellipses show 90% confidence regions, and

purple diamonds mark group centroids. Shape outlines adjacent to the axes illustrate humeral

changes associated with CV1 and CV2.

Figure 4. Evolutionary rates of hummingbird humerus shape show little association with

ecological or morphological predictors. Relationships between estimated humerus shape

evolutionary rates (log-transformed) and ecological and morphological predictors across 78

hummingbird species. Boxplots summarize group-level comparisons for categorical

predictors, and scatterplots with fitted regression lines illustrate associations with continuous

predictors. We do not detect associations between humerus shape evolutionary rates and any

predictors. Body mass was analyzed as the natural logarithm of mass in grams, and

geographic range size as the natural logarithm of area in km². Humerus size is derived from

landmark coordinates, in relative (coordinate) units.

Figure 5. Proximal and distal regions of the hummingbird humerus are integrated,

supporting a functional linkage rather than modular independence. The proximal and

distal regions of the humerus exhibit morphological integration, indicating coordinated

evolutionary changes rather than independent modular evolution. (a) Relationship between

PLS1 scores of the proximal region and PLS2 scores of the distal region (r-PLS = 0.899, Z =

4.2219, p = 0.001). Each point represents a species (n = 78), and the shaded region shows the

95% confidence interval for this relationship. (b) The *a priori* modularity hypothesis (upper

panel), representing the expected division of the humerus into proximal and distal functional

units, contrasted with the data-driven minimum-covariation modular structure (lower panel).

Nodes represent landmarks and edges represent the presence of covariation links among

landmarks.

791