Foraging behavior and ecology drive the evolution of humeral shape in hummingbirds

Juan Camilo Ríos-Orjuela[1,2,*], Carlos Daniel Cadena[1], Alejandro Rico-Guevara[3,4], Ilias

Berberi[5], Lauren Miner[5] & Roslyn Dakin[5]

[1]Laboratorio de Biología Evolutiva de Vertebrados, Departamento de Ciencias Biológicas,

Universidad de los Andes, Bogotá, Colombia

[2]Grupo de Morfología y Ecología Evolutiva, Instituto de Ciencias Naturales, Universidad

Nacional de Colombia, Sede Bogotá, Colombia

[3]Department of Biology, University of Washington, Seattle, WA, U.S.A.

[4]Burke Museum of Natural History and Culture, University of Washington, Seattle, WA,

U.S.A.

[5]Department of Biology, Carleton University, Ottawa, Ontario, Canada

* Corresponding author (jcriosor@gmail.com)

Orcid: JCR: 0000-0001-6976-9131, DC: 0000-0003-4530-2478, ARG: 0000-0003-4067-

5312, IB: 0000-0003-3051-8145, LM: 0009-0008-1991-7472, RD: 0000-0002-3140-3975.

Author contributions: JCR: Conceptualization, Methodology, Software, Formal analysis,

Investigation, Data curation, Writing – Original draft, Writing – Review & Editing,

Visualization. **RD:** Conceptualization, Validation, Investigation, Resources, Writing –

Original draft, Writing – Review & Editing, Supervision, Funding acquisition. **CDC:**

Conceptualization, Writing – Review & Editing, Supervision, Funding acquisition. **IB:**

Investigation, Data curation, Writing – Review & Editing. **LM:** Investigation, Data curation.

ARG: Conceptualization, Investigation, Writing – Review & Editing, Supervision.

Funding: This research was supported by an NSERC Discovery Grant (Canada), the Faculty

of Sciences at Universidad de los Andes (Bogotá, Colombia), and the Emerging Leaders in

the Americas Program (Global Affairs Canada).

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgments: We thank Steven Cardiff and Nick Mason for their support during

fieldwork and specimen access, as well as the Louisiana State University Museum of Natural

Science for facilitating access to their ornithological collections.

Abstract

Skeletal morphology underlies avian flight performance and ecological diversification. In

hummingbirds, the humerus is central to power generation and maneuverability, yet its

macroevolutionary dynamics remain poorly understood. Here, we integrate geometric

morphometrics, ecological data, and phylogenetic comparative methods to examine humeral

shape evolution across 80 species. Humeral shape varied significantly with body mass,

centroid size, foraging strategy, geographic range size, and latitude, reflecting the joint

influence of scaling, spatial ecology, and behavioral demands. Traplining, territorial, and

opportunistic species occupied partially distinct regions of morphospace, and foraging

strategy exerted a significant effect on evolutionary tempo, with trapliners evolving humeral

morphology at consistently faster rates. Continuous ecological predictors, however, did not

explain variation in evolutionary rates. Phylogenetic partial least squares revealed strong

evolutionary integration between proximal and distal humeral regions, indicating that

morphological changes are tightly coordinated across the bone. Together, these results show

that humeral evolution in hummingbirds is shaped by the interaction of ecological

differentiation, biomechanical constraints, and strong internal integration, illustrating how

functionally complex structures diversify under multiple selective pressures.

Keywords: Adaptive evolution, Allometry, Biomechanics, Bird flight, Geometric

morphometrics, Phenotypic modularity and integration.

Teaser text

How do tiny hummingbirds balance agility, power, and efficiency in flight? By analyzing humeral shape across 80 species, we uncover how body size, foraging behavior, and geographic distribution shape this key wing bone. Traplining and territorial species exhibit partially distinct humeral morphologies consistent with their contrasting flight demands, and trapliners evolve humeral form at significantly faster rates. Strong evolutionary integration between proximal and distal regions shows that changes in one part of the bone propagate through the entire structure. Our findings reveal how ecology, biomechanics, and evolutionary history interact to shape the hummingbird skeleton.

Introduction

Understanding the evolution of morphological diversity remains a central topic in evolutionary biology. Organisms diversify morphologically in response to ecological pressures, functional constraints, and historical contingencies, and disentangling these influences requires integrative approaches combining morphology, ecology, and phylogeny [1–3]. Birds, among the most ecologically and morphologically varied vertebrates, offer an insightful system to explore these dynamics of trait diversification, especially in traits shaped by the demands of powered flight [4,5].

Among skeletal components, the avian humerus plays a pivotal role in flight mechanics and ecological adaptation. It connects the axial skeleton to the forelimb and anchors major flight muscles responsible for the downstroke and upstroke of the wingbeat [6]. In addition, humerus morphology is understood to influence wing kinematics, which in turn likely affect the magnitude and direction of aerodynamic forces and aspects of maneuverability [7]. Despite its biomechanical importance, the humerus remains underexplored in comparative studies of avian flight, which have often emphasized external features such as wing aspect ratio [8]. While such metrics effectively capture overall aerodynamic properties, they do not reflect underlying skeletal variation, potentially overlooking functionally significant musculoskeletal morphology.

In hummingbirds, the humerus is distinctively short, robust, and exhibits morphological adaptations that support their specialized hovering and maneuverable flight [7]. Unlike the long, slender humeri typical of birds adapted for gliding or forward-powered flight [9], the hummingbird humerus is shortened to perform exceptionally high wingbeat frequencies and rapid rotation of the wing on each upstroke [10]. These changes allow hummingbirds to generate lift during both downstroke and upstroke and exert fine control during rapid aerial

maneuvers [7]. To meet these challenges, the humerus functions as a rigid, force-transmitting

lever for enlarged flight muscles, and exhibits structural specializations such as increased

cortical thickness, a relatively straight shaft, and pronounced muscle attachment sites at the

deltopectoral crest. This morphology reflects a biomechanical compromise: a shorter, sturdier

humerus reduces inertial resistance and facilitates maneuverability, but may constrain aspects

of aerodynamic efficiency (the lift-to-drag (L/D) ratio) or range of motion, requiring precise

tuning to maintain energetic economy during sustained flight. With over 300 species, the

hummingbird clade presents a rich system for comparative study, offering insight into how

skeletal structures evolve under varying ecological demands and locomotor constraints. Yet

despite its biomechanical importance, the hummingbird humerus remains underexplored in

evolutionary studies.

Hummingbirds exhibit diverse foraging strategies that reflect their ecological niches and

influence their flight behavior. Some species defend floral territories aggressively, engaging in

rapid, maneuver-demanding flights to repel intruders, while others adopt traplining behaviors,

following predictable expansive foraging circuits that require sustained travel across larger

areas and efficient navigation between patchy floral resources [11,12]. These differing

foraging strategies are expected to impose distinct aerodynamic and biomechanical demands,

which may drive morphological divergence in the skeleton. Species engaging in frequent,

high-intensity maneuvers—such as territorial defense—may benefit from shorter, more robust

humeri that facilitate rapid changes in direction and powerful wing strokes [7], In contrast,

species relying on traplining may favor longer and more slender humeri that enhance flight

efficiency and reduce energetic costs over longer distances, even if they still require precise

control for floral approach and handling [13,14].

Beyond foraging behavior, ecological factors such as those varying with elevation also shape

hummingbird morphology. Species occupying high-altitude environments face hypoxia and

reduced air density, which increase the biomechanical demands of flight [15,16]. Prior studies have documented phenotypic divergence along elevational gradients, particularly in wing shape, body size, and metabolic traits [17,18]. However, skeletal elements like the humerus remain understudied in this context, despite their critical role in flight mechanics.

Additional ecological and morphological factors may further influence humeral evolution. Species occurring at higher latitudes or occupying broader geographic ranges often encounter greater environmental heterogeneity and dispersal demands, potentially favoring either morphological flexibility or specialization [19,20]. Body mass also imposes biomechanical constraints, as larger species must support greater gravitational and aerodynamic forces [21]. Finally, overall skeletal size may influence scaling relationships critical for wing loading and flight efficiency [22].

Together, these patterns suggest that humeral morphology in hummingbirds is likely shaped by a complex interplay of biomechanical constraints, ecological demands, and evolutionary history. Yet, the extent to which ecological variables predict morphological divergence—and whether they are also associated with evolutionary rates—remains underexplored [23,24]. Evaluating evolutionary rates is important, as accelerated shape evolution may reflect ecological opportunity, functional innovation, or release from constraints [25,26]. In hummingbirds, one may expect that body size, ecological and geographical factors could all influence not only humeral shape but also the tempo of its evolution. By testing for associations among such variables, one may clarify how ecological and morphological factors jointly shape both the patterns and pace of morphological change in this key flight structure.

The hummingbird humerus, with its distinct proximal and distal functions, also provides an appropriate system to investigate phenotypic integration. The proximal region primarily serves as the attachment site for the pectoralis and supracoracoideus muscles, which are

responsible for generating the downstroke and upstroke during flight, respectively [6,27]. The

distal portion of the humerus articulates with the radius and ulna, transmitting the forces

generated by these muscles into wing motion and modulating the wing's orientation and

amplitude. Understanding whether these regions evolve in a coordinated manner offers

insights into developmental constraints and adaptive flexibility [28–30]. Morphological traits,

especially those underpinning locomotion, rarely evolve independently [31]. Instead, they

exhibit integration and modularity due to shared developmental pathways and functional

covariation [32,33]. While many studies of phenotypic integration focus on patterns across

multiple bones or skeletal regions (e.g., skull and mandible, or entire limbs), exploring

integration within a single element can reveal how distinct functional regions remain

evolutionarily coordinated despite localized biomechanical specializations.

Here, we investigate how foraging strategy, ecological pressures, and evolutionary processes

collectively shape humeral morphology across a broad sample hummingbird species. First, we

hypothesized that species differing in foraging strategies would exhibit divergent humeral

morphologies, reflecting adaptations to their specific flight demands and ecological contexts.

We also explored whether these strategies are linked to variation in the evolutionary rates of

humeral shape. Rather than assuming a single directional outcome, we framed this analysis as

a hypothesis-generating approach informed by the contrasting demands of different foraging

behaviors. For example, traplining—characterized by repeated, sustained flight along spatially

extensive routes—may select for morphologies that enhance energetic efficiency and

maneuverability across heterogeneous environments. Territorial behavior, by contrast,

involves frequent short-distance accelerations and complex aerial contests, potentially

selecting for structural robustness and high maneuverability. Whether such behaviors promote

morphological innovation or constrain it through stabilizing selection likely depends on

evolutionary history, ecological variability, and biomechanical trade-offs. Given that

evolutionary rates can reflect diverse processes, including directional selection, stabilizing

selection, or release from constraint [34], we interpret our rate comparisons within a

framework that considers multiple evolutionary scenarios, rather than presupposing a single

outcome.

Second, we predicted that humeral shape variation and evolutionary rates would be influenced

by a combination of ecological and morphological factors. Specifically, species inhabiting

higher elevations may exhibit distinctive humeral morphologies, as high-altitude

environments impose greater aerodynamic and metabolic constraints that may drive adaptive

changes. However, whether these conditions lead to accelerated evolutionary rates remains

uncertain. In some cases, strong stabilizing selection or habitat filtering may constrain

morphological evolution, resulting in low rates despite ecological pressure [34]. Similarly,

while broad geographic ranges or occurrence at higher latitudes have been associated with

increased environmental heterogeneity and dispersal challenges, the relationship with

evolutionary rate is context-dependent and may vary across clades and traits. Regarding body

mass, we predicted that larger-bodied species would exhibit corresponding shifts in humeral

morphology, reflecting biomechanical scaling effects on flight performance and energetic

costs. Although allometric relationships are expected, examining whether these scaling

patterns vary across ecological contexts could provide insights into how size-related

constraints interact with ecological specialization. Finally, we expected overall skeletal size to

correlate with humeral shape variation due to its impact on wing loading and flight efficiency.

Lastly, we hypothesized that the humerus exhibits strong internal phenotypic integration, with

coordinated morphological evolution between its proximal and distal regions, given their

shared role in force transmission and wing mechanics. Such integration is expected because

modifications in one region (e.g., altering muscle attachment sites proximally) would

necessitate compensatory changes distally to maintain flight performance. However, examples

from other systems suggest that bones can evolve in a modular fashion when regions fulfill

distinct functions or face divergent selective pressures [31,35,36]. Whether such modularity

also occurs within the hummingbird humerus remains an open question but would imply that

evolutionary changes could be targeted to specific regions without affecting the entire

structure. By integrating ecological categories and morphometric analyses within a

phylogenetic framework, our objective is to provide a comprehensive assessment of the

factors driving morphological evolution in the hummingbird humerus.

Methods

Specimens and image acquisition

We analyzed 80 hummingbird species, representing approximately 22% of extant species

diversity in the family Trochilidae [37]. To capture morphological variation of the humerus,

we photographed the right humerus in cranial view from male specimens in the collection of

the Louisiana State University Museum of Natural Science (LSUMNS). We focused on males

to reduce potential shape variation due to sexual dimorphism and to ensure consistency across

species, as male specimens are more frequently available and taxonomically reliable in

museum collections. Images were taken with a Canon EOS M50 (6000 × 4000 px) using a 28

mm macro lens. Because specimens were photographed in strict orthogonal orientation near

the center of the frame, geometric distortion at this focal length is minimal, and no additional

distortion correction was applied. The camera was mounted on a copy stand at a fixed

distance from the specimens to maintain consistent magnification and perspective. Each

photograph included an embedded scale bar (10 mm with 0.1 mm divisions) for accurate

scaling in subsequent analyses. This standardized view facilitated consistent comparative

analysis of skeletal structure across species [38].

A total of 25 anatomical landmarks were digitized on each humerus, focusing primarily on

points along the contour to capture overall shape (Figure 1A-B). Landmarks were chosen

based on biomechanical relevance and anatomical consistency across taxa [22,39]. No

semilandmarks were employed, because we aimed to focus on homologous and discrete

anatomical points rather than continuous curves. A detailed description of landmarks and

specimens list is provided in Table S1. Landmark digitization was conducted using TPSDig2

version 2.32 [40].

Given the rarity of skeletal material for many hummingbird species, sample sizes varied

across taxa. For approximately 87% of species (70 of 80), only a single male specimen was

available. For the remaining 10 species, where multiple specimens (two to four per species)

were available, we calculated the species-specific mean shape by averaging landmark

configurations across individuals [41].

To assess the potential influence of geometric uncertainty on the morphometric dataset, we

considered several non-statistical sources of measurement error. Pixel resolution of the Canon

EOS M50 (6000 × 4000 px) introduces at most 1–2 pixels of uncertainty in landmark

position, corresponding to ~0.1% of humeral length in our images. Intra-observer variation in

landmark placement in similar studies is typically 0.5–1.5%, and because all specimens were

digitized by the same observer, inter-observer error is likely below 1%. Anatomical ambiguity

in a subset of landmarks may contribute an additional 1–3% variability in local point

placement. Together, these factors suggest an overall geometric uncertainty on the order of 2–

4% of inter-landmark distances, well below the magnitude of interspecific shape differences

captured by the Procrustes analysis. Consequently, these sources of error are unlikely to

influence the main axes of humeral shape variation or the phylogenetic patterns recovered in

our analyses.

Morphometric data preparation

All data preparation and statistical analyses were conducted in R version 4.4.2 [42].

Landmark data were aligned using a Generalized Procrustes Analysis (GPA) to remove non-

shape variation related to position, orientation, and scale [43]. GPA was performed using the

gpagen() function in the *geomorph* package [44]. Following alignment, we calculated squared

Euclidean distances from each specimen to the mean shape configuration to identify and

exclude outliers. We retained 97.5% of specimens closest to the mean shape, a typical

threshold in morphometric studies. Euclidean distances were computed by converting the

aligned 3D landmark array to a two-dimensional matrix, followed by row-wise distance

calculations. Two specimens—*Calypte anna* and *Goethalsia bella*—were identified as outliers

for their unusually high morphological deviation from the overall mean shape configuration,

due to preservation or landmarking artifacts, and were thus removed from further

investigation. Following this filtering, 78 species were retained for downstream analyses. A

PCA analysis of the aligned coordinates was performed exclusively for graphical

representation of the shape variation and morphospace visualization (Figure 1C).

Species traits

For each species, we recorded a suite of ecological and morphological traits. Foraging

strategy was classified as territorial, trapliner, or opportunist, based on typical foraging

behavior and resource use [45]. We also recorded a species' maximum elevation as a proxy

for elevational environment [46]. Geographic range size (km²) was obtained as we expect

species occupying larger ranges to experience broader environmental heterogeneity. Body

mass was included as a proxy for biomechanical constraints associated with size, given that

larger species typically experience different mechanical stresses during flight [11]. Range

size, latitude at the centroid of the species' range, and body mass data were retrieved from

Tobias et al. (2022). Additionally, as a structure-specific indicator of skeletal size, we

measured humerus centroid size from the landmark data. While centroid size is

mathematically derived from the distribution of landmark coordinates, it is widely recognized

as a robust proxy for overall size in morphometric analyses [22].

Phylogenetic signal

To assess whether humeral shape variation exhibited phylogenetic signal, we quantified

Blomberg's multivariate K statistic [30,48] using the physignal() function in *geomorph*, based

on 1,000 permutations. We also computed a Z-transformed effect size to contextualize the

observed K values relative to a null distribution. An ultrametric phylogeny was obtained from

McGuire et al. (2014) for all comparative analyses.

Phylogenetic modeling of humeral shape variation

To evaluate how ecological and morphological predictors influence humeral shape, we

implemented a phylogenetic generalized least squares (PGLS) approach using the

procD.pgls() function in *geomorph* for high-dimensional shape data. Because distance-based

methods do not estimate Pagel's λ by maximum likelihood [50], we fixed λ at 0.99, a value

consistent with the strong phylogenetic signal detected in humeral shape (K = 0.309, p =

0.001). Significance testing was conducted using 1,000 permutations of residuals under a

Brownian correlation structure.

We evaluated a set of *a priori* candidate ecological models (Table S2) and then fitted a full

multivariate model including foraging strategy (categorical), body mass, humerus centroid

size, maximum elevation, geographic range size, and absolute centroid latitude. None of the

predictors showed problematic collinearity (Pearson's |r| < 0.36), and all variance inflation

factors remained low (VIF < 1.5), indicating that the model structure was appropriate for

multivariate estimation.

Because procD.pgls uses Type-I sequential sums of squares, we implemented a principled

ordering of predictors to maximize model stability and ensure biological interpretability. First,

we performed an exhaustive permutation analysis of all 720 possible predictor sequences.

This analysis consistently ranked centroid size as the strongest predictor, followed by body

mass, with environmental predictors emerging as the weakest contributors. Based on this

result, and following best practices for sequential sum-of-squares models, we ordered

predictors as follows: (1) centroid size, (2) body mass, (3) foraging strategy, (4) range size, (5)

absolute latitude, and (6) maximum elevation. This ordering minimizes instability due to

predictor order and reflects both biological expectations and empirical model performance.

The final PGLS model was then used to generate multivariate effect estimates, partial

residuals, and projected PC1 relationships shown for visualization. Although PC1 is used for

illustrative purposes, all hypothesis tests were based on the full multivariate shape data.

Discriminant analysis of humeral shape by foraging strategy

To evaluate whether humeral shape differs among foraging strategies, we performed a

Canonical Variates Analysis (CVA) using size-corrected Procrustes coordinates. Allometric

effects were removed by regressing aligned shape coordinates on log-transformed centroid

size, and the resulting residuals were used as inputs for all subsequent analyses [22]. CVA was

conducted using the CVA() function in the *Morpho* package [51], with foraging strategy

specified as the grouping factor. Statistical support for group separation was assessed via

1,000 permutation tests, which evaluate whether between-group differences exceed those

expected under random assignment of specimens to groups. To visualize phylogenetic

structure in the discriminant space, we projected species relationships onto the CVA axes by

connecting species scores according to the tree topology using *phytools* [52]. This allows

assessment of whether phylogenetic relatedness aligns with group clustering in canonical

morphospace. All analyses were conducted on the same dataset and grouping scheme used in

the main phylogenetic regression models to ensure consistency across analytical frameworks.

Evolutionary rates of humeral shape

To test whether foraging strategies differed in their evolutionary tempo, we used the

compare.evol.rates() function, specifying foraging strategy as the grouping factor and

performing 999 permutations to assess statistical significance under a null model of equal

rates. This analysis evaluates whether the mean evolutionary rate of each group departs from

expectations given the phylogeny and the distribution of shape change across species.

For continuous predictors, we tested associations between evolutionary rates and ecological or

morphological variables using the search.trend() function in RRphylo [53], which fits linear

regressions between branch-specific rate estimates and predictor values while accounting for

phylogeny. Because branch-specific rate estimates can be influenced by extreme values in

individual species, we applied standard RRphylo diagnostics to identify potential outliers or

leverage points using the built-in functions RRphylo::outliers() and RRphylo::search.trend().

No species were removed, but diagnostics are reported in Table S3. To evaluate time-

dependent patterns in evolutionary tempo, we extracted ancestral rate estimates from RRphylo

and regressed them against node ages, allowing us to assess whether rate magnitudes varied

systematically across the temporal depth of the phylogeny.

We additionally modeled species-specific evolutionary rates as a univariate response using

phylogenetic generalized least squares (PGLS). These models were fitted using the gls()

function in the *nlme* package, with a Brownian-motion correlation structure (corBrownian)

derived from the same ultrametric tree used in the RRphylo analyses. Each ecological or

morphological predictor (body mass, centroid size, geographic range size, absolute latitude,

and maximum elevation) was evaluated in a separate univariate PGLS model. Predictor

variables were log-transformed when appropriate to improve linearity and homoscedasticity.

This complementary framework allowed us to test whether continuous predictors showed

phylogenetically structured associations with evolutionary tempo that were consistent with the

patterns suggested by RRphylo's rate estimates.

Phenotypic modularity and integration

To evaluate the degree of phenotypic integration within the humerus, we partitioned the

Procrustes-aligned landmark data into two functional modules defined a priori based on avian

flight anatomy. The proximal module included the humeral head, tubercles, and associated

articular surfaces involved in shoulder rotation and load transfer during both hovering and

flapping. The distal–shaft module comprised the diaphysis, trochlea, capitulum, and distal

muscle insertion sites, which primarily mediate force transmission, wingbeat mechanics, and

resistance to bending loads during forward flight. This subdivision reflects the well-

established biomechanical differentiation between the articulation region of the humerus and

its load-bearing shaft [27,54] and was selected to test a specific functional hypothesis rather

than to infer empirical modular structure.

We quantified evolutionary covariation between these two modules using phylogenetic partial

least squares (pPLS) implemented in the phylo.integration() function of *geomorph*. This

method estimates the correlation between paired blocks of shape variables while explicitly

accounting for phylogenetic non-independence, thereby capturing integration patterns that

reflect coordinated evolutionary changes between modules. Statistical significance was

assessed using 1,000 phylogenetically informed permutations. We did not perform alternative

modularity searches (e.g., minimum covariation partitions or EMMLi-based module

optimization), because our goal was to test a focused, anatomically motivated prediction

regarding integration between proximal articulation structures and the distal shaft, rather than

to explore the full modular landscape of the humerus.

Results

Phylogenetic signal

Humeral shape exhibited a significant phylogenetic signal (K = 0.309, p = 0.001), indicating

that closely related hummingbird species tend to share similar humeral morphologies. This

result is further supported by a large effect size (Z = 41.1), underscoring the strength of

phylogenetic structuring in shaping bone morphology. The geometric configuration used to

quantify humeral shape reveals consistent variation in landmark positions across species,

highlighting major axes of morphological differentiation (Figure 1A–C). The distribution of

phylogenetic signal and its placement on the hummingbird phylogeny are shown in Figure

1D.

Predictors of humeral shape variation

The multivariate PGLS identified several significant predictors of humeral shape variation

(Table 1; Figure 2). Humerus size showed the strongest effect (R^2 = 0.081, Z = 3.08, p =

0.002), consistent with strong allometric contributions to humeral morphology. Range size

also emerged as a significant predictor (R^2 = 0.031, Z = 2.52, p = 0.008), indicating that

geographically widespread species tend to occupy distinct regions of humeral morphospace.

Body mass showed a moderate but significant association with shape (R^2 = 0.022, Z = 1.89, p

= 0.031), suggesting that mass-related biomechanical demands influence humeral

configuration. Importantly, foraging strategy also had a significant effect on humeral shape

(R^2 = 0.035, Z = 1.87, p = 0.035), reflecting detectable morphological differentiation among

territorial, traplining, and opportunistic species. Absolute latitude exhibited only a marginal

trend ($R^2 = 0.016$, $Z = 1.46$, $p = 0.070$), whereas maximum elevation showed no significant

influence on variation in shape ($R^2 = 0.013$, $Z = 1.25$, $p = 0.102$). Overall, the final model

explained approximately 20% of the total multivariate variation in humeral shape.

Morphospace differentiation by foraging strategy

The principal component analysis (PCA), used to visualize patterns of shape variation,

revealed differences in humeral morphology associated with foraging strategies (Figure 3A).

Notably, the morphospaces of traplining and territorial species were largely contained within

the broader morphospace occupied by opportunistic species, as illustrated by the convex hulls

outlining each group in Figure 3A. This suggests that opportunistic foragers may occupy a

broader and more variable humeral morphospace, although the overall overlap among groups

was substantial. The projected phylogenetic relationships onto the morphospace showed some

local clustering of related species, consistent with the significant phylogenetic signal detected

in the shape data. However, the overall pattern lacked clear structure, with multiple lineages

dispersed throughout the morphospace, indicating that a substantial proportion of shape

variation is explained by ecological differentiation rather than shared ancestry. The Canonical

Variate Analysis (CVA) further reinforced the role of foraging strategy in shaping humeral

morphology (Figure 3C). The first canonical axis (CV1) primarily distinguished trapliners

from both territorial and opportunistic species, while CV2 separated territorial from

opportunistic species. Cross-validated classification accuracy was low (43.6%), only modestly

exceeding random expectations, and the canonical morphospace showed broad overlap among

strategies. Traplining species displayed slightly higher correct assignment (47.6%) than

territorial or opportunistic species (40.6–44.0%), but all groups exhibited extensive

intermingling. These modest patterns of group separation are consistent with the significant

but relatively small effect of foraging strategy detected in the multivariate regression models.

Evolutionary rates

Estimates of evolutionary rates derived from RRphylo revealed a strongly right-skewed distribution, with most hummingbird species exhibiting very low rates of humeral shape change. Across the 78 species analyzed, median rate was 0.0063, with the 25th and 75th percentiles at 0.0030 and 0.0120, respectively. Only a small subset of species displayed markedly elevated rates (Table S3). These high-rate branches represent isolated episodes of accelerated morphological divergence, whereas most species retained relatively conserved humeral configurations. Phylogenetic signal in evolutionary rates was weak ($\lambda = 0.027$) and not significantly different from zero (LR = 0.041, p = 0.840), indicating that closely related species did not evolve at more similar rates than expected under phylogenetic independence. This demonstrates that rate variation is not phylogenetically structured and arises idiosyncratically across the tree.

Group-wise comparisons using compare.evol.rates() detected a statistically strong effect of foraging strategy on evolutionary tempo ($R^2 = 0.062$, Z = 3.465, p = 0.001). Traplining species exhibited the highest mean evolutionary rates (approximately 2.5 times faster than territorial species) followed by opportunistic taxa (Figure 4). When examining temporal patterns, rates across all strategies declined through time, consistent with an early burst followed by a deceleration of humeral shape evolution (Figure 4, upper left panel).

To assess whether continuous ecological or morphological traits predicted evolutionary rates, we fitted univariate PGLS models using a Brownian correlation structure. None of the predictors showed statistically significant associations with rate estimates: latitude (p = 0.132), body mass (p = 0.476), humerus centroid size (p = 0.925), geographic range size (p = 0.678), and maximum elevation (p = 0.890) all produced weak and non-significant effects (Table 2). Although the predictors themselves exhibited moderate phylogenetic signal ($\lambda =$

0.57–0.63), this structure did not translate into variation in evolutionary tempo. These results

indicate that foraging strategy is the only ecological factor with a detectable association with

evolutionary rates in the humerus, while other variables show no consistent influence under a

phylogenetic regression framework.

Phenotypic integration

The phylogenetic PLS analysis revealed strong phenotypic integration between the proximal

and distal regions of the humerus (r-PLS = 0.899, Z = 4.22, p = 0.001). This indicates that

morphological changes in one region are closely mirrored by changes in the other, reflecting

coordinated evolution across the bone (Figure 5). This high level of integration is consistent

with the functional coupling between articulation surfaces and the load-bearing shaft, where

evolutionary changes in joint morphology necessarily co-vary with adjustments in shaft

curvature, thickness, or distal anatomy.

436   **Discussion**

Phylogenetic signal

Our results reveal a clear phylogenetic signal in humeral morphology among hummingbirds

(Figure 1). This finding indicates that evolutionary history continues to shape patterns of

morphological similarity across species, consistent with general trends reported for other

avian skeletal traits [55,56]. The presence of a strong phylogenetic structure suggests that

humeral shape has evolved in a way that is partly structured by shared ancestry throughout the

radiation of hummingbirds. Rather than indicating strict evolutionary constraint, this pattern

may reflect the retention of biomechanically effective configurations across lineages, as well

as the influence of phylogenetic inertia during adaptive divergence [27,34,57].

Predictors of shape variation

Our multivariate phylogenetic regression identified several significant predictors of species differences in humeral shape, supporting the hypothesis that ecological and morphological factors contribute to skeletal morphology in hummingbirds. Humerus size showed the strongest effect in the model (Table 1), consistent with a marked allometric component in shape variation. As humerus size increased, species exhibited broader proximal epiphyses and straighter, shorter diaphyses. This pattern likely reflects biomechanical adaptations: larger humeri require expanded articular surfaces for muscle attachment (notably for the pectoralis and supracoracoideus) and reduced shaft curvature to minimize bending stress under greater loading [27,58,59]. These shape changes are consistent with general scaling principles observed in avian flight bones, where increased size leads to disproportionate reinforcement of load-bearing regions [27,54].

Body mass also significantly predicted humeral shape. Larger-bodied species displayed more robust humeral shapes, particularly with increased shaft thickness and straighter diaphyses. Biomechanically, this likely compensates for the higher aerodynamic forces and inertial loads encountered during both flapping and hovering flight in heavier hummingbirds [17]. These findings align with prior studies documenting allometric effects on limb bone robustness across volant birds [13].

Geographic range size was another significant ecological predictor. Species with larger geographic ranges tended to exhibit humeri with a more slender mid-shaft, characterized by reduced mediolateral breadth relative to anteroposterior depth (Figure 2). Wider ranges may be associated with longer average movement distances, whether through dispersal, exploratory behavior, or seasonal shifts, which in turn could favor morphologies that enhance flight efficiency and reduce the energetic costs of sustained flight (i.e. longer overall wing (via

elongation of the ulna, manus, or primary feathers), increased wing area, and reduced wing

loading; Claramunt et al., 2012; Drury et al., 2024). Alternatively, occupying a wide range of

habitats may select for versatile skeletal configurations capable of performing effectively

across diverse environmental conditions.

Latitude showed a marginal but notable trend, with species distributed farther from the

equator exhibiting more robust humeri, characterized by expanded proximal epiphyses and

straighter shafts. While this pattern could reflect biomechanical adaptations to greater

environmental seasonality and structurally complex habitats, conditions that may require

enhanced power generation and maneuverability during flight [61,62], it may also relate to

broader-scale movement ecology. This pattern is biomechanically consistent with the

demands of lift production in hummingbirds: a shorter diaphysis reduces the moment of

inertia of the wing, enabling faster angular accelerations, while deeper epiphyses provide

larger areas for the shoulder musculature to generate torque. Species at higher latitudes often

engage in longer-distance dispersal or seasonal movements, which could select for more

mechanically robust skeletal traits that support sustained flight performance in variable

conditions. In contrast, elevation did not show a significant effect, though the trend toward

shorter diaphyses and deeper epiphyses in high-elevation species is biomechanically

consistent with enhanced lift production under low air density and hypoxia [17,63]. Similar

morphological adjustments to elevation have been reported for wing shape and muscle

architecture in other avian taxa [15].

Foraging strategy showed a significant but modest effect on humeral shape (p = 0.035), and

the differences among strategies exhibited overlap in morphospace. Nonetheless, the observed

trends were biomechanically coherent. Traplining species tended to exhibit longer and more

slender humeri, a morphology likely supporting sustained, energy-efficient flight over

extended foraging circuits. A more elongated humerus can increase wingspan and reduce

wingbeat frequency, enhancing aerodynamic efficiency (approximated here as a higher lift-to-

drag ratio) and decreasing energetic costs during long-distance movement between floral

resources. Insofar as a longer proximal wing segment increases the span and reduces induced

drag during forward flight. [13,14,64]. In contrast, territorial species exhibited shorter, thicker,

and more curved humeri, traits associated with enhanced maneuverability and force

production. Shorter bones with greater curvature can reduce the moment of inertia around the

shoulder joint by keeping mass closer to the body's center of rotation, enabling faster angular

acceleration and tighter turning radii. These features are advantageous during rapid chases,

evasive maneuvers, and sudden vertical ascents used in territorial defense [11,12,62].

Additionally, a more robust humerus provides increased surface area for the attachment of

flight muscles such as the pectoralis and supracoracoideus, which are critical for explosive

bursts of power [64]. These divergent morphological configurations reflect adaptive tuning of

the flight apparatus to meet the specific aerodynamic and behavioral demands imposed by

alternative foraging strategies.

Importantly, elongation of the humerus is not the only anatomical pathway for increasing

wing span or improving L/D ratio; elongation of primary feathers could theoretically achieve

similar aerodynamic benefits with lower inertial costs. Our interpretation therefore does not

assume a one-to-one link between humeral length and wing length. Rather, we propose that

changes in the bony proximal wing segment may interact with distal elements (ulna, manus,

feathers) to modify overall wing planform in ways relevant to forward-flight efficiency.

Humeral elongation should therefore be viewed as a plausible, but not exclusive,

morphological route contributing to differences in flight performance among foraging

strategies.

Together, these findings underscore that the humerus in hummingbirds is shaped by a

complex combination of size-related biomechanics, ecological space use, and behavioral

flight demands. The moderate explanatory power of the model (~20% of total variation)

reflects the multifaceted nature of morphological evolution under interacting selective

pressures and phylogenetic constraints.

Morphospace differentiation by foraging strategy

The principal component analysis (PCA) showed that species using territorial and traplining

strategies occupied partially distinct areas of morphospace. In contrast, opportunistic species

encompassed much of the variation found in the other two groups (Figure 3A). This

distribution suggests that opportunistic species occupy a broader and more variable region of

morphospace, although all three strategies show partial overlap.

Building on these findings, the CVA revealed modest differentiation among groups, with

overlap in canonical morphospace despite statistically detectable effects. (Figure 3C). CV1

and CV2 effectively distinguished trapliners, territorial, and opportunistic species, which is

consistent with the significant but modest effect of foraging strategy detected in the

multivariate regression. Moreover, the distribution of species in the phylomorphospace

(Figure 3A) revealed that some distantly related species converge on similar humeral

configurations, suggesting that similar flight demands may drive morphological convergence

across the hummingbird phylogeny despite evolutionary distance. This pattern highlights the

role of ecological function in shaping skeletal form beyond phylogenetic constraints.

The 90% confidence ellipses showed partial overlap, particularly between territorial and

opportunistic species, while trapliners occupied a more distinct region of morphospace. These

patterns likely arise from the differing balance of maneuverability and energetic efficiency

required by each strategy, producing partially convergent solutions rather than discrete

morphotypes: territorial species must defend floral resources through rapid, aggressive flights,

while opportunistic species exploit transient feeding opportunities, potentially favoring more

generalized morphologies [12,65]. Notably, this group-level morphological differentiation

persisted even after accounting for phylogenetic relatedness and allometric scaling, indicating

that foraging strategy imposes a measurable and adaptive influence on humeral shape. These

results corroborate the directional trends identified in the multivariate regression models and

underscore the biomechanical relevance of humeral variation in relation to ecological

demands.

Evolutionary rates

The analysis of evolutionary rates provided a nuanced picture of how ecological factors might

shape the tempo of humeral evolution in hummingbirds. Traplining species exhibited

evolutionary rates approximately 2.5 times higher than those of territorial and opportunistic

species (Figure 4A) and this difference was statistically significant ($R^2 = 0.062$, $Z = 3.465$, $p =$

0.001). Traits associated with traplining, such as sustained flight along extended foraging

circuits, may impose distinct biomechanical and energetic demands that contribute to the

observed rate acceleration [11,12,66], although the precise functional mechanisms warrant

further investigation. These results indicate that foraging behavior is a consistent predictor of

evolutionary tempo in the humerus, with traplining lineages exhibiting reliably faster rates of

morphological change than territorial and opportunistic species.

Continuous ecological and morphological predictors showed no significant association with

evolutionary rate in univariate PGLS models, indicating that variables such as body mass,

elevation, latitude, centroid size, and geographic range size do not systematically predict

evolutionary tempo under a phylogenetic regression framework.

Although evolutionary rates showed a temporal decline consistent with an early burst model

(Figure 4A), species-level estimates mapped onto the phylogeny showed scattered pockets of

elevated rates (Figure 1B), especially near the tips. These patterns are not necessarily

contradictory: temporal trends capture average changes over macroevolutionary time, while

tip-specific estimates reflect recent shifts that may be driven by clade-specific adaptations or

environmental changes. Such dual patterns have been observed in other studies of

morphological evolution [26] and underscore the complexity of interpreting rate heterogeneity

in macroevolutionary datasets, where isolated pulses of accelerated divergence coexist with

long periods of relative stasis.

Phenotypic integration

Our analysis revealed a very high degree of morphological integration between the proximal

and distal regions of the humerus (Figure 5), supporting the prediction that these regions

evolve in a coordinated manner. The strong integration observed indicates that the humerus

behaves as a single functional unit, with evolutionary changes in the proximal region tightly

associated with coordinated adjustments along the shaft and distal end, reflecting the

mechanical continuity required during hovering and forward flight [14].

Such integration is expected in highly loaded locomotor structures, where functional

covariation and shared developmental pathways limit the potential for independent

morphological change [33,67]. While this integration may limit the range of morphological

trajectories available to hummingbirds, it likely facilitates the evolution of coordinated

adaptations to ecological and biomechanical demands. This is consistent with broader patterns

observed in vertebrate skeletal systems, where modularity and integration frequently emerge

as key features of evolutionary flexibility and constraint [68,69]. The strength of this

integration also suggests that evolutionary covariation dominates over any individual-level

plasticity associated with adult bone remodeling.

In hummingbirds, the persistence of a strongly integrated humeral structure across species

suggests that even as humeral shape diverges in response to ecological demands, internal

morphological coordination remains evolutionarily conserved. This high integration likely

constrains the directions along which humeral morphology can evolve, channeling change

along coordinated trajectories rather than allowing independent diversification of proximal

and distal regions. As a result, while ecological pressures may drive diversification in overall

humeral form, such change is shaped by internal structural coherence, constrained by it. Our

findings thus highlight that phenotypic integration can both enable functional refinement and

limit morphological disparity, shaping how form evolves under ecological and biomechanical

selection. Thus, even when ecological pressures promote divergence in overall humeral

morphology, internal structural coherence remains evolutionarily conserved.

603   **Conclusion**

Our results show that humeral evolution in hummingbirds reflects an interplay between

ecological demands, biomechanical constraints, and phylogenetic history. Although humeral

shape displays moderate phylogenetic structuring, evolutionary rates exhibit little

phylogenetic signal, indicating that tempo and mode are decoupled. Shape variation is

associated with foraging behavior, spatial distribution, and body size, and traplining species

evolve humeral morphology at significantly faster rates, consistent with the demands of

sustained circuit-foraging. Geographic range size further predicts differences in humeral

shape, although it does not influence evolutionary tempo.

Strong evolutionary integration between proximal and distal humeral regions highlights the

cohesive functional design of the bone. This integration implies that evolutionary

modifications are channeled along coordinated, biomechanically viable trajectories rather than

occurring independently across regions.

Together, these findings demonstrate that even within a lineage constrained by flight

performance, ecological and behavioral variation leaves detectable signatures on skeletal

design. The combined analysis of morphometrics, ecology, and phylogenetics provides a

clearer understanding of how integrated locomotor structures evolve under multiple selective

pressures.

Acknowledgments

We thank Steven Cardiff and Nick Mason for their support during fieldwork and specimen

access, as well as the Louisiana State University Museum of Natural Science for facilitating

access to their ornithological collections. This research was supported by funding from an

NSERC Discovery Grant (Canada), the Faculty of Sciences at Universidad de los Andes

(Bogotá, Colombia), the Walt Halperin Endowed Professorship and the Washington Research

Foundation as Distinguished Investigator to A.R.-G., and the Emerging Leaders in the

Americas Program (Global Affairs Canada).

References

1.   Sayol F, Reijenga BR, Tobias JA, Pigot AL. 2025 Ecophysical constraints on avian adaptation and diversification. *Current Biology* **35**, 1326-1336.e6. (doi:10.1016/j.cub.2025.02.015)

2.   Pigot AL *et al.* 2020 Macroevolutionary convergence connects morphological form to ecological function in birds. *Nat Ecol Evol* **4**, 230–239. (doi:10.1038/s41559-019-1070-4)

3.   Losos JB. 2011 Convergence, adaptation, and constraint. *Evolution (N Y)* **65**, 1827–1840. (doi:10.1111/j.1558-5646.2011.01289.x)

4.   Dudley R. 2002 Mechanisms and Implications of Animal Flight Maneuverability. *Integr Comp Biol* **42**, 135–140. (doi:10.1093/icb/42.1.135)

5.   Norberg UM. 1990 *Vertebrate Flight*. Berlin, Heidelberg: Springer Berlin Heidelberg. (doi:10.1007/978-3-642-83848-4)

6.   Tobalske BW, Biewener AA. 2008 Contractile properties of the pigeon supracoracoideus during different modes of flight. *Journal of Experimental Biology* **211**, 170–179. (doi:10.1242/jeb.007476)

7.   Agrawal S, Tobalske BW, Anwar Z, Luo H, Hedrick TL, Cheng B. 2022 Musculoskeletal wing-actuation model of hummingbirds predicts diverse effects of primary flight muscles in hovering flight. *Proceedings of the Royal Society B: Biological Sciences* **289**. (doi:10.1098/rspb.2022.2076)

8.   Hieronymus TL. 2015 Qualitative skeletal correlates of wing shape in extant birds (Aves: Neoaves). *BMC Evol Biol* **15**, 30. (doi:10.1186/s12862-015-0303-7)

9.   Dial KP. 1992 Avian Forelimb Muscles and Nonsteady Flight: Can Birds Fly without Using the Muscles in Their Wings? *Auk* **109**, 874–885. (doi:10.2307/4088162)

10.   Warrick DR, Tobalske BW, Powers DR. 2005 Aerodynamics of the hovering hummingbird. *Nature* **435**, 1094–1097. (doi:10.1038/nature03647)

11.   Altshuler DL, Stiles FG, Dudley R. 2004 Of Hummingbirds and Helicopters: Hovering Costs, Competitive Ability, and Foraging Strategies. *Am Nat* **163**, 16–25. (doi:10.1086/380511)

12.   Sargent AJ, Groom DJE, Rico-Guevara A. 2021 Locomotion and Energetics of Divergent Foraging Strategies in Hummingbirds: A Review. *Integr Comp Biol* **61**, 736–748. (doi:10.1093/icb/icab124)

13.   Nudds RL, Dyke GJ, Rayner JM V. 2007 Avian brachial index and wing kinematics: putting movement back into bones. *J Zool* **272**, 218–226. (doi:10.1111/j.1469-7998.2006.00261.x)

14. Sievwright H, Macleod N. 2012 Eigensurface analysis, ecology, and modelling of
morphological adaptation in the falconiform humerus (Falconiformes: Aves). *Zool J*
Linn Soc **165**, 390–419. (doi:10.1111/j.1096-3642.2012.00818.x)

15. Altshuler DL. 2006 The physiology and biomechanics of avian flight at high altitude.
Integr Comp Biol **46**, 62–71. (doi:10.1093/icb/icj008)

16. Scott GR, Milsom WK. 2006 Flying high: A theoretical analysis of the factors limiting
exercise performance in birds at altitude. *Respir Physiol Neurobiol* **154**, 284–301.
(doi:10.1016/j.resp.2006.02.012)

17. Altshuler DL, Dudley R, McGuire JA. 2004 Resolution of a paradox: Hummingbird
flight at high elevation does not come without a cost. *Proceedings of the National*
Academy of Sciences **101**, 17731–17736. (doi:10.1073/pnas.0405260101)

18. Stiles FG. 2008 Ecomorphology and phylogeny of Hummingbirds: Divergence and
Convergence in Adaptations to High Elevations. *Ornitol Neotrop* **19**, 511–519.

19. Yang J, Yang C, Lin H, Lees AC, Tobias JA. 2024 Elevational constraints on flight
efficiency shape global gradients in avian wing morphology.
(doi:10.1101/2024.07.12.603304)

20. Claramunt S, Derryberry EP, Remsen J V., Brumfield RT. 2012 High dispersal ability
inhibits speciation in a continental radiation of passerine birds. *Proceedings of the*
Royal Society B: Biological Sciences **279**, 1567–1574. (doi:10.1098/rspb.2011.1922)

21. Blackburn TM, Ruggiero A. 2001 Latitude, elevation and body mass variation in
Andean passerine birds. *Global Ecology and Biogeography* **10**, 245–259.
(doi:10.1046/j.1466-822X.2001.00237.x)

22. Klingenberg CP. 2016 Size, shape, and form: concepts of allometry in geometric
morphometrics. *Dev Genes Evol* **226**, 113–137. (doi:10.1007/s00427-016-0539-2)

23. Chira AM, Cooney CR, Bright JA, Capp EJR, Hughes EC, Moody CJA, Nouri LO,
Varley ZK, Thomas GH. 2018 Correlates of rate heterogeneity in avian
ecomorphological traits. *Ecol Lett* **21**, 1505–1514. (doi:10.1111/ele.13131)

24. Teplitsky C *et al.* 2014 Assessing Multivariate Constraints to Evolution across Ten
Long-Term Avian Studies. *PLoS One* **9**, e90444. (doi:10.1371/journal.pone.0090444)

25. Harmon LJ *et al.* 2010 Early bursts of body size and shape evolution are rare in
comparative data. *Evolution (N Y)* , no-no. (doi:10.1111/j.1558-5646.2010.01025.x)

26. Cooney CR, Bright JA, Capp EJR, Chira AM, Hughes EC, Moody CJA, Nouri LO,
Varley ZK, Thomas GH. 2017 Mega-evolutionary dynamics of the adaptive radiation
of birds. *Nature* **542**, 344–347. (doi:10.1038/nature21074)

27. Poore SO, Ashcroft A, Sánchez-Haiman A, Goslow GE. 1997 The Contractile
Properties of the M. Supracoracoideus in the Pigeon and Starling: a Case for Long-

Axis Rotation of the Humerus. *Journal of Experimental Biology* **200**, 2987–3002.
(doi:10.1242/jeb.200.23.2987)

28.   Hallgrímsson B, Katz DC, Aponte JD, Larson JR, Devine J, Gonzalez PN, Young NM,
Roseman CC, Marcucio RS. 2019 Integration and the Developmental Genetics of
Allometry. *Integr Comp Biol* **59**, 1369–1381. (doi:10.1093/icb/icz105)

29.   Felice RN, Watanabe A, Cuff AR, Noirault E, Pol D, Witmer LM, Norell MA,
O'Connor PM, Goswami A. 2019 Evolutionary Integration and Modularity in the
Archosaur Cranium. *Integr Comp Biol* **59**, 371–382. (doi:10.1093/icb/icz052)

30.   Adams DC, Felice RN. 2014 Assessing Trait Covariation and Morphological
Integration on Phylogenies Using Evolutionary Covariance Matrices. *PLoS One* **9**,
e94335. (doi:10.1371/journal.pone.0094335)

31.   Orkney A, Bjarnason A, Tronrud BC, Benson RBJ. 2021 Patterns of skeletal integration
in birds reveal that adaptation of element shapes enables coordinated evolution
between anatomical modules. *Nat Ecol Evol* **5**, 1250–1258. (doi:10.1038/s41559-021-
01509-w)

32.   Hallgrímsson B, Jamniczky H, Young NM, Rolian C, Parsons TE, Boughner JC,
Marcucio RS. 2009 Deciphering the Palimpsest: Studying the Relationship Between
Morphological Integration and Phenotypic Covariation. *Evol Biol* **36**, 355–376.
(doi:10.1007/s11692-009-9076-5)

33.   Klingenberg CP. 2008 Morphological Integration and Developmental Modularity. *Annu*
Rev Ecol Evol Syst **39**, 115–132. (doi:10.1146/annurev.ecolsys.37.091305.110054)

34.   Revell LJ, Harmon LJ, Collar DC. 2008 Phylogenetic Signal, Evolutionary Process,
and Rate. *Syst Biol* **57**, 591–601. (doi:10.1080/10635150802302427)

35.   Felice RN, Goswami A. 2018 Developmental origins of mosaic evolution in the avian
cranium. *Proceedings of the National Academy of Sciences* **115**, 555–560.
(doi:10.1073/pnas.1716437115)

36.   Navalón G, Marugán-Lobón J, Bright JA, Cooney CR, Rayfield EJ. 2020 The
consequences of craniofacial integration for the adaptive radiations of Darwin's finches
and Hawaiian honeycreepers. *Nat Ecol Evol* **4**, 270–278. (doi:10.1038/s41559-019-
1092-y)

37.   Gill F, Donsker D, Rasmussen P. 2023 IOC World Bird List 13.1. *IOC World Bird List*
Datasets. (doi:10.14344/IOC.ML.13.1)

38.   Zelditch ML, Swiderski DL, Sheets HD, Fink WL. 2012 *Geometric Morphometrics for*
Biologists. 2nd edn. Elsevier. (doi:10.1016/B978-0-12-778460-1.X5000-5)

39.   Bookstein FL. 1992 *Morphometric Tools for Landmark Data*. Cambridge University
Press. (doi:10.1017/CBO9780511573064)

40. Rohlf FJ. 2015 The tps series of software. *Hystrix* **26**, 1–4. (doi:10.4404/hystrix-26.1-
11264)

41. Cardini A, Elton S. 2007 Sample size and sampling error in geometric morphometric
studies of size and shape. *Zoomorphology* **126**, 121–134. (doi:10.1007/s00435-007-
0036-2)

42. R Core Team. 2025 R: A language and environment for statistical computing.

43. Rohlf FJ, Slice D. 1990 Extensions of the Procrustes Method for the Optimal
Superimposition of Landmarks. *Syst Zool* **39**, 40. (doi:10.2307/2992207)

44. Adams D, Collyer M, Kaliontzopoulou A, Baken E. 2012 geomorph: Geometric
Morphometric Analyses of 2D and 3D Landmark Data. *CRAN: Contributed Packages*.
(doi:10.32614/CRAN.package.geomorph)

45. Billerman SM, Keeney BK, Kirwan GM, Medrano F, Sly ND, Schmidt MG. 2025
Birds of the World. *Cornell Laboratory of Ornithology, Ithaca, NY, USA.*
https://birdsoftheworld.org/bow/ (doi:https://doi.org/10.2173/bow)

46. Quintero I, Jetz W. 2018 Global elevational diversity and diversification of birds.
Nature **555**, 246–250. (doi:10.1038/nature25794)

47. Tobias JA *et al.* 2022 AVONET: morphological, ecological and geographical data for
all birds. *Ecol Lett.* **25**, 581–597. (doi:10.1111/ele.13898)

48. Blomberg SP, Garland T, Ives AR. 2003 Testing for phylogenetic signal in comparative
data: behavioral traits are more labile. *Evolution (N Y)* **57**, 717–745.
(doi:10.1111/j.0014-3820.2003.tb00285.x)

49. McGuire JA, Witt CC, Remsen JV, Corl A, Rabosky DL, Altshuler DL, Dudley R. 2014
Molecular Phylogenetics and the Diversification of Hummingbirds. *Current Biology*
24, 910–916. (doi:10.1016/j.cub.2014.03.016)

50. Collyer ML, Baken EK, Adams DC. 2022 A standardized effect size for evaluating and
comparing the strength of phylogenetic signal. *Methods Ecol Evol* **13**, 367–382.
(doi:10.1111/2041-210X.13749)

51. Schlager S. 2017 Morpho and Rvcg – Shape Analysis in R. In *Statistical Shape and
Deformation Analysis*, pp. 217–256. Elsevier. (doi:10.1016/B978-0-12-810493-
4.00011-0)

52. Revell LJ. 2012 Phytools: an R package for phylogenetic comparative biology (and
other things). *Methods Ecol Evol* **3**, 217–223. (doi:10.1111/j.2041-210X.2011.00169.x)

53. Castiglione S, Tesone G, Piccolo M, Melchionna M, Mondanaro A, Serio C, Di
Febbraro M, Raia P. 2018 A new method for testing evolutionary rate variation and
shifts in phenotypic evolution. *Methods Ecol Evol* **9**, 974–983. (doi:10.1111/2041-
210X.12954)

54. Biewener AA. 1982 Bone Strength in Small Mammals and Bipedal Birds: Do Safety
Factors Change With Body Size? *Journal of Experimental Biology* **98**, 289–301.
(doi:10.1242/jeb.98.1.289)

55. Bright JA, Marugán-Lobón J, Cobb SN, Rayfield EJ. 2016 The shapes of bird beaks
are highly controlled by nondietary factors. *Proceedings of the National Academy of*
Sciences **113**, 5352–5357. (doi:10.1073/pnas.1602683113)

56. Shatkovska O V., Ghazali M. 2020 Integration of skeletal traits in some passerines:
impact (or the lack thereof) of body mass, phylogeny, diet and habitat. *J Anat* **236**,
274–287. (doi:10.1111/joa.13095)

57. Holmes J, Sustaita D, Hertel F. 2022 Geometric morphometric analysis of the humerus
in New and Old World vultures. *J Morphol* **283**, 379–394. (doi:10.1002/jmor.21449)

58. Skandalis DA *et al.* 2017 The biomechanical origin of extreme wing allometry in
hummingbirds. *Nat Commun* **8**, 1047. (doi:10.1038/s41467-017-01223-x)

59. Bleiweiss R. 2002 Patagial complex evolution in hummingbirds and swifts
(Apodiformes): a molecular phylogenetic perspective. *Biological Journal of the*
Linnean Society **77**, 211–219. (doi:10.1046/j.1095-8312.2002.00105.x)

60. Drury JP, Clavel J, Tobias JA, Rolland J, Sheard C, Morlon H. 2024 Limited ecological
opportunity influences the tempo of morphological evolution in birds. *Current Biology*
34, 661-669.e4. (doi:10.1016/j.cub.2023.12.055)

61. Sonne J *et al.* 2020 Ecological mechanisms explaining interactions within plant–
hummingbird networks: morphological matching increases towards lower latitudes.
Proceedings of the Royal Society B: Biological Sciences **287**, 20192873.
(doi:10.1098/rspb.2019.2873)

62. Dakin R, Segre PS, Straw AD, Altshuler DL. 2018 Morphology, muscle capacity, skill,
and maneuvering ability in hummingbirds. *Science (1979)* **359**, 653–657.
(doi:10.1126/science.aao7104)

63. Lim MCW, Bi K, Witt CC, Graham CH, Dávalos LM. 2021 Pervasive Genomic
Signatures of Local Adaptation to Altitude Across Highland Specialist Andean
Hummingbird Populations. *Journal of Heredity* **112**, 229–240.
(doi:10.1093/jhered/esab008)

64. Zusi RL. 2013 Introduction to the Skeleton of Hummingbirds (Aves: Apodiformes,
Trochilidae) in Functional and Phylogenetic Contexts. *Ornithol Monogr* **77**, 1–94.
(doi:10.1525/om.2013.77.1.1)

65. Tiebout HM. 1992 Comparative energetics of divergent foraging modes: a doubly
labelled water experiment on hummingbird competition. *Anim Behav* **44**, 895–906.
(doi:10.1016/S0003-3472(05)80585-8)

812 66. Rombaut LMK, Capp EJR, Hughes EC, Varley ZK, Beckerman AP, Cooper N, Thomas
813   GH. 2022 The evolution of the traplining pollinator role in hummingbirds:
814   specialization is not an evolutionary dead end. *Proceedings of the Royal Society B:*
815   *Biological Sciences* **289**. (doi:10.1098/rspb.2021.2484)

816 67. Goswami A, Randau M, Polly PD, Weisbecker V, Bennett CV, Hautier L, Sánchez-
817   Villagra MR. 2016 Do Developmental Constraints and High Integration Limit the
818   Evolution of the Marsupial Oral Apparatus? *Integr Comp Biol* **56**, 404–415.
819   (doi:10.1093/icb/icw039)

820 68. Randau M, Goswami A. 2017 Unravelling intravertebral integration, modularity and
821   disparity in Felidae (Mammalia). *Evol Dev* **19**, 85–95. (doi:10.1111/ede.12218)

822 69. Fabre A-C *et al.* 2020 Metamorphosis shapes cranial diversity and rate of evolution in
823   salamanders. *Nat Ecol Evol* **4**, 1129–1140. (doi:10.1038/s41559-020-1225-3)

826　　　　　　　　　　　　　　　　　　　　**Tables**

Table 1. Ecological and morphological predictors explain variation in hummingbird

humeral shape, with size, range, and body mass showing the strongest effects. Humerus

shape variation is significantly associated with humerus size, geographic range size, body

mass, and foraging strategy. The table reports effect sizes and significance values from the

final multivariate PGLS model, including degrees of freedom (df), coefficient of

determination (R^2), F-statistics, Z-scores, and p-values for each predictor. Asterisks indicate

significance levels (* $p < 0.05$, ** $p < 0.01$).

Predictor	df	R^2	F	Z	p-value	Significance
Foraging strategy	2	0.035	1.536	1.872	0.035	*
Latitude	1	0.016	1.445	1.458	0.070	.
Body mass	1	0.022	1.927	1.894	0.031	*
Humerus size	1	0.081	7.120	3.078	0.002	**
Range size	1	0.031	2.774	2.520	0.008	**
Elevation	1	0.013	1.209	1.251	0.102	

Table 2. Foraging strategy shows significantly different evolutionary rates of humeral

shape across regimes, whereas continuous predictors show no detectable univariate

effects. Foraging strategy explains significant differences in evolutionary rates of humeral

morphology when tested using compare.evol.rates, whereas RRphylo-based univariate

regressions show no meaningful effects for latitude, body mass, humerus size, range size, or

elevation. The permutation ANOVA (1000 residual randomizations) reports Df, R^2, F, Z, and

p-values for the categorical test. The Kappa statistic summarizes overall model fit. The

bottom section reports univariate PGLS results based on RRphylo tip rates, including

parameter estimates, standard errors, maximum-likelihood λ, and p-values. Significance

codes: *** $p < 0.001$.

Analysis of variance using residual randomization (1000 permutations)						
Predictor	Df	R^2	F	Z	Pr(>F)	Significance
Foraging strategy	2	0.062	2.517	3.465	0.001	***
Residuals	75	0.937				
Kappa statistic	0.143					
Univariate PGLS values						
Predictor	Estimate	Std. Error	Lambda (ML)	p-value		
Latitude	-0.0001	8.92E-05	0.630	0.132		
Body mass	0.000	0.00048	0.571	0.476		
Humerus size	-0.038	0.406	0.572	0.925		
Range size	1.49E-10	3.56E-10	0.568	0.678		
Elevation	-8.95E-08	6.47E-07	0.570	0.890		

 Figures

 Figure 1. Hummingbird humeral shape shows strong phylogenetic structure and

substantial evolutionary rate heterogeneity across major clades. Phylogenetic structure in

humeral morphology is high (K = 0.309), and evolutionary rates vary markedly across

hummingbird clades, with several lineages showing accelerated morphological evolution. (A)

Twenty-five anatomical landmarks used to quantify humerus shape. (B) Landmark dispersion

across specimens illustrating within-family morphological diversity. (C) Morphological

deformation grids for PC1 and PC2 showing major axes of shape variation. (D) Null

distribution of phylogenetic signal with the observed value (red arrow). The circular tree

shows the phylogeny of the 78 species analyzed; branch colors depict species-specific

evolutionary rates, and radial bars reflect normalized ecological and morphological predictors.

The lower-left inset shows the distribution of log-transformed mean evolutionary rates.

Hummingbird illustrations were AI-generated (ChatGPT-4o) and refined in Adobe Illustrator.

Figure 2. Humeral shape (PC1) is strongly associated with humerus size, range size,

body mass, and foraging strategies. Humeral shape (PC1) shows significant associations

with foraging strategy, body mass, humerus size, and geographic range size, consistent with

the multivariate PGLS results. Each panel shows the relationship between PC1 and a predictor

included in the model. Continuous predictors are displayed as scatterplots with fitted

regression lines and 95% confidence intervals (shaded areas), and foraging strategy is shown

as boxplots with point distributions. PC1 reflects a gradient from shorter, more robust humeri

(higher values) to longer, more slender humeri (lower values). Although PC1 is shown here

for visualization, the full multivariate analysis evaluates shape variation across the entire

morphometric space.

Figure 3. Hummingbird foraging strategies occupy distinct regions of humeral

morphospace, with clear group separation in both PCA and CVA analyses. Foraging

strategies differ in their occupation of humeral morphospace, with territorial, traplining, and

opportunistic species forming partially distinct clusters in both unsupervised (PCA) and

supervised (CVA) shape spaces. (A) PCA of Procrustes-aligned humeral shapes, with PC1 and

PC2 explaining 17.97% and 16.39% of variation, respectively. Points are colored by foraging

strategy; shaded convex hulls denote group morphospaces, and thin black lines represent

phylogenetic relationships projected into shape space. (B) Mean humeral shape variation

across specimens visualized as a thin-plate spline deformation of the consensus configuration;

red points denote landmarks. (C) Canonical Variate Analysis maximizing group separation

based on foraging strategy. Dashed ellipses show 90% confidence regions, and purple

diamonds mark group centroids. Shape outlines adjacent to the axes illustrate humeral

changes associated with CV1 and CV2.

Figure 4. Evolutionary rates of humeral shape vary among foraging strategies and show

no detectable associations with continuous ecological predictors. Foraging strategies differ

in their average evolutionary rates of humeral shape, while RRphylo-derived rates show no

significant relationships with body mass, humerus size, range size, elevation, or latitude. The

top-left panel displays group-level evolutionary rates estimated using compare.evol.rates,

illustrating differences among territorial, traplining, and opportunistic species. All remaining

panels show RRphylo tip rates plotted against continuous predictors; points represent species-

level $\log_{10}$-transformed evolutionary rates, and purple lines indicate fitted PGLS regressions

with 95% confidence intervals (shaded areas). "Time from root" corresponds to the relative

phylogenetic distance of each tip and is shown for visualization only.

Figure 5. Proximal and distal regions of the hummingbird humerus are strongly integrated, supporting a functional linkage rather than modular independence. The proximal and distal regions of the humerus exhibit strong morphological integration, indicating coordinated evolutionary changes rather than independent modular evolution. The scatterplot shows the relationship between PLS1 scores of the proximal region and PLS2 scores of the distal region, with a high correlation (r-PLS = 0.899) and strong statistical support (Z = 4.2219, p = 0.001). The fitted regression line and 95% confidence interval illustrate the magnitude and direction of covariation across species. The diagrams on the right depict the *a priori* modularity hypothesis (upper panel), representing the expected division of the humerus into proximal and distal functional units, contrasted with the data-driven minimum-covariation modular structure (lower panel). Nodes represent landmarks and edges represent strength of covariation. The strong positive PLS correlation demonstrates that evolutionary changes in the proximal humerus (e.g., deltopectoral crest and articular surface geometry) are tightly linked to changes in the distal region (e.g., condyles and shaft curvature), consistent with shared biomechanical demands during hovering and forward flight in hummingbirds.