Foraging behavior and ecology drive the evolution of humeral shape in hummingbirds

Juan Camilo Ríos-Orjuela[1,2,*], Carlos Daniel Cadena[1], Alejandro Rico-Guevara[3, 4], Ilias

Berberi[5], Lauren Miner[5] & Roslyn Dakin[5]

[1]Laboratorio de Biología Evolutiva de Vertebrados, Departamento de Ciencias Biológicas,

Universidad de los Andes, Bogotá, Colombia

[2]Grupo de Morfología y Ecología Evolutiva, Instituto de Ciencias Naturales, Universidad

Nacional de Colombia, Sede Bogotá, Colombia

[3]Department of Biology, University of Washington, Seattle, WA, U.S.A.

[4]Burke Museum of Natural History and Culture, University of Washington, Seattle, WA,

U.S.A.

[5]Department of Biology, Carleton University, Ottawa, Ontario, Canada

* Corresponding author (jcriosor@gmail.com)

Orcid: JCR: 0000-0001-6976-9131, DC: 0000-0003-4530-2478, ARG: 0000-0003-4067-

5312, IB: 0000-0003-3051-8145, LM: 0009-0008-1991-7472, RD: 0000-0002-3140-3975.

Author contributions: JCR: Conceptualization, Methodology, Software, Formal analysis,

Investigation, Data curation, Writing – Original draft, Writing – Review & Editing,

Visualization. **RD:** Conceptualization, Validation, Investigation, Resources, Writing –

Original draft, Writing – Review & Editing, Supervision, Funding acquisition. **CDC:**

Conceptualization, Writing – Review & Editing, Supervision, Funding acquisition. **IB:**

Investigation, Data curation, Writing – Review & Editing. **LM:** Investigation, Data curation.

ARG: Conceptualization, Investigation, Writing – Review & Editing, Supervision.

Funding: This research was supported by an NSERC Discovery Grant (Canada), the Faculty

of Sciences at Universidad de los Andes (Bogotá, Colombia), and the Emerging Leaders in

the Americas Program (Global Affairs Canada).

Conflict of Interest: The authors declare no conflict of interest.

Acknowledgments: We thank Steven Cardiff and Nick Mason for their support during

fieldwork and specimen access, as well as the Louisiana State University Museum of Natural

Science for facilitating access to their ornithological collections.

32   **Abstract**

Skeletal morphology is central to avian flight performance and ecological adaptation. In

hummingbirds, the humerus is critical for power generation and wing control, yet its

evolutionary diversification remains poorly understood. We integrate geometric

morphometrics, phylogenetic comparative methods, and ecological data to study humeral

shape evolution across a diverse sample of 80 hummingbird species. Humeral shape variation

was associated with ecological and geographical variables, including body mass, foraging

strategy, range size and latitude. Larger body mass or smaller humeri corresponded to more

robust, curved shapes, consistent with biomechanical demands of supporting greater forces. In

contrast, broader geographic ranges are associated with more slender shapes, possibly

reflecting aerodynamic efficiency across variable environments. Humerus shape also changed

with foraging strategy: traplining species, which visit flowers along dispersed routes, exhibit

elongated humeri favoring efficient, sustained flight, whereas territorial species showed

shorter, robust shapes suited to rapid maneuvers. Morphological evolution rates differed

among strategies, with trapliners showing highest rates, and increased significantly with range

size. Strong phenotypic integration within the humerus indicated that evolutionary changes

are tightly coordinated across the bone regions. These findings demonstrate how ecological

pressures, scaling effects, and developmental constraints together shape flight-related

morphology, underscoring the adaptive flexibility of the avian skeleton.

Keywords: Adaptive evolution, Allometry, Biomechanics, Bird flight, Geometric

morphometrics, Phenotypic modularity and integration.

Teaser text

How do tiny hummingbirds balance agility, power, and efficiency in flight? By analyzing humeral shapes from 80 species, we reveal how body size, foraging strategy, and geography shape this key wing bone. Trapliners and territorial species show distinct morphologies linked to their flight demands, while broader geographic ranges correlate with faster evolutionary rates. Strong integration within the humerus suggests that changes in one region influence the entire bone. Our findings uncover the interplay of ecology, biomechanics, and evolutionary history in shaping the avian skeleton.

Introduction

Understanding the evolution of morphological diversity remains a central topic in

evolutionary biology. Organisms diversify morphologically in response to ecological

pressures, functional constraints, and historical contingencies, and disentangling these

influences requires integrative approaches combining morphology, ecology, and phylogeny

(Losos, 2011; Pigot *et al.*, 2020; Sayol *et al.*, 2025). Birds, among the most ecologically and

morphologically varied vertebrates, offer an insightful system to explore these dynamics of

trait diversification, especially in traits shaped by the demands of powered flight (Norberg,

1990; Dudley, 2002).

Among skeletal components, the avian humerus plays a pivotal role in flight mechanics and

ecological adaptation. It connects the axial skeleton to the forelimb and anchors major flight

muscles responsible for the downstroke and upstroke of the wingbeat (Tobalske & Biewener,

2008). In addition, humerus morphology affects lift, wing loading, and maneuverability

(Agrawal *et al.*, 2022). Despite its biomechanical importance, the humerus remains

underexplored in comparative studies of avian flight, which have often emphasized external

features such as wing aspect ratio (Hieronymus, 2015). While such metrics effectively capture

overall aerodynamic properties, they do not reflect underlying skeletal variation, potentially

overlooking functionally significant musculoskeletal morphology.

In hummingbirds, the humerus is distinctively short, robust, and exhibits morphological

adaptations that support their specialized hovering and maneuverable flight (Agrawal *et al.*,

2022). Unlike the long, slender humeri typical of birds adapted for gliding or forward-

powered flight (Dial, 1992), the hummingbird humerus is shortened to perform exceptionally

high wingbeat frequencies and rapid rotation of the wing on each upstroke (Warrick *et al.*,

2005). These changes allow hummingbirds to generate lift during both downstroke and

upstroke and exert fine control during rapid aerial maneuvers (Agrawal *et al.*, 2022). To meet

these challenges, the humerus functions as a rigid, force-transmitting lever for enlarged flight

muscles, and exhibits structural specializations such as increased cortical thickness, a

relatively straight shaft, and pronounced muscle attachment sites at the deltopectoral crest.

This morphology reflects a biomechanical compromise: a shorter, sturdier humerus reduces

inertial resistance and facilitates maneuverability, but may constrain aspects of aerodynamic

efficiency or range of motion, requiring precise tuning to maintain energetic economy during

sustained flight. With over 300 species, the hummingbird clade presents a rich system for

comparative study, offering insight into how skeletal structures evolve under varying

ecological demands and locomotor constraints. Yet despite its biomechanical importance, the

hummingbird humerus remains underexplored in evolutionary studies.

Hummingbirds exhibit diverse foraging strategies that reflect their ecological niches and

influence their flight behavior. Some species defend floral territories aggressively, engaging in

rapid, maneuver-demanding flights to repel intruders, while others adopt traplining behaviors,

following predictable expansive foraging circuits that require sustained travel across larger

areas and efficient navigation between patchy floral resources (Altshuler *et al.*, 2004b;

Sargent *et al.*, 2021). These differing foraging strategies are expected to impose distinct

aerodynamic and biomechanical demands, which may drive morphological divergence in the

skeleton. Species engaging in frequent, high-intensity maneuvers—such as territorial

defense—may benefit from shorter, more robust humeri that facilitate rapid changes in

direction and powerful wing strokes (Agrawal *et al.*, 2022), In contrast, species relying on

traplining may favor longer and more slender humeri that enhance flight efficiency and reduce

energetic costs over longer distances, even if they still require precise control for floral

approach and handling (Nudds *et al.*, 2007; Sievwright & Macleod, 2012).

Beyond foraging behavior, ecological factors such as those varying with elevation also shape

hummingbird morphology. Species occupying high-altitude environments face hypoxia and

reduced air density, which increase the biomechanical demands of flight (Altshuler, 2006;

Scott & Milsom, 2006). Prior studies have documented phenotypic divergence along

elevational gradients, particularly in wing shape, body size, and metabolic traits (Altshuler *et*

al., 2004a; Stiles, 2008). However, skeletal elements like the humerus remain understudied in

this context, despite their critical role in flight mechanics.

Additional ecological and morphological factors may further influence humeral evolution.

Species occurring at higher latitudes or occupying broader geographic ranges often encounter

greater environmental heterogeneity and dispersal demands, potentially favoring either

morphological flexibility or specialization (Claramunt *et al.*, 2012; Yang *et al.*, 2024). Body

mass also imposes biomechanical constraints, as larger species must support greater

gravitational and aerodynamic forces (Blackburn & Ruggiero, 2001). Finally, overall skeletal

size may influence scaling relationships critical for wing loading and flight efficiency

(Klingenberg, 2016).

Together, these patterns suggest that humeral morphology in hummingbirds is likely shaped

by a complex interplay of biomechanical constraints, ecological demands, and evolutionary

history. Yet, the extent to which ecological variables predict morphological divergence—and

whether they are also associated with evolutionary rates—remains underexplored (Teplitsky

et al., 2014; Chira *et al.*, 2018). Evaluating evolutionary rates is important, as accelerated

shape evolution may reflect ecological opportunity, functional innovation, or release from

constraints (Harmon *et al.*, 2010; Cooney *et al.*, 2017). In hummingbirds, one may expect that

body size, ecological and geographical factors could all influence not only humeral shape but

also the tempo of its evolution. By testing for associations among such variables, one may

clarify how ecological and morphological factors jointly shape both the patterns and pace of

morphological change in this key flight structure.

The hummingbird humerus, with its distinct proximal and distal functions, also provides an

appropriate system to investigate phenotypic integration. The proximal region primarily

serves as the attachment site for the pectoralis and supracoracoideus muscles, which are

responsible for generating the downstroke and upstroke during flight, respectively (Poore *et*

al., 1997; Tobalske & Biewener, 2008). The distal portion of the humerus articulates with the

radius and ulna, transmitting the forces generated by these muscles into wing motion and

modulating the wing's orientation and amplitude. Understanding whether these regions evolve

in a coordinated manner offers insights into developmental constraints and adaptive flexibility

(Adams & Felice, 2014; Felice *et al.*, 2019; Hallgrímsson *et al.*, 2019). Morphological traits,

especially those underpinning locomotion, rarely evolve independently (Orkney *et al.*, 2021).

Instead, they exhibit integration and modularity due to shared developmental pathways and

functional covariation (Klingenberg, 2008; Hallgrímsson *et al.*, 2009). While many studies of

phenotypic integration focus on patterns across multiple bones or skeletal regions (e.g., skull

and mandible, or entire limbs), exploring integration within a single element can reveal how

distinct functional regions remain evolutionarily coordinated despite localized biomechanical

specializations.

Here, we investigate how foraging strategy, ecological pressures, and evolutionary processes

collectively shape humeral morphology across a broad sample hummingbird species. First, we

hypothesized that species differing in foraging strategies would exhibit divergent humeral

morphologies, reflecting adaptations to their specific flight demands and ecological contexts.

We also explored whether these strategies are linked to variation in the evolutionary rates of

humeral shape. Rather than assuming a single directional outcome, we framed this analysis as

a hypothesis-generating approach informed by the contrasting demands of different foraging

behaviors. For example, traplining—characterized by repeated, sustained flight along spatially

extensive routes—may select for morphologies that enhance energetic efficiency and

maneuverability across heterogeneous environments. Territorial behavior, by contrast,

involves frequent short-distance accelerations and complex aerial contests, potentially

selecting for structural robustness and high maneuverability. Whether such behaviors promote

morphological innovation or constrain it through stabilizing selection likely depends on

evolutionary history, ecological variability, and biomechanical trade-offs. Given that

evolutionary rates can reflect diverse processes, including directional selection, stabilizing

selection, or release from constraint (Revell *et al.*, 2008), we interpret our rate comparisons

within a framework that considers multiple evolutionary scenarios, rather than presupposing a

single outcome.

Second, we predicted that humeral shape variation and evolutionary rates would be influenced

by a combination of ecological and morphological factors. Specifically, species inhabiting

higher elevations may exhibit distinctive humeral morphologies, as high-altitude

environments impose greater aerodynamic and metabolic constraints that may drive adaptive

changes. However, whether these conditions lead to accelerated evolutionary rates remains

uncertain. In some cases, strong stabilizing selection or habitat filtering may constrain

morphological evolution, resulting in low rates despite ecological pressure (Revell *et al.*,

2008). Similarly, while broad geographic ranges or occurrence at higher latitudes have been

associated with increased environmental heterogeneity and dispersal challenges, the

relationship with evolutionary rate is context-dependent and may vary across clades and traits.

Regarding body mass, we predicted that larger-bodied species would exhibit corresponding

shifts in humeral morphology, reflecting biomechanical scaling effects on flight performance

and energetic costs. Although allometric relationships are expected, examining whether these

scaling patterns vary across ecological contexts could provide insights into how size-related

constraints interact with ecological specialization. Finally, we expected overall skeletal size to

correlate with humeral shape variation due to its impact on wing loading and flight efficiency.

Lastly, we hypothesized that the humerus exhibits strong internal phenotypic integration, with

coordinated morphological evolution between its proximal and distal regions, given their

shared role in force transmission and wing mechanics. Such integration is expected because

modifications in one region (e.g., altering muscle attachment sites proximally) would

necessitate compensatory changes distally to maintain flight performance. However, examples

from other systems suggest that bones can evolve in a modular fashion when regions fulfill

distinct functions or face divergent selective pressures (Felice & Goswami, 2018; Navalón *et*

al., 2020; Orkney *et al.*, 2021). Whether such modularity also occurs within the hummingbird

humerus remains an open question but would imply that evolutionary changes could be

targeted to specific regions without affecting the entire structure. By integrating ecological

categories and morphometric analyses within a phylogenetic framework, our objective is to

provide a comprehensive assessment of the factors driving morphological evolution in the

hummingbird humerus.

202   **Methods**

Specimens and image acquisition

We analyzed 80 hummingbird species, representing approximately 22% of extant species

diversity in the family Trochilidae (Gill *et al.*, 2023). To capture morphological variation of

the humerus, we photographed the right humerus in cranial view from male specimens in the

collection of the Louisiana State University Museum of Natural Science (LSUMNS). We

focused on males to reduce potential shape variation due to sexual dimorphism and to ensure

consistency across species, as male specimens are more frequently available and

taxonomically reliable in museum collections. Images were taken using a Canon EOS M50

camera equipped with a 28 mm macro lens. The camera was mounted on a copy stand at a

fixed distance from the specimens to maintain consistent magnification and perspective. Each

photograph included an embedded scale bar (10 mm with 0.1 mm divisions) for accurate

scaling in subsequent analyses. This standardized view facilitated consistent comparative

analysis of skeletal structure across species (Zelditch *et al.*, 2012).

A total of 25 anatomical landmarks were digitized on each humerus, focusing primarily on

points along the contour to capture overall shape (Figure 1A-B). Landmarks were chosen

based on biomechanical relevance and anatomical consistency across taxa (Bookstein, 1992;

Klingenberg, 2016). No semilandmarks were employed, because we aimed to focus on

homologous and discrete anatomical points rather than continuous curves. A detailed

description of landmarks and specimens list is provided as supplementary material. Landmark

digitization was conducted using TPSDig2 version 2.32 (Rohlf, 2015).

Given the rarity of skeletal material for many hummingbird species, sample sizes varied

across taxa. For approximately 87% of species (70 of 80), only a single male specimen was

available. For the remaining 10 species, where multiple specimens (two to four per species)

were available, we calculated the species-specific mean shape by averaging landmark

configurations across individuals (Cardini & Elton, 2007).

Morphometric data preparation

All data preparation and statistical analyses were conducted in R version 4.4.2 (R Core Team,

2024). Landmark data were aligned using a Generalized Procrustes Analysis (GPA) to remove

non-shape variation related to position, orientation, and scale (Rohlf & Slice, 1990). GPA was

performed using the gpagen() function in the *geomorph* package (Adams *et al.*, 2012).

Following alignment, we calculated squared Euclidean distances from each specimen to the

mean shape configuration to identify and exclude outliers. We retained 97.5% of specimens

closest to the mean shape, a typical threshold in morphometric studies. Euclidean distances

were computed by converting the aligned 3D landmark array to a two-dimensional matrix,

followed by row-wise distance calculations. Two specimens—*Calypte anna* and *Goethalsia*

bella—were identified as outliers for their unusually high morphological deviation from the

overall mean shape configuration, due to preservation or landmarking artifacts, and were thus

removed from further investigation. Following this filtering, 78 species were retained for

downstream analyses. A PCA analysis of the aligned coordinates was performed exclusively

for graphical representation of the shape variation and morphospace visualization (Figure 1C).

Species traits

For each species, we recorded a suite of ecological and morphological traits. Foraging

strategy was classified as territorial, trapliner, or opportunist, based on typical foraging

behavior and resource use (Billerman *et al.*, 2022). We also recorded a species' maximum

elevation as a proxy for elevational environment (Quintero & Jetz, 2018). Geographic range

size (km²) was obtained as we expect species occupying larger ranges to experience broader

environmental heterogeneity. Body mass was included as a proxy for biomechanical

constraints associated with size, given that larger species typically experience different

mechanical stresses during flight (Altshuler *et al.*, 2004b). Range size, latitude at the centroid

of the species' range, and body mass data were retrieved from Tobias et al. (2022).

Additionally, as a structure-specific indicator of skeletal size, we measured humerus centroid

size from the landmark data. While centroid size is mathematically derived from the

distribution of landmark coordinates, it is widely recognized as a robust proxy for overall size

in morphometric analyses (Klingenberg, 2016).

Phylogenetic signal

To assess whether humeral shape variation exhibited phylogenetic signal, we quantified

Blomberg's multivariate K statistic (Blomberg *et al.*, 2003; Adams & Felice, 2014) using the

physignal() function in *geomorph*, based on 1,000 permutations. We also computed a Z-

transformed effect size to contextualize the observed K values relative to a null distribution.

An ultrametric phylogeny was obtained from McGuire et al. (2014) for all comparative

analyses.

Phylogenetic modeling of humeral shape variation

To evaluate how ecological and morphological predictors influence humeral shape, we

implemented a phylogenetic generalized least squares (PGLS) approach using the

procD.pgls() function in *geomorph* for high-dimensional shape data. We first optimized the

lambda (λ) parameter for the phylogenetic correlation structure using likelihood profiling. The

optimization converged on $\lambda = 0.99$, indicating strong phylogenetic structure consistent with

Brownian motion. Permutation-based significance testing was conducted using 1,000

iterations, and effect sizes were calculated as Cohen's f^2.

The multivariate model included foraging strategy as a categorical predictor, alongside

continuous ecological and morphological variables: maximum elevation, absolute latitude,

range size, body mass, and humerus size. All predictors were incorporated simultaneously in

the same phylogenetic generalized least squares (PGLS) regression framework. None of the

variables exhibited strong collinearity (Pearson's $r < 0.36$), and all variance inflation factors

were within acceptable limits (VIF < 1.5).

Discriminant analysis of humeral shape by foraging strategy

To further assess whether humeral shape is differentiated among the three major foraging

strategies, we conducted a Canonical Variates Analysis (CVA). Prior to the CVA, we corrected

allometric effects by regressing Procrustes-aligned shape coordinates against log-transformed

centroid size, following standard procedures (Klingenberg, 2016). The residuals of this

regression represent shape variation independent of humerus size. We then analyzed these

residuals using the CVA() function in the *Morpho* package (Schlager, 2017), specifying

foraging strategy as the grouping variable. Statistical robustness of group discrimination was

assessed with 1,000 permutations. We also projected phylogenetic relationships into the

canonical morphospace using the phylomorphospace() function from *phytools* (Revell, 2012).

Evolutionary rates of humeral shape

To test whether the evolutionary rate of humeral shape varied among ecological categories,

we compared rates among foraging strategies using the compare.evol.rates() function in

geomorph, with 1,000 permutations. We further examined whether continuous predictors were

associated with variation in evolutionary rates. For each species, we calculated Procrustes

mean shape and reconstructed ancestral shapes using the fastAnc() function in *phytools*.

Morphological change along terminal branches was computed as the Euclidean distance

between ancestral and descendant shapes, standardized by branch length obtained from the

ultrametric phylogeny. To minimize the influence of extreme values, we excluded the top 10%

of evolutionary rates. We then modeled evolutionary rates as a function of five continuous

predictors—maximum elevation, range size, absolute latitude, body mass, and humerus size—

using a multivariate generalized additive model (GAM) implemented via the *mgcv* package in

R. This approach allowed us to flexibly account for potential nonlinear relationships between

predictors and evolutionary rates.

Phenotypic modularity and integration

We assessed phenotypic integration within the humerus by partitioning the landmark data into

proximal and distal regions. The proximal region included the humeral head and articular

surfaces, whereas the distal region encompassed the shaft and insertion points. We estimated

phylogenetic partial least squares (PLS) correlations between these modules using the

phylo.integration() function (Adams & Felice, 2014), with statistical significance assessed

through 1,000 permutations.

310   **Results**

Phylogenetic signal

Humeral shape exhibited a significant phylogenetic signal ($K = 0.309$, $p = 0.001$), indicating

that closely related hummingbird species tend to share similar humeral morphologies. This

result is further supported by a large effect size ($Z = 41.1$), underscoring the strength of

phylogenetic structuring in shaping bone morphology. The geometric configuration used to

quantify humeral shape reveals consistent variation in landmark positions across species,

highlighting major axes of morphological differentiation (Figure 1A–C). The distribution of

phylogenetic signal and its placement on the hummingbird phylogeny are shown in Figure

1D.

Predictors of humeral shape variation

The multivariate PGLS identified several significant predictors of humeral shape variation

(Table 1; Figure 2). Humerus size had the strongest effect on shape ($R^2 = 0.062$, $Z = 3.29$, $p =$

0.002), consistent with expectations of allometric scaling influencing skeletal morphology.

Body mass also emerged as a significant predictor ($R^2 = 0.032$, $Z = 2.25$, $p = 0.015$),

suggesting that mass-associated biomechanical demands contribute to shape differences.

Range size had a similarly significant effect ($R^2 = 0.031$, $Z = 2.51$, $p = 0.010$), indicating that

species occupying larger geographic range distributions tend to exhibit distinct humeral

configurations. Latitude, modeled as absolute distance from the equator, showed a marginal

trend ($R^2 = 0.021$, $Z = 1.35$, $p = 0.073$), hinting at potential shape shifts across environmental

gradients, though this effect did not reach significance. Maximum elevation showed no

statistically significant influence ($R^2 = 0.014$, $Z = 1.24$, $p = 0.103$). Foraging strategy was also

marginally non-significant ($R^2 = 0.040$, $Z = 1.51$, $p = 0.068$), though its directional effects

remained consistent with the results from group-wise comparisons. Overall, the model

accounted for 27% of total humeral shape variation.

Morphospace differentiation by foraging strategy

The principal component analysis (PCA), used to visualize patterns of shape variation,

revealed differences in humeral morphology associated with foraging strategies (Figure 3A).

Notably, the morphospaces of traplining and territorial species were largely contained within

the broader morphospace occupied by opportunistic species, as illustrated by the convex hulls

outlining each group in Figure 3A. This suggests that opportunistic foragers may exhibit more

generalist humeral shapes, while the other groups occupy more restricted morphological

configurations. The projected phylogenetic relationships onto the morphospace showed some

local clustering of related species, consistent with the significant phylogenetic signal detected

in the shape data. However, the overall pattern lacked clear structure, with multiple lineages

dispersed throughout the morphospace, indicating that a substantial proportion of shape

variation is explained by ecological differentiation rather than shared ancestry. The Canonical

Variate Analysis (CVA) further reinforced the role of foraging strategy in shaping humeral

morphology (Figure 3C). The first canonical axis (CV1) primarily distinguished trapliners

from both territorial and opportunistic species, while CV2 separated territorial from

opportunistic species. Classification accuracy was 71.2%, significantly exceeding random

expectations (permutation test, $p = 0.001$). The 90% confidence ellipses showed limited

overlap, particularly for trapliners, which occupied a distinct region of morphospace. This

group-level separation aligns with the differentiation detected in the multivariate regression

models.

Evolutionary rates

Traplining species exhibited approximately 2.5 times faster rates of humeral shape evolution

compared to territorial species (Figure 4), though this difference was not statistically

significant (p = 0.138). Across all foraging strategies, evolutionary rates declined through

time, consistent with an early burst followed by a deceleration of morphological change

(Figure 4, upper left panel). When modeling the association between evolutionary rates and

continuous predictors using a multivariate generalized additive model (GAM), only

geographic range size showed a statistically significant effect. The relationship was nonlinear

and smooth (edf = 2.69, F = 6.85, p < 0.001), suggesting that species with larger geographic

ranges tended to evolve humeral morphology at faster rates. In contrast, maximum elevation

(p = 0.488), body mass (p = 0.774), latitude (p = 0.466), and humerus size (p = 0.662) did not

show significant associations with evolutionary rate estimates. These results indicate that

among the ecological and morphological factors examined, geographic range size was the

most consistent predictor of evolutionary tempo, while other variables exhibited no detectable

effects under the current analytical framework.

Phenotypic integration

The modularity analysis revealed strong phenotypic integration between the proximal and

distal regions of the humerus (r-PLS = 0.899, Z = 4.22, p = 0.001). This indicates that

morphological changes in one region are closely mirrored by changes in the other, reflecting

coordinated evolution across the bone (Figure 5).

Discussion

Phylogenetic signal

Our results reveal a clear phylogenetic signal in humeral morphology among hummingbirds

(Figure 1). This finding indicates that evolutionary history continues to shape patterns of

morphological similarity across species, consistent with general trends reported for other

avian skeletal traits (Bright *et al.*, 2016; Shatkovska & Ghazali, 2020). The presence of a

strong phylogenetic structure suggests that humeral shape has evolved in a way that is partly

structured by shared ancestry throughout the radiation of hummingbirds. Rather than

indicating strict evolutionary constraint, this pattern may reflect the retention of

biomechanically effective configurations across lineages, as well as the influence of

phylogenetic inertia during adaptive divergence (Poore *et al.*, 1997; Revell *et al.*, 2008;

Holmes *et al.*, 2022).

Predictors of shape variation

Our multivariate phylogenetic regression identified several significant predictors of species

differences in humeral shape, supporting the hypothesis that ecological and morphological

factors contribute to skeletal morphology in hummingbirds. Humerus size showed the

strongest effect in the model (Table 1), consistent with a marked allometric component in

shape variation. As humerus size increased, species exhibited broader proximal epiphyses and

straighter, shorter diaphyses. This pattern likely reflects biomechanical adaptations: larger

humeri require expanded articular surfaces for muscle attachment (notably for the pectoralis

and supracoracoideus) and reduced shaft curvature to minimize bending stress under greater

loading (Poore *et al.*, 1997; Bleiweiss, 2002; Skandalis *et al.*, 2017). These shape changes are

consistent with general scaling principles observed in avian flight bones, where increased size

leads to disproportionate reinforcement of load-bearing regions (Biewener, 1982; Poore *et al.*,

1997).

Body mass also significantly predicted humeral shape. Larger-bodied species displayed more

robust humeral shapes, particularly with increased shaft thickness and straighter diaphyses.

Biomechanically, this likely compensates for the higher aerodynamic forces and inertial loads

encountered during both flapping and hovering flight in heavier hummingbirds (Altshuler *et*

al., 2004a). These findings align with prior studies documenting allometric effects on limb

bone robustness across volant birds (Nudds *et al.*, 2007).

Geographic range size was another significant ecological predictor. Species with broader

geographic distributions tended to exhibit more streamlined humeral shapes, particularly with

narrower shafts and reduced epiphyseal breadth (Figure 2). Wider ranges may be associated

with longer average movement distances, whether through dispersal, exploratory behavior, or

seasonal shifts, which in turn could favor morphologies that enhance aerodynamic efficiency

and reduce the energetic costs of sustained flight (Claramunt *et al.*, 2012; Drury *et al.*, 2024).

Alternatively, occupying a wide range of habitats may select for versatile skeletal

configurations capable of performing effectively across diverse environmental conditions.

Latitude showed a marginal but notable trend, with species distributed farther from the

equator exhibiting more robust humeri, characterized by expanded proximal epiphyses and

straighter shafts. While this pattern could reflect biomechanical adaptations to greater

environmental seasonality and structurally complex habitats, conditions that may require

enhanced power generation and maneuverability during flight (Dakin *et al.*, 2018; Sonne *et*

al., 2020), it may also relate to broader-scale movement ecology. Specifically, species at

higher latitudes often engage in longer-distance dispersal or seasonal movements, which

could select for more mechanically robust skeletal traits that support sustained flight

performance in variable conditions. In contrast, elevation did not show a significant effect,

though the trend toward shorter diaphyses and deeper epiphyses in high-elevation species is

biomechanically consistent with enhanced lift production under low air density and hypoxia

(Altshuler *et al.*, 2004a; Lim *et al.*, 2021). Similar morphological adjustments to elevation

have been reported for wing shape and muscle architecture in other avian taxa (Altshuler,

2006).

Foraging strategy showed a marginal association with humeral shape (p = 0.068), yet the

observed trends were biomechanically coherent. Traplining species tended to exhibit longer

and more slender humeri, a morphology that likely supports sustained, energy-efficient flight

over extended foraging circuits. A more elongated humerus can increase wing span and

reduce wingbeat frequency, enhancing aerodynamic efficiency and decreasing energetic costs

during long-distance movement between floral resources (Nudds *et al.*, 2007; Sievwright &

Macleod, 2012; Zusi, 2013). In contrast, territorial species exhibited shorter, thicker, and more

curved humeri, traits associated with enhanced maneuverability and force production. Shorter

bones with greater curvature can reduce the moment of inertia around the shoulder joint by

keeping mass closer to the body's center of rotation, enabling faster angular acceleration and

tighter turning radii. These features are advantageous during rapid chases, evasive maneuvers,

and sudden vertical ascents used in territorial defense (Altshuler *et al.*, 2004b; Dakin *et al.*,

2018; Sargent *et al.*, 2021). Additionally, a more robust humerus provides increased surface

area for the attachment of flight muscles such as the pectoralis and supracoracoideus, which

are critical for explosive bursts of power (Zusi, 2013). These divergent morphological

configurations reflect adaptive tuning of the flight apparatus to meet the specific aerodynamic

and behavioral demands imposed by alternative foraging strategies.

Together, these findings underscore that the humerus in hummingbirds is shaped by a

complex combination of size-related biomechanics, ecological space use, and behavioral

flight demands. The moderate explanatory power of the model (R^2 = 0.27) reflects the

multifaceted nature of morphological evolution under interacting selective pressures and

phylogenetic constraints

Morphospace differentiation by foraging strategy

Morphospace differentiation by foraging strategy

The principal component analysis (PCA) showed that species using territorial and traplining

strategies occupied partially distinct areas of morphospace. In contrast, opportunistic species

encompassed much of the variation found in the other two groups (Figure 3A). This

distribution suggests that opportunistic species tend to exhibit more generalized humeral

morphologies, whereas territorial and traplining species display more specialized shapes

adapted to their respective flight demands.

Building on these findings, the canonical variate analysis (CVA) demonstrated statistically

significant morphological separation among the three foraging groups (Figure 3C). CV1 and

CV2 effectively distinguished trapliners, territorial, and opportunistic species, confirming the

shape differentiation patterns identified in the multivariate regression. The low overlap

between groups, particularly for trapliners, reinforces that foraging strategies correspond to

distinctive humeral morphologies, likely reflecting the biomechanical demands associated

with each behavioral mode. Moreover, the distribution of species in the phylomorphospace

(Figure 3A) revealed that some distantly related species converge on similar humeral

configurations, suggesting that similar flight demands may drive morphological convergence

across the hummingbird phylogeny despite evolutionary distance. This pattern highlights the

role of ecological function in shaping skeletal form beyond phylogenetic constraints.

The reduced overlap of 90% confidence ellipses between groups in Figure 3B further

underscores the functional and evolutionary relevance of humeral shape variation. While

some convergence was observed between territorial and opportunistic species along CV2, the

separable centroids indicate that their humeral morphologies remain distinct. This

differentiation likely arises from the differing balance of maneuverability and energetic

efficiency required by each strategy: territorial species must defend floral resources through

Morphospace differentiation by foraging strategy

The principal component analysis (PCA) showed that species using territorial and traplining

strategies occupied partially distinct areas of morphospace. In contrast, opportunistic species

encompassed much of the variation found in the other two groups (Figure 3A). This

distribution suggests that opportunistic species tend to exhibit more generalized humeral

morphologies, whereas territorial and traplining species display more specialized shapes

adapted to their respective flight demands.

Building on these findings, the canonical variate analysis (CVA) demonstrated statistically

significant morphological separation among the three foraging groups (Figure 3C). CV1 and

CV2 effectively distinguished trapliners, territorial, and opportunistic species, confirming the

shape differentiation patterns identified in the multivariate regression. The low overlap

between groups, particularly for trapliners, reinforces that foraging strategies correspond to

distinctive humeral morphologies, likely reflecting the biomechanical demands associated

with each behavioral mode. Moreover, the distribution of species in the phylomorphospace

(Figure 3A) revealed that some distantly related species converge on similar humeral

configurations, suggesting that similar flight demands may drive morphological convergence

across the hummingbird phylogeny despite evolutionary distance. This pattern highlights the

role of ecological function in shaping skeletal form beyond phylogenetic constraints.

The reduced overlap of 90% confidence ellipses between groups in Figure 3B further

underscores the functional and evolutionary relevance of humeral shape variation. While

some convergence was observed between territorial and opportunistic species along CV2, the

separable centroids indicate that their humeral morphologies remain distinct. This

differentiation likely arises from the differing balance of maneuverability and energetic

efficiency required by each strategy: territorial species must defend floral resources through

rapid, aggressive flights, while opportunistic species exploit transient feeding opportunities,

potentially favoring more generalized morphologies (Tiebout, 1992; Sargent *et al.*, 2021).

Notably, this group-level morphological differentiation persisted even after accounting for

phylogenetic relatedness and allometric scaling, indicating that foraging strategy imposes a

measurable and adaptive influence on humeral shape. These results corroborate the directional

trends identified in the multivariate regression models and underscore the biomechanical

relevance of humeral variation in relation to ecological demands.

Evolutionary rates

The analysis of evolutionary rates provided a nuanced picture of how ecological factors might

shape the tempo of humeral evolution in hummingbirds. Traplining species exhibited

evolutionary rates approximately 2.5 times higher than those of territorial and opportunistic

species (Figure 4A), although this difference did not reach statistical significance ($p = 0.138$).

While some aspects of traplining, such as sustained flight over long circuits, may impose

biomechanical demands (Altshuler *et al.*, 2004b; Sargent *et al.*, 2021; Rombaut *et al.*, 2022),

it remains unclear whether this strategy is consistently more selective than the rapid,

acceleration-intensive flights involved in territorial defense. Given this uncertainty and the

lack of statistical support, we suggest interpreting these rate differences with caution.

Among continuous predictors, geographic range size was the only factor significantly

associated with evolutionary rate estimates, as revealed by generalized additive models

(Figure 4B). This nonlinear association suggests that species distributed across larger

geographic areas may experience greater morphological change in the humerus, potentially

due to exposure to diverse environmental conditions, ecological heterogeneity and flight

contexts (Webb & Gaston, 2000; Pie *et al.*, 2021). However, the underlying mechanisms

remain speculative.

Although evolutionary rates appeared to decline over time across foraging groups (Figure

4A), consistent with an early burst model, species-level estimates mapped onto the phylogeny

(Figure 1B) showed scattered pockets of elevated rates, especially near the tips. These

patterns are not necessarily contradictory: temporal trends capture average changes over

macroevolutionary time, while tip-specific estimates reflect recent shifts that may be driven

by clade-specific adaptations or environmental changes. Such dual patterns have been

observed in other studies of morphological evolution (Cooney *et al.*, 2017) and underscore the

complexity of linking evolutionary rates to ecological scenarios.

Phenotypic integration

Our analysis revealed a very high degree of morphological integration between the proximal

and distal regions of the humerus (Figure 5), supporting the prediction that these regions

evolve in a coordinated manner. The strong integration observed suggests that evolutionary

modifications in one region likely require compensatory changes in the other to preserve

flight mechanics and biomechanical efficiency (Sievwright & Macleod, 2012).

Such integration is expected in highly loaded locomotor structures, where functional

covariation and shared developmental pathways limit the potential for independent

morphological change (Klingenberg, 2008; Goswami *et al.*, 2016). While this integration may

limit the range of morphological trajectories available to hummingbirds, it likely facilitates

the evolution of coordinated adaptations to ecological and biomechanical demands. This is

consistent with broader patterns observed in vertebrate skeletal systems, where modularity

and integration frequently emerge as key features of evolutionary flexibility and constraint

(Randau & Goswami, 2017; Fabre *et al.*, 2020).

In hummingbirds, the persistence of a strongly integrated humeral structure across species

suggests that even as humeral shape diverges in response to ecological demands, internal

523 morphological coordination remains evolutionarily conserved. This integration likely imposes

524 constraints on the directions and combinations of shape changes that are viable, channeling

525 morphological evolution along coordinated trajectories rather than permitting independent

526 variation of individual regions. As a result, while ecological pressures may drive

527 diversification in overall humeral form, such change is shaped by internal structural

528 coherence, not unconstrained by it. Our findings thus highlight that phenotypic integration can

529 both enable functional refinement and limit morphological disparity, shaping how form

530 evolves under ecological and biomechanical selection.

532              **Conclusion**

533 Our results reveal that the evolution of humeral morphology in hummingbirds is driven by an

534 interplay between ecological demands, biomechanical constraints, and evolutionary history.

535 Despite strong phylogenetic signal, humeral shape is not purely conserved; instead, it exhibits

536 clear associations with foraging behavior, spatial distribution, and body size. In particular,

537 traplining species show faster evolutionary rates and more elongate humeri, suggesting that

538 energetic efficiency under sustained locomotion can drive morphological divergence.

539 Meanwhile, the influence of geographic range size on both shape and evolutionary tempo

540 points to the role of spatial ecological heterogeneity as a driver of morphological

541 diversification. At the same time, our finding of high phenotypic integration between

542 proximal and distal humeral regions highlights the developmental and functional cohesion of

543 this element, implying that adaptive evolution in one region must be coordinated with changes

544 in the other. This internal integration likely acts as both a constraint and a scaffold, channeling

545 morphological evolution along biomechanically viable pathways.

Taken together, this study underscores the value of integrating morphometrics, ecology, and

phylogenetics to understand how complex structures evolve under multiple simultaneous

pressures. It also highlights how even within a morphologically conservative lineage such as

hummingbirds, fine-scale variation in ecology and behavior can leave clear imprints on

skeletal design. Our results contribute to a broader understanding of how functionally

integrated traits evolve in response to ecological and evolutionary forces, offering a model for

investigating the dynamics of form and function across vertebrates.

554   **Acknowledgments**

We thank Steven Cardiff and Nick Mason for their support during fieldwork and specimen

access, as well as the Louisiana State University Museum of Natural Science for facilitating

access to their ornithological collections. This research was supported by funding from an

NSERC Discovery Grant (Canada), the Faculty of Sciences at Universidad de los Andes

(Bogotá, Colombia), and the Emerging Leaders in the Americas Program (Global Affairs

Canada).

562   **References**

Adams, D., Collyer, M., Kaliontzopoulou, A. & Baken, E. (2012) geomorph: Geometric
Morphometric Analyses of 2D and 3D Landmark Data. *CRAN: Contributed Packages*.

Adams, D.C. & Felice, R.N. (2014) Assessing Trait Covariation and Morphological
Integration on Phylogenies Using Evolutionary Covariance Matrices. *PLoS ONE*, **9**, e94335.

Agrawal, S., Tobalske, B.W., Anwar, Z., Luo, H., Hedrick, T.L. & Cheng, B. (2022)
Musculoskeletal wing-actuation model of hummingbirds predicts diverse effects of primary
flight muscles in hovering flight. *Proceedings of the Royal Society B: Biological Sciences*,
289.

Altshuler, D.L. (2006) The physiology and biomechanics of avian flight at high altitude.
Integrative and Comparative Biology, **46**, 62–71.

Altshuler, D.L., Dudley, R. & McGuire, J.A. (2004a) Resolution of a paradox: Hummingbird
flight at high elevation does not come without a cost. *Proceedings of the National Academy of*
Sciences, **101**, 17731–17736.

Altshuler, D.L., Stiles, F.G. & Dudley, R. (2004b) Of Hummingbirds and Helicopters:
Hovering Costs, Competitive Ability, and Foraging Strategies. *The American Naturalist*, **163**,
16–25.

Biewener, A.A. (1982) Bone Strength in Small Mammals and Bipedal Birds: Do Safety
Factors Change With Body Size? *Journal of Experimental Biology*, **98**, 289–301.

Billerman, S.M., Keeney, B.K., Rodewald, P.G. & Schulenberg, T.S. (2022) Birds of the
World [WWW Document]. *Cornell Laboratory of Ornithology, Ithaca, NY, USA.*
https://birdsoftheworld.org/bow/. URL [accessed on 2022].

Blackburn, T.M. & Ruggiero, A. (2001) Latitude, elevation and body mass variation in
Andean passerine birds. *Global Ecology and Biogeography*, **10**, 245–259.

Bleiweiss, R. (2002) Patagial complex evolution in hummingbirds and swifts (Apodiformes):
a molecular phylogenetic perspective. *Biological Journal of the Linnean Society*, **77**, 211–
219.

Blomberg, S.P., Garland, T. & Ives, A.R. (2003) Testing for phylogenetic signal in
comparative data: behavioral traits are more labile. *Evolution*, **57**, 717–745.

Bookstein, F.L. (1992) *Morphometric Tools for Landmark Data*. Cambridge University Press.

Bright, J.A., Marugán-Lobón, J., Cobb, S.N. & Rayfield, E.J. (2016) The shapes of bird beaks
are highly controlled by nondietary factors. *Proceedings of the National Academy of Sciences*,
113, 5352–5357.

Cardini, A. & Elton, S. (2007) Sample size and sampling error in geometric morphometric
studies of size and shape. *Zoomorphology*, **126**, 121–134.

Chira, A.M., Cooney, C.R., Bright, J.A., Capp, E.J.R., Hughes, E.C., Moody, C.J.A., *et al.*
(2018) Correlates of rate heterogeneity in avian ecomorphological traits. *Ecology Letters*, **21**,
1505–1514.

Claramunt, S., Derryberry, E.P., Remsen, J. V. & Brumfield, R.T. (2012) High dispersal ability
inhibits speciation in a continental radiation of passerine birds. *Proceedings of the Royal*
Society B: Biological Sciences, **279**, 1567–1574.

Cooney, C.R., Bright, J.A., Capp, E.J.R., Chira, A.M., Hughes, E.C., Moody, C.J.A., *et al.*
(2017) Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature*, **542**, 344–347.

Dakin, R., Segre, P.S., Straw, A.D. & Altshuler, D.L. (2018) Morphology, muscle capacity,
skill, and maneuvering ability in hummingbirds. *Science*, **359**, 653–657.

Dial, K.P. (1992) Avian Forelimb Muscles and Nonsteady Flight: Can Birds Fly without
Using the Muscles in Their Wings? *The Auk*, **109**, 874–885.

Drury, J.P., Clavel, J., Tobias, J.A., Rolland, J., Sheard, C. & Morlon, H. (2024) Limited
ecological opportunity influences the tempo of morphological evolution in birds. *Current*
Biology, **34**, 661-669.e4.

Dudley, R. (2002) Mechanisms and Implications of Animal Flight Maneuverability.
Integrative and Comparative Biology, **42**, 135–140.

Fabre, A.-C., Bardua, C., Bon, M., Clavel, J., Felice, R.N., Streicher, J.W., *et al.* (2020)
Metamorphosis shapes cranial diversity and rate of evolution in salamanders. *Nature Ecology*
& Evolution, **4**, 1129–1140.

Felice, R.N. & Goswami, A. (2018) Developmental origins of mosaic evolution in the avian
cranium. *Proceedings of the National Academy of Sciences*, **115**, 555–560.

Felice, R.N., Watanabe, A., Cuff, A.R., Noirault, E., Pol, D., Witmer, L.M., *et al.* (2019)
Evolutionary Integration and Modularity in the Archosaur Cranium. *Integrative and*
Comparative Biology, **59**, 371–382.

Gill, F., Donsker, D. & Rasmussen, P. (2023) IOC World Bird List 13.1. *IOC World Bird List*
Datasets.

Goswami, A., Randau, M., Polly, P.D., Weisbecker, V., Bennett, C.V., Hautier, L., *et al.* (2016)
Do Developmental Constraints and High Integration Limit the Evolution of the Marsupial
Oral Apparatus? *Integrative and Comparative Biology*, **56**, 404–415.

Hallgrímsson, B., Jamniczky, H., Young, N.M., Rolian, C., Parsons, T.E., Boughner, J.C., *et*
al. (2009) Deciphering the Palimpsest: Studying the Relationship Between Morphological
Integration and Phenotypic Covariation. *Evolutionary Biology*, **36**, 355–376.

Hallgrímsson, B., Katz, D.C., Aponte, J.D., Larson, J.R., Devine, J., Gonzalez, P.N., *et al.*
(2019) Integration and the Developmental Genetics of Allometry. *Integrative and*
Comparative Biology, **59**, 1369–1381.

Harmon, L.J., Losos, J.B., Jonathan Davies, T., Gillespie, R.G., Gittleman, J.L., Bryan
Jennings, W., *et al.* (2010) Early bursts of body size and shape evolution are rare in
comparative data. *Evolution*, no-no.

Hieronymus, T.L. (2015) Qualitative skeletal correlates of wing shape in extant birds (Aves:
Neoaves). *BMC Evolutionary Biology*, **15**, 30.

Holmes, J., Sustaita, D. & Hertel, F. (2022) Geometric morphometric analysis of the humerus
in New and Old World vultures. *Journal of Morphology*, **283**, 379–394.

Klingenberg, C.P. (2008) Morphological Integration and Developmental Modularity. *Annual
Review of Ecology, Evolution, and Systematics*, **39**, 115–132.

Klingenberg, C.P. (2016) Size, shape, and form: concepts of allometry in geometric
morphometrics. *Development Genes and Evolution*, **226**, 113–137.

Lim, M.C.W., Bi, K., Witt, C.C., Graham, C.H. & Dávalos, L.M. (2021) Pervasive Genomic
Signatures of Local Adaptation to Altitude Across Highland Specialist Andean Hummingbird
Populations. *Journal of Heredity*, **112**, 229–240.

Losos, J.B. (2011) Convergence, adaptation, and constraint. *Evolution*, **65**, 1827–1840.

McGuire, J.A., Witt, C.C., Remsen, J.V., Corl, A., Rabosky, D.L., Altshuler, D.L., *et al.*
(2014) Molecular Phylogenetics and the Diversification of Hummingbirds. *Current Biology*,
24, 910–916.

Navalón, G., Marugán-Lobón, J., Bright, J.A., Cooney, C.R. & Rayfield, E.J. (2020) The
consequences of craniofacial integration for the adaptive radiations of Darwin's finches and
Hawaiian honeycreepers. *Nature Ecology & Evolution*, **4**, 270–278.

Norberg, U.M. (1990) *Vertebrate Flight*. Springer Berlin Heidelberg, Berlin, Heidelberg.

Nudds, R.L., Dyke, G.J. & Rayner, J.M. V. (2007) Avian brachial index and wing kinematics:
putting movement back into bones. *Journal of Zoology*, **272**, 218–226.

Orkney, A., Bjarnason, A., Tronrud, B.C. & Benson, R.B.J. (2021) Patterns of skeletal
integration in birds reveal that adaptation of element shapes enables coordinated evolution
between anatomical modules. *Nature Ecology & Evolution*, **5**, 1250–1258.

Pie, M.R., Divieso, R. & Caron, F.S. (2021) Do Geographic Range Sizes Evolve Faster in
Endotherms? *Evolutionary Biology*, **48**, 286–292.

Pigot, A.L., Sheard, C., Miller, E.T., Bregman, T.P., Freeman, B.G., Roll, U., *et al.* (2020)
Macroevolutionary convergence connects morphological form to ecological function in birds.
Nature Ecology & Evolution, **4**, 230–239.

Poore, S.O., Ashcroft, A., Sánchez-Haiman, A. & Goslow, G.E. (1997) The Contractile
Properties of the M. Supracoracoideus in the Pigeon and Starling: a Case for Long-Axis
Rotation of the Humerus. *Journal of Experimental Biology*, **200**, 2987–3002.

Quintero, I. & Jetz, W. (2018) Global elevational diversity and diversification of birds.
Nature, **555**, 246–250.

R Core Team. (2024) R: A language and environment for statistical computing.

Randau, M. & Goswami, A. (2017) Unravelling intravertebral integration, modularity and
disparity in Felidae (Mammalia). *Evolution & Development*, **19**, 85–95.

Revell, L.J. (2012) Phytools: an R package for phylogenetic comparative biology (and other
things). *Methods in Ecology and Evolution*, **3**, 217–223.

Revell, L.J., Harmon, L.J. & Collar, D.C. (2008) Phylogenetic Signal, Evolutionary Process,
and Rate. *Systematic Biology*, **57**, 591–601.

Rohlf, F.J. (2015) The tps series of software. *Hystrix*, **26**, 1–4.

Rohlf, F.J. & Slice, D. (1990) Extensions of the Procrustes Method for the Optimal
Superimposition of Landmarks. *Systematic Zoology*, **39**, 40.

Rombaut, L.M.K., Capp, E.J.R., Hughes, E.C., Varley, Z.K., Beckerman, A.P., Cooper, N., *et
al.* (2022) The evolution of the traplining pollinator role in hummingbirds: specialization is
not an evolutionary dead end. *Proceedings of the Royal Society B: Biological Sciences*, **289**.

Sargent, A.J., Groom, D.J.E. & Rico-Guevara, A. (2021) Locomotion and Energetics of
Divergent Foraging Strategies in Hummingbirds: A Review. *Integrative and Comparative
Biology*, **61**, 736–748.

Sayol, F., Reijenga, B.R., Tobias, J.A. & Pigot, A.L. (2025) Ecophysical constraints on avian
adaptation and diversification. *Current Biology*, **35**, 1326-1336.e6.

Schlager, S. (2017) Morpho and Rvcg – Shape Analysis in R. In *Statistical Shape and
Deformation Analysis*. Elsevier, pp. 217–256.

Scott, G.R. & Milsom, W.K. (2006) Flying high: A theoretical analysis of the factors limiting
exercise performance in birds at altitude. *Respiratory Physiology & Neurobiology*, **154**, 284–
301.

Shatkovska, O. V. & Ghazali, M. (2020) Integration of skeletal traits in some passerines:
impact (or the lack thereof) of body mass, phylogeny, diet and habitat. *Journal of Anatomy*,
236, 274–287.

Sievwright, H. & Macleod, N. (2012) Eigensurface analysis, ecology, and modelling of
morphological adaptation in the falconiform humerus (Falconiformes: Aves). *Zoological
Journal of the Linnean Society*, **165**, 390–419.

Skandalis, D.A., Segre, P.S., Bahlman, J.W., Groom, D.J.E., Welch, K.C., Witt, C.C., *et al.*
(2017) The biomechanical origin of extreme wing allometry in hummingbirds. *Nature
Communications*, **8**, 1047.

Sonne, J., Vizentin-Bugoni, J., Maruyama, P.K., Araujo, A.C., Chávez-González, E., Coelho,
A.G., *et al.* (2020) Ecological mechanisms explaining interactions within plant–hummingbird

networks: morphological matching increases towards lower latitudes. *Proceedings of the Royal Society B: Biological Sciences*, **287**, 20192873.

Stiles, F.G. (2008) Ecomorphology and phylogeny of Hummingbirds: Divergence and Convergence in Adaptations to High Elevations. *Ornitología Neotropical*, **19**, 511–519.

Teplitsky, C., Tarka, M., Møller, A.P., Nakagawa, S., Balbontín, J., Burke, T.A., *et al.* (2014) Assessing Multivariate Constraints to Evolution across Ten Long-Term Avian Studies. *PLoS ONE*, **9**, e90444.

Tiebout, H.M. (1992) Comparative energetics of divergent foraging modes: a doubly labelled water experiment on hummingbird competition. *Animal Behaviour*, **44**, 895–906.

Tobalske, B.W. & Biewener, A.A. (2008) Contractile properties of the pigeon supracoracoideus during different modes of flight. *Journal of Experimental Biology*, **211**, 170–179.

Tobias, J.A., Sheard, C., Pigot, A.L., Devenish, A.J.M., Yang, J., Sayol, F., *et al.* (2022) AVONET: morphological, ecological and geographical data for all birds. *Ecology Letters*.

Warrick, D.R., Tobalske, B.W. & Powers, D.R. (2005) Aerodynamics of the hovering hummingbird. *Nature*, **435**, 1094–1097.

Webb, T.J. & Gaston, K.J. (2000) Geographic range size and evolutionary age in birds. *Proceedings of the Royal Society of London. Series B: Biological Sciences*, **267**, 1843–1850.

Yang, J., Yang, C., Lin, H., Lees, A.C. & Tobias, J.A. (2024) Elevational constraints on flight efficiency shape global gradients in avian wing morphology.

Zelditch, M.L., Swiderski, D.L., Sheets, H.D. & Fink, W.L. (2012) *Geometric Morphometrics for Biologists*. 2nd edn. Elsevier.

Zusi, R.L. (2013) Introduction to the Skeleton of Hummingbirds (Aves: Apodiformes, Trochilidae) in Functional and Phylogenetic Contexts. *Ornithological Monographs*, **77**, 1–94.

Tables

731 **Table 1. Summary of the phylogenetic generalized least squares (PGLS) model**

732 **evaluating humeral shape variation in relation to hummingbird species traits.** The table

733 presents the results of multivariate phylogenetic regression. Reported values include degrees

734 of freedom (df), proportion of variance explained (R²), F-statistic, effect size (Z), and

735 empirical p-values based on 1,000 permutations.

Predictor	df	R^2	F	Z	p-value	Significance
Foraging strategy	2	0.04	1.73	1.51	0.068	.
Latitude	1	0.021	1.85	1.35	0.073	.
Body mass	1	0.032	2.82	2.25	0.015	*
Humerus size	1	0.062	5.46	3.29	0.002	**
Range size	1	0.031	2.74	2.51	0.01	*
Elevation	1	0.014	1.21	1.24	0.103	

Table 2. Summary of generalized additive models relating evolutionary rate of humeral

shape to species traits in hummingbirds. This table presents the results of univariate

generalized additive models (GAMs) assessing the relationship between species-specific

evolutionary rates of humeral shape and each continuous predictor. For each variable, the

table reports the estimated degrees of freedom (edf), reference degrees of freedom (df), F-

statistic, p-value, and level of statistical significance.

Predictor	edf	df	F	p-value	Significance
Range size	2.691	3.311	6.847	< 0.001	***
Elevation	1.430	1.745	0.977	0.488	
Body mass	1.365	1.638	0.159	0.774	
Latitude	1.000	1.000	0.541	0.466	
Humerus size	1.000	1.001	0.193	0.662	

 Figures

Figure 1. Morphological evolution of the hummingbird humerus in a phylogenetic

context. (A) Anatomical landmarks used to quantify humerus shape (n = 25), shown on a

representative specimen. (B) Landmark variation across specimens illustrating the geometric

morphometric dataset. (C) Shape deformation grids for the first two principal component axes

(PC1 and PC2), showing the main axes of humeral shape variation and the anatomical regions

most contributing to shape change. (D) Null distribution of phylogenetic signal (K) in

humeral morphology across species with the observed value (red arrow) indicating significant

phylogenetic structuring (K = 0.309, p = 0.001). The circular phylogenetic tree depicts

evolutionary relationships among the 78 species analyzed. Branch colors represent species-

specific evolutionary rates of humeral shape (from lower rates in dark purple to higher rates in

bright yellow). Radial bars around the tree show normalized values of predictor variables and

continuous traits. The lower left inset shows the frequency distribution of evolutionary rates

across the tree (Log [mean rate]). Hummingbird illustrations were AI generated using Chat

GPT-4o and corrected with Adobe Illustrator.

Figure 2. Humeral shape variation in relation to species traits. Each panel depicts the

association between the first principal component (PC1) of humeral shape variation and one

of the ecological and morphological predictors included in the PGLS model. PC1 represents a

gradient from shorter, more robust humeri (lower PC1 values) to longer, more slender humeri

(higher PC1 values), as illustrated along the y-axis. Continuous predictors are displayed as

scatterplots with fitted phylogenetic regression lines and 95% confidence intervals (shaded

areas). Foraging strategy, a categorical variable, is shown as boxplots with overlaid data

points. While PC1 is shown here for illustrative purposes, the full multivariate PGLS model

considers variation across the entire shape space. Note that latitude represents the absolute

value at the centroid of the species' range.

Figure 3. Morphospace of humeral shape variation in relation to foraging strategies. (A)

Principal Component Analysis (PCA) of Procrustes-aligned humeral shapes across all

specimens, colored by foraging strategy. PC1 and PC2 account for 17.97% and 16.39% of

shape variation, respectively, capturing the major axes of morphological variation in the

dataset. Shaded convex hulls delimit the morphospace occupied by each foraging group.

Black lines represent phylogenetic relationships projected into the morphospace. (B) Mean

humeral shape variation across all specimens, visualized as a thin-plate spline deformation

grid relative to the consensus shape. Red dots represent each quantified landmark. (C)

Canonical Variate Analysis (CVA) focusing on foraging strategies, which maximizes

discrimination between groups. Dashed ellipses indicate 90% confidence intervals for each

group, and purple diamonds mark the centroid of each group. Shape diagrams adjacent to the

axes represent humeral shape changes associated with variation along CV1 and CV2.

Figure 4. Ecological and geographic predictors of evolutionary rates in humeral

morphology. (Top left) Temporal dynamics of evolutionary rates of humeral shape across

foraging strategies. Each line represents a smoothed trajectory of rate variation through

relative time (distance from the root), colored by foraging strategy. (Other panels)

Relationships between the evolutionary rate of humeral shape and individual ecological and

morphological predictor variables included in univariate regression models. Shaded areas

represent 95% confidence intervals. Note that latitude represents the absolute value at the

centroid of the species' range.

Figure 5. Phenotypic integration and modular organization of the hummingbird

humerus. (Left) Partial Least Squares (PLS) analysis of shape covariation between the

proximal and distal regions of the humerus. Each point represents a species' score on the first

PLS axis for each region. The strong positive correlation (r-PLS = 0.899, p = 0.001) indicates

high phenotypic integration, suggesting coordinated morphological evolution between

regions. (Right) Hypothesized (top) and empirically derived (bottom) modular configurations

of the humerus based on minimum covariation. Nodes represent landmarks, and colors denote

module assignment. The empirical partition largely recovers the expected proximal–distal

modular structure, reinforcing the developmental and functional differentiation of these

anatomical regions.

810