Integrated multi-omics reveals coordinated *Staphylococcus aureus* metabolic, iron transport,

and stress responses to human serum

Warasinee Mujchariyakul[1], Calum J. Walsh[1,2], Stefano Giulieri[1,2,3], Kim-Ahn LêCao[4], Timothy P.

Stinear[1,2], Benjamin P. Howden[1,2,5*], Romain Guérillot[1,2*] and Abderrahman Hachani[1,2*]

Author addresses:

[1] Department of Microbiology and Immunology, The University of Melbourne at the Peter Doherty Institute for Infection

and Immunity, Melbourne, Victoria, Australia.

[2] Centre for Pathogen Genomics, The University of Melbourne, Melbourne, Victoria, Australia.

[3] Victorian Infectious Diseases Service, Royal Melbourne Hospital, The Peter Doherty Institute for Infection and

Immunity, Melbourne, Victoria, Australia

[4] Melbourne Integrative Genomics, School of Mathematics and Statistics, University of Melbourne, Melbourne,

Victoria, 3000, Australia.

[5] Microbiological Diagnostic Unit Public Health Laboratory, Department of Microbiology and Immunology, Doherty

Institute, University of Melbourne, Melbourne, Victoria, Australia.

* Joint senior authors

Keywords: *Staphylococcus aureus*, Multi-omics data integration, Metabolism, iron acquisition, stress

responses, bacteraemia

Running title: ***Staphylococcus aureus* adaptive responses to human serum**

ABSTRACT

Bloodstream infections caused by *Staphylococcus aureus* remain a leading cause of mortality worldwide. Our understanding of *S. aureus* survival and persistence in human serum, a cell-free fraction of blood hostile for bacteria, is still limited. Here, we applied multivariate data integration methods and network analysis to a multi-omic dataset generated from five clinically prevalent *S. aureus* genotypes exposed to human serum. We observed and then confirmed using isogenic mutants, the significant roles of *gapdhB*, *sucA*, *sirA*, *sstD*, and *perR* in bacterial survival in serum. These data show that metabolic versatility in carbon source usage, iron transport and resistance to oxidative stress are interlinked and central to *S. aureus* fitness in serum, representing potential *S. aureus* vulnerabilities that could be exploited therapeutically.

IMPORTANCE

Bloodstream infections caused by *Staphylococcus aureus* are associated with mortality rates of up to 30%. However, the molecular mechanisms that enable this pathogen to survive in human serum, a nutrient-limited and immunologically hostile environment remain poorly understood. By integrating multi-omic data from five clinically relevant *S. aureus* genotypes and validating key signatures using mutants, we identified conserved genetic determinants critical for bacterial survival in serum. Our findings highlight the interconnected roles of carbohydrate metabolic flexibility, iron acquisition, and oxidative stress resistance in shaping *S. aureus* adaptation to serum. This work advances our understanding of microbial strategies to survive in the bloodstream and demonstrate the potential of multi-omic integration to uncover therapeutic vulnerabilities in bacterial pathogens.

INTRODUCTION

Staphylococcus aureus is a significant opportunistic human pathogen (Howden et al., 2023). As a commensal, *S. aureus* coexists with other members of the human microbiome in up to 30% of the human population, typically colonizing the skin or nares, with the potential to become a lethal pathogen (Räz et al., 2023). *S. aureus* bloodstream infections are greater than 10 in 100,000 people per year, with the severity of infections compounded by the widespread emergence of resistance to last-line antibiotics such as vancomycin, and mortality rates reaching 20% (Murray et al., 2022).

During bloodstream infections, professional phagocytes and non-cellular immune components, including antimicrobial peptides (AMPs), immunoglobulins and the complement system, target invasive pathogens for clearance (Guerra et al., 2017; Hornef et al., 2002). While serum does not contain immune cells, it represents a potent bactericidal environment for blood-invasive bacterial pathogens (Kuehl et al., 2020; Lopatkin et al., 2019; Mathew et al., 2023); an environment against which *S. aureus* has evolved multiple defence mechanisms (Howden et al., 2023; Kwiecinski & Horswill, 2020; Price & Boyd, 2020). The bacterium can evade humoral immunity by directly interfering with antibody functions to prevent opsonization for phagocytosis (Bear et al., 2023; de Jong et al., 2019), or by degrading complement components (Paiva et al., 2023). *S. aureus* can inactivate AMPs through proteolytic degradation (Joo & Otto, 2015), modification of its surface charge, or remodelling both the organization of its cell wall peptidoglycan and the abundance of cardiolipin in its plasma membrane. The latter two processes also contribute to *S. aureus* resistance to cell wall-targeting antibiotics (Ledger et al., 2022). The host can also impose nutritional immunity by sequestering essential nutrients, primarily trace metals like iron, zinc, and manganese to restrict bacterial growth. Bacterial pathogens have evolved mechanisms to circumvent nutritional immunity (Murdoch & Skaar, 2022). For example, *S. aureus* has multiple trace metal-sequestering strategies that also contribute to its survival in blood (Cassat & Skaar, 2012).

Elucidating the metabolic reprogramming of *S. aureus* in response to serum-derived stressors—such as nutrient limitation and acellular immune factors—can uncover fundamental metabolic pathways

that facilitate its adaptation, survival, and persistence in the bloodstream. These insights may inform

the development of targeted antimicrobial strategies and rationally designed interventions to mitigate

invasive and chronic *S. aureus* infections.

We previously reported a comprehensive multi-omics analysis of four clinically significant sepsis-

causing bacterial pathogens (*Escherichia coli*, *Klebsiella pneumoniae, S. aureus*, and *Streptococcus*

pyogenes) following their exposure to human serum (Mu et al., 2023). This work revealed a set of

conserved adaptive responses across these phylogenetically diverse species.

Notably, all four pathogens exhibited a coordinated upregulation of lipid metabolic pathways that

was associated with extensive remodelling of the cell envelope. This adaptation likely reflects a

shared osmoprotective strategy, enhancing bacterial survival in the hostile and nutrient-limited

environment of human serum. Building on our multi-omics dataset, this study examines the serum

response of *Staphylococcus aureus* across five clinically relevant clonal lineages—BPH2760 (ST1),

BPH2819 (ST5), BPH2900 (ST22), BPH2947 (ST239), and BPH2986 (ST8). The aim is to identify

a conserved set of molecular features that drives species-specific adaptation to serum during

septicaemia.

With a focused analysis on *S. aureus*, we refined here our analytical workflow. In our previous study

(Mu et al., 2023), we used a late-integration approach, independently analysing transcriptomic,

proteomic, and metabolomic datasets before mapping significant features to metabolic pathways and

Gene Ontology terms. While this revealed broad functional patterns, it lacked the resolution to detect

coordinated molecular changes across omics layers or to uncover direct cross-omic relationships. To

address these limitations, we adopted the DIABLO framework (Data Integration Analysis for

Biomarker discovery using Latent cOmponents) from the mixOmics R package, which applies a

sparse, multiblock partial least squares (PLS) method, optimised to integrate all omics datasets within

a single multivariate model. By selecting a minimal set of features that co-vary across data types and

distinguish between serum- and RPMI-exposed conditions, DIABLO enables the identification of

cross-omic interactions, reduces the statistical burden of separate analyses, and preserves the

integrated biological context of the *S. aureus* serum response (Rohart et al., 2017; Singh et al., 2019).

We further interrogated these signatures using differential expression and network-based analyses,

and identified interconnected metabolic pathways involved in serum adaptation through over-

representation analysis (ORA) and Gene Set Enrichment Analysis (GSEA).

Our integrated omic analysis show that *S. aureus* maintains optimal fitness in serum by undergoing

changes in key carbohydrate metabolic processes, iron acquisition, and resistance to oxidative stress

pathways. This combined response likely enables *S. aureus* to withstand host stressors such as

nutritional immunity and promote its survival in human serum.

RESULTS

Coordinated transcriptomic, proteomic, and metabolomic adaptation of *S. aureus* to serum

We performed multi-omics analysis to find the shared molecular responses to human serum of five *S. aureus* clinical isolates (BPH2760, BPH2819, BPH2900, BPH2947, and BPH2986) (**Figure 1A**). Principal Component Analysis (PCA), a dimensionality reduction method, was applied to each omics layer, revealing clear separation between serum- and RPMI-exposed *S. aureus*. In each omic dataset, the first principal component explained approximately 30% of the total variance, capturing the dominant condition-specific signal (**Figure 1B**). We used DIABLO to integrate the datasets into a unified predictive model of *S. aureus* exposure to human serum. DIABLO is a supervised multiblock method based on sparse Partial Least Squares Discriminant Analysis (sPLS-DA) (Rohart et al., 2017; Singh et al., 2019). DIABLO enabled simultaneous feature selection (a 'feature' is defined here as a molecular variable derived from transcriptomic, proteomic, or metabolomic datasets) and identification of highly correlated molecular signatures that distinguished *S. aureus* responses to serum versus the control condition (RPMI) (**Figure 1C-D**, **Figure S1A**). The sample plot showed that the first component captured most of the common serum responses across all five *S. aureus* strains, while the second component reflected strain-specific variation (**Figure S1B**). The final DIABLO model used 60 multi-omic features (see methods) that optimally discriminated serum from RPMI exposure on the first component (**Table S1**, **Figure S1C**).

A coordinated serum response signature was indicated by highly correlated latent components across the omics layers, including transcriptomic, proteomic and metabolomic, (Pearson's r > 0.95), (**Figure 1C**). Loading values from the DIABLO model (**Figure 1D**) show the specific contribution of individual variables, with higher absolute values marking greater discriminative power.

To explore consistent patterns between omics data, a similarity matrix was calculated across all components and selected features (**Table S1**) and visualized as a network (**Figure 1E)** (González et

al., 2012). The network revealed interactions across transcriptomic, proteomic and metabolomic layers, with features grouped by function and showing both positive and negative correlations and suggesting a coordinated response to serum. The staphylococcal pathways enriched among the discriminant and correlated features identified by DIABLO included cell wall biosynthesis, carbohydrate metabolism, iron transport, and defence against host immunity (**Figure 1D, 1E and Table 1)**. CDS and their products central to these pathways exhibited strong positive correlations: i) *glmS*, pivotal in the cell wall biosynthesis and lipid metabolism (via linoleic and oleic acids, and cholesterol metabolites); ii) *gapdhB* and *sucA*, involved in carbohydrate metabolism, and *sstD*, *scdA*, and *isdB*, contributing to iron uptake; and iii) *perR* and *sbi*, supporting oxidative stress resistance mechanisms and immune evasion, respectively. In contrast, genes involved in nucleotide metabolism (*mtaB*, *miaB*, *guaC*) were negatively correlated with carbohydrate metabolism genes (*gapdhB* and s*ucA*) and lipids (linoleic and oleic acids).

The observed patterns show that *S. aureus* undergoes substantial metabolic adaptation in response to serum, engaging uptake and utilization of serum-derived amino acids, carbohydrates, fatty acids, nucleotides, and their derivatives. Notably, the analysis revealed a marked increase in the abundance of staphylococcal iron acquisition proteins upon serum exposure. In the host environment, extracellular iron is limited and typically sequestered by high-affinity iron-binding proteins such as transferrin, lactoferrin, and haemoglobin within erythrocytes. The elevated levels of both siderophore-dependent (HtsA, SirA, SstA) and siderophore-independent (IsdB) iron acquisition proteins underscore the critical role of iron sequestration in *S. aureus* survival (**Figure 1D**). The integration of multi-omic data reveal the responses that support metabolic resilience, enabling *S. aureus* to maintain fitness while countering intrinsic serum stressors.

Figure 1. Multi-omics identification of molecular signatures of *S. aureus* adaptation to serum

(A) Schematic workflow of multi-omics analysis to predict *S. aureus* signature molecules in response

to human serum; (B) Unsupervised analysis using Principal Component Analysis (PCA). Sample plot

from PCA illustrating the consistency of sample clustering across datasets (transcriptomics,

proteomics, and metabolomics (GC-MS and LC-MS)), with samples labelled based on media conditions (SERUM and RPMI); (C) Supervised analysis using multiblock sPLS-DA. Diagnostic plot (sample scatter plot), according to media conditions (SERUM and RPMI). The numbers in each block below the diagonal represent correlation coefficients between the first components of each dataset; (D) Loading plot depicting features selected as optimally discriminatory between conditions by PLS-DA from the first component in each dataset; (E) Correlation network visualizing pairwise correlations (> |0.9|) between variables (transcripts, proteins, metabolites). Edge connections represent correlations, with edge colors indicating positive (orange) or negative (black) correlations. Node sizes are proportional to the number of interactions, and node colors represent data types: purple (transcripts), pink (proteins), green (GC-MS metabolites), and yellow (LC-MS metabolites).

Table 1. Functions of selected variables shown in **Figure 1E**

Variable Name	Function	Reference
acnA/ citB (SAUSA300_RS06765)	TCA cycle	(Cox & Nelson, 2008)
Alpha/beta hydrolase (SAUSA300_RS03435)	Putative lipase/esterase Maintenance of bacterial homeostasis and mediation of host-pathogen interactions	(Fuchs et al., 2018; Simon & Cravatt, 2010)
cntA (SAUSA300_RS13350)	Transport of multiple metals such as nickel, cobalt, zinc, iron, copper, and manganese	(Ghssein et al., 2016; Grim et al., 2017; Remy et al., 2013)
ecb (SAUSA300_RS05670)	Complement binding	(Jongerius et al., 2010)
gapdhB (SAUSA300_RS08910)	Gluconeogenesis	(Purves et al., 2010)
glpQ (SAUSA300_RS04655)	Utilization of glycerophosphodiesters (GroPC, GroPI, GroPS, GroPE, and GroPG) and release of glycerol-3-phosphate	(Jorge et al., 2017)
glnA (SAUSA300_RS06485)	Synthesis of L-glutamine from L-glutamate	(Gustafson et al., 1994)
guaC (SAUSA300_RS06695)	Conversion of GMP to IMP	(Wolff et al., 2022)
hstA (SAUSA300_RS11760)	Substrate-binding protein in siderophore-dependent iron acquisition pathways	(Beasley, Marolda, et al., 2011)
Hypothetical protein/ hisR (SAUSA300_RS10330)	Repressor in Histidine biosynthesis	(Fuchs et al., 2018)
isdB (SAUSA300_RS05535)	Substrate-binding protein in siderophore-independent iron acquisition pathways (heme–iron uptake pathway), interaction with von Willebrand factor, promotion of adherence to endothelial cells	(Alfeo et al., 2021; Mathelié-Guinlet et al., 2020)
sbi (SAUSA300_RS13060)	Complement binding	(Burman et al., 2008)
scdA (SAUSA300_RS01350)	Repair of oxidative and nitrosative damage to iron-sulphur centres and role in cell wall metabolism	(Brunskill et al., 1997; Overton et al., 2008)
serA (SAUSA300_RS09115)	Biosynthesis of L-serine	(Zhang et al., 2017)
sirA (SAUSA300_RS00605)	Substrate-binding protein in siderophore-dependent iron acquisition pathways	(Beasley, Marolda, et al., 2011)
sirTM (SAUSA300_RS01740)	Regulation of oxidative stress	(Rack et al., 2015)
sucA (SAUSA300_RS07105)	TCA cycle	(Cox & Nelson, 2008)
sucB	TCA cycle	(Cox & Nelson, 2008)

(SAUSA300_RS07100)		
sucD (SAUSA300_RS06165)	TCA cycle	(Cox & Nelson, 2008)
sstD (SAUSA300_RS03880)	Substrate-binding protein in siderophore-dependent iron acquisition pathways	(Beasley, Marolda, et al., 2011)
miaB (SAUSA300_RS06405)	Modification of tRNA	(Arragain et al., 2010)
mtaB (SAUSA300_RS08375)	Modification of tRNA	(Arragain et al., 2010)
mtlD (SAUSA300_RS11620)	Conversion of mannitol-1-P to fructose-6-P	(Murphey & Rosenblum, 1964)
perR (SAUSA300_RS10060)	Sensing of Hydrogen Peroxide (H_2O_2)	(Ji et al., 2015)
pflA (SAUSA300_RS01160)	Role in Formate metabolism	(Leibig et al., 2011)
pflB (SAUSA300_RS01155)	Role in Formate metabolism and promotion of biofilm persistence	(Bertrand et al., 2022; Leibig et al., 2011)
Polyisoprenoid-binding protein (SAUSA300_RS14560)	Role in the electron transport system	(Handa et al., 2005; Wolff et al., 2022)
Protein-ADP-ribose hydrolase (SAUSA300_RS01735)	Reversal of ADP-ribosylation in posttranslational modification	(Rack et al., 2020)
pruA/rocA (SAUSA300_RS13840)	Catabolism of proline to glutamate	(Christgen & Becker, 2019)

Network and enrichment analyses reveal *S. aureus* shared and strain-specific responses to serum

Differentially expressed (DE) genes or proteins for each strain were identified and further subjected to an intersection analysis. This analysis showed that at the transcript level, 143 genes were up-regulated, and 65 genes were down-regulated in response to serum. At the proteomic level, there were 21 up-regulated and 17 down-regulated proteins in response to serum (**Figure S2A-B, Table S2**). We performed a Pathway Enrichment Analysis (PEA), including an overrepresentation analysis (ORA) and a gene set enrichment analysis (GSEA), to reduce the complexity of data and find overrepresented pathways elicited upon exposure to serum. ORA is widely used to identify over-represented biological functions in DE gene sets, performing best when handling wide gene expression differences. Genes were filtered using criteria of log2 fold change (log2FC) > 1 and adj-P value < 0.05. To improve the granularity of PEA outputs, a GSEA analysis was also applied to capture pathways where gene expression changes were more subtle but consistently enriched, enabling the detection of significant, but lower-magnitude responses. Using the similarity in gene subsets shared between pathways, we also conducted a network analysis to find interactions among the enriched

pathways across the omics layers (**Figure 2A, Figure S3A-J, Table S3**). Consistent with previous DIABLO findings, ORA and GSEA identified four distinct functional modules comprising carbohydrate metabolism, ribosome-associated pathways, iron acquisition, and nucleotide metabolism (**Figure 2B-E**).

The carbohydrate metabolism module was composed of the glycolysis/gluconeogenesis pathway, along with other carbohydrate-processing routes such as phosphotransferase systems (PTS) (**Figure 2B-C**). Upregulated carbohydrate transporter genes, *uhpT*, *mtlA*, *fruA*, *glcC*, *lacE*, SAUSA300_RS01760, and *rbsU,* support the usage of various saccharides, including glucose, fructose, lactose, D-mannitol, and ribose (**Table S2**). Once transported and phosphorylated by PTS, carbohydrates such as glucose and fructose, are processed by glycolysis and converted into ribose-5-phosphate, glyceraldehyde-3-phosphate, and fructose-6-phosphate intermediates. These key metabolites bridge glycolysis to nucleotide and lipid metabolism (Kim et al., 2023; Richardson et al., 2015) (further detailed in **Figure 4)**, thereby enhancing the staphylococcal metabolic flexibility in serum. In contrast to other metabolic modules, our network analysis revealed that ribosome-associated pathways were largely isolated, exhibiting no direct connectivity with broader metabolic networks (**Figure 2C, Figure S3A-D**). Transcriptome profiling indicated a suppression of ribosome biogenesis following serum exposure, characterized by the downregulation of genes encoding 30S and 50S ribosomal subunit proteins (**Table S2 and S3)** Expression of the translational GTPase *typA* (also known as *bipA*), a critical regulator of ribosome assembly and translational fidelity under stress conditions (Gibbs et al., 2020; Njenga et al., 2023), was also significantly reduced (**Table S2**). These findings show that the protein biogenesis is downregulated as part of a stress-adaptive response of *S. aureus* to serum.

Among the proteomic changes induced by serum exposure, iron acquisition and nucleotide metabolism pathways showed the most substantial alterations (**Figure S1C**). Notably, there was a

marked increase in the abundance of iron transport proteins, including SirA, FhuC, SstC, SstD, HtsA,

IsdB, IsdE, and IsdI. In addition, elevated levels of ScdA and CntA (Esquilin-Lebron et al., 2021) -

proteins involved in iron-sulfur cluster binding - alongside the downregulation of *miaB* (Barrault et

al., 2024), suggest the activation of iron-sparing strategies by *S. aureus* under serum conditions

(**Tables S2** and **S3**). The significant increase in levels of ScdA and CntA, iron cluster binding proteins

concomitant with the downregulation of *miaB*, point to staphylococcal iron-sparing strategies being

switched on during serum exposure (**Table S2** and **S3**). Thus, *S. aureus* coordinates enhanced iron

uptake with the regulation of iron incorporation into metalloproteins to efficiently manage iron

availability during serum exposure (**Figure 2D**, **Figure S1B** and **S3F**).

Proteins involved in nucleotide metabolism, particularly those associated with the inosine

monophosphate (IMP) biosynthetic pathway, were prominently increased across all *S. aureus* isolates

(**Figure 2E** and **Figure S3H**). This pathway is central to the synthesis of purine nucleotides such as

adenosine monophosphate (AMP) and guanosine monophosphate (GMP), critical molecules for DNA

replication, RNA transcription, and cellular proliferation. These shifts underscore the prioritization

of iron homeostasis and nucleotide biosynthesis as key adaptive strategies employed by *S. aureus* to

support survival and growth in serum.

Figure 2. Clustering of transcriptomics and proteomics data into function-related modules

(A) Workflow overview for network analysis of enriched pathways identified from over-representation analysis (ORA) and gene set enrichment analysis (GSEA), with nodes representing pathways and edges representing pairwise similarity (Sim) between enriched terms or pathways (PW), calculated using Jaccard's similarity coefficient. (B-C) ORA and GSEA of transcriptomic data. (D-E) ORA and GSEA of proteomic data. Dot sizes and colors represent the number of strains sharing the same enriched pathways. Edge widths indicate the number of strains with interactions between

pathways, while edge lengths reflect pathway similarity. Yellow shading highlights denote

functionally related modules.

Experimental validation of predicted molecular signatures of *S. aureus* responses to human serum

To confirm findings from the multi-omic integration and pathway enrichment analyses, we assessed the fitness of *S. aureus* transposon mutants in genes involved in carbon metabolism, iron acquisition, and oxidative stress resistance, relative to their parental wild-type USA300 JE2 strain, in the presence of serum (**Figures 1E and 3A**). We initially verified the genetic relatedness of the five *S. aureus* bloodstream isolates to the laboratory strain JE2, used for this phenotypic validation. We observed a high conservation of the *S. aureus* core genome (2,222 core genes- representing 71% of JE2 genome- and 903 accessory genes). Strain JE2 was most closely related to the clinical isolate BPH2986, with both belonging to the ST8 USA300 lineage and differing by only 73 single nucleotide polymorphisms (SNPs) (**Figure S4A-D**).

The DIABLO integrative model identified *gapdhB* and *sucA* as key contributors to response to serum, exhibiting the highest loading values (**Figure 1D, Table S1**). Disruption of either gene resulted in a significant reduction in bacterial fitness in serum, highlighting their critical roles in response to serum exposure (**Figure 3B and S5A**).

GapdhB, a gluconeogenic enzyme, and SucA, an enzyme of the tricarboxylic acid (TCA) cycle, represent distinct nodes in the carbon metabolism pathway. This functional difference suggests that their contributions to fitness may be contingent upon the availability and type of carbon substrates. To test this, we compared the growth and respiration phenotypes of *gapdhB* and *sucA* mutants using a panel of 190 sole carbon sources (Biolog Inc., Hayward, CA, USA), to that of the wild type S. aureus JE2 strain. These carbon sources fall into several broad categories, including carbohydrates (mono-, di-, and oligosaccharides), carboxylic acids (short-chain fatty acids like acetate, dicarboxylic

acids like fumarate, and hydroxy acids like lactate), amino acids, alcohols, nucleosides and other organic compounds that do not fit in the aforementioned classes. The metabolic activities of JE2, *gapdhB* and *sucA* were more strongly influenced by individual carbon sources, irrespective of their broader carbon classification (**Figure S6A**).

Among the tested substrates, methyl pyruvate, a carboxylic acid and α-D-glucose, D-glucosamine, and other hexoses significantly enhanced respiration in both WT and mutant strains, whereas glyoxylic acid and itaconic acid suppressed respiration (**Figure 3C, Figure S6** and **Table S4**). Notably, methyl pyruvate supplementation partially rescued the respiratory deficiency of the *sucA* mutant, surpassing WT levels, while it failed to restore *gapdhB* mutant activity to WT levels (**Figures 3C and S6B**). These findings suggest that while both *gapdhB* and *sucA* are critical for optimal fitness in serum, the encoded enzymes play distinct roles in pyruvate catabolism. Methyl pyruvate, a membrane permeable pyruvate analog, enter cells where it is converted to acetyl-coa. In absence of *sucA,* supplementation with methyl pyruvate provides an alternative carbon source that bypasses the glycolytic block caused by the absence of conversion of 2-oxaloglutarate to succinyl-coA in the TCA cycle. GapdhB catalyzes the conversion of glyceraldehyde-3-phosphate (G3P) to 1,3-biphosphoglycerate in the gluconeogenic direction (Purves et al., 2010). In a *gapdhB* mutant, the gluconeogenic pathway is blocked, so cells cannot convert methyl pyruvate (see primary metabolism in **Figure 4**).

To investigate the contribution of iron acquisition systems to *Staphylococcus aureus* fitness in serum, we evaluated four transposon mutants (*htsA*, *sirA*, *sstD*, and *isdB*). The *htsA* and *sirA* genes encode substrate-binding proteins associated with the siderophores staphyloferrin A (SA) and staphyloferrin B (SB), respectively, both of which facilitate iron extraction from transferrin in serum (Beasley, Marolda, et al., 2011). The gene *sstD* encodes a component of the Sst system, which mediates iron uptake from host-derived catechols (Beasley, Marolda, et al., 2011). In contrast, *isdB* encodes a cell wall-anchored haemoglobin receptor within the Isd (iron-regulated surface determinant) system, a

siderophore-independent pathway that enables heme iron acquisition (**Figure 3D**). Proteomics

revealed significantly elevated levels of HtsA, SirA, SstD, and IsdB following serum exposure,

indicating their upregulation in response to iron-limited conditions (**Figure S1C**). *S. aureus* mutants

with transposon insertions in *sirA* and *sstD* had significantly impaired fitness in serum, consistent

with the importance of SB-mediated and catechol-mediated iron uptake under these conditions

(**Figure 3E**). In contrast, disruption of *htsA* did not significantly affect fitness, aligned with previous

findings that SB biosynthesis, unlike SA, is independent of TCA cycle activity and may be

preferentially utilized in iron-restricted environments (Sheldon et al., 2014). The reduced fitness

observed in the *sirA* mutant likely reflects the impaired utilization of SB as an iron capture process in

serum.

Interestingly, the *isdB* mutant exhibited no fitness defect in commercially sourced serum, likely due

to the absence of haemoglobin or heme (Pishchany et al., 2014), which are typically sequestered

within red blood cells and thus not present in commercial cell-free serum preparations. These findings

underscore the context-dependent utilization of iron acquisition systems in *S. aureus* and highlight

the critical role of siderophore- and catechol-mediated pathways in supporting staphylococcal

survival in iron-limited host environments.

Oxidative stress constitutes a fundamental component of the host immune response, primarily through

the generation of reactive oxygen species (ROS) aimed at neutralizing invading pathogens. To

counteract ROS, bacteria have evolved intricate oxidative stress response mechanisms (Imlay, 2008;

Toledano et al., 2004). Central to these systems are redox-sensitive transcriptional regulators such as

OxyR, PerR, and OhrR, which enable precise detection of specific ROS and orchestrate the

staphylococcal defensive gene expression programs (Dubbs & Mongkolsuk, 2012). In *S. aureus*,

PerR functions as a principal peroxide sensor and belongs to the ferric uptake regulator (Fur) family

of metalloregulatory proteins, which also includes Fur and Zur (Horsburgh et al., 2001). These

regulators require metal cofactors for activity, underscoring the metal-dependent nature of oxidative

stress sensing. In our analysis, PerR emerged as a key molecular determinant of oxidative stress

resistance in *S. aureus*, a finding that was substantiated through experimental validation (**Figure 3F–**

G). The *per* mutant exhibited significant fitness defect in serum supplemented with 0.25- and 0.5 mM

hydrogen peroxide (H_2O_2), with the most pronounced impairment observed after 20 hours of exposure

to 0.5 mM H_2O_2. These results underscore the critical role of PerR in facilitating *S. aureus* survival

under oxidative stress conditions, highlighting its importance in maintaining bacterial fitness in ROS-

rich host environments.

Figure 3. Phenotypic validation of molecular signatures of serum adaptation and metabolic characterization

(A) Workflow of phenotypic validation experiments to confirm the predicted molecular signatures of *S. aureus* exposed to human serum. (B) Box plots comparing Area-under-the-curve (AUC) values of *gapdhB* and *sucA* mutants to wild-type *S. aureus* (WT-JE2) grown in presence of serum, for 20 hours. (C) Fitness profiling of *S. aureus* represented by the rate of bacterial respiration in presence of different carbon sources (corresponding to metabolic activity inferred from the reduction of tetrazolium). Plots comparing Area-under-the-curve (AUC) values of *gapdhB* and *sucA* mutants to wild-type *S. aureus* (WT-JE2) grown in presence of sole carbon sources, for 24 hours. (D) Representation of iron acquisition pathways in *S. aureus.* (E) Box plots comparing AUC values of iron transport-deficient mutants (*htsA, sirA, sstD, isdB*) to WT-JE2 cultured in serum for 20 hours. (F) Box plots comparing AUC values of *perR* mutant to WT-JE2 grown in serum with different concentrations of H_2O_2 for 20 hours, demonstrating the role of *perR* in oxidative stress resistance. (G) Representative growth curves for the *S. aureus* WT- JE2 and *perR* mutant grown in serum alone (inset plot) and serum containing H_2O_2 (large plot) over 40 hours. All data points represent mean values for at least three independent biological replicates, with error bars indicating the corresponding standard deviations from the means.

Statistical significance between the AUC of WT-JE2 and isogenic mutants was determined using the **Wilcoxon paired test**. Significance levels are indicated by asterisks: $*p < 0.05$, $**p < 0.01$, and $***p < 0.001$.

Integrated metabolic responses of *S. aureus* to human serum

To conceptualize the dynamic remodelling of *S. aureus* metabolism in response to serum, we constructed a scaffolded metabolic network based on curated literature (Beasley, Cheung, et al., 2011; Carvalho et al., 2017; Gaupp et al., 2012; Halsey et al., 2017; Hines et al., 2020; Imada et al., 1977; Kenny et al., 2013; Kriegeskorte et al., 2014; Nolan et al., 2023; O'Connor et al., 2024; Parsons &

Rock, 2013; Potter et al., 2020; Qi et al., 2023; Sachla & Helmann, 2019). This network was overlaid

with the significant changes in key enzymes identified through single omics (genes upregulated in

orange, downregulated in light blue) and by DIABLO integration (genes and metabolites in red),

combined with network analyses (**Figure 4, Tables S2–S5**). This integrative approach revealed that

S. aureus mounts a broad and adaptable metabolic response to the challenges posed by the serum

environment, particularly in the utilization of diverse carbon sources including carbohydrates, amino

acids, and lipids. Multi-omics data (**Figure S7A–D**) highlighted consistent upregulation of

gluconeogenic and TCA cycle enzymes at both transcript and protein levels, indicating a metabolic

shift toward energy-efficient carbon utilization upon exposure to serum.

Carbohydrate metabolism was organized around three key metabolic nodes—glucose-6-

phosphate/fructose-6-phosphate (G6P/F6P), glyceraldehyde-3-phosphate/dihydroxyacetone

phosphate (GA3P/DHAP), and pyruvate (**Figure 4**, purple nodes embedded in

glycolysis/gluconeogenesis)—which serve as convergence points for nutrient assimilation. The

upregulation of carbohydrate transporter genes (*uhpT, mtlA, fruA, glcC, lacE, SAUSA300_RS01760,*

rbsU) in serum aligns with our findings that saccharides, including the hexose monosaccharides α-

D-glucose, D-fructose, D-mannose and disaccharides D-maltose, and sucrose, significantly enhanced

the metabolic activity of *S. aureus* (**Figures 3C and S6A**). Notably, saccharides such as G6P,

fructose, and mannose—but not glucose—have been shown to influence virulence gene expression

(Seo et al., 2021). Our transcriptomic data support this, revealing increased expression of γ-

hemolysins genes (*hlgA, hlgB, hlgC*) and immune evasion genes (*ecb, sbi, vraX*) across five *S. aureus*

strains in serum. In particular, strain BPH2986 (ST8) exhibited strong upregulation of *lukF-*

PV and *lukS-PV* genes, encoding Panton-Valentine leucocidins (**Figure S8A**). These findings suggest

that carbohydrates, such as fructose and mannose, present in serum, support the metabolic activity of

S. aureus while increasing virulence gene expression, in contrast to the glucose-rich RPMI.

Amino acids and peptides from serum further contribute to the metabolic adaptability of *S. aureus* by entering carbohydrate metabolism and the TCA cycle through key intermediates such as pyruvate, 2-oxoglutarate, and oxaloacetate. These entry points define three functional groups of amino acids: (1) pyruvate-yielding, (2) 2-oxoglutarate-yielding, and (3) oxaloacetate-yielding (**Figure 4**). Our transcriptomic data revealed significant upregulation of enzymes involved in amino acid catabolism, including *pckA* (encoding pyruvate carboxykinase) and *gudB* (encoding glutamate dehydrogenase), consistent with elevated intracellular levels of phosphoenolpyruvate (PEP) and 2-oxoglutarate (**Figure S7C–E**). Genes encoding enzymes for the degradation of glutamate (*gudB*), proline (*pruA/rocA*, *putA*), arginine (*rocD/argD*), and histidine (*hutI, hutG*) were also upregulated, indicating preferential catabolism of these amino acids during serum exposure. These findings are consistent with previous reports identifying glutamate and its precursors as major carbon sources supporting *S. aureus* proliferation (Halsey et al., 2017), positioning glutamate as a central amino acid in the adaptive response to serum.

Exposure to human serum triggered a marked upregulation of *S. aureus* genes involved in lipid and phospholipid degradation, including *glpQ* and *gehA*. This suggests an enhanced capacity in using host-derived lipids—such as cholesterol, linoleic acid (C18:2), and oleic acid (C18:1)—as nutrient sources. These lipids are metabolized into glycerol-3-phosphate (G3P), which can be then funnelled into the carbohydrate metabolism via the GA3P/DHAP node. Interestingly, our transcriptomic data revealed downregulation of *tpiA*, which encodes triosephosphate isomerase, an enzyme responsible for the reversible interconversion of GA3P and DHAP. This suggests a potential bottleneck at this metabolic junction, which may constrain glycolytic flux and influence the efficiency of carbon utilization from lipids present in serum.

Counteracting innate immune defences is essential for *S. aureus* survival during bloodstream infection. Upon exposure to human serum, *S. aureus* exhibited robust transcriptional upregulation of

a broad range of virulence-associated genes involved in immune evasion, host colonization, and

pathogenesis. These include genes encoding complement-evasion proteins (*ecb, sbi, vraX*), pore-

forming cytotoxins (*hlgA, hlgB, hlgC*), superantigen-like proteins (*ssl13, ssl14*), and the fibrinogen-

binding adhesin *clfA*. This coordinated immune evasion strategy underscores the pathogen's capacity

to rapidly adapt to immune pressures and enhance its invasive potential in the bloodstream.

Serum exposure also elicited a pronounced oxidative stress response in *S. aureus*. DIABLO analysis

identified *perR,* which encodes the peroxide-sensing transcriptional regulator PerR, as a key

molecular signature of staphylococcal adaptation to serum. Consistent with this, genes involved in

carotenoid biosynthesis —*crtM, crtN, crtQ* —were upregulated, suggesting enhanced anti-oxidative

defence through increased production of staphyloxanthin, a carotenoid pigment known to protect

against reactive oxygen species (Clauditz et al., 2006). In parallel, elevated expression of *scdA*, a gene

involved in iron-sulphur cluster repair, indicates activation of protein maintenance pathways essential

for preserving enzymatic and cellular functions under oxidative stress (Chang et al., 2006; Overton

et al., 2008).

Collectively, our integrative approach highlighted the strong adaptive capacity of *S. aureus* to

withstand the survival challenges imposed by human serum. Through the coordinated activation of

nutrient acquisition systems, oxidative stress defences, and virulence-associated mechanisms, *S.*

aureus enhances its ability to survive and persist during systemic infection.

Figure 4. Integrated metabolic and virulence adaptation of *S. aureus* in response to serum

Systems-level metabolic reprogramming of *S. aureus* upon exposure to serum, integrating single- and

multi-omics data. The depicted key metabolic pathways include glycolysis/gluconeogenesis (upright

rectangle shaded in light green), the pentose phosphate pathway (PPP), the tricarboxylic acid (TCA)

cycle (circle shaded in light green), nucleotide biosynthesis, ribosome biogenesis, thiamine (vitamin

B1) biosynthesis and transport, cell wall biosynthesis, and lipid metabolism. Host-responsive processes such as immune evasion, metal ion acquisition, and oxidative stress responses are also represented. The name of enzyme is colored to indicate changes in gene expression following serum exposure identified by single omics (compared to RPMI), with orange: increased expression, light blue: decreased expression. Variables predicted by DIABLO (multi-omics analysis) are in red, indicating their status as signature molecules of *S. aureus* serum response.

Discussion

Whilst a leading cause of lethal bacteraemia, *S. aureus* must contend with the nutrient-limited and immunologically hostile conditions of the bloodstream to establish infection and cause disease. In the bloodstream, the bacterium uses a large panoply of virulence factors to evade clearance by cellular and humoral immunity (Cheung et al., 2021; Howden et al., 2023; Soe et al., 2021). Notwithstanding the bactericidal activities of immune cells patrolling the bloodstream, serum remains a challenging environment for *S. aureus* due to the paucity of free forms of essential nutrients, often complexed to large proteins (e.g. iron ions complexed to transferrin) (Ledala et al., 2014). Several studies have shown that *S. aureus* has evolved multiple mechanisms, not related to its cytotoxicity, to overcome the nutritional constrains and acellular insults of the serum (Murdoch &Skaar, 2022). While informing on discrete *S. aureus* coping mechanisms to survive nutritional stress, these studies focused on staphylococcal genes and pathways in isolation and did not consider the concomitant adaptive responses to the serum environment. Building on our previous work showing that four phylogenetically diverse blood invasive pathogens, including *S. aureus*, adapted to serum by remodelling their cell wall envelopes (Mu et al., 2023), we employed a multi-omics framework to determine the commonly shared *S. aureus* adaptive response to serum across five distinct clinical isolates. Such approach offers a systems-level view of microbial adaptation in response to environmental stimuli, by linking molecular changes across genomic, transcriptomic, proteomic, and metabolomic layers (Fouché & Zinovyev, 2023; Maghsoudi et al., 2022; Vahabi & Michailidis,

2022). Using integrative machine learning-based multivariate analyses, differential expression

profiling, and network-based approaches, our study identified staphylococcal reprogramming in cell

wall biosynthesis, carbon metabolism, iron transport, and defence mechanisms in response to serum.

The experimental validation of the dominant outputs identified by our integrative framework

confirmed the importance of *gapdhB*, *sucA*, *sirA*, *sstD*, and *perR* genes for staphylococcal fitness in

serum (**Figure 3**). These results align with the positive co-variation patterns observed in the DIABLO

model, suggesting a functional interdependence among the pathways involving these genes (**Figure**

1C). Whilst critical for optimal fitness in serum, *gapdhB* and *sucA* have distinct functions in carbon

metabolism. Metabolic profiling revealed that hexose saccharides (e.g., α-D-glucose, D-glucosamine)

enhanced the metabolic activity in WT and *gapdhB* and *sucA* mutants, through glycolysis,

gluconeogenesis, PPP, and TCA cycle. While methyl pyruvate partially restored bacterial respiration

in the *sucA* mutant, likely via conversion to acetyl-CoA and oxaloacetate (Campbell et al., 2021), the

gapdhB mutant remained metabolically impaired, reflecting incomplete gluconeogenesis and

insufficient PPP intermediate generation (**Figures 3B** and **S6**). While highlighting the importance of

gluconeogenesis for *S. aureus* during serum exposure, these results further underscore the non-

redundant roles of GAPDH homologs (*gapdhA*, *gapdhB*) in maintaining metabolic flux (Purves et al.,

2010). Interestingly, pyruvate was shown to activate *S. aureus* regulatory networks controlling the

production of leucocidins, and thus, potentiating the cytotoxicity of *S. aureus* (Harper et al., 2018).

Conversely, exposure to glyoxylate and itaconate inhibited *S. aureus* metabolism **(Figures 3C** and

S6). The bacterium lacks a glyoxylate shunt, rendering glyoxylate toxic due to metabolic imbalance.

Moreover, glyoxylate is limited in serum, as humans lack an active glyoxylate cycle (Halsey et al.,

2017). Itaconate, an immunometabolite produced by macrophages, further exacerbates stress by

directly inhibiting glycolysis (Tomlinson et al., 2021; Tomlinson & Riquelme, 2021). These findings

highlight the potential of *S. aureus* metabolic vulnerabilities that could be exploited therapeutically.

We constructed a metabolic scaffold of *S. aureus* that integrates the predicted signature molecules

elicited/repressed in serum (**Figure 4, Table S4**). This illustrates the impressive flexibility of the

pathogen in using diverse carbon sources in serum, such as hexose saccharides, amino acids yielding

2-oxoglutarate (e.g., Glu, Pro, Arg, His), and host lipids and fatty acids (e.g., cholesterol, linoleic acid

(C18:2), and oleic acid (C18:1)) to support its fitness and virulence. Hexose saccharides have been

shown to regulate virulence gene expression (Seo et al., 2021). Peptides and amino acids present in

serum contribute to *S. aureus* metabolism by entering the glycolysis via pyruvate or intermediates of

the TCA cycle (e.g., 2-oxoglutarate, oxaloacetate). Transcriptomic analysis revealed significant

upregulation of key enzymes involved in amino acid catabolism, including pyruvate carboxykinase

(*pckA*) and glutamate dehydrogenase (*gudB*), phosphoenolpyruvate (PEP) and 2-oxoglutarate

(**Figure S7C-E**). Notably, we found a significant increase in gene expression of enzymes (e.g. *gudB*,

pruA(*rocA*)*, putA, rocD(argD), hutI,* and *hutG*) converting amino acids (e.g., Glu, Pro, Arg, His) to

2-oxoglutarate in amino acid catabolism, indicating the usage of these amino acids in serum. This is

consistent with the observation that glutamate, and the amino acids that can be converted into

glutamate (Pro, Arg, and His), provide most of the carbon required for cell proliferation (Halsey et

al., 2017). Several upregulated metabolic pathways facilitating glutamate production, driven by *pruA*

(converting proline to glutamate) and *putA* (proline dehydrogenase) underline glutamate as a key

intermediate metabolite in the adaptive response to serum. Concurrently, the downregulation of

glutamine synthesis, through decreased expression of GlnA, suggests a reduction in the utilization of

glutamate. Although excess levels of glutamate did not significantly influence the respiration or

fitness of WT-JE2 (**Table S4**), several studies have highlighted the important role of glutamate in

protecting bacteria from cell death under stress (Ramond et al., 2014; Tavares et al., 2011; Yee et al.,

2015; Yee et al., 2019).

Iron is an important determinant for *S. aureus* fitness and virulence (Ledala et al., 2014). Our analyses

further confirm the engagement of multiple iron acquisition mechanisms to cope with iron-limited

host environments such as the serum. Interestingly, mutants lacking *htsA* and *isdB*, both involved in

iron acquisition, displayed fitness levels comparable to that of the wild-type (WT-JE2), suggesting

their potential redundancy within the iron uptake systems and adaptation to the serum's low free iron

levels (**Figure 3E**). This is consistent with prior findings reporting the upregulation of iron acquisition

systems such as Isd and SirABC in *S. aureus* cultured in serum or blood (Malachowa et al., 2011).

Further studies are necessary to understand the specificity of iron-acquisition modules in iron-limited

environments in the context of bacterial metabolic reprograming during bloodstream infection.

Serum exposure also elicited the engagement of *S. aureus* defence mechanisms against acellular

immunity. Our analytical framework highlighted the upregulation of virulence-related genes (**Figures**

4 and S8). These include complement-binding proteins (e.g., *ecb*, *sbi*, *vraX*) (Haupt et al., 2008;

Jongerius et al., 2010; Yan et al., 2017), pore-forming toxins (e.g., the γ-haemolysins *hlgA*, *hlgB*,

hlgC) (Spaan et al., 2014), superantigen-like proteins (*ssl13*, *ssl14*) (Zhao et al., 2018), and the

fibrinogen-binding protein *clfA* (Higgins et al., 2006). Such responses would prime *S. aureus* against

clearance by immune cells. The identification of immunoglobulin-binding protein (Sbi) and

complement convertase inhibitor (Ecb), both enabling evasion from host antibodies and the

complement system (Jongerius et al., 2010; Smith et al., 2011) as signatures by DIABLO reinforce

the pertinence of using multi-omics analysis for the identification of bacterial responses relevant to

environmental conditions. The *S. aureus* sensing and response to host oxidative stress were also

highlighted as essential for optimal fitness in serum (Horsburgh et al., 2001; Ji et al., 2015; Rack et

al., 2015; Rack et al., 2020).

Conclusion and perspectives

This study exploited multi-omics analyses, including machine learning-based approaches, DE

analysis and network analysis, to uncover key *S. aureus* signature molecules responding to human

serum. Our findings highlight the concomitant activation of nutrient acquisition systems, oxidative

stress defences, and virulence-associated mechanisms, underlining the strong adaptive capacity of *S. aureus* to withstand the multifactorial challenges imposed by human serum. While serum is acellular and does not reflect the breadth of host-pathogen interactions during bloodstream infections, this matrix is a useful proxy for understanding how invading bacteria survive in blood. Our multi-omic analytical approach identified key staphylococcal pathways that could be further studied to continue dissecting staphylococcal adaptive responses and to identify bacterial (and host) targets for new therapeutic approaches.

Limitations of the study

The integrative analysis performed here was highly sensitive and detected subtle patterns linking different omic datasets. However, the metabolic flexibility and functional redundancy within key bacterial pathways - particularly central carbon metabolism and iron acquisition – are likely not limited to individual contributions of specific genes, thus making genes contributing to these pathways difficult to confirm experimentally using single isogenic mutants. Furthermore, the individual contributions of metabolites such as citrate, an intermediate of the TCA cycle, can repress the activity of the transcriptional activity of genes involved in iron homeostasis (Chen et al., 2024), underscoring the cross-talks between pathways that could alter single -omic readouts, and thus multi-omics analyses. This study used commercially sourced serum for phenotypic confirmation of the staphylococcal signatures. Whilst such reagent offers more control, such as batch control, the concentrations of carbohydrates, metal ions, and reactive oxygen species are unknown, which may influence the significance of phenotypic outputs.

METHODS

S. aureus genome and pangenome analyses

The complete genomes of five clinical *S. aureus* strains, BPH2760, BPH2819, BPH2900, BPH2947, BPH2986 (Mu et al., 2023), and of the strain USA300 JE2 were downloaded from the NCBI database (https://www.ncbi.nlm.nih.gov/bioproject/PRJEB29881/, https://www.ncbi.nlm.nih. gov/bioproject/PRJNA381486).

Genomic analysis was conducted using Bohra v2.3.6 (https://github.com/MDU-PHL/bohra). Bohra provided information such as Single Nucleotide Polymorphisms (SNPs), phylogeny, multi-locus sequence typing (MLST), resistome, virulome, and Pan-genome. We used Phandango to visualize the pangenome profiles (https://jameshadfield.github.io/phandango/#/).

The AureoWiki database (Fuchs et al., 2018) was used to annotate *S. aureus* genes/proteins and to define orthologous relationships, through a reciprocal best hit (RBH) search using MMseqs2 with default parameters (https://github.com/soedinglab/MMseqs2) between the five studied genomes (FAA format) and the *S. aureus* reference genome and proteome (USA300_FPR3757_NCBI_2017 and USA300_FPR3757_NCBI_UniProt_2013).

Confirmation of *S. aureus* transposon mutants' genome sequence

We used *S. aureus* transposon mutants of signature genes identified by our analytical pipeline, to verify their fitness in serum. These mutants were selected from the Nebraska transposon library (NTL), including NE275, NE400, NE547, NE665, NE1102, NE1343, and NE1767 (Fey et al., 2013). The mutants are expected to contain a precise and non-polar transposon insertions, without additional mutations. All mutants were genome sequenced on the Oxford Nanopore platform using Min114 flow cells. Base calling, demultiplexing, and adapter removal were performed in real-time using MinKNOW. Mutant NE1102 was sequenced on a GridION and processed using Guppy. Remaining mutants were sequenced on a MinION and processed using Dorado. FASTQ reads were assembled

using Trycycler version 0.5.3 (Wick et al., 2021). The assembled genomes were compared to the cognate wild-type reference genome using the Mauve whole-genome aligner plugin tool in the Geneious Prime (version 2022.2.2) (https://www.ncbi.nlm.nih.gov/pmc/ articles/PMC442156/).

S. aureus fitness in serum

We used batch-controlled human serum (Male AB plasma, Sigma Aldrich) to reduce variations arising from different donors. Human serum was clarified from lipids by centrifugation at 3,000 x g during 5 min at 4°C. Overnight cultures of *S. aureus* JE2 (WT), and NTL mutants of *gapdhB*, *sucA*, *htsA*, *sirA*, *sstD*, *isdB* and *perR* were grown from single colonies in 10 ml Brain Heart Infusion media (BHI, BD BACTO^{TM}) at 37 °C with orbital shaking. Overnight cultures were centrifuged for 5 min at 5,000 x g, and pellets were washed twice with sterile PBS. The cells were then standardized to an OD 600 nm = 0.5 (0.5 UOD) in RPMI (Gibco^{TM}). Bacteria equivalent to 0.01 UOD were inoculated in 180 μL of BHI, RPMI or 50% serum (diluted in RPMI) in clear 96-well plate (Costar), in triplicate to a final volume of 200 μL. Plates were incubated at 37°C in ClariostarPLUS microplate reader (BMG Labtech) and bacterial growth was measured every 10 min for 20 h at 600 nm.

Phenotypic analysis of *S. aureus* carbohydrate metabolism

Phenotypic profiling was performed using the Biolog Phenotype MicroArray™ (PM) system with PM1 and PM2 plates (Biolog Inc., Hayward, CA, USA), to determine the *S. aureus* fitness in presence of 190 distinct carbon sources. Overnight *S. aureus* cultures were grown in BHI at 37 °C with shaking. Bacterial suspensions were prepared according to the manufacturer's protocol for Gram-positive organisms. The inocula of *gapdhB*, *sucA* and parental WT strain were loaded into PM1 and PM2 plates and incubated at 37 °C for 24 hours in the Biolog Odin™ system, which enables the simultaneous monitoring of growth and respiration (by redox dye reduction). Metabolic activity was assessed by measuring the reduction of tetrazolium-based redox dyes, which serve as indicators of electron transport chain activity in response to specific carbon sources. Dye reduction results in the

623   formation of purple formazan, with absorbance recorded every 20 minutes at 590 nm. The area under

624   the curve (AUC) for each well was calculated to quantify metabolic activity, enabling comparative

625   analysis of carbon source utilization across conditions or strains.

Principal component analysis

Principal Component Analysis (PCA) was independently performed on each omics dataset—

transcriptomics, proteomics, GC-MS metabolomics, and LC-MS metabolomics—to explore global

data structure and identify the primary sources of variation between serum- and RPMI-exposed

samples. PCA was conducted using the *pca()* function from the mixOmics package (v6.24.0) in R

(v4.2.3), with data scaled to unit variance. The optimal number of components was determined using

the tune.pca() function, which indicated that the first three components captured approximately 50%

of the total variance across datasets.

Multi-omics data integration

We implemented the MixOmics data analytic pipeline known as Multiblock (s)PLS-DA or DIABLO,

a multivariate statistical method for classifying and separating different categorical classes (Rohart et

al., 2017; Singh et al., 2019). This supervised multi-omics integrative method adds sparsity to the

model, improving result interpretability by selecting the most relevant variables and setting others'

coefficients to zero. DIABLO treats each omics dataset as a separate "block" and identifies co-

variation patterns across these blocks, to determine interactions between, and changes by, biological

molecules under various conditions. The DIABLO model was designed with a selected design matrix

weight of 0.5 to balance discrimination between conditions and correlation across omics layers,

ensuring that feature selection prioritised both inter-omics agreement and RPMI-Serum separation.

An initial model trained with five components revealed that the first component alone was sufficient

to clearly discriminate between serum and RPMI conditions. To assess model robustness and guide

component selection, we performed leave-one-out (LOO) cross-validation, which confirmed that a

single component provided perfect classification performance. To optimize feature selection for each omics layer, we performed parameter tuning using the tune.block.splsda function. A grid of candidate values for the number of features to retain per block was defined: 20–100 (step size 20) for transcriptomics and proteomics, and 10–100 (step size 10) for both metabolomics datasets (GC-MS and LC-MS). The model was tuned using 5-fold cross-validation repeated 10 times and centroid distance as the classification metric. The number of features retained per block was: 20 for transcriptomics, 20 for proteomics, 10 for GC-MS, and 10 for LC-MS, totalling 60 features. Using these parameters, the final DIABLO model was built using block.splsda function with one component (as determined during component tuning).(Rohart et al., 2017; Singh et al., 2019)

Enrichment analysis

To identify staphylococcal pathways prominently represented following exposure to serum, feature lists derived from all omics data underwent pathway enrichment analysis, using over-representation analysis (ORA) and Gene Set Enrichment Analysis (GSEA). Pathway enrichment analysis was performed with package clusterProfiler v4.6.2. The obtained feature lists were mapped to the associated biological annotation terms (e.g. GO terms or KEGG pathways) and statistical tests for enrichment probability (enrichment P-value) were calculated to obtain the enrichment of the feature sets. In ORA, the preselected features were filtered by a criterion of log2FC > 1 and p-value < 0.05. For GSEA, all features were ranked based on their log2FC values, meaning features with the largest positive and negative expression changes are positioned at opposite ends of the ranked list, while those with smaller or no changes are located near the middle.

Network analysis

The biological relationships among the list of enriched pathways were identified by selecting the significant pathways with p.adjust < 0.05 (except for metabolomic datasets; p-value < 0.05). Further network analysis, using pairewise_termsim function in the enrichplot package in R, calculated the

pairwise similarity of the enriched terms using Jaccard's similarity coefficient or the similarity of

gene subsets sharing between pathways. Conversion to enrichment map to dataframes provided

similarity scores and number of strains sharing individual enriched pathways, represented as node

and edge, respectively, for networks' visualization.

Data availability

All codes are available on GitHub (https://github.com/warasinee/Multiomics_Analyses_2024).

Oxford Nanopore reads of all strains and mutants generated are deposited in NCBI SRA (BioProject

ID PRJNA1284587).

Acknowledgements

This research was supported by the Development and Promotion of Science and Technology Talents

Project (DPST) from the Thai Government (WM), the National Health and Medical Research Council

of Australia to BPH (GNT1196103), TPS (GNT1194325), AH, SGG, RG (GNT2018880).

References

Alfeo, M. J., Pagotto, A., Barbieri, G., Foster, T. J., Vanhoorelbeke, K., De Filippis, V., Speziale,
P., & Pietrocola, G. (2021). Staphylococcus aureus iron-regulated surface determinant
B (IsdB) protein interacts with von Willebrand factor and promotes adherence to
endothelial cells. *Sci Rep, 11*(1), 22799. https://doi.org/10.1038/s41598-021-02065-w
Arragain, S., Handelman, S. K., Forouhar, F., Wei, F. Y., Tomizawa, K., Hunt, J. F., Douki, T.,
Fontecave, M., Mulliez, E., & Atta, M. (2010). Identification of eukaryotic and
prokaryotic methylthiotransferase for biosynthesis of 2-methylthio-N6-
threonylcarbamoyladenosine in tRNA. *J Biol Chem, 285*(37), 28425-28433.
https://doi.org/10.1074/jbc.M110.106831
Barrault, M., Leclair, E., Kumeko, E. K., Jacquet, E., & Bouloc, P. (2024). Staphylococcal sRNA
IsrR downregulates methylthiotransferase MiaB under iron-deficient conditions.
Microbiol Spectr, 12(10), e0388823. https://doi.org/10.1128/spectrum.03888-23
Bear, A., Locke, T., Rowland-Jones, S., Pecetta, S., Bagnoli, F., & Darton, T. C. (2023). The
immune evasion roles of Staphylococcus aureus protein A and impact on vaccine
development. *Front Cell Infect Microbiol, 13*, 1242702.
https://doi.org/10.3389/fcimb.2023.1242702
Beasley, F. C., Cheung, J., & Heinrichs, D. E. (2011). Mutation of L-2,3-diaminopropionic acid
synthase genes blocks staphyloferrin B synthesis in Staphylococcus aureus. *BMC*
Microbiol, 11, 199. https://doi.org/10.1186/1471-2180-11-199

Beasley, F. C., Marolda, C. L., Cheung, J., Buac, S., & Heinrichs, D. E. (2011). Staphylococcus aureus transporters Hts, Sir, and Sst capture iron liberated from human transferrin by Staphyloferrin A, Staphyloferrin B, and catecholamine stress hormones, respectively, and contribute to virulence. *Infect Immun*, *79*(6), 2345-2355. https://doi.org/10.1128/iai.00117-11

Bertrand, B. P., Heim, C. E., West, S. C., Chaudhari, S. S., Ali, H., Thomas, V. C., & Kielian, T. (2022). Role of Staphylococcus aureus Formate Metabolism during Prosthetic Joint Infection. *Infect Immun*, *90*(11), e0042822. https://doi.org/10.1128/iai.00428-22

Brunskill, E. W., de Jonge, B. L. M., & Bayles, K. W. (1997). The Staphylococcus aureus scdA gene: a novel locus that affects cell division and morphogenesis. *Microbiology (Reading)*, *143 (Pt 9)*, 2877-2882. https://doi.org/10.1099/00221287-143-9-2877

Burman, J. D., Leung, E., Atkins, K. L., O'Seaghdha, M. N., Lango, L., Bernadó, P., Bagby, S., Svergun, D. I., Foster, T. J., Isenman, D. E., & van den Elsen, J. M. (2008). Interaction of human complement with Sbi, a staphylococcal immunoglobulin-binding protein: indications of a novel mechanism of complement evasion by Staphylococcus aureus. *J Biol Chem*, *283*(25), 17579-17593. https://doi.org/10.1074/jbc.M800265200

Campbell, C., Fingleton, C., Zeden, M. S., Bueno, E., Gallagher, L. A., Shinde, D., Ahn, J., Olson, H. M., Fillmore, T. L., Adkins, J. N., Razvi, F., Bayles, K. W., Fey, P. D., Thomas, V. C., Cava, F., Clair, G. C., & O'Gara, J. P. (2021). Accumulation of Succinyl Coenzyme A Perturbs the Methicillin-Resistant Staphylococcus aureus (MRSA) Succinylome and Is Associated with Increased Susceptibility to Beta-Lactam Antibiotics. *mBio*, *12*(3), e0053021. https://doi.org/10.1128/mBio.00530-21

Carvalho, S. M., de Jong, A., Kloosterman, T. G., Kuipers, O. P., & Saraiva, L. M. (2017). The Staphylococcus aureus α-Acetolactate Synthase ALS Confers Resistance to Nitrosative Stress. *Front Microbiol*, *8*, 1273. https://doi.org/10.3389/fmicb.2017.01273

Cassat, J. E., & Skaar, E. P. (2012). Metal ion acquisition in Staphylococcus aureus: overcoming nutritional immunity. *Semin Immunopathol*, *34*(2), 215-235. https://doi.org/10.1007/s00281-011-0294-4

Chang, W., Small, D. A., Toghrol, F., & Bentley, W. E. (2006). Global transcriptome analysis of Staphylococcus aureus response to hydrogen peroxide. *J Bacteriol*, *188*(4), 1648-1659. https://doi.org/10.1128/jb.188.4.1648-1659.2006

Chen, F., Zao, Q., Yang, Z., Chen, R., Pan, H., Wang, Y., Liu., H., Cao, Q., Gan, J., Liu, X, Zhang, N., Liang., H &Lan, L. (2024) PLoS Pathogen, e1012425. https://doi.org/10.1371/journal.ppat.1012425

Cheung, G. Y. C., Bae, J. S., & Otto, M. (2021). Pathogenicity and virulence of Staphylococcus aureus. *Virulence*, *12*(1), 547-569. https://doi.org/10.1080/21505594.2021.1878688

Christgen, S. L., & Becker, D. F. (2019). Role of Proline in Pathogen and Host Interactions. *Antioxid Redox Signal*, *30*(4), 683-709. https://doi.org/10.1089/ars.2017.7335

Clauditz, A., Resch, A., Wieland, K. P., Peschel, A., & Götz, F. (2006). Staphyloxanthin plays a role in the fitness of Staphylococcus aureus and its ability to cope with oxidative stress. *Infect Immun*, *74*(8), 4950-4953. https://doi.org/10.1128/iai.00204-06

Cox, M., & Nelson, D. (2008). *Lehninger Principles of Biochemistry* (5 ed.). W. H. Freeman and Company.

de Jong, N. W. M., van Kessel, K. P. M., & van Strijp, J. A. G. (2019). Immune Evasion by Staphylococcus aureus. *Microbiol Spectr*, *7*(2). https://doi.org/10.1128/microbiolspec.GPP3-0061-2019

Dubbs, J. M., & Mongkolsuk, S. (2012). Peroxide-sensing transcriptional regulators in bacteria. *J Bacteriol*, *194*(20), 5495-5503. https://doi.org/10.1128/jb.00304-12

Esquilin-Lebron, K., Dubrac, S., Barras, F., & Boyd, J. M. (2021). Bacterial Approaches for Assembling Iron-Sulfur Proteins. *mBio*, *12*(6), e0242521. https://doi.org/10.1128/mBio.02425-21

Fey, P. D., Endres, J. L., Yajjala, V. K., Widhelm, T. J., Boissy, R. J., Bose, J. L., & Bayles, K. W. (2013). A genetic resource for rapid and comprehensive phenotype screening of nonessential Staphylococcus aureus genes. *mBio*, *4*(1), e00537-00512. https://doi.org/10.1128/mBio.00537-12

Fuchs, S., Mehlan, H., Bernhardt, J., Hennig, A., Michalik, S., Surmann, K., Pané-Farré, J., Giese, A., Weiss, S., Backert, L., Herbig, A., Nieselt, K., Hecker, M., Völker, U., & Mäder, U. (2018). AureoWiki -The repository of the Staphylococcus aureus research and annotation community. *Int J Med Microbiol*, *308*(6), 558-568. https://doi.org/10.1016/j.ijmm.2017.11.011

Gaupp, R., Ledala, N., & Somerville, G. A. (2012). Staphylococcal response to oxidative stress. *Front Cell Infect Microbiol*, *2*, 33. https://doi.org/10.3389/fcimb.2012.00033

Ghssein, G., Brutesco, C., Ouerdane, L., Fojcik, C., Izaute, A., Wang, S., Hajjar, C., Lobinski, R., Lemaire, D., Richaud, P., Voulhoux, R., Espaillat, A., Cava, F., Pignol, D., Borezée-Durant, E., & Arnoux, P. (2016). Biosynthesis of a broad-spectrum nicotianamine-like metallophore in Staphylococcus aureus. *Science*, *352*(6289), 1105-1109. https://doi.org/10.1126/science.aaf1018

Gibbs, M. R., Moon, K. M., Warner, B. R., Chen, M., Bundschuh, R., Foster, L. J., & Fredrick, K. (2020). Functional Analysis of BipA in E. coli Reveals the Natural Plasticity of 50S Subunit Assembly. *J Mol Biol*, *432*(19), 5259-5272. https://doi.org/10.1016/j.jmb.2020.07.013

Grim, K. P., San Francisco, B., Radin, J. N., Brazel, E. B., Kelliher, J. L., Párraga Solórzano, P. K., Kim, P. C., McDevitt, C. A., & Kehl-Fie, T. E. (2017). The Metallophore Staphylopine Enables Staphylococcus aureus To Compete with the Host for Zinc and Overcome Nutritional Immunity. *mBio*, *8*(5). https://doi.org/10.1128/mBio.01281-17

Guerra, F. E., Borgogna, T. R., Patel, D. M., Sward, E. W., & Voyich, J. M. (2017). Epic Immune Battles of History: Neutrophils vs. Staphylococcus aureus. *Front Cell Infect Microbiol*, *7*, 286. https://doi.org/10.3389/fcimb.2017.00286

Gustafson, J., Strässle, A., Hächler, H., Kayser, F. H., & Berger-Bächi, B. (1994). The femC locus of Staphylococcus aureus required for methicillin resistance includes the glutamine synthetase operon. *J Bacteriol*, *176*(5), 1460-1467. https://doi.org/10.1128/jb.176.5.1460-1467.1994

Halsey, C. R., Lei, S., Wax, J. K., Lehman, M. K., Nuxoll, A. S., Steinke, L., Sadykov, M., Powers, R., & Fey, P. D. (2017). Amino Acid Catabolism in Staphylococcus aureus and the Function of Carbon Catabolite Repression. *mBio*, *8*(1). https://doi.org/10.1128/mBio.01434-16

Handa, N., Terada, T., Doi-Katayama, Y., Hirota, H., Tame, J. R., Park, S. Y., Kuramitsu, S., Shirouzu, M., & Yokoyama, S. (2005). Crystal structure of a novel polyisoprenoid-binding protein from Thermus thermophilus HB8. *Protein Sci*, *14*(4), 1004-1010. https://doi.org/10.1110/ps.041183305

Harper, L., Balasubramanian, D., Ohneck, E. A., Sause, W. E., Chapman, J., Mejia-Sosa, B., Lhakhang, T., Heguy, A., Tsirigos, A., Ueberheide, B., Boyd, J. M., Lun, D. S., & Torres, V. J. (2018). Staphylococcus aureus Responds to the Central Metabolite Pyruvate To Regulate Virulence. *mBio*, *9*(1). https://doi.org/10.1128/mBio.02272-17

Haupt, K., Reuter, M., van den Elsen, J., Burman, J., Hälbich, S., Richter, J., Skerka, C., & Zipfel, P. F. (2008). The Staphylococcus aureus protein Sbi acts as a complement

inhibitor and forms a tripartite complex with host complement Factor H and C3b. *PLoS*
Pathog, *4*(12), e1000250. https://doi.org/10.1371/journal.ppat.1000250

Higgins, J., Loughman, A., van Kessel, K. P., van Strijp, J. A., & Foster, T. J. (2006). Clumping
factor A of Staphylococcus aureus inhibits phagocytosis by human
polymorphonuclear leucocytes. *FEMS Microbiol Lett*, *258*(2), 290-296.
https://doi.org/10.1111/j.1574-6968.2006.00229.x

Hines, K. M., Alvarado, G., Chen, X., Gatto, C., Pokorny, A., Alonzo, F., 3rd, Wilkinson, B. J., &
Xu, L. (2020). Lipidomic and Ultrastructural Characterization of the Cell Envelope of
Staphylococcus aureus Grown in the Presence of Human Serum. *mSphere*, *5*(3).
https://doi.org/10.1128/mSphere.00339-20

Hornef, M. W., Wick, M. J., Rhen, M., & Normark, S. (2002). Bacterial strategies for overcoming
host innate and adaptive immune responses. *Nat Immunol*, *3*(11), 1033-1040.
https://doi.org/10.1038/ni1102-1033

Horsburgh, M. J., Clements, M. O., Crossley, H., Ingham, E., & Foster, S. J. (2001). PerR
controls oxidative stress resistance and iron storage proteins and is required for
virulence in Staphylococcus aureus. *Infect Immun*, *69*(6), 3744-3754.
https://doi.org/10.1128/iai.69.6.3744-3754.2001

Howden, B. P., Giulieri, S. G., Wong Fok Lung, T., Baines, S. L., Sharkey, L. K., Lee, J. Y. H.,
Hachani, A., Monk, I. R., & Stinear, T. P. (2023). Staphylococcus aureus host
interactions and adaptation. *Nat Rev Microbiol*, *21*(6), 380-395.
https://doi.org/10.1038/s41579-023-00852-y

Imada, A., Nozaki, Y., Kawashima, F., & Yoneda, M. (1977). Regulation of glucosamine
utilization in Staphylococcus aureus and Escherichia coli. *J Gen Microbiol*, *100*(2),
329-337. https://doi.org/10.1099/00221287-100-2-329

Imlay, J. A. (2008). Cellular defenses against superoxide and hydrogen peroxide. *Annu Rev*
Biochem, *77*, 755-776. https://doi.org/10.1146/annurev.biochem.77.061606.161055

Ji, C. J., Kim, J. H., Won, Y. B., Lee, Y. E., Choi, T. W., Ju, S. Y., Youn, H., Helmann, J. D., & Lee,
J. W. (2015). Staphylococcus aureus PerR Is a Hypersensitive Hydrogen Peroxide
Sensor using Iron-mediated Histidine Oxidation. *J Biol Chem*, *290*(33), 20374-20386.
https://doi.org/10.1074/jbc.M115.664961

Jongerius, I., Garcia, B. L., Geisbrecht, B. V., van Strijp, J. A. G., & Rooijakkers, S. H. M. (2010).
Convertase inhibitory properties of Staphylococcal extracellular complement-binding
protein. *J Biol Chem*, *285*(20), 14973-14979. https://doi.org/10.1074/jbc.M109.091975

Joo, H. S., & Otto, M. (2015). Mechanisms of resistance to antimicrobial peptides in
staphylococci. *Biochim Biophys Acta*, *1848*(11 Pt B), 3055-3061.
https://doi.org/10.1016/j.bbamem.2015.02.009

Jorge, A. M., Schneider, J., Unsleber, S., Göhring, N., Mayer, C., & Peschel, A. (2017).
Utilization of glycerophosphodiesters by Staphylococcus aureus. *Mol Microbiol*,
103(2), 229-241. https://doi.org/10.1111/mmi.13552

Kenny, J. G., Moran, J., Kolar, S. L., Ulanov, A., Li, Z., Shaw, L. N., Josefsson, E., & Horsburgh,
M. J. (2013). Mannitol utilisation is required for protection of Staphylococcus aureus
from human skin antimicrobial fatty acids. *PLoS One*, *8*(7), e67698.
https://doi.org/10.1371/journal.pone.0067698

Kim, J., Kim, G. L., Norambuena, J., Boyd, J. M., & Parker, D. (2023). Impact of the pentose
phosphate pathway on metabolism and pathogenesis of Staphylococcus aureus.
PLoS Pathog, *19*(7), e1011531. https://doi.org/10.1371/journal.ppat.1011531

Kriegeskorte, A., Block, D., Drescher, M., Windmüller, N., Mellmann, A., Baum, C., Neumann,
C., Lorè, N. I., Bragonzi, A., Liebau, E., Hertel, P., Seggewiss, J., Becker, K., Proctor, R.
A., Peters, G., & Kahl, B. C. (2014). Inactivation of thyA in Staphylococcus aureus

attenuates virulence and has a strong impact on metabolism and virulence gene
expression. *mBio*, *5*(4), e01447-01414. https://doi.org/10.1128/mBio.01447-14

Kuehl, R., Morata, L., Meylan, S., Mensa, J., & Soriano, A. (2020). When antibiotics fail: a
clinical and microbiological perspective on antibiotic tolerance and persistence of
Staphylococcus aureus. *J Antimicrob Chemother*, *75*(5), 1071-1086.
https://doi.org/10.1093/jac/dkz559

Kwiecinski, J. M., & Horswill, A. R. (2020). Staphylococcus aureus bloodstream infections:
pathogenesis and regulatory mechanisms. *Curr Opin Microbiol*, *53*, 51-60.
https://doi.org/10.1016/j.mib.2020.02.005

Ledala, N., Zhang, B., Seravalli, J., Powers, R., & Somerville, G. A. (2014). Influence of iron and
aeration on Staphylococcus aureus growth, metabolism, and transcription. *J
Bacteriol*, *196*(12), 2178-2189. https://doi.org/10.1128/jb.01475-14

Ledger, E. V. K., Mesnage, S., & Edwards, A. M. (2022). Human serum triggers antibiotic
tolerance in Staphylococcus aureus. *Nat Commun*, *13*(1), 2041.
https://doi.org/10.1038/s41467-022-29717-3

Leibig, M., Liebeke, M., Mader, D., Lalk, M., Peschel, A., & Götz, F. (2011). Pyruvate formate
lyase acts as a formate supplier for metabolic processes during anaerobiosis in
Staphylococcus aureus. *J Bacteriol*, *193*(4), 952-962.
https://doi.org/10.1128/jb.01161-10

Lopatkin, A. J., Stokes, J. M., Zheng, E. J., Yang, J. H., Takahashi, M. K., You, L., & Collins, J. J.
(2019). Bacterial metabolic state more accurately predicts antibiotic lethality than
growth rate. *Nat Microbiol*, *4*(12), 2109-2117. https://doi.org/10.1038/s41564-019-
0536-0

Malachowa, N., Whitney, A. R., Kobayashi, S. D., Sturdevant, D. E., Kennedy, A. D.,
Braughton, K. R., Shabb, D. W., Diep, B. A., Chambers, H. F., Otto, M., & DeLeo, F. R.
(2011). Global changes in Staphylococcus aureus gene expression in human blood.
PLoS One, *6*(4), e18617. https://doi.org/10.1371/journal.pone.0018617

Mathelié-Guinlet, M., Viela, F., Alfeo, M. J., Pietrocola, G., Speziale, P., & Dufrêne, Y. F. (2020).
Single-Molecule Analysis Demonstrates Stress-Enhanced Binding between
Staphylococcus aureus Surface Protein IsdB and Host Cell Integrins. *Nano Lett*,
20(12), 8919-8925. https://doi.org/10.1021/acs.nanolett.0c04015

Mathew, J., Sankar, P., & Varacallo, M. (2023). Physiology, Blood Plasma. In *StatPearls*.
StatPearls Publishing
Copyright © 2023, StatPearls Publishing LLC.

Mu, A., Klare, W. P., Baines, S. L., Ignatius Pang, C. N., Guérillot, R., Harbison-Price, N.,
Keller, N., Wilksch, J., Nhu, N. T. K., Phan, M. D., Keller, B., Nijagal, B., Tull, D.,
Dayalan, S., Chua, H. H. C., Skoneczny, D., Koval, J., Hachani, A., Shah, A. Walker, M.
J. (2023). Integrative omics identifies conserved and pathogen-specific responses of
sepsis-causing bacteria. *Nat Commun*, *14*(1), 1530. https://doi.org/10.1038/s41467-
023-37200-w

Murdoch, C. C., & Skaar, E. P. (2022). Nutritional immunity: the battle for nutrient metals at
the host-pathogen interface. *Nat Rev Microbiol*, *20*(11), 657-670.
https://doi.org/10.1038/s41579-022-00745-6

Murphey, W. H., & Rosenblum, E. D. (1964). MANNITOL CATABOLISM BY STAPHYLOCOCCUS
AUREUS. *Arch Biochem Biophys*, *107*, 292-297. https://doi.org/10.1016/0003-
9861(64)90332-7

Murray, C. J., Ikuta, K. S., Sharara, F., Swetschinski, L., Aguilar, G. R., Gray, A., Han, C.,
Bisignano, C., Rao, P., & Wool, E. (2022). Global burden of bacterial antimicrobial
resistance in 2019: a systematic analysis. *The lancet*, *399*(10325), 629-655.

Njenga, R., Boele, J., Öztürk, Y., & Koch, H. G. (2023). Coping with stress: How bacteria fine-
tune protein synthesis and protein transport. *J Biol Chem*, *299*(9), 105163.
https://doi.org/10.1016/j.jbc.2023.105163

Nolan, A. C., Zeden, M. S., Kviatkovski, I., Campbell, C., Urwin, L., Corrigan, R. M., Gründling,
A., & O'Gara, J. P. (2023). Purine Nucleosides Interfere with c-di-AMP Levels and Act as
Adjuvants To Re-Sensitize MRSA To β-Lactam Antibiotics. *mBio*, *14*(1), e0247822.
https://doi.org/10.1128/mbio.02478-22

O'Connor, M. J., Bartler, A. V., Ho, K. C., Zhang, K., Casas Fuentes, R. J., Melnick, B. A.,
Huffman, K. N., Hong, S. J., & Galiano, R. D. (2024). Understanding Staphylococcus
aureus in hyperglycaemia: A review of virulence factor and metabolic adaptations.
Wound Repair Regen, *32*(5), 661-670. https://doi.org/10.1111/wrr.13192

Overton, T. W., Justino, M. C., Li, Y., Baptista, J. M., Melo, A. M., Cole, J. A., & Saraiva, L. M.
(2008). Widespread distribution in pathogenic bacteria of di-iron proteins that repair
oxidative and nitrosative damage to iron-sulfur centers. *J Bacteriol*, *190*(6), 2004-2013.
https://doi.org/10.1128/jb.01733-07

Paiva, T. O., Geoghegan, J. A., & Dufrêne, Y. F. (2023). High-force catch bonds between the
Staphylococcus aureus surface protein SdrE and complement regulator factor H drive
immune evasion. *Commun Biol*, *6*(1), 302. https://doi.org/10.1038/s42003-023-04660-
1

Parsons, J. B., & Rock, C. O. (2013). Bacterial lipids: metabolism and membrane
homeostasis. *Prog Lipid Res*, *52*(3), 249-276.
https://doi.org/10.1016/j.plipres.2013.02.002

Pishchany, G., Sheldon, J. R., Dickson, C. F., Alam, M. T., Read, T. D., Gell, D. A., Heinrichs, D.
E., & Skaar, E. P. (2014). IsdB-dependent hemoglobin binding is required for
acquisition of heme by Staphylococcus aureus. *J Infect Dis*, *209*(11), 1764-1772.
https://doi.org/10.1093/infdis/jit817

Potter, A. D., Butrico, C. E., Ford, C. A., Curry, J. M., Trenary, I. A., Tummarakota, S. S.,
Hendrix, A. S., Young, J. D., & Cassat, J. E. (2020). Host nutrient milieu drives an
essential role for aspartate biosynthesis during invasive Staphylococcus aureus
infection. *Proc Natl Acad Sci U S A*, *117*(22), 12394-12401.
https://doi.org/10.1073/pnas.1922211117

Price, E. E., & Boyd, J. M. (2020). Genetic Regulation of Metal Ion Homeostasis in
Staphylococcus aureus. *Trends Microbiol*, *28*(10), 821-831.
https://doi.org/10.1016/j.tim.2020.04.004

Purves, J., Cockayne, A., Moody, P. C., & Morrissey, J. A. (2010). Comparison of the regulation,
metabolic functions, and roles in virulence of the glyceraldehyde-3-phosphate
dehydrogenase homologues gapA and gapB in Staphylococcus aureus. *Infect Immun*,
78(12), 5223-5232. https://doi.org/10.1128/iai.00762-10

Qi, Z., Sun, N., & Liu, C. (2023). Glyoxylate cycle maintains the metabolic homeostasis of
Pseudomonas aeruginosa in viable but nonculturable state induced by chlorine stress.
Microbiol Res, *270*, 127341. https://doi.org/10.1016/j.micres.2023.127341

Rack, J. G., Morra, R., Barkauskaite, E., Kraehenbuehl, R., Ariza, A., Qu, Y., Ortmayer, M.,
Leidecker, O., Cameron, D. R., Matic, I., Peleg, A. Y., Leys, D., Traven, A., & Ahel, I.
(2015). Identification of a Class of Protein ADP-Ribosylating Sirtuins in Microbial
Pathogens. *Mol Cell*, *59*(2), 309-320. https://doi.org/10.1016/j.molcel.2015.06.013

Rack, J. G. M., Palazzo, L., & Ahel, I. (2020). (ADP-ribosyl)hydrolases: structure, function, and
biology. *Genes Dev*, *34*(5-6), 263-284. https://doi.org/10.1101/gad.334631.119

Ramond, E., Gesbert, G., Rigard, M., Dairou, J., Dupuis, M., Dubail, I., Meibom, K., Henry, T.,
Barel, M., & Charbit, A. (2014). Glutamate utilization couples oxidative stress defense

and the tricarboxylic acid cycle in Francisella phagosomal escape. *PLoS Pathog*, *10*(1), e1003893. https://doi.org/10.1371/journal.ppat.1003893

Räz, A. K., Andreoni, F., Boumasmoud, M., Bergada-Pijuan, J., Schweizer, T. A., Mairpady Shambat, S., Hasse, B., Zinkernagel, A. S., & Brugger, S. D. (2023). Limited Adaptation of Staphylococcus aureus during Transition from Colonization to Invasive Infection. *Microbiol Spectr*, *11*(4), e0259021. https://doi.org/10.1128/spectrum.02590-21

Remy, L., Carrière, M., Derré-Bobillot, A., Martini, C., Sanguinetti, M., & Borezée-Durant, E. (2013). The Staphylococcus aureus Opp1 ABC transporter imports nickel and cobalt in zinc-depleted conditions and contributes to virulence. *Mol Microbiol*, *87*(4), 730-743. https://doi.org/10.1111/mmi.12126

Richardson, A. R., Somerville, G. A., & Sonenshein, A. L. (2015). Regulating the Intersection of Metabolism and Pathogenesis in Gram-positive Bacteria. *Microbiol Spectr*, *3*(3). https://doi.org/10.1128/microbiolspec.MBP-0004-2014

Rohart, F., Gautier, B., Singh, A., & KA, L. C. (2017). mixOmics: An R package for 'omics feature selection and multiple data integration. *PLoS Comput Biol*, *13*(11), e1005752. https://doi.org/10.1371/journal.pcbi.1005752

Sachla, A. J., & Helmann, J. D. (2019). A bacterial checkpoint protein for ribosome assembly moonlights as an essential metabolite-proofreading enzyme. *Nat Commun*, *10*(1), 1526. https://doi.org/10.1038/s41467-019-09508-z

Seo, K. S., Park, N., Rutter, J. K., Park, Y., Baker, C. L., Thornton, J. A., & Park, J. Y. (2021). Role of Glucose-6-Phosphate in Metabolic Adaptation of Staphylococcus aureus in Diabetes. *Microbiol Spectr*, *9*(2), e0085721. https://doi.org/10.1128/Spectrum.00857-21

Sheldon, J. R., Marolda, C. L., & Heinrichs, D. E. (2014). TCA cycle activity in Staphylococcus aureus is essential for iron-regulated synthesis of staphyloferrin A, but not staphyloferrin B: the benefit of a second citrate synthase. *Mol Microbiol*, *92*(4), 824-839. https://doi.org/10.1111/mmi.12593

Simon, G. M., & Cravatt, B. F. (2010). Activity-based proteomics of enzyme superfamilies: serine hydrolases as a case study. *J Biol Chem*, *285*(15), 11051-11055. https://doi.org/10.1074/jbc.R109.097600

Singh, A., Shannon, C. P., Gautier, B., Rohart, F., Vacher, M., Tebbutt, S. J., & KA, L. C. (2019). DIABLO: an integrative approach for identifying key molecular drivers from multi-omics assays. *Bioinformatics*, *35*(17), 3055-3062. https://doi.org/10.1093/bioinformatics/bty1054

Smith, E. J., Visai, L., Kerrigan, S. W., Speziale, P., & Foster, T. J. (2011). The Sbi protein is a multifunctional immune evasion factor of Staphylococcus aureus. *Infect Immun*, *79*(9), 3801-3809. https://doi.org/10.1128/iai.05075-11

Soe, Y. M., Bedoui, S., Stinear, T. P., & Hachani, A. (2021). Intracellular Staphylococcus aureus and host cell death pathways. *Cell Microbiol*, *23*(5), e13317. https://doi.org/10.1111/cmi.13317

Spaan, A. N., Vrieling, M., Wallet, P., Badiou, C., Reyes-Robles, T., Ohneck, E. A., Benito, Y., de Haas, C. J., Day, C. J., Jennings, M. P., Lina, G., Vandenesch, F., van Kessel, K. P., Torres, V. J., van Strijp, J. A., & Henry, T. (2014). The staphylococcal toxins γ-haemolysin AB and CB differentially target phagocytes by employing specific chemokine receptors. *Nat Commun*, *5*, 5438. https://doi.org/10.1038/ncomms6438

Tavares, A. F., Teixeira, M., Romão, C. C., Seixas, J. D., Nobre, L. S., & Saraiva, L. M. (2011). Reactive oxygen species mediate bactericidal killing elicited by carbon monoxide-releasing molecules. *J Biol Chem*, *286*(30), 26708-26717. https://doi.org/10.1074/jbc.M111.255752

Toledano, M. B., Delaunay, A., Monceau, L., & Tacnet, F. (2004). Microbial H2O2 sensors as archetypical redox signaling modules. *Trends Biochem Sci*, *29*(7), 351-357. https://doi.org/10.1016/j.tibs.2004.05.005

Tomlinson, K. L., Lung, T. W. F., Dach, F., Annavajhala, M. K., Gabryszewski, S. J., Groves, R. A., Drikic, M., Francoeur, N. J., Sridhar, S. H., Smith, M. L., Khanal, S., Britto, C. J., Sebra, R., Lewis, I., Uhlemann, A. C., Kahl, B. C., Prince, A. S., & Riquelme, S. A. (2021). Staphylococcus aureus induces an itaconate-dominated immunometabolic response that drives biofilm formation. *Nat Commun*, *12*(1), 1399. https://doi.org/10.1038/s41467-021-21718-y

Tomlinson, K. L., & Riquelme, S. A. (2021). Host-bacteria metabolic crosstalk drives S. aureus biofilm. *Microb Cell*, *8*(5), 106-107. https://doi.org/10.15698/mic2021.05.749

Wick, R. R., Judd, L. M., Cerdeira, L. T., Hawkey, J., Méric, G., Vezina, B., Wyres, K. L., & Holt, K. E. (2021). Trycycler: consensus long-read assemblies for bacterial genomes. *Genome Biol*, *22*(1), 266. https://doi.org/10.1186/s13059-021-02483-z

Wolff, D. W., Deng, Z., Bianchi-Smiraglia, A., Foley, C. E., Han, Z., Wang, X., Shen, S., Rosenberg, M. M., Moparthy, S., Yun, D. H., Chen, J., Baker, B. K., Roll, M. V., Magiera, A. J., Li, J., Hurley, E., Feltri, M. L., Cox, A. Nikiforov, M. A. (2022). Phosphorylation of guanosine monophosphate reductase triggers a GTP-dependent switch from pro- to anti-oncogenic function of EPHA4. *Cell Chem Biol*, *29*(6), 970-984.e976. https://doi.org/10.1016/j.chembiol.2022.01.007

Yan, J., Han, D., Liu, C., Gao, Y., Li, D., Liu, Y., & Yang, G. (2017). Staphylococcus aureus VraX specifically inhibits the classical pathway of complement by binding to C1q. *Mol Immunol*, *88*, 38-44. https://doi.org/10.1016/j.molimm.2017.05.018

Yee, R., Cui, P., Shi, W., Feng, J., & Zhang, Y. (2015). Genetic Screen Reveals the Role of Purine Metabolism in Staphylococcus aureus Persistence to Rifampicin. *Antibiotics (Basel)*, *4*(4), 627-642. https://doi.org/10.3390/antibiotics4040627

Yee, R., Feng, J., Wang, J., Chen, J., & Zhang, Y. (2019). Identification of Genes Regulating Cell Death in Staphylococcus aureus. *Front Microbiol*, *10*, 2199. https://doi.org/10.3389/fmicb.2019.02199

Zhang, W., Zhang, M., Gao, C., Zhang, Y., Ge, Y., Guo, S., Guo, X., Zhou, Z., Liu, Q., Zhang, Y., Ma, C., Tao, F., & Xu, P. (2017). Coupling between d-3-phosphoglycerate dehydrogenase and d-2-hydroxyglutarate dehydrogenase drives bacterial l-serine synthesis. *Proc Natl Acad Sci U S A*, *114*(36), E7574-e7582. https://doi.org/10.1073/pnas.1619034114

Zhao, Y., van Kessel, K. P. M., de Haas, C. J. C., Rogers, M. R. C., van Strijp, J. A. G., & Haas, P. A. (2018). Staphylococcal superantigen-like protein 13 activates neutrophils via formyl peptide receptor 2. *Cell Microbiol*, *20*(11), e12941. https://doi.org/10.1111/cmi.12941

A

Pathway (PW)	# Strain
PW1	5
...	...

Node

From	To	#Strain	Similarity
PW1	PW2	5	0.7
...	...	...	...

Edge

Node (Pathway)
Edge (Similarity)

Overrepresentation Analysis (ORA)

Gene Set Enrichment Analysis (GSEA)

Transcriptome

B   Carbohydrate Metabolism

Ribosome

Phosphotransferase system (PTS)

Microbial metabolism in diverse environment

Ala,Asp,Glu metabolism

PPP

Methane metabolism

Ascorbate/aldarate metabolism

Inositol phosphate metabolism

Glycolysis/Gluconeogenesis

Carbon metabolism

Biosynthesis of antibiotics

Biosynthesis of secondary metabolites

Metabolic pathways

C   Carbohydrate Metabolism

Microbial metabolism in diverse environment

Propanoate metabolism

Glycerolipid metabolism

Arg,Pro metabolism

Carbon metabolism.

Pentose and glucuronate interconversions

Metabolic pathways

PPP

Ribosome

Fructose and mannose metabolism

PTS

Ascorbate and aldarate metabolism

Nitrogen metabolism

Galactose metabolism

Proteome

D

iron-sulfur cluster binding

Iron coordination entity transport

metal ion homeostasis

iron assimilation

cofactor transport

metal ion transport

iron ion homeostasis

Metal cluster binding

Iron ion transport

Metal Ion Metabolism

'de novo' IMP biosynthetic process

IMP metabolic process

IMP biosynthetic process

E

iron ion transport

transition metal ion transport

Nucleotide Metabolism

ribonucleoside monophosphate metabolic process

'de novo' IMP biosynthetic process

IMP biosynthetic process

purine ribonucleoside monophosphate biosynthetic process

ribonucleoside monophosphate biosynthetic process

purine nucleoside monophosphate metabolic process

purine nucleoside monophosphate biosynthetic process

IMP metabolic process

purine ribonucleoside monophosphate metabolic process

Number of strain — 1 — 2 — 3 — 4 — 5

Number of strain   1 2 3 4 5

A DIABLO Results → Genes of Interest → Nebraska Library (Transposon Mutant Library) → Fitness Assay → Cellular Metabolic Characterization

Primary Metabolism

B

C

Metal Ion Transport

D Siderophore-dependent | Siderophore-independent

E

Defense Mechanism

F

G Growth in serum with 0.5 mM H_2O_2

WORKFLOW

Input
DEG
Signature molecules
Enriched pathways

Annotation Database
aureowiki TIGRFAMs, TheSEED
EggNOG COG catogory

Literature

Staphylococcus Pathways

Primary metabolism

Carbohydrate metabolism

Transporters
fruA lacD mtlA
glcA lacE mtlD
glcB lacG mtlR
glcC uhpT ulaA
rbsU SAUSA300_RS01760
SAUSA300_RS01765
SAUSA300_RS01770

Glycolysis/Gluconeogenesis

Glucose
G6P
F6P
fbp
FBP
GA3P — tpiA — DHAP
gapdh-A   gapdh-B
BPG
pgk
3PG
pgm
2PG
PEP
pckA
Pyruvate
pflA
pflB
Formate

PPP R5P tal

Nucleotide Metabolism

Purines/Pyrimidines
adk guaC pbuG pbuX nupC pdp
purD purE purN xpt Thymine

DNA metabolism/Transcription
SAUSA300_RS09325 mcsA
SAUSA300_RS14660 Thymidine
gapR glvR mgrA topA rpoA
rpoC xerD SAUSA300_RS09835

Ribosome Biogenesis
typA rimM rpmD
rpmG rpmI rplGB
rplQ rplR rplY
rpsH rpsK rpsL
rpsM rpsN rpsP rpsT

Translation/Modification
gid infA lspA miaB
mtaB secY metK groL
SAUSA300_RS05800
SAUSA300_RS07215
SAUSA300_RS09320
SAUSA300_RS11465
SAUSA300_RS01735
SAUSA300_RS01990
SAUSA300_RS12805
SAUSA300_RS13730

Cofactor metabolism

Thiamine/Vitamin B1   Lipoate
tenA thiD thiE thiM   lipA
thiV thiW thiX   lplA2
Pyridoxine/Vitamin B6   Other
pdxT pdxS   sufA

Cell wall metabolism

cwrA lrgA lrgB cidB
pbp4 sle1

Neu5Ac → ManNAc
nanT
nanA
Cell wall ← Teichoic acid ← UDP-GlcNAc
Gln Glu
glmS
GlcN-6P → GlcN-1P → GlcNAc-1P

Glycerophosphodiesters
glpQ
Acetyl-G3P → Phosphatidic acid
G3P
DHA
dhaKLM
Glycerol
Membrane lipids
gehA → FAs
Triglycerides
Acyl-PO₄ ← Exogenous FAs
Cholesterol
Linoleic acid
Oleic acid
Palmitoleic acid
FAS II
Acyl-ACP
bmrU pnbA gehE
SAUSA300_RS12575

Lipid metabolism

cidC
Acetyl-CoA → Acetyl-PO₄ → Acetate
acsA

Ala ← Cys
Ser ← Gly ← Thr
gcvPA
gcvPB
Asp ← Asn

TCA cycle
gltA (citZ)
Citrate
Aconitate
Oxaloacetate
Malate — Glyoxylate — Isocitrate
icd (citC)
Fumarate
2-oxoglutarate
Succinate sucC sucD
Succinyl-CoA
sucA sucB
gudB

acuA acuC adh1 aldA butA deoC1
SAUSA300_RS01660 SAUSA300_RS12485
SAUSA300_RS09875 SAUSA300_RS14010
SAUSA300_RS12450

Arg
His   Ornithine
hutI   argF
hutG   rocD
           pruA(rocA)
Glu   P5C
glnA   PutA
Gln   Pro

Amino acid metabolism
aimA arcC gcvH-L gcvT   L-Lysine
brnQ SAUSA300_RS00995   L-Alanine
hisR SAUSA300_RS05765   Taurine
proP SAUSA300_RS10855
serA

Defense mechanism

Complement   Pore-forming
binding   toxins
ecb   hlgA
sbi   hlgB
vraX   hlgC

Superantigen-like protein
ssl13 ssl14

Fibrinogen-binding
clfA

Others
asp23 amaP uspA1 opuD2

Virulence

Glu
Glutathione
NADP +
Antioxidants
perR ohR
sirTM katA

ROS

Antioxidant System

staphyloxanthin
crtM
crtN
crtP
crtQ

Pigmentation

DNA repair
recJ
Protein repair
scdA

Repair System

Metal ion metabolism

Metal ion uptake
cntA isdE   sirA
fhuC isdI   sstC
htsA iruO   sstD
isdB ntrA   ftnA
SAUSA300_RS12980
SAUSA300_RS13280

Staphyloferrin A
sfnaB

Staphyloferrin B
sbnB sbnF

Single-omics Analysis
DEG: Upregulation
DEG: Downregulation

DIABLO
Signature Molecules

Functional Annotation
Major group
Subgroup