Insights gained into Vascular Streak Dieback, an Emerging Disease of Woody

Ornamentals in the United States, using Metagenomic Sequencing

Kassaye H. Belay.[1,2], Sahar Abdelrazek[3], Devin Bily[4], Tashi Gyatso[4], Farhat A. Avin[5], John

Bonkowski[6], Prabha Liyanapathiranage[5,7], Lina Rodriguez Salamanca[2], Lenwood S. Heath[8],

Fulya Baysal-Gurel[5], Boris A. Vinatzer[2*]

[1] Graduate Program in Genetics, Bioinformatics, and Computational Biology, Virginia Tech,

Blacksburg, VA 24061, USA

[2] School of Plant and Environmental Sciences, Virginia Tech, Blacksburg, VA 24061, USA

[3] Department of Biomedical Sciences and Pathobiology, Virginia-Maryland College of

Veterinary Medicine, Virginia Tech, Blacksburg, VA 24061, USA

[4] Virginia Department of Agriculture and Consumer Services, 102 Governor Street, Richmond,

VA 23219, USA

[5] Otis L. Floyd Nursery Research Center, Tennessee State University, McMinnville, TN 37110,

USA

[6] Department of Botany and Plant Pathology, Purdue University, West Lafayette, IN 47907,

USA

[7] Tennessee Department of Agriculture, Ellington Agricultural Center, Nashville, TN 37220,

USA

[8] Department of Computer Science, Virginia Tech, Blacksburg, VA 24061, USA

*Corresponding author: vinatzer@vt.edu ORCID: 0000-0003-4612-225X

Summary

● Woody ornamentals are an integral part of home gardens and urban landscapes and are

key to the economic vitality of the nursery industry. Movement of plants infected with

emerging pathogens can quickly lead to wide geographic spread of new diseases

impacting ecosystem health and leading to economic losses.

● Using metagenomic sequencing, we investigated the microbial species associated with
vascular streak dieback (VSD), an emerging disease of woody ornamentals in the United
States, analyzing 106 samples obtained from 34 different species in seven states.
● The fastidious fungal pathogen *Ceratobasidium* sp. (*Csp*), previously found to be
associated with VSD, was the only putative pathogen consistently identified. Assembly
of 17 high quality draft genomes followed by phylogenetic and pangenome analyses
revealed *Csp* to form a tight genetic cluster closely related to, yet distinct in gene
content, from genomes of *Ceratobasidium theobromae*, a pathogen of cacao, avocado,
and cassava in Southeast Asia.
● These results reveal the unparalleled ability of metagenomics for identification and
characterization of fastidious fungal pathogens and suggest the recent introduction of a
single genetic lineage of *Csp* into the U.S., possibly from Southeast Asia, that is
characterized by a wide host range and that quickly spread geographically.
43

44   **Key words**
45   Emerging disease, metagenomic sequencing, woody ornamentals, phylogenetics, pangenome
46   analysis
47
48

Introduction
Plant pathogenic fungi in the genus *Ceratobasidium* are known to cause severe vascular diseases
in Southeast Asia, notably vascular streak dieback (VSD) in cacao (*Theobroma cacao*) and
avocado (*Persea americana*) (Keane et al., 1972; Guest & Keane, 2007) and witches' broom
disease in cassava (*Manihot esculenta*) (Leiva et al., 2023).
Recently, VSD has also been reported in woody ornamental species across the United
States. A *Ceratobasidium* sp. (*Csp*) was first detected in eastern redbud (*Cercis canadensis* L.)
with VSD symptoms in Tennessee in 2019, while similar symptoms were observed in maple
(*Acer* spp.) in North Carolina as early as 2018 (McClellan, 2023; Liyanapathiranage et al.,
2024). To date, the disease has been reported in over 12 states and 50 woody plant genera, with
the highest incidence in eastern redbud (*Cercis canadensis* L.), followed by maple (*Acer* spp.)
and dogwood (*Cornus* spp.) (Lauderdale, 2023; Baysal-Gurel & Liyanapathiranage; 2024, Bily

et al., 2025; Liyanapathiranage et al., 2024). Woody ornamentals play a crucial role in landscaping, habitat restoration, and ecological conservation, and their cultivation and sale significantly contribute to the economic viability of the U.S. nursery industry. The continued spread of VSD is thus a major concern from both an ecological and an economic perspective (Liyanage et al., 2025).

Ceratobasidium sp. detection in the U.S. is currently based on a PCR assay that uses primers specific to the *Csp* ITS region (Avin et al., 2025). Phylogenetic reconstruction has been performed using the ITS, LSU, ATP6, TEF1, and RPB2 genes and revealed a close genetic relationship between *Csp* in the U.S. and *C. theobromae* (*Ct*) from cacao and cassava in Southeast Asia but since *Csp* sequences form a distinct clade, they may represent a separate, previously unrecognized species (Liyanapathiranage et al., 2024). Also, the 100% sequence identity in the ITS region between the U.S. *Csp* sequences and *Ct* sequences from Japanese honeysuckle (*Lonicera japonica*) in China suggest that *Csp* in the U.S may be more closely related to *Ct* in China than to *Ct* in Southeast Asia (Liyanapathiranage et al., 2024). Koch's postulates have not been completed because of the fastidious nature of *Csp*. Mycelia grow out of plant tissue but, so far, is recalcitrant to sub-culturing. Therefore, it cannot be excluded that other biotic or abiotic factors contribute to the disease.

The genera *Ceratobasidium* and *Rhizoctonia* are closely related basidiomycete fungi within the family *Ceratobasidiaceae*, exhibiting overlapping morphological and ecological traits that complicate their differentiation (Moliszewska et al., 2023). Historically, many *Ceratobasidium* species were identified as the sexual (teleomorphic) stage of *Rhizoctonia* (González García et al., 2006; Gónzalez et al., 2016). Recent phylogenetic investigations have further blurred genus boundaries, revealing that some species formerly classified under *Ceratobasidium* are more accurately placed within *Rhizoctonia* (Moliszewska et al., 2023; Gónzalez et al., 2016). Regarding pathogenicity, *Rhizoctonia* is notorious for causing a broad range of diseases such as damping-off and root rot (Moliszewska et al., 2023). In contrast, *Ceratobasidium* species generally adopt a more specialized biotrophic lifestyle, maintaining a more intimate and often less aggressive interaction with their hosts (Keane et al., 1972; Guest & Keane, 2007).

Genomic studies of *Ct* have greatly advanced understanding of this pathogen's biology and evolution (Ali et al., 2019; Gil-Ordóñez et al., 2024). These studies revealed compact

genome sizes of approximately 33–34 Mb encoding 9,000–10,000 genes. Comparative genomic

analyses identified around 800 genes unique to *Ct* compared to its closest relatives potentially

underpinning *Ct*'s vascular pathogenicity, including extensive secretomes, plant cell wall

degrading enzymes, and virulence-associated factors such as toxin biosynthesis and effector

proteins (Ali et al., 2019; Gil-Ordóñez et al., 2024).

A comprehensive understanding of the emergence, distribution, transmission, and

evolutionary dynamics of emerging plant diseases, such as VSD, is essential for developing

effective disease containment or management strategies (Rasmussen and Grünwald, 2021;

Woolhouse et al., 2005). Today, for culturable pathogens, this information can be obtained by

whole genome sequencing (WGS) with long-read platforms, like PacBio and Oxford Nanopore

Technologies, which facilitate accurate assembly and characterization of complex fungal

genomes (Wang et al., 2021; Sun et al., 2022). However, WGS becomes challenging or

impossible to use in the case of fastidious or non-culturable pathogens, like *Csp*. Metagenomic

sequencing, *i.e.*, the direct sequencing and analysis of all genetic material obtained from an

environmental or host sample without requiring cultivation (Handelsman et al., 2004), can

potentially provide the same information as WGS. It can even identify mixed infections.

However, metagenomics is faced with several challenges. In the case of plant samples,

overabundance of host sequences limits the number of pathogen sequences that can be obtained.

Also, the presence of many other microbes may make it challenging to separate pathogen

sequences from non-pathogen sequences, which limits the possibility to obtain a complete

pathogen genome with minimal contamination. Therefore, metagenome-assembled genomes

(MAGs) of plant pathogens have so far mainly been obtained for bacteria (Mande et al., 2012;

Meyer et al., 2022; Johnson et al., 2022; Roman-Reyna and Crandall., 2024; Abdelrazek et al.,

2025).

To start answering the open questions about VSD on woody ornamentals in the U.S.,

here, we sequenced 106 samples from plants with VSD symptoms that had been confirmed to be

infected with *Csp* with a PCR assay (Liyanapathiranage et al., 2024; Avin et al., 2025),

including 61 symptomatic vascular tissue samples and 45 mycelia from 34 different woody plant

species collected in seven U.S. states. The main objectives were to (i) perform a survey of

bacterial, fungal, and oomycete pathogens present in symptomatic vascular tissues to identify

the microbial diversity in VSD symptomatic tissue, including the detection of *Csp* and other co-

existing microorganisms, (ii) assemble *Csp* genomes from plant metagenomic and mycelium
samples to elucidate evolutionary relationships among *Csp* sequences from the U.S. and with *Ct*
sequences from Southeast Asia, and (iii) assess differences in gene content, including predicted
virulence genes, between *Csp* and *Ct* to start unraveling potential adaptation of *Csp* to woody
ornamentals and the various regional climates in North America.

Methods

Plant Sample Collection, Storage, and Mycelium Growth
Survey sites included wholesale and retail nurseries, along with landscape sites with newly
planted trees. Symptomatic native plants located near landscape trees or nursery stock at each
survey site were also sampled. One to three plant samples were chosen from a block, group, or
cultivar of plants that best represented the symptoms observed, which included leaf scorching,
leaf chlorosis, stunting, wilt, vascular discoloration, and dieback. Symptomatic plant parts or the
entire plant with roots were placed in sealed plastic bags at room temperature and were either
shipped overnight or delivered the same day to the laboratory.

Upon receipt, the foliage was removed, stems were cut into 15 to 20 cm pieces, surface
sterilized in 70% ethanol and incubated at ≤90% humidity in a sealed plastic bag with a moist
paper towel at $22 \pm 2°C$ in the dark for 24 to 96 h before 0.1 g of discolored xylem tissue was
cut out lengthwise with a scalpel or sliced into discs from smaller, debarked shoots. If parts of
the plant were dead, any discolored living tissue at the margin between dead or wilted parts was
selected, making sure that any tissue colonized by canker-causing pathogens was excluded. If
the sample lacked dead tissue or vascular discoloration, four pieces (0.025g each) of xylem
tissue would be selected every 5 cm and combined, making sure to select from multiple growth
rings. **Supplementary Table 1** provides comprehensive information including host plant
species, geographic collection locations, and isolation dates.

DNA extraction
Plant tissue was macerated in CD1 buffer with a BioSpec MiniBeadbeater-16 (Bartlesville, OK)
or MP BiomedicaFastPrep-24™ 5G (Solon, OH), followed by DNA extraction using the
Qiagen DNeasy Plant Pro Kit (Germantown, MD). Additionally, DNA was extracted from

mycelia from 45 samples that had been harvested from a Petri dish or incubated cut stems using
the same kit. If the stem had sufficient moisture content, culturing from vascular tissue was
performed, where a 4 to 5 mm piece of discolored xylem tissue was surface sterilized with 2.5%
NaOCl, again with 70% ethanol, and air-dried before being placed on water agar medium (pH
7.1) and incubated in the dark at $22 \pm 2°C$ for as long as 15 d, where 0.1 g of *Rhizoctonia*-like
mycelia was swabbed with a sterile needle. Mycelia were also harvested under a dissection
microscope with a sterile needle from cut stems incubated for as long as 96 h and placed into an
extraction tube with CD1 buffer.
162

Computational Resources

164 All bioinformatic analyses were performed on the OWL's Nest and TinkerCliffs high-
165 performance computing clusters (Advanced Research Computing, ARC) at Virginia Tech.
166 Analyses typically used 20–40 CPUs and up to 256 GB of RAM per job, and all major software
167 tools and pipelines were installed and managed through centrally maintained conda
168 environments.
169

Metagenomic sequencing

Long-read sequencing was performed on an Oxford Nanopore Technologies (ONT)
PromethION sequencer. Sequencing libraries were prepared following the manufacturer's
protocol for SQK-NBD114.24, provided by ONT. Briefly, nucleic acid extracted from each
sample went through an FFPE repair step and then purified using AMPure XP beads (Beckman
Coulter Life Sciences, Indianapolis, IN). Following purification, sequencing adapters were
ligated to each sample, then subjected to a final purification step using AMPure XP beads. For
each sequencing library that was prepared, barcoded and adapter-ligated samples were pooled
together and eluted in 25 µl of ONT elution buffer. PromethION flow cells (FLO-PRO114M)
were primed, and the sequencing library concentration and quality were assessed using Qubit
and TapeStation, then adjusted to meet PromethION requirements before being loaded into the
primed flow cells for sequencing. The resulting FASTQ files from each flow cell were
concatenated into a single file per sample for downstream analysis. All reads that passed ONT
quality control (fastq_pass) were utilized in subsequent analyses. After concatenation, each
FASTQ file was evaluated to determine read statistics, including the total number of reads, total

base count (the sum of all read lengths), and the average read length (**Supplementary Table 1**).

These metrics were calculated using a custom Bash script that parsed the FASTQ files by

counting sequence lines and summing their lengths to generate the summary statistics.

Taxonomic Profiling

To identify plant host sequences, the read mapping tool Minimap2 (Li, 2018) was used,

choosing as reference genomes the plant genome assembly available in NCBI that was most

closely related to the host of isolation (accession numbers are listed in **Supplementary Table**

1). Minimap2 was also used to identify *Csp* sequences by mapping reads against the two

available *Ct* genomes from cacao: CT2 (accession GCA_009078325.1) and Gudang 4 (accession

GCA_012932095.1). Reads that mapped against both genomes were de-duplicated before

downstream analysis.

The taxonomic profiler Kraken2 (Breitwieser et al., 2018; Wood et al., 2019; Lu et al.,

2023) was employed to assign sequencing reads to bacterial, fungal, and oomycete taxa. Two

databases were used for identification of bacterial and fungal taxa, respectively: (1) the standard

k2_pluspfp_20240605 database, comprising RefSeq archaea, bacteria, viral, plasmid, human,

protozoa, fungi, plants, and UniVec_Core (downloaded June 20, 2024, from

https://benlangmead.github.io/aws-indexes/k2), and (2) a custom fungi-oomycete database

(https://github.com/vinatzer-lab/Csp-custom-sourmash-and-kraken-databases) to address the

limited representation of these taxa in RefSeq. To generate the custom database, 18,644 fungal

and 513 oomycete genome assemblies were downloaded from NCBI on July 24, 2024. These

assemblies were processed using LINflow (Tian et al., 2021) to assign Life Identification

Numbers (LINs). Subsequently, representative genomes were selected at an average nucleotide

identity (ANI) threshold of 99.5%, resulting in a non-redundant, comprehensive Kraken2

database containing 8,797 fungal and 342 oomycete genomes.

Additionally, metagenomic sequence similarity, containment analyses, and taxonomic

profiling were performed using Sourmash v4.5.0 (Brown and Irber, 2016; Irber et al., 2024).

Raw sequencing reads were converted into FracMinHash sketches with a k-mer size of 31 and

scaled parameter of 1000. Bacterial community composition was determined using the sourmash

`gather` command against the GTDB-RS207 reference database. To analyze fungal

communities, the genomes used above for the fungi-oomycete Kraken2 database were used to

build a custom Sourmash database with the same FracMinHash parameters as for bacteria.

Csp **Genome Assembly and Analysis**

Samples containing more than 10,000 reads mapping to the two *Ct* reference genomes

(GCA_009078325.1 and GCA_012932095.1) were selected for genome assembly. Each selected

sample was assembled independently using Flye (Kolmogorov et al., 2019), specifying a

genome size parameter of 34 Mb based on the *Ct* reference genome size. Five rounds of iterative

polishing were conducted to enhance assembly accuracy.

To capture *Csp*-specific genomic regions potentially missed by aligning exclusively to *Ct*

genomes, we constructed a comprehensive assembly by pooling the five metagenomic samples

containing the highest abundance of *Ct*-specific reads. Host-derived sequences were first

removed using Minimap2. The remaining non-host reads were combined and subjected to *de*

novo assembly using Flye (Kolmogorov et al., 2019). Following this assembly, initial contig

filtering was conducted using Kraken2 with both standard and custom fungi-oomycete

databases. Contigs were further analyzed using BLASTn searches (Altschul et al., 1990) against

the NCBI core nucleotide database with relaxed alignment parameters, maximizing detection of

divergent sequences. Additionally, read-depth consistency analyses were performed on each

contig using Samtools (Li et al., 2009), CoverM (Aroney et al., 2025), and mosdepth (Pedersen

et al., 2017). Only contigs exhibiting high sequence similarity to non-*Ceratobasidiaceae* taxa or

inconsistent read-depth patterns were excluded, to minimize the risk of excluding genuine *Csp*

sequences.

Phylogenetic Analysis

For core-genome construction, genome completeness was first assessed using BUSCO v5.2.2

with the Agaricomycetes_odb10 dataset, comprising 2,898 single-copy orthologs (Manni et al.,

2021). Conserved single-copy orthologs present in all genomes used for tree construction were

aligned using MUSCLE (Edgar, 2004), and alignments were trimmed with TrimAl (Capella-

Gutiérrez et al., 2009) to remove poorly aligned regions that could introduce artifacts in

phylogenetic reconstruction. A maximum likelihood tree was then inferred using IQ-TREE

(Nguyen et al., 2015). Branch support was evaluated using ultrafast bootstrap approximations

implemented in IQ-TREE. The final phylogenetic trees were visualized and annotated in R

version 4.2.3 using the ggtree package (Yu et al., 2017).

For the single nucleotide polymorphism (SNP) analysis, reads were aligned to the

reference genome using minimap2 v2.29 (Li, 2018) with the `-ax map-ont` preset, and

alignments were sorted and indexed in a single□step pipeline with samtools (v1.19.1). Variants

were called per isolate using NanoCaller (v3.6.2) in haploid mode (`--mode all, --`

`sequencing ul_ont,--snp_model ONT-HG002, --indel_model ONT-`

`HG002`), producing one VCF containing SNPs and indels per sample. VCFs were

coordinated□sorted, BGZF□compressed and indexed with `bcftools` and `tabix`, then

merged into a single multi□sample VCF (`bcftools merge --force-samples`). The

merged VCF was filtered to retain only high□confidence SNPs (per□site QUAL $\geq$ 20 and

per□sample depth DP $\geq$ 10) using bcftools `view -i 'QUAL>=20 && FMT/DP>=10'`. The

resulting filtered VCF was converted to a multi□FASTA alignment in PHYLIP format using a

custom vcf2phylip script. A maximum□likelihood phylogeny was inferred in IQ-TREE

(v2.2.0.3) (Nguyen et al., 2015) under the `GTR+Γ` model with 1,000 ultrafast bootstrap

replicates. The final phylogenetic tree was visualized and annotated in R version 4.2.3 using the

ggtree package (Yu et al., 2017).

Gene Prediction and Comparative Genomics

Gene prediction was performed using AUGUSTUS v3.5.0 (Stanke et al., 2006) with an *ab initio*

approach trained on the *Rhizoctonia solani AG-1 IA* genome (GenBank: GCF_016906535.1).

The training pipeline began by initializing species-specific configuration files using

`new_species.pl` to establish baseline parameters, followed by automated gene model

training with `autoAugTrain.pl` utilizing multicore optimization. Pre-processing of

annotations retained only mRNA and CDS features to ensure AUGUSTUS training

compatibility, resulting in optimized gene model parameters for accurate prediction in *Csp*

genomes. Following gene prediction, protein-coding sequences were extracted from GFF3

annotations using `gffread` (Pertea et al., 2020) and underwent quality control curation.

OrthoFinder (Emms et al., 2019) performed orthologous clustering using the DIAMOND

(Buchfink et al., 2021) algorithm in a multi-threaded configuration, while custom Python scripts

generated presence/absence matrices to define core and accessory orthogroups with gene IDs

mapped to their respective genomes. Core orthogroup protein sequences were concatenated. The concatenated core protein sequences underwent MAFFT (Kazutaka et al., 2019) alignment using 40 threads for optimal performance, followed by TrimAl processing to remove poorly aligned regions that could interfere with phylogenetic signals. Seqmagick converted the trimmed alignment into PHYLIP format for compatibility with downstream analysis tools. IQ-TREE performed maximum likelihood phylogenetic reconstruction using the VT+F+I+I+R6 substitution model, with branch support assessed through both ultrafast bootstrap and SH-aLRT (1000 replicates each), ultimately providing evolutionary relationship insights among *Csp* isolates.

Functional Genome Annotation

Protein sequences from core and accessory genomes were functionally annotated using InterProScan (Quevillon et al., 2005). Before annotation, sequences were quality-checked to retain only valid amino acid residues. InterProScan analyses utilized multiple protein databases (Pfam, SMART, CDD, etc.) to annotate protein domains, GO terms, and metabolic pathways. InterProScan was executed on high-performance computing resources, allocating 150 GB RAM and 16 CPU cores. Outputs included TSV, XML, and GFF3 formats, providing extensive functional annotations suitable for downstream enrichment and comparative analyses. Quick summary tables were generated to count the frequency of InterPro domains, GO terms, and associated pathways, facilitating functional enrichment analyses to identify potential host-specific adaptations and evolutionary innovations in *Csp* genomes.

Prediction and Characterization of Putative Effector Proteins

Putative effector proteins were identified from the predicted accessory proteome following a multi-step computational pipeline integrating secretion, localization, and functional prediction tools (Logachev et al., 2024). Candidate protein sequences derived from accessory orthogroups were first screened for the presence of N-terminal signal peptides using SignalP v6.0 (Teufel et al., 2022) in eukaryotic mode with fast processing settings. Signal peptide-positive sequences were then analyzed with TMHMM v2.0c (Krogh et al., 2001; Käll et al 2007) in short output mode to identify proteins containing transmembrane helices. Proteins predicted with one or more helices were excluded to only retain putative soluble secreted proteins. Proteins lacking

transmembrane helices were further screened using TargetP v2.0 (Armenteros et al., 2019) in
non-plant eukaryotic mode to identify and exclude proteins predicted to contain mitochondrial
targeting peptides (mTP). The remaining secretome candidates were scanned for endoplasmic
reticulum retention signals (PS00014 motif) using ScanProsite (de Castro et al., 2006). Proteins
carrying this motif were excluded to avoid ER-retained proteins. The refined secretome was
finally evaluated for glycosylphosphatidylinositol (GPI) anchor signals using NetGPI v1.1
(Gíslason et al., 2021). Candidate proteins were split into batches and submitted to the web
server for prediction. Proteins identified as GPI-anchored were excluded from the secretome.
The filtered secretome was submitted to EffectorP v3.0 (Sperschneider & Dodds, 2022) in
fungal mode to predict candidate effector proteins. EffectorP provided classification outputs
distinguishing effector and non-effector proteins, along with localization categories as
apoplastic, cytoplasmic, or dual-localized effectors. EffectorP prediction outputs were parsed to
separate candidate effectors by localization class. Protein sequences corresponding to each
effector class were extracted using seqkit (Shen et al., 2016) and compiled into dedicated
FASTA files.
323
324 **Results**
325

Collection of 106 symptomatic *Csp*-positive samples from 34 plant species in seven U.S.
states
Plant tissue specimens exhibiting VSD symptoms were collected from various woody
ornamental hosts across seven U.S. states. Sampling criteria included the presence of vascular
discoloration, foliar chlorosis progressing to necrosis, and terminal dieback consistent with
previously documented VSD symptoms (**Figure 1**). A total of 106 samples were obtained and
tested positive for *Csp* based on a diagnostic PCR assay (Avin et al., 2025). Virginia contributed
the largest sample cohort (n=64, 58.2%), followed by Tennessee (n=25, 22.7%), North Carolina
(n=7, 6.4%), Indiana (n=4, 3.6%), Texas (n=3, 2.7%), Maryland (n=2, 1.8%), and Missouri
(n=1, 0.9%) (**Figure 2A**). Within Virginia, specimens were collected from 14 counties with the
greatest concentration in the central Piedmont region (Orange, Louisa, Fauquier, Nelson,
Loudoun and Albemarle) (**Figure 2B**), which contains the highest density of commercial woody
ornamental nurseries in the state.

Figure 1. Representative vascular streak dieback (VSD) symptoms observed in *Csp*-positive

woody ornamental hosts: (A) *Acer rubrum* 'October Glory'; (B) *Cercis canadensis* 'Flame

Thrower'; (C) *Cornus kousa* 'Greensleeves'. Symptoms include vascular discoloration, foliar

chlorosis progressing to necrosis, and terminal dieback.

The samples originated from 34 woody ornamental host species belonging to 19 different

plant families (**Supplementary Table 1**), with eastern redbud (*Cercis canadensis*, n=28),

Freeman maple (*Acer × freemanii*, n=18), and flowering dogwood (*Cornus florida*, n=16)

comprising the majority of specimens. Additional hosts included, but were not limited to,

Japanese maple (*Acer palmatum*), southern wax myrtle (*Morella cerifera*), Chinese dogwood

(*Cornus kousa*), crepe myrtle (*Lagerstroemia indica*), and various oak species (*Quercus* spp.).

Figure 2. Geographic distribution of *Csp* sample collection sites across the United States. **A** map of the U.S. with the states from which samples were obtained highlighted in purple. The numbers indicate the number of samples collected from each state. **B** A map of Virginia showing counties and number of samples that were collected.

Deep metagenomic sequencing provides insight into microbial communities associated with VSD

Sequencing was performed on DNA either extracted from vascular tissue (61 samples) or from mycelium cultured directly from vascular tissue (45 samples). Sequencing yielded a cumulative total of 188,856,320 reads across all samples. The sequencing depth per sample was highly variable, ranging from 14,364 to 7,555,390 reads, with an average of 1,760,656 reads (see detailed sequencing results for each sample in **Supplementary Table 1**). The mean read length for all sequencing runs was 2,940 base pairs (bp) with the longest obtained read being 396,120 bp long, resulting in a combined total of approximately 893 gigabases (Gb) of sequence data. The high sequencing depth in many samples promised to enable comprehensive profiling of microbial communities and robust detection of *Csp* even in samples with low *Csp* abundance.

Figure 3. Boxplots of log$_{10}$-transformed normalized read counts (reads per million) for plant, fungal, Oomycota, and bacterial taxa, comparing Mycelium-derived (left) and plant-derived (right) samples.

Not surprisingly, plant reads represented the majority of reads in sequenced plant samples (67.8% on average with a range from 50.01-99.01%), but even in some mycelium samples (averaging 31.7% with a range of 0.3-88.2%), possibly due to inclusion of some plant material during sample preparation (**Figure 3**). As expected, fungal and bacterial read counts were higher in mycelium compared to plant samples (approximately 10-fold) because of the lower average number of plant reads. However, in both sample types, bacterial read and fungal read numbers were similar to each other. Reads assigned to oomycetes were overall much lower, but relatively higher in plant samples compared to mycelium samples.

Previous investigations of VSD in woody ornamentals in the U.S. primarily focused on the fungal pathogen *Csp* (Liyanapathiranage et al., 2024). Here, metagenomic sequencing allowed us to investigate the entire microbial communities associated with VSD and determine if any other microbe consistently co-occurs with *Csp*. Fungal taxonomic profiling using the Kraken2 tool (Breitwieser et al., 2018; Wood et al., 2019; Lu et al., 2023) with a custom fungal and oomycete database revealed distinct fungal community patterns between mycelium and plant samples. In mycelium samples, *Csp* was the dominant taxon, ranking first in 23 of the 30 samples with relative abundance from 19.6% to 91.9%, exceeding half of all non-host reads in 20 samples. The number of *Csp* reads was independently confirmed by mapping reads against *Ct* genomes, which largely aligned with the taxonomic profiling results (**Figure 4** and **Supplementary Table 1**). Based on Kraken2, *Fusarium falciforme*, a pathogen that causes wilt disease and root rots in many herbaceous and some wooden species (Thao et al., 2021; Siddig et al., 2023), was the dominant taxon in the remaining mycelium samples. Plant samples exhibited greater variability regarding the most highly abundant fungal species than mycelium samples. Only in 16 of 76 plant samples was *Csp* the most abundant fungus with relative abundances ranging from 6.33% to 82.46% out of all non-host reads. As in mycelium samples, when *Csp* was not dominant, *F. falciforme* was often the most dominant species based on Kraken2. Other fungal species detected included the endophyte *Tetracladium marchalianum*, the broad host-range pathogen *Fusarium oxysporum,* the environmental fungus *Penicillium bialowiezense,* the powdery mildew pathogen *Erysiphe pulchra,* the mushroom *Psilocybe semilanceata*, and the endophyte (and occasionally pathogen) *Trichoderma* spp.

To validate the fungal species identified by Kraken2, we performed additional analyses on the samples for which Kraken2 had found them to be most abundant (**Supplementary Table**

2). As pointed out above, while read mapping had largely confirmed Kraken2 results for *Csp*,

this was not the case for the other fungal species. For example, on average, only 0.3% of reads

identified by Kraken2 as *F. falciforme* mapped to *F. falciforme* reference genomes. When these

reads were assembled into contigs, the contigs could not be confirmed by BLAST searches

against a custom *Fusarium* database. Further validation using the FracMinHash-based sourmash

tool with a custom fungi and oomycete database containing the same genomes used for the

Kraken2 database also failed to detect *F. falciforme* and several other fungal species while only

Fusarium oxysporum was confirmed but was detected at much lower abundance. This strongly

suggests that Kraken2 likely misclassified and overestimated the abundance of other fungal

species than *Csp*.

Figure 4. Boxplots of log□□-transformed normalized read counts comparing Kraken2 (red) and

Minimap2 (blue) across mycelium-derived (left) and plant-derived (right) samples. In both

sample types, Minimap2 shows a slightly higher median and tighter distribution of read counts

than Kraken2.

Oomycetes represented a minor component of the non-host reads in both sample types

based on Kraken2 using our custom fungi-oomycetes database accounting on average for only

1.08% of all non-host reads in plant samples and for only 0.36% in mycelium samples.
Plasmopara halstedii and *Plasmopara obducens* were the most frequently detected species
across both sample types (**Supplementary Table 1**). Other regularly detected oomycetes
included *Pseudoperonospora cubensis* and various *Phytophthora* species. We thus conclude that
oomycetes are unlikely to play a role in VSD.

Metagenomic analysis of bacteria revealed thirty-seven genera among the top three most
abundant bacterial taxa across samples (**Supplementary Table 1**). Relative abundances varied
between plant samples and mycelium samples. In plant samples, members of the
Enterobacteriaceae including *Enterobacter hormaechei*, *Escherichia coli*, and *Klebsiella*
quasipneumoniae were among the most prevalent, while mycelium-derived samples were
characterized by higher relative abundances of *Stenotrophomonas maltophilia, Clostridium*
perfringens, and *Achromobacter xylosoxidans* (**Figure 5**, **Supplementary Table 1**). To validate
these taxonomic assignments, we performed read mapping against reference genomes for the
five most abundant bacterial species. Of these, only assignments to *Achromobacter xylosoxidans*
and *Stenotrophomonas maltophilia* were confirmed and enough reads were found to assemble
the respective genomes, whose identities were confirmed using genomeRxiv (Pritchard et al.,
2022). For *Escherichia coli*, results varied significantly across samples, prompting additional
verification using the FracMinHash-based Sourmash tool, which indicated that most reads
assigned to *E. coli* by Kraken2 more closely matched genomes of various environmental and
commensal taxa, such as *Methylobacterium* sp. *Bradyrhizobium sp.*, and *Curtobacterium* sp.
suggesting potential misclassification by Kraken2. Likewise, reads assigned by Kraken2 to
Clostridium perfringens and *Mycobacterium branderi* aligned to other taxa in the Sourmash
analysis (**Supplementary Table 3**).

In summary, besides *Csp*, no other pathogenic bacterial, fungal or oomycete species was
consistently identified across samples making it unlikely that other pathogens than *Csp*
contribute to VSD on woody ornamentals in the U.S.

Figure 5. Two-panel boxplot of log₁₀-transformed normalized reads for the top 10 bacterial species identified by Kraken2, comparing mycelium-derived (left) and plant-derived (right) samples. Species marked with an asterisk (*) in the legend were not confirmed by read mapping or by analysis with Sourmash, indicating likely false-positive assignments by Kraken2.

Genome Assemblies of *Ceratobasidium species* (*Csp*) isolates

Csp reads identified by mapping against the two *Ct* reference genomes were used to construct metagenome-assembled genomes (MAGs). Mapped reads were extracted, deduplicated, and assembled. High-quality draft genome assemblies were obtained from 17 samples (4 from plant samples and 13 from mycelium samples), comparable in size and completeness to the *Ct* reference genomes based on BUSCO completeness scores (**Table 1**). Comprehensive statistics including read counts, genome completeness, genome sizes, and BUSCO scores are detailed in **Supplementary Table 4**. Twenty-five other samples yielded partial genome assemblies due to lower read counts (fewer than 59,918 reads) with genomes sizes ranging from 2.3 to 30.2, while 46 samples had such a small number of *Csp* reads (fewer than 14,177) that no meaningful assemblies were obtained. Detailed assembly statistics are reported in **Supplementary Table 4**.

Since construction of the *Csp* MAGs relied exclusively on reads mapped to the *Ct*
reference genomes, we were concerned that we may have failed to capture genomic regions
potentially present in *Csp* but absent from *Ct*. Therefore, we conducted a second, independent
assembly approach to include these potential regions unique to *Csp*. To do this, we selected the
five samples with the highest *Ct*-aligned read counts. Instead of assembling mapped reads, we
removed host sequences and then assembled all other reads (2,520,850 reads with a total length
of 12.1Gb). The assembly resulted in 522 contigs. We then taxonomically classified these
contigs using BLASTn and Kraken2 and removed 4 contigs that had best hits to non-fungal
organisms (**Supplementary Table 5**). While detailed assembly results are shown in
Supplementary Figure 1, **Table 1** shows that the comprehensive assembly is very similar in
quality to the best individual assemblies. Importantly, the number of annotated genes in the
comprehensive assembly (10,962) fell within the range of annotated genes in the individual
assemblies (10,442 to 11,141) and, based on our pangenomes analysis (see more details below),
only five singletons were found in the comprehensive assembly suggesting that the best
individual *Csp* assemblies were not missing a significant number of genes.
482
483   **Table 1:** Summary statistics of *Csp* genome assemblies and *Ct* reference genomes.

Sample ID	Assembly Length (bp)	N50 (bp)	No. of Contigs	Annotated Genes	No. of Singletons	BUSCO (Agaricomycetes _odb10)	BUSCO (Fungi_odb 10)
23_02068_EasternRedbud_IN	32.97	21,383	2,325	10,442	356	72.7	83.3
FBG5639_EasternRedbud_TN	36.94	53,266	2,040	11,655	567	82.0	91.9
RT23_468_FloweringDogwood_VA	38.11	28,656	2,614	10,971	320	78.1	89.3
RT23_372_FloweringDogwood_VA	36.84	161,947	786	10,990	35	83.6	94.2
RT23_384_FloweringDogwood_VA	36.78	460,389	562	10,936	8	84.0	94.6
RT23_392_SouthernCatalpa_VA	36.86	549,311	483	10,779	2	83.7	94.5
RT23_402_RedOsierDogwood_VA	36.01	643,614	475	10,887	0	83.9	94.6
RT23_423_ChineseDogwood_VA	36.65	390,318	543	10,678	9	83.9	94.6
RT23_424_ChineseDogwood_VA	35.37	139,030	915	10,919	38	83.6	94.2
RT23_428_ChineseDogwood_VA	37.00	616,318	539	10,939	1	83.8	94.5
RT23_429_ChineseDogwood_VA	36.86	700,228	485	10,880	3	83.9	94.5
RT23_433_FloweringDogwood_VA	36.53	816,925	401	11,141	5	83.9	94.6
VSD24_268_Sourwood_VA	32.89	37,465	1,409	10,217	311	72.2	82.7
VSD24_269_RedOsierDogwood_VA	36.25	266,961	559	10,863	24	83.6	94.6
VSD24_270_FloweringDogwood_VA	36.11	97,071	669	10,913	177	80.4	91.7

VSD24_DSB_110M_SouthernWaxMyrtle _VA	36.82	379,450	293	10,779	28	83.6	94.6
VSVD24_368_EasternRedbud_VA	36.18	320,509	504	10,792	13	83.9	94.6
Comprehensive_assembly	37.75	824,000	518	10,962	5	83.7	94.3
Ceratobasidium theobromae CT2	31.20	86,100	1,240	9,763	104	83.6	94.5
Ceratobasidium theobromae Gudang 4	38.60	39,200	4,661	13,311	1,201	82.0	92.2

Phylogeny to resolve evolutionary relationships between *Csp* samples from woody ornamentals in the U.S. and related species and genera

To resolve the evolutionary relationships between *Csp* genomes from the U.S., other publicly available *Csp* and *Ct* genomes, and related *R. solani* genomes, a core phylogenetic tree was constructed based on 76 conserved single-copy orthologs. Only the 13 U.S. *Csp* genome assemblies with the highest BUSCO scores were used. The resulting maximum likelihood tree (**Figure 6A**) revealed a well-supported monophyletic clade composed exclusively of U.S. *Csp* isolates. The relationships among most of these isolates remains unresolved with bootstrap support below 50, leading to a striking polytomy. The Southeast Asian *Ct* genomes (CT2 and Gudang4 from cacao) formed a separate, well-supported clade distinct from the U.S. *Csp* lineage. The distinct branch of the single LAO1 genome from cassava suggests that it may be the only representative of a yet undersampled additional *Ceratobasidium* clade. These genomes all shared a most recent common ancestor (MRCA) distinct from a sister clade of all other *Ceratobasidium* species. Three *R. solani* genomes, chosen to root the tree, formed yet another, well-separated clade. In summary, the core genome tree confirms that the *Csp* isolates from woody species in the U.S. only recently diverged from a common ancestor and that they are closely related to, but distinct from, the *Ct* isolates from cacao and cassava in Southeast Asia.

Figure 6A. Maximum likelihood phylogenetic tree constructed using 76 single-copy orthologous genes from assembled U.S. *Csp* genomes, publicly available *Ceratobasidium* genomes, and *R. solani* isolates used as an outgroup to root the tree (best model: VT+F+I+I+R6). Branch tips are labeled with the sample IDs. Bootstrap support values are shown at each node.

Phylogeny to resolve evolutionary relationships among *Csp* samples from woody ornamentals in the U.S.

A finer-scale analysis focusing on the 17 U.S. *Csp* assemblies of the highest quality and the three publicly available *Ct* genomes based on 1,473 core genes confirmed the genetic coherence of the U.S. *Csp* population (**Figure 6B**). Bootstrap support below 50 at the base of the U.S. *Csp* clade still leads to multifurcation confirming a recent radiation event from a common ancestor. However, compared to the 76-gene tree in **Figure 6A**, the U.S. *Csp* clade in the 1,473-gene tree has more resolution and a few subclades with high bootstrap support become evident. In most cases, these clades include isolates from different host species such as redbud, dogwood, and wax myrtle, indicating the absence of host adaptation. Since only one isolate from Indiana and

519   one from Tennessee provided enough reads for high quality assemblies and all other isolates

520   came from Virginia, nothing can be concluded about regional differentiation.

Figure 6B: Core genome phylogenetic tree of *Csp* genomes recovered from woody ornamental

hosts in the United States (best model: VT+F+I+I+R6). A total of 1,473 core genes were used

to construct the tree, which was rooted on genomes of *Ct* isolates from cacao and cassava in

Southeast Asia. Branch tips are labeled with the sample IDs.

527   Since we only obtained high quality genome assemblies from 17 out of the 106 total

samples, our ability to infer evolutionary relationships of U.S. *Csp* isolates using a core genome

approach was limited. Thus, we also tried to use unassembled *Ceratobasidium* reads of a larger

number of samples: the 17 samples that had provided high quality draft assemblies and 14

samples that had provided partial genome assemblies, which included two *Ct* samples from

cacao from Indonesia. These reads were aligned against the *Ct* LOA1 genome, the genome of

533   the closest available relative of the U.S. *Csp* lineage based on the above core genome analyses.

534   1,317,177 SNPs were identified and used for phylogenetic reconstruction (**Figure 7)**.

Figure 7. Maximum☐likelihood phylogeny constructed under a `GTR+Γ` model from 32 *Ceratobasidium* isolates using 1.32 million SNPs identified by the NanoCaller tool by aligning reads against the *Ct* LOA1 genome, which was chosen as root. Ultrafast bootstrap values (≥ 50%) are shown at nodes.

The two Indonesian cacao isolates formed a separate clade with 100% bootstrap support,

confirming the deep evolutionary split between U.S. *Csp* isolates and Southeast Asian *Ct*

isolates seen in the two core genome trees (**Figure 6**). All U.S. *Csp* isolates form a single clade,

also with 100% bootstrap support and thus also in agreement with the core genome trees. The

only North Carolina isolate branched off from the most basal node of this clade, which is in line

with the epidemiological finding that trees with VSD symptoms were first observed in North

Carolina in 2018, and *Csp* may have spread from there to the other U.S. states. Intriguingly, all

but two Virginia isolates form a single well supported clade and the majority of Tennessee

isolates form a second well supported clade suggesting that *Csp* populations in these two states

may have started to genetically separate from each other. As already seen in the core genome

tree, there are no host-specific lineages suggesting no adaptation to different host species.

Exploring gene content differences between genomes of U.S. *Csp* isolates and other

***Ceratobasidium* and *Rhizoctonia* isolates and among U.S. *Csp* isolates**

All phylogenetic trees showed that the genomes of the U.S. *Csp* isolates form a single clade that

is clearly distinct from the genomes of the Southeast Asian *Ct* isolates from cacao and cassava

and all public genomes of other *Ceratobasidium* and *Rhizoctonia* isolates. However, just how

different is this U.S. clade compared to all other isolates? To help answer this question, we

performed a pangenome analysis. We found a total of 176,561 proteins in 12 *Ceratobasidium*

and 3 *R. solani* genomes. This dataset incorporated the five highest quality individual genome

assemblies of our U.S. samples, our comprehensive assembly, the genomes of the three *Ct*

isolates from cacao and cassava, three public genomes of *Csp* isolates with the highest BUSCO

scores, and the genomes of the three *R. solani* genomes used as outgroup in our core genome

phylogeny (**Figure 6A**).

The pangenome analysis identified 10,913 non-redundant orthologous protein groups

(orthogroups). Of these, 5,211 orthogroups (47.8%) were classified as core, being shared across

every single analyzed genome. Additionally, 4,278 orthogroups (39.2%) were categorized as

accessory, present in some but not all genomes, reflecting genomic variability within the

species. Singletons, representing 13.1% of the orthogroups (1,424), consisted of genes unique to

individual genomes, either due to differences in genome quality or representing true differences

between individual isolates (**Figure 8A**). The gene-accumulation curve (**Figure 8B**) plots

orthogroup counts against the number of genomes considered. Counts generally decline as *N*

increases, with a pronounced spike in the rightmost bin (15 genomes) that reflects the core

orthogroups shared by nearly all genomes, while the spread across intermediate bins indicates an

open pangenome with substantial accessory variation among isolates.

The UpSet plot (**Figure 8C**) shows how the accessory orthogroups are distributed.
Importantly, 269 orthogroups were shared among the five genomes derived from US samples
and the comprehensive assembly but absent from all other analyzed genomes, while 161
orthogroups were shared by the two *Ct* genomes from cacao but absent from all other analyzed
genomes. This difference in gene content underscores the evolutionary separation between these
two clades. An additional 86 genes were shared among the U.S. *Csp* genomes and the *Ct* LAO1
genome from cassava but absent from all other genomes while fewer than 50 genes were shared
between the *Ct* LAO1 genome and the two *Ct* genomes from cacao or between the U.S. *Csp*
genomes and the two *Ct* genomes from cacao. This further confirms the closer relationship
between the U.S. *Csp* isolates with the *Ct* lineage from cassava than with the *Ct* genomes from
cacao, which was also evident in the core genome tree in **Figure 6A**. Per-genome composition
plots (**Figure 8D**) further indicate broadly similar proportions of core, accessory, and singleton
content across isolates, with modest variation in singleton burden that likely reflects both
assembly completeness and genuine lineage-specific gene gain/loss.

To hone in on the genomic diversity among the U.S. *Csp* genomes and their comparison
with the *Ct* genomes, we conducted a second pangenome analysis of all 17 *Csp* genomes and 3
Ct genomes from the core-genome phylogeny (**Figure 6B**). We also included the comprehensive
assembly because of the confidence we have in its completeness. Clustering 227,262 proteins
from these 20 genomes resulted in 9,902 orthologous groups: 65.1% core, 31.7% accessory, and
3.2% singletons. The UpSet plot (**Supplementary Figure 2**) reveals 6,446 core orthogroups
found in all genomes. One can notice that each of the lowest quality genomes, such as *Csp*
VSD24-268_Sourwood_VA or 23_02068_EasternRedbud_IN, are missing between 200 and 400
genes, probably because of their low genome completeness. Therefore, the true core genome of
the analyzed genomes probably exceeds 7,000 orthogroups. Since we had observed two well-
supported clades in **Figure 6B**, we were particularly curious to see if these two clades could be
distinguished based on orthologue presence or absence, but no such trend is visible suggesting
that the two clades have only diverged minimally so far.

Figure 8. Pangenome analysis. **A.** Composition of core and accessory orthogroups and singleton genes in the *Ceratobasidium*/*Rhizoctonia* pan-genome. **B.** Gene accumulation curve (orthogroup count vs. number of genomes): each bar represents how many gene families are present when considering a specific number of genomes. The numbers decrease as more genomes are included, with a sharp increase near the rightmost bars (particularly at 15 genomes), reflecting the core gene families shared by nearly or all genomes. This curve illustrates how the pangenome grows and highlights both unique and shared gene content across genomes. **C.** UpSet plot showing intersections of orthogroups across 15 *Ceratobasidium* and *R. solani* genomes. Bar heights indicate the number of orthogroups for each presence/absence pattern (only intersections with ≥50 OGs are shown), with the connected dots below identifying the genomes included in each intersection. **D.** Composition of core, accessory and singleton genes per genome.

As for the differences between the U.S. *Csp* genomes and the *Ct* genomes from Southeast Asia, this focused *Csp*/*Ct* pangenome analysis confirmed the substantial differences in orthogroup content found between the two groups in the wider pangenome analysis from **Figure**

8, only that the U.S. *Csp* genomes now shared slightly fewer orthogroups, 214 versus 269,

probably because of the inclusion of the lower quality genomes with lower completeness.

Prediction of effector genes in accessory genomes

For effector prediction, we targeted the accessory orthogroups identified in the 15 genomes used

in the phylogenetic analysis (**Figure 6A**) and the pangenome analysis (**Figure 8**). Accessory

proteins per genome ranged from 2,184 in *Ct* LAO1 to 8,485 in *Csp* AG_Ba_Jn (**Table 3**). Out

of the 16,294 accessory proteins, SignalP 6.0 predicted 1,431 to have N-terminal signal peptides.

To exclude membrane-bound proteins from this group, we applied TMHMM v2.0c to remove

sequences with at least one predicted transmembrane (TM) helix. After this filtering step, 1,231

proteins were retained. One of these proteins was removed because it contained a mitochondrial

transit peptide identified by TargetP 2.0. None were found to have ER-retention motifs by

ScanProsite (PS00014) or GPI-anchor signals by NetGPI 1.1. Therefore, 1230 secretome

candidates were finally screened with EffectorP 3.0 in fungal mode, which partitioned them into

effectors and non☐effectors and further sub☐classified effector candidates as apoplastic,

cytoplasmic, or dual☐localized (**Table 3**).

Table 3. Summary of accessory gene content and effector predictions for *Ceratobasidium* and *R.*

solani genomes.

Sample	Accessory Genes	Secreted	Effectors	Apoplastic	Cytoplasmic	Dual	Non☐effectors
Ceratobasidium_cereale_R0301.hap1	3,067	191	88	50	24	14	103
Ceratobasidium_cereale_R0301.hap2	3,078	195	87	50	25	12	108
Ceratobasidium_sp.AG_Ba_JN	8,485	514	207	105	57	45	307
Ceratobasidium_theobromae_CT2	2,241	171	79	32	20	27	92
Ceratobasidium_theobromae_Gudang4	4,073	322	149	64	48	37	173
Ceratobasidium_theobromae_LAO1	2,184	178	82	39	26	17	96
csp_comp_assembly	3,281	691	312	144	83	85	379
Rhizoctonia_solani_AG_1_IA	2,211	161	73	28	23	22	88
Rhizoctonia_solani_AG1_IA_HG81	2,355	170	74	33	23	18	96
Rhizoctonia_solani_YS_517	2,607	191	93	45	25	23	98
RT23_384_FloweringDogwood_VA	2,787	200	94	45	26	23	106
RT23_392_SouthernCatalpa_VA	2,723	203	94	46	28	20	109
RT23_423_ChineseDogwood_VA	2,712	198	95	40	31	24	103

VSD24_269_RedOsierDogwood_VA	2,690	203	90	41	21	28	113
VSD24_DSB_110M_SouthernWaxMyrtle_ VA	2,691	219	98	53	21	24	121

To identify clade-specific effectors, we analyzed the predicted effector repertoires across
the major clades identified in our core-genome phylogeny (**Figure 6A**). The non-U.S. *Csp* clade
contained the highest number of unique predicted effectors, with 150 effectors specific to this
group. The U.S. *Csp* clade harbored a distinct set of 34 unique effectors, while the *Ct*
(cacao/cassava) clade had 52 unique effectors. The *R. solani* group displayed the lowest effector
count, with only 10 identified effectors. These results reveal substantial differences in effector
content between the US *Csp* clade and its sister clades, yielding a distinct set of effector
candidates that can be further investigated for their potential roles in host specificity and
virulence.

Discussion
Since the first detection of *Csp* on an eastern redbud with VSD symptoms in Tennessee in 2019,
VSD has quickly spread across 13 U.S. states and *Csp* has been detected molecularly on 50
genera of woody ornamentals (Liyanapathiranage et al., 2025; McClellan 2023). However, the
fastidious nature of *Csp* has prevented Koch's postulates to be completed and to establish *Csp* as
the sole causative agent of VSD (Ali et al., 2019; Fredricks & Relman., 1996, Liyanapathiranage
et al. 2024). Therefore, the first objective of this study was to determine if any other bacterial,
fungal, or oomycete pathogen besides *Csp* is consistently associated with VSD and may
contribute to the disease.

Metagenomic sequencing is a promising approach to answer this question since it does
not require culturing and can potentially identify all microbes in a sample to the species rank
and, if enough sequencing data are obtained, MAGs can be obtained and used in phylogenetic
reconstruction (Asnicar et al., 2020; Czech et al., 2022). However, while metagenomics has
become relatively established for the identification of bacterial pathogens, including plant
pathogens (Duan et al., 2009; Piombo et al., 2021; Navgire et al., 2022), identification of fungi from
metagenomic data is challenging and is still in its infancy (Donovan et al., 2018; Gökdemir et al.,
2022; Avershina et al., 2025). One main issue is that taxonomic profiling of metagenomic
sequences heavily relies on the availability of correctly classified reference genomes and only

659 fungal genomes are currently (July 2025) included in NCBI's manually curated RefSeq
database compared to 435,623 prokaryotic genomes (Goldfarb et al., 2025). Therefore, instead of
using the standard RefSeq-based reference database provided by the developers of the
taxonomic profiler Kraken2, we built a custom fungal and oomycete Kraken2 database using all
genomes from NCBI's Assembly database. However, since curation by NCBI of the Assembly
database is limited, genomes may be misclassified, leading to errors in the Kraken2 database
constructed from these genomes, and, ultimately, giving false negative and positive results. This
is probably the reason that several fungal species identified in our metagenomes by Kraken2
were not confirmed when using other tools (sourmash and minimap2). The same problem of
false positives even happened when using Kraken2 with a standard RefSeq-based database for
identification of bacterial species, possibly a result of misclassified genomes even presents in
NCBI's RefSeq database (Bagheri et al., 2020; Chorlton SD., 2024; et al., 2024) and/or limitations
inherent to the Kraken2 algorithm (Breitwieser et al., 2018; Wood et al., 2019; Lu et al., 2023).
Therefore, improved Kraken2 reference databases are urgently needed, and it is key not to rely
on a single taxonomic profiler when analyzing metagenomic data.

Importantly though, we were able to identify sequencing reads as *Csp* in all samples
using both Kraken2 in combination with our custom fungal database (which included *Ct*
genomes while the standard RefSeq-based Kraken2 database did not) and mapping to two *Ct*
reference genomes using minimap2. We obtained a similar number of reads with both tools
giving us confidence that the number of false positive and negative reads assigned to *Csp* by
these tools were limited. However, independently of the bioinformatics tools, we cannot exclude
that a small number of *Csp* sequences were the results of barcode hopping, whereby some DNA
molecules from one sample get erroneously labeled with the barcode from another sample, as
has been reported to occur at low frequency with nanopore sequencing (Ezpeleta et al., 2022).

Since no bacterial or fungal plant pathogen besides *Csp* was consistently detected in the
plant samples, our results confirmed that no other pathogen beyond *Csp* plays a significant role
in VSD. Even *Stenotrophomonas maltophilia*, sometimes reported as plant pathogen (Hu et al.,
2021), was mainly identified in the mycelium samples but not in the plant samples. However,
we were surprised by the low number of *Csp* reads in many of the plant samples. We speculate
that in some cases samples may have been taken from tissue with low inoculum or late during
infection (with *Csp* DNA already degraded by the time of sampling) or that the employed DNA

extraction protocols were relatively inefficient in releasing DNA from *Csp* cells compared to

releasing DNA from other fungal or bacterial cells. Moreover, metabolites present in woody

tissue are known to interfere with DNA extraction (Rizzo et al., 2022). Thus, for sensitive

detection of *Csp*, it is important to still improve DNA extraction protocols.

After we had confirmed *Csp* as the VSD pathogen with the metagenomic results, we

attempted assembly of *Csp* genomes from as many samples as possible for downstream core

genome phylogenetic and pangenome analyses. For including genomes in downstream analysis,

we used a minimum completeness score threshold of 70%, based on the Agaricomycetes

BUSCO dataset (Manni et al., 2021). This may seem a relatively low completeness score but even

our comprehensive assembly with a genome coverage of 250X (*i.e.*, every region of the

assembly was covered on average by 250 reads making it very unlikely that some genomic

regions were not covered at all) only reached a completeness score of 83.7. This suggests that

the *Ceratobasidium* genus may be missing some of the genes present in most of the other

sequenced Agaricomycetes genomes or that some genomic regions either escaped nanopore

sequencing or could not be assembled. However, since our average read length was just shy of

3,000bp and nanopore sequencing is particularly well suited for assembling structurally complex

regions and repeat regions (Zheng et al., 2020; Wang et al., 2021; Kinkar et al., 2021), this seems

unlikely. Also, since the pangenome analysis only identified 1 singleton gene in the

comprehensive assembly and only between 0 and 38 singletons in the 12 best individual

assemblies (which had completeness scores of 83-84% based on BUSCO

agaricomycetes_odb10), we conclude that our best individual assemblies have higher

completeness than what the BUSCO scores suggest. Moreover, the pangenome results suggest

that the genome assemblies present very low levels of contamination with other microbial

sequences or plant sequences, and that sequencing errors did not significantly affect gene

calling. Finally, the highly similar gene content between the best individual *Csp* assemblies

(assembled from reads that mapped to genomes of the related, yet distinct, Southeast Asian *Ct*

lineage) and the comprehensive assembly (which was based on all non-plant reads from five

samples) shows that when using long reads, assembling genomes from reads mapped to a related

genome has a low risk of accidentally excluding genomic regions that are possibly absent from

the related genome used for mapping.

Once the genome assemblies were obtained, we built one 76-gene core genome tree to
investigate the evolutionary relationships between the *Csp* genomes from the U.S. and related
Ceratobasidium and *Rhizoctonia* species from other world regions, and one 1,473-gene core
genome tree to unravel the evolutionary relationships among the *Csp* genomes from the U.S.
The results from the first tree mainly confirmed a previously published tree based on the ITS,
LSU, ATP6, TEF1, and RPB2 genes (Liyanapathiranage et al., 2024). However, probably
because of the higher number of loci used (76 compared to 5), a single node at the base of all
Ceratobasidium genomes had high bootstrap support confirming that the *Ceratobasidium* genus
is monophyletic while the 5-gene tree could not separate the *Ceratobasidium* genomes from the
Rhizoctonia genomes. Moreover, the 76-gene tree subdivided the *Ceratobasidium* clade into two
strongly supported subclades: one consisting of the Southeast Asian *Ct* genomes and the U.S.
Csp genomes and the other consisting of all other *Ceratobasidium* species. Interestingly, both
the 76-gene tree and the 1,473-gene tree provide strong support for the *Csp* genomes from the
U.S. sharing a more recent common ancestor with the *Ct* LAO1 genome from cassava in Laos
than with the *Ct* genomes from cacao in Indonesia. Because of the small number of genomes
that were available, this may just be a coincidence. However, the result aligns with the earlier
finding that the ITS regions of *Csp* isolates in the U.S. are identical to ITS sequences of
Ceratobasidium isolates from Japanese honeysuckle in China (Liyanapathiranage et al., 2024),
of which Laos is a direct neighbor, further suggesting that the U.S. *Csp* lineage may have its
origin in that geographic region.

Regarding the evolutionary relationships among U.S. isolates, the 76-gene tree did not
provide support for any subclades. This is not surprising since their most recent common
ancestor (MRCA) may have existed not much longer than 7 years ago when the first trees with
VSD symptoms were observed. Therefore, few mutations may have accumulated in the 76 core
genes since then. The 1,473-gene tree did reveal two well-supported individual clades, both
including samples from several different plant genera suggesting no adaptation to different
hosts. Because only two clades had strong bootstrap support and since most of our best
assemblies were of samples collected in one geographic region of Virginia, we could not make
any conclusions about possible dissemination pathways. Therefore, we attempted construction
of a SNP-based phylogeny based on alignment of unassembled individual reads against the *Ct*
LAO1 genome. Since assembly was not required, this allowed us to include many more samples.

While ONT significantly reduced their sequencing error rate over the years, it is still around 1%

(Delahaye et al., 2020; Sahlin et al., 2021). Probably because of this relatively high error rate,

even when using the most stringent SNP-calling parameters, an excessively high number of 1.32

million SNPs were called. Random sequencing errors are expected to occur independently in

each genome and are not expected to falsely group genomes together into clades. However, this

high number of SNPs still reduced our confidence in the obtained SNP tree (even though

bootstrap support for most clades was very high). Therefore, although the location of the only

North Carolina sample at the very base of the U.S. clade (suggesting that it is most similar to the

MRCA of all U.S. isolates) agrees with the finding that VSD was first observed in North

Carolina and *Csp* may have spread to the rest of the U.S. from there, this result will need to be

followed up on. Additional sampling and sequencing, ideally using technologies with lower

error rates, such as Illumina or PacBio HiFi, should be performed. Similarly, the possible

separation of *Csp* into a Virginia and a Tennessee sub-population also still needs to be

confirmed.

The final question we wanted to answer was: are there any signs in the genomes of the

U.S. *Csp* isolates that suggest that the *Csp* lineage adapted to woody ornamentals and North

American climate and, possibly, represents a new *Ceratobasidium* species distinct from *Ct*? Our

pangenome and effector gene prediction analyses found substantial differences in gene content

between the genomes in the U.S. *Csp* clade and genomes in its sister clades. The U.S. *Csp* clade

contained 214 to 269 unique genes, depending on which genomes were compared. Of these, up

to 34 genes were predicted to encode secreted *Csp*-specific effectors, which possibly contribute

to virulence by suppressing plant immunity or degrading plant cell walls, typical mechanisms

used by fungal plant pathogens (de Sain and Rep, 2015; Jashni et al., 2015). Unfortunately,

following up on these findings with functional studies of pathogenicity and host specificity (Gil-

Ordóñez et al., 2024; De La Fuente et al., 2022) will be complicated because of the fastidious

nature of *Csp*, making in unlikely to unravel the basis of the exceptional wide host range of the

U.S. *Csp* lineage any time soon.

Interestingly, the *Ct* lineage presented by the only available LAO1 genome from cassava

in Laos is not only more closely related to the U.S. *Csp* lineage based on phylogeny but also

shares more genes with the U.S. *Csp* lineage than the cacao lineage, further suggesting that the

U.S. *Csp* lineage may have originated from a pathogen population located in geographic

proximity to Laos. Therefore, the question about speciation is more complex than we
anticipated. We may either have just one *Ct* species including *Ct* from cacao and from cassava
and *Csp* from ornamentals in the U.S., two species (either grouping *Ct* from cassava in Laos
with the U.S. *Csp* lineage or with the *Ct* lineage from cacao), or three separate species. Only a
more in-depth taxonomic analysis including additional genome-sequenced isolates, and possibly
phenotypic analyses, will be able to answer this question.

In conclusion, this study revealed how metagenomics using long read sequencing can
overcome the challenges involved with the absence of pure cultures when dealing with
fastidious fungal pathogens. While we encountered many issues with false positives during
taxonomic profiling, the use of multiple, parallel approaches and a custom database still allowed
us to confidently conclude that *Csp* is the only causative agent of VSD in U.S. ornamentals.
Also, the long reads allowed assembly of several high-quality draft genomes giving insight into
the relationships between the U.S. *Csp* lineage and related pathogens. On the other hand, the still
relatively high error rate of ONT reduced our confidence in the obtained SNP-based tree, which
has the potential to give additional insights into pathogen dissemination patterns. Also, our high-
quality genomes allowed us to perform a solid pangenome analysis providing the basis for a
future taxonomic investigation, functional studies of pathogenicity and host range, and
development of new molecular markers for highly specific, high throughput detection assays.

Data Availability
The metagenomic sequencing data generated in this study have been deposited in the NCBI
Sequence Read Archive (SRA) under BioProject accession number PRJNA1290818. A total of
106 sequence datasets is available, with associated BioSample metadata provided in the SRA
submission. Genome assemblies for 17 isolates are being deposited in GenBank and will be
made publicly available upon completion of the ongoing submission process.

References

Abdelrazek S, Salamanca LR, Vinatzer BA. Metagenomic Sequencing of Tomato Plants with
Wilt Symptoms Allows for Strain-Level Pathogen Identification and Genome-Based
Characterization. Phytopathology. 2025 Apr;115(4):354-366. doi: 10.1094/PHYTO-09-24-
0279-R. Epub 2025 Apr 25. PMID: 39752554.

Ali SS, Asman A, Shao J, Firmansyah AP, Susilo AW, Rosmana A, McMahon P, Junaid M,
Guest D, Kheng TY, Meinhardt LW, Bailey BA. Draft genome sequence of fastidious
pathogen *Ceratobasidium theobromae*, which causes vascular-streak dieback in
Theobroma cacao. Fungal Biol Biotechnol. 2019 Sep 30;6:14. doi: 10.1186/s40694-019-
0077-6. PMID: 31583107; PMCID: PMC6767637.

Almagro Armenteros JJ, Salvatore M, Emanuelsson O, Winther O, von Heijne G, Elofsson A,
Nielsen H. Detecting sequence signals in targeting peptides using deep learning. Life Sci
Alliance. 2019 Sep 30;2(5):e201900429. doi: 10.26508/lsa.201900429. PMID: 31570514;
PMCID: PMC6769257.

Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J
Mol Biol. 1990 Oct 5;215(3):403-10. doi: 10.1016/S0022-2836(05)80360-2. PMID:
2231712.

Asnicar, F., Thomas, A.M., Beghini, F. *et al.* Precise phylogenetic analysis of microbial isolates
and genomes from metagenomes using PhyloPhlAn 3.0. *Nat Commun* **11**, 2500 (2020).
https://doi.org/10.1038/s41467-020-16366-7

Avershina, E., Qureshi, A.I., Winther-Larsen, H.C. et al. Challenges in capturing the mycobiome
from shotgun metagenome data: lack of software and databases. Microbiome 13, 66
(2025). https://doi.org/10.1186/s40168-025-02048-3

Avin FA, Liyanapathiranage P, Bonkowski J, Bily D, Rodriguez Salamanca L, Olson J, Wallace
S, McConnell M, Bec S, Baysal-Gurel F. Development of Molecular Markers for the
Detection of *Ceratobasidium* sp. D.P. Rogers Associated with Vascular Streak Dieback of
Woody Ornamentals in the United States. Plant Dis. 2025 May 30. doi: 10.1094/PDIS-01-
25-0005-RE. Epub ahead of print. PMID: 40445165.

Bagheri H, Severin AJ, Rajan H. Detecting and correcting misclassified sequences in the large-
scale public databases. Bioinformatics. 2020 Sep 15;36(18):4699-4705. doi:
10.1093/bioinformatics/btaa586. PMID: 32579213; PMCID: PMC7821992.

Banos, S., Lentendu, G., Kopf, A. et al. A comprehensive fungi-specific 18S rRNA gene
sequence primer toolkit suited for diverse research issues and sequencing platforms. BMC
Microbiol 18, 190 (2018). https://doi.org/10.1186/s12866-018-1331-4

Baysal-Gurel, F., and Liyanapathiranage, P. 2024. Vascular streak dieback is an emerging threat
to the Redbud nursery production in the southeastern United States (ANR-PATH-1-2024).

Tennessee State University Extension Publications. Retrieved March 8, 2024, from
https://digitalscholarship.tnstate.edu/cgi/viewcontent.cgi?article=1180&context=extension

Bhunjun CS, Phillips AJL, Jayawardena RS, Promputtha I, Hyde KD. Importance of Molecular
Data to Identify Fungal Plant Pathogens and Guidelines for Pathogenicity Testing Based
on Koch's Postulates. Pathogens. 2021 Aug 28;10(9):1096. doi:
10.3390/pathogens10091096. PMID: 34578129; PMCID: PMC8465164.

Bily D, Gyatso T, Avin FA, Bonkowski J, Liyanapathiranage P, Rodriguez Salamanca L,
Vinatzer B, Baysal-Gurel F. A *Ceratobasidium* sp. D.P. Rogers associated with vascular
streak dieback of woody ornamental plants in Virginia, U.S.A. Plant Dis. 2025 Jun 5. doi:
10.1094/PDIS-02-25-0375-RE. Epub ahead of print. PMID: 40471229.

Breitwieser, F.P., Baker, D.N. & Salzberg, S.L. KrakenUniq: confident and fast metagenomics
classification using unique k-mer counts. Genome Biol 19, 198 (2018).
https://doi.org/10.1186/s13059-018-1568-0

Brown et al, (2016), sourmash: a library for MinHash sketching of DNA, Journal of Open-
Source Software, 1(5), 27, doi:10.21105/joss.00027

Buchfink, B., Reuter, K. & Drost, HG. Sensitive protein alignments at tree-of-life scale using
DIAMOND. Nat Methods 18, 366–368 (2021). https://doi.org/10.1038/s41592-021-01101-
x

Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment
trimming in large-scale phylogenetic analyses. Bioinformatics. 2009 Aug 1;25(15):1972-3.
doi: 10.1093/bioinformatics/btp348. Epub 2009 Jun 8. PMID: 19505945; PMCID:
PMC2712344.

Català S, Pérez-Sierra A, Abad-Campos P. The use of genus-specific amplicon pyrosequencing
to assess phytophthora species diversity using eDNA from soil and water in Northern
Spain. PLoS One. 2015 Mar 16;10(3):e0119311. doi: 10.1371/journal.pone.0119311.
PMID: 25775250; PMCID: PMC4361056.

Chklovski, A., Parks, D.H., Woodcroft, B.J. et al. CheckM2: a rapid, scalable and accurate tool
for assessing microbial genome quality using machine learning. Nat Methods 20, 1203–
1212 (2023). https://doi.org/10.1038/s41592-023-01940-w

Chorlton SD. Ten common issues with reference sequence databases and how to mitigate them.
Front Bioinform. 2024 Mar 15;4:1278228. doi: 10.3389/fbinf.2024.1278228. PMID:
38560517; PMCID: PMC10978663.

Clarridge JE 3rd. Impact of 16S rRNA gene sequence analysis for identification of bacteria on
clinical microbiology and infectious diseases. Clin Microbiol Rev. 2004 Oct;17(4):840-62,
table of contents. doi: 10.1128/CMR.17.4.840-862.2004. PMID: 15489351; PMCID:
PMC523561.

Csardi G, Nepusz T (2006). "The igraph software package for complex network research."
InterJournal, Complex Systems, 1695. https://igraph.org.

Czech L, Stamatakis A, Dunthorn M, Barbera P. Metagenomic Analysis Using Phylogenetic
Placement-A Review of the First Decade. Front Bioinform. 2022 May 26;2:871393. doi:
10.3389/fbinf.2022.871393. PMID: 36304302; PMCID: PMC9580882.

De La Fuente, L., Merfa, M. V., Cobine, P. A. & Coleman, J. J. Pathogen adaptation to the
xylem environment. Annu. Rev. Phytopathol. 60, 161–186 (2022).

Delahaye C, Nicolas J. Sequencing DNA with nanopores: Troubles and biases. PLoS One. 2021
Oct 1;16(10):e0257521. doi: 10.1371/journal.pone.0257521. PMID: 34597327; PMCID:
PMC8486125.

Dinghua Li, Chi-Man Liu, Ruibang Luo, Kunihiko Sadakane, Tak-Wah Lam, MEGAHIT: an
ultra-fast single-node solution for large and complex metagenomics assembly via succinct
de Bruijn graph, Bioinformatics, Volume 31, Issue 10, May 2015, Pages 1674–1676,
https://doi.org/10.1093/bioinformatics/btv033

Donovan PD, Gonzalez G, Higgins DG, Butler G, Ito K. Identification of fungi in shotgun
metagenomics   datasets.   PLoS   One.   2018   Feb   14;13(2):e0192898.   doi:
10.1371/journal.pone.0192898. PMID: 29444186; PMCID: PMC5812651.

Duan Y, Zhou L, Hall DG, Li W, Doddapaneni H, Lin H, Liu L, Vahling CM, Gabriel DW,
Williams KP, Dickerman A, Sun Y, Gottwald T. Complete genome sequence of citrus
huanglongbing bacterium, *'Candidatus Liberibacter asiaticus'* obtained through
metagenomics. Mol Plant Microbe Interact. 2009 Aug;22(8):1011-20. doi: 10.1094/MPMI-
22-8-1011. PMID: 19589076.

Edgar, R.C. MUSCLE: a multiple sequence alignment method with reduced time and space
complexity. BMC Bioinformatics 5, 113 (2004). https://doi.org/10.1186/1471-2105-5-113

Edouard de Castro, Christian J. A. Sigrist, Alexandre Gattiker, Virginie Bulliard, Petra S.
Langendijk-Genevaux, Elisabeth Gasteiger, Amos Bairoch, Nicolas Hulo, ScanProsite:
detection of PROSITE signature matches and ProRule-associated functional and structural
residues in proteins, Nucleic Acids Research, Volume 34, Issue suppl_2, 1 July 2006,
Pages W362–W365, https://doi.org/10.1093/nar/gkl124

Ezpeleta J, Garcia Labari I, Villanova GV, Bulacio P, Lavista-Llanos S, Posner V, Krsticevic F,
Arranz S, Tapia E. Robust and scalable barcoding for massively parallel long-read
sequencing. Sci Rep. 2022 May 10;12(1):7619. doi: 10.1038/s41598-022-11656-0. PMID:
35538127; PMCID: PMC9090787.

Emms, D.M., Kelly, S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol 20, 238 (2019). https://doi.org/10.1186/s13059-019-1832-y

Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, Smit AF. RepeatModeler2 for
automated genomic discovery of transposable element families. Proc Natl Acad Sci U S A.
2020 Apr 28;117(17):9451-9457. doi: 10.1073/pnas.1921046117. Epub 2020 Apr 16.
PMID: 32300014; PMCID: PMC7196820.

Fredricks DN, Relman DA. Sequence-based identification of microbial pathogens: a
reconsideration of Koch's postulates. Clin Microbiol Rev. 1996 Jan;9(1):18-33. doi:
10.1128/CMR.9.1.18. PMID: 8665474; PMCID: PMC172879.

Geneve, R. (1991). Eastern Redbud (Cercis canadensis L.) and Judas Tree (*Cercis siliquastrum*
L.). , 142-151. https://doi.org/10.1007/978-3-662-13231-9_8.

Gil-Ordóñez, A., Pardo, J.M., Sheat, S. et al. Isolation, genome analysis and tissue localization
of *Ceratobasidium theobromae*, a new encounter pathogen of cassava in Southeast Asia.
Sci Rep 14, 18139 (2024). https://doi.org/10.1038/s41598-024-69061-8

Gíslason, M. H., Nielsen, H., Almagro Armenteros, J. J., & Johansen, A. R. (2021). Prediction
of GPI-anchored proteins with pointer neural networks. Current Research in
Biotechnology, 3, 6–13. https://doi.org/10.1016/j.crbiot.2021.01.001

Gökdemir FŞ, İşeri ÖD, Sharma A, Achar PN, Eyidoğan F. Metagenomics Next Generation
Sequencing (mNGS): An Exciting Tool for Early and Accurate Diagnostic of Fungal
Pathogens in Plants. J Fungi (Basel). 2022 Nov 13;8(11):1195. doi: 10.3390/jof8111195.
PMID: 36422016; PMCID: PMC9699264.

Goldfarb T, Kodali VK, Pujar S, Brover V, Robbertse B, Farrell CM, Oh DH, Astashyn A,
Ermolaeva O, Haddad D, Hlavina W, Hoffman J, Jackson JD, Joardar VS, Kristensen D,
Masterson P, McGarvey KM, McVeigh R, Mozes E, Murphy MR, Schafer SS, Souvorov
A, Spurrier B, Strope PK, Sun H, Vatsan AR, Wallin C, Webb D, Brister JR, Hatcher E,
Kimchi A, Klimke W, Marchler-Bauer A, Pruitt KD, Thibaud-Nissen F, Murphy TD.
NCBI RefSeq: reference sequence standards through 25 years of curation and annotation.
Nucleic Acids Res. 2025 Jan 6;53(D1):D243-D257. doi: 10.1093/nar/gkae1038. PMID:
39526381; PMCID: PMC11701664.

Gónzalez D, Rodriguez-Carres M, Boekhout T, Stalpers J, Kuramae EE, Nakatani AK, Vilgalys
R, Cubeta MA. Phylogenetic relationships of *Rhizoctonia* fungi within the Cantharellales.
Fungal Biol. 2016 Apr;120(4):603-619. doi: 10.1016/j.funbio.2016.01.012. Epub 2016 Jan
29. PMID: 27020160; PMCID: PMC5013834.

Gonzalez Garcia, V., Portal Onco, M. A. & Rubio Susan, V. Review. Biology and systematics
of the genus *Rhizoctonia*. *Span. J. Agric. Res.* **4**(1), 55–79.
https://doi.org/10.5424/sjar/2006041-178 (2006).

Grünwald, N. (2012). Novel insights into the emergence of pathogens: the case of chestnut
blight. Molecular Ecology, 21. https://doi.org/10.1111/j.1365-294X.2012.05597.x.

Guest, D., & Keane, P. (2007). Vascular-Streak Dieback: A New Encounter Disease of Cacao in
Papua New Guinea and Southeast Asia Caused by the Obligate Basidiomycete
Oncobasidium theobromae. Phytopathology, 97 12, 1654-7 .
https://doi.org/10.1094/PHYTO-97-12-1654

Handelsman J. Metagenomics: application of genomics to uncultured microorganisms.
Microbiol Mol Biol Rev. 2004 Dec;68(4):669-85. doi: 10.1128/MMBR.68.4.669-
685.2004. PMID: 15590779; PMCID: PMC539003.

Horta MAC, Steenwyk JL, Mead ME, Dos Santos LHB, Zhao S, Gibbons JG, Marcet-Houben
M, Gabaldón T, Rokas A, Goldman GH. Examination of Genome-Wide Ortholog
Variation in Clinical and Environmental Isolates of the Fungal Pathogen *Aspergillus*
fumigatus. mBio. 2022 Aug 30;13(4):e0151922. doi: 10.1128/mbio.01519-22. Epub 2022
Jun 29. PMID: 35766381; PMCID: PMC9426589.

Hu M, Li C, Xue Y, Hu A, Chen S, Chen Y, Lu G, Zhou X, Zhou J. Isolation, Characterization,
and Genomic Investigation of a Phytopathogenic Strain of *Stenotrophomonas maltophilia*.

Phytopathology. 2021 Nov;111(11):2088-2099. doi: 10.1094/PHYTO-11-20-0501-R.
Epub 2021 Nov 11. PMID: 33759550.

Irber et al. (2024). sourmash v4: A multitool to quickly search, compare, and analyze genomic
and metagenomic data sets. Journal of Open-Source Software, 9(98), 6830.
https://doi.org/10.21105/joss.06830

Jacques F, Bolivar P, Pietras K, Hammarlund EU. Roadmap to the study of gene and protein
phylogeny and evolution-A practical guide. PLoS One. 2023 Feb 24;18(2):e0279597. doi:
10.1371/journal.pone.0279597. PMID: 36827278; PMCID: PMC9955684.

Jesus Lozano-Fernandez, A Practical Guide to Design and Assess a Phylogenomic Study,
Genome Biology and Evolution, Volume 14, Issue 9, September 2022, evac129,
https://doi.org/10.1093/gbe/evac129

Johnson MA, Liu H, Bush E, Sharma P, Yang S, Mazloom R, Heath LS, Nita M, Li S, Vinatzer
BA. Investigating plant disease outbreaks with long-read metagenomics: sensitive
detection and highly resolved phylogenetic reconstruction applied to Xylella fastidiosa.
Microb Genom. 2022 May;8(5):mgen000822. doi: 10.1099/mgen.0.000822. PMID:
35584001; PMCID: PMC9465077.

Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, Wang Z. MetaBAT 2: an adaptive binning
algorithm for robust and efficient genome reconstruction from metagenome assemblies.
PeerJ. 2019 Jul 26;7:e7359. doi: 10.7717/peerj.7359. PMID: 31388474; PMCID:
PMC6662567.

Kazutaka Katoh, John Rozewicki, Kazunori D Yamada, MAFFT online service: multiple
sequence alignment, interactive sequence choice and visualization, Briefings in
Bioinformatics, Volume 20, Issue 4, July 2019, Pages 1160–1166,
https://doi.org/10.1093/bib/bbx108

Keane P, Flentje N, Lamb K. Investigation of vascular-streak dieback of cacao in Papua New
Guinea. Aust J Biol Sci. 1972;25:553–64.

Kinkar L, Gasser RB, Webster BL, Rollinson D, Littlewood DTJ, Chang BCH, Stroehlein AJ,
Korhonen PK, Young ND. Nanopore Sequencing Resolves Elusive Long Tandem-Repeat
Regions in Mitochondrial Genomes. Int J Mol Sci. 2021 Feb 11;22(4):1811. doi:
10.3390/ijms22041811. PMID: 33670420; PMCID: PMC7918261.

Kolmogorov, M., Yuan, J., Lin, Y. et al. Assembly of long, error-prone reads using repeat graphs. Nat Biotechnol 37, 540–546 (2019). https://doi.org/10.1038/s41587-019-0072-8

Krogh A, Larsson B, von Heijne G, Sonnhammer EL. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. J Mol Biol. 2001 Jan 19;305(3):567-80. doi: 10.1006/jmbi.2000.4315. PMID: 11152613.

Lauderdale, D. (2023). New vascular streak dieback fact sheet. North Carolina Cooperative Extension.

Leiva AM, Pardo JM, Arinaitwe W, Newby J, Vongphachanh P, Chittarath K, Oeurn S, Thi Hang L, Gil-Ordóñez A, Rodriguez R, Cuellar WJ. *Ceratobasidium* sp. is associated with cassava witches' broom disease, a re-emerging threat to cassava cultivation in Southeast Asia. Sci Rep. 2023 Dec 15;13(1):22500. doi: 10.1038/s41598-023-49735-5. PMID: 38110543; PMCID: PMC10728180.

Li H, Handsaker B, Wysoker A, Fennell T, Ruan J, Homer N, Marth G, Abecasis G, Durbin R; 1000 Genome Project Data Processing Subgroup. The Sequence Alignment/Map format and SAMtools. Bioinformatics. 2009 Aug 15;25(16):2078-9. doi: 10.1093/bioinformatics/btp352. Epub 2009 Jun 8. PMID: 19505943; PMCID: PMC2723002.

Li H. Minimap2: pairwise alignment for nucleotide sequences. Bioinformatics. 2018 Sep 15;34(18):3094-3100. doi: 10.1093/bioinformatics/bty191. PMID: 29750242; PMCID: PMC6137996.

Liyanage, K. H. E., Liyanapathiranage, P., & Baysal-Gurel, F. (2025). Investigating the Economic Impact of Emerging Vascular Streak Dieback Threat to Redbuds in the US Nursery Industry. HortScience, 60(7), 1244-1250. https://www.doi.org/10.21273/HORTSCI18589-25

Liyanapathiranage P, Avin FA, Bonkowski J, Beckerman JL, Munster M, Hadziabdic D, Trigiano RN, Baysal-Gurel F. Vascular Streak Dieback: A Novel Threat to Redbud and Other Woody Ornamental Production in the United States. Plant Dis. 2025 Mar 3:PDIS04240905FE. doi: 10.1094/PDIS-04-24-0905-FE. Epub ahead of print. PMID: 39115954.

Lu J, Rincon N, Wood DE, Breitwieser FP, Pockrandt C, Langmead B, Salzberg SL, Steinegger M. Metagenome analysis using the Kraken software suite. Nat Protoc. 2022

Dec;17(12):2815-2839. doi: 10.1038/s41596-022-00738-y. Epub 2022 Sep 28. Erratum in:

Nat Protoc. 2024 Aug 29. doi: 10.1038/s41596-024-01064-1. PMID: 36171387; PMCID:

PMC9725748.

Lukas Käll, Anders Krogh, Erik L.L. Sonnhammer, Advantages of combined transmembrane

topology and signal peptide prediction—the Phobius web server, Nucleic Acids Research,

Volume 35, Issue suppl_2, 1 July 2007, Pages W429–W432,

https://doi.org/10.1093/nar/gkm256

Mande SS, Mohammed MH, Ghosh TS. Classification of metagenomic sequences: methods and

challenges. Brief Bioinform. 2012 Nov;13(6):669-81. doi: 10.1093/bib/bbs054. Epub 2012

Sep 8. PMID: 22962338.

Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO Update: Novel and

Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for

Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. Mol Biol Evol. 2021 Sep

27;38(10):4647-4654. doi: 10.1093/molbev/msab199. PMID: 34320186; PMCID:

PMC8476166.

Manni M, Berkeley MR, Seppey M, Zdobnov EM. BUSCO: Assessing Genomic Data Quality

and Beyond. Curr Protoc. 2021 Dec;1(12):e323. doi: 10.1002/cpz1.323. PMID: 34936221.

Mcclellan, M. 2023. The redbud problem. https://www.nurserymag.com/article/the-

redbudproblem/

McMahon P, Purwantara A. Vascular streak dieback (*Ceratobasidium theobromae*): history and

biology. In: Bailey BA, Meinhardt LW, editors. Cacao diseases: a history of old enemies

and new encounters. New York: Springer International Publishing; 2016. p. 307–35.

Meyer, F., Fritz, A., Deng, ZL. et al. Critical Assessment of Metagenome Interpretation: the

second round of challenges. Nat Methods 19, 429–440 (2022).

https://doi.org/10.1038/s41592-022-01431-4

Moliszewska E, Maculewicz D, Stępniewska H. Characterization of three-nucleate Rhizoctonia

AG-E based on their morphology and phylogeny. Sci Rep. 2023 Oct 13;13(1):17328. doi:

10.1038/s41598-023-44448-1. PMID: 37833315; PMCID: PMC10575891.

Navgire, G.S., Goel, N., Sawhney, G. et al. Analysis and Interpretation of metagenomics data:

an approach. Biol Proced Online 24, 18 (2022). https://doi.org/10.1186/s12575-022-

00179-7

Nguyen LT, Schmidt HA, von Haeseler A, Minh BQ. IQ-TREE: a fast and effective stochastic
algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol. 2015
Jan;32(1):268-74. doi: 10.1093/molbev/msu300. Epub 2014 Nov 3. PMID: 25371430;
PMCID: PMC4271533.

Nurk S, Meleshko D, Korobeynikov A, Pevzner PA. metaSPAdes: a new versatile metagenomic
assembler. Genome Res. 2017 May;27(5):824-834. doi: 10.1101/gr.213959.116. Epub
2017 Mar 15. PMID: 28298430; PMCID: PMC5411777.

Paradis E, Schliep K. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses   in   R.   Bioinformatics.   2019   Feb   1;35(3):526-528.   doi:
10.1093/bioinformatics/bty633. PMID: 30016406.

Pedersen BS, Quinlan AR. Mosdepth: quick coverage calculation for genomes and exomes.
Bioinformatics. 2018 Mar 1;34(5):867-868. doi: 10.1093/bioinformatics/btx699. PMID:
29096012; PMCID: PMC6030888.

Pertea G, Pertea M. GFF Utilities: GffRead and GffCompare. F1000Res. 2020 Apr 28;9:ISCB
Comm   J-304.   doi: 10.12688/f1000research.23297.2.   PMID: 32489650;   PMCID:
PMC7222033.

Piombo E, Abdelfattah A, Droby S, Wisniewski M, Spadaro D, Schena L. Metagenomics
Approaches for the Detection and Surveillance of Emerging and Recurrent Plant
Pathogens. Microorganisms. 2021 Jan 16;9(1):188. doi: 10.3390/microorganisms9010188.
PMID: 33467169; PMCID: PMC7830299.

Pritchard, L., Sharma, P., Mazloom, R., Pierce, T., Irber, L., Harrington, B., Heath, L., Brown,
C.T., Vinatzer, B., 2022. genomeRxiv: a microbial whole-genome database and diagnostic
marker design resource for classification, identification, and data sharing. Access
Microbiology, 4(5).

Quevillon E, Silventoinen V, Pillai S, Harte N, Mulder N, Apweiler R, Lopez R. InterProScan:
protein domains identifier. Nucleic Acids Res. 2005 Jul 1;33(Web Server issue):W116-20.
doi: 10.1093/nar/gki442. PMID: 15980438; PMCID: PMC1160203.

Rasmussen DA, Grünwald NJ. Phylogeographic Approaches to Characterize the Emergence of
Plant Pathogens. Phytopathology. 2021 Jan;111(1):68-77. doi: 10.1094/PHYTO-07-20-
0319-FI. Epub 2020 Nov 21. PMID: 33021879.

Rizzo D, Da Lio D, Bartolini L, Francia C, Aronadio A, Luchi N, Campigli S, Marchi G, Rossi
E. DNA Extraction Methods to Obtain High DNA Quality from Different Plant Tissues.
Methods Mol Biol. 2022;2536:91-101. doi: 10.1007/978-1-0716-2517-0_5. PMID:
35819599.

Roman-Reyna V and Crandall SG (2024) Seeing in the dark: a metagenomic approach can
illuminate the drivers of plant disease. Front. Plant Sci. 15:1405042. doi:
10.3389/fpls.2024.1405042

Sahlin, K., Medvedev, P. Error correction enables use of Oxford Nanopore technology for
reference-free transcriptome analysis. Nat Commun 12, 2 (2021).
https://doi.org/10.1038/s41467-020-20340-8

Samuel T N Aroney, Rhys J P Newell, Jakob N Nissen, Antonio Pedro Camargo, Gene W
Tyson, Ben J Woodcroft, CoverM: read alignment statistics for metagenomics,
Bioinformatics, Volume 41, Issue 4, April 2025, btaf147,
https://doi.org/10.1093/bioinformatics/btaf147

Samuels GJ, Ismaiel A, Rosmana A, Junaid M, Guest D, McMahon P, Keane P, Purwantara A,
Lambert S, Rodriguez-Carres M, Cubeta MA. Vascular Streak Dieback of cacao in
Southeast Asia and Melanesia: in planta detection of the pathogen and a new taxonomy.
Fungal Biol. 2012 Jan;116(1):11-23. doi: 10.1016/j.funbio.2011.07.009. Epub 2011 Jul 23.
PMID: 22208598.

Shen, W., Le, S., Li, Y., & Hu, F. (2016). SeqKit: A Cross-Platform and Ultrafast Toolkit for
FASTA/Q File Manipulation. PLoS ONE, 11(10), e0163962.
https://doi.org/10.1371/journal.pone.0163962

Skousen, J., Monteleone, A., Tyree, M., Swab, R., Groninger, J., Adams, M., Buckley, D.,
Wood, Smit AF. Repeat-Masker Open-3.0. 2004. [Google Scholar]

Sperschneider, J. & Dodds, P. N. EffectorP 3.0: Prediction of apoplastic and cytoplasmic
effectors in fungi and oomycetes. Mol. Plant-Microb. Interact. 35(2), 146–156 (2022).

Stanke M, Keller O, Gunduz I, Hayes A, Waack S, Morgenstern B. AUGUSTUS: ab initio
prediction of alternative transcripts. Nucleic Acids Res. 2006 Jul 1;34(Web Server
issue):W435-9. doi: 10.1093/nar/gkl200. PMID: 16845043; PMCID: PMC1538822.

Sun K, Liu Y, Zhou X, Yin C, Zhang P, Yang Q, Mao L, Shentu X, Yu X. Nanopore sequencing
technology and its application in plant virus diagnostics. Front Microbiol. 2022 Jul

25;13:939666. doi: 10.3389/fmicb.2022.939666. PMID: 35958160; PMCID: PMC9358452.

Tang B, Hu X, Yue M. Contaminated bacterial genome data in the public domains: Evidence and solution. J Infect. 2024 Dec;89(6):106369. doi: 10.1016/j.jinf.2024.106369. Epub 2024 Nov 26. PMID: 39608578.

Teufel F, Almagro Armenteros JJ, Johansen AR, Gíslason MH, Pihl SI, Tsirigos KD, Winther O, Brunak S, von Heijne G, Nielsen H. SignalP 6.0 predicts all five types of signal peptides using protein language models. Nat Biotechnol. 2022 Jul;40(7):1023-1025. doi: 10.1038/s41587-021-01156-3. Epub 2022 Jan 3. PMID: 34980915; PMCID: PMC9287161.

Tian L, Huang C, Mazloom R, Heath LS, Vinatzer BA. LINbase: a web server for genome-based identification of prokaryotes as members of crowdsourced taxa. Nucleic Acids Res. 2020 Jul 2;48(W1):W529-W537. doi: 10.1093/nar/gkaa190. PMID: 32232369; PMCID: PMC7319462.

Wang, Y., Zhao, Y., Bollas, A. et al. Nanopore sequencing technology, bioinformatics and applications. Nat Biotechnol 39, 1348–1365 (2021). https://doi.org/10.1038/s41587-021-01108-x

Wang, Y., Zhao, Y., Bollas, A. et al. Nanopore sequencing technology, bioinformatics and applications. Nat Biotechnol 39, 1348–1365 (2021). https://doi.org/10.1038/s41587-021-01108-x

Wood, D.E., Lu, J. & Langmead, B. Improved metagenomic analysis with Kraken 2. Genome Biol 20, 257 (2019). https://doi.org/10.1186/s13059-019-1891-0

Woolhouse ME, Gowtage-Sequeria S. Host range and emerging and reemerging pathogens. Emerg Infect Dis. 2005 Dec;11(12):1842-7. doi: 10.3201/eid1112.050997. PMID: 16485468; PMCID: PMC3367654.):68-77. doi: 10.1094/PHYTO-07-20-0319-FI. Epub 2020 Nov 21. PMID: 33021879.

Wouter De Coster, Svenn D'Hert, Darrin T Schultz, Marc Cruts, Christine Van Broeckhoven, NanoPack: visualizing and processing long-read sequencing data, Bioinformatics, Volume 34, Issue 15, August 2018, Pages 2666–2669, https://doi.org/10.1093/bioinformatics/bty149

Yu G, Smith D, Zhu H, Guan Y, Lam TT (2017). "ggtree: an R package for visualization and
annotation of phylogenetic trees with their covariates and other associated data." Methods
in Ecology and Evolution, 8, 28-36. doi:10.1111/2041-210X.12628

Zheng P, Zhou C, Ding Y, Liu B, Lu L, Zhu F, Duan S. Nanopore sequencing technology and its
applications. MedComm (2020). 2023 Jul 10;4(4):e316. doi: 10.1002/mco2.316. PMID:
37441463; PMCID: PMC10333861.