Dengue virus harnesses mosquito Syntenin to load and secrete viral RNA into salivary

exosomes

Florian Rachenne[1], Laura Di Pietro[1], Félix Rey-Cadilhac[1], Idalba Serrato-Pomar[1], Norman

Schneider[1], Lauryne Pruvost[1], Jacques Dainat[1], Audrey Vernet[2], Chantal Cazevieille[3], Aurélie

Ancelin[4], Joséphine Lai Kee Him[4], Sander Jansen[5], Jérémy Fraering[6], Laurianne Simon[7],

Dorothée Missé[1,8], Marie Morille[7], Philippe Hammann[6], Martial Seveno[9], Kai Dallmeier[5],

Éric Marois[10], Julien Pompon[1,11,#]

1 MIVEGEC, Univ. Montpellier, IRD, CNRS, Montpellier, France

2 PHIM, Univ. Montpellier, IRD, Montpellier, France

3 INM, Electronic Microscopy Platform, Saint Eloi Hospital, Montpellier, France

4 CBS, Electronic Microscopy Platform, CNRS, Montpellier, France

5 KU Leuven Department of Microbiology, Immunology and Transplantion, Rega Institute,

Laboratory of Molecular Vaccinology & Vaccine Discovery (MVVD), Leuven, Belgium

6 Plateforme Protéomique Strasbourg-Esplanade, CNRS UAR1589, Institut de Biologie

Moléculaire et Cellulaire, Université de Strasbourg, France

7 ICGM, Univ. Montpellier, CNRS, ENSCM, Montpellier, France.

8 International Joint Laboratory PRESTO, Chiang Mai University, Thailand

9 Proteomic platform, PPM, Montpellier, France

10 IBMC, Univ. Strasbourg, CNRS, Inserm, Strasbourg, France.

11 Institute of Molecular Biosciences, Mahidol University, Thailand

corresponding author: Julien.pompon@ird.fr

Abstract

Viruses exploit extracellular vesicles (EVs) to transfer infection-enhancing viral RNAs.

However, mechanisms underlying viral RNA loading remain elusive. We leveraged our

previous discovery that dengue virus secretes transmission-enhancing subgenomic flaviviral

RNA (sfRNA) into mosquito salivary EVs to investigate viral RNA loading mechanism. We

demonstrate that sfRNA promotes the secretion of sfRNA-containing EVs marked by the

mosquito EV biogenesis protein AeSyntenin, by applying microscopy and viral genetic

editing in *in vitro* and *in vivo* models. SfRNA interacts intracellularly with mosquito

AeSyntenin and this interaction is selectively maintained within EVs as shown by

complementary RNA-affinity chromatography and RNA immunoprecipitation, and AI-based

prediction. Finally, we used systemic and salivary gland-specific protein depletion to establish

a functional role for mosquito AeSyntenin in exosome production and salivary secretion of

sfRNA. We propose that sfRNA binds AeSyntenin to drive its selective packaging and release

into exosomes, elucidating a mechanism for viral RNA incorporation into EVs.

Introduction

Viruses exploit extracellular vesicles (EVs) to facilitate infection[1,2]. EVs are non-replicative,

lipid bilayer structures secreted into the extracellular space and are broadly classified into two

main categories: microvesicles (also known as ectosomes), which bud directly from the

plasma membrane[3,4], and exosomes, which originate from the endosomal pathway[3,5–7].

Exosome biogenesis involves Syntenin-1-mediated recruitment via two PDZ domains of the

Endosomal Sorting Complex Required for Transport (ESCRT) to form intraluminal vesicles

(ILVs) within multivesicular bodies (MVBs)[8–10], which subsequently fuse with the plasma

membrane to release exosomes[11,12]. EVs are selectively loaded with RNA and act as

intercellular carriers, transferring functional RNA molecules between cells[13–16]. Viruses divert

EV carriers to package and transfer viral RNAs (complete or fragment viral genome) into

other cells, activating various strategies to enhance infection[1]. Although the presence of viral

RNA within EVs is well-documented for multiple virus families[1,17,18], the mechanism(s)

governing viral RNA loading into EVs has not been elucidated.

Our team discovered that mosquito-borne dengue viruses (DENV) – Flaviviridae

family - secrete a subgenomic flaviviral RNA (sfRNA) inside EVs from mosquito saliva[19].

SfRNA is an RNA decay product, resulting from partial degradation of the viral RNA genome

(gRNA) by 5'-3' exoribonucleases which stall on secondary structures in the 3'UTR[20].

Secondary structures include two stem loops (i.e., xrRNA1 and xrRNA2), two dumbbells (i.e.,

DB1 and DB2) and a final 3' stem loop (3'SL). SfRNA is abundant in salivary glands[21],

which produce saliva, and is deposited via salivary EVs at the mosquito bite site[19,22]. In the

bitten skin, sfRNA anti-immune properties[23] then enhance local infection and facilitate

transmission[19,22], highlighting sfRNA-containing EVs as critical enhancers of pathogen

dissemination. Here, we report how viral sfRNA is loaded within exosomes. We combined

cell biology, molecular biology and cutting-edge genome editing using *in vitro* and *in vivo*

approaches to show that sfRNA harnesses mosquito Syntenin homolog (AeSyntenin) to

induce secretion of cellular and salivary exosomes loaded with sfRNA. Our results reveal a

molecular mechanism for viral RNA packaging into EVs.

Results

DENV infection stimulates EV secretion in mosquitoes

To evaluate the impact of DENV infection on EV production, we infected mosquito cells with

low and high multiplicity of infection (MOI), and quantified EVs at 72 hours post-infection

(hpi) (Fig. 1a). Compared to the low inoculum, the high inoculum produced higher infection

intensity and quantity of secreted virions, as indicated by intracellular and extracellular gRNA

copies, respectively - and higher quantity of produced and secreted sfRNA, as indicated by

intracellular and extracellular sfRNA quantities, respectively (Fig. S1a-d). To determine the

effect on EVs, we observed that two established mosquito EV markers, AeSyntenin and the

homolog of human CD63 tetraspanin (hCD63)[3,24], increased in the cell media with infection

in a dose-dependent manner (Fig. 1b-d; Fig. S2a), indicating enhanced EV marker release. In

contrast, intracellular levels of these markers at the protein (Fig. 1e-g; Fig. S2b) and mRNA

(Fig. S3a) levels remained unchanged. Nanoparticle tracking analysis (NTA) analysis

confirmed an increase in secreted particles (Fig. 1h) and the average size (Fig. 1i) upon

infection, with a uniform concentration increase across EV subpopulations (Fig. 1j,k) and

higher proportions of larger EVs (Fig. 1l,m).

We examined the effect of DENV infection on EVs *in vivo* by infecting mosquitoes

and analyzing tissues and saliva 10 days post infection (Fig. 1o), when infection was

established and sfRNA produced in carcasses and salivary glands (Fig. S1e-h). Both

AeSyntenin and hCD63 levels were elevated in infected carcasses (Fig. 1p-r; Fig. S2c),

indicative of enhanced EV biogenesis, and in infected salivary glands (Fig. 1s-u; Fig. S2d),

suggesting an increased production of salivary EVs. Consistent with *in vitro* findings,

AeSyntenin mRNA levels remained unchanged in both carcasses and salivary glands (Fig.

S3b, c). Finally, we used TEM to assess EVs in saliva from infected mosquitoes (Fig. S4).

Salivary EVs were mostly smaller than 40 nm (Fig. S5) and to avoid biasing EV

quantification by counting virions, we did not count EVs between 40 and 60 nm. TEM

analysis showed that infection increased significantly EV secretion in saliva (Fig. 1v).

Together, these results indicate that DENV infection enhances salivary EV production, likely

by activating secretion instead of biogenesis, as intracellular EV markers remained stable..

DENV sfRNA induces secretion of EV-packaged sfRNA

To determine whether sfRNA is required for the infection-induced EV secretion, we infected

mosquito cells with a DENV infectious clone carrying deletions in the RNA pseudo-knot 1

(ΔPk1), which confers exoribonuclease resistance[25,26], to impair sfRNA biogenesis (Fig. 2a).

The ΔPk1 infectious clone had a reduced sfRNA concentration, as quantified by the

diminished sfRNA:gRNA ratio compared to wild-type infectious clone (Fig. 2b).

Interestingly, infection with ΔPk1 reduced and abrogated the infection-induced secretion of

AeSyntenin and hCD63, respectively (Fig. 2c-e; Fig. S6a). Intracellular EV marker levels

remained unchanged by infection (Fig. 2f-h; Fig. S6b), consistent with prior observations

upon infection. To assess whether sfRNA is sufficient to induce EV production, we

transfected *in vitro*-transcribed sfRNA or a size-matched viral control RNA (Ctl. RNA) and

quantified EVs 72h later (Fig. 2i). Both sfRNA and control RNA were efficiently delivered

intracellularly (Fig. S7a) and were similarly reduced by half over the three day experiment

duration (Fig. S7b,c). Strikingly, sfRNA transfection increased AeSyntenin secretion, with no

effect on hCD63 (Fig. 2j-l; Fig. S8a), while intracellular levels of both EV markers remained

unchanged (Fig. 2m-o; Fig. S8b). Although NTA analysis of ultracentrifugated EVs showed

no major change in total particle number (Fig. 2p) and mean size (Fig. 2q), size distribution

analysis revealed a specific increase in the proportion of small particles (< 60 nm) following

sfRNA transfection (Fig. 2r,s).

We next investigated whether transfected sfRNA is secreted inside lipid-based

vesicles. We submitted sfRNA present in cell media at 72h post-transfection to a RNase

resistance assay – a method previously applied to detect sfRNA packaging within EVs[19,22].

Despite multiple post-transfection washes, residual sfRNA was consistently detected in the

media immediately after transfection. This may reflect rapid secretion or contamination from

transfection reagents. To ensure that residual transfecting sfRNA did not bias the RNase

protection evaluation for secreted sfRNA, we showed that sfRNA encapsulated in transfecting

liposomes was not protected from RNase digestion (Fig. S9a). We then observed that sfRNA

from transfected cell media was partially degraded by RNase, consistent with free unprotected

sfRNA[19,22]. However, pre-treatment with detergent favored RNase degradation (Fig. 2t),

indicating that a fraction of sfRNA was protected within lipid membranes. To test whether

this packaging is specific to sfRNA, we repeated the RNase assay with Ctl. RNA, which was

similarly unprotected by liposomes (Fig. S9b). In contrast to sfRNA, Ctl. RNA was similarly

degraded with or without detergent (Fig. 2u), suggesting it was not vesicle-associated.

Together, these results suggest that sfRNA alone is sufficient to drive the secretion of

AeSyntenin-positive EVs and to be selectively packaged into EVs.

DENV sfRNA interacts with AeSyntenin

We hypothesized that sfRNA is packaged into EVs through interactions with EV-associated

proteins and investigated this by performing RNA-affinity chromatography[19,27] on mosquito

EV lysates (Fig. 3a). Compared to a size-matched Ctl. RNA, MS analysis identified 50

proteins that were enriched more than 1.5-fold with sfRNA, including proteins involved in EV

biogenesis and previously detected within EVs (Fig. 3b; Dataset S1). Notably, AeSyntenin

was enriched with sfRNA, and its interaction with sfRNA was validated by Western blot

(WB) (Fig. 3c; Fig. S10a,b). Repeating the RNA-affinity chromatography on cell lysates, we

also detected the sfRNA-AeSyntenin interaction (Fig. 3d; Fig. S10c,d), suggesting that the

interaction is initiated intracellularly. We further confirmed this interaction using RNA-

immunoprecipitation (RIP) with anti-AeSyntenin antibody and detected sfRNA bound to

AeSyntenin in both EVs from infected cells (Fig. 3f,g; Fig. S10e,f) and in DENV-infected

cells (Fig. 3h,i; Fig. S10g,h). Interestingly, gRNA was also enriched in AeSyntenin RIP

eluates from both EVs and cells (Fig. 3j,k). To evaluate AeSyntenin binding specificity, we

measured enrichment of *AeActin*, *AeRPS7*, and *AeSyntenin* mRNAs. While all three mRNAs

interacted with AeSyntenin intracellularly (Fig. S11a-c), none were recovered in RIP eluates

from EVs (Fig. S11d-f), suggesting a selective EV loading mechanism. AI-based prediction of

protein-RNA interactions with RoseTTAFoldNA[28] identified putative interactions between

the AeSyntenin PDZ domains (aa 150-331) and the RNA stems in xrRNA1, xrRNA2 and 3'

SL structures of sfRNA (Fig. S12). Collectively, these results demonstrate that DENV sfRNA

binds AeSyntenin intracellularly and that this interaction persists within secreted EVs.

AeSyntenin is required for sfRNA secretion in saliva

We started by evaluating functional homology between AeSyntenin and Human Syntenin-1.

Although we previously identified sequence homology in the functional PDZ domains

between the two proteins[24], we assessed structural homology by using AlphaFold[29] and

ScanProsite[30]. Blast between AeSyntenin and the human Syntenin-1 showed a match in the

globular structure corresponding to the PDZ domains, which were predicted with high

confidence (Fig. S13). We further assessed AeSyntenin functional homology by identifying

interacting partners in mosquito EVs using co-immunoprecipitation (co-IP) on EVs secreted

from DENV-infected mosquito cells, for relevance to AeSyntenin's role in sfRNA loading.

Among several EV-associated proteins identified (Fig. S14; Dataset S2), we detected the

mosquito homolog of human ALIX, a known partner with which Syntenin induces EV

biogenesis and cargo loading[9,10]. Our structural and interactome results suggest functional

homology between AeSyntenin and Syntenin-1.

To assess whether AeSyntenin is required for sfRNA secretion in saliva, we silenced

AeSyntenin in mosquitoes by injecting dsRNA, followed four days later, by intrathoracic

DENV inoculation. Ten days post-infection, we quantified gRNA and sfRNA in salivary

glands and saliva (Fig. 4a). While mosquito survival was unaffected at the time of inoculation,

it was reduced at day 10 in the AeSyntenin-depleted, DENV-infected group (Fig. S15a,b),

suggesting a fitness cost of AeSyntenin depletion upon infection. At the time of saliva

collection, we confirmed AeSyntenin depletion in carcasses (Fig. 4b) and salivary glands (Fig.

4c,d). In salivary glands, AeSyntenin depletion reduced gRNA levels (Fig. 4e), suggesting a

pro-viral function potentially linked to its interaction with gRNA (Fig. 3j,k). SfRNA levels

were also reduced (Fig. 4f), consistent with lower gRNA template availability; however, the

sfRNA:gRNA ratio remained unchanged (Fig. 4g), indicating that AeSyntenin does not

influence sfRNA biogenesis. In contrast, in saliva, while gRNA levels were unaffected (Fig.

4h), sfRNA levels were significantly decreased following AeSyntenin depletion (Fig. 4i),

resulting in a marked reduction in the sfRNA:gRNA ratio (Fig. 4j).

To account for the reduced salivary gland infection following AeSyntenin depletion -

which could confound the analysis of sfRNA biogenesis and secretion - we repeated the

experiment by compensating with a 140-fold higher viral inoculum in *AeSyntenin*-silenced

mosquitoes (Fig. 4k). This approach restored infection, yielding comparable levels of gRNA,

sfRNA, and sfRNA:gRNA ratios, as well as infectious particles in salivary glands (Fig. 4l-o)

and whole mosquitoes (Fig. S16), irrespective of AeSyntenin status. As observed previously

(Fig. 4h), salivary gRNA levels remained unaffected (Fig. 4p), whereas salivary sfRNA levels

were reduced upon AeSyntenin depletion (Fig. 4q), resulting in a significantly decreased

sfRNA:gRNA ratio (Fig. 4r). Supporting a lack of AeSyntenin involvement in viral particle

secretion, AeSyntenin depletion had no impact on the quantity of infectious particles in saliva

(Fig. 4s). Altogether, these results demonstrate that AeSyntenin plays a key role in secretion

of salivary sfRNA.

AeSyntenin in salivary glands is required for EV production and sfRNA secretion in

saliva

To further investigate AeSyntenin function, we genetically abrogated its expression in

mosquitoes' salivary glands. Female mosquitoes expressing Cas9 under a salivary gland-

specific promoter[31] were crossed with male mosquitoes carrying multiple guide RNAs under

ubiquitous promoters (Fig. 5a). We generated three female lines - each derived from a

different gene insertion event and labeled with different fluorophores (i.e., RFP, EGFP and

YFP) - to cross with the same male line, yielding three different progenies. Compared to the

maternal Cas9 lines, AeSyntenin levels were successfully reduced in salivary glands but not

in carcasses (Fig. 5b,c; Fig. S17a-e). Partial AeSyntenin deletion may result from the

promoter's restricted expression to the distal lateral lobes of salivary glands[31]. TEM revealed

that salivary gland-specific AeSyntenin deletion significantly lowered EV concentrations in

saliva, as estimated by EV counts per image area (Fig. 5f; Fig. S18, S19a). Notably, EVs in

the 30–40 nm range were selectively depleted (Fig. 5g,h; Fig. S19b,c).

To evaluate the impact of salivary gland-specific AeSyntenin deletion on sfRNA

salivary secretion, we infected transgenic mosquitoes by injecting the high DENV inoculum

as previously and quantified gRNA and sfRNA 10 days post-infection. Unlike systemic

RNAi, tissue-specific genetic deletion of AeSyntenin did not affect survival (Fig. S20a-c),

infection or sfRNA levels in salivary glands (Fig. 5i,j; Fig. S21a,b), resulting in comparable

sfRNA:gRNA ratios across Cas9 parental lines and most progeny (Fig. 5k; Fig. S21c) – with

the exception of one progeny (derived from the YFP Cas9 line) which had a reduced infection

and increased sfRNA:gRNA ratio (Fig. S21d-f), highlighting variability among transgenic

lines. In saliva, infection levels remained unchanged in two progeny (Fig. 5l; Fig. S21h) and

moderately increased in one (Fig. S21g), correlating with slightly higher salivary gland

infection (p = 0.07; Fig. S21a). Importantly, despite progeny variation, most lines exhibited

lower sfRNA levels (Fig. 5m; Fig. S21i,j) and all had reduced sfRNA:gRNA ratios (Fig. 5n-p)

in saliva. Together, these findings demonstrate that AeSyntenin in salivary glands specifically

contributes to EV-associated sfRNA secretion in saliva.

Discussion

Our study elucidates the first mechanism for selective loading and secretion of viral RNA into

exosomes. First, we demonstrate that infection-induced secretion of salivary EVs is partially

dependent on the production of sfRNA. Second, we confirm that sfRNA actively promotes

EV secretion by showing that sfRNA alone is sufficient to drive its own packaging into small

EVs. Third, we provide evidence that sfRNA binds both intracellularly and extracellularly to

AeSyntenin, which we identified as a key mediator of small EV production in mosquito

saliva. Finally, we establish that AeSyntenin in salivary glands is required for the secretion of

sfRNA-containing exosomes in saliva. Collectively, these findings support a model in which

sfRNA interacts with AeSyntenin to mediate its loading into exosomes, inducing their

secretion in mosquito saliva.

Although mechanisms underlying viral RNA loading into EVs remain poorly defined,

several pathways have been proposed for the selective incorporation of host RNA into

EVs[32,33]. One well-supported model involves RNA-binding proteins (RBPs) that recognize

specific RNA motifs and subsequently interact with EV biogenesis proteins to mediate cargo

sorting[34,35]. The involvement of RBPs is supported by the detection of multiple RBPs within

EVs, along with the enrichment of their inferred RNA targets. For example, hnRNPA2B1,

which contains four RNA-binding domains,[36] and the La protein[37] have been shown to bind

specific microRNAs and facilitate their inclusion into EVs. In addition, ALG-2-interacting

protein X (ALIX)—a key regulator of EV biogenesis—interacts with RBPs such as Dicer-2 to

control the selective loading of miRNAs[38]. Alternatively, direct electrostatic interactions

between RNA and membrane lipids[39] or indirect interactions through lipid-binding RBPs[40]

can drive RNA incorporation within EV membranes. In this study, we demonstrate that the

interaction between sfRNA and AeSyntenin - a key player in EV biogenesis - drives sfRNA

secretion into EVs. Based on the small size of AeSyntenin-dependent EVs and the homology

with human Syntenin-1[9], we suggest that the sfRNA-containing EVs are exosomes. While our

previous work identified sfRNA-interacting RBPs using mosquito cell lysates[19], the

AeSyntenin co-IP did not yield RBPs in the current study. This suggests that AeSyntenin-

mediated loading of sfRNA into EVs occurs independently of RBPs. Supported by the

capacity of Syntenin-1 to load host and viral proteins into exosomes[41,42], our results show that

viral RNA harnesses EV biogenesis proteins for secretion.

Overall, we used mosquito vectors and a global viral pathogen as models to delineate

the first mechanism of viral RNA loading into exosomes. Leveraging homology in EV

biogenesis, this knowledge could help understand how other viruses exploit EVs in multiple

animals.

Limitations of the study

We leveraged our expertise in mosquito-borne viral diseases to study EV loading. However,

our study system suffers from several technical limitations. The nanoliter-size volumes of

mosquito saliva preclude the use of biochemical methods usually applied for EV isolation.

Additionally, although we generated and previously characterized[24] a mosquito-specific

antibody against EV markers, we did not have access to multiple EV-targeted antibodies,

reducing our capacity to characterize mosquito EVs. Moreover, sfRNA sequence identity with

gRNA did not allow its direct quantification, and instead we relied on a well-supported[19,21,22]

indirect subtractive sfRNA quantification.

Acknowledgements

We are grateful to the VectoPole team in Montpellier, particularly Bethsabée Scheid and

Carole Ginibre for providing mosquito eggs. We thank Sarah Debaveye for help with

generating virus constructs. The authors acknowledge the ISO 9001 certified IRD i-trop HPC

(member of the South Green Platform) at IRD Montpellier for providing HPC resources that

have contributed to the research results reported within this paper. URL: https://bioinfo.ird.fr/-

http://www.southgreen.fr.

Disclosure and competing interests statement

The authors declare that they have no conflict of interest.

Funding

PhD scholarship for FR was provided by the French ministry of research and higher education

(CBS2 doctoral school) on a competitive basis. Support for this research came from French

Agence Nationale pour la Recherche (ANR-20-CE15-0006) and EU HORIZON-HLTH-2023-

DISEASE-03-18 (#101137006) to JP, from the Research Foundation Flanders (FWO) under

the Excellence of Science (EOS) program (no. 30981113; VirEOS) and the EU RIA Health

Action (no. 101137459; Yellow4FLAVI) to KD.

Material and methods

Cells and viruses. *Aedes aegypti* Aag2 and *Aedes albopictus* C6/36 (ATCC - CRL1660) were grown in Roswell Park Memorial Institute (RPMI) medium (Gibco) and Dulbecco's Modified Eagle Medium (DMEM) (Gibco), respectively, with 10% heat-inactivated fetal bovine serum (FBS) (Gibco), 1X non-essential amino acids (Gibco), and 1% penicillin/streptomycin (Gibco) at 28°C with 5% CO_2. Hamster kidney BHK-21 cells (ATCC – CL10) were grown in RPMI medium with 5% heat-inactivated FBS and 1% penicillin/streptomycin at 37°C, 5% CO_2.

DENV-2 New Guinea C strain (NGC) was obtained from ATCC (VR-1584). Yellow fluorescent protein (mCitrine)-tagged NGC DENV2-ΔPK1 mutant reporter virus was generated using standard molecular biology techniques, i.e. side-directed PCR mutagenesis (KapaHF HotStart ReadyMix, Roche) and homologous recombination in yeast[43], to modify infectious molecular clone pShuttle-DV2/mCherry[44]. To abolish sfRNA expression, a three-nt base exchange was introduced in the DENV2 3'-UTR cDNA impairing the intramolecular base-paring between the top of stem-loop (SL II) and its proximal downstream RNA required for the formation of pseudoknot structure 1 (pk1)[20]. Recombinant virus was rescued by plasmid DNA transfection of BHK21 cells using TransIT-LT1 transfection reagent (MirusBio), amplified by passage in C6/36 mosquito cells and quantified by plaque titration as previously described[43]. Wild-type NGC infectious clone were similarly produced with mCherry sequence at the same position as mCitrine.

Mosquitoes. *Aedes aegypti* mosquitoes from the Bora Bora colony[45] were maintained in BugDorm cages (BugDorm) at 27 ± 1°C, with 70 ± 5% relative humidity, under a 12 h:12 h

light:dark cycle, and provided with *ad libitum* access to a 10% sucrose solution. Larvae were

reared at 28°C in Milli-Q water supplemented with fish food flakes (TetraMin) until pupation.

Cell infection. For IP, 1.8 x 10^7 Aag2 cells were incubated with DENV at a MOI of 1 in 5 ml

of FBS-free cell media for 1 h. For WB and RT-qPCR analyses, 2.5 x 10^5 Aag2 cells were

incubated with DENV at a MOI of 0.05 or 1 in 150 µl of FBS-free cell media for 1 h. For

NTA, 8 x 10^6 Aag2 cells were incubated with DENV at a MOI of 0.05 or 1 for 1 h in 2 ml of

FBS-free cell media for 1 h. After removing the inoculum, cells were incubated with complete

media containing 2% EV-free FBS for 72 h.

Quantification of gRNA, sfRNA and Ctl RNA. Total RNA was extracted using E.Z.N.A.

Total RNA Kit I and eluted in DEPC-treated water. gRNA, wild-type sfRNA/3'UTR, Pk1

mutated sfRNA/3'UTR and Ctl RNA were quantified by RT-qPCR using the iTaq SYBR

GREEN One-Step Kit (Bio-Rad) with primers detailed in Table S1. A total reaction volume

of 10 µl containing 0.3 µl of 300 nM of forward primer, 0.3 µl of 300 nM reverse primer, 5 µl

of iScript master mix, 2.275 µl of water nuclease-free water and 2 µl of RNA extract was

amplified using LightCycler 96 (Roche) with the following thermal profile: 50°C for 15 min,

95°C for 1 min, followed by 40 cycles of 95°C for 10 sec and 60°C for 15 sec, with a final

melting curve analysis.

Absolute quantification was achieved by amplifying the qPCR templates using

forward T7-tagged primers detailed in table S1. Templates were transcribed using the

MegaScript T7 Kit and purified with the E.Z.N.A. Total RNA Extraction Kit I. RNA

concentrations were measured using a NanoDrop spectrophotometer (Thermo Fisher

Scientific) and serial dilutions were used to generate absolute standard curves.

SfRNA copies were calculated by subtracting gRNA copies (quantified using primers

targeting the envelop) from the combined sfRNA and 3'UTR copy numbers (quantified using

sfRNA/3'UTR primers), as previously developed[21]. For samples containing detectable levels

of sfRNA, sfRNA:gRNA ratio was calculated by dividing sfRNA to gRNA copy numbers.

Western Blot (WB). WB analyses were conducted on eluates from RNA affinity

chromatography and IP, and on lysates from Aag2 cells, cell media, whole mosquitoes and

salivary glands. Cell lysate was obtained by adding RIPA buffer IV (Bio Basic),

centrifugating the lysate at $20,000 \times$ g for 20 min and collecting the supernatant. Cell media

was pre-cleared by centrifugation at 3,500 x g for 10 min and lysed in RIPA buffer IV.

Mosquito and salivary gland lysates were obtained by homogenizing 5 carcasses and 10

salivary glands in 250 or 150 µl of RIPA buffer IV, respectively, with silica beads (BioSpec)

using a BeadBeater homogenizer (MP Biomedicals) and collecting supernatants after

centrifugation at 14,000 rpm for 10 min at 4°C. Proteins from eluates and lysates were

quantified using Qubit protein assay (ThermoFisherScientific). Homogenized quantities of

proteins were diluted in reducing Laemmli SDS buffer, heated to 95°C for 10 min, separated

on NuPAGE 4-12% acrylamide gel (Invitrogen) and transferred onto a 0.2 µm nitrocellulose

membrane (Bio-Rad) using Trans-Blot turbo Transfer System (Bio-Rad). Membranes were

washed with 0.1% Tween 20 in PBS, blocked with 5% milk (Régilait) for 1 h at RT,

incubated overnight at 4°C with primary antibodies diluted in 0.1% Tween 20 in PBS.

Antibodies included 1:500 rabbit anti-human CD63 (Ab134045,clone EPR5702, Abcam),

1:500 mouse anti-human pan-actin (MA5-11869, clone ACTN05, Thermo Scientific), and

1:500 rabbit custom-made anti-AeSyntenin[46]. Secondary antibody staining was performed

with 1:2,000 HRP-conjugated anti-rabbit IgG (7074, Cell Signaling Technology) or 1:2,000

HRP-conjugated anti-mouse IgG (7076, Cell Signaling Technology) in 0.1% Tween 20 5%

milk in PBS for 1 h at RT. Blots were revealed with SuperSignal West Pico PLUS

Chemiluminescent Substrate (Thermo Scientific) or SuperSignal West Femto

Chemiluminescent Substrate (Thermo Scientific) using a ChemiDoc MP Imaging System

(Bio-Rad) and analyzed with ImageLab Software (Bio-Rad). Blot intensity was estimated

with ImageLab and normalized to actin level.

Nanoparticle Tracking Assay (NTA). Cell media was diluted by half in PBS and analyzed

using a NanoSight NS300 instrument (Malvern) equipped with a 488 nm laser. The capture

settings were as follows: camera type sCMOS, laser type Blue488, camera level 12, slider

shutter 1206, slider gain 366, frame rate (FPS) 25.0, number of frames 1498, temperature 25.0

°C, viscosity 0.9 cP (for water), and syringe pump speed 40. Five 1-minute videos were

recorded, and particle analysis was conducted using NanoSight NTA software version 3.4

(build 3.4.4, Malvern). NTA analyses were performed at the Charles Gerhardt Institute,

Montpellier, France.

Infection of mosquitoes. Three-to-five day-old female mosquitoes were intrathoracically

microinjected with 5 FFU of wild-type DENV in a single injection of 69 nl using Nanoject II

(Drummond). An equivalent volume of RPMI was injected as control. DsRNA-injected

female mosquitoes were microinjected with 5 or 700 FFU of wild-type DENV at 4 days post

dsRNA injection. Mosquitoes were maintained on a 10% sugar solution for an additional 10

days before analysis.

Extracellular vesicle (EV) isolation. EV-free FBS was produced by ultracentrifugating heat-

inactivated FBS at $100,000 \times g$ for 18 h using an S58-A rotor (k-factor 50, Thermo Scientific)

in a Sorvall MX Plus Series Floor Model Micro-Ultracentrifuge (Thermo Scientific). EV-free

FBS was collected by leaving approximately 1 ml of residual volume at the bottom of the 8

ml tube, filtered through a 0.22 µm filter (Sartorius), and stored at 4°C. 1.8 x 10^6 Aag2 cells

per T175 flask were maintained in complete medium supplemented with 2% EV-free FBS for

3 days. Cell media was then centrifuged at 3,500 × g for 5 min to remove cell debris. The

clarified supernatant was ultracentrifuged at 100,000 × g for 3 h to obtain an EV-

concentrated pellet. For NTA, the EV-concentrated pellet was washed in PBS and

ultracentrifuged one more time at 100,000 x g for 3h before resuspension in PBS.

Collection of mosquito saliva and tissues. Mosquitoes were starved for 24h, cold-

anesthetized, and immobilized by removing their wings and legs. The proboscis of each

mosquito was individually inserted into a 20 µl pipette tip containing 10 µl of pre-warmed

DMEM supplemented with 0.5 mM ATP (Invitrogen) and 1% Erioglaucine (25 mM,

Sigma-Aldrich) for 1 h[22]. Mosquitoes with blue-colored abdomens, indicating ingestion of

the dye, were considered as having salivated. For TEM observation, saliva was collected in

bulk. Hundreds of mosquitoes were let to feed on artificial feeding system (Hemotek)

containing 3 ml of DMEM supplemented with 0.5 mM ATP and 1% Erioglaucine covered

with a silicone membrane for 2 h. The DMEM media was then precleared by centrifugation at

1,500 g for 10 min at 4°C, ultracentrifuged at 100,000 g for 3 h at 4°C and resuspended in

30 µl of PBS. Number of salivating mosquitoes was estimated by counting the number of

blue-abdomen mosquitoes, indicating salivation and quantities of EVs were normalized per

mosquitoes between infected and non-infected samples. For RNA quantification, individual

saliva samples were transferred into 200 µl of TRK lysis buffer. For FFU assays, pooled

saliva samples from 10 mosquitoes was combined with 100 µl of RPMI medium

supplemented with 1% P/S and 1% antibiotic-antimycotic solution (Gibco). Following saliva

collection, salivary glands were dissected using fine forceps. For RNA quantification, 5

salivary glands were pooled in 200 µl of TRK lysis buffer. For FFU assays, 5 salivary glands

were pooled in 150 µl of RPMI 1% P/S and 1% antibiotic-antimycotic. For WB, 5 or 10

salivary glands were pooled in 150 µl of RIPA buffer IV. Remaining carcasses were

collected and processed similarly by pools of five. Tissues and saliva samples were

homogenized using silica beads with a BeadBeater homogenizer. RNA was extracted using

the EZNA Total RNA Kit I. For FFU assays, homogenized tissues and saliva were sterilized

by filtration through a 0.22 µm filter (Millex-GV). For WB, lysates from salivary glands and

carcasses were pre-cleared by centrifugation at 3,500 × g for 10 min.

Transmission electron microscopy (TEM). For uninfected and infected EV saliva, 3 µl of

ultracentrifuged EV solution was absorbed on a glow-discharged 300 mesh formvar carbon-

coated grid (Electron Microscopy Science) and stained with 1% uranyl acetate in two steps.

Grids were observed using a JEOL 1400 Flash transmission electron microscope at 120kV.

Micrographs were recorded using a One View camera (Gatan Inc.). Experiment was

performed at the Center of Structural Biology of Montpellier, CNRS, France. For EV saliva

from transgenic mosquitoes, 7 µl of ultracentrifuged salivary EV solution was loaded onto a

formvar-coated copper grid 100 mesh (EMS FCF100-Cu-50), left to settle for 2 min, and

dried with a blotting paper. The grid was stained with 1% uranyl acetate and analyzed using a

TEMF20 operating at 120 KV (Tecnai) with a Veleta numeric camera (Olympus) at the

Neuroscience institute of Montpellier, INSERM U1298, France.

Production of sfRNA and control RNA (Ctl RNA). sfRNA and Ctl RNA, corresponding to

a DENV NS2 fragment[27], were amplified from viral cDNA using T7 promoter-tagged

forward primers. sfRNA and Ctl. RNA primers are detailed in table S1. PCR products were

size-validated by agarose gel electrophoresis, purified using the QIAquick PCR Purification

Kit, and *in vitro* transcribed using MEGAscript T7 Kit. Transcripts were purified with the

E.Z.N.A. Total RNA Kit I and quantified via NanoDrop. RNA fragments were treated with

RNA 5' polyphosphatase (Biosearch Technologies), and re-purified and quantified as above.

Samples were incubated at 95 °C for 5 min and gradually cooled to 4 °C for folding.

Transfection of sfRNA and Ctl RNA. 10^{11} and 10^{12} copies of monophosphorylated folded

sfRNA and Ctl RNA, respectively, were transfected for 1 h using TransIT mRNA

Transfection Kit (Euromedex) into 2.5 x 10^5 Aag2 cells for WB analysis or into 1.2 x 10^6

Aag2 cells for 1 h for NTA analysis. Difference between sfRNA and Ctl RNA amounts was to

compensate for a lower transfection efficiency of Ctl RNA (Fig. S7a). Transfection media was

removed and cells were washed three times with pre-warmed RPMI medium. After three days

of incubation with 2% EV-free FBS 1% P/S 1X NEAA RPMI, media was collected, pre-

cleared by centrifugation at 3,500 x g for 10 min and lysed in RIPA buffer IV. For NTA

analysis, pre-cleared cell media was ultracentrifuged at 100,000 × g for 3 h, the pellet

washed, ultracentrifuged at 100,000 x g for 3 h, and resuspended in 500 µl PBS. Aag2 cells

were lysed in TRK lysis buffer (EZNA Total RNA Kit I) for RNA quantification or in 100 µl

of RIPA buffer IV for WB.

RNAse resistance assay. Three days post sfRNA and Ctl RNA transfection, 1 ml of cell

media was pre-cleared at 3,500 x g for 10 min, divided into equal volumes of 4 x 200 µl and

treated with different conditions. Samples were treated with 2.5 µl of 0.1 % triton X 100

(Sigma or PBS as control and incubated at 4°C at 30 min. Then, 5 µl of RNase A/T1

(Thermofisher Scientific) or PBS as control was added and samples were incubated at 37°C

for 30 min before RNA extraction using E.Z.N.A. Total RNA extraction Kit I.

RNA affinity chromatography. RNA affinity chromatography was performed as previously

described[27] with minor modifications. 1 ml of NHS Mag Sepharose bead (Cytiva) flurry was

washed four times with 1 ml of 1 mM HCl (Carlo Erba) and incubated in 950 µl of coupling

buffer [0.2 M NaHCO□ (Sigma), 0.5 M NaCl (Sigma), pH 8.3] with 50 µl of 5 mM

tobramycin (Sigma) overnight at 4°C under gentle rotation. Tobramycin beads were blocked

in 1 ml of blocking buffer [20 mM Tris pH 7.0 (Sigma), 1 mM $CaCl_2$ (Sigma), 1 mM $MgCl_2$

(Sigma), 300 mM KCl (Sigma), 0.1 mg/ml of tRNA (Ambion), 0.5 mg/ml BSA (Sigma), 0.2

mM Dithiothreitol (Biorad), 0.01% NP-40 (MP Biomedical)] for 1 h at 4°C, washed three

times with PBS and resuspended in 500 µl of PBS. DENV sfRNA and a same-size control

RNA fragment both flanked by a tobramycin aptamer sequence were amplified from

plasmids[27,47] using T7-tagged forward primers detailed in table S1. After size validation on

agarose gel, PCR products were purified using the QIAquick PCR Purification Kit (Qiagen)

and transcribed using MEGAscript T7 Kit (Thermo Fisher). Transcribed RNA was extracted

with EZNA Total RNA Kit I (Omega) and quantified using Nanodrop. 30 µg of *in vitro*-

transcribed sfRNA or control RNA in 200 µl of RNA binding buffer (20 mM Tris pH 7.0, 1

mM CaCl□, 1 mM MgCl□, 75 mM NaCl, 145 mM KCl, 0.1 mg/ml yeast tRNA, 0.2 mM

Dithiothreitol) was heated to 95°C for 5 min and slowly cooled to RT to enable RNA folding.

The RNA solution was incubated with 30 µl of tobramycin-coupled beads for 1 h at 4°C

under end-over-end rotation. Beads were washed once with 250 µl of RNA washing buffer

(20 mM Tris pH 7.0, 1 mM CaCl□, 1 mM MgCl□, 75 mM NaCl, 145 mM KCl, 0.1% NP-40,

0.2 mM Dithiothreitol).

1.8×10^7 uninfected Aag2 cells or isolated EVs were lysed in 1 ml and 500 µl of lysis

buffer [50 mM Tris pH 7.5 (Sigma), 150 mM NaCl, 1% NP-40, 1 X protease and phosphatase

inhibitor (Sigma)], respectively. 417 µl of cell lysate in 500 µl of 1 X modified protein buffer

(2 mM $CaCl_2$, 2 mM $MgCl_2$, 290 mM KCl, 4 mM Dithiothreitol) or 210 µl of EV lysate in

300 µl of 1 X modified protein buffer were pre-cleared with 30 µl of tobramycin bead flurry

during 1 h at 4 °C and then incubated with beads from 30 µl of RNA-bead flurry for 2 h at

4°C with end-over-end rotation. Beads were washed four times in 500 µl of 145 mM protein

washing buffer (20 mM Tris pH 7.0, 1 mM CaCl☐, 1 mM MgCl☐, 75 mM NaCl, 145 mM

KCl, 0.5% NP-40). Elution for MS analysis was performed by incubating beads in 20 µl of

elution buffer (20 mM Tris pH 7.0, 1 mM CaCl☐, 3 mM MgCl☐, 145 mM KCl, 5 mM

tobramycin, 0.2 mM Dithiothreitol) for 5 min on ice and elution for WB analyses was

performed in 25 µl of Laemmli SDS buffer (Thermo Scientific) and heated to 95 °C for 5 min.

Mass spectrometry (MS). Eluates from RNA affinity chromatography were analyzed at the

PPM laboratory (Pôle Protéome de Montpellier, France). Briefly, peptide samples were

reconstituted in 10 µl of buffer A (0.1% formic acid) and injected into nano-flow HPLC

system (Ultimate 3000 RSLC, Thermo Fisher Scientific) coupled to a mass spectrometer

equipped with a nanospray source (Q-Exactive HF, Thermo Fisher Scientific). Peptides were

separated on a capillary column (0.075 mm × 500 mm, Acclaim PepMap 100, reverse-phase

C18, NanoViper, Thermo Fisher Scientific) using a gradient of 2-40% B over 120 minutes

with A = 0.1% formic acid and B = 0.1% formic acid in 80% acetonitrile, at a flow rate of 300

nl/min (total runtime: 150 min). Spectra were recorded using Xcalibur 4.1 software (Thermo

Fisher Scientific) with the method file 120LowQtt.meth. Spectral data were analyzed with

MaxQuant v1.6.10.43, including the script leading FPP v3.2. The following databases were

used: RefProteome_Aedes_aegypti-cano_2021_01_UP000008820_7159.fasta,

Uniprot_SPOFR_2020_02.fasta,   VectorBase-

52_AaegyptiLVP_AGWG_AnnotatedProteins.fasta,   and   contaminants_fpp_180320.fasta.

Data validation was performed using a 1% FDR filter at both peptide and protein levels.

Proteins with more than 1.5 fold enrichment with LFQ ratio and 2 fold enrichment with

MS/MS pseudoRatio were considered significantly enriched. The mass spectrometry

proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE[48]

partner repository with the dataset identifier PXD065152. (Username:

reviewer_pxd065152@ebi.ac.uk , Password: jQGpLX9honIs).

Eluates from AeSyntenin co-IP were analyzed by the Proteomic Platform of

Strasbourg-Esplanade (PPSE) at the Institut de Biologie Moléculaire et Cellulaire (IBMC).

Briefly, proteins were first precipitated overnight using 0.1 M ammonium acetate in 100%

methanol. After centrifugation at 12,000g and 4°C for 15 min, the pellets were washed twice

with 0.1 M ammonium acetate in 80% methanol. This precipitation step was repeated once

more. Then, proteins were dried using a Speed-Vac concentrator. The dried pellets were

resuspended in 100 µL of 50 mM ammonium bicarbonate, reduced with 5 mM dithiothreitol

at 95°C for 10 minutes, and alkylated with 10 mM iodoacetamide at room temperature for 30

min in the dark. Samples were digested with 150 ng sequencing-grade trypsin (Promega) at

37°C overnight. Peptides were resuspended in 0.1% formic acid, and 3/64 of the sample was

loaded for nLC-MS/MS analysis. Samples were analyzed using a timsTOF Pro 2 coupled with

a NanoElute 2 (Bruker Daltonik GmbH) in Data-Independent Acquisition and Parallel

Accumulation Serial Fragmentation (DIA-PASEF) mode. Peptides were separated on an

IonOpticks Aurora Elite column (25 cm × 75 µm, 1.7 µm particle size, and 120 Å pore size;

AUR3–15075C18-CSI) with a 45-minute elution gradient. Data were searched against the

Aedes aegypti database from UniProt (v2021_03, 33 998 sequences) using DIANN 1.9

software with a library-free approach keeping most of the default parameters. One missed

cleavage and one variable modification were allowed (M-Ox, Ac-Nterm), and match between

runs was enabled. Prostar software (1.34.5) was used for statistical analyses. Imputation of

missing values was based on the 1% detection quantile, and a LIMMA statistical test was

used for each comparison. The mass spectrometry proteomics data from the

immunoprecipitations have been deposited to the ProteomeXchange Consortium via the

PRIDE partner repository with the dataset identifier PXD065851 (Username:

reviewer_pxd065851@ebi.ac.uk , Password: GmeflV8jgwL9).

Immunoprecipitation (IP) and co-immunoprecipitation (co-IP). Infected Aag2 cells and

EVs from infected Aag2 cells were lysed in 500 µl of NT2 buffer [200 mM KCl, 20 mM

HEPES pH 7.2, 2% N-dodecyl-β-D-maltoside (Thermofisher Scientific), 1% Igepal (Sigma),

100 U/mL Murine RNase inhibitor (Biolabs)] and supernatant was collected after

centrifugation at 13,000 × g for 13 min at 4°C. 100 µl of SureBeads Protein G Magnetic

Beads (Bio-Rad) were washed with 500 µl of 0.01% Tween 20 in PBS, blocked with 50 µl of

1% BSA 0.01% Tween 20 in PBS for 10 min at 4°C, washed with PBS and incubated for 1 h

at 4°C with 500 µl of 0.01% Tween 20 in PBS containing 2.5 µg of custom-made anti-

AeSyntenin antibody[46] on a disk rotator. Beads were similarly incubated with anti-rabbit IgG

antibody (Invitrogen) as control. After three washes with 500 µl of 0.01% Tween 20 in PBS,

beads were incubated with homogenized volumes of lysed proteins for 2 h at 4°C on a rotator.

1 % of initial input was kept for analysis by WB as control. After washes, beads diluted in

500 µl of 0.01% Tween 20 in PBS were separated into aliquots of 100 µl for WB analysis and

400 µl for RNA quantification. WB samples were resuspended in 30 µl of Laemmli SDS

buffer and warmed at 95°C for 5 min. RNA quantification samples were resuspended in 350

µl of TRK lysis buffer before extraction with EZNA Total RNA extraction Kit I. For co-IP,

all beads were resuspended in 50 µl of Laemmli SDS buffer and heated at 95°C for 5 min.

Elution was repeated once with 50 µl of Laemmli SDS buffer to obtain a final elution volume

of 100 µl.

Affinity of the anti-AeSyntenin antibody for sfRNA was assessed by coupling

SureBeads Protein G Magnetic Beads with either anti-rabbit IgG or rabbit anti-AeSyntenin

antibody as above and incubating them with $10^{1\square}$ copies of *in vitro*-transcribed sfRNA for 2

h at 4$\square$°C on a disk rotator. Elution for RNA extraction was performed as above.

RoseTTAFold2NA prediction. Interaction predictions between the 3D structures of

xrRNA1, xrRNA2, xrRNA1-2, SL1-SL2, DB1, DB2, 3'SL and the AeSyntenin was

performed on RoseTTAFold2NA version 0.2[49]. Molecular visualization was performed with

Molstar[50].

Quantification of mRNA. Total RNA was extracted using the EZNA Total RNA Kit I,

cleared of genomic DNA using DNAse following protocol from gDNA Clear cDNA

Synthesis kit (Biorad), quantified with a NanoDrop, normalized, and reverse transcribed using

iScript gDNA Clear cDNA Synthesis Kit (Bio-Rad). *AeSyntenin* mRNA was quantified in a

total reaction of 10 µl, containing 2 µl of Hot FirePol SolisGreen qPCR mix (Euromedex), 0.3

µl of 300 nM forward, 0.3 µl of 300 nM reverse primers (Table S1), 5.4 µl of nuclease-free

water and 2 µl of cDNA. qPCR thermal profile consisted of 95°C for 5 min, followed by 40

cycles of 95°C for 15 sec, 60°C for 20 sec, and 72°C for 20 sec, with a final melting curve

analysis on a LightCycler 96 (Roche). *Actin* and *RPS7* mRNA were quantified as

housekeeping genes and the geometric mean of their Ct values was used for normalization

using the delta-delta Ct method.

RNAi-mediated silencing. Target sequences were amplified from Aag2 cDNA using GoTaq

Master Mix (Promega) and 400 nM T7-flanked primer pairs (Table S1) designed via the E-

RNAi website[51] with the following thermal profile: 95°C for 5 min; 45 cycles of 95°C for 15

sec., 55°C for 15 sec., and 72°C for 45 sec.; 72°C for 10 min. After product size confirmation

on agarose gel, PCR products were purified using the QIAquick PCR Purification Kit and

transcribed overnight with the MEGAscript T7 Kit. RNA was extracted with the EZNA Total

RNA Kit I, adjusted to 14 µg/µl, and annealed by heating at 95°C for 5 min followed by

gradual cooling. Negative control dsRNA targeting LacZ[52] was similarly produced (Table

S1). Four-day-old female mosquitoes were cold-anesthetized and injected with 138 nl of

dsRNA using Nanoject II microinjector (Drummond Scientific). Mosquitoes were maintained

on a 10 % sugar solution.

Focus forming unit (FFU) assay. Samples were subjected to a four-fold serial dilution in

RPMI medium and 150 µl of each dilution was incubated with 1.5×10^5 BHK-21 cells for 1 h

on a rocking platform. After removing the inoculum, cells were overlaid with 500 µl of 1%

carboxymethyl cellulose (CMC) (Sigma) in 2% FBS (Gibco) RPMI medium. After 5 days,

cells were fixed with 4% paraformaldehyde (VWR) in PBS, permeabilized with 0.3% Triton

X-100 (Euromedex), blocked with 1% FBS in PBS, and stained using a 1:200 mouse

monoclonal anti-envelope antibody (4G2, provided by Dr Subhash Vasudevan, Duke-NUS

Medical School, Singapore) and 1:500 secondary anti-mouse IgG Alexa Fluor 488

(Invitrogen). FFU were counted using the EVOS m5000 microscope (Invitrogen) in three

replicates per dilution, and average FFU per sample was calculated per ml.

Salivary gland-specific abrogation of AeSyntenin. Golden Gate Cloning was used to

construct a first piggyBac transgenesis vector with Cas9 under the control of the 30Ka

promoter[31]. In a second piggyBac transgenesis vector, different guide RNAs targeting

AeSyntenin were placed under the control of four different U6 promoters: protospacer

GACGAGTCCTTAACGATGG(CGG) under control of U6 promoter AAEL017774;

GTCACCAACGGAATTCGAG(AGG)   under   U6   AAEL017763;

GATACTTCCGGTTGGTAAG(CGG)   under   U6   AAEL017774,   and

GGGATTGGAGCTCAGCCGTG(AGG) under U6 AAEL017702. The sequences of the

transgenesis plasmids are provided in Fig. S22 and DataSet S3 and S4. Transgenic

mosquitoes were obtained by injecting embryos of the Liverpool strain with 240 ng/l of

piggyBac plasmid mixed with 80 ng/µl of transposase helper plasmid[53]. Injected mosquitoes

reaching adulthood and showing transient expression of the fluorescence marker were

backcrossed to wild-types. In the next generation (G1), transgenic larvae were selected based

on expression of the plasmid's fluorescence markers[54]. Single G1 females were crossed to

wild-type males. Transgenes showing about 50% inheritance, indicative of probable single

insertions, were retained and the G2 positives of each progeny were self-crossed to establish

lines. In the G3 generation, homozygous larvae were selected by their stronger fluorescence to

establish stable lines. Three lines with Cas9 expression were produced each with a different

selection fluorescent marker (i.e., RFP, EGFP, YFP), while one line with gRNA was

produced. From the guide RNA lines, seven homozygous males were mass-crossed with

twenty virgin Cas9-expressing females. The resulting eggs were allowed to develop for 3 to 5

days before hatching. Individual containers were provided with blood meals. Blood-fed

females were then transferred to separate boxes for oviposition. After egg-laying, the egg

papers were removed, and the eggs were allowed to mature for at least five days before

hatching the cohorts selected for experiments.

Statistics. One-tailed T-test was applied to test differences in log-transformed gRNA, sfRNA

copy numbers and fold changes, in non-transformed sfRNA:gRNA ratio, AeSyntenin and

hCD63 protein quantities. Variations in concentration and proportion of particles by size were

tested using X^2 test. Differences in RNA quantities after RNase assay and in AeSyntenin and

hCD63 protein quantities upon different infection MOI were assessed with Fisher's LSD

multiple comparison test. Statistical analyses were conducted using Prism v8 (GraphPad).

References

1.   Rey-Cadilhac, F., Rachenne, F., Missé, D. & Pompon, J. Viral Components Trafficking with(in) Extracellular Vesicles. *Viruses* **15**, 2333 (2023).

2.   Zhu, X., Lin, X., Hu, L., Wang, L. & Zhu, Q. Harnessing crosstalk between extracellular vesicles and viruses for disease diagnostics and therapeutics. *Extracell Vesicles Circ Nucleic Acids* **5**, 458–60 (2024).

3.   Théry, C. *et al.* Minimal information for studies of extracellular vesicles 2018 (MISEV2018): a position statement of the International Society for Extracellular Vesicles and update of the MISEV2014 guidelines. *Journal of Extracellular Vesicles* **7**, 1535750 (2018).

4.   Couch, Y. *et al.* A brief history of nearly EV□erything – The rise and rise of extracellular vesicles. *J of Extracellular Vesicle* **10**, e12144 (2021).

5.   Jadli, A. S., Ballasy, N., Edalat, P. & Patel, V. B. Inside(sight) of tiny communicator: exosome biogenesis, secretion, and uptake. *Mol Cell Biochem* **467**, 77–94 (2020).

6.   Zhang, Y., Liu, Y., Liu, H. & Tang, W. H. Exosomes: biogenesis, biologic function and clinical potential. *Cell Biosci* **9**, 19 (2019).

7.   Van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of extracellular vesicles. *Nat Rev Mol Cell Biol* **19**, 213–228 (2018).

8.   Zimmermann, P. *et al.* Characterization of Syntenin, a Syndecan-binding PDZ Protein, as a Component of Cell Adhesion Sites and Microfilaments. *MBoC* **12**, 339–350 (2001).

9.   Baietti, M. F. *et al.* Syndecan–syntenin–ALIX regulates the biogenesis of exosomes. *Nat Cell Biol* **14**, 677–685 (2012).

10.   Ghossoub, R. *et al.* Syntenin-ALIX exosome biogenesis and budding into multivesicular bodies are controlled by ARF6 and PLD2. *Nat Commun* **5**, 3477 (2014).

11. Friand, V., David, G. & Zimmermann, P. Syntenin and syndecan in the biogenesis of

exosomes: Syndecan-syntenin pathway in exosome biogenesis. *Biol. Cell* **107**, 331–341

(2015).

12. Van Niel, G., D'Angelo, G. & Raposo, G. Shedding light on the cell biology of

extracellular vesicles. *Nat Rev Mol Cell Biol* **19**, 213–228 (2018).

13. Valadi, H. *et al.* Exosome-mediated transfer of mRNAs and microRNAs is a novel

mechanism of genetic exchange between cells. *Nat Cell Biol* **9**, 654–659 (2007).

14. Mittelbrunn, M. *et al.* Unidirectional transfer of microRNA-loaded exosomes from T

cells to antigen-presenting cells. *Nat Commun* **2**, 282 (2011).

15. Pegtel, D. M. *et al.* Functional delivery of viral miRNAs via exosomes. *Proc. Natl.*

Acad. Sci. U.S.A. **107**, 6328–6333 (2010).

16. Zomer, A. *et al.* In Vivo Imaging Reveals Extracellular Vesicle-Mediated

Phenocopying of Metastatic Behavior. *Cell* **161**, 1046–1057 (2015).

17. Bukong, T. N., Momen-Heravi, F., Kodys, K., Bala, S. & Szabo, G. Exosomes from

Hepatitis C Infected Patients Transmit HCV Infection and Contain Replication Competent

Viral RNA in Complex with Ago2-miR122-HSP90. *PLoS Pathog* **10**, e1004424 (2014).

18. Caobi, A., Nair, M. & Raymond, A. D. Extracellular Vesicles in the Pathogenesis of

Viral Infections in Humans. *Viruses* **12**, 1200 (2020).

19. Yeh, S.-C. *et al.* The anti-immune dengue subgenomic flaviviral RNA is present in

vesicles in mosquito saliva and is associated with increased infectivity. *PLoS Pathog* **19**,

e1011224 (2023).

20. Pijlman, G. P. *et al.* A Highly Structured, Nuclease-Resistant, Noncoding RNA

Produced by Flaviviruses Is Required for Pathogenicity. *Cell Host & Microbe* **4**, 579–591

(2008).

21.   Pompon, J. *et al.* Dengue subgenomic flaviviral RNA disrupts immunity in mosquito

salivary glands to increase virus transmission. *PLoS Pathog* **13**, e1006535 (2017).

22.   Serrato-Pomar, I. *et al.* Multiple orthoflaviviruses secrete sfRNA in mosquito saliva to

promote transmission by inhibiting MDA5-mediated early interferon response. Preprint at

https://doi.org/10.1101/2025.01.21.634113 (2025).

23.   Slonchak, A. & Khromykh, A. A. Subgenomic flaviviral RNAs: What do we know

after the first decade of research. *Antiviral Research* **159**, 13–25 (2018).

24.   Rey-Cadilhac, F. *et al.* Characterization of size distribution and markers for mosquito

extracellular vesicles. *Front. Cell Dev. Biol.* **13**, 1497795 (2025).

25.   MacFadden, A. *et al.* Mechanism and structural diversity of exoribonuclease-resistant

RNA structures in flaviviral RNAs. *Nat Commun* **9**, 119 (2018).

26.   Chapman, E. G. *et al.* The Structural Basis of Pathogenic Subgenomic Flavivirus RNA

(sfRNA) Production. *Science* **344**, 307–310 (2014).

27.   Ward, A. M. *et al.* Quantitative mass spectrometry of DENV-2 RNA-interacting

proteins reveals that the DEAD-box RNA helicase DDX6 binds the DB1 and DB2 3' UTR

structures. *RNA Biology* **8**, 14.

28.   Baek, M. *et al.* Accurate prediction of protein–nucleic acid complexes using

RoseTTAFoldNA. *Nat Methods* **21**, 117–121 (2024).

29.   Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold. *Nature*

596, 583–589 (2021).

30.   De Castro, E. *et al.* ScanProsite: detection of PROSITE signature matches and

ProRule-associated functional and structural residues in proteins. *Nucleic Acids Research* **34**,

W362–W365 (2006).

31.   Mathur, G. *et al.* Transgene☐mediated suppression of dengue viruses in the salivary

glands of the yellow fever mosquito, *Aedes aegypti. Insect Molecular Biology* **19**, 753–763

(2010).

32.   Martin, C., Ligat, G. & Malnou, C. E. The Yin and the Yang of extracellular vesicles

during viral infections. *Biomedical Journal* **47**, 100659 (2024).

33.   Dellar, E. R., Hill, C., Melling, G. E., Carter, D. R. F. & Baena☐Lopez, L. A.

Unpacking extracellular vesicles: RNA cargo loading and function. *J of Extracellular Bio* **1**,

e40 (2022).

34.   Corrado, C., Barreca, M. M., Zichittella, C., Alessandro, R. & Conigliaro, A.

Molecular Mediators of RNA Loading into Extracellular Vesicles. *Cells* **10**, 3355 (2021).

35.   Fabbiano, F. *et al.* RNA packaging into extracellular vesicles: An orchestra of

RNA☐binding proteins? *J of Extracellular Vesicle* **10**, e12043 (2020).

36.   Villarroya-Beltri, C. *et al.* Sumoylated hnRNPA2B1 controls the sorting of miRNAs

into exosomes through binding to specific motifs. *Nat Commun* **4**, 2980 (2013).

37.   Temoche-Diaz, M. M. *et al.* Distinct mechanisms of microRNA sorting into cancer

cell-derived extracellular vesicle subtypes. *eLife* **8**, e47544 (2019).

38.   Iavello, A. *et al.* Role of Alix in miRNA packaging during extracellular vesicle

biogenesis. *International Journal of Molecular Medicine* **37**, 958–966 (2016).

39.   Janas, T., Janas, P., Sapoń, K. & Janas, T. Binding of RNA Aptamers to Membrane

Lipid Rafts: Implications for Exosomal miRNAs Transfer from Cancer to Immune Cells.

IJMS **21**, 8503 (2020).

40.   Hagiwara, K., Katsuda, T., Gailhouste, L., Kosaka, N. & Ochiya, T. Commitment of

Annexin A2 in recruitment of microRNAs into extracellular vesicles. *FEBS Letters* **589**,

4071–4078 (2015).

41.   Nkosi, D. *et al.* Epstein-Barr Virus LMP1 Promotes Syntenin-1- and Hrs-Induced

Extracellular Vesicle Formation for Its Own Secretion To Increase Cell Proliferation and

Migration. *mBio* **11**, e00589-20 (2020).

42.   Cone, A. S. *et al.* Alix and Syntenin-1 direct amyloid precursor protein trafficking into

extracellular vesicles. *BMC Mol and Cell Biol* **21**, 58 (2020).

43.   Li, L.-H. *et al.* A dengue type 2 reporter virus assay amenable to high-throughput

screening. *Antiviral Research* **183**, 104929 (2020).

44.   Li, L.-H. *et al.* Multiplexed multicolor antiviral assay amenable for high-throughput

research. *Nat Commun* **15**, 42 (2024).

45.   Kuno, G. Early History of Laboratory Breeding of *Aedes aegypti* (Diptera: Culicidae)

Focusing on the Origins and Use of Selected Strains. *jnl. med. entom.* **47**, 957–971 (2010).

46.   Rey-Cadilhac, F. *et al.* Characterization of size distribution and markers for mosquito

extracellular vesicles. *Front. Cell Dev. Biol.* **13**, 1497795 (2025).

47.   Liao, K.-C. *et al.* Identification and characterization of host proteins bound to dengue

virus 3′ UTR reveal an antiviral role for quaking proteins. *RNA* **24**, 803–814 (2018).

48.   Perez-Riverol, Y. *et al.* The PRIDE database at 20 years: 2025 update. *Nucleic Acids*

Research **53**, D543–D553 (2025).

49.   Baek, M. *et al.* Accurate prediction of protein–nucleic acid complexes using

RoseTTAFoldNA. *Nat Methods* **21**, 117–121 (2024).

50.   Sehnal, D. *et al.* Mol* Viewer: modern web app for 3D visualization and analysis of

large biomolecular structures. *Nucleic Acids Research* **49**, W431–W437 (2021).

51.   Horn, T. & Boutros, M. E-RNAi: a web application for the multi-species design of

RNAi reagents—2010 update. *Nucleic Acids Research* **38**, W332–W339 (2010).

52.   Blandin, S. A., Marois, E. & Levashina, E. A. Antimalarial Responses in Anopheles

gambiae: From a Complement-like Protein to a Complement-like Pathway. *Cell Host &*

Microbe **3**, 364–374 (2008).

53.   Olmo, R. P. *et al.* Control of dengue virus in the midgut of *Aedes aegypti* by ectopic

expression of the dsRNA-binding protein Loqs2. *Nat Microbiol* **3**, 1385–1393 (2018).

54.   Marois, E. Techniques for Identifying and Sorting Transgenic Mosquito Larvae. *Cold*

Spring Harb Protoc **2024**, pdb.top107694 (2024).

Fig. 1 | DENV infection stimulates secretion of EVs in cells and mosquitoes. a Assessing the effect of DENV infection on EV secretion in mosquito cells. **b-d** Representative WB (b) and quantification of AeSyntenin (c) and hCD63 (d) in media from cells infected with MOI 0.05 or 1. **e-g** Representative WB (e) and quantification of AeSyntenin (f) and hCD63 (g) in cells infected with MOI 0.05 or 1. Actin was used a loading control. **h,i** Particle concentration (h) and size (i) in media from cells infected with MOI 1 by NTA. **j-m** Size-range (20 nm width) particle concentration in media from cells mock-infected (j) and infected with MOI 1 (k), and size-range proportion in media from cells mock-infected (l) and infected with MOI 1 (m) by NTA. **o** Assessing the effect of DENV infection on EV secretion in mosquitoes. **p-r** Representative WB (p) and quantification of AeSyntenin (q) and hCD63 (r) in mock and infected carcasses. Actin was used as a loading control. **s-u** Representative WB (s) and quantification of AeSyntenin (t) and hCD63 (u) in mock and infected salivary glands. Five pairs of salivary glands were combined in each sample. Actin was used as a loading control. **v** EVs per cm² of pictures. n pictures, 40 and 35 for mock and infected saliva, respectively. Particles from 40-60 nm were not counted. c,d,f-m,q,r,t,u,v Bars represent mean ± s.e.m. and dots indicate repeats. j-m N, 3. c,d,f,g,h,i,q,r,t,u,v *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$, as determined by Fisher's LSD test or T-test. j-m *, $p < 0.05$ as determined by T-test in comparison with same size range in the corresponding mock condition.

Fig. 2 | sfRNA stimulates secretion of EVs. a Assessing the effect of sfRNA-deficient

DENV (ΔPk1 DENV) infection on EV secretion in mosquito cells. Control cells were infected

with a wild-type infectious clone. **b** Levels of intracellular sfRNA:gRNA ratio. **c-e**

Representative WB (c) and quantification of AeSyntenin (d) and hCD63 (e) in media from

cells infected with mock, ΔPk1 and WT DENV. **f-h** Representative WB (f) and quantification

of AeSyntenin (g) and hCD63 (h) in cells infected with mock, ΔPk1 and WT DENV. Actin

was used as a loading control.. **i** Assessing the effect of sfRNA transfection on EV secretion

in mosquito cells. A same-size viral RNA fragment was transfected as control (Ctl. RNA). **j-l**

Representative WB (j) and quantification of AeSyntenin (k) and hCD63 (l) in media from

cells transfected with sfRNA or Ctl. **m-o** Representative WB (m) and quantification of

AeSyntenin (n) and hCD63 (o) in cells transfected with sfRNA or Ctl. RNA. Actin was used

as a loading control. **p-q** Particle concentration (p) and size (q) in EVs isolated from cell

media after transfection with sfRNA and Ctl. RNA. N, 3. **r,s** Size-range (20 nm width)

particle proportion in media from cells transfected with Ctl. RNA (r) or sfRNA (s) by NTA.

N, 3. **t,u** Levels of secreted RNase-resistant sfRNA (t) and Ctl. RNA (u) with or without

Triton X-100 pre-treatment. b,d,e,g,h,k,l,n-u Bars show mean ± s.e.m. and dots indicate

repeats. Ld., protein size ladder. g,h *, $p < 0.05$; **, $p <0.01$; ***, $p < 0.001$, as determined by

Fisher's LSD Test. N **, $p < 0.01$ as determined by T test. t,u Different letters indicate

significant differences as determined by Fisher's LSD test.

Fig. 3 | sfRNA interacts with AeSyntenin.

a Identification of sfRNA-interacting proteins by RNA-affinity chromatography. A same-

sized viral control RNA (Ctl RNA) fragment was used as control. **b** Label free quantification

(LFQ) and pseudo ratio for sfRNA-interacting proteins as determined by MS analysis.

Proteins with putative functions associated with EVs, EV biogenesis, RNA binding or Golgi

trafficking are identified with different colors. **c, d** AeSyntenin detection in flow-through and

eluate from RNA-affinity chromatography conducted with EV (c) and cell (d) lysates. **e**

Identification of AeSyntenin-interacting viral RNAs in EVs and cells infected with DENV by

RNA immunoprecipitation. **f-h** Syntenin detection (f) in flow-through and RNA

immunoprecipitates (RIP), sfRNA (g) and gRNA (h) levels precipitated with anti-AeSyntenin

and IgG control in EV lysates. **i-k** AeSyntenin detection (i) in flow-through and RNA

immunoprecipitates (RIP), sfRNA (j) and gRNA (k) levels precipitated with anti-AeSyntenin

and IgG control in cell lysates. g,h,j,k Bars represent mean ± s.e.m. Dots indicate repeats. **,

$p < 0.01$; ***, $p < 0.001$, as determined by T test.

Fig. 4 | AeSyntenin depletion in mosquitoes reduces sfRNA secretion in saliva. a

Assessing the effect of AeSyntenin depletion on gRNA and sfRNA levels in carcasses,

salivary glands (SG) and saliva. Mosquitoes were injected with dsRNA against AeSyntenin or

LacZ as control before inoculation with 5 FFU of DENV. **b,c** AeSyntenin levels in carcass (b)

and in SG (c). Actin was used as a loading control. **d** *AeSyntenin* mRNA levels in SG. **e-g**

Levels of gRNA (e), sfRNA (f) and sfRNA:gRNA ratio (g) in SG. **h-j** Levels of gRNA (h),

sfRNA (i) and sfRNA:gRNA ratio (j) in saliva. **k** Assessing the effect of AeSyntenin

depletion on gRNA and sfRNA levels in carcasses, salivary glands (SG) and saliva after

compensating for AeSyntenin effect on SG infection levels. Mosquitoes injected with

dsAeSyntenin were inoculated with 700 FFU of DENV, whereas mosquitoes injected with

dsLacZ were injected with 5 FFU. **k-o** Levels of gRNA (k), sfRNA (m), sfRNA:gRNA ratio

(n) and infectious particles (o) in SG. **p-s** Levels of gRNA (p), sfRNA (q), sfRNA:gRNA ratio

(r) and infectious particles (s) in saliva. d,g,j,n,o,r,s Bars represent mean ± s.e.m. and dots

indicate repeats. e,f,h,u,l,m,p,q Bars represent geometric mean ± 95% C.I. and dots indicate

repeats. FFU, focus forming unit. *, $p < 0.05$; **, $p < 0.01$; ***, $p < 0.001$, as determined by

T-test.

Fig. 5 | AeSyntenin genetic deletion in salivary glands reduces EVs and sfRNA in saliva.

a Production of SG-specific AeSyntenin deletion mosquitoes. **b,c** AeSyntenin detection in SG

(b) and carcass (c) from control RFP and RFP x guide RNA (guRNA) progeny. Bar graphs

show AeSyntenin fold change normalized to Actin. **d-f** Representative TEM pictures of saliva

from RFP (d) and RFP x guRNA progeny (e), and EVs per cm² of pictures (f). Scale bar, 100

nm. Red arrows indicate EVs. EV concentration was calculated from 34 and 35 pictures for

RFP and RFP x guRNA, respectively. **g,h** Distribution of EVs per size range in RFP (g) and

RFP x guRNA progeny (h). N EVs for g, 120; and for h, 44. **i-k** Levels of gRNA (i), sfRNA

(j) and sfRNA:gRNA ratio (k) in SG from infected RFP and RFP x guRNA progeny. **l-n**

Levels of gRNA (l), sfRNA (m) and sfRNA:gRNA ratio (n) in saliva from infected RFP and

RFP x guRNA progeny. **o,p** Levels of sfRNA:gRNA ratio in saliva from infected EGFP and

EGFP x guRNA progeny (o) and YFP and YFP x guRNA progeny (p). f,k,n,o,p Bars

represent mean ± s.e.m. and dots indicate repeats. g,h Bars show proportion ± SD. i,j,l,m Bars

represent geometric mean ± 95% C.I. and dots indicate repeats. f,m-p *, $p < 0.05$; *, $p < 0.01$;

***, $p < 0.001$ as determined by T test. h *, $p < 0.05$ as determined by Z test with same size

range in RFP.

862