Contrasting ecological communities in rice paddy fields under conventional and no-fertilizer farming practices

Masayuki Ushio[1,*], Hiroki Saito[2], Yoshiko Shimono[3], Kazunori Okada[4], Rika Ozawa[5], Kaori Shiojiri[6], Junji Takabayashi[5]

[1]Department of Ocean Science, The Hong Kong University of Science and Technology, Clear Water Bay, Kowloon, Hong Kong SAR; [2]Tropical Agriculture Research Front, Japan International Research Center for Agricultural Sciences, 1091-1 Maezato-Kawarabaru, Ishigaki, Okinawa 907-0002, Japan; [3]Graduate School of Agriculture, Kyoto University, Oiwake-cho, Kitashirakawa, Sakyo-ku, Kyoto 606-8502, Japan; [4]Agro-Biotechnology Research Center, the University of Tokyo, 113-8657, Tokyo, Japan; [5]Center for Ecological Research, Kyoto University, 2-509-3 Hirano, Otsu, Shiga 520-2113, Japan; [6]Faculty of Agriculture, Ryukoku University, 1-5 Yokotani, Seta Oe-cho, Otsu, Shiga 520-2194, Japan, *Corresponding author, ong8181@gmail.com

SUMMARY: Global crop production is essential for providing energy and nutrients to humans, with rice (*Oryza sativa*) being one of the most important crops globally. However, agricultural systems contribute substantially to greenhouse gas emissions and other environmental challenges, making it crucial to find sustainable methods for crop production that minimize environmental impacts. Paddy fields serve as habitats for numerous species and ecological communities, which play a critical role in rice performance and can be leveraged to promote sustainable crop production. In this study, we conducted extensive field monitoring of two rice paddy fields in Kyoto Prefecture, Japan, where a rice variety, *Hinohikari*, was cultivated, to investigate how ecological communities influence rice growth and yields under two contrasting farming practices: conventional farming and no-fertilizer farming. The no-fertilizer paddy field has been maintained without fertilizers and pesticides since 1951, yet it still maintains sufficient grain yields. We have three objectives in this study: (1) to monitor rice growth and ecological dynamics to create a comprehensive ecological time series using quantitative eDNA metabarcoding and complementary monitoring methods, (2) to identify ecological variables that causally affect rice performance and quantify these effects using nonlinear time series analysis, and (3) to examine the effects of the two farming practices on ecological variables and rice performance in real paddy fields. As expected, the no-fertilizer paddy field showed lower rice growth and yields but still produced 40–50% of the yields of the conventional field. Ecological monitoring revealed contrasting ecological communities between the two fields, particularly among plant species on the paddy ridges and microbial taxa. Our causal detection analysis indicated that 25 taxa had statistically significant causal influences on rice performance indicators such as growth, SPAD readings, or the number of rice stems. While the per-abundance influences of the causal taxa were largely similar in both paddy fields, their abundances were different between the two paddy fields, contributing to the differences in the overall effects of these taxa on the rice performance indicators. These findings suggest that the abundance of causal taxa, along with nutrient conditions, may drive differences in rice growth between the two paddy fields.

Keywords: Agricultural system; Ecological monitoring; Ecological variables; Environmental DNA; Rice; Time-series analysis; Unified information-theoretic causality

1. INTRODUCTION

Global production of crops provides energy and nutrients for humans (Godfray et al., 2010), and rice (*Oryza sativa*) is one of the most important crops, along with maize, wheat, sugar cane, and oil palm, comprising 8% of world crop production (FAO, 2024). As with many other crops, improving yields and production efficiency such as resource use efficiency and pathogen resistance is one of the major concerns in rice production (e.g., Zhang, 2007), which is essential to maintain the sustainability of human society. However, agriculture systems are significant contributors to greenhouse gas emissions and other environmental issues such as nutrient runoff as nonpoint source pollution (Hua et al., 2017; Menegat et al., 2022; Wang et al., 2014). Greenhouse gas emissions (CH_4 and N_2O) from paddy fields can affect the global climate (Qian et al., 2023) while nutrient runoff impacts surrounding ecosystems (Ko et al., 2021), both of which have deteriorating effects on the sustainability of the agriculture systems. Thus, finding ways to achieve sustainable rice crop production while minimizing environmental impacts is a key challenge in agricultural science.

Rice cultivation occurs in paddy fields, and rice growth, yields, and gene expression patterns can be influenced by various biotic and abiotic factors. Previous studies have examined how meteorological and endogenous variables (e.g., plant age and genotype) affect gene expression patterns (transcriptome dynamics) of rice in fluctuating field environments (Kashima et al., 2021; Nagano et al., 2012). These studies suggest that transcriptome dynamics are primarily driven by plant age, genotypes, and meteorological variables such as air temperature and solar radiation. Under these circumstances, genomics-based breeding, alongside advanced technologies such as high-throughput field-based phenotyping, is one of the most promising strategies to accommodate the complex interactions between rice and abiotic variables for improving rice yields and production efficiency (Wing et al., 2018).

Paddy fields are a habitat for a large number of species (e.g., more than 5000 species have been recorded in paddy fields in Japan; Natuhara, 2013), and ecological communities such as insect pests, aquatic animals, and microbial mutualists/pathogens significantly influence the rice performance (Ji et al., 2023; Li et al., 2025; Savary et al., 2019). For example, recurring outbreaks of an insect pest, brown planthoppers, can cause serious physical and physiological damages to rice (Bottrell & Schoenly, 2012). Co-culturing rice with aquatic animals such as fish, crabs, crayfish, and turtles can improve the rice yields, reduce insect pests, and increase soil carbon and nitrogen contents (Ji et al., 2023). Also, in paddy fields, fish can modify habitat structures through their foraging and swimming behaviour and excrement, and Li et al. (2025) showed that this effect could change the abundance of other key species such as earthworms and even paddy ecosystem functions, which could influence rice yields. Furthermore, not only organisms but also organic compounds such as green leaf volatiles (GLVs) can influence plant performance such as pathogen resistance (Matsui & Koeduka, 2016). Common GLVs, (*E*)-2-hexenal and (*Z*)-3-hexenal, induce defenses against pathogenic fungi (Matsui & Koeduka, 2016). Also, plant volatiles emitted by weeds on paddy ridges can influence rice physiological states and yields (Shiojiri et al., 2021). Thus, effects of ecological communities in a paddy field on rice should not be underestimated.

However, the dynamics of an ecological community are often challenging to monitor

because of the large number of species (Natuhara, 2013) and the complex, state-dependent, and nonlinear nature of the dynamics (e.g., Hsieh et al., 2005). Furthermore, understanding the influence of ecological communities on rice performance, or interspecific interactions between rice and ecological communities, is even more difficult due to possible confounding factors under field conditions, despite the potential of such understanding for achieving sustainable agriculture (Ji et al., 2023; Toju et al., 2018). As a result, effects of a relatively small number of species and/or variables on rice have been studied at one time in paddy fields (Ji et al., 2023; Li et al., 2025) and how a large number of ecological variables simultaneously affect the rice performance in paddy fields has remained unclear.

Environmental DNA (eDNA) analysis is one of the promising methods for overcoming the above-noted issues about comprehensive biodiversity monitoring under field conditions. eDNA refers to DNA directly extracted from environmental samples such as water and soil (Taberlet et al., 2018), which includes both microbial and macrobial DNA. Previous studies have shown that eDNA metabarcoding, an approach to comprehensively amplify and sequence marker regions of DNA belonging to a target taxa in an environmental sample, is a cost- and time-effective means to detect a large number of species (e.g., Deiner et al., 2016; Miya et al., 2015), and that the eDNA-based community monitoring is especially informative when it is performed quantitatively (Ushio, Murakami, et al., 2018; e.g., sequencing with internal spike-in DNAs: Ushio, 2022). It enables tracking the detailed dynamics of a speciose ecological community (Ushio, 2022), and it can be even more effective when used with other traditional methods such as direct capture and visual census, as they can provide complementary information about biodiversity (He et al., 2023; Krol et al., 2019).

To investigate whether and how ecological variables influence the rice performance under field conditions, we previously conducted eDNA-based, intensive monitoring of ecological community dynamics in five artificial rice plots (Ushio, 2022; Ushio et al., 2023). We collected water samples from artificial rice plots daily for 122 days and analyzed microbes and macrobes using a quantitative eDNA metabarcoding approach. To overcome another issue, that is, effects of confounding factors under fields condition (i.e., pseudo-correction or "mirage" correlation issue), we applied a time series-based causality detection method (Chang et al., 2021; Osada et al., 2023; Sugihara et al., 2012) to the time series, and identified species that affected the rice growth (i.e., an Oomycetes species, *Globisporangium nunn*, and a midge species, *Chironomus kiiensis*). Further, the effects of these two species on the rice growth and gene expression were validated using manipulative experiments. The findings of that study demonstrated that ecological community members indeed have effects on the rice growth performance and gene expression. However, the system used in Ushio et al. (2023) was artificial: small rice plots (about 1 m × 1m plot) and the farming practice was different from that used for real paddy fields. Also, only eDNA metabarcoding was used to monitor the ecological communities, which could have missed potentially important ecological variables such as flying insects, plants on paddy ridges, and plant specialized metabolites.

In the present study, we conducted field monitoring of two rice paddy fields in Kyoto Prefecture, Japan, where rice variety *Hinohikari* was planted, to understand whether and how ecological communities affect the rice growth and yields in real paddy fields. To investigate a

wider range of environmental gradients, we targeted paddy fields with different farming
practices and different backgrounds. The first one was a paddy field with conventional
farming practice; namely, farmers applied basal nitrogen (N), phosphorus (P), and potassium
(K) fertilizers and used pesticides to reduce plants on the paddy ridge. The second one was a
paddy field with no fertilizer treatment. The farming practice was unique, as it has not
employed any basal fertilizers or pesticides since 1951 but still maintain rice yields equal to
about 70–80% of the yields obtained with the conventional management practice (e.g.,
Okumura, 2002). How the "no-fertilizer" paddy field maintains its yield without nutrient
inputs is unknown, but investigating the effects and dynamics of ecological communities may
shed light on the mechanism of the maintenance of the yields. Given this background, the
objectives of this study were: (1) to monitor rice growth and ecological dynamics to generate
extensive time series using quantitative eDNA metabarcoding and other complementary
monitoring approaches, (2) to identify ecological variables that causally influence rice
performance and to quantify the causal effects using time series-based causality detection
methods, and (3) to clarify the effects of the two contrasting management practices on the
ecological variables and the rice performance, in the real paddy fields.
148
149

2. METHODS

2.1 Study site

151
Our monitoring was performed at paddy fields in Uji, Kyoto, Japan (34°54′09″N,
135°46′29″E) from May to September in 2017 (Fig. 1a). Climate data during the field
monitoring was retrieved from the Japan Meteorological Agency website
(https://www.data.jma.go.jp/risk/obsdl/index.php, accessed on 1 November 2024; "daily
mean values", "Kyoto city", and "2017" were selected in the webpage) (Fig. 1b–d), and the
daily mean air temperatures were from 16.9°C to 31.2°C. The field monitoring was
performed in two rice fields with different management practices: rice farming with fertilizer
and pesticides ("conventional" farming; Fig. 1e) and that without fertilizer and pesticides
("no-fertilizer" farming; Fig. 1f). Hereafter, we refer to these fields as "conventional" and "no
fertilizer" paddy fields, respectively. The conventional paddy field occupies about 580 m^2 and
the no-fertilizer paddy field occupies about 945 m^2, and they are separated by about 15 m
(Fig. S1a). In the conventional field, basal N-P-K fertilizer at 24–30 kg N ha^{-1} year^{-1} was
added in May, and another 18 kg N ha^{-1} year^{-1} was subsequently added in mid-August. Three
to four days after the rice planting, pesticide (ST Longet Floable, Sumitomo Chemical
Company, Ltd., Tokyo, Japan) at 6,700 ml ha^{-1} was applied in the conventional field. The no-
fertilizer paddy field has been maintained by a non-profit organization "Research Group on
No Fertilizer and No Pesticide Farming" (https://muhiken.or.jp/). This paddy soil was
initially located in Ritto, Shiga Prefecture (next to Kyoto Prefecture; about 25 km from the
location of the no-fertilizer paddy field), Japan, where farming without fertilizers or
pesticides commenced in 1951 (https://muhiken.or.jp/activities/history/, accessed as of 15
July 2025). In 2006, after farming in Ritto ceased, the no-fertilizer paddy soil was moved to
Uji, Kyoto, where no-fertilizer rice farming has continued. Even without fertilizers, the no-

fertilizer field maintained production of about 80% of the typical grain yield when rice
variety *Beniasahi* was cultivated (Okumura, 2002). Furthermore, previous research has
suggested differences in microbial community composition and nutrient uptake efficiency
between the conventional and no-fertilizer fields (Kamata et al., 1991; Okumura, 2002).

Figure 1 | Study site, climate variables, and rice paddy fields with "conventional" and "no-fertilizer" farming practices
(**a**) Location of the study site. The yellow marker indicates the site, while the red rectangle in the inset highlights the study region (Maps created using the ggmap package of R; ©Stadia Maps, ©Stamen Design, ©OpenMapTiles and ©OpenStreetMap). (**b**) Mean daily air temperature (°C). (**c**) Daily rainfall (mm). (**d**) Daily sunlight hours. Climate data were obtained from the weather station in Kyoto City, managed by the Japan Meteorological Agency. (**e**) "Conventional" paddy field. (**f**) "No-fertilizer" paddy field. The yellow panel in the right side of the image was used for collecting insect samples. Field photographs were taken on 12 July 2017 (©M. Ushio).

2.2 Sampling design

179
In each paddy field, we conducted weekly monitoring of rice growth and ecological
community dynamics from 19 May 2017 to 30 August 2017. The first monitoring was
conducted on 19 May 2017 (Friday) before planting rice, var. *Hinohikari*, and the other
monitoring was conducted in the morning (between 8:30 and 10:40) every Wednesday (see
the sample metadata in Table S1). Although var. *Beniasahi* had previously been cultivated in
the no-fertilizer paddy field, var. *Hinohikari* was selected for the present study to align with
the variety grown by farmers in the conventional paddy field and to ensure consistency across
the field types for evaluating ecological effects on rice growth. In total, we conducted the
field survey 15 times, including the first one before the rice planting. The monitoring was
ceased on 30 August because the rice field was drained in preparation for harvesting rice.

We set five rice monitoring points in each paddy field and five sampling points on each
paddy ridge (Fig. S1a). In each paddy field, five 3 m × 3 m plots, which were 2 m apart from
the paddy ridge, were set to monitor the rice performance, including rice heights, SPAD
values, and the number of rice stems (see the section "*2.3 Rice growth monitoring and yield*
survey"). In each paddy ridge, we set sticky panels for collecting flying insects (Fig. S1b) and
50 cm × 50 cm quadrats (0.25 m^2) for monitoring wild weeds (Fig. S1c) in each of five
sampling points. At the same monitoring points in paddy fields, we collected rice leaves for
analyzing plant specialized metabolites. Water samples for eDNA-based ecological
community monitoring were collected from the rice monitoring points for quantitative
monitoring of microbes and macrobes (see section "*2.4 Monitoring of ecological variables*
and communities"). The rice was harvested on 29 September 2017, and the rice yield survey
was conducted (Fig. S1d, e).
202

2.3 Rice growth monitoring and yield survey

Rice growth was monitored by measuring rice leaf height of target individuals at each
monitoring point every week using a ruler (the highest leaf heights were measured). The
average height of the three individuals was used as a representative rice height of each
monitoring point, and thus we had five representative values for each plot for each week.
Leaf SPAD was also measured every week for the same leaf whose height was measured
using a SPAD meter (SPAD-502Plus, KONICA-MINOLTA, Inc., Tokyo, Japan). The number
of rice stems was also counted. After we harvested rice, we counted the number of fertile and
sterile grains and measured their wet and dry weights as indices of rice yields.
212

2.4 Monitoring of ecological variables and communities

2.4.1 Plant specialized metabolites
Rice leaves were collected for analyzing plant specialized metabolites and kept at –20°C
during the transport to the laboratory. The volatile collection system using Gerstel-Twister
(polydimethylsiloxane [PDMS] coated stir bar, film thickness 0.5 mm, 10 mm length, Gerstel
GmbH & Co. KG) was used to collect volatile compounds emitted from the frozen rice
leaves. Headspace volatiles emitted from four leaf pieces, each 4 cm long, were collected on
two Twisters in a closed glass vial (30 ml) in a laboratory room (25 ± 1°C) for 3 hours.
Before starting the volatile collection, 0.1 µg of tridecane on a piece of filter paper was added
to the glass bottle as an internal standard. The collected volatile compounds were analyzed by
GC-MS with an HP-5MS capillary column as described previously (Rim et al., 2018). The
peak area of each compound was normalized by dividing it by the peak area of the internal
standard and the leaf weight (g). The normalized peak area was calculated as "relative peak
area."
Momilactone was quantified using rice leaf samples. Each leaf sample (10–400 mg
FW) was immersed in 1 ml of 70% methanol and incubated at 4°C for 24 hours. A 2-µl
aliquot of the extract was analyzed by LC–MS/MS (Sciex QTRAP 3200) with the following
selected reaction monitoring (SRM) transitions (Riemann et al., 2013): momilactone A, m/z
315→271; momilactone B, m/z 331→269; phytocassanes A and E, m/z 317→299;
phytocassane B, m/z 335→317; and phytocassane C, m/z 319→301. The total phytoalexin

level was calculated as the sum of all compounds, reflecting the diterpenoid content. We
detected momilactone A, momilactone B, phytocassane B, and phytocassane B3, and their
concentrations were aggregated as "Diterpenoids (μg/g)" for statistical analysis.
236
2.4.2 Insect collections
Yellow sticky traps (Horiver; 257 mm × 100 mm; Arysta LifeScience Corporation, Tokyo,
Japan) were installed at the five sampling locations on the paddy ridges during the
monitoring period. One Horiver trap was mounted on a plastic stick at each sampling point.
The traps were replaced with new ones every week. The collected traps were examined under
a stereomicroscope, and the numbers of thrips (Thysanoptera), leafhoppers (Cicadellidae),
and other planthoppers (mainly Homoptera) were recorded.
244
2.4.3 Plant cover survey on the ridge
Each 50 cm × 50 cm quadrat was subdivided into 25 cells of 10 cm × 10 cm by strings
stretched across the frame (Fig. S1c). Photographs of the entire quadrat were taken from
directly above the quadrat every week from June to August and the percentage cover of each
major species was calculated as visual estimates from the photographs. On June 28 and July
26, direct vegetation surveys were also conducted on-site and confirmed that there were no
discrepancies with the judgments made based on the photographs taken.
252
2.4.4 Water sampling for eDNA analysis
Water samples in each paddy field were collected for eDNA-based ecological community
analysis. The water was filtered using Sterivex filter cartridges (Merck Millipore, Darmstadt,
Germany). Due to the high density of sediment in the rice paddy fields, the volume of water
we could filter was relatively small (30–100 ml for each sampling point). Two types of filter
cartridges were used to filter water samples: to detect microorganisms, φ0.22-μm Sterivex
(SVGV010RS) filter cartridges that included zirconia beads inside were used to filter 30 ml
of water (Ushio, 2019); and to detect macroorganisms, φ0.45-μm Sterivex (SVHV010RS)
filter cartridges were used to filter 30–100 ml of water. After filtration, 2 ml of RNAlater
solution (ThermoFisher Scientific, Waltham, Massachusetts, USA) were added to each filter
cartridge to prevent DNA degradation during storage. The same volume of MilliQ water was
filtered using the two types of filter cartridges as field negative controls. The filter cartridges
were stored in a cooler box until they were taken back to the laboratory. They were stored at
–20°C until further processing.
Although we conducted weekly water sampling from 19 May 2017 to 30 August 2017,
water sampling was impossible for some sampling events because mid-season drainage was
practiced in the rice fields to improve rice performance in August. In total, we collected 266
filter cartridges (118 field water samples × 2 filter cartridges + 15 field-level negative
controls × 2 filter cartridges) from the two rice fields. Details about the water samples are
available in Table S1.
273

2.5 Quantitative environmental DNA metabarcoding
2.5.1 DNA extraction and library preparation for quantitative eDNA metabarcoding

Detailed protocols of the quantitative eDNA metabarcoding are described in the
supplementary methods and Ushio (2022). Briefly, DNA was extracted and purified using a
DNeasy Blood & Tissue kit (Qiagen, Hilden, Germany) (Ushio, 2019). After the purification,
DNA was eluted using 100 µl of the elution buffer and stored at −20°C until further
processing.
A two-step PCR approach was adopted for the library preparation. The first-round PCR
(first PCR) was carried out with the internal standard DNAs to amplify metabarcoding
regions using primers specific to prokaryotes (515F and 806R), eukaryotes (Euk_1391f and
EukBr), fungi (ITS1-F-KYO1 and ITS2-KYO2) and animals (mlCOIintF and HCO2198).
Sequences for primers and internal standard DNAs and primer references are available in the
Supplementary methods. To monitor cross-contamination during the library preparation, 16
PCR-level negative controls with standard DNAs and 4 PCR-level negative controls without
standard DNAs were included. In total, 153 DNA samples were analyzed for each type of
filter cartridge (118 field water samples + 15 field-level negative controls + 20 PCR-level
negative controls). The second-round PCR (second PCR) was carried out to append indices
for different samples for sequencing. The DNA library was sequenced on the MiSeq platform
(Illumina, San Diego, CA, USA) using the 250PE reagent kit for 16S, COI, and ITS, and the
150PE reagent kit for 18S.
294
2.5.2 Sequence data processing
The raw sequence data were converted into FASTQ files using the bcl2fastq program
provided by Illumina (bcl2fastq v2.18). The FASTQ files were then demultiplexed using the
command implemented in Claident (http://www.claident.org) (Tanabe & Toju, 2013).
Demultiplexed FASTQ files were then analyzed using the Amplicon Sequence Variant (ASV)
method implemented in the "dada2" package (Callahan et al., 2016) of R (R Core Team,
2024). ASVs detected in the three experiments were merged and clustered into OTUs at 97%
similarity using the DECIPHER package of R (Wright, 2016), which converted the ASV-
sample matrix into the OTU-sample matrix. We converted ASVs to OTUs because ASVs
often detected intraspecific variations and the focus of this experiment was fish community
compositions at the species level (not at the intra-species level). Taxonomic identification
was performed using Claident v0.9.2024.03.04 (Tanabe & Toju, 2013). Note that Claident
employs conservative species assignment criteria (i.e., it does not assign species name if the
reliability is low), which we intentionally utilized to minimize taxonomic misidentification.
For all analyses after this subsection, the free statistical environment R was used (R
Core Team, 2024). The procedure used to estimate DNA copy numbers consisted of two
parts, following previous studies (Ushio, Murakami, et al., 2018; Ushio, 2019). Briefly, we
performed (i) linear regression analysis to examine the relationship between sequence reads
and the copy numbers of the internal standard DNAs for each sample, and (ii) conversion of
sequence reads of non-standard DNAs to estimate the copy numbers using the result of the
linear regression for each sample. The regression equation was: Sequence reads = sample-
specific regression slope × the number of standard DNA copies [/µl]. Then, the estimated
copy numbers per µl extracted DNA (copies/µl) were converted to DNA copy numbers per
ml water in the rice plot (copies/ml water). After the conversion to DNA copy numbers, rare

OTUs (their concentrations were lower than half of the minimum concentration of standard
DNA; 1,250 copies/ml for 16S, 12.5 copies/ml for 18S, 1.0625 copies/ml for COI, and 10
copies/ml for ITS) and infrequently detected OTUs (detected in less than 10% of total
monitoring events) were removed to mitigate the effects of sequencing errors.
323
2.6 Statistical analysis and visualization of ecological community dynamics
The monitoring data on rice performance and ecological communities were initially
visualized through time-series plots created with the ggplot2 package in R (Wickham, 2009).
Subsequently, the effects of time (sampling date) and farming practices were analyzed using
generalized additive mixed models (GAMM) for rice performance, plant specialized
metabolites, insect abundance, plant cover, and eDNA abundance and diversity, incorporating
random effects for rice individuals (when applicable) and sampling locations. GAMM
analysis was performed using the "mgcv" package in R (Wood, 2004). The influence of
farming practices on the ecological variables affecting rice performance (see the next section)
was assessed using linear mixed models (LMM) with random effects for sampling locations,
utilizing the "nlme" package in R (Pinheiro et al., 2009). Additionally, the effect of
management practices on rice yields was evaluated using analysis of variance (ANOVA).
Non-metric Multi-dimensional Scaling (NMDS) was performed to visualize the
differences in plant cover on the ridge, plant specialized metabolites, and ecological
communities between the two paddy fields. The `adonis2()` function was utilized to assess
whether time (i.e., the number of weeks) and farming practices had statistically significant
effects. In the eDNA community composition data, taxa that differentiated the two paddy
fields were identified using the `envfit()` function. The "vegan" package in R (Oksanen et
al., 2008) was employed for both NMDS and `adonis2()` analyses. Throughout the statistical
analysis, we use the phrase "statistically clear" instead of "significant" in this study to clarify
the meaning of the word "significant" by following Dushoff et al. (2019).
345
2.7 Time series analysis-based causality detection and quantification of
influences from ecological variables on rice growth
To identify factors influencing the rice performance, we quantified information flow between
the rice performance and ecological variables by the unified information-theoretic causality
(UIC) method (Osada et al., 2023) implemented in the "rUIC" v0.9.13 package (Osada &
Ushio, 2021) of R. UIC tests the statistical clarity of information flow between variables in
terms of TE (Schreiber, 2000) computed by nearest neighbor regression based on time-delay
embedding of explanatory variables (i.e., cross mapping; Sugihara et al. 2012). In contrast to
the standard method used to measure TE, UIC quantifies information flow as follows:

$$TE = \frac{1}{N}\sum_{t=1}^{N} \log\left(\frac{p\left(y_{t+tp} \mid x_t, x_{t-\tau}, \cdots, x_{t-(E-1)\tau}, z_t\right)}{p\left(y_{t+tp} \mid x_{t-\tau}, x_{t-2\tau}, \cdots, x_{t-(E-1)\tau}, z_t\right)}\right), \qquad \dots [1]$$

where x, y, and z represent an effect variable, a potential causal variable, and a conditional
variable (if available), respectively. $p(A|B)$ represents conditional probability: the
probability of A conditioned on B. t, tp, τ, and E represent the time index, time step, a unit of
time lag, and the optimal embedding dimension, respectively. N is the total number of points

in the reconstructed state space (this is equivalent to the total number of time points – the
optimal embedding dimension + 1). For example, if $tp = -1$ in Eqn. [1], UIC tests the causal
effect from y_{t-1} to x_t. Optimal E was selected using a TE version of simplex projection
(Sugihara & May, 1990) implemented in the rUIC package. In the present study, the causality
time-lag (tp) up to -2 (equivalent to two-week time-lag) was tested. Statistical clarity was
tested by bootstrapping data after embedding (the statistical clarity levels were set to 0.05).
Because most of our time-series data exhibited a clear trend (e.g., rice height showed
an increasing trend until August), we took the first-difference of the original time-series to
make the time series stationary for the UIC analysis. To determine the variables that causally
influence rice performance, we quantified TE between the indices of rice performance (i.e.,
growth rate, SPAD, and the number of stems) and climate and ecological variables. Given the
large number of statistical tests required (the total number of UIC tests = [rice performance
indices] × tp [0, –1, or –2] × [climate and ecological variables]), we calculated the false
discovery rate (FDR) (Benjamini & Hochberg, 1995) for each category (climate, plant cover,
plant specialized metabolites, and ecological communities). We determined that a variable
has a statistically clear causal effect on rice performance if the FDR is below 0.05. We first
performed the UIC analysis between the climate variables and the rice performance. Then,
we included climate variables that have statistically clear effects on the rice performance in
the embedding in the UIC analysis to reduce the possibility of misidentification of causality
from ecological variables on the rice performance due to the strong effect of climate
variables.
381
2.8 Nonlinear time series analysis to quantify interaction strengths
We quantified the effects of ecological variables on the rice performance using an improved
version of the sequential locally weighted global linear map (S-map) (Sugihara, 1994), called
the multiview distance regularized S-map (MDR S-map) (Chang et al., 2021). Consider a
system that has E different interacting variables and assume that the state space at time t is
given by $\boldsymbol{X}_t = \{x_{1,t}, x_{2,t}, \ldots, x_{E,t}\}$. For each target time point t^*, the S-map method produces
a local linear model that predicts the future value x_{1,t^*+p} from the multivariate reconstructed
state space vector $\boldsymbol{X}_{t^*} = \{x_{1,t^*}, x_{2,t^*}, \ldots, x_{E,t^*}\}$. That is,

$$\hat{x}_{1,t^*+p} = C_0 + \sum_{j=1}^{E} C_j \, x_{j,t^*} \qquad \ldots [2]$$

where $\hat{x}_{1,t^*+p}$ is a predicted value of x_1 at time $t^* + p$, C_j is a regression coefficient and
interpreted as interaction strength (or called S-map coefficient), and C_0 is an intercept of the
linear model. Note that, in real analysis, some of x_{j,t^*} may be time-delayed coordinates of an
interacting variable. The linear model is fit to the other vectors in the state space, but points
that are close to the target point, $\boldsymbol{X}_{t^*}$, are given greater weighting (i.e., locally weighted
linear regression). In the standard S-map, the distances between the target point, $\boldsymbol{X}_{t^*}$, and
other points are measured by Euclidean distance. However, Euclidean distance cannot be a
good measure if the dimension of the state space is high. In the MDR S-map, the distance
between the target point and other points is measured by the multiview distance (Chang et al.,
2021), which can be calculated by ensembling distances measured in various low-

dimensional state spaces (multiview embedding; see Ye & Sugihara, 2016). In addition, to
reduce the possibility of overestimation and to improve forecasting skill, regularization (i.e.,
ridge regression) is also applied (Cenci et al., 2019). As in the UIC analysis, we included
climate variables as conditional variables if they had statistically clear influence on the rice
performance. In the present study, we implemented the MDR S-map in our custom R
package, "macam" v0.0.10 (Ushio, 2022) and used it for computation.
407
408

3. RESULTS

3.1 Rice growth trajectory and yield survey
During the monitoring period in 2017, the daily mean air temperatures were about 20°C in
May, peaked at 30°C in August, and decreased to 20–25°C in September (Fig. 1b). In 2017,
increased rainfall was observed due to seasonal frontal activity in June and typhoons in July,
August, and September, which was negatively correlated with daily sunlight hours (Fig. 1c,
d). In general, daily air temperatures during the monitoring period were similar or slightly
higher than those during the normative year, while the rainfall was greater than during the
normative year because of the seasonal frontal activity and typhoon activity.
The rice growth rate, SPAD, and the number of stems were consistently higher in the
conventional paddy field than in the no-fertilizer paddy field, except for the first-differenced
SPAD values (the original values; Fig. 2a–c: the first-differenced values; Fig. 2d–f) ($P <$
0.0001, Table S2). Large increases in the rice height and the number of stems were observed
in late June and early July (Fig. 2a, c, d, f), while a large increase in SPAD value was
observed in June (Fig. 2b, e). The total number of heads per rice individual and the total wet
weight and total dry weight were also higher in the conventional paddy field than in the no-
fertilizer paddy field ($P < 0.0001$; Fig. 2e, g and Table S3). These patterns were well reflected
in the rice grain yields. The numbers of (total or fertile) grain counts, wet weight, and dry
weight were higher in the conventional paddy field than in the no-fertilizer paddy field ($P <$
0.005; Fig. S2 and Table S3). Despite the lack of input of fertilizer, the no-fertilizer paddy
field still produced (compared to the conventional paddy field production taken as 100%):
total number of heads per 12 individuals: 56%, total grain count per head: 76%, and total
grain dry weight per head: 77%, which corresponds to about 43% of total grain dry weight
per individual (= 56% × 77%).
433

Figure 2 | Rice growth patterns and yields
(**a**) Rice height (cm). (**b**) SPAD values. (**c**) The number of rice stems. (**d**) Changes in rice height per week. (**e**) Changes in SPAD values per week. (**f**) Changes in the number of rice stems per week. (**g**) The total number of rice heads per rice individual. (**h**) The total wet or dry weight (g) per 12 rice individuals. Solid lines indicate mean values. Green color and and filled circle indicate data from the paddy fields using the conventional farming methods, and yellow color and filled triangle indicate data from the paddy fields using no fertilizers. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$. "N.S." indicates that the difference between the two farming practices is not statistically clear.

3.2 Plant specialized metabolites, insect abundance, and plant cover on the

ridge

We detected eight dominant plant specialized metabolites: (*Z*)-3-hexenal, (*E*)-2-hexenal, (*Z*)-
3-hexen-1-ol (leaf alcohol), 2-4-heptadienal, linalool, β-caryophyllene, methyl salicylate
(MeSA), and diterpenoids (Fig. 3a). The concentrations of (*Z*)-3-hexenal and (*E*)-2-hexenal
were higher than those of the other metabolites. The levels of (*Z*)-3-hexenal and β-
caryophyllene were greater in the early period (June) compared to the late period, while 2-4-
heptadienal showed higher concentrations in the late period. (*E*)-2-hexenal, (*Z*)-3-hexen-1-ol,
and MeSA were detected throughout the monitoring period, although their temporal patterns
were unclear. Linalool and diterpenoids were observed only sporadically. Overall, the
differences between the conventional and no-fertilizer paddy fields were relatively unclear,
but we identified statistically clear effects of the farming practice on (*Z*)-3-hexenal, (*E*)-2-
hexenal, and methyl salicylate (Fig. 3a; Table S2).
Thrips (Thysanoptera) were the most commonly observed insect taxon during the
monitoring period (Fig. 3b), primarily in the first half of the monitoring (June to July).
Leafhoppers (Cicadellidae) were also frequently observed, though in smaller numbers than
thrips, except in late June. Other planthoppers were observed almost exclusively in early

June. Statistically clear effects of the farming practice were detected only for other
planthoppers ($P < 0.001$; Fig. 3b; Table S2).

Figure 3 | Plant specialized metabolites, insect abundance, and plant cover in the paddy fields
(**a**) Plant specialized metabolites. Each panel indicates a different type of plant specialized metabolites; (*Z*)-
3-hexenal, (*E*)-2-hexenal, (*Z*)-3-hexen-1-ol, 2-4-heptadienal, linalool, *β*-caryophyllene, and methyl salicylate
(MeSA) indicate plant volatile (relative peak area/fresh leaf weight), while diterpenoids (µg/g fresh leaf
weight) indicates phytoalexin. (**b**) Insect abundance trapped by the sticky plate method set on the ridge
between paddy fields. Each panel shows a different insect group. (**c**) Plant coverage (%) on the ridge area
between the paddy fields. Each panel shows a different plant type. Solid lines indicate mean values. Green
color and and filled circle indicate data from the conventional paddy field, and yellow color and filled tri-
angle indicate data from the no-fertilizer paddy field. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.
"N.S." indicates that the difference between the two farming practices is not statistically clear.

A total of 14 plant species from six families were recorded on the paddy ridge (Figs. 3c
and S3). Species from the Cyperaceae family were more dominant in the no-fertilizer paddy
field ($P < 0.01$; Fig. 3c; Table S2). No statistically clear difference was found in Poacea
between the two paddy fields. In contrast, other plant species were more dominant in the
conventional paddy field, particularly in the later part of the monitoring period ($P < 0.05$; Fig.
3c; Table S2). In the conventional paddy field, an annual graminoid, *Digitaria ciliaris*, was
the most dominant species in June. From July to August, an annual graminoid, *Dinebra*
chinensis, and two annual forbs, *Euphorbia maculata* and *Portulaca oleracea*, became
dominant. In the no-fertilizer rice field, a perennial forb, *Ixeris japonica*, dominated from
June to early July, followed by two perennial sedges, *Fimbristylis diphylloides* and *Cyperus*
brevifolius, which dominated in late July to August.

The overall patterns of plant cover and specialized metabolites were analyzed using
NMDS (Fig. 4). Insect community compositions were not compared, as only three insect
groups were examined in this study. No statistically clear differences were found in the
composition of specialized metabolites between the two paddy fields (number of weeks, *P* =
0.0147; farming practice, *P* > 0.05; interaction, *P* > 0.05). However, a statistically clear
difference was observed in plant cover composition between the conventional and no-
fertilizer paddy fields (number of weeks, *P* < 0.0001; farming practice, *P* < 0.0001;
interaction, *P* < 0.0001), which aligns with the patterns observed in the individual metabolites
and plant species presented in Figs. 3 and S3.

Figure 4 | Non-metric dimensional scaling (NMDS) of plant communities on the paddy ridges and plant specialized metabolites
(**a**) NMDS of the plant communities on the paddy ridges based on Bray-Curtis dissimilarity (stress value = 0.1899). (**b**) NMDS of the plant specialized metabolites based on Bray-Curtis dissimilarity (stress value = 0.1568). The density of each color indicates the number of weeks since the rice planting in the rice fields. Green and and filled circle indicate data from the conventional paddy field, and yellow and filled triangle indicate data from the no-fertilizer paddy field.

3.3 Ecological community dynamics detected by quantitative eDNA metabarcoding

Quantitative eDNA metabarcoding was conducted using water samples from the paddy field.
A total of 21,638,562 high-quality sequence reads (>Q30 = 94.17%) were obtained from four
MiSeq runs (7,274,858 reads with >Q30 = 89.19% for 16S; 7,352,339 reads with >Q30 =
98.09% for 18S; 7,749,778 reads with >Q30 = 97.13% for COI; and 2,899,016 reads with
>Q30 = 87.25% for ITS). Generally, we detected negligible non-target DNA sequence reads
from negative controls (Fig. S4). While some negative controls detected non-target DNA

sequences, these were mostly specific to the negative controls and did not affect subsequent
analyses. Thus, we concluded that no serious contamination occurred during our sampling
and library preparation processes. After DADA2 processing, 20,067,623 reads remained
(Table S4). Following the conversion to copy numbers per ml of water and the filtering of
very rare and infrequently detected OTUs, we identified 1,213 OTUs (Table S5). Table S6
provides a summary of eDNA copy numbers, sample metadata, and taxonomic information.
Regarding the temporal trend of total eDNA concentration, we observed contrasting
patterns between the two paddy fields; total eDNA concentration decreased in the
conventional paddy field while it increased in the no-fertilizer rice field during the
monitoring period (Fig. 5a). A statistically clear effect of the farming practice on total eDNA
concentration was found ($P < 0.0001$; Fig. 5a).
Bacteria were the most dominant taxa detected (Figs. 5a), with dominant genera
including *Paenibacillus* sp., *Flavobacterium* sp., *Pedobacter* sp., and several species
belonging to Actinomycetes and Comamonadaceae (Figs. S5, S6; Table S7). Most major
bacterial and fungal phyla showed higher abundance in the conventional paddy field
compared to the no-fertilizer paddy field (Fig. S6). However, Actinomycetota and Bacillota
were more abundant in the no-fertilizer paddy field after July–August. Additionally,
Fusobacteriota showed increased abundance during June–July in the no-fertilizer paddy field.
The dominant non-bacterial taxa included green algae (Desmidiales), freshwater algae
(*Gomphonema parvulum*), freshwater ostracods (Cyprididae), and small crustaceans (*Monia*
sp.) (Table S7), with generally higher abundances in the conventional paddy field than in the
no-fertilizer paddy field. Photosynthetic taxa, including Bacillariophyta, Chlorophyta, and
Streptophyta, were also more abundant in the conventional paddy field compared to the no-
fertilizer paddy field (Fig. S7).
The number of OTUs detected, which serves as an index of species richness, was
consistently higher in the conventional paddy field than in the no-fertilizer paddy field ($P <$
0.0001; Fig. 5b; Table S2). In the conventional paddy field, the number of detected OTUs
ranged from approximately 600 to 800, consistently exceeding that in the no-fertilizer paddy
field.
The ecological community compositions identified through the quantitative eDNA
metabarcoding were also analyzed using NMDS (Fig. 5c). There was a statistically clear
difference between the two paddy fields, as well as a clear seasonal pattern (number of
weeks, $P < 0.0001$; farming practice, $P < 0.0001$; interaction, $P < 0.0001$). In the
conventional paddy field, the ecological community compositions were characterized by
species belonging to Actinomycetes, γ-Proteobacteria, β-Proteobacteria, Verrucomicrobia,
Oligoflexia, and Sphingobacteriia (Table S8; Fig. S8). The abundances of most of these
species were higher during the first half of the monitoring period in the conventional paddy
field, with differences between the two paddy fields diminishing after August. In the no-
fertilizer paddy field, the ecological community composition was predominantly
characterized by the dominance of Actinomycetes (PRO_Taxa00038; Table S8; Fig. S8). The
abundance of PRO_Taxa00038 was higher in the no-fertilizer paddy field than in the
conventional paddy field until August.
527

Figure 5 | Environmental DNA in water samples collected from the two paddy fields (**a**) The abundance (eDNA copies/ml water) of the ecological communities in the water samples. Each panel indicates the conventional or no-fertilizers paddy field. Colors indicate different taxonomic groups. (**b**) The number of OTUs detected in the water samples. Solid lines indicate mean values.(**c**) NMDS of the eDNA-based ecological community data based on Bray-Curtis dissimilarity (stress value = 0.1924). Green and and filled circle indicate data from the conventional paddy fields, and yellow and filled triangle indicate data from the no-fertilizer paddy field. *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$. "N.S." indicates that the difference between the two farming practices is not statistically clear.

3.4 Effects of ecological variables on the rice performance

Based on the time series data, we detected causal variables that affect rice performance,

specifically height growth (Fig. 6 and Table S9) (For results on SPAD values and stem

counts, see Figs. S9, S10 and Tables S10, S11). Using UIC, we detected 16 ecological

variables that causally impacted rice growth. The OTUs with the strongest positive effects on

rice height growth were Undetermined taxon (ITS_Taxa00138), β-Proteobacteria

(PRO_Taxa00019), and Bacteria (PRO_Taxa00332) (Fig. 6a). These species were more

abundant in the conventional rice field (Table S9 and Fig. S11), contributing to improved rice
growth in that environment. While our analysis included both microbes and macrobes (e.g.,
invertebrate taxa), only *Digitaria ciliaris* (Poaceae) on the ridge had statistically clear causal
effects on rice growth. Several bacterial species from Bacteroidota, Pseudomonadota,
Burkholderiales, *Flavobacterium* sp., and *Polynucleobacter cosmopolitanus* also showed
clear causal influences on rice growth. Additionally, a species from Chlamidomonadales
affected rice growth. Species from Cochliopodium, Chlorophyceae, Ploima, Actinomycetes,
eukaryota, and bacteria significantly influenced SPAD values (Fig. S9), while only two
species from Silvanigrellaceae and Chlorophyceae had significant effects on stem counts
(Fig. S10).

546   Importantly, the impact of these ecological members per abundance on overall rice
performance was largely consistent between the two paddy fields (Figs. 6a, S9a, and S10a
and Tables S9–S11). Differences in their overall contributions to rice growth between the two
fields (Figs. 6c, S9c, and S10c and Tables S9–S11) were primarily attributed to variations in
the abundance of these causal species (Figs. 6b, S9b, and S10b and Tables S9–S11). Most
causally influential species were more abundant in the conventional paddy field compared to
the no-fertilizer paddy field (Fig. S11). The abundances of ecological community members
positively influencing rice growth were generally higher in the conventional rice field than in
the no-fertilizer rice field.

Figure 6 | Effects of ecological variables on rice height growth in the conventional and no-fertilizer paddy fields
Each panel shows (**a**) the effects of ecological variables on overall rice height growth per unit variable (i.e., coefficients of the MDR S-map), (**b**) the abundance of ecological variables, and (**c**) the overall effects of ecological variables on rice height growth (calculated by "the MDR S-map coefficient × Abundance"), respectively. Green and and filled circle indicate data from the conventional paddy fields, and yellow and filled triangle indicate data from the no-fertilizer paddy fields. The y-axis labels include assigned taxa names, taxa code, and effect delay (tp; e.g., $tp = -1$ indicates the taxon has effects on the rice performance one week later).

4. DISCUSSION

In the present study, we conducted thorough monitoring to identify ecological variables that affect rice growth performance in the two paddy fields with different farming practices. We observed clear differences in rice performance regarding height growth, SPAD values, stem counts, and yields. Despite the lack of fertilizer inputs, the no-fertilizer paddy field still produced approximately 40–50% of the yields produced in the conventional paddy field. The rice yield in the no-fertilizer field was lower than that reported in previous research (e.g., Okumura, 2002), likely due to differences in rice variety (*Hinohikari* in the present study versus *Beniasahi* in the previous research). Ecological variables, including plant species on the paddy ridge and ecological communities identified using the quantitative eDNA metabarcoding, also showed clear differences between the fields. Using time series analysis, we identified ecological variables that influence rice performance.

4.1 Contrasting ecological communities in the two paddy fields

First, we observed statistically clear differences in ecological communities between the two paddy fields. The conventional paddy field exhibited a lower abundance of Cyperaceae species on the ridge (Fig. 3), a higher total eDNA concentration in June and July (Fig. 5a), and greater OTU richness (Fig. 5b). Specific OTUs from γ-proteobacteria, Chlorophyceae, Verrucomicrobiota, Actinomycetes, and other taxa displayed distinct differences between the two fields (Fig. S8 and Table S7). Given that the climate variables were nearly identical in the two paddy fields, the farming practices, namely the use of chemical fertilizers and pesticides, were the primary factors driving these differences.

4.1.1 Plant specialized metabolites and insect abundance

We observed only minor differences in plant specialized metabolites and insect abundance between the two paddy fields (Fig. 3). Among the metabolites measured, (*Z*)-3-hexenal, (*E*)-2-hexenal, and methyl salicylate showed slightly higher concentrations in the rice leaves of the no-fertilizer paddy field compared to those of the conventional field (Fig. 3a and Table S2). (*Z*)-3-hexenal and (*E*)-2-hexenal are common green leaf volatiles (GLVs) that can be released in response to mechanical damage and can activate defense genes against certain pathogenic fungi (Matsui & Koeduka, 2016). Methyl salicylate is a type of insect-induced plant volatile and can be emitted by rice seedlings in response to insect herbivory (Zhao et al., 2010). Since the no-fertilizer paddy field did not receive any pesticides, rice plants there may have been more frequently affected by arthropodal pests and microbial pathogens. The abundance of an OTU belonging to Actinomycetes (PRO_Taxa00038) was approximately 100 times higher in the no-fertilizer paddy field (Fig. S8), which may have contributed to the increased emission of the two GLVs.

　　　The abundances of the three insect groups were generally similar (Figs. 3, 4), but a statistically clear difference was found in the category of "other planthoppers," likely due to their higher abundance in the no-fertilizer paddy field during the first monitoring event. Interestingly, the increased abundance of "other planthoppers" in early June coincided with higher concentrations of (*Z*)-3-hexenal and (*E*)-2-hexenal, suggesting that herbivory damage

may contribute to the elevated levels of GLVs in the no-fertilizer paddy field. Overall,
however, the differences in plant specialized metabolites and insect groups were subtle, as
indicated by the NMDS (Fig. 4a). This suggests that fertilizers and pesticides have a lesser
impact on plant specialized metabolites and insect groups compared to their effects on plant
communities on the paddy ridge and other ecological communities (Figs. 4b and 5), at least in
the two paddy fields studied. The lesser effects of pesticides on the insect communities may
also have resulted from the fact that our survey plots were located within 50–70 m of each
other, so insects could migrate between the conventional paddy field and the no-fertilizer
paddy field (Fig. S1a).
608
4.1.2 Plant communities on the paddy ridge
In the conventional paddy field, all dominant species were annuals, while perennials were
more prevalent in the no-fertilizer paddy field. The plant species observed on the paddy
ridges are species commonly found in agricultural fields across Japan. Pesticide application
for weed control eliminates existing plants, allowing fast-growing annuals to dominate. In
contrast, mowing does not disrupt the underground structures, allowing perennials with low
growing points or those that reproduce vegetatively via rhizomes or stolons to thrive. In
particular, two perennial species from the Cyperaceae family, *Cyperus brevifolia* and
Fimbristylis diphylloides, were particularly dominant in the no-fertilizer paddy field. Many
Cyperaceae species are well adapted to open, sunny, moist environments, which constitute
major components of paddy ecosystems (Bryson & Carter, 2008). Specifically, *Cyperus*
brevifolia develops a robust rhizome network, enabling it to quickly establish a dense, space-
occupying mat (Shaw et al., 2011).
622
4.1.3 Ecological community compositions detected by eDNA metabarcoding
The increased total eDNA concentration in the conventional paddy field, primarily due to
higher microbial abundance, was likely influenced by fertilizer inputs. While nitrogen
addition to the soil can lead to a reduction in microbial biomass (Treseder, 2008), it can also
enhance phytoplankton biomass in freshwater ecosystems (Søndergaard et al., 2017). In the
present study, major photosynthetic taxa, including Chlorophyta, were more abundant in the
conventional paddy field (Fig. S7), likely due to the higher nitrogen input. The greater
abundance of these photosynthetic taxa contributes to increased carbon input into the system,
which may boost microbial biomass (and total eDNA concentration) in the conventional
paddy field. Additionally, the higher productivity of rice further adds to the carbon input.
Indeed, Kamata et al. (1991) reported a greater concentration of soil carbon in conventional
paddy soil compared to no-fertilizer paddy soil (in the no-fertilizer paddy soil studied in
Ritto, Shiga Prefecture, Japan).
The nitrogen fertilizer and increased carbon input were likely the primary factors
driving the differences in community composition (Figs. 5 and S6–S8). Changes in
community composition in response to nutrient addition or carbon input have frequently been
reported in paddy fields (e.g., Dong et al., 2014; Z. Zhao et al., 2019). Actinomycetes
(PRO_Taxa00038), which contributed to the differences between the two paddy fields, share
the same 16S sequence fragment with those found in Neotropical floodplain lakes (Targueta

et al., 2023), suggesting a potential association with nutrient-poor conditions typical of
natural ecosystems rather than agricultural environments. Several Chlorophyceae OTUs
(EUK_Taxa00281, EUK_Taxa00423, and EUK_Taxa00170) were more abundant in the
conventional paddy fields than in the no-fertilizer fields, and Chlorophyceae have been
detected in rice paddy fields where light and nutrient availability are high (Sahu, 2012).
647

4.2 Influences of ecological communities on the rice performance

Our time series analysis detected ecological variables that can causally influence rice
performance (Figs. 6, S9–S11 and Tables S9–S11). Among the causal taxa affecting rice
height growth, some, such as ITS_Taxa00138 and ITS_Taxa0007, showed much higher
abundances in the conventional paddy field than in the no-fertilizer field. Although we could
not assign species names based on the OTU sequences, a common characteristic of these taxa
was their dominance during the early monitoring period, particularly in June (Fig. S11). This
pattern also applied to the only causal plant species on the paddy ridge, *Digitaria ciliaris*
(Fig. S3). While we could not identify the genus or species for most taxa, exceptions included
Flavobacterium spp. (PRO_Taxa00037 and COI_Taxa00221) and *Polynucleobacter*
cosmopolitanus (PRO_Taxa00024), which were also dominant in the conventional paddy
field in June. *Flavobacterium* is a genus of Gram-negative bacteria found in various
ecosystems, and a related species to COI_Taxa00221, *F. keumense*, was isolated from
freshwater ecosystems in South Korea (Ekwe et al., 2017). *Polynucleobacter cosmopolitanus*
is a free-living, cosmopolitan freshwater bacterioplankton frequently detected in ecosystems
in Japan (Watanabe et al., 2012). Unfortunately, physiological and ecological information
about these species is limited, making it difficult to infer the exact mechanisms by which
these taxa influence rice height growth in this study. Future research could leverage the
spatio-temporal data on these beneficial taxa, i. e., their general dominance in June and
higher detection rates in conventional paddy fields, to focus on identifying their ecological
and physiological roles.
Similarly, while we detected statistically clear causal effects of several species (e.g.,
Cochliopodium, Ploima, Actinomycetes, Silvanigrellaceae, and Chlorophyceae) on SPAD
values and the number of stems (Figs. S9, S10), it remains difficult to determine the exact
mechanisms by which they influence rice performance. Interestingly, some of these species
showed increased abundance during the middle of the monitoring period (from late June to
July). Thus, the mechanisms through which these taxa affect rice performance may differ
from those influencing rice height growth. Focusing on taxa that demonstrate clear temporal
patterns, such as PRO_Taxa00050 (Bacteria) and COI_Taxa00134 (Ploima, a rotifer species)
(Fig. S11), could provide insights into their roles in the paddy fields.
678

4.3 Potential factors that contribute to the maintenance of the rice performance in the paddy fields

While we could not determine the exact mechanisms by which ecological communities
influence rice performance in the paddy fields, our analysis provide insights into the
differences between the two fields. Importantly, the per-abundance effects of key ecological
community members on overall rice performance were generally similar across both paddy

fields (Figs. 6a, S9a–S10a). Statistically clear differences in the effects between the farming
practices were observed in only a few taxa (Tables S9–S11; three out of 16 species, zero out
of seven species, and one out of two species). In contrast, 13 of 16 species, four of seven
species, and both species in one category showed statistically clear differences in abundance
between the two paddy fields, with most showing lower abundance in the no-fertilizer field
than in the conventional field (Fig. S11). These findings suggest that, despite the differences
in rice yields, growth trajectories, and ecological communities between the two paddy fields,
the mechanisms or interactions between rice and ecological communities were largely
consistent. The abundance of influential species appears to contribute to the differences in
rice growth performance and yields. While it is clear that nutrient inputs (i.e., N-P-K
fertilizers) enhanced rice growth and yield in the conventional paddy fields, the abundance
and composition of ecological community members may also support rice growth. This
indicates that introducing these species could potentially improve rice growth and yields in
the no-fertilizer paddy field.
699

4.4 Advantages, limitations, implications and future direction

In this study, we integrated monitoring of rice performance, collection of plant specialized
metabolites, plant cover surveys on the ridge, insect survey, eDNA-based ecological
community monitoring, and a nonlinear time series causality detection to understand the
impact of ecological variables in actual rice paddy fields. The integration of field-based
monitoring with advanced causality detection methods enabled the detection of potential
causal ecological variables influencing rice growth in the real paddy fields.
707   One advantage of our research framework is its ability to elucidate ecological
phenomena and complex causal interactions under field conditions. Manipulative
experiments are among the most effective methods for understanding causal interactions in
biological systems. However, while laboratory experiments can be conducted (provided that
certain causal microbial species can be isolated and cultivated, or that other macrobial species
can be easily maintained in the laboratory), the interactions observed in controlled settings
may not necessarily reflect those in natural environments, as ecological interactions and
resulting dynamics are often context-dependent and nonlinear (Hsieh et al., 2005; Sugihara et
al., 2012; Ushio, 2022; Ushio, Hsieh, et al., 2018). In this context, field-based monitoring
combined with time series analysis for causal detection and quantification of interaction
strengths can provide a more realistic understanding of natural systems.
718   Nonetheless, we must address the technical limitations of our study for future research.
First, while quantitative eDNA metabarcoding offers a comprehensive and quantitative
ecological time series (Ushio, 2022; Ushio, Murakami, et al., 2018), several challenges
remain, particularly regarding abundance estimation and species identification. The
relationship between eDNA concentrations and species abundance, especially for macro-
organisms, remains an open question (Harrison et al., 2021). eDNA concentrations should be
considered "an index" of species abundance, as the rates of eDNA release and degradation
vary by species and environmental conditions (Dejean et al., 2011). For microbial species,
variations in the copy numbers of target genes, particularly 18S rRNA (Godhe et al., 2008),
complicate the estimation of abundance or biomass from metabarcoding data. Additionally,

assigning species names using short-read amplicon data is often challenging. Long-read
sequencing-based eDNA metabarcoding or shotgun metagenomic analysis may address these
issues associated with short-read amplicon sequencing, as they can provide more detailed
information, including intra-specific and individual variations within ecological communities
(Baloğlu et al., 2021; Nousias et al., 2025).
Another limitation is the number of time points in our time series. Although we made
substantial efforts to measure multiple ecological variables in the two paddy fields, our
monitoring was conducted weekly. While we detected "statistically clear" causal effects of
ecological variables, our sampling design did not allow us to capture dynamics occurring
within a timeframe shorter than one week. For instance, rice gene expression dynamics can
vary across multiple time scales, including seasonal, daily, hourly, and even shorter intervals
(Nagano et al., 2012). Utilizing more frequent and high-throughput field monitoring
technologies, such as unmanned aerial vehicles (UAVs) for rice monitoring (Chen et al.,
2024) and high-throughput, low-cost RNA-seq methods (Kamitani et al., 2019), would
greatly enhance the efficiency and accuracy of detecting causal ecological variables in field
conditions.
744

4.5 Conclusions

In the present study, we integrated various ecological monitoring methods and a time series-
based causality detection approach to explore the role of ecological variables in sustaining
rice performance in both conventional and no-fertilizer paddy fields in Japan. Our time series
analysis revealed several previously unrecognized ecological community members that
causally influenced rice performance. Interestingly, these members showed similar per-
abundance effects on rice in both fields but their abundance differed, suggesting that similar
mechanisms underlie the differing rice growth patterns, with abundance potentially driving
some of the differences observed. While we could not fully elucidate the detailed
mechanisms by which these communities affect rice performance, we demonstrated that our
framework—integrating field-based monitoring with advanced time series analysis—
facilitates an understanding of the role of ecological communities in rice paddy fields. Future
studies that incorporate more advanced monitoring and data analysis techniques will enhance
our understanding and management of rice paddy fields, promoting more sustainable crop
production.
760

References:

Baloğlu, B., Chen, Z., Elbrecht, V., Braukmann, T., MacDonald, S., & Steinke, D. (2021). A workflow for accurate metabarcoding using nanopore MinION sequencing. *Methods in Ecology and Evolution*, *12*(5), 794–804. https://doi.org/10.1111/2041-210X.13561

Benjamini, Y., & Hochberg, Y. (1995). Controlling the False Discovery Rate: A Practical and Powerful Approach to Multiple Testing. *Journal of the Royal Statistical Society: Series B (Methodological)*, *57*(1), 289–300. https://doi.org/10.1111/j.2517-6161.1995.tb02031.x

Bottrell, D. G., & Schoenly, K. G. (2012). Resurrecting the ghost of green revolutions past: The brown planthopper as a recurring threat to high-yielding rice production in tropical Asia. *Journal of Asia-Pacific Entomology*, *15*(1), 122–140. https://doi.org/10.1016/j.aspen.2011.09.004

Bryson, C. T., & Carter, R. (2008). The significance of Cyperaceae as weeds. In *Sedges: Uses, Diversity, and Systematics of the Cyperaceae* (Vol. 108, p. 309). Missouri Botanical Garden Press. https://www.mbgpress.org/product-p/msb-108.htm

Callahan, B. J., McMurdie, P. J., Rosen, M. J., Han, A. W., Johnson, A. J. A., & Holmes, S. P. (2016). DADA2: High-resolution sample inference from Illumina amplicon data. *Nature Methods*, *13*(7), 581–583. https://doi.org/10.1038/nmeth.3869

Cenci, S., Sugihara, G., & Saavedra, S. (2019). Regularized S-map for inference and forecasting with noisy ecological time series. *Methods in Ecology and Evolution*, *10*(5), 650–660. https://doi.org/10.1111/2041-210X.13150

Chang, C.-W., Miki, T., Ushio, M., Lu, H.-P., Shiah, F.-K., & Hsieh, C. (2021). Reconstructing large networks with time-varying interactions. *arXiv:2102.04069 [q-Bio]*. http://arxiv.org/abs/2102.04069

Chen, R., Lu, H., Wang, Y., Tian, Q., Zhou, C., Wang, A., Feng, Q., Gong, S., Zhao, Q., & Han, B. (2024). High-throughput UAV-based rice panicle detection and genetic mapping of heading-date-related traits. *Frontiers in Plant Science*, *15*, 1327507. https://doi.org/10.3389/fpls.2024.1327507

Deiner, K., Fronhofer, E. A., Mächler, E., Walser, J.-C., & Altermatt, F. (2016). Environmental DNA reveals that rivers are conveyer belts of biodiversity information. *Nature Communications*, *7*, 12544.

Dejean, T., Valentini, A., Duparc, A., Pellier-Cuit, S., Pompanon, F., Taberlet, P., & Miaud, C. (2011). Persistence of Environmental DNA in Freshwater Ecosystems. *PLOS ONE*, *6*(8), e23398.

Dong, W.-Y., Zhang, X.-Y., Dai, X.-Q., Fu, X.-L., Yang, F.-T., Liu, X.-Y., Sun, X.-M., Wen, X.-F., & Schaeffer, S. (2014). Changes in soil microbial community composition in response to fertilization of paddy soils in subtropical China. *Applied Soil Ecology*, *84*, 140–147. https://doi.org/10.1016/j.apsoil.2014.06.007

Dushoff, J., Kain, M. P., & Bolker, B. M. (2019). I can see clearly now: Reinterpreting statistical significance. *Methods in Ecology and Evolution*, *10*(6), 756–759. https://doi.org/10.1111/2041-210X.13159

Ekwe, A. P., Ahn, J., & Kim, S. B. (2017). Flavobacterium keumense sp. Nov., isolated from

freshwater. *International Journal of Systematic and Evolutionary Microbiology*, *67*(7),
2166–2170. https://doi.org/10.1099/ijsem.0.001920

FAO. (2024). *World Food and Agriculture – Statistical Yearbook 2024*. FAO ;
https://doi.org/10.4060/cd2971en

Godfray, H. C. J., Beddington, J. R., Crute, I. R., Haddad, L., Lawrence, D., Muir, J. F., Pretty,
J., Robinson, S., Thomas, S. M., & Toulmin, C. (2010). Food security: The challenge
of feeding 9 billion people. *Science*, *327*(5967), 812–818.
https://doi.org/10.1126/science.1185383

Godhe, A., Asplund, M. E., Härnström, K., Saravanan, V., Tyagi, A., & Karunasagar, I. (2008).
Quantification of Diatom and Dinoflagellate Biomasses in Coastal Marine Seawater
Samples by Real-Time PCR. *Applied and Environmental Microbiology*, *74*(23), 7174–
7182. https://doi.org/10.1128/AEM.01298-08

Harrison, J. G., John Calder, W., Shuman, B., & Alex Buerkle, C. (2021). The quest for absolute
abundance: The use of internal standards for DNA-based community ecology.
Molecular Ecology Resources, *21*(1), 30–43. https://doi.org/10.1111/1755-0998.13247

He, X., Jeffery, N. W., Stanley, R. R. E., Hamilton, L. C., Rubidge, E. M., & Abbott, C. L.
(2023). eDNA metabarcoding enriches traditional trawl survey data for monitoring
biodiversity in the marine environment. *ICES Journal of Marine Science*, *80*(5), 1529–
1538. https://doi.org/10.1093/icesjms/fsad083

Hsieh, C., Glaser, S. M., Lucas, A. J., & Sugihara, G. (2005). Distinguishing random
environmental fluctuations from ecological catastrophes for the North Pacific Ocean.
Nature, *435*(7040), 336–340. https://doi.org/10.1038/nature03553

Hua, L., Liu, J., Zhai, L., Xi, B., Zhang, F., Wang, H., Liu, H., Chen, A., & Fu, B. (2017). Risks
of phosphorus runoff losses from five Chinese paddy soils under conventional
management practices. *Agriculture, Ecosystems & Environment*, *245*, 112–123.
https://doi.org/10.1016/j.agee.2017.05.015

Ji, Z.-J., Zhao, L.-F., Zhang, T.-J., Dai, R.-X., Tang, J.-J., Hu, L.-L., & Chen, X. (2023).
Coculturing rice with aquatic animals promotes ecological intensification of paddy
ecosystem. *Journal of Plant Ecology*, *16*(6), rtad014.
https://doi.org/10.1093/jpe/rtad014

Kamata, M., Kawamura, S., & Sugino, M. (1991). Annual Changes in Soil Nitrogen and
Microflora in Paddy Fields not Fertilized Long-term and Fertilized Paddy Fields.
Memoirs of the Faculty of Agriculture of Kinki University, *24*, 1–13.

Kamitani, M., Kashima, M., Tezuka, A., & Nagano, A. J. (2019). Lasy-Seq: A high-throughput
library preparation method for RNA-Seq and its application in the analysis of plant
responses to fluctuating temperatures. *Scientific Reports*, *9*(1), Article 1.
https://doi.org/10.1038/s41598-019-43600-0

Kashima, M., Sakamoto, R. L., Saito, H., Ohkubo, S., Tezuka, A., Deguchi, A., Hashida, Y.,
Kurita, Y., Iwayama, K., Adachi, S., & Nagano, A. J. (2021). Genomic Basis of
Transcriptome Dynamics in Rice under Field Conditions. *Plant and Cell Physiology*,
62(9), 1436–1445. https://doi.org/10.1093/pcp/pcab088

Ko, C.-Y., Asano, S., Lin, M.-J., Ikeya, T., Peralta, E. M., Triño, E. M. C., Uehara, Y., Ishida,

T., Iwata, T., Tayasu, I., & Okuda, N. (2021). Rice paddy irrigation seasonally impacts stream benthic macroinvertebrate diversity at the catchment level. *Ecosphere*, *12*(5), e03468. https://doi.org/10.1002/ecs2.3468

Krol, L., Van der Hoorn, B., Gorsich, E. E., Trimbos, K., Bodegom, P. M. van, & Schrama, M. (2019). How Does eDNA Compare to Traditional Trapping? Detecting Mosquito Communities in South-African Freshwater Ponds. *Frontiers in Ecology and Evolution*, *7*. https://doi.org/10.3389/fevo.2019.00260

Li, S.-X., Jiang, J., Lv, W.-G., Siemann, E., Woodcock, B. A., Wang, Y.-Q., Cavalieri, A., Bai, N.-L., Zhang, J.-Q., Zheng, X.-Q., Zhang, H.-L., Zhang, H.-Y., Zhang, Y., & Wan, N.-F. (2025). Rice-fish co-culture promotes multiple ecosystem services supporting increased yields. *Agriculture, Ecosystems & Environment*, *381*, 109417. https://doi.org/10.1016/j.agee.2024.109417

Matsui, K., & Koeduka, T. (2016). Green Leaf Volatiles in Plant Signaling and Response. *Sub-Cellular Biochemistry*, *86*, 427–443. https://doi.org/10.1007/978-3-319-25979-6_17

Menegat, S., Ledo, A., & Tirado, R. (2022). Greenhouse gas emissions from global production and use of nitrogen synthetic fertilisers in agriculture. *Scientific Reports*, *12*(1), 14490. https://doi.org/10.1038/s41598-022-18773-w

Miya, M., Sato, Y., Fukunaga, T., Sado, T., Poulsen, J. Y., Sato, K., Minamoto, T., Yamamoto, S., Yamanaka, H., Araki, H., Kondoh, M., & Iwasaki, W. (2015). MiFish, a set of universal PCR primers for metabarcoding environmental DNA from fishes: Detection of more than 230 subtropical marine species. *Royal Society Open Science*, *2*(7), 150088. https://doi.org/10.1098/rsos.150088

Nagano, A. J., Sato, Y., Mihara, M., Antonio, B. A., Motoyama, R., Itoh, H., Nagamura, Y., & Izawa, T. (2012). Deciphering and Prediction of Transcriptome Dynamics under Fluctuating Field Conditions. *Cell*, *151*(6), 1358–1369. https://doi.org/10.1016/j.cell.2012.10.048

Natuhara, Y. (2013). Ecosystem services by paddy fields as substitutes of natural wetlands in Japan. *Ecological Engineering*, *56*, 97–106. https://doi.org/10.1016/j.ecoleng.2012.04.026

Nousias, O., McCauley, M., Stammnitz, M. R., Farrell, J. A., Koda, S. A., Summers, V., Eastman, C. B., Duffy, F. G., Duffy, I. J., Whilde, J., & Duffy, D. J. (2025). Shotgun sequencing of airborne eDNA achieves rapid assessment of whole biomes, population genetics and genomic variation. *Nature Ecology & Evolution*, *9*(6), 1043–1060. https://doi.org/10.1038/s41559-025-02711-w

Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G. L., Solymos, P., Stevens, M. H. H., & Wagner, H. (2008). *vegan: Community Ecology Package* (R package). http://cran.r-project.org/, http://vegan.r-forge.r-project.org/

Okumura, T. (2002). Rice Production in Unfertilized Paddy Field: Mechanism of grain production as estimated from nitrogen economy. *Plant Production Science*, *5*(1), 83–88. https://doi.org/10.1626/pps.5.83

Osada, Y., & Ushio, M. (2021). *R package for Unified Information-theoretic Causality* (Version 0.1.5) [Computer software]. Zenodo. https://doi.org/10.5281/zenodo.5163234

Osada, Y., Ushio, M., & Michio, K. (2023). A unified framework for nonparametric causality
detection. *bioRxiv*, 2023.04.20.537743. https://doi.org/10.1101/2023.04.20.537743

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., & R Development Core Team. (2009). *nlme:*
Linear and Nonlinear Mixed Effects of Models (R package).

Qian, H., Zhu, X., Huang, S., Linquist, B., Kuzyakov, Y., Wassmann, R., Minamikawa, K.,
Martinez-Eixarch, M., Yan, X., Zhou, F., Sander, B. O., Zhang, W., Shang, Z., Zou, J.,
Zheng, X., Li, G., Liu, Z., Wang, S., Ding, Y., … Jiang, Y. (2023). Greenhouse gas
emissions and mitigation in rice agriculture. *Nature Reviews Earth & Environment*,
4(10), 716–732. https://doi.org/10.1038/s43017-023-00482-1

R Core Team. (2024). *R: A language and environment for statistical computing* [Computer
software]. R Foundation for Statistical Computing. https://www.R-project.org/

Riemann, M., Haga, K., Shimizu, T., Okada, K., Ando, S., Mochizuki, S., Nishizawa, Y.,
Yamanouchi, U., Nick, P., Yano, M., Minami, E., Takano, M., Yamane, H., & Iino, M.
(2013). Identification of rice Allene Oxide Cyclase mutants and the function of
jasmonate for defence against Magnaporthe oryzae. *The Plant Journal*, *74*(2), 226–238.
https://doi.org/10.1111/tpj.12115

Rim, H., Uefune, M., Ozawa, R., & Takabayashi, J. (2018). An omnivorous arthropod,
Nesidiocoris tenuis, induces gender-specific plant volatiles to which conspecific males
and females respond differently. *Arthropod-Plant Interactions*, *12*(4), 495–503.
https://doi.org/10.1007/s11829-018-9612-2

Sahu, A. K. and R. (2012). Diversity of Algae (Cholorophyceae) in Paddy Fields of Lalgutwa
Area, Ranchi, Jharkhand. *Journal of Applied Pharmaceutical Science*, *2*(11), 092–095.
https://doi.org/10.7324/JAPS.2012.21116

Savary, S., Willocquet, L., Pethybridge, S. J., Esker, P., McRoberts, N., & Nelson, A. (2019).
The global burden of pathogens and pests on major food crops. *Nature Ecology &*
Evolution, *3*(3), Article 3. https://doi.org/10.1038/s41559-018-0793-y

Schreiber, T. (2000). Measuring Information Transfer. *Physical Review Letters*, *85*(2), 461–
464. https://doi.org/10.1103/PhysRevLett.85.461

Shaw, D. A., Wilen, C. A., Cudney, D. W., & Elmore, C. L. (2011). *Green Kyllinga* (pp. 1–4).
UC Statewide, Integrated Pest Management Program.
https://ipm.ucanr.edu/legacy_assets/PDF/PESTNOTES/pngreenkyllinga.pdf

Shiojiri, K., Ozawa, R., Uefune, M., & Takabayashi, J. (2021). Field-Grown Rice Plants
Become More Productive When Exposed to Artificially Damaged Weed Volatiles at the
Seedling Stage. *Frontiers in Plant Science*, *12*.
https://doi.org/10.3389/fpls.2021.692924

Søndergaard, M., Lauridsen, T. L., Johansson, L. S., & Jeppesen, E. (2017). Nitrogen or
phosphorus limitation in lakes and its impact on phytoplankton biomass and submerged
macrophyte cover. *Hydrobiologia*, *795*(1), 35–48. https://doi.org/10.1007/s10750-017-
3110-x

Sugihara, G. (1994). Nonlinear Forecasting for the Classification of Natural Time Series.
Philosophical Transactions of the Royal Society A: Mathematical, Physical and
Engineering Sciences, *348*(1688), 477–495. https://doi.org/10.1098/rsta.1994.0106

Sugihara, G., & May, R. M. (1990). Nonlinear forecasting as a way of distinguishing chaos
from measurement error in time series. *Nature*, *344*(6268), 734–741.
https://doi.org/10.1038/344734a0

Sugihara, G., May, R., Ye, H., Hsieh, C., Deyle, E., Fogarty, M., & Munch, S. (2012). Detecting
causality in complex ecosystems. *Science*, *338*(6106), 496–500.
https://doi.org/10.1126/science.1227079

Taberlet, P., Bonin, A., Zinger, L., & Coissac, E. (2018). *Environmental DNA: For Biodiversity
Research and Monitoring*. OUP Oxford.

Tanabe, A. S., & Toju, H. (2013). Two new computational methods for universal DNA
barcoding: A benchmark using barcode sequences of bacteria, archaea, animals, fungi,
and land plants. *PloS One*, *8*(10), e76910.
https://doi.org/10.1371/journal.pone.0076910

Targueta, C. P., Antunes, A. M., Machado, K. B., Fernandes, J. G., Telles, M. P. D. C., Vieira,
L. C. G., Logares, R., Nabout, J. C., & Soares, T. N. (2023). Diversity of eukaryotic
and prokaryotic microbiota revealed by metabarcoding in Neotropical floodplain lakes.
Anais Da Academia Brasileira de Ciências, *95*, e20201578.
https://doi.org/10.1590/0001-3765202320201578

Toju, H., Peay, K. G., Yamamichi, M., Narisawa, K., Hiruma, K., Naito, K., Fukuda, S., Ushio,
M., Nakaoka, S., Onoda, Y., Yoshida, K., Schlaeppi, K., Bai, Y., Sugiura, R., Ichihashi,
Y., Minamisawa, K., & Kiers, E. T. (2018). Core microbiomes for sustainable
agroecosystems. *Nature Plants*, *4*(5), 247–257. https://doi.org/10.1038/s41477-018-
0139-4

Treseder, K. K. (2008). Nitrogen additions and microbial biomass: A meta-analysis of
ecosystem studies. *Ecology Letters*, *11*(10), 1111–1120. https://doi.org/10.1111/j.1461-
0248.2008.01230.x

Ushio, M. (2019). Use of a filter cartridge combined with intra-cartridge bead-beating
improves detection of microbial DNA from water samples. *Methods in Ecology and
Evolution*, *10*(8), 1142–1156. https://doi.org/10.1111/2041-210X.13204

Ushio, M. (2022). Interaction capacity as a potential driver of community diversity.
Proceedings of the Royal Society B: Biological Sciences, *289*(1969), 20212690.
https://doi.org/10.1098/rspb.2021.2690

Ushio, M. (2022). *macam: A collection of miscellaneous functions for ecology in R* [R].
https://github.com/ong8181/macam (Original work published 2022)

Ushio, M., Hsieh, C., Masuda, R., Deyle, E. R., Ye, H., Chang, C.-W., Sugihara, G., & Kondoh,
M. (2018). Fluctuating interaction network and time-varying stability of a natural fish
community. *Nature*, *554*, 360–363.

Ushio, M., Murakami, H., Masuda, R., Sado, T., Miya, M., Sakurai, S., Yamanaka, H.,
Minamoto, T., & Kondoh, M. (2018). Quantitative monitoring of multispecies fish
environmental DNA using high-throughput sequencing. *Metabarcoding and
Metagenomics*, *2*, e23297. https://doi.org/10.3897/mbmg.2.23297

Ushio, M., Saito, H., Tojo, M., & Nagano, A. J. (2023). An ecological network approach for
detecting and validating influential organisms for rice growth. *eLife*, *12*, RP87202.

https://doi.org/10.7554/eLife.87202

Wang, J., Wang, D., Zhang, G., Wang, Y., Wang, C., Teng, Y., & Christie, P. (2014). Nitrogen
and phosphorus leaching losses from intensively managed paddy fields with straw
retention. *Agricultural Water Management*, *141*, 66–73.
https://doi.org/10.1016/j.agwat.2014.04.008

Watanabe, K., Komatsu, N., Kitamura, T., Ishii, Y., Park, H.-D., Miyata, R., Noda, N.,
Sekiguchi, Y., Satou, T., Watanabe, M., Yamamura, S., Imai, A., & Hayashi, S. (2012).
Ecological niche separation in the Polynucleobacter subclusters linked to quality of
dissolved organic matter: A demonstration using a high sensitivity cultivation-based
approach. *Environmental Microbiology*, *14*(9), 2511–2525.
https://doi.org/10.1111/j.1462-2920.2012.02815.x

Wickham, H. (2009). *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.
http://ggplot2.org

Wing, R. A., Purugganan, M. D., & Zhang, Q. (2018). The rice genome revolution: From an
ancient grain to Green Super Rice. *Nature Reviews Genetics*, *19*(8), Article 8.
https://doi.org/10.1038/s41576-018-0024-z

Wood, S. N. (2004). Stable and efficient multiple smoothing parameter estimation for
generalized additive models. *Journal of the American Statistical Association*, *99*, 673–
686.

Wright, E. S. (2016). Using DECIPHER v2.0 to analyze big biological sequence data in R. *The
R Journal*, *8*(1), 352–359.

Ye, H., & Sugihara, G. (2016). Information leverage in interconnected ecosystems:
Overcoming the curse of dimensionality. *Science*, *353*(6302), 922 LP – 925.

Zhang, Q. (2007). Strategies for developing Green Super Rice. *Proceedings of the National
Academy of Sciences*, *104*(42), 16402–16409.
https://doi.org/10.1073/pnas.0708013104

Zhao, N., Guan, J., Ferrer, J.-L., Engle, N., Chern, M., Ronald, P., Tschaplinski, T. J., & Chen,
F. (2010). Biosynthesis and emission of insect-induced methyl salicylate and methyl
benzoate from rice. *Plant Physiology and Biochemistry: PPB*, *48*(4), 279–287.
https://doi.org/10.1016/j.plaphy.2010.01.023

Zhao, Z., Ge, T., Gunina, A., Li, Y., Zhu, Z., Peng, P., Wu, J., & Kuzyakov, Y. (2019). Carbon
and nitrogen availability in paddy soil affects rice photosynthate allocation, microbial
community composition, and priming: Combining continuous 13C labeling with PLFA
analysis. *Plant and Soil*, *445*(1), 137–152. https://doi.org/10.1007/s11104-018-3873-5

1007

1008

Acknowledgments

We thank Itsuo Kitagawa and Keisuke Katsura for assistance in the field survey, Asako Kawai for assistance in the DNA library preparations, and Akifumi Sugiyama for assistance in the field sampling. This research was supported by KAKENHI (B) 17H03811 and 20H02922 (to KO), PRESTO (JPMJPR16O2) from the Japan Science and Technology Agency (JST), the Hakubi Project in Kyoto University (to MU), and The Hong Kong University of Science and Technology Startup Fund (to MU). We used generative AIs to improve the clarity of the manuscript text. We reviewed and edited all AI-generated content to ensure accuracy and appropriateness.

Author Contributions

MU, HS, RO, KS, and JT conceived and designed research; MU, HS, YS, KO, RO and KS performed field monitoring and analyzed samples; MU performed time-series analysis; MU summarized and visualized the data; MU wrote the first draft; all authors discussed the results and completed the manuscript.

Conflicts of Interest

The authors declare no conflict of interest.

Data Availability Statement

All scripts and raw data except for the sequence data used in the present study are available on Github (https://github.com/ong8181/ogura-rice2017). High-throughput sequence data was deposited in DDBJ Sequence Read Archives (DRA) (BioProject ID = PRJDB9326, DRA Run ID = DRR709379-DRR709530 [16S rRNA], DRR709531-DRR709682 [18S rRNA], DRR709683-DRR709834 [COI], and DRR709835-DRR709986 [ITS]).