Temporal Clonal Tracing and Functional Perturbation Reveal Niche-Adaptive and Tumor-Intrinsic IFNγ Dependencies Driving Ovarian Cancer Metastasis

Emilija Aleksandrovic[1,3], Shaneann R. Fross[1,3], Samantha M. Golomb[1,3], Xiyu Liu[1,3], Zhuo Zhao[2,3], Nikitha M. Das[1,3], Tanner C. Reese[1,3], Wei Ma[1,3], Jacqueline Lopez[4], M. Sharon Stack[5,6], Min Zhao[1,3], Siyuan Zhang[1, 3]

Affiliations:

1. Department of Pathology, UT Southwestern Medical Center, Dallas, TX, 75390, USA
2. Lyda Hill Department of Bioinformatics, School of Public Health, UT Southwestern Medical Center, Dallas, TX, 75390, USA
3. Harold C. Simmons Comprehensive Cancer Center, Dallas, TX, 75390, USA
4. Department of Biological Sciences, College of Science, University of Notre Dame, Notre Dame, IN 46556, USA
5. Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA
6. Mike and Josie Harper Cancer Research Institute, University of Notre Dame, 1234 N. Notre Dame Avenue, South Bend, IN 46617, USA

* Correspondence to: Siyuan Zhang, M.D., Ph.D., Department of Pathology, UT Southwestern Medical Center, ND6.200A, 6001 Forest Park Rd., Dallas, TX 75235. E-mail: Siyuan.Zhang@UTSouthwestern.edu; Telephone: 214-648-6537

Key Words: Metastasis, Single cell analysis, Barcoding, Lineage tracing, CRISPR screening, Interferon, Ovarian Cancer, Omentum, Ascites.

SUMMARY

Metastasis is an emergent continuum, driven by evolving reciprocal adaptations between continuously disseminating tumor cells (DTCs) and the specialized metastatic niches of distant organs. The interplay between intrinsic and niche-driven mechanisms that enables DTCs to survive and home to distant organs remains incompletely understood. Here, using MetTag, a single-cell barcoding and transcriptome profiling approach with time-stamped batch identifiers (BC.IDs), we mapped temporal, clonal dynamics of DTCs and the immune cell landscape across ovarian cancer metastatic niches. Deep sequencing of barcodes revealed preferred enrichment of early-disseminated clones across metastatic niches. Mechanistically, single-cell RNA sequencing (scRNA-seq) coupled with velocity analyses in ascites and metastasis-bearing omenta uncovered an emergent, distinct interferon-gamma (IFNγ) centric transcriptional trajectory, enriched among early seeding clones. Moreover, *in vivo* CRISPR/Cas9 screening of metastatic niche-specific signatures demonstrated that genes belonging to the ascites IFNγ signature, including *Marco*, *Gbp2b, and Slfn1,* are functionally important for peritoneal metastasis. Knockout of IFNγ receptor 1 (*Ifngr1*) in tumor cells significantly reduced metastatic burden and extended survival, underscoring the importance of tumor cell intrinsic IFNγ signaling in ovarian cancer metastasis. Furthermore, we identified that the tumor intrinsic IFNγ response and ascites-derived tumor-associated macrophages (TAMs) protect cancer cells from anoikis-mediated death within the IFNγ-rich ascites environment. Our study resolves temporal dynamics of disseminating tumor cells and highlights an ascites-driven, IFNγ program as a necessary pro-metastatic adaptation in the ovarian metastasis cascade.

INTRODUCTION

Cancer metastasis encompasses continuous metastatic dissemination from the primary tumor, followed by passive clonal selection, and active epigenetic niche adaptation in response to the metastatic niche[1]. Unlike most cancer types, where hematogenous metastasis is common, high-grade serous ovarian carcinoma (HGSOC) exhibits a unique form of transcoelomic, local spreading within the peritoneal cavity[2]. In HGSOC,

detached cancer cells from the fallopian tube are carried by the physiological flow of

peritoneal fluid to implant into intraperitoneal tissue, particularly into the omentum[3]. This

unique mode of metastatic dissemination leads to continuous and widespread

intraperitoneal metastases in the peritoneal cavity before colonizing distant organs[3].

Resolving the clonal, temporal, and microenvironmental determinants along the ovarian

cancer metastatic continuum has long been a challenge in the field.

Malignant ascites formed in HGSOC patients is a known "reservoir" of floating cancer

cells and cell clusters that can seed the omentum or other organs in the peritoneal

cavity[4,5]. Aside from cancer cells, malignant ascites contains immune cells and acellular

factors that influence pro-metastatic programs[6]. Notably, malignant ascites are enriched

in pro-metastatic cytokines, which increase cancer cell migration and correlate with

reduced disease-free progression[7,8]. In ascites, after shedding from the primary tumor,

ovarian cancer cells acquire resistance to anoikis, a critical mechanism for surviving

loss of attachment and enabling spread to other organs[9]. A key knowledge gap remains

in how cellular and acellular factors in malignant ascites regulate anoikis resistance and

shape the timing of successful colonization of intraperitoneal compartments.

Here, we conducted comprehensive lineage tracing, single-cell RNA sequencing

(scRNA-seq), *in vivo* CRISPR screening, and Cre-based niche-tracing analysis to

dissect the complex interplay between continuously arriving DTCs and ovarian

metastatic niche microenvironments. We identified that early DTCs in ascites are

dependent on niche-adaptive IFNγ response genes for omental colonization. We

additionally found that the interferon-rich ascites microenvironment harbors a

macrophage population capable of protecting cancer cells from anoikis-mediated death

via direct contact. Together, our data provide novel insights into the temporal dynamics

of ovarian cancer clonal dynamics and highlight ascites-specific factors that prime

cancer cells for successful omental colonization.

RESULTS

MetTag Captures Clonal Lineage Across Metastatic Sites and Across Time

To capture the transcriptional dynamics across ovarian cancer metastasis, we employed an RNA-level barcoding system based on the LARRY (<u>L</u>ineage <u>A</u>nd <u>R</u>NA <u>R</u>ecover<u>y</u>) barcoding construct, enabling single-cell transcriptome-level lineage tracing[10]. In brief, we subcloned the publicly available highly diverse LARRY library downstream of a U6 promoter in a CROP-seq backbone. We next added a unique batch identifier "BC.ID" to the LARRY barcode for distinguishing early from later-arriving DTCs. In our "MetTag" system, early DTCs refer to the first two BC.ID batches and late DTCs represent the last two DTC batches. Individually barcoded and BC.ID batch-coded cells were later injected across six different time points to model early, mid, and late DTCs. Within the LARRY transcript, a Chromium 10x capture sequence allows for simultaneous capture of the LARRY transcript with mRNA transcripts (**Fig.1A, S1A**). Diversity of each BC.ID library batch was confirmed by deep next-generation sequencing (NGS). We observed a relatively consistent diversity across the six BC.ID library batches comparable with the original depositor LARRY library (**Fig.S1B-D**).

Next, we generated six unique isogenic ID8 p53$^{-/-}$ murine ovarian cancer pools (BC.ID1-6) with MetTag-LARRY-GFP lentiviral libraries harboring BC.IDs (**Fig.1A**). To model continuous seeding from the primary tumor, transduced cells were IP injected in a fractionated manner at low doses. The injections were separated by 48 hours to allow immune and stromal responses to occur after exposure to each DTC batch (**Fig.1B**). Four weeks after the final DTC exposure, ascitic fluid (Asc), omenta (Om), and peritoneal fat pads (FPs) were extracted. These organs and compartments consistently developed metastatic foci (**Fig.S1E**). This experimental scheme was performed under several immune cell depletion conditions, including wild type C57BL/6J (WT) mice, immunocompromised NOD.Cg-Rag1^{tm1Mom}IL2rg^{tm1Wjl}/SzJ (NRG) mice (lacking functional T, B, and NK cells), and under conditions of macrophage depletion via chlodrosome administration. Temporally spaced DTC exposure resulted in robust metastasis formation in fully immunocompetent mice, evidenced by ascites accumulation and positive GFP signal in the omentum and peritoneal adipose tissue (**Fig.S1E top**). After grouping BC.IDs into early timepoint "TP" (*BC.ID1, BC.ID2*), mid

(*BC.ID3, BC.ID4*), and late (*BC.ID5, BC.ID6*) arrivers, most BC.IDs detected using

amplicon DNA sequencing represent the early timepoint fraction (**Fig.1C-E**). These

results are consistent with clinical studies indicating that early cancer cell dissemination

is a common occurrence in ovarian and breast cancer metastasis [11–14]. Moreover, the

same tumor cell dosing and injection schema yielded reduced frequencies of GFP+

cancer cells in ascites fluids in macrophage-depleted mice compared with WT

C57BL/6J mice, indicating lower metastatic burden (**Fig.S1G**), suggesting an expected

role of the immune system in promoting metastasis. Interestingly, despite decreased

metastases burden, amplicon DNA sequencing revealed a near identical BC.ID

distribution across all three anatomical locations in both wild type and NRG mice

(**Fig.1C-1E**), as well as chlodrosome depletion in omenta (**Fig.1C**), suggesting a

negligible role of niche-specific tumor immune microenvironment (TIME) in the overall

dominance of early arriving DTCs.

To understand if later-arriving cancer cells are dispensable for overall metastases

burden, we evaluated tumor burden in mice with a single injection of tumor cells (TP-1

only) compared to mice with fractionated dosing (TP1-6) as described in Fig.1B in WT

mice with luciferase-tagged ID8 $p53^{-/-}$ cells. At the experimental endpoint of 4 weeks

post final cancer cell injection, tumor burden levels were comparable (**Fig.1F-G**),

measured by *in vivo* bioluminescent imaging (BLI). This indicates that early DTCs are

not dependent on later DTCs for metastatic colonization and outgrowth. Given that the

disseminating ID8 $p53^{-/-}$ cancer cells are isogenic and have the same genomic

alterations in each batch, this preliminary data strongly suggests that early DTC survival

could be attributed to transcriptional adaptations facilitating increased fitness in a niche-

specific manner.

To gain a better understanding of the transcriptional programs necessary for early DTCs

to colonize and subsequently assess their clonal expansion status, we performed CITE-

seq (scRNA-seq with HTO sample multiplexing) coupled with direct capture of U6-driven

MetTag transcripts via capture sequence 1 (CS1) (**Fig. 1H**). Sequential injections were

performed in the same manner as in Fig.1B with a reverse barcode injection schema to

ensure no unintended bias towards certain barcodes gained over the cell selection

period. Single cell analysis of MetTag BC.IDs of metastasis-bearing tissues showed that

early DTCs (*TP1-2*) were enriched in overt metastases (**Fig.1H-I**), consistent with gDNA

sequencing data in Fig. 1C and 1D. Moreover, integrated analysis of ascites and

omentum single-cell datasets showed a prominent trend towards early DTC dominance,

especially in ascites (**Fig.1I-J**). Differential gene expression (DEG) analysis between

early and late timepoint cancer cells confirmed that early DTC "TP-1" cells adopted

unique transcriptional programs (**Fig. 1K**). Notably, the top enriched gene in TP-1 cells,

small nucleolar RNA host gene 18 (Snhg18), is a lncRNA with known roles in

modulating expression of genes involved in invasion and cell proliferation[15,16].

Clonal Transcriptomics Analysis of DTCs Reveals Shared Evolutionary Path

Towards Inflammatory Signature

To further investigate characteristics of early and late DTCs, we assessed whether

clonal dynamics differed between niches. Specifically, as most DTCs do not survive and

only rare clones successfully adapt to metastatic niches[17], we asked whether tumor

cells disseminated over time evolve along diverse trajectories or adopt a convergent

path leading to symptomatic overt metastases. To explore this, we performed a deeper

analysis on the cancer cell compartment, integrating clonal barcode information with

transcriptional profiles across ascites and omental niches. We first annotated cell

clusters with higher levels of keratins and absence of CD45 transcript expression

(*Krt18+*, *Krt8+*, *Ptprc-*) as the tumor cell-enriched cluster on the Seurat[18] UMAP (**Fig. 2A**

and S2A). After reclustering *Krt18+/Krt8+/Ptprc-* cells, cell cluster states were

determined through a combination of cluster marker genes and pathway-level analysis

using the Escape R package[19] (**Fig. S2B – S2E**). Ascites-derived cancer cells (AscMet)

displayed distinct transcriptional states from those in the omentum (OmMet) (**Fig. 2B-**

2D). Single Cell Gene Set Enrichment Analysis (ssGSEA) further showed that ascites-

derived cells were strongly enriched for interferon alpha (IFNα) and gamma (IFNγ)

signaling, as opposed to more heterogeneous pathways enriched in the metastatic

omentum (**Fig. 2E**). OmMet consisted of 4 distinct subclusters/modules: proliferative,

epithelial-mesenchymal transitioning (EMT), matrisome, and a hypoxic, inflammatory

subcluster (**Fig. 2D-2E**).

Cellular barcoding approaches have been widely used to reveal clonal dynamics driven

by both genetic predisposition and epigenetic changes in response to environmental

cues[20]. Integrating LARRY BC with single-cell transcriptome readouts, we asked

whether the most expanded clones would 1) cluster together based on transcriptional

status, suggesting genetically pre-determined cell-autonomous programs that drive

ovarian cancer metastasis; 2) occupy heterogeneous transcriptional states (clusters),

indicating diverse transcriptional adaptation to different metastatic niche-derived cues

(**Fig. 2F**). Interestingly, we observed minimal overlap between dominant clones across

each HTO/mouse in both anatomical locations (**Fig. 2G**) - while one clone, BC-1, was

shared across ascites and omentum, the most of highly expanded clones are distinct

between metastatic niches. These results suggest that successful clonal outgrowth

requires unique adaptations reflective of the specific microenvironment.

Given that omentum metastases showed diverse single cell transcriptional states (**Fig.**

2D-E), we projected clonal identity of the most expanded clones from the omentum onto

the cancer cell UMAP and examined similarities between clones. At both the

transcriptome cluster and pathway levels, dominant clones were enriched for several

defined signatures (**Fig. 2H-2I, S2F**). Apart from the proliferative subcluster 4, expanded

clones consistently occupied the hypoxic, inflammatory subcluster 0 (**Fig. 2I**),

suggesting diverse, yet directed evolutionary trajectories within the same clonal lineage.

To understand the lineage relationships among different subclusters, we applied RNA

velocity and pseudotime analysis using scVelo and CellRank[21]. Paga velocity and

pseudotime analyses revealed a clear trajectory originating from the "primitive" ascites

compartment subcluster 3, leading to proliferative status, then further converging to

subcluster 0, a terminal inflammatory state in the omentum (**Fig. 2J & 2K, Fig. S2G**).

Notably, IFNγ, but not the IFNα signature, was significantly enriched in early DTCs (**Fig.**

2L, S2H). Together, the integrated clonal and transcriptional analysis suggests that

successful metastatic clones occupy several niche-dependent transcriptional states.

These evolved phenotypes appear to originate from the ascites compartment, which

encompasses an IFN-signaling-rich module that may prime these later adaptations.

***In vivo* CRISPR screen identifies functional essentiality of IFNγ signaling in**

omental colonization

As ovarian cancer cells initially disseminate into the peritoneal cavity to form ascites, we

next asked whether transcriptional adaptation of cancer cells to the ascites environment

is essential for subsequent intraperitoneal seeding and successful omental metastasis.

Here, we hypothesized that the ascites gene signature adopted by early DTCs –

characterized by limited nutrient availability and low attachment – is (1) induced by initial

detachment from a primary tumor and (2) necessary for subsequent peritoneal

metastasis. To test this hypothesis, we performed differential gene expression (DEG)

analysis and over-representation pathway analysis (ORA) with the clusterProfiler R

package[22] to compare gene expression in metastatic tumor cells residing in the two

distinct niche compartments (Asc vs Om). Compared to OmMet, DEGs of AscMet were

enriched in hemoglobin genes (*Hbb-bs, Hba-a1, Hba-a2, Hbb-bt*) and interferon-

stimulated genes (*Ifi27l2a* and *Irf7*). In contrast, OmMets were enriched in immediate-

early response genes (*Atf3, Egr1, Fos, Fosb, Jun*)[23,24], adaptive immune

immunoglobulin components (*Igkc, Igha, Jchain*)[25], and angiogenesis (*Vegfa*) (**Fig. 3A**).

Next, IFNγ/IFNα module scoring and ORA analysis confirmed a dominant interferon-rich

inflammatory signature in AscMet, as opposed to more heterogeneous programs in

OmMet (**Fig. 3B, Fig. S3A & B**).

To assess the functional significance of AscMet-OmMet DEGs, we designed a custom *in*

vivo CRISPR/Cas9 screen targeting 100 DEGs identified (**Fig. 3C,** "In Vivo Ovarian

Metastatic Niche Signature Screen"). We introduced SpCas9 into ID8 p53$^{-/-}$ ovarian

cancer cells via PiggyBac (PB) transposon system and validated Cas9 expression (**Fig.**

S3C). Cas9 function was validated by assessing loss of mCherry fluorescence over 5

days, after transfection of sgRNA targeting mCherry (**Fig. S3D**). After constructing the

curated sgRNA library containing 1,030 different guides, ID8.Cas9 cells were

transduced at an MOI of 0.15. After 7 days of puromycin selection, 2.4 million cells

(~2000X representation of sgRNA library) were i.p. injected to a whole-body Cas9-

expressing host (Rosa26-Cas9 knockin on B6J, Jax 026179) (**Fig. 3C**). This host has

central tolerance to Cas9, decreasing immunogenicity towards Cas9-expressing tumor

cells[26]. Six weeks post-injection, metastasis-bearing omenta and ascites were

harvested for genomic DNA (gDNA) extraction and NGS sequencing, followed by

MAGeCK[27] analysis (**Fig. 3C**). The positive controls for the screen, *Pten* sgRNAs, were

significantly enriched in metastases derived from both anatomical locations (**Fig. 3D &**

3E). Compared to the initial injected cell pool sgRNA distribution, sgRNAs targeting

eight genes in OmMet and ten genes in AscMet were significantly depleted (*p < 0.05*)

(**Fig. 3D**). These results were further visualized by plotting the distribution of each

sgRNA for top hits (**Fig. 3E**). The top depleted (dropout) genes – *CD300lb*, *Ms4a8a*,

Marco, *Gbp2b* (OmMet), and *Slfn1* (AscMet) – were all upregulated DEGs identified in

AscMet, suggesting a necessity for transcriptome adaptation to the ascites niche - the

first niche DTCs encounter - for ultimate metastasis success. Furthermore, GSEA

analysis of dropout genes using the GO Biological Function (GOBP) gene sets revealed

a shared enrichment of apoptotic cell clearance processes in both Asc and Om

compartments and enrichments of programs related to interferon−gamma and

interleukin−1 response (**Fig. 3F**). Despite high IFNγ signaling in AscMet which was

described in Fig.2 and Fig.3B, OmMet expressed a higher level of *Ifngr1* compared to

AscMet (**Fig. S3E**), consistent with IFNγ response dependency observed in the *in vivo*

CRISPR screen (**Fig. 3F**). These data suggest that, despite the heterogeneity of

transcriptional programs revealed by scRNA-seq, the (1) shared essentiality of apoptotic

pathway regulation across ascites and omentum, and (2) IFNγ response first activated

in ascites, later necessary for omental metastasis, are both critical DTC adaptive

cellular processes that collectively drive ovarian metastases.

Loss of IFNγ-response in DTCs significantly decreases peritoneal metastases and

extends overall survival

To experimentally validate IFN-γ inducibility of screen hits in cancer cells, we treated ID8

p53[-/-] cells *in vitro* with recombinant IFN-γ and tested the inducibility of *Marco*, *Gbp2b*,

and *Slfn1* via RT-qPCR, where *Stat1* and *Irf1* transcription factors served as positive

controls for IFN-γ response[28] (**Fig. 3G**). By 24 hours of treatment, all three genes

displayed significant IFN-γ inducibility (**Fig. 3G**). *Gbp2b* (*GBP1* gene in humans) is an

interferon-inducible GTPase and part of the large guanylate-binding protein (GBP)

family with known roles in innate immune cells such as promoting antimicrobial

immunity, inflammatory response, and cell death regulation[29]. Although this protein was

shown to have both pro- and anti-tumor effects in various cancer contexts, in ovarian

cancer high Gbp2b expression mediates chemotherapy resistance[30,31]. With Gbp2b

being the most statistically significant hit in the screen, we also validated its inducibility

on the protein level via Western Blot (**Fig. 3H**).

Given that the top hits from the CRISPR screen were IFNγ-inducible, we reasoned that

IFNγ may serve as a primary regulator that exerts broader pro-metastatic adaptive

cellular programs in DTCs. Thus, instead of focusing on a single downstream gene

target, we sought to disable the entire cascade. To assess the functional role of the IFNγ

signaling pathway in tumor cells during ovarian cancer metastasis, we genetically

knocked out the IFNγ receptor 1, *Ifngr1,* in ID8 p53$^{-/-}$ cells. Successful knockout of *Ifngr1*

was validated via flow cytometry (**Fig. S3F**) and ablation of the IFN-γ transcriptional

downstream targets was validated via western blot probing for Stat1 and Gbp2b (**Fig.**

S3G). We next I.P. injected Cas9-expressing mice with either ID8.Luciferase-sgNTC or

ID8.Luciferase-sgIfngr1. Tumor burden was monitored by BLI intensity throughout the 6-

week experimental window (**Fig. 3I**). sgIfngr1 groups showed a significant reduction of

overall metastasis burden compared with sgNTC group (**Fig. 3J**). In contrast to sgNTC

group, ~80% of mice in the sgIfngr1 groups remained alive at the end of the survival

study (**Fig. 3K**). Furthermore, in human ovarian cancer microarray gene expression

data[32], patients with ovarian cancer expressing lower levels of *IFNGR1* exhibited longer

progression-free survival compared to the higher expressing group (**Fig. S3I**).

IRF1 targets mediate pro-survival and anti-proliferative programs in DTCs

We next sought to examine mechanisms by which IFNγ signaling promotes metastasis.

Ifngr1 knockout did not alter cancer cell fitness under standard adherent culture

conditions in the presence or absence of IFNγ treatment, evidenced by comparable

colony count formation (**Fig. 4A**). IFN-γ had similarly no impact on cancer cell survival

as measured by CellTiter-Glo luminescent cell viability assay under regular culture

conditions (**Fig. 4B**). Interestingly, IFNγ treatment had a protective effect and increased

cell viability in sgNTC cells under ultra-low attachment (ULA) conditions (**Fig. 4C**). This

protective effect was ablated in the *sgIfngr1* group (**Fig. 4C**). Moreover, we observed

that knockout of downstream *Stat1* and *Irf1* transcription factors (**Fig. S4A & S4B**)

phenocopied *Ifngr1* knockout under ULA and IFNγ treatment conditions (**Fig. 4D**). Irf1 is

a transcription factor downstream from Ifngr1/Stat1 signaling. As Irf1 knockout did not

rescue the survival phenotype (**Fig. 4D**), this suggests that (1) Irf1 activation is

necessary for the pro-survival phenotype and (2) the genes necessary for increased

survival under low attachment conditions are likely downstream of Irf1. To investigate

this further, we performed RNA-based gating in the scRNA-seq dataset and defined

Irf1[high] and *Irf1*[low] cancer cell populations (**Fig. 4D**). We created custom module scores

that reflected common pathways cancer cells use to resist anoikis-mediated death:

autophagy, stress, and pro-survival/anti-apoptosis[33] (**Fig. 4D**). Out of the three modules

tested, *Irf1*[high] cells most significantly upregulated the pro-survival/anti-apoptosis module

(**Fig. 4F**, right panel). One of the top upregulated genes in the *Irf1*[high] population was

Stat3 (belonging to the pro-survival module) (**Fig. 4G, Fig. S4C**). The IL-6/JAK/STAT3

signaling axis is known to promote cancer cell survival by regulating the expression of

anti-apoptotic genes[34]. Indeed, anti-apoptotic Stat3 target genes, *Mcl-1* and *Bcl2*[35],

were significantly upregulated in *Irf1*[high] cells (**Fig. 4H**). Socs3 is a known marker of

active Stat3 signaling and is responsible for dampening the JAK/STAT pathway[36]. We

confirmed *Socs3* upregulation in this population of cells (**Fig. S4D**). Next, we

experimentally tested pStat3 activation under ultra-low attachment and IFNγ treatment

conditions. As predicted based on the scRNA-seq modules (**Fig. 4F**), SQSTM1

(autophagy marker) levels were unaltered with IFNγ treatment in WT cells, while pStat3

increased (**Fig. 4I**).

Interestingly, we observed increased *Btg2* expression, an anti-proliferative gene, in

Irf1[high] cells (**Fig. S4E**). As DTCs often exhibit reduced proliferation[37], we tested *Irf1*[high]

cells for an anti-proliferative/dormancy module. The dormancy module was significantly

upregulated in *Irf1*[high] cells (**Fig. 4J**). Consistent with this observation, AscMet exhibited

reduced barcode expansion (**Fig. S4F**) and reduced *Top2a* expression (**Fig. S4G**)

compared to OmMet. Moreover, module scoring in patient TCGA-OV dataset[32] yielded

moderate positive correlations of pro-survival and dormancy programs to IFNγ response

(**Fig. S4H**). Collectively, these results suggest that, while the loss of *Ifngr1* does not

affect cancer cell growth *in vitro*, under low attachment and IFN-γ-rich conditions that

mimic ascites environment, DTCs upregulate an Irf1/Stat3-driven pro-survival module

and are phenotypically less proliferative.

Pro-inflammatory TAMs in ascites are distinct from omental TAMs

Next, we considered whether ascites priming of cancer cells occurs via autocrine

signaling or through interactions with the metastatic niche microenvironment. To identify

IFNγ producers and responders within the ovarian metastatic niche, we first examined

the expression patterns of IFNγ and IFN-responsive genes identified in our CRISPR

screen. Analysis across all cells (**Fig. 5A, S5A & S5B**) revealed that the T/NK cell

populations are the main producers of IFNγ, while the downstream CRISPR hits

(*Gbp2b*, *Marco*, *Ms4a8a*, *CD300lb*, and *Slfn1*) are mainly expressed in the myeloid

fraction, at higher levels compared to tumor cells (**Fig. 5A**). CellChat[38] analysis

identified a significant increase in both the number and strength of cell-cell interactions

between cells from naïve and metastasis-derived samples (**Fig. S5C**). Tumor cells

primarily interacted with myeloid cells in both ascites in omentum (**Fig. 5B**). Additionally,

stromal interactions were enriched in the metastatic omentum (**Fig. 5B**). These results

suggest that interaction between IFNγ-responsive tumor cells and macrophages may be

a key axis of communication in ovarian metastasis. Focusing on the myeloid

compartment, we grouped Seurat clusters into broader cell identity categories (**Fig. S5A**

& B) and found a proportional increase in myeloid cells in both metastatic sites

compared to their naïve counterparts (**Fig. 5C**). Subclustering the myeloid population

revealed that the major contribution to the proportional increase are subclusters 0,1, 5,

and 6 (**Fig. 5D, Fig. S5D**). These macrophage subclusters were identified as tumor-

associated macrophages (TAMs) based on high expression of *Adgre1*, *Cd68*, *Csf1r*,

Itgam, *Mertk*, and *Trem2* (**Fig. S5E**). RNA velocity analysis revealed a dynamic

transition from the ascites-derived TAM clusters 1 and 5 toward omental TAM clusters 0

and 6 (**Fig. 5E**). Omental clusters 0 and 6 were characterized by increased expression

of costimulatory molecules and interleukin 1 family genes as compared to ascites

clusters 1 and 5 (**Fig. 5F, Fig. S5F**). Collectively, the RNA velocity and DEG analyses

suggest that ascites-derived TAMs may represent a less differentiated or "primitive"

state relative to omental TAMs.

Hybrid functional state of omental TAMs is accompanied by an increase in

activated and regulatory T cells

Given the apparent macrophage niche-dependent differentiation states, we next asked

whether omental TAMs reflect a polarized, terminal state within the metastatic niche. If

so, increased TAM polarization would likely shape the T cell compartment[39,40]. Indeed,

while the ascites-derived T/NK compartment was dominated by a mixture of naïve

CD4+ and CD8 +T cells, omental metastases and niche tissue contained a broader and

more complex repertoire of activated, exhausted, and regulatory T cells (**Fig. 5G**)

enriched in pathways related to regulation of immune cell activation (**Fig. S5G**). We next

subsetted macrophages ($Cd68^{high}$/$Adgre1^{high}$, **Fig. S5H**) and scored them against M1

and M2 gene modules. Compared to ascites-derived TAMs, omental TAMs exhibited

higher hybrid functional state scores (**Fig. S5I**), reflecting an immunomodulatory TAM

cellular state[41] that cannot be fully captured by binning into a strictly immune-promoting

or suppressing category. Unlike the immunomodulatory and ECM-related omental

signatures, ascites-derived TAMs were enriched in pathways associated with cellular

stress responses and NRF2-dependent redox metabolism (**Fig. 5H & 5I**). On the gene

level, ascites TAM signatures were reflected in elevated expression of key stress-related

genes – including *Anxa1* and *Hmox1* – as well as *Ndufb8* and *Cox5a*, components of

mitochondrial respiration (**Fig. S5J**). These findings indicate that ascites TAMs may

represent an earlier, less polarized yet metabolically active state, while omental TAMs

have transitioned into an immunomodulatory, polarized phenotype.

To characterize prevalent interactions between omental TAMs and T/NK cells compared

to ascites, we applied NicheNetR, a computational tool used to predict receptor-ligand

interactions[42] (**Fig. S6A**). NicheNetR comparison between the two anatomical locations

revealed *Cd80* interacting with *Cd28*, *Cd274* (PD-L1 transcript), and *Ctla4* as prevalent

interactions between myeloid cells and T/NK cells in the metastatic omentum (**Fig.**

S6B). *Cd80* was strongly expressed and upregulated in omental compared to ascites

TAMs (**Fig. S6C**). Cd80 engagement with Cd28 and Ctla4 are necessary interactions for

co-stimulation and co-inhibition of T cell activation, respectively[43]. Cd80 interaction with

Cd274 is a less characterized interaction that can occur bidirectionally and leads to T

cell inhibition, similar to the PD-1/PD-L1 signaling axis[44]. Next, we utilized flow

cytometry to validate immune cell dynamics observed in the scRNA-seq dataset (**Fig.**

S6D-H). GFP+ cancer cells were detected in both metastases-bearing ascites and

omentum samples (**Fig. S6D**). While there was a significant increase in

CD11b+/CD11c+ macrophages as well as CD11b-/CD11c+ DCs in ascites (**Fig.S6E &**

F), metastatic omentum samples were significantly enriched in mature F4/80+ TAMs

(**Fig. S6G**) and cytotoxic CD8+ T cells (**Fig. S6H**). These findings further implicate the

omental environment in fostering a highly immunomodulatory population of TAMs.

Peritoneal macrophages protect cancer cells against anoikis-mediated cell death

Although TAM-induced immunomodulation is a well-established, significant metastasis

driver, less is known about the role of direct TAM-tumor cell interactions. To identify cells

that are in close contact with ovarian metastases, we developed a Cre-based iteration

of the "NicheTracing" system to genetically mark and trace microenvironmental contacts

in vivo over a prolonged timeframe. In this system, cancer cells expressing Cre.ERT2

were injected into Ai6 (RCL-zsGreen) Cre reporter mice. Upon cytoplasmic and nuclear

entry of Cre into neighboring cells, Cre-mediated recombination turns on the zsGreen

cassette in host cells (**Fig. 6A**). After validating Cre.ERT2 expression in transduced cell

lines (**Fig. S7A**), we benchmarked this system using *in vitro* coculture (**Fig. S7B-S7F**)

and in an independent model of triple-negative breast cancer (TNBC) lung metastasis

established using the E0771 cell line (**Fig. S7G & S7H**). In the ovarian metastasis

model, we readily detected zsGreen+ (ZsG) niche cells in both metastasis-bearing

omentum and ascites at the experimental endpoint six weeks post cancer cell injection

(**Fig. 6A**). We performed flow cytometry analysis on ascites and gated cells into ZsG^low

and ZsG^high (**Fig. 6B**, top panel). Unlike ZsG cells detected in lung metastases (**Fig.**

S7H, right panel), over 90% of ZsG^high host cells were F4/80$^+$/Ly6G^{low} TAMs (**Fig. 6B,**

bottom panel, **Fig. 6D**).

We next asked if peritoneal TAMs actively promote ovarian cancer metastasis.

Macrophage depletion via direct i.p. liposomal chlodronate injections (**Fig. 6E**)

demonstrated that peritoneal TAMs are pro-metastatic: depleting TAMs significantly

reduced omentum weights (**Fig. 6F, Fig. S7I**) and CD45 negative cancer cell

frequencies in ascites at experimental endpoint (**Fig. 6G, Fig. S8A** flow gating strategy).

We next explored whether macrophages could promote cancer cell survival under low

attachment conditions, akin to the effects observed with IFNγ treatment (**Fig. 4F**). To

test this, we performed a macrophage-cancer cell coculture under ultra-low attachment

(ULA) conditions. Macrophages from peritoneal lavage were enriched using F4/80

magnetic beads (**Fig. 6H**, **Fig. S8B & S8C** flow gating strategy) and cocultured with

three different murine ovarian cancer cell lines; ID8 p53$^{-/-}$, ID8 p53$^{-/-}$ BRCA2$^{-/-}$, and

PPNM[45] (**Fig. 6I-K, S8D** flow gating strategy). After a 48-hour coculture period, flow

cytometry analysis revealed that cancer cell viability, as measured by live/dead staining

(ZombieAqua), decreased under ULA conditions compared to regular attachment

plates. Most importantly, survival was partially rescued with macrophage coculture

across all three cell lines (**Fig. 6I-K**). These results indicate that peritoneal macrophages

directly support cancer cell survival under conditions mimicking the low attachment,

ascites environment.

Patient ascites-derived TAM-Tumor interactions are enriched in pro-survival

pathways

To determine if tumor-supportive macrophage interactions occur in human ovarian

cancer, we analyzed a human-derived single-cell dataset containing over 200,000 single

cells from 14 ovarian cancer patients and collected from five locations: peripheral blood

(PB), peripheral lymph nodes (PLN), malignant ascites (Asc), primary tumor (Pri.OT),

and matched metastatic tumor (Met.Ome)[46]. As observed in our murine dataset, cancer

cell-containing compartments, including Asc/Pri.OT/Met.Ome, had increased

proportions of macrophages compared to PB and PLN (**Fig. 7A, S9A & S9B**). DEG

analysis between TAMs at metastatic sites (Fig. 7B) and subsequent GO pathway

analysis conducted on the DEG list revealed immune-modulatory pathways (cytokine,

chemokine, allograft rejection), EMT, TNFα, and KRAS signaling upregulated in

Met.Ome TAMs (**Fig. 7D, S9D**). In Asc TAMs, the pathways were predominantly

metabolism and cargo receptor activity related (**Fig. 7E, S9E**).

We applied NicheNetR analysis to further characterize TAM-to-tumor cell interactions in

ascites. We implemented this strategy to both our murine dataset as well as the patient

dataset (**Fig. 7F**). After defining myeloid cells (Broad cell ID) as the sender cell type in

the murine dataset and macrophage cluster as the sender in the human dataset, we

compared the top predicted tumor-interacting ligands in each dataset. We found 4

overlapping sender ligands across species (**Fig. 7F**) - *TGFB1*, *IL10*, *TNF, and IL1B* -

that were the most significant ligands predicted to interact with cancer cell receptors

(**Fig. 7G**). Next, we examined the top predicted downstream target genes in cancer

cells affected by top TAM ligands (**Fig. S9F**). ORA pathway analysis of the predicted

downstream genes yielded several pathways involved in cell-cell adhesion, response to

oxygen levels, and other tumor-promoting pathways (**Fig. 7I**). Together, these data

suggest that patient-derived Asc TAMs may function similarly to murine TAMs in directly

supporting ascites cancer cells.

DISCUSSION

Early DTC dominance phenotype in ovarian cancer metastasis

DTC-niche interactions at each stage of the metastatic cascade fundamentally

determine disease progression and patient outcomes. Following the initial wave of

metastatic dissemination, early DTCs inevitably reshape the metastatic niche

microenvironment, irrespective of whether they ultimately survive or perish upon arrival.

The niche remodeling by pioneer DTCs likely exerts more profound effects on

subsequent colonization than distant signals from the primary tumor. While the pre-

metastatic niche (PMN) has been extensively characterized[47,48] , this dynamic process

following DTC seeding creates what we term the Post-Seeding Metastatic Niche

(PSMN). The multi-cellular dynamics resulting from progressive DTC accumulation at

newly colonized PSMN mirrors a well-established ecological concept: the Allee effect,

which describes how population growth rates depend on population density through

competition or cooperation. Although the Allee effect has been demonstrated in

metastatic contexts using in vitro models[49], how continuously arriving DTCs dynamically

alter PSMN properties and influence the fate of later-arriving tumor cells within the intact

physiology of living systems remains unstudied.

Leveraging our MetTag barcoding and staggered injection model, the data presented

here consistently demonstrated a skewed overrepresentation of early disseminated

tumor cells (DTCs): TP-1 and TP-2 DTCs overwhelmingly dominate the metastatic

burden at day 40, whereas cells introduced at later days (TP-6) showed minimal

contributions to final overt metastases. This pattern persists across immunocompetent,

immunodeficient (NRG), and macrophage-depleted hosts (**Fig. 1**). This skewness in

early DTC dominance cannot simply be attributed to inherent, longer chronological

growth times of early DTCs, as there remains an ample growth window for TP-6 cells

(from Day 11 to Day 40). We put forward several non-mutually exclusive hypotheses

that could explain early DTC dominance in our study. First, the omentum might contain

a limited number of permissive micro-niches that favor DTC colonization. Pioneer DTCs

might rapidly saturate these micro-niches and monopolize spatial resources – a "first

come, first served" model. Second, the immune-independent nature of skewed

distribution of temporally disseminated DTCs suggests a role of non-immune stromal

elements in dictating colonization success. Early DTCs may actively remodel the niche,

rendering it unfavorable for later DTCs. Comprehensive spatiotemporally resolved

imaging and genomics analyses are required to further explore these hypotheses.

Duality of IFNγ signaling-driven metastatic adaptation

Beyond the "seed and soil" hypothesis, it has been increasingly appreciated that

metastasis is not merely passive fitness-based selection of DTC clones at the niche, but

also an active epigenetic DTC-niche mutual adaptation process. Our single-cell

transcriptome analysis, coupled with barcoding clonal analysis, provided us with clonal

resolution of the underlying genetic and epigenetic determinants governing DTC

behavior. Interestingly, instead of occupying unique transcriptome clusters - which

would likely suggest distinctive genetic mutational features - the expanded clones form

similar coordinated adaptive "mini-ecosystems" encompassing multifaceted

transcriptome features that converge toward an inflammatory/IFNγ response (**Fig. 2H-**

L). Our CRISPR/Cas9 *in vivo* screen results (**Fig. 3**) further supported the notion that

cancer cell fitness in the peritoneal cavity and subsequent omentum metastasis

depends on rapid transcriptional-level adaptation to activate IFNγ-responsive programs

in ascites. Our findings extend the conventional view of genetically pre-determined

clonal fitness in the metastatic cascade. In this "first come, first served" model, early

metastatic niche sensing and epigenetic adaptation in the race towards occupying

permissive micro-niches ultimately determines metastasis success.

Although the findings presented here support a tumor cell-centric pro-metastatic role of

IFNγ, it is important to note that IFNγ in metastatic niches can act in complex ways. In

some cancers, IFNγ reduces metastasis through activation of effector T and NK cells,

thereby enhancing the effectiveness of immunotherapy[50,51]. Furthermore, in some

contexts, IFNγ can induce apoptosis in regulatory T cells and inhibit angiogenesis[52].

Specifically in ovarian cancer cells, both anti-metastatic and pro-metastatic roles have

been reported[53,54]. For the latter, IFNγ has been previously reported to upregulate PD-

L1 and IL-8 in ovarian cancer cells, leading to increased cell migration and

proliferation[54]. We did not observe any proliferation differences via colony formation

assay *in vitro* (**Fig. S4F**) after a 7-day IFNγ exposure, and our scRNA-seq data suggest

that IFNγ programs are associated with dormancy *in vivo* (**Fig. 4J & S4**). Based on the

data presented here, we believe that, although both effector immune cells and cancer

cells could be impacted by IFNγ, IFNγ response by cancer cells (intrinsically) leads to an

overall pro-metastatic effect that outweighs the anti-metastatic roles in ovarian cancer.

In particular, although both ascites and omental TAMs in our study closely resembled a

Trem2/Apoe-high state described elsewhere[55], we found that they exhibited niche-

specific adaptations promoting divergent functions (**Fig. 5**).

In summary, our study dissected the complex landscape of ovarian cancer metastasis through a systematic characterization of both cancer cells and local immune factors using MetTag temporal, clonal and scRNA-seq analysis. Our findings highlight the importance of ascites as a functional reservoir for these factors and the early acquisition of IFNγ response as a necessary adaptation step for DTCs to survive and colonize at the omentum.

ACKNOWLEDGMENTS

This study was partially funded by the NIH T32GM075762 training grant (E. A.), NIH R01 CA194697-01(S.Z.), NIH R01 CA255064-01A1(S.Z.), NIH R21 CA263798-01 (S.Z.), and the Cancer Prevention and Research Institute of Texas (CPRIT) Scholar Award RR220024 (S.Z.). We are additionally grateful to the Dee Family endowment (S.Z.) and to the following core facilities: Notre Dame Freimann Life Sciences Center, Notre Dame Genomics and Bioinformatics Core Facility, Indiana University School of Medicine Center for Medical Genomics, Indiana University Simon Cancer Center, Simmons Comprehensive Cancer Center, UT Southwestern (UTSW) Animal Resource Center, UTSW Flow Cytometry Core, and UTSW Small Animal Imaging Core Facility. The computational arm of this research project was, in part, supported by the BioHPC supercomputing facility at UTSW.

AUTHOR CONTRIBUTIONS

Conceptualization and experiment design, E.A., M.S.S. and S.Z.; Data analysis, E.A., S.Z., X.L., Z.Z.; Experiments and methodology, E.A., S.Z., S.R.F., S.M.G., J.L., W.M., N.D., T.C.R., M.Z.; Manuscript writing and revision, E.A., S.M.G., and S.Z.; Study supervision, S.Z. We attest that all coauthors reviewed and pre-approved the manuscript before submission.

DECLARATION OF INTERESTS

We do not have any competing interests to declare.

FIGURE LEGENDS

Figure 1: Tracking Clonal Lineage in Ovarian Cancer Metastasis

(**A**) Schematic depicting main elements of the MetTag library barcoding construct. (**B**) Schematic of injection and tissue collection timeline for the staggered injection experimental model using MetTag-barcoded ID8 p53$^{-/-}$ ovarian cell lines. Schematic was partially created with BioRender.com. (**C**) Genomic DNA (gDNA)-level barcode distributions in the omentum graphed as frequency under WT, NRG, and MAC depletion conditions. (**D-E**) gDNA-level barcode distributions in ascites and pelvic fat pads under WT and NRG conditions. In (C-E) panels, BC.ID1 = TP-1, BC.ID2 = TP-2, BC.ID3 = TP-3, BC.ID4 = TP-4, BC.ID5 = TP-5, BC.ID6 = TP-6 for clarity. (**F**) Injection schema for comparing one-time (400,000 total, n = 10) cell dose and fractionated (2,400,000 total, n = 10) cell injections (left) and representative BLI images at endpoint (right). (**G**) Bar graph depicting total flux emission in each group at experimental endpoint. (**H**) Schematic of scRNA-seq experiment conducted using a reversed BC.ID injection order. LARRY barcodes are readily captured by Chromium 10x Feature Barcoding. (**I**) Pie charts of RNA-level barcode frequencies in each location grouped as early (BC.ID5, BC.ID6), mid (BC.ID4, BC.ID3), and late (BC.ID2, BC.ID1). (**J**) Heatmap of sample level barcode frequencies, BC.ID6 = TP-1, BC.ID5 = TP-2, BC.ID4 = TP-3, BC.ID3 = TP-4, BC.ID2 = TP-5, BC.ID1 = TP-6. (**K**) Volcano plot of DEGs between early and late timepoint barcoded cancer cells where threshold for significance was defined at $p < 0.05$ and log2FC < -1.0 for downregulation or >1.0 for upregulation. Genes labeled are the top 30 genes based on significance. Data in Figure 1G was analyzed using a two-tailed Student's t test and graphed as mean with standard deviation (SD). In Figure 1K, the DEG table was generated using the Seurat FindMarkers() function and p values calculated via a Wilcoxon rank-sum test. For all Wilcoxon rank-sum tests, p values graphed represent the adjusted p value.

Figure 2: Clonal Analysis Reveals Shared Evolutionary Path Towards Inflammatory Signature

(**A**) Split UMAP depicting cluster distributions across the four experimental conditions and identifying cancer cell-enriched clusters in ascites and omentum. (**B**) Reclustered cancer cell UMAP colored by the two metastasis-bearing anatomical locations. (**C**) Stacked bar chart depicting the frequency of Seurat subclusters within each metastatic location. (**D**) Seurat cluster UMAP of reclustered cancer cells with transcriptional program labels. (**E**) Dot plot depicting Hallmark pathways across all subclusters used to define each cluster subset gene program in (D). (**F**) Schematic representation of the possible outcomes of LARRY clonal analysis based on transcriptome patterns of enriched clones. (**G**) Heatmap of normalized clonal LARRY barcode counts across the two anatomical locations and individual mice (HTOs). (**H**) UMAP overlay of top 3 expanded clones in the omentum. (**I**) Bar chart depicting the relative subcluster proportions present in each of the top 3 barcode clones. (**J**) scVelo-generated PAGA velocity graph indicating trajectory arrows and inferred cluster lineage. (**K**) scVelo and CellRank-based simulated path from start to endpoint clusters. (**L**) Violin plot depicting RNA-level contribution of each timepoint to the IFNγ signature. Data in Figure 2L was analyzed by first defining the IFNγ signature using canonical mouse genes, followed by directly applying the wilcox.test() function in Seurat to compare TP-1 and TP-6 expression.

Figure 3: Activation of IFNγ Response in Ascites is Necessary for Successful Omental Colonization

(**A**) DEG-based volcano plot of gene expression differences between OmMet and AscMet, where threshold for significance was defined at $p < 0.05$ and log2FC < -1.0 for downregulation or >1.0 for upregulation. (**B**) Boxplots comparing IFN signature modules between AscMet and OmMet. (**C**) Schematic describing *in vivo* ovarian metastatic niche signature screen. Partially created with BioRender.com. (**D**) *In vivo* CRISPR screen RankView plots generated using MAGeCKFlute. OmMet (n=5) and AscMet (n=4)

samples were compared to the original pool of ID8 p53$^{-/-}$ cells. (**E**) Ranks of individual

sgRNAs targeting positive control *Pten* and depleted genes. (**F**) Lollipop plot of depleted

pathways associated with depleted genes in Fig. 3E. (**G**) IFNγ treatment time course

qPCR plots of select CRISPR hits and positive controls. (**H**) IFNγ treatment time course

western blot probing for Stat1/pStat1 and GBP1. (**I**) Schematic of *in vivo* experimental

metastasis study comparing sgNTC and sgIFNGR1.g1 groups. (**J**) Time course log10-

transformed total emission/flux values (left). Two independent groups pooled; cohort 1

NTC n = 8, KO n = 7; cohort 2 NTC n = 5, KO n = 5. log10-transformed flux plot for

week six with p value (right). (**K**) Kaplan Meier survival curve generated for cohort 1.

Data in Figure 3B was analyzed by first defining IFN signatures, followed by applying

wilcox.test() in Seurat to compare AscMet and OmMet. CRISPR screen related data

(3D-F) was first analyzed using MAGeCK. Count summary, gRNA summary, and gene

summary output files were loaded into RStudio to generate rankview plots using

MAGeCKFlute. For Figure 3G, qPCR data was analyzed by averaging technical

replicates for each biological replicate, followed by calculating dCq by normalizing target

gene Cq to housekeeping gene control. Next, 2^(-ddCq) or fold change was calculated

by normalizing the values to the zero hour (PBS) treatment samples and plotted in

Prism GraphPad. Significance between groups was assessed using a two-tailed

Student's t test and data was plotted as mean with standard deviation (SD). Data in

figure 3J left panel was plotted as log transformed mean with 95% confidence interval

(CI) and a two-tailed Student's t test was performed at each timepoint. Data in 3J right

panel was plotted with median log transformed values.

Figure 4: IFN-γ Sensing Mediates Resistance to Anoikis and Activates Pro-

survival Signaling

(**A**) Colony formation assay performed without (left) or with (right) 100 ng/ml IFNγ. (**B-C**)

Cell-Titer Glo assay luminescence CTG values graphed as fold change for regular

culture conditions (B) and ultra-low attachment conditions (C), with and without IFNγ

treatment. Fold change was determined by normalizing each cell line's CTG value to the

regular attachment, no IFNγ treatment condition. (**D**) CTG flux value dot plots for Stat1-

KO and Irf1-KO cells under ultra-low attachment, with or without IFN-γ treatment. (**E**)

scRNA-seq based gating strategy used to identify Irf1-high and low cells for module

scoring. (**F**) Violin plots for autophagy, stress, and pro-survival (anti-apoptotic) scores in

Irf1-high vs. Irf1-low cells. Violin plots comparing Stat3 (**G**), Mcl-1, and Bcl2 (**H**)

transcript levels between Irf1-high, mid, and low cells. (**I**) Western blot probing for Stat3

activation and SQSTM1 expression under ultra-low attachment conditions, with or

without IFNγ treatments. (**J**) Violin plot of dormancy module comparing Irf1-high and low

cells. CTG assay luminescence data (4B-4D) was analyzed in Prism using student's t

test. Data in B and C was graphed as mean and standard deviation, data in D was

graphed as median values. Anoikis and dormancy modules, along with Stat3, Mcl-1,

and Bcl2 transcript expression levels were compared between Irf1-high and Irf1-low

cells by applying the wilcox.test() function in Seurat.

Figure 5: Ascites and Omental TAMs Differ in Their Immunomodulatory Potential

(**A**) Violin plots depicting CRISPR screen hits expression level across Broad cell ID

categories (cell types). (**B**) CellChat-generated circus plots of interaction strengths

between Broad cell ID categories in metastatic omenta (left) and ascites (right). (**C**)

Stacked bar chart of cell type proportions across the four experimental groups. (**D**)

UMAPs of reclustered myeloid cells, split by experimental condition. (**E**) scVelo RNA

velocity plot showing arrows pointing from ascites TAMs (clusters 1 and 5) to omentum

TAMs (clusters 0 and 6). (**F**) DEG volcano plot comparing gene expression differences

between ascites TAM cluster 1 and omentum TAM cluster 0. (**G**) Stacked bar chart of

subsetted T/NK cells comparing cell proportions in ascites and omentum. (**H-I**) ORA

generated Reactome pathways enriched in ascites (left) and omentum-derived TAMs

(right). Significance of DEGs in Figure 5F was evaluated by applying the FindMarkers()

and Wilcoxon test in Seurat to compare ascites and omental *Cd68*^{high}/*Adgre1*^{high}

macrophages.

Figure 6: NicheTracing Identifies Ascites-derived TAMs as Pro-Metastatic
Interactors

(**A**) Schematic describing the Cre-based NicheTracing labeling system where ID8 p53$^{-/-}$
and E0771 cancer cells express Cre.ERT2 which transfers to neighboring RCL-ZsGreen
niche cells. Upon nuclear entry of Cre.ERT2, labeled niche cells turn ZsGreen positive
(ZsG+). (**B**) Representative gating strategy depicting the major cell population tagged
with zsG. (**C-D**) Bar plots showing preference for F4/80+ macrophage labeling in
ascites. (**E**) Liposomal chlodronate-based macrophage depletion strategy and timeline
schematic with representative tissue images. n= 7 liposome control group and n = 7
chlodrosome treatment group. (**F**) Bar plot showing quantification of omental
metastases via omental weight measurement at endpoint. (**G**) Images of ascites
samples harvested at endpoint for flow cytometry analysis (left) and flow-based
quantification of CD45 negative cells (right). (**H**) Schematic of macrophage-cancer cell
coculture layout (top) and representative coculture images for one cell line (bottom). (**I-**
K) Plots of percent CD45 negative cancer cell viability post coculture. Data in Figures
6C-D and 6F-6H was plotted as mean with SD and differences between groups were
compared via a two-tailed unpaired Student's t test. Data in Figures 6I-K was similarly
analyzed using a Student's t test and plotted with median values.

Figure 7: Patient Ascites MAC/Tumor Interactions are Enriched for Tumor-
supporting Pathways

(**A**) Stacked bar chart of main cell type proportions captured in human scRNA-seq
dataset across five anatomical locations, using maintype-2 annotation provided by
publisher. (**B**) Heatmap depicting DEGs between Met.Ome and Ascites. (**C-D**) GO
Molecular Function pathway level differences between omental MACs (C) and ascites
MACs (D). (**E**) Schematic depicting NicheNetR analysis approach for finding top TAM

ligands in both murine and human datasets. This panel was partially generated using

BioRender.com. (**F**) Venn diagram of top NicheNet predicted ligands after filtering for

robust expression and overlapping with top interactions in our murine dataset. (**G**)

Heatmap showing top receptor-ligand interactions upregulated in patient ascites-MACs

compared to metastases. (**H**) GO enrichment dotplot generated from the gene list

containing top target genes in cancer cells that are predicted to respond to top ligands

in the human dataset. Data in Figure 7B was analyzed using the FindMarkers() function

in Seurat and Wilcoxon test to compare Asc and Met.Ome.

SUPPLEMENTAL INFORMATION (Including Supplemental Figures S1-S9)

Figure S1. MetTag Tracing Construct and Metastatic Model Validation

(**A**) Schematic depicting MetTag library barcoding plasmid, and modes of detection

upon plasmid integration: DNA level, Pol III U6 driven RNA level, and Pol II CROP-seq

based. (**B-C**) Barcode frequencies comparing relative eaviness of library distribution

across BC.ID batches (B) and original depositor LARRY library (C). (**D**) Table depicting

unique reads detected in each library batch after MiSeq sequencing and read filtering.

(E-F) Representative ascites images and dissecting scope images of GFP+ metastases

formed under the repeated injection schema. (F) Bar chart depicting macrophage

depletion in chlodrosome treatment group at experimental endpoint (right). (**G**)

Representative gating strategy (left) and quantification of GFP+ tumor cells in ascites

under macrophage depletion conditions (right). Data in Figures S1F and S1G was

plotted as mean with SD and differences between groups were compared via a two-

tailed unpaired Student's t test.

Figure S2. scRNA-seq-based Tumor Cell Identification and Cancer Cell Velocity

(**A**) Feature plot of main marker genes used to define and subset cancer cells. (**B**)

Heatmap of reclustered cancer cell marker genes. (**C**) Violin plots of top defining cluster

marker genes. (**D**) Escape generated Hallmark pathways each represented by a

different color. (**E**) Escape pathway-based violin plots of each pathway class in (D). (**F**)

UMAPs depicting main pathway signatures represented in the top three clonally

expanded barcoded cells. (**G**) CellRank-generated initial and terminal cellular states

overlayed on the cancer cell UMAP. (**H**). Violin plot of IFNα signature score across the

six different BC.ID timepoints. Data in Figure S2H was analyzed by first defining the

IFNα signature using Hallmark IFNα response genes, followed by directly applying the

wilcox.test() function in Seurat to compare TP-1 and TP-6 signature expression.

Figure S3. *In vivo* CRISPR Screening and IFN Signaling in AscMet and OmMet

(**A**) ORA-generated Hallmark pathways based on DEGs between AscMet and OmMet.

(**B**) Heatmap depicting IFN response gene scoring differences between AscMet and

OmMet. (**C**) Western blot validation of SpCas9 expression in ID8 p53$^{-/-}$ cells after

PiggyBac transposon transfection and selection with hygromycin. (**D**) Representative

flow cytometry generated histogram of mCherry fluorescence (left) and bar chart

depicting percent mCherry positive cells five days after lenti-guide transduction (right).

(**E**) Violin plot of *Ifngr1* expression levels in cancer cells *in vivo*. (**F**) Representative flow

cytometry generated histogram of IFNGR1 surface expression in knockout vs. wild type

cells (left) and bar chart with quantification of IFNGR1 mean fluorescent signal changes

(right). (**G**). Western blot probing for IFNγ downstream transcriptional targets, Stat1 and

Gbp2b in *Ifingr1* knockout cells. (**H**) Endpoint (week six) BLI images of sgIFNGR1

compared to sgNTC experiment cohort 1 mice. (**I**) KMPlotter-generated Kaplan Meier

progression free survival curve comparing low and high IFNGR1 expression at stage III

and stage IV. IFNα/IFNγ signatures in Fig. S3B were described in previous figures. Data

in (D) and (F) was analyzed in GraphPad Prism using a two-tailed Student's t-test and

graphed as mean with SD. Fig. S3E violin plot Ifngr1 expression levels were compared

using the wilcox.test() function in Seurat.

Figure S4. IFNγ-induced Pro-Survival and Dormancy Programs

(**A-B**) Western blots showing protein levels of Stat1, Irf1, and Gbp2b under Stat1 (A) or Irf1 (B) knockout, with IFNγ treatment. (**C**) Volcano plot of DEGs between Irf1-high and Irf1-low cancer cells. (**D-E**) Violin plots showing Stat3-related *Socs3* and *Btg2* transcript expression level differences between Irf1-high, mid, and low cancer cells. (**F**) Stacked bar plot binning AscMet and OmMet cancer cells into frequency categories based on LARRY barcode expansion score: freq. 1 = no clonal expansion, freq. 2-5 = moderate clonal expansion, and freq. 5+ = high clonal expansion. (**G**) Violin plot comparing Top2a expression levels between AscMet and OmMet. (**H**) Correlation plots generated from the TCGA-OV dataset; top panel represents the correlation between pro-survival and IFNγ modules, bottom panel represents the correlation between dormancy and IFNγ modules. DEG analysis in (C) was performed using the FindMarkers() function in Seurat. p-values in (D-E) and (G) were obtained from a wilcox.test(). For Fig. S4H, Pearson correlation coefficients and p-values were obtained after ssGSEA scoring. The correlation curves are plotted with a linear regression line with shaded 95% CI intervals.

Figure S5. Ascites and Omentum Immune Cell Characterization

(**A**) UMAP depicting all cells captured across four sequenced groups. (**B**) "Broad_cell.ID" classification of Seurat clusters into cell types based on cluster marker genes. (**C**) CellChat generated bar charts depicting number of interactions (left) and interaction strength (right) across the four sample groups. (**D**) Heatmap of myeloid TAM subclusters 1, 0, 5, and 6. (**E**) Violin plots showing gene expression of canonical TAM markers across myeloid subclusters. (**F**) Volcano plot of DEGs between myeloid cluster 5 and 6. (**G**) Dotplot of GO Biological Process pathways upregulated in omentum-derived T/NK cells based on the DEG list comparing to ascites-derived T/NK cells. (**H**) RNA-based gating strategy to subset out macrophages for subsequent analysis. (I) Biaxial scatter plot of M1 and M2 module scores with overlapped ascites and omental TAMs. (J) Violin plots comparing individual genes belonging to cellular response to

stress (Anxa1, Hmox1) and cellular respiration (Ndufb8, Cox5a) pathways between

omental and ascites TAMs. In (F), DEG analysis was performed using the FindMarkers()

function in Seurat and significance determined via Wilcoxon test. Fig. S4I hybrid score

and S4J violin plot expression levels were compared using the wilcox.test() function in

Seurat.

Figure S6. NicheNetR Receptor-Ligand Interaction and Immunotyping Analyses

(**A**) Schematic depicting cell types of interest for NicheNetR comparison between

omentum and ascites: receptors on T/NK cells and ligands secreted by myeloid cells.

(**B**) NicheNetR-generated heatmap of predicted receptor-ligand interactions. (**C**) Violin

plot comparing *Cd80* co-stimulatory molecule expression differences between

metastasis bearing omenta and ascites. (**D-H**) Flow cytometry analysis plots of cancer

cell and immune cell frequencies in naïve and metastatic settings: (**D**) Frequency of

GFP+ cancer cells, (**E**) Frequency of CD11b+/CD11c+ double positive myeloids, (**F**)

Frequency of CD11b-/CD11c+ dendritic cells, (**G**) Frequency of CD11b+/F4/80+

macrophages, and (**H**) Frequency of CD8+ T cells. S6C violin plot *Cd80* expression

levels were compared using the wilcox.test() function in Seurat. Flow cytometry data in

(D-H) was analyzed via two-tailed unpaired Student's t test and plotted with median

values.

Figure S7. NicheTracing In Vitro/In Vivo Validation and Macrophage Depletion

(**A**) Western blot validation of Cre expression in target cell lines: E0771 murine TNBC

line and ID8 p53$^{-/-}$ ovarian cancer cell line. (**B**) Schematic depicting Cre reporter used to

transfect HEK293T cells and Cre constructs expressed by target cells used for *in vitro*

assays. (**C-F**) Images generated after 72 hours of coculture with or without 2 uM 4-OHT

treatment. (**G**) Representative dissecting microscope images of negative control spleen

and lugs collected 2 weeks post E0771 retro-orbital injection. (**H**) Flow cytometry bar

plots at experimental endpoint showing percent switched immune cells (ZsG+) with or

without tamoxifen treatment (left) and within the tamoxifen treated group, frequency of

ZsG+ macrophages compared to ZsG- (right). (**I**) Metastasis-bearing omentum images

at experimental endpoint of chlodrosome-based macrophage depleted experiment. Flow

cytometry data in (H) was analyzed using a two-tailed unpaired Student's t test and

plotted as mean with SD.

Figure S8. Flow Cytometry Gating Strategies, Related to Main Figure 6

(**A**) Flow gating strategy for macrophage depletion study (left) and representative gates

for F4/80 macrophages in FMO, liposome control, and chlodrosome treated groups

(right). (**B**) Flow gating strategy for assessing MHC-II+/CD11b+ macrophage enrichment

following F4/80 magnetic bead isolation from peritoneal cavity. (**C**) Bar plot showing

relative enrichment of macrophage populations before the start of coculture. (**D**) Flow

gating strategy to assess CD45- cancer cell viability (Zombie Aqua negative

populations) after co-culture completion.

Figure S9. Ovarian Cancer Patient Macrophage and NicheNet Analysis

(**A**) UMAP depicting all (>200,000) single cells collected from five locations and

annotated by cell type ("Maintypes-2"). (**B**) Split UMAP of subsetted macrophage

clusters, split by anatomical location. (**C**) Bubble plot showing expression of top

NicheNet-predicted ligands in each cell type. (**D-E**) Dot plots depicting top Hallmark

pathways expressed in patient omental TAMs (D) and ascites TAMs (E). (**F**) NicheNet

heatmap depicting top ligands based on regulatory potential and downstream genes

affected by corresponding ligands.

METHODS

Experimental Models

Mice

All animal experiments were conducted according to IACUC protocols which were pre-approved by the University of Notre Dame and UT Southwestern Medical Center IACUC committees. Mice were housed with unlimited access to food and water in a temperature-controlled holding room with a standard 12-hour light/dark cycle. Health checks were performed daily during *in vivo* experiments. Wild type C57BL/6J (B6) mice were 8-10 weeks old at the experiment onset. Control and treated transgenic mice were age-matched, greater than 8 weeks of age. All experiments were carried out exclusively on female mice.

C57BL/6J (B6, 000664), NOD.Cg-$Rag1^{tm1Mom}Il2rg^{tm1Wjl}$/SzJ (NRG, 007799), B6.Cg-$Gt(ROSA)^{26Sortm6(CAG-ZsGreen1)Hze}$/J (Ai6, 007906), and B6J.129(Cg)-$Gt(ROSA)26Sor^{tm1.1(CAG-cas9*,-EGFP)Fezh}$/J (Cas9 026179) mice were obtained from Jackson Laboratory (Ben Harbor, ME).

Cell Lines

The TNBC E0771 cell line (CH3 Biosystems, Amherst, NY; Cat.No. 940001-Vial) was maintained in RPMI-1640 media supplemented with 10%FBS and 1% Penicilin/Streptomycin (Pen/Strep). ID8 p53$^{-/-}$, ID8 p53$^{-/-}$ BRCA2$^{-/-}$, and PPNM ovarian cancer cell lines, provided by the Stack lab at the University of Notre Dame, were maintained in DMEM low glucose media with 5% FBS, 1% Pen/Strep, and 1% Insulin-Transferrin-Selenium (ITS). PPNM cell media was additionally supplemented with 100 ng/ml Cholera toxin and 2ng/mL epidermal growth factor (EGF). HEK293T cells were cultured in DMEM high glucose supplemented with 10% FBS, 1% Pen/Strep, and 1% GlutaMAX (Gibco, 35050061). Cell cultures were grown at 37 ºC, 5% CO_2, and 95% humidity. All cell lines were tested and were negative for mycoplasma contamination using a routine mycoplasma PCR kit (ABM).

Establishment and Monitoring of Experimental Ovarian Metastasis

For the single-injection OvCa model, 4x10^5 ID8-LARRY.GFP or ID8-Luciferase were injected into the peritoneal cavity. To ensure adequate library coverage, 2.4x10^6 ID8-Cas9.Lib tagged cells were injected for CRISPR screening experiments (>1000x coverage). In the repeated injection model, mice received 4x10^5 cells per injection,

administered six times over the course of 12 days. Mice were typically sacrificed 6

weeks after initial dose of cancer cells. For monitoring metastasis in luciferase-

expressing models, 100 ul of D-luciferin (15 mg/mL stock, GoldBio cat#: LUCK-500)

was administered intraperitoneally and imaged using an AMI-HTX spectral imaging

instrument. The total emission (flux) data for each region of interest (ROI) was obtained

after image analysis using the Aura software.

Macrophage Depletion

Macrophage depletion was achieved using encapsulated liposomal chlodronate

(Encapsula Nano Sciences, SKU# CLD-8901). Mice received a pre-treatmeant with 200

ul chlodrosome or empty liposome nanoparticles i.p. 24 hours before cancer cell

injection. Following pre-treatment, 100 ul of chlodrosome was administered twice

weekly throughout the study duration.

Individual Gene Knockout Guide Cloning

The CROP-seq-V2 plasmid (Addgene # 127458) was first modified to include a P2A-

mCherry sequence downstream of the puromycin resistance gene (PuroR) through a

NEBuilder HiFi DNA Assembly (NEB, E5520S) reaction. Then, single-guide RNA

(sgRNA) sequences were designed using CRISPick (Mouse GRCm38 reference

genome), oligos synthesized by IDT with BsmBI overhangs, annealed, and cloned into

the modified CROP-seq-V2 plasmid using the NEBridge Golden Gate Assembly Kit

(NEB, E1602S). The resulting plasmids were first transformed into competent *E. coli*,

plated on LB-ampicilin (AMP) plates, and grown at 30 ºC, before plasmid extraction.

Below are the sgRNA sequences used for gene targeting, including the necessary

BsmBI overhangs:

sgNTC-1 Top oligo

5'- **CAC CG**A CCT GAT ACG TCG TCG CGT A -3'

sgNTC-1 Bottom oligo

5'- **AAA C**TA CGC GAC GAC GTA TCA GGT **C** -3'

sgNTC-2 Top oligo

5'- **CAC CG**G TAT TAC TGA TAT TGG TGG G -3'

sgNTC-2 Bottom oligo

5'- **AAA C**CC CAC CAA TAT CAG TAA TAC **C** -3'

sgIrf1 guide 1 Top oligo

5'- **CAC CG**T GGA AGC ACG CTG CTA AGC A -3'

sgIrf1 guide 1 Bottom oligo

5'- **AAA C**TG CTT AGC AGC GTG CTT CCA **C** -3'

sgIrf1 guide 2 Top oligo

5'- **CAC CG**C TGT AGG TTA TAC AGA TCA G -3'

sgIrf1 guide 2 Bottom oligo

5'- **AAA C**CT GAT CTG TAT AAC CTA CAG **C** -3'

sgStat1 guide 1 Top oligo

5'- **CAC CG**T ATC CTG TGG TAC AAC ATG C -3'

sgStat1 guide 1 Bottom oligo

5'- **AAA C**GC ATG TTG TAC CAC AGG ATA **C** -3'

sgStat1 guide 2 Top oligo

5'- **CAC CG**G CAA GCG TAA TCT CCA GGT A -3'

sgStat1 guide 2 Bottom oligo

5'- **AAA C**TA CCT GGA GAT TAC GCT TGC **C** -3'

sgIfngr1 guide 1 Top oligo

5'- **CAC CG**G GCT CGG AGA GAT TAC CCG A -3'

sgIfngr1 guide 1 Bottom oligo

5'- **AAA C**TC GGG TAA TCT CTC CGA GCC **C** -3'

sgIfngr1 guide 2 Top oligo

5'- **CAC CG**T ATA CCA ATA CGC AAA TAC C -3'

sgIfngr1 guide 2 Bottom oligo

5'- **AAA C**GG TAT TTG CGT ATT GGT ATA **C** -3'

Library Cloning

Six MetTag-LARRY-BC.ID libraries were generated via overhang PCR by amplifying the

region within the original pLARRY-eGFP plasmid containing random barcodes. Primers

were designed with overhangs containing: (1) a unique 6-nucleotide library identifier

(ID), (2) a chromium 10x capture sequence, and (3) BsmBI restriction sites flanking both

ends of the barcode (synthesized by Sigma-Aldrich). Amplified barcode fragments were

introduced into the pRG212 third-generation lentiviral plasmid using the NEB Golden

Gate Assembly Kit (NEB, E1602S) at a 2:1 insert-to-vector molar ratio. The cloning

reaction was transformed into Endura electrocompetent *E. coli* cells (Biosearch

Technologies, 60242-1) and plated overnight on LB-AMP plates. $3x10^6$-$2x10^7$ CFU

colonies were harvested by flushing prewarmed LB broth and cultured for 2 hours in

liquid LB. sgRNA oligo pools were synthesized as oPools (IDT), containing BsmBI

cloning compatible overhangs. The oligos were then annealed and introduced into the

CROP-seq-V2.mCherry lentiviral plasmid using the NEB Golden Gate Assembly Kit as

described above. Approximately 5x10^7 CFUs for OvCa library and 2x10^7 colonies for

MetTag library were harvested for maxiprep.

For both LARRY barcode and CRISPR libraries, plasmids were purified using the

ZymoPURE II Plasmid Maxiprep kit (Zymo Research, D4202). The MetTag-LARRY-

BC.ID vector map and CROP-seq-V2.mCherry vector map will be made available on

Addgene upon publication.

Plasmid Library Amplification and Diversity Validation

Plasmid DNA was amplified using two step PCR to first amplify and then add indexing

sequences to libraries. First round PCR was conducted using custom P5 and P7

adapters with overhangs compatible with Nextera indexing primers, followed by cleanup

with Agencourt AMPure XP beads (Beckman Coulter, NC9933872) at a 1.8x bead ratio

to remove primers. Index PCR reaction was conducted as outlined in the Illumina 16S

Metagenomic Sequencing Library Preparation manual (Illumina, Doc #15044223, page

11-12) using 2x KAPA HiFi HotStart ReadyMix (Roche, KK2601) and Nextera XT Index

kit v2 primers (Illumina, FC-131-1001). Final indexed libraries were quantified via the

Qubit HS assay kit. LARRY libraries were run on a Bioanalyzer instrument and

sequenced on an Illumina MiSeq instrument. CRISPR libraries were run on TapeStation

using the High Sensitivity D1000 assay (Agilent, 5067-5585, 5067-5594) and

sequenced on a NovaSeq-X instrument. Demultiplexed fastq files were filtered for high

quality reads, and the number of unique reads was quantified. For the CRISPR libraries,

a customized script was used to estimate the percentage coverage based on the

original guide RNA whitelist.

1005

Lentivirus Production and Target Cell Transduction

MetTag-LARRY-BC.ID, CROP-seq-V2.mCherry guide, or Lenti-Luciferase.P2A.Neo

(Addgene # 105621), and second-generation lentiviral packaging plasmids were

transfected into HEK293T cells along with the CalFectin transfection reagent

(SignaGen, SL100478). Lentivirus was harvested after 72 hours and concentrated

(Lenti-X Concentrator, Takara Bio, 631231). Concentrated lentivirus was added to target

cells along with 8 ug/ml polybrene. To achieve low MOI for CRISPR screening

experiments, ID8 p53$^{-/-}$ ovarian cancer cells were infected at a low multiplicity of

infection (MOI) of 0.15, as calculated by the percentage of mCherry-positive cells 48

hours post transduction. 48 hours post transfection, cells were selected with either

G418 (pRG212-LARRY; Sigma-Aldrich, 4727878001) or puromycin (CROP-seq-

V2.mCherry; Invivogen, ant-pr-1) for a minimum of 7 days.

1018

Genomic DNA (gDNA) Amplicon Library Preparation

Genomic DNA was extracted from cultured cells and digested tissues using either the

Quick-DNA Midiprep Kit (Zymo Research, D4075) or the Quick-DNA Miniprep Plus Kit

(Zymo Research, D4068), depending on the starting cell number. Extracted DNA yield

was quantified via a NanoDrop spectrophotometer, and samples were normalized to the

same, lowest input concentration prior to PCR. Two rounds of PCR amplification were

performed on > 200 ng starting gDNA material. The first round was utilized for amplicon

amplification with nested primers for 15-20 cycles using a high-fidelity hot start

polymerase, followed by purification with a 1.8x bead-to-sample AMPure XP bead ratio.

Eluted amplicon DNA was quantified using the Qubit high sensitivity (HS) assay kit

(Thermo Fisher Scientific, Q33231) and concentrations normalized prior to indexing.

Approximately 5 ng of DNA was carried into the index PCR rection, which was

conducted similarly to the plasmid library construction protocol described above. Index

products were purified using the same AMPure XP beads ratio to remove excess

primers. Library quantity was assessed via the Qubit HS assay kit and analyzed on a

TapeStation instrument using the High Sensitivity D1000 assay to confirm expected library size and distribution. LARRY gDNA libraries were sequenced in-house on an Illumina MiniSeq instrument (Mid output kit, 300 cycles) while CRISPR gDNA libraries (101 genes*10 guides per gene, 20 non targeting guides = 1,030 guides) were sequenced on a NovaSeq-X instrument. High quality reads were filtered and analyzed via two parallel approaches: LARRY barcode counts were extracted to quantify library batch identity (ID), while CRISPR libraries were processed using the MAGeCK pipeline to identify enriched or depleted guide RNAs. Primers used for amplification were as follows:

LARRY Round 1 PCR REV, Tm: 55.9:

5' – GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GTG GGG AAT TTC TAC TCT TGT AGA TCG – 3'

LARRY Round 1 PCR FWD Stagger, Tm: 59.0:

5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CTT GCT AGG ACC GGC CTT AAA GC -3'

CRISPR Round 1 PCR REV, Tm: 51.0:

5' – GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GTT CTT GGC TTT ATA TAT CTT GTG G– 3'

CRISPR Round 1 PCR FWD Stagger, Tm: 53.0:

5' – TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CTC AAG TTG ATA ACG GAC TAG C– 3'

PiggyBac Transposon Transfection

SpCas9 and CreERT2 coding sequences were cloned into PiggyBac transposon plasmids containing a hygromycin resistance cassette via Gateway cloning (ThermoFisher Scientific, 11791020 & 11789020). Target cells were plated in 6-well plates 24 hours prior to transfection. Five ug of the SpCas9 or 1 ug of CreERT2 PiggyBac plasmid was co-transfected with 0.5 ug of Super PiggyBac Transposase (System Biosciences, PB210PA-1), along with the Calfectin transfection reagent. The transfection mixture was added to cells in a dropwise manner. 12-16 hours post transfection complete growth media was replenished. Forty-eight hours post

1065 transfection, cells were selected with hygromycin B (Corning, 30-240-CR) for a

1066 minimum of 10 days to enrich for successful transgene integrated cells.

1067

Cell Preparation for Single Cell RNA Sequencing

Naïve and tumor bearing mice were sacrificed via cervical dislocation six weeks

following initial cancer cell injection. To collect peritoneal cells, 5 mL of cold 1xHBSS

was injected intraperitoneally using a 25G needle, followed by lavage and peritoneal

fluid retrieval using a 20G needle. Omenta were excised from the peritoneal cavity and

examined under a dissecting microscope to confirm the presence of macroscopic

metastases. Omental tissues were then minced and digested for 30 minutes at 37ºC in

RPMI-1640 medium containing 0.4% collagenase type II (Gibco, 17101015), 50 ug/ml

DNAseI, and 0.1% BSA. Following digestion, cell suspensions were passed through a

70-uM strainer and washed with cold 1xPBS. Lavage, ascites, and omentum samples

were resuspended in 1x RBC lysis buffer (BioLegend, 420301) to remove red blood

cells. To prepare for single-cell analysis, omental samples were stained with ZombieRed

(BioLegend, 423109) to exclude dead cells and sorted via FACS. Sorted cells were then

prepared for 10X Genomics Chromium Single Cell Gene Expression analysis. A detailed

protocol for sample multiplexing via hashtags can be found in the CITE-seq/cell hashing

protocol on the CITE-seq website (https://cite-seq.com/wp-

content/uploads/2019/02/cite-seq_and_hashing_protocol_190213.pdf). In brief, samples

were first blocked with FcR Blocking Reagent (Miltenyi Biotec, 130-092-575) in 50 uL of

cell staining buffer for 20 minutes on ice. Samples were then stained with TotalSeq™-A

hashing antibodies obtained from BioLegend, including:

M-HTO-1 (M1/42; 30-F11, 155801),

M-HTO-2 (M1/42; 30-F11, 155803),

M-HTO-3 (M1/42; 30-F11, 155805),

M-HTO-4 (M1/42; 30-F11, 155807),

M-HTO-5 (M1/42; 30-F11, 155809),

M-HTO-6 (M1/42; 30-F11, 155811),

M-HTO-7 (M1/42; 30-F11, 155813),

M-HTO-8 (M1/42; 30-F11, 155815).

Staining was performed for 25 minutes on ice, followed by three washes with staining

buffer (decreasing EDTA concentration per wash) and loading onto a 10X Genomics

Chromium Controller.

1099

scRNA-seq Library Preparation and Illumina Sequencing

Single-cell transcriptomic libraries were prepared as outlined in the Chromium Next

GEM Single Cell 3' Reagents Kits v3.1 with Feature Barcoding technology kit manual.

Kits necessary for transcriptome library preparation were: Chromium Next GEM Single

Cell 3' GEM, Library & Gel Bead Kit v3.1, 4 rxns (10x Genomics, PN-1000128),

Chromium Single Cell 3' Feature Barcode Library Kit, 16 rxns (10x Genomics, PN-

1000079), Chromium Next GEM Chip G Single Cell Kit, 16 rxns (10x Genomics, PN-

1000127), and Single Index Kit T Set A, 96 rxns (10x Genomics, PN-1000213).

Hashtag libraries were prepared using the standard CITE-seq protocol (https://cite-

seq.com/wp-content/uploads/2019/02/cite-seq_and_hashing_protocol_190213.pdf

). LARRY-BC.ID barcode libraries were constructed using a custom nested PCR

protocol followed by indexing. Primers required for LARRY-BC.ID library preparation

included:

Round 1 PCR FWD, Tm: 61.0

5' – *GCA GCG TCA GAT GTG TAT AAG AGA CAG* – 3'

Round 1 PCR REV: 45.0

5' – <u>CCT TGG CAC CCG AGA ATT CCA</u> aat ttc tac tct tgt aga t – 3'

Index PCR FWD, Tm: 61.0

5' – AAT GAT ACG GCG ACC ACC GAG ATC TAC ACT CGT CG*G CAG CGT CAG ATG*

TGT ATA AGA GAC AG – 3'

Index PCR REV (ADT-RPI-1), Tm 61.0:

5' – CAA GCA GAA GAC GGC ATA CGA GAT CGT GAT GTG ACT GGA GTT <u>CCT TGG</u>

<u>CAC CCG AGA ATT CCA</u> – 3'

{Partial Read 1N; LARRY transcript (lowercase), <u>P7 adaptor</u>*,* RPI1 **index***}*

1124

1125 The quality and quantity of the three individual libraries (transcriptome, HTO, and

1126 LARRY) were assessed via Qubit and TapeStation prior to sequencing. Libraries were

1127   pooled for sequencing on an Illumina NovaSeq X Plus platform with paired end reads

1128   (PE150).

1129

scRNA-seq Data Analysis

Raw fastq files generated from the sequencing runs (mRNA, HTO, LARRY outputs)

were processed through the 10x Genomics CellRanger pipeline v.7.2 to generate count

matrices. Gene expression (mRNA) reads were aligned to a custom mouse reference

genome containing fluorescent marker transcripts (mm10_3.0.0_FPs). Data from two

different chromium 10x lanes, ascites lane and omentum lane, were merged using the

SCT-normalized dataset integration method. Details related to integration steps can be

found at: https://satijalab.org/seurat/articles/. In brief, individual lane single cell data was

normalized for HTO (HTOdemux) followed by SCTransform prior to integration.

Downstream analyses including dimensional reduction via principal component analysis

(PCA), clustering and UMAP embedding, and differential gene expression (DEG)

analysis were performed in RStudio under the Seurat package (v4.0)[18]. Cell type

identities were assigned based on cluster marker genes. Subcluster level labels

indicating different cellular statuses were determined through a combination of cluster

marker genes and examining enriched pathways. RNA velocity analysis was performed

in Python using scVelo and CellRank. Loom files were generated for each lane and

merged, followed by Seurat object conversion to H5AD format. Metadata containing

clustering and HTO information was integrated into the SCANPY anndata object. More

details on the CellRank method can be found at:

https://cellrank.readthedocs.io/en/latest/index.html. NicheNetR and CellChat were

performed in RStudio. Sender and receiver cell types were selected prior to NicheNetR

analysis. scRNA-seq expression data was analyzed using the Wilcoxon rank sum test

either through the direct function wilcox.test() when assigning custom module scores or

through the FindMarkers() function as part of the standard DEG analysis. Adjusted p

values <0.05 were considered significant. Over-representation analysis (ORA) was

performed on a filtered list of upregulated or downregulated genes from DEG analysis

and the ReactomePA and clusterProfiler RStudio packages. org.Mm.eg.db package was

used to convert genes to Entrez ID symbols. Adjusted p values <0.05 and log fold

change > (+/-) 0.25 were considered significant. ORA uses Fisher's exact test to

determine enriched gene sets. For Single Cell Gene Set Enrichment Analysis

(ssGSEA), we utilized the Escape R package[19].

1161

Flow cytometry

Samples were extracted and pre-processed similarly as the scRNA-seq samples.

Following tissue digestion of lung or omentum tissues and RBC lysis, digested samples

were washed in 1x Flow Staining Buffer (Tonbo Biosciences, TNB-4222-L500). For

viability assessment, cells were incubated with ZombieAqua Fixable Viability Kit

(BioLegend, 423102) at a 1:750 dilution in 1xPBS for 20 minutes at room temperature.

Following viability staining, cells were blocked with 1x FcR Blocking Reagent (Miltenyi

Biotec, 130-092-575) to minimize nonspecific antibody staining. Cells were then stained

with a combination of the following fluorochrome-conjugated surface antibodies: Pacific

Blue™ anti-mouse CD45 (BioLegend, 157211, S18009F), APC/Fire™ 750 anti-mouse

CD3 (BioLegend, 100247, 17A2) or PE/Cyanine7 anti-mouse CD3 (BioLegend, 100219,

17A2), PE anti-TREM-2 (BioLegend, 824805, 6E9), APC anti-mouse CCR5 (BioLegend,

107011, HM-CCR5), Brilliant Violet 605™ anti-mouse/human CD11b (BioLegend,

101237, M1/70), APC anti-mouse I-A-I-E (BioLegend, 107613, M5/114.15.2), Brilliant

Violet 711™ anti-mouse CD4 (BioLegend, 100447, GK1.5), PE/Dazzle™ 594 anti-

mouse NK-1.1 (BioLegend, 156517, S17016D) or PE anti-mouse NK-1.1 (BioLegend,

156503, S17016D), Brilliant Violet 421™ anti-mouse PD-L1 (BioLegend, 124315,

10F.9G2), PE/Cyanine7 anti-mouse CD8a (BioLegend, 100721, 53-6.7) or

APC/Cyanine7 anti-mouse CD8a (BioLegend, 100713, 53-6.7), Brilliant Violet 785™

anti-mouse F4/80 (BioLegend, 123141, BM8). Cancer cells harvested from culture were

similarly stained with ZombieAqua, FcR blocked, and then incubated with IFNGR1

Recombinant Rabbit Monoclonal Antibody FITC (Invitrogen, MA5-46853, 062) with the

corresponding Rabbit IgG Isotype Control FITC, eBioscience™ (Invitrogen, 11-4614-

80). Cells were then washed with staining buffer, followed by analysis on a Cytek

Northern Lights flow cytometer. FCS files were exported and analyzed using FlowJo

(BD Biosciences).

1188

Western Blotting

Cells were plated at a density of 250,000-500,000 cells per 10 cm dish for regular attachment plate experiments. For anoikis assays, 3x 6 wells per sample were seeded with 300,000 cells each. The next day, cells were either treated with vehicle (1xPBS) or 100 ng/ml mouse recombinant IFNγ (PeproTech, AF-315-05-100UG). After treatment, cells were harvested, washed with 1xPBS, and lysed in M2 lysis buffer (50 nM Tris-HCl, pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.5 mM EDTA) with 1x Protease/Phosphatase Inhibitor Cocktail (CST, 5872). Cells were lysed for 1 hour on ice before centrifugation at high speed to collect protein lysates. Protein quantity was deterimed using the Pierce^{TM} BCA Protein Assay Kit (Thermo Scientific, 23225). Protein lysate samples utilized for GBP-1 detection were resolved using 4-20% Mini-PROTEAN TGX Precast Protein Gels (Bio-Rad, 4561094). Other immunoblots were performed on fixed concentration using 8-10% SDS-polyacrylamide gels. All gel transfers were performed onto nitrocellulose membranes (Fisher Scientific, 45-004-075). Nitrocellulose membranes were blocked in 5% BSA/1xTBST for 1 hour at room temperature, followed by primary antibody incubation overnight on a 4°C shaker. The following day, blots were washed with 1xTBST followed by incubation with rabbit or mouse-specific horseradish peroxidase (HRP)-conjugated secondary antibodies. Signals were developed with SuperSignal West Pico PLUS Chemiluminescent Substrate (Thermo Fisher Scientific, #34577) or SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific, #34094). Primary antibodies used for western blotting were:

Beta-Actin (13E5) Rabbit mAb, CST #4970, 1:5,000 dilution

GBP-1 Rabbit Polyclonal Ab, {Thermo, PA5-23509}, 1:1,000 dilution

Cas9 (S. pyogenes) (7A9-3A3) Mouse mAb, CST #14697, 1:1,000 dilution

IRF-1 (D5E4) XP® Rabbit mAb, CST #8478, 1:1,000 dilution

Cre Recombinase (D7L7L) XP® Rabbit mAb, CST #15036, 1:1,000 dilution

GAPDH (14C10) Rabbit mAb, CST #2118, 1:5,000 dilution

Stat1 (D1K9Y) Rabbit mAb, CST #14994, 1:1,000 dilution

p-Stat1 (Tyr701) (58D6) Rabbit mAb #9167, 1:1,000 dilution

Cleaved Caspase-3 (Asp175) Rabbit Ab #9661, 1:750 dilution

SQSTM1/p62 Rabbit Ab #5114, 1:1,000 dilution

STAT3 (124H6) Mouse mAb #9139, 1:1,000 dilution

IDO (D8W5E) Rabbit mAb #51851, 1:1,000 dilution

Phospho-STAT3 (Tyr705) (D3A7) XP® Rabbit mAb #9145, 1:1,000 dilution

Vinculin (E1E9V) XP® Rabbit mAb #13901, 1:5,000 dilution

Anti-rabbit IgG, HRP-linked Ab, CST #7074, 1:5,000 dilution

Anti-mouse IgG, HRP-linked Ab, CST #7076, 1:5,000 dilution

1226

Clinical dataset mining

Patients were stratified into IFNGR1 high and IFNGR1 low expression groups and

survival curve was generated using Kaplan-Meier Plotter (KMplotter,

https://www.kmplot.com). This database includes 1816 samples primarily from

GSE9891 and TCGA[32]. Correlation curves were drawn in RStudio upon downloading

the TCGA-OV patient dataset[56] and calculating IFNγ, dormancy, and pro-survival

scores. Patient scRNA-seq RDS file was obtained from Mendeley Data[57]

(https://doi.org/10.17632/rc47y6m9mp.1) and contained 14 patient samples including

PBMCs, lymph nodes, primary tumor, metastasis, and malignant ascites. The RNA data

was re-normalized and analyzed using the Seurat package in RStudio in similar ways to

the murine scRNA-seq dataset described above.

1238

Quantitative Reverse Transcriptase (RT) PCR

The IFNγ treatment time course was performed in 12 well plates. After seeding 1.0x105

cells per well, wells were treated with 100 ng/ml IFNγ for 0 hours (PBS control), 4 hours,

8 hours, or 24 hours. Total cell RNA was extracted from cells using the Direct-zol RNA

Miniprep kit with TriReagent (Zymo Research, R2051-A) following the manufacturer

protocol. One microgram of total RNA was used for reverse transcription to synthesize

single stranded cDNA using the iScript^{TM} Reverse Transcription Supermix for RT-qPCR

(Bio-Rad, 1708840). qPCR reactions were performed on a Bio-Rad CFX Opus

instrument with SsoAdvanced Universal SYBR Green Supermix (Bio-Rad, 1725270).

Data analysis was performed using the CFX Maestro Software. Cq values were used to

calculate relative gene expression levels. In brief, calculations were based on the $2^{-\Delta\Delta CT}$

method with housekeeping genes serving as internal controls and fold change was

calculated by normalizing to the zero-hour treatment timepoint Cq values. Statistical

analyses and plots were generated in GraphPad Prism. RT-qPCR primers used were as

follows:

Beta-actin (Mouse) Forward primer TM: 57.6

5'- GGC TGT ATT CCC CTC CAT CG -3'

Beta-actin (Mouse) Reverse primer TM: 55.9

5'- CCA GTT GGT AAC AAT GCC ATG T-3'

GAPDH (Mouse) Forward primer TM: 59.8

5'- CAT CAC TGC CAC CCA GAA GAC TG -3'

GAPDH (Mouse) Reverse primer TM: 61.6

5'- ATG CCA GTG AGC TTC CCG TTC AG -3'

Gbp2b (Mouse) Forward primer TM: 58.5

5'- ATC TGC TCA TTG CTC AGA CTT ACT GG -3'

Gbp2b (Mouse) Reverse primer TM: 57.4

5'- AAG TAT TTT CTC AGC ATC ACT TCA GCC -3'

Irf1 (Mouse) Forward primer TM: 60.4

5'- ACC AAA TCC CAG GGC TGA TCT GG -3'

Irf1 (Mouse) Reverse primer TM: 58.3

5'- GAA GTT GGC CTT CCA TGT CTT GGG-3'

Stat1 (Mouse) Forward primer TM: 55.1

5'- GCT GCC TAT GAT GTC TCG TTT -3'

Stat1 (Mouse) Reverse primer TM: 55.2

5'- TGC TTT TCC GTA TGT TGT GCT -3'

Marco (Mouse) Forward primer TM: 61.1

5'- ATC CAC CTG ATC CTG CTC ACG G -3'

Marco (Mouse) Reverse primer TM: 60.8

5'- CAC TGC AGC GAG AAG AAG GGC -3'

Slfn1 (Mouse) Forward primer TM: 60.9

5'- AGG GAA AGG TCA AGG CTC ACA TTA AAA ATC C -3'

Slfn1 (Mouse) Reverse primer TM: 60.1

5'- AGG CTT CCC AAC AGA GAC ATC TGG -3'

1282

Anoikis Assays

Cancer cells (1×10^4) were seeded into 96-well ultra-low attachment (ULA) plates

(Corning, CLS7007) to mimic anchorage-independent conditions in the peritoneal cavity

environment. After 24 hours, 100 ng/ml recombinant mouse IFNγ or vehicle control

(1xPBS) was added to appropriate wells. Twenty-four hours after treatment (i.e. 48hr

after start of culture), CellTiter-Glo Luminescent Cell Viability Assay reagent (Promega,

G7570) was applied to every assay well, and plates were incubated for 20 min at room

temperature. Luminescence, indicating cellular ATP levels, was measured using a

Tecan plate reader instrument and analyzed using GraphPad Prism software.

For macrophage coculture experiments, 1.0×10^4 cancer cells were seeded together with

1.0×10^4 macrophages per well in ULA plates. Macrophages were enriched from the

peritoneal cavity of wild type female mice using the MojoSort Mouse F4/80 Positive

Selection Kit (BioLegend, 480170), following manufacturer's protocol. After 48 hours of

coculture, cell suspensions were harvested, pelleted, and incubated in TrypLE (Thermo

Fisher Scientific, 12605028) for 10 minutes at 37 ºC to dissociate cell aggregates. Cell

suspensions were passed through a 40-uM strainer before flow cytometric staining as

described earlier.

1300

Colony Formation Assay

Two hundred ID8 p53$^{-/-}$ cells were plated onto 6 well plates at the start of the

experiment. For wells treated with IFN-γ, 100 ng/ml IFN-γ was maintained throughout

the 7-day colony formation assay period. Wells were washed with 1xPBS, fixed with

methanol, and stained with crystal violet solution (0.5% w/v) at room temperature for 20

minutes. Wells were then washed with deionized water and dried overnight. Colonies

were counted and quantified in GraphPad Prism.

1308

Validation of Cre-based NicheTracing

HEK293T cells were transfected with a loxp-dsRED-loxp-eGFP Cre reporter construct

(Addgene # 141148) and cocultured in a 1:1 ratio with Cre.ERT2 expressing ID8 p53$^{-/-}$

or E0771 cell lines in the presence or absence of 2 uM 4-hydroxytamoxifen (4-OHT,

MedChemExpress, Cat # HY-16950). Seventy-two hours after plating, wells were

assessed for GFP+ switched cells using a Nikon Ti2 microscope. For independent *in*

vivo validation, E0771-Cre.ERT2 cells were injected into the retro-orbital sinus of Ai6

mice. One week post injection, three consecutive tamoxifen or corn oil control

treatments were performed (75 mg/kg/dose) i.p.. Two weeks post cancer cell injection,

mice were euthanized, perfused intracardially, and lungs were extracted for imaging and

flow cytometry analysis. For the ovarian metastasis model, tamoxifen injections were

administered in five consecutive doses, two weeks post ID8-Cre.ERT2 cell i.p. injection.

1321

Data and Resource Availability

scRNA-seq data with LARRY barcode and HTO multiplexing was deposited at GEO:

GSE302619 and will be made publicly available upon publishing. Human scRNA-seq

data has been published elsewhere and is publicly available: GSE14764.

Code and any additional information required to reanalyze the data reported here can

be requested from the lead contact.

1328

1329

1330 **REFERENCES**

1331

1. Gerstberger, S., Jiang, Q., and Ganesh, K. (2023). Metastasis. Cell *186*, 1564–1579.
https://doi.org/10.1016/j.cell.2023.03.003.

2. Ritch, S.J., and Telleria, C.M. (2022). The Transcoelomic Ecosystem and Epithelial
Ovarian Cancer Dissemination. Front. Endocrinol. *13*, 886533.
https://doi.org/10.3389/fendo.2022.886533.

3. Motohara, T., Masuda, K., Morotti, M., Zheng, Y., El-Sahhar, S., Chong, K.Y., Wietek, N.,
Alsaadi, A., Carrami, E.M., Hu, Z., et al. (2019). An evolving story of the metastatic
voyage of ovarian cancer cells: cellular and molecular orchestration of the adipose-rich
metastatic microenvironment. Oncogene *38*, 2885–2898.
https://doi.org/10.1038/s41388-018-0637-x.

4. Latifi, A., Luwor, R.B., Bilandzic, M., Nazaretian, S., Stenvers, K., Pyman, J., Zhu, H.,
Thompson, E.W., Quinn, M.A., Findlay, J.K., et al. (2012). Isolation and Characterization
of Tumor Cells from the Ascites of Ovarian Cancer Patients: Molecular Phenotype of

Chemoresistant Ovarian Tumors. PLOS ONE *7*, e46858.
https://doi.org/10.1371/journal.pone.0046858.

5. Ahmed, N., and Stenvers, K. (2013). Getting to Know Ovarian Cancer Ascites:
Opportunities for Targeted Therapy-Based Translational Research. Front. Oncol. *3*.
https://doi.org/10.3389/fonc.2013.00256.

6. Almeida-Nunes, D.L., Mendes-Frias, A., Silvestre, R., Dinis-Oliveira, R.J., and Ricardo, S.
(2022). Immune Tumor Microenvironment in Ovarian Cancer Ascites. Int. J. Mol. Sci. *23*,
10692. https://doi.org/10.3390/ijms231810692.

7. Lane, D., Matte, I., Garde-Granger, P., Bessette, P., and Piché, A. (2018). Ascites IL-10
Promotes Ovarian Cancer Cell Migration. Cancer Microenviron. *11*, 115–124.
https://doi.org/10.1007/s12307-018-0215-3.

8. Mustea, A., Pirvulescu, C., Könsgen, D., Braicu, E.I., Yuan, S., Sun, P., Lichtenegger, W.,
and Sehouli, J. (2008). Decreased IL-1 RA concentration in ascites is associated with a
significant improvement in overall survival in ovarian cancer. Cytokine *42*, 77–84.
https://doi.org/10.1016/j.cyto.2008.01.011.

9. Yamanoi, K., Matsumura, N., Murphy, S.K., Baba, T., Abiko, K., Hamanishi, J.,
Yamaguchi, K., Koshiyama, M., Konishi, I., and Mandai, M. (2016). Suppression of
ABHD2 , identified through a functional genomics screen, causes anoikis resistance,
chemoresistance and poor prognosis in ovarian cancer. Oncotarget *7*, 47620–47636.
https://doi.org/10.18632/oncotarget.9951.

10. Lineage tracing on transcriptional landscapes links state to fate during differentiation |
Science https://www.science.org/doi/10.1126/science.aaw3381.

11. Śliwa, A., Szczerba, A., Pięta, P.P., Białas, P., Lorek, J., Nowak-Markwitz, E., and
Jankowska, A. (2024). A Recipe for Successful Metastasis: Transition and Migratory
Modes of Ovarian Cancer Cells. Cancers *16*, 783.
https://doi.org/10.3390/cancers16040783.

12. Schrijver, W.A.M.E., Selenica, P., Lee, J.Y., Ng, C.K.Y., Burke, K.A., Piscuoglio, S., Berman,
S.H., Reis-Filho, J.S., Weigelt, B., van Diest, P.J., et al. (2018). Mutation Profiling of Key
Cancer Genes in Primary Breast Cancers and Their Distant Metastases. Cancer Res.
78, 3112–3121. https://doi.org/10.1158/0008-5472.CAN-17-2310.

13. Müller, V., and Pantel, K. (2004). Bone marrow micrometastases and circulating tumor
cells: current aspects and future perspectives. Breast Cancer Res. *6*, 258–261.
https://doi.org/10.1186/bcr942.

14. Sänger, N., Effenberger, K.E., Riethdorf, S., Van Haasteren, V., Gauwerky, J., Wiegratz, I.,
Strebhardt, K., Kaufmann, M., and Pantel, K. (2011). Disseminated tumor cells in the

bone marrow of patients with ductal carcinoma in situ. Int. J. Cancer *129*, 2522–2526. https://doi.org/10.1002/ijc.25895.

15. Ke, M., Sun, N., Lin, Z., Zhang, P., Hu, Y., Wu, S., Zheng, Z., Lu, Y., and Jin, H. (2023). SNHG18 inhibits bladder cancer cell proliferation by increasing p21 transcription through destabilizing c-Myc protein. Cancer Cell Int. *23*, 48. https://doi.org/10.1186/s12935-023-02887-w.

16. Fan, H., Yuan, J., Li, Y., Jia, Y., Li, J., Wang, X., and Li, X. (2021). MKL1-induced lncRNA SNHG18 drives the growth and metastasis of non-small cell lung cancer via the miR-211-5p/BRD4 axis. Cell Death Dis. *12*, 128. https://doi.org/10.1038/s41419-021-03399-z.

17. Dasgupta, A., Lim, A.R., and Ghajar, C.M. (2017). Circulating and disseminated tumor cells: harbingers or initiators of metastasis? Mol. Oncol. *11*, 40–61. https://doi.org/10.1002/1878-0261.12022.

18. Satija, R., Farrell, J.A., Gennert, D., Schier, A.F., and Regev, A. (2015). Spatial reconstruction of single-cell gene expression data. Nat. Biotechnol. *33*, 495–502. https://doi.org/10.1038/nbt.3192.

19. Borcherding, N., Vishwakarma, A., Voigt, A.P., Bellizzi, A., Kaplan, J., Nepple, K., Salem, A.K., Jenkins, R.W., Zakharia, Y., and Zhang, W. (2021). Mapping the immune environment in clear cell renal carcinoma by single-cell genomics. Commun. Biol. *4*, 122. https://doi.org/10.1038/s42003-020-01625-6.

20. Cellular barcoding to decipher clonal dynamics in disease | Science https://www.science.org/doi/10.1126/science.abm5874?url_ver=Z39.88-2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%20%200pubmed.

21. CellRank for directed single-cell fate mapping | Nature Methods https://www.nature.com/articles/s41592-021-01346-6.

22. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. OMICS J. Integr. Biol. *16*, 284–287. https://doi.org/10.1089/omi.2011.0118.

23. Bahrami, S., and Drabløs, F. (2016). Gene regulation in the immediate-early response process. Adv. Biol. Regul. *62*, 37–49. https://doi.org/10.1016/j.jbior.2016.05.001.

24. Koivisto, E., Acosta, A.J., Moilanen, A.-M., Tokola, H., Aro, J., Pennanen, H., Säkkinen, H., Kaikkonen, L., Ruskoaho, H., and Rysä, J. (2014). Characterization of the Regulatory Mechanisms of Activating Transcription Factor 3 by Hypertrophic Stimuli in Rat Cardiomyocytes. PLOS ONE *9*, e105168. https://doi.org/10.1371/journal.pone.0105168.

25. Zhao, J., Peng, H., Gao, J., Nong, A., Hua, H., Yang, S., Chen, L., Wu, X., Zhang, H., and
Wang, J. (2021). Current insights into the expression and functions of tumor-derived
immunoglobulins. Cell Death Discov. *7*, 148. https://doi.org/10.1038/s41420-021-
00550-9.

26. Mehta, A., and Merkel, O.M. (2020). Immunogenicity of Cas9 Protein. J. Pharm. Sci. *109*,
62–67. https://doi.org/10.1016/j.xphs.2019.10.003.

27. Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S., Brown, M.,
and Liu, X.S. (2014). MAGeCK enables robust identification of essential genes from
genome-scale CRISPR/Cas9 knockout screens. Genome Biol. *15*, 554.
https://doi.org/10.1186/s13059-014-0554-4.

28. Shultz, D.B., Sandhya Rani, M.R., Fuller, J.D., Ransohoff, R.M., and Stark, G.R. (2009).
Roles of IKK-β, IRF1, and p65 in the Activation of Chemokine Genes by Interferon-γ. J.
Interferon Cytokine Res. *29*, 817–824. https://doi.org/10.1089/jir.2009.0034.

29. Fisch, D., Bando, H., Clough, B., Hornung, V., Yamamoto, M., Shenoy, A.R., and Frickel,
E. (2019). Human GBP1 is a microbe-specific gatekeeper of macrophage apoptosis and
pyroptosis. EMBO J. *38*, e100926. https://doi.org/10.15252/embj.2018100926.

30. Tailor, D., Garcia-Marques, F.J., Bermudez, A., Pitteri, S.J., and Malhotra, S.V. (2023).
Guanylate-binding protein 1 modulates proteasomal machinery in ovarian cancer.
iScience *26*. https://doi.org/10.1016/j.isci.2023.108292.

31. Honkala, A.T., Tailor, D., and Malhotra, S.V. (2020). Guanylate-Binding Protein 1: An
Emerging Target in Inflammation and Cancer. Front. Immunol. *10*, 3139.
https://doi.org/10.3389/fimmu.2019.03139.

32. Győrffy, B. (2023). Discovery and ranking of the most robust prognostic biomarkers in
serous ovarian cancer. GeroScience *45*, 1889–1898. https://doi.org/10.1007/s11357-
023-00742-4.

33. Kim, Y.-N., Koo, K.H., Sung, J.Y., Yun, U.-J., and Kim, H. (2012). Anoikis Resistance: An
Essential Prerequisite for Tumor Metastasis. Int. J. Cell Biol. *2012*, 306879.
https://doi.org/10.1155/2012/306879.

34. Johnson, D.E., O'Keefe, R.A., and Grandis, J.R. (2018). Targeting the IL-6/JAK/STAT3
signalling axis in cancer. Nat. Rev. Clin. Oncol. *15*, 234–248.
https://doi.org/10.1038/nrclinonc.2018.8.

35. Al Zaid Siddiquee, K., and Turkson, J. (2008). STAT3 as a target for inducing apoptosis in
solid and hematological tumors. Cell Res. *18*, 254–267.
https://doi.org/10.1038/cr.2008.18.

36. Qin, H., Yeh, W.-I., De Sarno, P., Holdbrooks, A.T., Liu, Y., Muldowney, M.T., Reynolds,
S.L., Yanagisawa, L.L., Fox, T.H., Park, K., et al. (2012). Signal transducer and activator
of transcription-3/suppressor of cytokine signaling-3 (STAT3/SOCS3) axis in myeloid
cells regulates neuroinflammation. Proc. Natl. Acad. Sci. *109*, 5004–5009.
https://doi.org/10.1073/pnas.1117218109.

37. Chemi, F., Mohan, S., Guevara, T., Clipson, A., Rothwell, D.G., and Dive, C. (2021). Early
Dissemination of Circulating Tumor Cells: Biological and Clinical Insights. Front. Oncol.
11. https://doi.org/10.3389/fonc.2021.672195.

38. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung, P.,
Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell communication using
CellChat. Nat. Commun. *12*, 1088. https://doi.org/10.1038/s41467-021-21246-9.

39. Chen, S., Saeed, A.F.U.H., Liu, Q., Jiang, Q., Xu, H., Xiao, G.G., Rao, L., and Duo, Y.
(2023). Macrophages in immunoregulation and therapeutics. Signal Transduct. Target.
Ther. *8*, 207. https://doi.org/10.1038/s41392-023-01452-1.

40. Ahmed, I., and Ismail, N. (2020). M1 and M2 Macrophages Polarization via mTORC1
Influences Innate Immunity and Outcome of Ehrlichia Infection. J. Cell. Immunol. *2*,
108–115. https://doi.org/10.33696/immunology.2.029.

41. O'Carroll, C., Fagan, A., Shanahan, F., and Carmody, R.J. (2014). Identification of a
Unique Hybrid Macrophage-Polarization State following Recovery from
Lipopolysaccharide Tolerance. J. Immunol. *192*, 427–436.
https://doi.org/10.4049/jimmunol.1301722.

42. Browaeys, R., Saelens, W., and Saeys, Y. (2020). NicheNet: modeling intercellular
communication by linking ligands to target genes. Nat. Methods *17*, 159–162.
https://doi.org/10.1038/s41592-019-0667-5.

43. Pentcheva-Hoang, T., Egen, J.G., Wojnoonski, K., and Allison, J.P. (2004). B7-1 and B7-2
Selectively Recruit CTLA-4 and CD28 to the Immunological Synapse. Immunity *21*, 401–
413. https://doi.org/10.1016/j.immuni.2004.06.017.

44. Butte, M.J., Keir, M.E., Phamduy, T.B., Sharpe, A.H., and Freeman, G.J. (2007).
Programmed Death-1 Ligand 1 Interacts Specifically with the B7-1 Costimulatory
Molecule to Inhibit T Cell Responses. Immunity *27*, 111–122.
https://doi.org/10.1016/j.immuni.2007.05.016.

45. Iyer, S., Zhang, S., Yucel, S., Horn, H., Smith, S.G., Reinhardt, F., Hoefsmit, E., Assatova,
B., Casado, J., Meinsohn, M.-C., et al. (2021). Genetically Defined Syngeneic Mouse
Models of Ovarian Cancer as Tools for the Discovery of Combination Immunotherapy.
Cancer Discov. *11*, 384–407. https://doi.org/10.1158/2159-8290.CD-20-0818.

46. Zheng, X., Wang, X., Cheng, X., Liu, Z., Yin, Y., Li, X., Huang, Z., Wang, Z., Guo, W.,
Ginhoux, F., et al. (2023). Single-cell analyses implicate ascites in remodeling the
ecosystems of primary and metastatic tumors in ovarian cancer. Nat. Cancer *4*, 1138–
1156. https://doi.org/10.1038/s43018-023-00599-8.

47. Patras, L., Shaashua, L., Matei, I., and Lyden, D. (2023). Immune determinants of the
pre-metastatic niche. Cancer Cell *41*, 546–572.
https://doi.org/10.1016/j.ccell.2023.02.018.

48. Peinado, H., Zhang, H., Matei, I.R., Costa-Silva, B., Hoshino, A., Rodrigues, G., Psaila,
B., Kaplan, R.N., Bromberg, J.F., Kang, Y., et al. (2017). Pre-metastatic niches: organ-
specific homes for metastases. Nat. Rev. Cancer *17*, 302–317.
https://doi.org/10.1038/nrc.2017.6.

49. Hershey, B.J., Barozzi, S., Orsenigo, F., Pompei, S., Iannelli, F., Kamrad, S., Matafora, V.,
Pisati, F., Calabrese, L., Fragale, G., et al. (2023). Clonal cooperation through soluble
metabolite exchange facilitates metastatic outgrowth by modulating Allee effect. Sci.
Adv. *9*, eadh4184. https://doi.org/10.1126/sciadv.adh4184.

50. Wawrzyniak, P., and Hartman, M.L. (2025). Dual role of interferon-gamma in the
response of melanoma patients to immunotherapy with immune checkpoint inhibitors.
Mol. Cancer *24*, 89. https://doi.org/10.1186/s12943-025-02294-x.

51. Remsik, J., Tong, X., Kunes, R.Z., Li, M.J., Estrera, R., Snyder, J., Thomson, C., Osman,
A.M., Chabot, K., Sener, U.T., et al. (2025). Interferon-γ orchestrates leptomeningeal
anti-tumour response. Nature, 1–10. https://doi.org/10.1038/s41586-025-09012-z.

52. Jorgovanovic, D., Song, M., Wang, L., and Zhang, Y. (2020). Roles of IFN-γ in tumor
progression and regression: a review. Biomark. Res. *8*, 49.
https://doi.org/10.1186/s40364-020-00228-x.

53. Zheng, H., Guan, X., Meng, X., Tong, Y., Wang, Y., Xie, S., Guo, L., and Lu, R. (2023). IFN-γ
in ovarian tumor microenvironment upregulates HLA-E expression and predicts a poor
prognosis. J. Ovarian Res. *16*, 229. https://doi.org/10.1186/s13048-023-01286-z.

54. Reddy, S.U., Sadia, F.Z., Vancura, A., and Vancurova, I. (2024). IFNγ-Induced Bcl3, PD-
L1 and IL-8 Signaling in Ovarian Cancer: Mechanisms and Clinical Significance.
Cancers *16*, 2676. https://doi.org/10.3390/cancers16152676.

55. Ghisoni, E., Benedetti, F., Minasyan, A., Desbuisson, M., Cunnea, P., Grimm, A.J., Fahr,
N., Capt, C., Rayroux, N., Carlo, F.D., et al. (2025). Myeloid cell networks govern re-
establishment of original immune landscapes in recurrent ovarian cancer. Cancer Cell
0. https://doi.org/10.1016/j.ccell.2025.07.005.

56. McLendon, R., Friedman, A., Bigner, D., Van Meir, E.G., Brat, D.J., M. Mastrogianakis, G.,
Olson, J.J., Mikkelsen, T., Lehman, N., Aldape, K., et al. (2008). Comprehensive genomic

characterization defines human glioblastoma genes and core pathways. Nature *455*,
1061–1068. https://doi.org/10.1038/nature07385.

57. Wang, X., and Zheng, X. (2022). scRNA-seq data in ovarian cancer. *1*.
https://doi.org/10.17632/rc47y6m9mp.1.

1524

1525

1526

Graphical abstract

Figure 1:
Tracking Clonal Lineage in Ovarian Cancer Metastasis

Figure 1: Tracking Clonal Lineage in Ovarian Cancer Metastasis

(**A**) Schematic depicting main elements of the MetTag library barcoding construct. (**B**) Schematic of injection and tissue collection timeline for the staggered injection experimental model using MetTag-barcoded ID8 ovarian cell lines. Schematic was partially created with BioRender.com. (**C**) Genomic DNA (gDNA)-level barcode distributions in the omentum graphed as frequency under WT, NRG, and MAC depletion conditions. (**D-E**) gDNA-level barcode distributions in ascites and pelvic fat pads under WT and NRG conditions. In (C-E) panels, BC.ID1 = TP-1, BC.ID2 = TP-2, BC.ID3 = TP-3, BC.ID4 = TP-4, BC.ID5 = TP-5, BC.ID6 = TP-6 for clarity. (**F**) Injection schema for comparing one-time (400,000 total, n = 10) cell dose and fractionated (2,400,000 total, n = 10) cell injections (left) and representative BLI images at endpoint (right). (**G**) Bar graph depicting total flux emission in each group at experimental endpoint. (**H**) Schematic of scRNA-seq experiment conducted using a reversed BC.ID injection order. LARRY barcodes are readily captured by Chromium 10x Feature Barcoding. (**I**) Pie charts of RNA-level barcode frequencies in each location grouped as early (BC.ID5, BC.ID6), mid (BC.ID4, BC.ID3), and late (BC.ID2, BC.ID1). (**J**) Heatmap of sample level barcode frequencies, BC.ID6 = TP-1, BC.ID5 = TP-2, BC.ID4 = TP-3, BC.ID3 = TP-4, BC.ID2 = TP-5, BC.ID1 = TP-6. (**K**) Volcano plot of DEGs between early and late timepoint barcoded cancer cells where threshold for significance was defined at p <0.05 and log2FC < -1.0 for downregulation or >1.0 for upregulation. Genes labeled are the top 30 genes based on significance. Data in Figure 1G was analyzed using a two-tailed Student's t test and graphed as mean with standard deviation (SD). In Figure 1K, the DEG table was generated using the Seurat FindMarkers() function and p values calculated via a Wilcoxon rank-sum test. For all Wilcoxon rank-sum tests, p values graphed represent the adjusted p value.

Figure 2:
Clonal Analysis Reveals Shared Evolutionary Path Towards Inflammatory Signature

Figure 2: Clonal Analysis Reveals Shared Evolutionary Path Towards Inflammatory Signature
(**A**) Split UMAP depicting cluster distributions across the four experimental conditions and identifying cancer cell-enriched clusters in ascites and omentum. (**B**) Reclustered cancer cell UMAP colored by the two metastasis-bearing anatomical locations. (**C**) Stacked bar chart depicting the frequency of Seurat subclusters within each metastatic location. (**D**) Seurat cluster UMAP of reclustered cancer cells with transcriptional program labels. (**E**) Dot plot depicting Hallmark pathways across all subclusters used to define each cluster subset gene program in (D). (**F**) Schematic representation of the possible outcomes of LARRY clonal analysis based on transcriptome patterns of enriched clones. (**G**) Heatmap of normalized clonal LARRY barcode counts across the two anatomical locations and individual mice (HTOs). (**H**) UMAP overlay of top 3 expanded clones in the omentum. (**I**) Bar chart depicting the relative subcluster proportions present in each of the top 3 barcode clones. (**J**) scVelo-generated PAGA velocity graph indicating trajectory arrows and inferred cluster lineage. (**K**) scVelo and CellRank-based simulated path from start to endpoint clusters. (**L**) Violin plot depicting RNA-level contribution of each timepoint to the IFN-gamma signature. Data in Figure 2L was analyzed by first defining the IFN-gamma signature using canonical mouse genes, followed by directly applying the wilcox.test() function in Seurat to compare TP-1 and TP-6 expression.

Figure 3:
Activation of IFNγ Response in Ascites is Necessary for Successful Omental Colonization

Figure 3: Activation of IFN-gamma Response in Ascites is Necessary for Successful Omental Colonization
(A) DEG-based volcano plot of gene expression differences between OmMet and AscMet, where threshold for significance was defined at p <0.05 and log2FC < -1.0 for downregulation or >1.0 for upregulation. (B) Boxplots comparing IFN signature modules between AscMet and OmMet. (C) Schematic describing in vivo ovarian metastatic niche signature screen. Partially created with BioRender.com. (D) In vivo CRISPR screen RankView plots generated using MAGeCKFlute. OmMet (n=5) and AscMet (n=4) samples were compared to the original pool of ID8 cells. (E) Ranks of individual sgRNAs targeting positive control Pten and depleted genes. (F) Lollipop plot of depleted pathways associated with depleted genes in Fig. 3E. (G) IFN-gamma treatment time course qPCR plots of select CRISPR hits and positive controls. (H) IFN treatment time course western blot probing for Stat1/pStat1 and GBP1. (I) Schematic of in vivo experimental metastasis study comparing sgNTC and sgIFNGR1.g1 groups. (J) Time course log10-transformed total emission/flux values (left). Two independent groups pooled; cohort 1 NTC n = 8, KO n = 7; cohort 2 NTC n = 5, KO n = 5. log10-transformed flux plot for week six with p value (right). (K) Kaplan Meier survival curve generated for cohort 1. Data in Figure 3B was analyzed by first defining IFN signatures, followed by applying wilcox-.test() in Seurat to compare AscMet and OmMet. CRISPR screen related data (3D-F) was first analyzed using MAGeCK.

Count summary, gRNA summary, and gene summary output files were loaded into RStudio to generate rankview plots using MAGeCKFlute. For Figure 3G, qPCR data was analyzed by averaging technical replicates for each biological replicate, followed by calculating dCq by normalizing target gene Cq to housekeeping gene control. Next, $2^{\wedge}(-ddCq)$ or fold change was calculated by normalizing the values to the zero hour (PBS) treatment samples and plotted in Prism GraphPad. Significance between groups was assessed using a two-tailed Student's t test and data was plotted as mean with standard deviation (SD). Data in figure 3J left panel was plotted as log transformed mean with 95% confidence interval (CI) and a two-tailed Student's t test was performed at each timepoint. Data in 3J right panel was plotted with median log transformed values.

Figure 4: IFN-gamma Sensing Mediates Resistance to Anoikis and Activates Pro-survival Signaling
(**A**) Colony formation assay performed without (left) or with (right) 100 ng/ml IFN-gamma. (**B-C**) Cell-Titer Glo assay luminescence CTG values graphed as fold change for regular culture conditions (B) and ultra-low attachment conditions (C), with and without IFN-gamma treatment. Fold change was determined by normalizing each cell line's CTG value to the regular attachment, no IFN-gamma treatment condition. (**D**) CTG flux value dot plots for Stat1-KO and Irf1-KO cells under ultra-low attachment, with or without IFN-gamma treatment. (**E**) scRNA-seq based gating strategy used to identify Irf1-high and low cells for module scoring. (**F**) Violin plots for autophagy, stress, and pro-survival (anti-apoptotic) scores in Irf1-high vs. Irf1-low cells. Violin plots comparing *Stat3* (**G**), *Mcl-1*, and *Bcl2* (**H**) transcript levels between Irf1-high, mid, and low cells. (**I**) Western blot probing for Stat3 activation and SQSTM1 expression under ultra-low attachment conditions, with or without IFN-gamma treatments. (**J**) Violin plot of dormancy module comparing Irf1-high and low cells. CTG assay luminescence data (4B-4D) was analyzed in Prism using student's t test. Data in B and C was graphed as mean and standard deviation, data in D was graphed as median values. Anoikis and dormancy modules, along with *Stat3*, *Mcl-1*, and *Bcl2* transcript expression levels were compared between Irf1-high and Irf1-low cells by applying the wilcox.test() function in Seurat.

Figure 5:
Ascites and Omental TAMs Differ in Their Immunomodulatory Potential

Figure 5: Ascites and Omental TAMs Differ in Their Immunomodulatory Potential
(**A**) Violin plots depicting CRISPR screen hits expression level across Broad cell ID categories (cell types). (**B**) CellChat-generated circus plots of interaction strengths between Broad cell ID categories in metastatic omentums (left) and ascites (right). (**C**) Stacked bar chart of cell type proportions across the four experimental groups. (**D**) UMAPs of reclustered myeloid cells, split by experimental condition. (**E**) scVelo RNA velocity plot showing arrows pointing from ascites TAMs (clusters 1 and 5) to omentum TAMs (clusters 0 and 6). (**F**) DEG volcano plot comparing gene expression differences between ascites TAM cluster 1 and omentum TAM cluster 0. (**G**) Stacked bar chart of subsetted T/NK cells comparing cell proportions in ascites and omentum. (**H-I**) ORA generated Reactome pathways enriched in ascital (left) and omentum-derived TAMs (right). Significance of DEGs in Figure 5F was evaluated by applying the FindMarkers() and Wilcoxon test in Seurat to compare ascites and omental *Cd68*high/*Adgre1*high macrophages.

Figure 6:
NicheTracing Identifies Ascites-derived TAMs as Pro-Metastatic Interactors

Figure 6: NicheTracing Identifies Ascites-derived TAMs as Pro-Metastatic Interactors
(**A**) Schematic describing the Cre-based NicheTracing labeling system where ID8 and E0771 cancer cells express Cre.ERT2 which transfers to neighboring RCL-ZsGreen niche cells. Upon nuclear entry of Cre.ERT2, labeled niche cells turn ZsGreen positive (ZsG+). (**B**) Representative gating strategy depicting the major cell population tagged with zsG. (**C-D**) Bar plots showing preference for F4/80+ macrophage labeling in ascites. (**E**) Liposomal chlodronate-based macrophage depletion strategy and timeline schematic with representative tissue images. n= 7 liposome control group and n = 7 chlodrosome treatment group. (**F**) Bar plot showing quantification of omental metastases via omental weight measurement at endpoint. (**G**) Images of ascites samples harvested at endpoint for flow cytometry analysis (left) and flow-based quantification of CD45 negative cells (right). (**H**) Schematic of macrophage-cancer cell coculture layout (top) and representative coculture images for one cell line (bottom). (**I-K**) Plots of percent CD45 negative cancer cell viability post coculture. Data in Figures 6C-D and 6F-6H was plotted as mean with SD and differences between groups were compared via a two-tailed unpaired Student's t test. Data in Figures 6I-K was similarly analyzed using a Student's t test and plotted with median values.

Figure 7:
Patient Ascites MAC/Tumor Interactions are Enriched for Tumor-supporting Pathways

Figure 7: Patient Ascites MAC/Tumor Interactions are Enriched for Tumor-supporting Pathways
(**A**) Stacked bar chart of main cell type proportions captured in human scRNA-seq dataset across five anatomical locations, using main-type-2 annotation provided by publisher. (**B**) Heatmap depicting DEGs between Met.Ome and Ascites. (**C-D**) GO Molecular Function pathway level differences between omental MACs (C) and ascites MACs (D). (**E**) Schematic depicting NicheNetR analysis approach for finding top TAM ligands in both murine and human datasets. This panel was partially generated using BioRender.com. (**F**) Venn diagram of top NicheNet predicted ligands after filtering for robust expression and overlapping with top interactions in our murine dataset. (**G**) Heatmap showing top receptor-ligand interactions upregulated in patient ascites-MACs compared to metastases. (**H**) GO enrichment dotplot generated from the gene list containing top target genes in cancer cells that are predicted to respond to top ligands in the human dataset. Data in Figure 7B was analyzed using the FindMarkers() function in Seurat and Wilcoxon test to compare Asc and Met.Ome.