Temporal Clonal Tracing Reveals Tumor-Intrinsic IFNγ-Dependencies Driving Niche Adaptation and Early Metastatic Colonization

Emilija Aleksandrovic[1,3], Shaneann R. Fross[1,3], Samantha M. Golomb[1,3], Xiyu Liu[1,3], Zhuo Zhao[2,3], Nikitha M. Das[1,3], Tanner C. Reese[1,3], Wei Ma[1,3], Jacqueline Lopez[4], Zhenyu Zhong[3,7], Min Zhao[1,3], M. Sharon Stack[5,6], Siyuan Zhang[1, 3]

Affiliations:
1. Department of Pathology, UT Southwestern Medical Center, Dallas, TX, 75390, USA
2. Lyda Hill Department of Bioinformatics, School of Public Health, UT Southwestern Medical Center, Dallas, TX, 75390, USA
3. Harold C. Simmons Comprehensive Cancer Center, Dallas, TX, 75390, USA
4. Department of Biological Sciences, College of Science, University of Notre Dame, Notre Dame, IN 46556, USA
5. Department of Chemistry and Biochemistry, University of Notre Dame, Notre Dame, IN 46556, USA
6. Mike and Josie Harper Cancer Research Institute, University of Notre Dame, 1234 N. Notre Dame Avenue, South Bend, IN 46617, USA
7. Department of Immunology, UT Southwestern Medical Center, Dallas, TX, 75390, USA

* Correspondence to: Siyuan Zhang, M.D., Ph.D., Department of Pathology, UT Southwestern Medical Center, ND6.200A, 6001 Forest Park Rd., Dallas, TX 75235. E-mail: Siyuan.Zhang@UTSouthwestern.edu; Telephone: 214-648-6537

Key Words: Metastasis, Disseminating Tumor Cells (DTCs), Clonal dynamics, Metastatic niches, IFN-gamma (IFNγ), Anoikis, Single-cell RNA sequencing, MetTag barcoding, Lineage tracing, In vivo CRISPR Screen.

SUMMARY

Metastasis is an emergent continuum, driven by evolving reciprocal adaptations between continuously disseminating tumor cells (DTCs) and the specialized metastatic niches of distant organs. The interplay between intrinsic and niche-driven mechanisms that enables DTCs to survive and home to distant organs remains incompletely understood. Here, using MetTag, a single-cell barcoding and transcriptome profiling approach with time-stamped batch identifiers (BC.IDs) and functional CRISPR screening, we resolved the clonality, temporal dynamics, and molecular determinates of DTC colonization success across evolving metastatic niches. Deep sequencing of barcodes revealed preferred enrichment of early-disseminated clones across metastatic niches. Single-cell RNA sequencing (scRNA-seq) coupled with RNA velocity analyses in ascites and metastasis-bearing omenta uncovered an emergent and distinct interferon-gamma (IFNγ) centric transcriptional trajectory, enriched among early seeding clones. *In vivo* CRISPR/Cas9 screening of metastatic niche-specific signatures demonstrated that genes belonging to the IFNγ response are functionally important for peritoneal metastasis. Knockout of IFNγ receptor 1 (*Ifngr1*) in the first batch of DTCs significantly reduced metastatic burden and extended survival, underscoring the importance of tumor cell intrinsic IFNγ signaling in shaping post-seeding metastatic niche (PSMN) and subsequent metastatic co-evolution. Mechanistically, tumor intrinsic IFNγ response and ascites-derived tumor-associated macrophages (TAMs) protect cancer cells from anoikis-mediated death by promoting pro-survival signaling. Our study defines temporal dynamics of disseminating tumor cells at metastatic niches and reveals a general "first come, first served" pro-metastatic adaptation principle of DTCs.

INTRODUCTION

Cancer metastasis encompasses continuous metastatic dissemination from the primary tumor, followed by passive clonal selection, and active epigenetic niche adaptation in response to the metastatic niche[1]. Resolving the clonal, temporal, and microenvironmental determinants along the cancer metastatic continuum has long been a challenge in the field. Unlike most cancer types, where hematogenous metastasis

predominates, high-grade serous ovarian carcinoma (HGSOC) exhibits a unique pattern

of transcoelomic, continuous local spreading within the peritoneal cavity, representing a

metastatic continuum[2]. In HGSOC, malignant ascites functions as a "reservoir" of

floating cancer cells and cell clusters – DTCs detached from the primary tumor – initially

seeding locally within the peritoneal cavity before disseminating to distant organs[3].

Different from immunologically cold solid tumors, malignant ovarian cancer ascites is

heavily enriched with immune cells and acellular factors known to influence tumor

progression. Consequently, ovarian cancer DTCs - continuously shed from the primary

tumor - must overcome immune surveillance and adapt to the unique intraperitoneal

microenvironment to establish colonization[4,5,6,7]. Notably, initial and subsequent DTC

colonization is not merely a passive adhesion to a distant organ, but an active

remodeling of the surrounding microenvironment, leading to a consistently evolving

post-seeding metastatic niche (PSMN) – a concept analogous to the pre-metastatic

niche. However, the underlying molecular determinants that modulate reciprocal

interplay between DTC seeding waves and PSMN evolution and drive the ultimate

metastasis outgrowth remains largely unknown.

Here, we conducted comprehensive lineage tracing, single-cell RNA sequencing

(scRNA-seq), *in vivo* CRISPR screening, and Cre-based niche-tracing analysis to

dissect the complex interplay between continuously arriving DTCs and PSMN. We

identified that early DTCs in ascites are dependent on anti-apoptotic niche-adaptive

IFNγ response genes for omental colonization. Additionally, the interferon-rich ascites

microenvironment harbors a macrophage population capable of protecting cancer cells

from anoikis-mediated death via paracrine signaling. Together, our data provide novel

insights into the temporal dynamics of ovarian cancer clonal dynamics and highlight

ascites-specific factors that prime cancer cells for successful metastatic colonization.

RESULTS

MetTag Captures Clonal Lineage Across Metastatic Sites and Across Time

To capture the transcriptional dynamics across ovarian cancer metastasis, we created a

"MetTag" tracing system by employing RNA-level barcoding based on the LARRY

(Lineage And RNA Recovery) barcoding construct, enabling single cell transcriptome-

level lineage tracing[8]. Importantly, we added a unique batch identifier "BC.ID" as a time

stamp for distinguishing time points (TP) of early or later-arriving DTCs. Within the U6-

driven MetTag sequence, a Chromium 10x capture sequence 1 (CS1) allows for

simultaneous capture of the MetTag transcript with mRNA transcripts (**Fig.1A, S1A-D**).

Diversity of each BC.ID library batch was confirmed by deep next-generation

sequencing (NGS). We observed a relatively consistent diversity across the six BC.ID

library batches comparable with the original depositor LARRY library (**Fig.S1B-D**).

Next, we generated six unique isogenic ID8 p53$^{-/-}$ murine ovarian cancer pools (BC.ID1-

6) with MetTag-GFP lentiviral libraries harboring BC.IDs (**Fig.1A**). To model the

continuous dissemination, individually LARRY barcoded (BC) and BC.ID batch-coded

cells were injected interperitoneally (i.p.) in a fractionated manner at low doses across

six different time points to model early, mid, and late waves of DTCs. Early DTCs refer

to the first two BC.ID batches and late DTCs represent the last two DTC batches

(**Fig.1B**). Four weeks after the final DTC exposure, ascitic fluid (Asc), omenta (Om), and

peritoneal fat pads (FPs) were extracted. This experimental scheme was performed

under several immune cell depletion conditions, including wild type C57BL/6J (WT)

mice, immunocompromised NOD.Cg-Rag1^{tm1Mom}IL2rg^{tm1Wjl}/SzJ (NRG) mice (lacking

functional T, B, and NK cells), and under conditions of macrophage (MAC) depletion via

chlodrosome administration. Temporally spaced DTC exposure resulted in robust

metastasis formation in fully immunocompetent mice, evidenced by ascites

accumulation and positive GFP signal in the omentum and peritoneal adipose tissue

(**Fig.S1E top**). Quantification of overall metastases burden in a separate cohort of ID8-

Luciferase injected mice indicated that lacking T, B, and NK cells lead to clearer ascites

and reduced omental weight (**Fig.S1E-F**). Moreover, chlodrosome-mediated

macrophage depletion in C57B6 mice resulted in reduced intraperitoneal metastases

burden under the fractionated injection schema (**Fig.S1G-H**), suggesting the potential

role of myeloid cell in ovarian cancer metastasis.

To dissect the relative contribution of each batch of DTCs continuous seeded at
intraperitoneal metastatic niches, we used MetTag amplicon DNA sequencing to
quantify the relative abundance of BC.IDs in final overt metastases: early timepoint
(*TP-1, TP-2*), mid (*TP-3, TP-4*), and late (*TP-5, TP-6*) arrivers (**Fig. 1B**, pie chart).
Interestingly, in the overt metastases collected from three distinct peritoneal niches
(omentum, ascites, and fat pad), we observed a consistent pattern of enrichment of
early BC.IDs, suggesting dominance of early arrivers in contributing to final metastasis
outcomes. Interestingly, we observed a near identical BC.ID distribution across all three
anatomical locations in both wild type and NRG mice (**Fig.1C-1E**), as well as
macrophage depleted host in omenta (**Fig.1C**), suggesting a negligible role of niche-
specific tumor immune microenvironment (TIME) in the overall dominance of early
arriving DTCs. To examine if later-arriving cancer cells are dispensable for overall
metastases burden, we evaluated tumor burden in mice with a single injection of tumor
cells (TP-1 only) compared to mice with fractionated dosing (TP1-6) as described in
Fig.1B in WT mice with luciferase-tagged ID8 p53$^{-/-}$ cells. At the experimental endpoint
of four weeks post final cancer cell injection, tumor burden levels were comparable
(**Fig.1F-G**), as measured by *in vivo* bioluminescent imaging (BLI). This data indicates
that early DTCs are not dependent on later DTCs for metastatic colonization and
outgrowth. Given the isogenic nature of disseminating ID8 p53-/- cancer cells and their
consistent genomic alterations, these results strongly suggest that early DTC
colonization reciprocally reshapes the PSMN, creating a less favorable tissue
environment for subsequent metastatic seeding.
144
To gain a better understanding of the transcriptional programs necessary for early DTCs
to colonize and subsequently assess their clonal expansion status, we performed CITE-
seq (scRNA-seq with HTO sample multiplexing) coupled with direct capture of U6-driven
MetTag transcripts via capture sequence 1 (CS1) (**Fig. 1H**). Sequential injections were
performed in the same manner as in Fig.1B with a reverse barcode injection schema to
ensure no unintended bias towards certain BC.IDs gained over the cell selection period.
Single cell analysis of MetTag BC.IDs of metastasis-bearing tissues revealed that early
DTCs (*TP1-2*) were enriched in overt metastases (**Fig.1I & 1J**). The prominent trend

towards early DTC dominance, especially in ascites (**Fig.1J**), was consistent with gDNA
sequencing data in Fig. 1C and 1D.

We next asked whether the dominance of early DTCs in overt metastases represents a
widespread niche-adaptive and PSMN conditioning feature of DTCs. We applied three
MetTag-BC.IDs in a triple negative breast cancer (TNBC) mouse model of lung
metastasis (Lung.Met): BC.ID1 representing early TP-1 and TP2, BC.ID2 representing
mid TP-3 and TP4, and BC.ID6 representing the late TP-5 and TP-6. In this model, we
identified almost exclusively early TP DTCs dominance (91.9%) in the overt lung
metastases, on the scRNA-seq level at experimental endpoint (**Fig. 1K**). Collectively,
this data suggests that early dissemination occurs in both hematogenous (TNBC) and
transcoelomic (ovarian) metastases.

Clonal Transcriptomics Analysis of DTCs Reveals Shared Evolutionary Path
Towards Inflammatory Signature
To further investigate characteristics of early and later arrived DTCs, we assessed
clonal dynamics in different metastatic niches. Specifically, as most DTCs do not survive
and only rare clones successfully adapt to metastatic niches[9], we examined whether
tumor cells disseminated over time evolve along diverse trajectories or converge on a
common path characterized by a limited set of adaptive transcriptome features. We first
integrated LARRY clonal barcode information with transcriptional profiles across ascites
and omental niches, followed by cell cluster annotation. We defined cells with high
levels of keratins and absence of CD45 transcript expression (*Krt18+*, *Krt8+*, *Ptprc-*) as
the tumor cell-enriched cluster on the UMAP (**Fig. 2A and S2A**). After reclustering
Krt18+/Krt8+/Ptprc- cells, cell cluster states were determined through a combination of
cluster marker genes and pathway-level analysis using the Escape R package[10] (**Fig.**
S2B-E). Ascites-derived cancer cells (AscMet) displayed distinct transcriptional states
from those in the omentum (OmMet) (**Fig. 2B-D**). Single Cell Gene Set Enrichment
Analysis (ssGSEA) further showed that ascites-derived cells were strongly enriched for
interferon alpha (IFNα) and gamma (IFNγ) signaling (cluster 3), as opposed to more
heterogeneous four distinct subclusters/modules in OmMet including proliferative,

epithelial-mesenchymal transitioning (EMT), matrisome, and a hypoxic, inflammatory

subclusters (**Fig. 2D & 2E**).

Cellular barcoding approaches have been widely used to reveal clonal dynamics driven

by both genetic predisposition and epigenetic changes in response to environmental

cues[11]. We asked whether the most expanded clones would 1) cluster together based

on transcriptional status, indicating strong genetically pre-determined cell-autonomous

programs driving metastasis, or 2) disperse into heterogeneous transcriptional states

(clusters) reflective of diverse transcriptional adaptation to different metastatic niche

cues (**Fig. 2F**). Interestingly, we observed minimal overlap between dominant

outgrowing clones across each HTO/mouse in both anatomical locations (**Fig. 2G**).

While one clone, BC-1, was shared across ascites and omenta, most of the highly

expanded clones are distinct between metastatic niches. These results suggest that

successful clonal outgrowth requires unique adaptations reflective of the specific

microenvironment.

Given that omentum metastases showed diverse single-cell transcriptional states (**Fig.**

2D-E), we projected clonal identity of the top expanded clones from omenta onto the

cancer cell UMAP and examined similarities between clones. At both the transcriptome

cluster and pathway levels, dominant clones were enriched for several defined

signatures (**Fig. 2H-2I, S2F**). Apart from the proliferative subcluster 4, expanded clones

consistently occupied the hypoxic, inflammatory subcluster 0 (**Fig. 2I**), suggesting

diverse, yet directed evolutionary trajectories within the same clonal lineage. To

understand the lineage relationships among different subclusters, we applied RNA

velocity and pseudotime analysis using scVelo and CellRank[12]. Paga velocity and

pseudotime analyses revealed a clear trajectory originating from the IFN signaling

enriched ascites compartment subcluster 3, leading to proliferative status, then further

converging to subcluster 0, a terminal inflammatory state in the omentum (**Fig. 2J & 2K,**

Fig. S2G top). Notably, IFNγ, but not the IFNα signature, was significantly enriched in

early DTCs (TP-1) (**Fig. 2L, S2G** bottom).

Similarly to our ovarian cancer dataset, we observed heterogeneous clonal distribution

between each mouse in the TNBC Lung.Met MetTag scRNA-seq dataset, suggesting

non-genetic, transcriptomic drivers of successful colonization (**Fig. S2H-S2I**). To

examine the generality of the enriched IFN signature in metastases, we performed a

separate scRNA-seq experiment comparing TNBC Lung.Met and in vitro cell culture

(**Fig. S3A**). scRNA-seq analyses revealed that, compared to cultured cells, *in vivo*

Lung.Met tumor expressed higher IFNγ and IFNα signatures (**Fig. S3B-S3E**).

Collectively, the ovarian and lung metastasis data suggests that transcriptional

adaptation towards IFN-signaling *in vivo* is a common trait of successfully colonized

DTCs. In ovarian metastasis specifically, successful metastatic clones in the omentum

occupy heterogeneous transcriptional states. These evolved phenotypes originate from

the ascites compartment, which encompasses an IFN-signaling-rich module that may

prime these later adaptations.

***In vivo* CRISPR screen identifies functional essentiality of IFNγ signaling in**

omental colonization

As ovarian cancer cells initially disseminate into the peritoneal cavity, we next asked

whether transcriptional adaptation of cancer cells to the ascites environment is essential

for subsequent intraperitoneal seeding and successful omental metastasis. We

hypothesized that the ascites gene signature adopted by early DTCs – characterized by

limited nutrient availability and low attachment – is (1) induced by initial detachment

from a primary tumor and (2) necessary for subsequent peritoneal metastasis. To test

this hypothesis, we performed differential gene expression (DEG) analysis and over-

representation pathway analysis (ORA) with clusterProfiler[13] to compare gene

expression in metastatic tumor cells residing in the two distinct niche compartments

(Asc vs Om). Compared to OmMet, DEGs of AscMet were enriched in hemoglobin

genes (*Hbb-bs, Hba-a1, Hba-a2, Hbb-bt*) and interferon-stimulated genes (*Ifi27l2a* and

Irf7). In contrast, OmMets were enriched in immediate-early response genes (*Atf3,*

Egr1, Fos, Fosb, Jun)[14,15], adaptive immune immunoglobulin components (*Igkc, Igha,*

Jchain)[16], and angiogenesis (*Vegfa*) (**Fig. 3A**). Next, IFNγ/IFNα module scoring and

pathway analysis further highlighted a dominant interferon-rich inflammatory signature

in AscMet, as opposed to more heterogeneous programs in OmMet (**Fig. 3B, Fig. S3F-**
S3H).

To assess the functional significance of AscMet-OmMet DEGs, we designed a custom *in*
vivo CRISPR/Cas9 screen targeting 100 DEGs identified (**Fig. 3C,** "In Vivo Ovarian
Metastatic Niche Signature Screen"). SpCas9 was introduced into ID8 p53$^{-/-}$ cancer
cells and validated (**Fig. S3I & S3J**). 2.4 million cells (~2000X representation of sgRNA
library) were i.p. injected to a whole-body Cas9-expressing host (**Fig. 3C**). This host has
central tolerance to Cas9, decreasing immunogenicity towards Cas9-expressing tumor
cells[17]. Six weeks post-injection, metastasis-bearing omenta and ascites were
harvested for gRNA target amplicon sequencing and analysis (**Fig. 3C**)[18]. The positive
controls for the screen, *Pten* gRNAs, were significantly enriched in metastases derived
from both anatomical locations (**Fig. 3D & 3E**). Compared to the initial injected cell pool
sgRNA distribution, sgRNAs targeting eight genes in OmMet and ten genes in AscMet
were significantly depleted ($p < 0.05$) (**Fig. 3D-E**). The top depleted (dropout) genes –
CD300lb, *Ms4a8a*, *Marco*, *Gbp2b* (OmMet), and *Slfn1* (AscMet) – were "upregulated"
DEGs associated with AscMet, suggesting a necessity of initial transcriptome adaptation
to the ascites niche - the first niche DTCs encounter - for subsequent metastasis.
Furthermore, GSEA analysis of dropout genes using the GO Biological Function
(GOBP) gene sets revealed a shared enrichment of apoptotic cell clearance processes
in both Asc and Om compartments and enrichment of programs related to IFNγ in
omentum (**Fig. 3F**). To experimentally validate whether IFNγ is required for the
transcriptional upregulation of CRISPR screen hits in cancer cells, we treated ID8 p53$^{-/-}$
cells *in vitro* with recombinant IFNγ. We tested the inducibility of *Marco*, *Gbp2b*, and
Slfn1 by IFNγ via RT-qPCR, where *Stat1* and *Irf1* transcription factors served as positive
controls for IFNγ response[19] (**Fig. 3G**). By 24 hours of treatment, all three genes were
significantly induced by IFNγ (**Fig. 3G**). The protein level of Gbp2b, an interferon-
inducible GTPase, was also significantly induced by IFNγ (**Fig. 3H**). Collectively, despite
the transcriptional heterogeneity revealed by scRNA-seq, consistent enrichment of
apoptotic pathway and the sequential activation of an IFNγ response (first in ascites,

then required for omental metastasis) suggest two functionally essential adaptive

mechanisms employed by DTCs to survive.

Loss of IFNγ response in DTCs significantly decreases peritoneal metastases and

extends overall survival

Because the top hits from our CRISPR screen are IFNγ-inducible genes, we

hypothesized that IFNγ may function as a primary regulator that exerts broader pro-

metastatic adaptive cellular programs in DTCs. Thus, instead of focusing on a single

downstream gene target, we sought to disable the entire cascade via IFNG receptor 1

knockout. We first confirmed robust expression of IFNγ receptor 1, *Ifngr1,* in both

AscMet and OmMet in vivo (**Fig. S3K**). Next, we genetically knocked out *Ifngr1* in ID8

p53$^{-/-}$ cells. Successful knockout of *Ifngr1* was validated via flow cytometry (**Fig. S3L**)

and ablation of the IFNγ transcriptional downstream targets was validated via western

blot probing for Stat1 and Gbp2b (**Fig. S3M**). We next i.p. injected ID8.Luciferase-

sgNTC or ID8.Luciferase-sgIfngr1 to Cas9-expressing mice. Tumor burden was

monitored by BLI intensity throughout the 6-week experimental window (**Fig. 3I**).

sgIfngr1 groups showed a significant reduction of overall metastasis burden compared

with sgNTC group (**Fig. 3J, Fig. S3N**). In contrast to sgNTC group, ~80% of mice in the

sgIfngr1 group remained alive at the end of the survival study (p = 0.004) (**Fig. 3K**). This

notion is consistent with the observation in human ovarian cancer microarray gene

expression data[20]. Patients with ovarian cancer expressing lower levels of *IFNGR1*

exhibited longer progression-free survival compared to the higher expressing group

(**Fig. S3O**).

Above results demonstrated that DTCs' intrinsic IFNγ signaling is essential for the

overall metastatic outcome. Given early DTCs (TP-1) showed a significantly higher IFNγ

signaling comparing with later DTCs (**Fig. 2L**), we next investigated whether the IFNγ

response in early DTCs influences the success of subsequent waves of DTC

colonization. We specifically perturbed IFNγ response (sgIfngr1) in the first batch of

early DTCs (G2), but not the immediate second wave of DTCs (sgNTC2) (**Fig. 3L**).

Interestingly, loss of Ifngr1 in the first batch of DTCs significantly reduced overall

metastasis burden at the endpoint (**Fig. 3M**). In the residual metastases in the sgIfngr1

group (G2), control (sgNTC2) cells outcompete the sgIfngr1 cells, as quantified through

sgRNA amplicon sequencing (**Fig. 3N**). Taken together, these results demonstrated that

(1) a robust IFNγ response in the initial seeded DTCs prevent the colonization of later

DTCs - potentially through conditioning of PSMN - and (2) are paradoxically required for

overall metastasis success.

IFNγ response mediates resistance to anoikis by activating pro-survival programs

We next sought to examine mechanisms by which IFNγ signaling promotes metastasis.

First, we observed that *Ifngr1* knockout did not alter cancer cell fitness under standard

adherent culture conditions in the presence or absence of IFNγ treatment, evidenced by

both CellTiter-Glo luminescent cell viability assay (**Fig. 4A**, left) and colony count

formation (**Fig. S4A**). Interestingly, IFNγ treatment had a protective effect and increased

cell viability in control (sgNTC) cells under ultra-low attachment (ULA) conditions. This

protective effect was ablated in the *sgIfngr1* group (**Fig. 4A**, right).

To investigate the mechanism underlying increased cancer cell survival under IFNγ

treatment and ULA conditions, we performed bulk RNA-seq (**Fig. S4B**) with wild type

(sgNTC) and single clone knockout line (sgIfngr1). First, principal component analysis

(PCA) and MA dispersion plot showed that ULA culture alone induced significant

transcriptomic shifts in both sgNTC and sgIfngr1 cells (**Fig. S4C-D**). ULA culture

condition induced pathways associated with TGFβ and platelet functions in sgNTC cells

(Fig. S4E). sgNTC cells that survived under ULA conditions showed an overall reduced

S/G2 signature (**Fig. S4F**). Next, we compared the transcriptomic changes in response

to IFNγ. PCA of all groups revealed variance between sgNTC cells treated with PBS and

IFNγ (**Fig. S4C**). This IFNγ induced variance was ablated in the sgIfngr1 (KO) line as

expected, independent of culture conditions (**Fig. S4C**). We analyzed differentially

expressed genes (DEGs) between cells with or without IFNγ treatment under both

regular and ULA conditions (**Fig. 4B**). Interestingly, despite lower proliferation, cells

grown under ULA condition exhibited transcriptionally diverse responses (**Fig. 4B**) and

were overall more responsive to IFNγ than under regular condition, rendering more

DEGs detected (**Fig. 4C and S4C**). Besides 25 common IFNγ responsive genes, we

discovered 30 IFNγ responsive genes uniquely up-regulated only under ULA condition

(**Fig. 4C & S4G**). We next grouped the unique 30 upregulated genes into broad classes.

As expected, most genes represented classical interferon stimulated genes (ISGs) with

known roles in viral defense and antigen presentation (**Fig. 4D**). In addition, we

discovered several non-canonical ISGs, such as the immunoproteasome 20S subunit

beta 10 (*Psmb10*), poly(ADP-ribose) polymerase family member 14 (*Parp14*), as well as

PD-L1 (*Cd274*), in response to IFNγ under ULA condition (**Fig. 4D**). Both the canonical

and non-canonical ISG signatures were reflected in pathway analysis where the most

significant Reactome pathways were related to antigen processing and presentation, as

well as proteasome assembly and endosomal trafficking (**Fig. 4E**). Importantly,

Psmb10, Parp14, and PD-L1 all have reported context dependent, cell autonomous pro-

survival functions[21–24]. Indeed, knocking out IFNGR1 downstream *Stat1* and *Irf1* ablated

IFN downstream signaling (**Fig. S4H**) and led to a loss of IFNγ-induced pro-survival

phenotype under ULA (**Fig. 4F-G**).

To validate *in vivo* relevance of our findings, we analyzed above DEGs detected in bulk

RNA-seq using our single cell RNA-seq data derived from ovarian metastases. *In vivo*

AscMet exhibited higher levels of immunoproteasome genes (*Psmb10*, *Psmb9*) as well

as poly(ADP-ribose) polymerase family members (*Parp12*, *Parp14*, and *Parp9*)

compared with OmMet (**Fig. 4H**). Unexpectedly, *Cd274* was more highly expressed in

OmMet (**Fig. 4H**). Next, we performed RNA-based gating and defined *Irf1*high and *Irf1*low

cancer cells to represent IFNγ downstream responsive levels of each cell (**Fig. 4I**).

*Irf1*high cells upregulated *Parp14*, but not *Cd274* and *Psmb10* (**Fig. 4J**). Parp14 is known

to inhibit apoptosis by regulating Jun N-terminal kinase 1 (JNK1) in multiple myeloma[23].

Moreover, in ovarian cancer microarray data, low PARP14 expression correlates with

prolonged progression free survival, particularly at earlier primary tumor stages (**Fig.**

S4I).

Myeloid cells are the major interacting partners to tumor cells

In ascites, DTC's IFNγ response under low attachment condition is likely influenced by

the crosstalk with ascites niche-specific cells. To identify IFNγ producers and responders

within the ovarian metastatic niche, we first examined the expression patterns of

IFNγ and IFN-responsive genes identified in our CRISPR screen across all cells

detected in our MetTag single cell dataset (**Fig. 5A, S5A & S5B**). While T/NK cell

populations are the main producers of IFNγ, a significant proportion of myeloid cells

expressed a high level of CRISPR hits detected in tumor cells (*Gbp2b*, *Marco*, *Ms4a8a*,

CD300lb, and *Slfn1*) (**Fig. 5A**), suggesting myeloid cell similarly respond to available

IFNγ in the metastatic niche.

CellChat[25] analysis identified a significant increase in both the number and strength of

cell-cell interactions under OmMet condition compared with naïve condition (**Fig. S5C**).

Specifically, tumor cells primarily interacted with myeloid cells in both ascites in

omentum, while interactions with stromal cells were evident in the metastatic omentum

(**Fig. 5B**). These results suggest macrophages may be a key axis of communication that

shapes DTCs' response to both IFNγ and detachment conditions in ascites. Focusing on

the myeloid compartment, we grouped Seurat clusters into broader cell identity

categories (**Fig. S5A-B**) and found a proportional increase of myeloid cells in both

metastatic sites compared to their naïve counterparts (**Fig. 5C**). Subclustering the

myeloid population revealed that the major contribution to the proportional increase are

subclusters 0,1, 5, and 6 (**Fig. 5D, Fig. S5D**). Those highly heterogenous macrophage

subclusters expressed diverse tumor-associated macrophage (TAM) markers, such as

Adgre1, *Cd68*, *Csf1r*, *Itgam*, *Mertk*, and *Trem2* (**Fig. S5E**). RNA velocity analysis

revealed a dynamic transition from the ascites-derived myeloid clusters 1 and 5 toward

omental-specific myeloid clusters 0 and 6 (**Fig. 5E**). Omental clusters 0 and 6 were

characterized by increased expression of costimulatory molecules and angiogenesis

genes as compared to ascites clusters 1 and 5 (**Fig. 5F, Fig. S5F**). Collectively, the RNA

velocity and DEG analyses suggest that ascites-derived TAMs may represent a less

differentiated state relative to omental TAMs.

Given the apparent macrophage niche-dependent differentiation states, we next asked

whether omental TAMs reflect a polarized, terminal state within the metastatic niche. If

so, increased TAM polarization would likely shape the T cell compartment[26,27]. Indeed,

while the ascites-derived T/NK compartment was dominated by a mixture of naïve

CD4+ and CD8 +T cells, omental metastases and niche tissue contained a broader and

more complex repertoire of activated, exhausted, and regulatory T cells (**Fig. 5G**)

enriched in pathways related to regulation of immune cell activation (**Fig. S5G**). We next

subsetted macrophages ($Cd68^{high}$/$Adgre1^{high}$, **Fig. S5H**) and scored them against M1

and M2 gene modules. Compared to ascites-derived TAMs, omental TAMs exhibited

higher hybrid functional state scores (**Fig. S5I**), reflecting an immunomodulatory TAM

cellular state[28] that cannot be captured by binning into a strictly immune-promoting or

suppressing category. Unlike the immunomodulatory and ECM-related omental

signatures, ascites-derived TAMs were enriched in pathways associated with cellular

stress responses and NRF2-dependent redox metabolism (**Fig. 5H & 5I**). On the

individual gene expression level, ascites TAM signatures were reflected in elevated

expression of key stress-related genes – including *Anxa1* and *Hmox1* – as well as

Ndufb8 and *Cox5a*, components of mitochondrial respiration (**Fig. S5J**). These findings

indicate that ascites TAMs may represent an earlier, less polarized yet metabolically

active state, while omental TAMs have transitioned into an immunomodulatory,

polarized phenotype. Finally, we utilized flow cytometry to validate immune cell

dynamics observed in the scRNA-seq dataset (**Fig. S5K**). GFP+ cancer cells were

detected in both metastases-bearing ascites and omental samples. While there was a

significant increase in CD11b+/CD11c+ macrophages as well as CD11b-/CD11c+ DCs

in ascites, metastatic omentum samples were significantly enriched in mature F4/80+

TAMs and cytotoxic CD8+ T cells (**Fig. S5K**). These findings further implicate the

omental environment in fostering a highly immunomodulatory population of TAMs.

Peritoneal macrophages are direct DTC pro-survival interactors

We first tested whether macrophages are functionally important in promoting ovarian

cancer metastasis. Macrophage depletion via direct i.p. liposomal chlodronate injections

(**Fig. 6A**) demonstrated that peritoneal TAMs are pro-metastatic: depleting TAMs

significantly reduced omental weights and CD45 negative cancer cell frequencies in

ascites at experimental endpoint (**Fig. 6A**, **Fig. SAB** flow gating strategy and depletion

validation). Although TAM-induced immunomodulation is a well-established metastasis

driver, less is known about the role of direct TAM-tumor cell interactions in ovarian

cancer metastasis. To identify cells that are in close contact or has been contacted with

ovarian metastases, we developed a non-SP-TATK Cre-based iteration of the

"NicheTracing" system to genetically trace microenvironmental contacts *in vivo* over a

prolonged timeframe. In this system, cancer cells expressing Cre.ERT2 were injected

into Ai6 (RCL-zsGreen) Cre reporter mice. Upon intercellular-cytoplasmic transfer and

Tamoxifen-mediated nuclear entry of Cre into neighboring cells, Cre-mediated

recombination turns on the zsGreen cassette in host cells (**Fig. 6B**). After validating

Cre.ERT2 expression in transduced cell lines (**Fig. S6A**), we benchmarked this system

using *in vitro* coculture (**Fig. S6B-S6F**) and in an independent model of TNBC lung

metastasis established using the E0771 cell line (**Fig. S6G-H**). In the ovarian metastasis

model, we readily detected zsGreen+ (ZsG) niche cells in both metastasis-bearing

omenta and ascites at the experimental endpoint six weeks post cancer cell injection

(**Fig. 6B**). We performed flow cytometry analysis on ascites and gated cells into ZsG^{low}

and ZsG^{high} (**Fig. 6C**). Unlike ZsG cells detected in lung metastases where only ~30%

ZsG^{high} cells were $F4/80^+$ macrophages (**Fig. S6H**, right panel), over 90% of ZsG^{high}

host cells were $F4/80^+/Ly6G^{low}$ TAMs (**Fig. 6C**).

Interestingly, a significant portion of ZsG^{high} $F4/80^+/Ly6G^{low}$ TAMs expressed a specific

macrophage marker Trem2 (**Fig. 6D, Fig. S6I-J**). To explore the transcriptome

differences between $Trem2^{high}$ and $Trem2^{low}$ myeloid cells, we specifically analyzed

those two groups of myeloid cells using our scRNA-seq dataset. $Trem2^{high}$ showed a

high expression of Olfml3, Tmem119, Mmmp13 genes, reflecting broad chemotaxis,

ERK pathway, and adhesion pathways (**Fig. 6E and Fig. S6L**). In contrast, significantly

enriched pathways in $Trem2^{low}$ cell were related to epithelial cell proliferation,

mesenchymal transition, and TGFβ pathways (**Fig. S6L**), demonstrating distinct roles of

these macrophage subsets. As Trem2+ macrophages are known to play lipid sensing,

anti-inflammatory and immunosuppressive tumor promoting roles[29,30,31], we next

investigated the functional significance of Trem2 signaling in ovarian cancer metastasis.

After induction of experimental ovarian metastasis in macrophage-specific (Lyz2-driven)

Trem2 KO mice (**Fig. 6F and Fig. S6M**), unexpectedly, we observed no significant effect

of the Trem2 KO on ovarian metastasis burden (**Fig. S6M**). As ablating Trem2 signaling

in macrophages was not sufficient to change the overall ovarian cancer metastasis

progression, these data suggest that peritoneal macrophages, even Trem2 low subsets,

could promote metastasis independently from known Trem2-mediated efferocytotic and

anti-inflammatory programs.

We next explored whether peritoneal macrophages could promote cancer cell survival

under low attachment conditions. We performed a macrophage-cancer cell coculture

under ULA conditions. Macrophages from peritoneal lavage were enriched using F4/80

magnetic beads (**Fig. 6H**, **Fig. S7B & S7C** flow gating strategy) and cocultured with

three different murine ovarian cancer cell lines; ID8 p53$^{-/-}$, ID8 p53$^{-/-}$ BRCA2$^{-/-}$, and

PPNM[32] (**Fig. 6I, S7D** flow gating strategy). After a 48-hour coculture period, flow

cytometry analysis revealed that cancer cell viability, as measured by live/dead staining

(ZombieAqua), decreased under ULA conditions compared to regular attachment

plates. Most importantly, survival was partially rescued with macrophage coculture

across all three cell lines (**Fig. 6I**). These results indicate that peritoneal macrophages

directly support cancer cell survival under low attachment conditions.

Patient and murine ascites-derived TAMs provide TNF to cancer cells

To determine if tumor-supportive macrophage interactions occur in human ovarian

cancer, we analyzed a human-derived single-cell dataset containing over 200,000 single

cells from 14 ovarian cancer patients and collected from five locations: peripheral blood

(PB), peripheral lymph nodes (PLN), malignant ascites (Asc), primary tumor (Pri.OT),

and matched metastatic tumor (Met.Ome)[33]. As observed in our murine scRNA-seq

dataset, cancer cell-containing compartments, including Asc/Pri.OT/Met.Ome, had

increased proportions of macrophages compared to PB and PLN (**Fig. 6J & Fig. S8A**).

DEG analysis between TAMs at metastatic sites (**Fig. S8B**) and subsequent GO

pathway analysis conducted on the DEG list revealed immune-modulatory pathways

(cytokine, chemokine, allograft rejection) were upregulated in Om.TAMs while cargo

receptor activity related genes were enriched in As.TAMs (**Fig. S8C**).

Our macrophage-tumor co-culture assay (**Fig. 6H-I**) suggests that macrophages could

support tumor cell through macrophage-derived factors. Therefore, we applied NicheNet

analysis to further characterize ligand-receptor interactions between macrophage

(ligand provider) and tumor cells (signaling receiver through receptor) (**Fig. 6K**). We

implemented this strategy to both our murine dataset as well as the patient dataset (**Fig.**

S8E). After defining myeloid cells (Broad cell ID) as the sender cell type in the murine

dataset and macrophage cluster as the sender in the human dataset, we compared the

top predicted tumor-interacting ligands in each dataset. We found four overlapping

sender ligands across species - *TGFB1*, *IL10*, *TNF, and IL-1β-* that were the most

significant ligands predicted to interact with cancer cell receptors (**Fig. 6K, Fig. S8D-E**).

Notably, these ligands were detected at a high-level in patient macrophages (**Fig. S8F**).

Next, we examined the top predicted downstream target genes in cancer cells affected

by top TAM ligands. Pathway analysis of the predicted downstream genes yielded

pathways involved in cell-cell adhesion, response to oxygen levels, and apoptosis

regulation (**Fig. S8G**). For the common ligands (*TGFB1, IL10, TNF, and IL-1β*) we

derived a list of possible corresponding receptors (common receptors) based on "prior

interaction potential" in the NicheNet analysis. Interestingly, cross-examination of

corresponding common receptor expression in the bulk RNA-seq data derived from ULA

culture condition (**Fig 4**) showed that a subset for receptors were significantly up-

regulated in response to ULA culture condition (e.g. *Tgfbr1/2*, *Il1r1*, and *Il10rb*). Such

ULA-dependent receptor up-regulation was largely independent of IFNγ treatment (**Fig.**

6L). In parallel to cancer cell intrinsic IFNγ programs, these data suggest that ascites-

derived TAMs provide a cocktail of ligands that provide additional pro-survival signals to

IFNγ-primed AscMet cells to survive.

DISCUSSION

Early DTC dominance phenotype in ovarian cancer metastasis

DTC-niche interactions at each stage of the metastatic cascade fundamentally determine disease progression and patient outcomes. Following the initial wave of metastatic dissemination, early DTCs inevitably reshape the metastatic niche microenvironment, irrespective of whether they ultimately survive or perish upon arrival. The niche remodeling by pioneer DTCs likely exerts more profound effects on subsequent colonization than distant signals from the primary tumor. While the pre-metastatic niche (PMN) has been extensively characterized[34,35], this dynamic process following DTC seeding creates what we term the Post-Seeding Metastatic Niche (PSMN). The multi-cellular dynamics resulting from progressive DTC accumulation at newly colonized PSMN mirrors a well-established ecological concept: the Allee effect, which describes how population growth rates depend on population density through competition or cooperation. Although the Allee effect has been demonstrated in metastatic contexts using in vitro models[36], how continuously arriving DTCs dynamically alter PSMN properties and influence the fate of later-arriving tumor cells within the intact physiology of living systems remains unstudied.

Leveraging our MetTag barcoding and staggered injection model, the data presented here consistently demonstrated a skewed overrepresentation of early disseminated tumor cells (DTCs): TP-1 and TP-2 DTCs overwhelmingly dominate the metastatic burden at day 40, whereas cells introduced at later days (TP-6) showed minimal contributions to final overt metastases. This pattern persists across immunocompetent, immunodeficient (NRG), and macrophage-depleted hosts, as well as TNBC lung metastasis model (**Fig. 1**). This skewness in early DTC dominance cannot simply be attributed to inherent, longer chronological growth times of early DTCs, as there remains an ample growth window for TP-6 cells (from Day 11 to Day 40). Additionally, these results are consistent with early cancer cell dissemination suggested by clinical tumor genetics studies in ovarian and breast cancer metastasis[37–40]. There are several non-mutually exclusive hypotheses that could explain early DTC dominance observed in our study. First, the omentum might contain a limited number of permissive micro-niches that favor DTC colonization. Pioneer DTCs might rapidly saturate these micro-niches

and monopolize spatial resources – a "first come, first served" model. Second, the

immune-independent nature of skewed distribution of temporally disseminated DTCs

suggests a role of non-immune stromal elements in dictating colonization success. Early

DTCs may actively remodel the niche, rendering it unfavorable for later DTCs.

Comprehensive spatiotemporally resolved imaging and genomics analyses are required

to further explore these hypotheses.

Duality of IFNγ signaling-driven metastatic adaptation

Beyond the "seed and soil" hypothesis, it has been increasingly appreciated that

metastasis is not merely passive fitness-based selection of DTC clones at the niche, but

also an active epigenetic DTC-niche mutual adaptation process. Our single-cell

transcriptome analysis, coupled with barcoding clonal analysis, provided us with clonal

resolution of the underlying genetic and epigenetic determinants governing DTC

behavior. Interestingly, instead of occupying unique transcriptome clusters - which

would likely suggest distinctive genetic mutational features - the expanded clones form

similar coordinated adaptive "mini-ecosystems" encompassing multifaceted

transcriptome features that converge toward an inflammatory/IFNγ response (**Fig. 2H-**

L). This data aligns with single-cell whole-genome sequencing (scWGS) of treatment

naïve ovarian cancer patients which details the existence of (1) significant clonal

heterogeneity within the same patient and (2) distinct IFNα/IFNγ expressing clones[54].

Our CRISPR/Cas9 *in vivo* screen results (**Fig. 3**) further support the notion that cancer

cell fitness in the peritoneal cavity and subsequent omental metastasis depend on rapid,

transcriptional-level adaptation to IFNγ-responsive programs. Intriguingly, our data

indicate that the early DTCs IFNγ response is required for subsequent metastatic

seeding (**Fig. 3L-N**). Knocking out IFNGR1 in the early batch of DTCs significantly

reduced the colonization success of later DTCs with intact IFNGR1, indicating a critical

temporal window in which early IFNγ responsiveness and active PSMN conditioning by

initial colonized DTCs jointly drive the subsequent metastatic continuum. Our findings

expand the conventional view of clonal fitness by revealing its temporal and niche-

dependent nature, where transient adaptive states at critical windows determine

metastatic success. In this "first come, first served" model, early metastatic DTC–niche

crosstalk through IFNγ sensing and epigenetic adaptation determines metastasis

success by enabling the earliest colonizers to occupy and remodel permissive micro-

niches while preventing later DTCs' entry.

Although the findings presented here support a tumor cell-centric pro-metastatic role of

IFNγ, it is important to note that IFNγ in metastatic niches can act in complex ways. In

some cancers, IFNγ reduces metastasis through activation of effector T and NK cells,

thereby enhancing the effectiveness of immunotherapy[41,42]. Furthermore, in some

contexts, IFNγ can induce apoptosis in regulatory T cells and inhibit angiogenesis[43].

Specifically in ovarian cancer cells, both anti-metastatic and pro-metastatic roles have

been reported[44,45]. IFNγ has been shown to synergize with beta glucan treatment to

reduce peritoneal metastasis and Ifngr1 knockout in host cells leading to a significant

increase in metastasis[46]. For pro-metastatic roles, IFNγ has been previously reported to

upregulate PD-L1 and IL-8 in ovarian cancer cells, leading to increased cell migration

and proliferation[45]. We did not observe any proliferation differences via colony formation

assay *in vitro* (**Fig. S4A**) after a 7-day IFNγ exposure. Given that PD-L1 and Psmb10

levels were not increased in Irf1^{high} cells *in vivo* (**Fig. 4J**), we nominate increased

Parp14 as an active pro-survival cellular adaptation maintained in ascites over time, as

opposed to an immediate physiological response to IFNγ signaling. Based on the data

presented here, we believe that, although both effector immune cells and cancer cells

could be impacted by IFNγ, IFNγ response in cancer cells (intrinsically) leads to an

overall pro-metastatic effect. This effect is an important mechanism driving metastasis

and it should be carefully considered when designing anti-metastasis treatment.

Macrophage plasticity and pro-survival ligands

Cells of the myeloid lineage are characterized by significant plasticity, especially in

disease[47,48]. However, recent studies point to niche-regulated plasticity restriction[49].

Upon infiltration of inflammatory monocytes into metastatic sites, they often exhibit

"commonly described" reprogrammed states such as increased checkpoint expression

or efferocytotic receptor upregulation due to cancer cell engulfment. In addition to these

classical reprogramming events, myeloid cells exhibit tissue-specific reprogramming

after prolonged residence at a specific niche[49]. Thus, although we observe Trem2^high

macrophages in metastatic omenta and ascites, depleting Trem2 signaling in those cells

did not significantly impact on metastasis phenotype (**Fig. 6F-G**), pointing the role of

other programs within the broad myeloid population that could provide support to cancer

cells. Although IFNγ treatment is sufficient to trigger expression of anti-apoptotic genes

such as *Parp14*, and loss of attachment triggers *IL-1β*, *TNF*, *TGFβ*, and *IL10* receptor

transcription (**Fig. 6L**), our data suggests that macrophages provide multiple pro-

survival signals to primed cancer cells. Specifically, TNF itself[50] and *IL-1β*[51] are known

amplifiers of NF-kB, a master-regulator of anti-apoptotic and anti-necrotic gene

expression[52]. TGFβ has been reported to activate the pro-survival P3-k/Akt pathway in

certain contexts[53,54]. The cancer cell-intrinsic roles of IL10 warrant more investigation.

Given the known roles of TNF, *IL-1β*, and TGFβ and increased receptor expression in

cancer cells when cultured under ULA conditions, we hypothesize that initial

IFNγ exposure and loss of attachment may prime DTCs to respond strongly to

macrophage-derived pro-survival cues and collectively amplified pro-survival programs

in DTCs.

In summary, our study dissected the complex landscape of ovarian cancer metastasis

through a systematic characterization of both cancer cells and local immune factors

using MetTag temporal, clonal and scRNA-seq analysis. Our findings highlight the

importance of ascites as a functional reservoir for these factors and that early

acquisition of IFNγ response and interaction with macrophages are necessary

deterministic adaptation steps for DTCs to survive and colonize.

ACKNOWLEDGMENTS

This study was partially funded by the NIH T32GM075762 training grant (E. A.), NIH

R01 CA194697-01 (S.Z.), NIH R01 CA255064-01A1 (S.Z.), NIH R21 CA263798-01

(S.Z.), Cancer Prevention and Research Institute of Texas (CPRIT) Scholar Award

RR220024 (S.Z.), CPRIT Individual Investigator Research Awards RP230261

(Z.Zhong), and NIH R35GM142654 (Z.Zhong). We are additionally grateful to the Dee

Family endowment (S.Z.) and to the following core facilities: Notre Dame Freimann Life

Sciences Center, Notre Dame Genomics and Bioinformatics Core Facility, Indiana

University School of Medicine Center for Medical Genomics, Indiana University Simon

Cancer Center, Simmons Comprehensive Cancer Center, UT Southwestern (UTSW)

Animal Resource Center, UTSW Flow Cytometry Core, and UTSW Small Animal

Imaging Core Facility. The computational arm of this research project was, in part,

supported by the BioHPC supercomputing facility at UTSW.

AUTHOR CONTRIBUTIONS

Conceptualization and experiment design, E.A., M.S.S. and S.Z.; Data analysis, E.A.,

S.Z., X.L., Z. Zhao; Experiments and methodology, E.A., S.Z., S.R.F., S.M.G., J.L.,

W.M., N.D., T.C.R., M.Z., Z. Zhong; Manuscript writing and revision, E.A., S.M.G., and

S.Z.; Study supervision, S.Z. We attest that all coauthors reviewed and pre-approved

the manuscript before submission.

DECLARATION OF INTERESTS

We do not have any competing interests to declare.

FIGURE LEGENDS

Figure 1: Tracking Clonal Lineage in Ovarian Cancer Metastasis

(**A**) Schematic depicting main elements of the MetTag library barcoding construct. (**B**)

Schematic of injection and tissue collection timeline for the staggered injection

experimental model using MetTag-barcoded ID8 ovarian cell lines. Schematic was

partially created with BioRender.com. (**C**) Genomic DNA (gDNA)-level barcode

distributions in the omentum graphed as frequency under WT, NRG, and MAC depletion

conditions. (**D-E**) gDNA-level barcode distributions in ascites and pelvic fat pads under

WT and NRG conditions. In (C-E) panels, BC.ID1 = TP-1, BC.ID2 = TP-2, BC.ID3 = TP-

3, BC.ID4 = TP-4, BC.ID5 = TP-5, BC.ID6 = TP-6 for clarity. (**F**) Injection schema for

comparing one-time (400,000 total, n = 10) cell dose and fractionated (2,400,000 total, n

= 10) cell injections (left) and representative BLI images at endpoint (right). (**G**) Bar

graph depicting total flux emission in each group at experimental endpoint. (**H**) Schematic of scRNA-seq experiment conducted using a reversed BC.ID injection order. LARRY barcodes are readily captured by Chromium 10x Feature Barcoding. (**I**) Pie charts of RNA-level barcode frequencies in each location grouped as early (BC.ID5, BC.ID6), mid (BC.ID4, BC.ID3), and late (BC.ID2, BC.ID1). (**J**) Heatmap of sample level barcode frequencies, BC.ID6 = TP-1, BC.ID5 = TP-2, BC.ID4 = TP-3, BC.ID3 = TP-4, BC.ID2 = TP-5, BC.ID1 = TP-6. (**K**) Schematic (top) and MetTag BC.ID distribution (bottom) for independent TNBC lung metastasis model depicting enrichment of scRNA-seq derived early TP DTCs in the metastatic lung at endpoint. Data in Figure 1G was analyzed using a two-tailed Student's t test and graphed as mean with standard deviation (SD).

Figure 2: Clonal Analysis Reveals Shared Evolutionary Path Towards Inflammatory Signature

(**A**) Split UMAP depicting cluster distributions across the four experimental conditions and identifying cancer cell-enriched clusters in ascites and omentum. (**B**) Reclustered cancer cell UMAP colored by the two metastasis-bearing anatomical locations. (**C**) Stacked bar chart depicting the frequency of Seurat subclusters within each metastatic location. (**D**) Seurat cluster UMAP of reclustered cancer cells with transcriptional program labels. (**E**) Dot plot depicting Hallmark pathways across all subclusters used to define each cluster subset gene program in (D). (**F**) Schematic representation of the possible outcomes of LARRY clonal analysis based on transcriptome patterns of enriched clones. (**G**) Heatmap of normalized clonal LARRY barcode counts across the two anatomical locations and individual mice (HTOs). (**H**) UMAP overlay of top 3 expanded clones in the omentum. (**I**) Bar chart depicting the relative subcluster proportions present in each of the top 3 barcode clones. (**J**) scVelo-generated PAGA velocity graph indicating trajectory arrows and inferred cluster lineage. (**K**) scVelo and CellRank-based simulated path from start to endpoint clusters. (**L**) Violin plot depicting RNA-level contribution of each timepoint to the IFNγ signature. Data in Figure 2L was analyzed by first defining the IFNγ signature using canonical mouse genes, followed by

directly applying the wilcox.test() function in Seurat to compare TP-1 and TP-6

expression.

Figure 3: Activation of IFNγ Response in Ascites is Necessary for Successful

Omental Colonization

(**A**) DEG-based volcano plot of gene expression differences between OmMet and

AscMet, where threshold for significance was defined at $p < 0.05$ and log2FC < -1.0 for

downregulation or >1.0 for upregulation. (**B**) Boxplots comparing IFN signature modules

between AscMet and OmMet. (**C**) Schematic describing in vivo ovarian metastatic niche

signature screen. Partially created with BioRender.com. (**D**) In vivo CRISPR screen

RankView plots for all 100 genes generated using MAGeCKFlute. OmMet (n=5) and

AscMet (n=4) samples were compared to the original pool of ID8 cells. (**E**) Ranks of

individual sgRNAs targeting positive control Pten and depleted genes. (**F**) Lollipop plot

of depleted pathways associated with depleted genes in Fig. 3E. (**G**) IFNγ treatment

time course qPCR plots of select CRISPR hits and positive controls. (**H**) IFNγ treatment

time course western blot probing for Stat1/pStat1 and GBP1. (**I**) Schematic of in vivo

experimental metastasis study comparing sgNTC and sgIFNGR1.g1 groups. (**J**) Time

course log10-transformed total emission/flux values (left). Two independent groups

pooled; cohort 1 NTC; n = 8, KO; n = 7; cohort 2 NTC; n = 5, KO; n = 5. log10-

transformed flux plot for week six with p value (right). (**K**) Kaplan Meier survival curve

generated for cohort 1. (**L**) Experimental schematic for staggered sgIfngr1 (n = 4) or

sgNTC1 (n = 3) followed by sgNTC2-cell injection. (**M**) Representative endpoint BLI

images of metastasis bearing mice in each group (left) and endpoint BLI quantification

comparing tumor burden between sgIfngr1 and sgNTC first injection (right). (**N**)

Heatmap showing enrichment of first DTC seed - either sgNTC1 or sgIfngr1 - and 2nd

DTC seed, sgNTC2. DEG data in Figure 3A was analyzed using the FindMarkers()

function. Data in Figure 3B was analyzed by first defining custom IFN signatures,

followed by applying wilcox.test() in Seurat to compare module scores between AscMet

and OmMet. CRISPR screen related data (3D-3F) was analyzed using MAGeCK and

MAGeCKFlute. Data in 3G, 3J, 3K, and 3M was analyzed and graphed in Prism. 3G,

SJ, and 3M were analyzed using a standard t test; 3G and 3M data was plotted as

mean with SD, while 3J was plotted as median. Survival data in 3K was computed using

a standard log-rank test.

Figure 4: IFNγ Pathway Mediates Resistance to Anoikis and Activates TNF

Signaling and Dormancy Programs

(**A**) Cell-Titer Glo (CTG) assay luminescence values graphed as fold change for regular

culture conditions (left) and ultra-low attachment (ULA) conditions (right), with and

without IFNγ treatment. (**B**) Volcano plots generated from bulk RNA-seq depicting top

DEGs between sgNTC (wild type) cells under regular culture conditions with or without

IFNγ treatment (left) and under ULA conditions with and without IFNγ treatment (right).

(**C**) Venn diagram showing 25 overlapping and 30 unique upregulated genes in sgNTC

cells treated with IFNγ under ULA conditions. (**D**) Heatmap of vst-normalized gene

counts for ULA-specific upregulated genes. (**E**) Bar plot depicting top 15 Reactome

pathways upregulated under ULA conditions in cells treated with IFNγ. (**F-G**) CTG flux

dot plots for Stat1-KO (F) and Irf1-KO (G) cells under ultra-low attachment, with or

without IFNγ treatment. (**H**) Violin plots comparing expression level of

immunoproteasome (Psmb10, Psmb9), DNA repair/apoptosis (Parp12, Parp14, Parp9),

and immune checkpoint (Cd274) levels between OmMet and AscMet. (**I**) scRNA-seq

based gating strategy used to identify Irf1^{high} and Irf1^{low} cells for module scoring. (**J**)

Violin plots comparing expression level of Cd274 (left), Psmb10 (middle), and Parp14

(right) between Irf1^{high} and Irf1^{low} cells. Fold change in F4A was determined by

normalizing each cell line's CTG value to the regular attachment, without IFNγ

treatment. CTG assay luminescence data (4A) was graphed as mean and SD, data in

4F and 4G was graphed as median values. scRNA-seq based transcript expression

levels were compared between Irf1^{high} and Irf1^{low} cells and OmMet/AscMet cells by

applying the wilcox.test() function in Seurat. RNA-seq data was analyzed using DESeq2

and visualized as variance-stabilized counts.

Figure 5: Niche-specific Transcriptional States of Omental and Ascites-Derived

TAMs

(**A**) Violin plots depicting CRISPR screen hits expression level across Broad cell ID

categories (cell types). (**B**) CellChat-generated circus plots of interaction strengths

between Broad cell ID categories in metastatic omenta (left) and ascites (right). (**C**)

Stacked bar chart of cell type proportions across the four experimental groups. (**D**)

UMAPs of reclustered myeloid cells, split by experimental condition. (**E**) UMAP of

subclustered myeloid cells overlayed with scVelo RNA velocity arrows. Arrows are

pointing from ascites TAMs (clusters 1 and 5) to omentum TAMs (clusters 0 and 6). (**F**)

DEG volcano plot comparing gene expression differences between ascites TAM cluster

1 and omentum TAM cluster 0. (**G**) Stacked bar chart of subsetted T/NK cells comparing

cell proportions in ascites and omentum. (**H-I**) Dot plots depicting Reactome pathways

enriched in ascites (H) and omentum-derived TAMs (I). Significance of DEGs in Figure

5F was evaluated by applying the FindMarkers() and Wilcoxon test in Seurat to

compare ascites and omental $Cd68^{high}$/$Adgre1^{high}$ macrophages.

Figure 6: Peritoneal Macrophages are Direct DTC Pro-Survival Interactors

(**A**) Liposomal chlodronate-based macrophage depletion strategy and timeline

schematic with representative tissue images. Omental tumor burden quantification and

flow-based ascites quantification (bottom). Control group; n= 7, and Chlodrosome; n =

7. (**B**) Schematic describing Cre-based NicheTracing labeling system where ID8 and

E0771 cancer cells express Cre.ERT2. Upon nuclear entry of Cre.ERT2 into Ai6

reporter niche cells, labeled niche cells turn ZsGreen positive. (**C**) Representative gating

strategy depicting the major cell population tagged with zsG. (**D**) Quantification of

F4/80+ and TREM2 mean fluorescent intensity (MFI) of contacted, ZsGreen high cells.

(**E**) Volcano plot comparing DEGs between metastasis-infiltrating Trem2-low and

Trem2-high macrophages. (**F**) Schematic depicting mouse model for macrophage-

specific Trem2 knockout. (**G**) Log transformed BLI-derived flux values in Trem2fl/fl (wild

type; n = 5) and Lys2-Cre/Trem2fl/fl (knockout; n = 5) mice at metastasis endpoint. Mice

were injected with ID8 p53-/- Luciferase cells. (**H**) Schematic of macrophage-cancer cell

coculture layout (top) and representative coculture images for one cell line (bottom). (**I**)

Plots of percent CD45 negative cancer cell viability (ZombieAqua negative) post

coculture in three cell lines. (**J**) Stacked bar chart of main cell type proportions captured

in human scRNA-seq dataset across five anatomical locations, using maintype-2 annotation provided by publisher. (**K**) Venn diagram of top NicheNet predicted ligands after filtering for robust expression and overlapping with top ligands in our murine dataset. (**L**) Heatmap comparing enrichment of a curated list of receptors in bulk RNA-seq comparing ULA-cultured cells treated with IFNγ and regular attachment conditions. Data in Figures 6D was plotted as mean with SD and differences between groups were compared via a two-tailed unpaired Student's t test. Data in Figures 6I was similarly analyzed using a Student's t test and plotted with median values. Data in Figure 6E was analyzed using the FindMarkers() function in Seurat and Wilcoxon test to compare gated Trem2-low and Trem2-high cells. RNA-seq heatmap data in 6L was analyzed using DESeq2 and visualized as variance-stabilized counts.

SUPPLEMENTAL INFORMATION (Including Supplemental Figures S1-S9)

Figure S1. MetTag Tracing Construct and Metastatic Model Validation

(A) Schematic depicting MetTag library barcoding plasmid, and modes of detection upon plasmid integration: DNA level, Pol III U6 driven RNA level, and Pol II CROP-seq based. (B-C) Barcode frequencies comparing relative evenness of library distribution across BC.ID batches (B) and original depositor LARRY library (C). (D) Table depicting unique reads detected in each library batch after MiSeq sequencing and read filtering. (E) Representative ascites images and dissecting scope images of GFP+ metastases formed under the repeated injection schema in wild type (n = 4) and NRG mice (n = 4). (F) Separate cohort of mice were subjected to ID8 p53-/- Luciferase injection and omental burden was quantified via ex vivo BLI imaging (top) and by measuring omental weights (bottom). (G) Representative ascites images and dissecting scope images of GFP+ metastases formed under the repeated injection schema in liposome (n = 4) and chlodrosome treated mice (n = 4). (H) Barchart depicting macrophage depletion efficiency (left) and quantification of GFP+ cells (right) in ascites under macrophage. Data in Figures S1H and S1F was analyzed in GraphPad Prism and plotted as mean

with SD and differences between groups were compared via a two-tailed unpaired

Student's t test.

Figure S2. scRNA-seq-based Tumor Cell Identification and Cancer Cell Velocity

(**A**) Feature plot of main marker genes used to define and subset cancer cells. (**B**)

Heatmap of reclustered cancer cell marker genes. (**C**) Violin plots of top defining cluster

marker genes. (**D**) Escape generated Hallmark pathways each represented by a

different color. (**E**) Escape pathway-based violin plots of each pathway class in (D). (**F**)

UMAPs depicting main pathway signatures represented in the top three clonally

expanded barcoded cells. (**G**) CellRank-generated initial and terminal cellular states

overlayed on the cancer cell UMAP (top) and violin plot of IFNα signature score across

the six different BC.ID timepoints (bottom). (**H**) UMAP of captured single cells in TNBC

lung metastasis model (left) and RNA-based gating strategy to subset non-immune cells

for subsequent barcode analysis. (**I**) Heatmap depicting enriched clonal LARRY

barcodes in each lung samples. IFNα signature data in Figure S2G was plotted by first

defining the IFNα signature using Hallmark IFNα response genes, followed by directly

applying the wilcox.test() function in Seurat to compare expression level differences

between TP-1 and TP-6 groups.

Figure S3. IFN Signaling in Metastatic Niches & *In vivo* CRISPR Screening

(A) UMAP of captured cancer cells colored by in vivo (Lung.Met) and in vitro

(E0771.cell) cell line. (B) Hallmark pathway analysis on DEG list comparing in vivo vs. in

vitro. Module scores comparing IFNγ (C) and IFNα (D) genes between in vivo and in

vitro. (E) VlnPlots of individual IFNγ response related genes. (F-G) Dot plots of Hallmark

pathways based on DEGs between AscMet and OmMet. (H) Heatmaps depicting IFN

response gene scoring differences between AscMet and OmMet. (I) Western blot

validation of SpCas9 expression in ID8 p53-/- cells after PiggyBac transposon

transfection and selection with hygromycin. (J) Representative flow cytometry generated

histogram of mCherry fluorescence (left) and bar chart depicting percent mCherry positive cells five days after lenti-guide transduction (right). (K) Violin plot of Ifngr1 expression levels in cancer cells in vivo. (L) Representative flow cytometry generated histogram of IFNGR1 surface expression in knockout vs. wild type cells (left) and bar chart with quantification of IFNGR1 mean fluorescent signal changes (right). (M) Western blot probing for IFNγ downstream transcriptional targets, Stat1 and Gbp2b, in Ifingr1 knockout cells. (N) Endpoint (week six) BLI images of sgIfngr1 (n = 7) compared to sgNTC experiment mice (n = 8). (O) KMPlotter-generated Kaplan Meier progression free survival curve comparing low and high IFNGR1 expression at stage III and stage IV. IFNγ/IFNα signatures in Fig. S3C and S3D were described in previous figures. Data in S3J and S3L was analyzed in GraphPad Prism using a two-tailed Student's t-test and graphed as mean with SD. Figure 3C-D module scores and S3K transcript level violin plot values were compared using the wilcox.test() function in Seurat.

Figure S4. IFNγ-induced Pro-Survival and Dormancy Programs

(A) Clonogenic assays in absence and presence of recombinant IFNγ, with or without IFNGR1 knockout. (B) Schematic depicting in vitro bulk RNA-seq experiment comparing ULA and regular culture conditions under IFNγ treatment. (C) PCA plot of sequenced RNA-seq groups. Each dot represents a different sample; color represents different treatment or culture condition. (D) MA plot showing differentially expressed genes (DEGs) between ULA and regular culture conditions. Blue points indicate significant genes (padj < 0.05, log2FC>+/- 1). (E) Bar plot depicting top Reactome pathways enriched in ULA compared to regular culture conditions. (F) S and G2/M cell cycle gene heatmap comparing sgNTC cells cultured under ULA and regular culture conditions. (G) Heatmap of top DEGs (both unique and overlapped with regular culture) between ULA and regular culture. (H) Western blots showing protein levels of Stat1, Irf1, and Gbp2b under Stat1 (left) or Irf1 (right) knockout, with IFNγ treatment. (I) Kaplan-Meier survival curves for ovarian cancer patients stratified by early stage (left) and late stage (right) and by high and low expression of PARP14. Data in (A) was plotted in GraphPad Prism and analyzed using a Student's t test. The PCA plot was generated from variance-

stabilized transformed (VST) counts. Progression free survival Kaplan-Meier curves

were generated using KMplotter.

Figure S5. Related to Main Figure 5; Additional macrophage and T/NK cell

analysis

(A) UMAP depicting all cells captured across four sequenced groups. (B)

"Broad_cell.ID" classification of Seurat clusters into cell types based on cluster marker

genes. (C) CellChat generated bar charts depicting number of interactions (left) and

interaction strength (right) across the four sample groups. (D) Heatmap of myeloid TAM

subclusters 1, 0, 5, and 6. (E) Violin plots showing gene expression of canonical TAM

markers across myeloid subclusters. (F) Volcano plot of DEGs between myeloid cluster

5 and 6. (G) Dot plot of GO Biological Process pathways upregulated in omentum-

derived T/NK cells based on the DEG list comparing to ascites-derived T/NK cells. (H)

RNA-based gating strategy to subset out macrophages for subsequent analysis. (I)

Biaxial scatter plot of M1 and M2 module scores with overlapped ascites and omental

TAMs. (J) Violin plots comparing individual genes belonging to cellular response to

stress (Anxa1, Hmox1) and cellular respiration (Ndufb8, Cox5a) pathways between

omental and ascites TAMs. (K) Flow cytometry analysis plots of cancer and immune cell

frequencies in naïve and metastatic settings: frequency of GFP+ cancer cells,

CD11b+/CD11c+ double positive myeloids, CD11b-/CD11c+ dendritic cells,

CD11b+/F4/80+ macrophages, and frequency of CD8+ T cells. In (F), DEG analysis

was performed using the FindMarkers() function in Seurat and significance determined

via Wilcoxon test. Fig. S4I hybrid score and S4J violin plot expression levels were

compared using the wilcox.test() function in Seurat. Flow cytometry data in S5K was

analyzed via two-tailed unpaired Student's t test and plotted with median values.

Figure S6. Related to Main Figure 6; NicheTracing Validation & Trem2-knockout

(**A**) Western blot validation of Cre expression in target cell lines: E0771 murine TNBC

line and ID8 ovarian cancer cell line. (**B**) Schematic depicting Cre reporter used to

transfect HEK293T cells and Cre constructs expressed by target cells for in vitro color switch assays. (**C-F**) Images generated after 72 hours of coculture with or without 2 uM 4-OHT treatment. (**G**) Representative dissecting microscope images of negative control spleen and lugs collected 2 weeks post E0771 retro-orbital injection. (**H**) Flow cytometry bar plots at experimental endpoint showing percent switched immune cells (ZsG+) with or without tamoxifen treatment (left) and within the tamoxifen treated group, frequency of ZsG+ macrophages compared to ZsG- (right) in metastatic lungs. (**I**) Bar plot showing frequency of ZsG+ CD11b myeloids compared to ZsG- in ascites. (**J**) Representative flow cytometry gating for TREM2 macrophage enrichment in ZsG+ fraction. (**K**) scRNA-seq gating strategy to subset TREM2-high and low TAMs. (**L**) Go Biological Processes pathways upregulated in Trem2-high (left) and Trem2-low (right) TAMs. (**M**) Macrophage-specific Trem2 depletion validation. Flow cytometry data in (S6H-I) was analyzed using a two-tailed unpaired Student's t test and plotted as mean with SD.

Figure S7. Related to Main Figures 5 & 6; Immunotyping and Flow Cytometry Gating Strategies

(**A**) Flow gating strategy for macrophage depletion study (left) and representative gates for F4/80 macrophages in FMO, liposome control, and chlodrosome treated groups (right). (B-**C**) Flow gating strategy for assessing MHC-II+/CD11b+ macrophage enrichment following F4/80 magnetic bead isolation from peritoneal cavity. Bar plot showing relative enrichment of macrophage populations before the start of coculture. (**D**) Flow gating strategy to assess CD45- cancer cell viability (Zombie Aqua negative populations) after co-culture completion. Flow cytometry data in (S7A) was analyzed via two-tailed unpaired Student's t test and plotted with median values.

Figure S8. Related to Main Figure 6; Patient & Murine NicheNet Analysis

(A) UMAP depicting all (>200,000) single cells collected from five locations and annotated by cell type ("Maintypes-2"). (B) Heatmap of DEGs between Ascites and Met.Ome-derived macrophages. (C) Dot plots depicting top GO molecular function pathways expressed in patient omental TAMs (left) and ascites TAMs (right). (D) NicheNet heatmap depicting top receptor-ligands pairs in murine ascites. (E) NicheNet

heatmap depicting top receptor-ligands pairs in patient ascites. (F) Bubble plot depicting

NicheNet-derived ligand expression among Maintypes-2 clusters. (G) Dot plot of GO

pathways enriched in cancer cells based on regulatory potential and downstream genes

affected by macrophage ligands. DEG analysis in Figure S8B was performed using the

FindMarkers() function in Seurat.

METHODS

Experimental Models

Mice

All animal experiments were conducted according to IACUC protocols which were pre-

approved by the University of Notre Dame and UT Southwestern Medical Center IACUC

committees. Mice were housed with unlimited access to food and water in a

temperature-controlled holding room with a standard 12-hour light/dark cycle. Health

checks were performed daily during *in vivo* experiments. Wild type C57BL/6J (B6) mice

were 8-10 weeks old at the experiment onset. Control and treated transgenic mice were

age-matched, greater than 8 weeks of age. All experiments were carried out exclusively

on female mice.

C57BL/6J (B6, 000664), NOD.Cg-$Rag1^{tm1Mom}Il2rg^{tm1Wjl}$/SzJ (NRG, 007799), B6.Cg-

$Gt(ROSA)^{26Sortm6(CAG-ZsGreen1)Hze}$/J (Ai6, 007906), and B6J.129(Cg)-

$Gt(ROSA)26Sor^{tm1.1(CAG-cas9*,-EGFP)Fezh}$/J (Cas9 026179) mice were obtained from

Jackson Laboratory (Ben Harbor, ME). We generated Lyz2-Cre(+/-)/Trem2(fl/fl) double

transgenic mice by breeding Lyz2-Cre heterozygous (+/-) mice with Trem2(fl/fl). These

lines were generous gifts from the Zhong Lab at UT Southwestern Medical Center.

Cell Lines

The TNBC E0771 cell line (CH3 Biosystems, Amherst, NY; Cat.No. 940001-Vial) was

maintained in RPMI-1640 media supplemented with 10%FBS and 1%

Penicillin/Streptomycin (Pen/Strep). ID8 p53$^{-/-}$, ID8 p53$^{-/-}$ BRCA2$^{-/-}$, and PPNM ovarian

cancer cell lines, provided by the Stack lab at the University of Notre Dame, were

maintained in DMEM low glucose media with 5% FBS, 1% Pen/Strep, and 1% Insulin-

Transferrin-Selenium (ITS). PPNM cell media was additionally supplemented with 100 ng/ml Cholera toxin and 2ng/mL epidermal growth factor (EGF). HEK293T cells were cultured in DMEM high glucose supplemented with 10% FBS, 1% Pen/Strep, and 1% GlutaMAX (Gibco, 35050061). Cell cultures were grown at 37 ºC, 5% CO_2, and 95% humidity. All cell lines were tested and were negative for mycoplasma contamination using a routine mycoplasma PCR kit (ABM).

Establishment and Monitoring of Experimental Metastasis

For the single-injection OvCa model, 4×10^5 ID8-LARRY.GFP or ID8-Luciferase were injected into the peritoneal cavity. To ensure adequate library coverage, 2.4×10^6 ID8-Cas9.Lib tagged cells were injected for CRISPR screening experiments (>1000x coverage). In the repeated injection model, mice received 4×10^5 cells per injection, administered six times over the course of 12 days. Mice were typically sacrificed 6 weeks after initial dose of cancer cells. 6×10^5 E0771-LARRY.GFP or E0771-Luciferase tagged cells were injected into the retro-orbital sinus of mice for one time injection lung metastasis model or 1.5×10^5 cells in six separate injections for the repeated injection model. Mice were sacrificed between 14-18 days in both experimental models. For monitoring metastasis in luciferase-expressing models, 100 ul of D-luciferin (15 mg/mL stock, GoldBio cat#: LUCK-500) was administered intraperitoneally and imaged using an AMI-HTX spectral imaging instrument. The total emission (flux) data for each region of interest (ROI) was obtained after image analysis using the Aura software.

Macrophage Depletion

Macrophage depletion was achieved using encapsulated liposomal chlodronate (Encapsula Nano Sciences, SKU# CLD-8901). Mice received a pre-treatmeant with 200 ul chlodrosome or empty liposome nanoparticles i.p. 24 hours before cancer cell injection. Following pre-treatment, 100 ul of chlodrosome was administered twice weekly throughout the study duration.

Individual Gene Knockout Guide Cloning

The CROP-seq-V2 plasmid (Addgene # 127458) was first modified to include a P2A-

mCherry sequence downstream of the puromycin resistance gene (PuroR) through a

NEBuilder HiFi DNA Assembly (NEB, E5520S) reaction. Then, single-guide RNA

(sgRNA) sequences were designed using CRISPick (Mouse GRCm38 reference

genome), oligos synthesized by IDT with BsmBI overhangs, annealed, and cloned into

the modified CROP-seq-V2 plasmid using the NEBridge Golden Gate Assembly Kit

(NEB, E1602S). The resulting plasmids were first transformed into competent *E. coli*,

plated on LB-ampicilin (AMP) plates, and grown at 30 ºC, before plasmid extraction.

Below are the sgRNA sequences used for gene targeting, including the necessary

BsmBI overhangs:

sgNTC-1 Top oligo

5'- **CAC CG**A CCT GAT ACG TCG TCG CGT A -3'

sgNTC-1 Bottom oligo

5'- **AAA C**TA CGC GAC GAC GTA TCA GGT **C** -3'

sgNTC-2 Top oligo

5'- **CAC CG**G TAT TAC TGA TAT TGG TGG G -3'

sgNTC-2 Bottom oligo

5'- **AAA C**CC CAC CAA TAT CAG TAA TAC **C** -3'

sgIrf1 guide 1 Top oligo

5'- **CAC CG**T GGA AGC ACG CTG CTA AGC A -3'

sgIrf1 guide 1 Bottom oligo

5'- **AAA C**TG CTT AGC AGC GTG CTT CCA **C** -3'

sgIrf1 guide 2 Top oligo

5'- **CAC CG**C TGT AGG TTA TAC AGA TCA G -3'

sgIrf1 guide 2 Bottom oligo

5'- **AAA C**CT GAT CTG TAT AAC CTA CAG **C** -3'

sgStat1 guide 1 Top oligo

5'- **CAC CG**T ATC CTG TGG TAC AAC ATG C -3'

sgStat1 guide 1 Bottom oligo

5'- **AAA C**GC ATG TTG TAC CAC AGG ATA **C** -3'

sgStat1 guide 2 Top oligo

5'- **CAC CG**G CAA GCG TAA TCT CCA GGT A -3'

sgStat1 guide 2 Bottom oligo

5'- **AAA C**TA CCT GGA GAT TAC GCT TGC **C** -3'

sgIfngr1 guide 1 Top oligo

5'- **CAC CG**G GCT CGG AGA GAT TAC CCG A -3'

sgIfngr1 guide 1 Bottom oligo

5'- **AAA C**TC GGG TAA TCT CTC CGA GCC **C** -3'

sgIfngr1 guide 2 Top oligo

5'- **CAC CG**T ATA CCA ATA CGC AAA TAC C -3'

sgIfngr1 guide 2 Bottom oligo

5'- **AAA C**GG TAT TTG CGT ATT GGT ATA **C** -3'

1049

Library Cloning

Six MetTag-LARRY-BC.ID libraries were generated via overhang PCR by amplifying the

region within the original pLARRY-eGFP plasmid containing random barcodes. Primers

were designed with overhangs containing: (1) a unique 6-nucleotide library identifier

(ID), (2) a chromium 10x capture sequence, and (3) BsmBI restriction sites flanking both

ends of the barcode (synthesized by Sigma-Aldrich). Amplified barcode fragments were

introduced into the pRG212 third-generation lentiviral plasmid using the NEB Golden

Gate Assembly Kit (NEB, E1602S) at a 2:1 insert-to-vector molar ratio. The cloning

reaction was transformed into Endura electrocompetent *E. coli* cells (Biosearch

Technologies, 60242-1) and plated overnight on LB-AMP plates. $3x10^6$-$2x10^7$ CFU

colonies were harvested by flushing prewarmed LB broth and cultured for 2 hours in

liquid LB. sgRNA oligo pools were synthesized as oPools (IDT), containing BsmBI

cloning compatible overhangs. The oligos were then annealed and introduced into the

CROP-seq-V2.mCherry lentiviral plasmid using the NEB Golden Gate Assembly Kit as

described above. Approximately 5x10^7 CFUs for OvCa library and 2x10^7 colonies for

MetTag library were harvested for maxiprep. For both LARRY barcode and CRISPR

libraries, plasmids were purified using the ZymoPURE II Plasmid Maxiprep kit (Zymo

Research, D4202). The MetTag-LARRY-BC.ID vector map and CROP-seq-V2.mCherry

vector map will be made available on Addgene upon publication.

1069

Plasmid Library Amplification and Diversity Validation

Plasmid DNA was amplified using two step PCR to first amplify and then add indexing

sequences to libraries. First round PCR was conducted using custom P5 and P7

adapters with overhangs compatible with Nextera indexing primers, followed by cleanup

with Agencourt AMPure XP beads (Beckman Coulter, NC9933872) at a 1.8x bead ratio

to remove primers. Index PCR reaction was conducted as outlined in the Illumina 16S

Metagenomic Sequencing Library Preparation manual (Illumina, Doc #15044223, page

11-12) using 2x KAPA HiFi HotStart ReadyMix (Roche, KK2601) and Nextera XT Index

kit v2 primers (Illumina, FC-131-1001). Final indexed libraries were quantified via the

Qubit HS assay kit. LARRY libraries were run on a Bioanalyzer instrument and

sequenced on an Illumina MiSeq instrument. CRISPR libraries were run on TapeStation

using the High Sensitivity D1000 assay (Agilent, 5067-5585, 5067-5594) and

sequenced on a NovaSeq-X instrument. Demultiplexed fastq files were filtered for high

quality reads, and the number of unique reads was quantified. For the CRISPR libraries,

a customized script was used to estimate the percentage coverage based on the

original guide RNA whitelist.

1086

Lentivirus Production and Target Cell Transduction

MetTag-LARRY-BC.ID, CROP-seq-V2.mCherry guide, or Lenti-Luciferase.P2A.Neo

(Addgene # 105621), and second-generation lentiviral packaging plasmids were

transfected into HEK293T cells along with the CalFectin transfection reagent

(SignaGen, SL100478). Lentivirus was harvested after 72 hours and concentrated

(Lenti-X Concentrator, Takara Bio, 631231). Concentrated lentivirus was added to target

cells along with 8 ug/ml polybrene. To achieve low MOI for CRISPR screening

experiments, ID8 p53$^{-/-}$ ovarian cancer cells were infected at a low multiplicity of

infection (MOI) of 0.15, as calculated by the percentage of mCherry-positive cells 48

hours post transduction. 48 hours post transfection, cells were selected with either

G418 (pRG212-LARRY; Sigma-Aldrich, 4727878001) or puromycin (CROP-seq-

V2.mCherry; Invivogen, ant-pr-1) for a minimum of 7 days.

1099

Genomic DNA (gDNA) Amplicon Library Preparation

Genomic DNA was extracted from cultured cells and digested tissues using either the Quick-DNA Midiprep Kit (Zymo Research, D4075) or the Quick-DNA Miniprep Plus Kit (Zymo Research, D4068), depending on the starting cell number. Extracted DNA yield was quantified via a NanoDrop spectrophotometer, and samples were normalized to the same, lowest input concentration prior to PCR. Two rounds of PCR amplification were performed on > 200 ng starting gDNA material. The first round was utilized for amplicon amplification with nested primers for 15-20 cycles using a high-fidelity hot start polymerase, followed by purification with a 1.8x bead-to-sample AMPure XP bead ratio. Eluted amplicon DNA was quantified using the Qubit high sensitivity (HS) assay kit (Thermo Fisher Scientific, Q33231) and concentrations normalized prior to indexing. Approximately 5 ng of DNA was carried into the index PCR rection, which was conducted similarly to the plasmid library construction protocol described above. Index products were purified using the same AMPure XP beads ratio to remove excess primers. Library quantity was assessed via the Qubit HS assay kit and analyzed on a TapeStation instrument using the High Sensitivity D1000 assay to confirm expected library size and distribution. LARRY gDNA libraries were sequenced in-house on an Illumina MiniSeq instrument (Mid output kit, 300 cycles) while CRISPR gDNA libraries (101 genes*10 guides per gene, 20 non targeting guides = 1,030 guides) were sequenced on a NovaSeq-X instrument. High quality reads were filtered and analyzed via two parallel approaches: LARRY barcode counts were extracted to quantify library batch identity (ID), while CRISPR libraries were processed using the MAGeCK pipeline to identify enriched or depleted guide RNAs. Primers used for amplification were as follows:

LARRY Round 1 PCR REV, Tm: 55.9:

5' – GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GTG GGG AAT TTC TAC TCT TGT AGA TCG – 3'

LARRY Round 1 PCR FWD Stagger, Tm: 59.0:

5'-TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CTT GCT AGG ACC GGC CTT AAA GC -3'

CRISPR Round 1 PCR REV, Tm: 51.0:

1131 5' – GTC TCG TGG GCT CGG AGA TGT GTA TAA GAG ACA GTT CTT GGC TTT ATA

1132 TAT CTT GTG G– 3'

1133 CRISPR Round 1 PCR FWD Stagger, Tm: 53.0:

1134 5' – TCG TCG GCA GCG TCA GAT GTG TAT AAG AGA CAG CTC AAG TTG ATA ACG

1135 GAC TAG C– 3'

1136

PiggyBac Transposon Transfection

SpCas9 and CreERT2 coding sequences were cloned into PiggyBac transposon

plasmids containing a hygromycin resistance cassette via Gateway cloning

(ThermoFisher Scientific, 11791020 & 11789020). Target cells were plated in 6-well

plates 24 hours prior to transfection. Five ug of the SpCas9 or 1 ug of CreERT2

PiggyBac plasmid was co-transfected with 0.5 ug of Super PiggyBac Transposase

(System Biosciences, PB210PA-1), along with the Calfectin transfection reagent. The

transfection mixture was added to cells in a dropwise manner. 12-16 hours post

transfection complete growth media was replenished. Forty-eight hours post

transfection, cells were selected with hygromycin B (Corning, 30-240-CR) for a

minimum of 10 days to enrich for successful transgene integrated cells.

1148

Cell Preparation for Single-Cell RNA Sequencing

Naïve and tumor bearing mice were sacrificed via cervical dislocation six weeks

following initial cancer cell injection. To collect peritoneal cells, 5 mL of cold 1xHBSS

was injected intraperitoneally using a 25G needle, followed by lavage and peritoneal

fluid retrieval using a 20G needle. Omenta were excised from the peritoneal cavity and

examined under a dissecting microscope to confirm the presence of macroscopic

metastases. Omental tissues were then minced and digested for 30 minutes at 37ºC in

RPMI-1640 medium containing 0.4% collagenase type II (Gibco, 17101015), 50 ug/ml

DNAseI, and 0.1% BSA. Following digestion, cell suspensions were passed through a

70-uM strainer and washed with cold 1xPBS. Lavage, ascites, and omental samples

were resuspended in 1x RBC lysis buffer (BioLegend, 420301) to remove red blood

cells. To prepare for single-cell analysis, omental samples were stained with ZombieRed

(BioLegend, 423109) to exclude dead cells and sorted via FACS. Sorted cells were then

prepared for 10X Genomics Chromium Single Cell Gene Expression analysis. For

TNBC lung sample scRNA-seq, mice were perfused with 5mL of ice cold 1xPBS, lungs

were collected and imaged using the GFP channel on a dissecting scope. Lungs with

detectable tumor burden were cut into small pieces and processed into single cell

suspensions by digestion with 1x collagenase/hyaluronidase (STEMCELL Technologies,

#07912) and 0.15 mg/ml DNase I in serum free RPMI 1640 medium. After passing the

digested samples through a 70-uM strainer and washing with serum free RPMI, the

samples were lysed for RBCs and processed similarly to OvCa samples downstream. A

detailed protocol for sample multiplexing via hashtags can be found in the CITE-seq/cell

hashing protocol on the CITE-seq website (https://cite-seq.com/wp-

content/uploads/2019/02/cite-seq_and_hashing_protocol_190213.pdf). In brief, samples

were first blocked with FcR Blocking Reagent (Miltenyi Biotec, 130-092-575) in 50 uL of

cell staining buffer for 20 minutes on ice. Samples were then stained with TotalSeq™-A

hashing antibodies obtained from BioLegend, including:

M-HTO-1 (M1/42; 30-F11, 155801),

M-HTO-2 (M1/42; 30-F11, 155803),

M-HTO-3 (M1/42; 30-F11, 155805),

M-HTO-4 (M1/42; 30-F11, 155807),

M-HTO-5 (M1/42; 30-F11, 155809),

M-HTO-6 (M1/42; 30-F11, 155811),

M-HTO-7 (M1/42; 30-F11, 155813),

M-HTO-8 (M1/42; 30-F11, 155815).

Staining was performed for 25 minutes on ice, followed by three washes with staining

buffer (decreasing EDTA concentration per wash) and loading onto a 10X Genomics

Chromium Controller.

1187

1188   **scRNA-seq Library Preparation and Illumina Sequencing**

1189   Single-cell transcriptomic libraries were prepared as outlined in the Chromium Next

1190   GEM Single Cell 3' Reagents Kits v3.1 with Feature Barcoding technology kit manual.

1191   Kits necessary for transcriptome library preparation were: Chromium Next GEM Single

1192   Cell 3' GEM, Library & Gel Bead Kit v3.1, 4 rxns (10x Genomics, PN-1000128),

Chromium Single Cell 3' Feature Barcode Library Kit, 16 rxns (10x Genomics, PN-1000079), Chromium Next GEM Chip G Single Cell Kit, 16 rxns (10x Genomics, PN-1000127), and Single Index Kit T Set A, 96 rxns (10x Genomics, PN-1000213). Hashtag libraries were prepared using the standard CITE-seq protocol (https://cite-seq.com/wp-content/uploads/2019/02/cite-seq_and_hashing_protocol_190213.pdf). LARRY-BC.ID barcode libraries were constructed using a custom nested PCR protocol followed by indexing. Primers required for LARRY-BC.ID library preparation included:

Round 1 PCR FWD, Tm: 61.0

5' – *GCA GCG TCA GAT GTG TAT AAG AGA CAG* – 3'

Round 1 PCR REV: 45.0

5' – <u>CCT TGG CAC CCG AGA ATT CCA</u> aat ttc tac tct tgt aga t – 3'

Index PCR FWD, Tm: 61.0

5' – AAT GAT ACG GCG ACC ACC GAG ATC TAC ACT CGT CG*G CAG CGT CAG ATG*

TGT ATA AGA GAC AG – 3'

Index PCR REV (ADT-RPI-1), Tm 61.0:

5' – CAA GCA GAA GAC GGC ATA CGA GAT CGT GAT GTG ACT GGA GTT <u>CCT TGG</u>

<u>CAC CCG AGA ATT CCA</u> – 3'

{Partial Read 1N; LARRY transcript (lowercase), <u>P7 adaptor</u>, RPI1 **index**}

The quality and quantity of the three individual libraries (transcriptome, HTO, and LARRY) were assessed via Qubit and TapeStation prior to sequencing. Libraries were pooled for sequencing on an Illumina NovaSeq X Plus platform with paired end reads (PE150).

scRNA-seq Data Analysis

Raw fastq files generated from the sequencing runs (mRNA, HTO, LARRY outputs) were processed through the 10x Genomics CellRanger pipeline v.7.2 to generate count matrices. Gene expression (mRNA) reads were aligned to a custom mouse reference genome containing fluorescent marker transcripts (mm10_3.0.0_FPs). Data from two different chromium 10x lanes, ascites lane and omentum lane, were merged using the

SCT-normalized dataset integration method. Details related to integration steps can be found at: https://satijalab.org/seurat/articles/. In brief, individual lane single cell data was normalized for HTO (HTOdemux) followed by SCTransform prior to integration. Downstream analyses including dimensional reduction via principal component analysis (PCA), clustering and UMAP embedding, and differential gene expression (DEG) analysis were performed in RStudio under the Seurat package (v4.0)[18]. Cell type identities were assigned based on cluster marker genes. Subcluster level labels indicating different cellular statuses were determined through a combination of cluster marker genes and examining enriched pathways. RNA velocity analysis was performed in Python using scVelo and CellRank. Loom files were generated for each lane and merged, followed by Seurat object conversion to H5AD format. Metadata containing clustering and HTO information was integrated into the SCANPY anndata object. More details on the CellRank method can be found at: https://cellrank.readthedocs.io/en/latest/index.html. NicheNetR and CellChat were performed in RStudio. Sender and receiver cell types were selected prior to NicheNetR analysis. scRNA-seq expression data was analyzed using the Wilcoxon rank sum test either through the direct function wilcox.test() when assigning custom module scores or through the FindMarkers() function as part of the standard DEG analysis. Adjusted p values <0.05 were considered significant. Over-representation analysis (ORA) was performed on a filtered list of upregulated or downregulated genes from DEG analysis and the ReactomePA and clusterProfiler RStudio packages. org.Mm.eg.db package was used to convert genes to Entrez ID symbols. Adjusted p values <0.05 and log fold change > (+/-) 0.25 were considered significant. ORA uses Fisher's exact test to determine enriched gene sets. For Single Cell Gene Set Enrichment Analysis (ssGSEA), we utilized the Escape R package.

Flow cytometry

Samples were extracted and pre-processed similarly as the scRNA-seq samples. Following tissue digestion of lung or omentum tissues and RBC lysis, digested samples were washed in 1x Flow Staining Buffer (Tonbo Biosciences, TNB-4222-L500). For viability assessment, cells were incubated with ZombieAqua Fixable Viability Kit

(BioLegend, 423102) at a 1:750 dilution in 1xPBS for 20 minutes at room temperature.

Following viability staining, cells were blocked with 1x FcR Blocking Reagent (Miltenyi

Biotec, 130-092-575) to minimize nonspecific antibody staining. Cells were then stained

with a combination of the following fluorochrome-conjugated surface antibodies: Pacific

Blue™ anti-mouse CD45 (BioLegend, 157211, S18009F), APC/Fire™ 750 anti-mouse

CD3 (BioLegend, 100247, 17A2) or PE/Cyanine7 anti-mouse CD3 (BioLegend, 100219,

17A2), PE anti-TREM-2 (BioLegend, 824805, 6E9), APC anti-mouse CCR5 (BioLegend,

107011, HM-CCR5), Brilliant Violet 605™ anti-mouse/human CD11b (BioLegend,

101237, M1/70), APC anti-mouse I-A-I-E (BioLegend, 107613, M5/114.15.2), Brilliant

Violet 711™ anti-mouse CD4 (BioLegend, 100447, GK1.5), PE/Dazzle™ 594 anti-

mouse NK-1.1 (BioLegend, 156517, S17016D) or PE anti-mouse NK-1.1 (BioLegend,

156503, S17016D), Brilliant Violet 421™ anti-mouse PD-L1 (BioLegend, 124315,

10F.9G2), PE/Cyanine7 anti-mouse CD8a (BioLegend, 100721, 53-6.7) or

APC/Cyanine7 anti-mouse CD8a (BioLegend, 100713, 53-6.7), Brilliant Violet 785™

anti-mouse F4/80 (BioLegend, 123141, BM8). Cancer cells harvested from culture were

similarly stained with ZombieAqua, FcR blocked, and then incubated with IFNGR1

Recombinant Rabbit Monoclonal Antibody FITC (Invitrogen, MA5-46853, 062) with the

corresponding Rabbit IgG Isotype Control FITC, eBioscience™ (Invitrogen, 11-4614-

80). Cells were then washed with staining buffer, followed by analysis on a Cytek

Northern Lights flow cytometer. FCS files were exported and analyzed using FlowJo

(BD Biosciences).

1276

Western Blotting

Cells were plated at a density of 250,000-500,000 cells per 10 cm dish for regular

attachment plate experiments. For anoikis assays, 3x 6 wells per sample were seeded

with 300,000 cells each. The next day, cells were either treated with vehicle (1xPBS) or

100 ng/ml mouse recombinant IFNγ (PeproTech, AF-315-05-100UG). After treatment,

cells were harvested, washed with 1xPBS, and lysed in M2 lysis buffer (50 nM Tris-HCl,

pH 7.4, 150 mM NaCl, 1% Triton X-100, 0.5 mM EDTA) with 1x Protease/Phosphatase

Inhibitor Cocktail (CST, 5872). Cells were lysed for 1 hour on ice before centrifugation at

high speed to collect protein lysates. Protein quantity was deterimed using the Pierce™

BCA Protein Assay Kit (Thermo Scientific, 23225). Protein lysate samples utilized for

GBP-1 detection were resolved using 4-20% Mini-PROTEAN TGX Precast Protein Gels

(Bio-Rad, 4561094). Other immunoblots were performed on fixed concentration using 8-

10% SDS-polyacrylamide gels. All gel transfers were performed onto nitrocellulose

membranes (Fisher Scientific, 45-004-075). Nitrocellulose membranes were blocked in

5% BSA/1xTBST for 1 hour at room temperature, followed by primary antibody

incubation overnight on a 4ºC shaker. The following day, blots were washed with

1xTBST followed by incubation with rabbit or mouse-specific horseradish peroxidase

(HRP)-conjugated secondary antibodies. Signals were developed with SuperSignal

West Pico PLUS Chemiluminescent Substrate (Thermo Fisher Scientific, #34577) or

SuperSignal West Femto Maximum Sensitivity Substrate (Thermo Fisher Scientific,

#34094). Primary antibodies used for western blotting were:

Beta-Actin (13E5) Rabbit mAb, CST #4970, 1:5,000 dilution

GBP-1 Rabbit Polyclonal Ab, {Thermo, PA5-23509}, 1:1,000 dilution

Cas9 (S. pyogenes) (7A9-3A3) Mouse mAb, CST #14697, 1:1,000 dilution

IRF-1 (D5E4) XP® Rabbit mAb, CST #8478, 1:1,000 dilution

Cre Recombinase (D7L7L) XP® Rabbit mAb, CST #15036, 1:1,000 dilution

GAPDH (14C10) Rabbit mAb, CST #2118, 1:5,000 dilution

Stat1 (D1K9Y) Rabbit mAb, CST #14994, 1:1,000 dilution

p-Stat1 (Tyr701) (58D6) Rabbit mAb #9167, 1:1,000 dilution

Vinculin (E1E9V) XP® Rabbit mAb #13901, 1:5,000 dilution

Anti-rabbit IgG, HRP-linked Ab, CST #7074, 1:5,000 dilution

Anti-mouse IgG, HRP-linked Ab, CST #7076, 1:5,000 dilution

1309

Clinical dataset mining

Patients were stratified into IFNGR1 high and IFNGR1 low expression groups and

survival curve was generated using Kaplan-Meier Plotter (KMplotter,

https://www.kmplot.com). This database includes 1816 samples primarily from

GSE9891 and TCGA[20]. Patient scRNA-seq RDS file was obtained from Mendeley

Data[55] (https://doi.org/10.17632/rc47y6m9mp.1) and contained 14 patient samples

including PBMCs, lymph nodes, primary tumor, metastasis, and malignant ascites. The

RNA data was re-normalized and analyzed using the Seurat package in RStudio in

similar ways to the murine scRNA-seq dataset described above.

1319

Quantitative Reverse Transcriptase (RT) PCR

The IFNγ treatment time course was performed in 12 well plates. After seeding 1.0x105

cells per well, wells were treated with 100 ng/ml IFNγ for 0 hours (PBS control), 4 hours,

8 hours, or 24 hours. Total cell RNA was extracted from cells using the Direct-zol RNA

Miniprep kit with TriReagent (Zymo Research, R2051-A) following the manufacturer

protocol. One microgram of total RNA was used for reverse transcription to synthesize

single stranded cDNA using the iScript^{TM} Reverse Transcription Supermix for RT-qPCR

(Bio-Rad, 1708840). qPCR reactions were performed on a Bio-Rad CFX Opus

instrument with SsoAdvanced Universal SYBR Green Supermix (Bio-Rad, 1725270).

Data analysis was performed using the CFX Maestro Software. Cq values were used to

calculate relative gene expression levels. In brief, calculations were based on the $2^{-\Delta\Delta CT}$

method with housekeeping genes serving as internal controls and fold change was

calculated by normalizing to the zero-hour treatment timepoint Cq values. Statistical

analyses and plots were generated in GraphPad Prism. RT-qPCR primers used were as

follows:

Beta-actin (Mouse) Forward primer TM: 57.6

5'- GGC TGT ATT CCC CTC CAT CG -3'

Beta-actin (Mouse) Reverse primer TM: 55.9

5'- CCA GTT GGT AAC AAT GCC ATG T-3'

GAPDH (Mouse) Forward primer TM: 59.8

5'- CAT CAC TGC CAC CCA GAA GAC TG -3'

GAPDH (Mouse) Reverse primer TM: 61.6

5'- ATG CCA GTG AGC TTC CCG TTC AG -3'

Gbp2b (Mouse) Forward primer TM: 58.5

5'- ATC TGC TCA TTG CTC AGA CTT ACT GG -3'

Gbp2b (Mouse) Reverse primer TM: 57.4

5'- AAG TAT TTT CTC AGC ATC ACT TCA GCC -3'

Irf1 (Mouse) Forward primer TM: 60.4

5'- ACC AAA TCC CAG GGC TGA TCT GG -3'

Irf1 (Mouse) Reverse primer TM: 58.3

5'- GAA GTT TGC CTT CCA TGT CTT GGG-3'

Stat1 (Mouse) Forward primer TM: 55.1

5'- GCT GCC TAT GAT GTC TCG TTT -3'

Stat1 (Mouse) Reverse primer TM: 55.2

5'- TGC TTT TCC GTA TGT TGT GCT -3'

Marco (Mouse) Forward primer TM: 61.1

5'- ATC CAC CTG ATC CTG CTC ACG G -3'

Marco (Mouse) Reverse primer TM: 60.8

5'- CAC TGC AGC GAG AAG AAG GGC -3'

Slfn1 (Mouse) Forward primer TM: 60.9

5'- AGG GAA AGG TCA AGG CTC ACA TTA AAA ATC C -3'

Slfn1 (Mouse) Reverse primer TM: 60.1

5'- AGG CTT CCC AAC AGA GAC ATC TGG -3'

1363

Anoikis Assay

Cancer cells ($1x10^4$) were seeded into 96-well ultra-low attachment (ULA) plates

(Corning, CLS7007) to mimic anchorage-independent conditions in the peritoneal cavity

environment. After 24 hours, 100 ng/ml recombinant mouse IFNγ or vehicle control

(1xPBS) was added to appropriate wells. Twenty-four hours after treatment (i.e. 48hr

after start of culture), CellTiter-Glo Luminescent Cell Viability Assay reagent (Promega,

G7570) was applied to every assay well, and plates were incubated for 20 min at room

temperature. Luminescence, indicating cellular ATP levels, was measured using a

Tecan plate reader instrument and analyzed using GraphPad Prism software.

For macrophage coculture experiments, $1.0x10^4$ cancer cells were seeded together with

$1.0x10^4$ macrophages per well in ULA plates. Macrophages were enriched from the

peritoneal cavity of wild type female mice using the MojoSort Mouse F4/80 Positive

Selection Kit (BioLegend, 480170), following manufacturer's protocol. After 48 hours of

coculture, cell suspensions were harvested, pelleted, and incubated in TrypLE (Thermo

Fisher Scientific, 12605028) for 10 minutes at 37 ºC to dissociate cell aggregates. Cell

suspensions were passed through a 40-uM strainer before flow cytometric staining as
described earlier.

1381

Colony Formation Assay

Two hundred ID8 p53$^{-/-}$ cells were plated onto 6 well plates at the start of the

experiment. For wells treated with IFNγ, 100 ng/ml IFNγ was maintained throughout the

7-day colony formation assay period. Wells were washed with 1xPBS, fixed with

methanol, and stained with crystal violet solution (0.5% w/v) at room temperature for 20

minutes. Wells were then washed with deionized water and dried overnight. Colonies

were counted and quantified in GraphPad Prism.

1389

Validation of Cre-based NicheTracing

HEK293T cells were transfected with a loxp-dsRED-loxp-eGFP Cre reporter construct

(Addgene # 141148) and cocultured in a 1:1 ratio with Cre.ERT2 expressing ID8 p53$^{-/-}$

or E0771 cell lines in the presence or absence of 2 uM 4-hydroxytamoxifen (4-OHT,

MedChemExpress, Cat # HY-16950). Seventy-two hours after plating, wells were

assessed for GFP+ switched cells using a Nikon Ti2 microscope. For independent *in*

vivo validation, E0771-Cre.ERT2 cells were injected into the retro-orbital sinus of Ai6

mice. One week post injection, three consecutive tamoxifen or corn oil control

treatments were performed (75 mg/kg/dose) i.p.. Two weeks post cancer cell injection,

mice were euthanized, perfused intracardially, and lungs were extracted for imaging and

flow cytometry analysis. For the ovarian metastasis model, tamoxifen injections were

administered in five consecutive doses, two weeks post ID8-Cre.ERT2 cell i.p. injection.

1402

Bulk RNA-seq

3.0×10^5 Ifngr1-KO or sgNTC ID8 p53$^{-/-}$ cells were plated onto 6-well ULA and regular

treatment plates. Recombinant IFNγ (100 ng/ml) or vehicle (1xPBS) was added to

corresponding wells 24 hours post initial seeding. Cells were harvested 24 hours post

IFNγ treatment. ULA samples were subjected to dead cell removal (Miltenyi Biotec, 130-

090-101) and total RNA was extracted using Direct-zol RNA Miniprep kit with

TriReagent (Zymo Research, R2051-A) following the manufacturer protocol. Purified

RNA was subjected to library preparation. In brief, mRNA was purified using poly-T oligo
magnetic beads, fragmented, and reverse transcribed. Resulting cDNA was subjected
to amplification and indexing. Final libraries were sequenced on a NovaSeq-X
instrument. Raw reads were trimmed and aligned to the mouse reference genome
(mm10). Following transcript level quantification, subsequent count matrices were
analyzed in RStudio. DESeq2 was used for downstream analysis and variance-
stabilized counts (VSTs) were used for PCA and visualization. Pathway analysis
(Hallmark, Reactome, Go Biological Processes) was conducted through the
clusterProfiler R package.
1419

Data and Resource Availability
scRNA-seq datasets with LARRY barcode and HTO multiplexing was deposited at
GEO: GSE302619 (ovarian) and GSE309120 (lung) and will be made publicly available
upon publishing. Reviewer access tokens for the datasets presented here are:
qvevaiacnnarbwr (ovarian) and obyficmwzdmnnkh (lung). Human scRNA-seq data has
been published elsewhere and is publicly available: GSE14764. Code and any
additional information required to reanalyze the data reported here can be requested
from the lead contact.
1428

1429 **Declaration of generative AI and AI-assisted technologies in the writing process**
1430 During the preparation of this work the author(s) used ChatGPT in order to improve
1431 writing clarity. After using this tool/service, the author(s) reviewed and edited the content
1432 as needed and take(s) full responsibility for the content of the publication.
1433

1434 **REFERENCES:**
1435

1. Gerstberger, S., Jiang, Q., and Ganesh, K. (2023). Metastasis. Cell *186*, 1564–1579.
https://doi.org/10.1016/j.cell.2023.03.003.

2. Ritch, S.J., and Telleria, C.M. (2022). The Transcoelomic Ecosystem and Epithelial
Ovarian Cancer Dissemination. Front. Endocrinol. *13*, 886533.
https://doi.org/10.3389/fendo.2022.886533.

3. Motohara, T., Masuda, K., Morotti, M., Zheng, Y., El-Sahhar, S., Chong, K.Y., Wietek,
N., Alsaadi, A., Carrami, E.M., Hu, Z., et al. (2019). An evolving story of the
metastatic voyage of ovarian cancer cells: cellular and molecular orchestration of
the adipose-rich metastatic microenvironment. Oncogene *38*, 2885–2898.
https://doi.org/10.1038/s41388-018-0637-x.

4. Almeida-Nunes, D.L., Mendes-Frias, A., Silvestre, R., Dinis-Oliveira, R.J., and
Ricardo, S. (2022). Immune Tumor Microenvironment in Ovarian Cancer Ascites. Int.
J. Mol. Sci. *23*, 10692. https://doi.org/10.3390/ijms231810692.

5. Lane, D., Matte, I., Garde-Granger, P., Bessette, P., and Piché, A. (2018). Ascites IL-
10 Promotes Ovarian Cancer Cell Migration. Cancer Microenviron. *11*, 115–124.
https://doi.org/10.1007/s12307-018-0215-3.

6. Mustea, A., Pirvulescu, C., Könsgen, D., Braicu, E.I., Yuan, S., Sun, P.,
Lichtenegger, W., and Sehouli, J. (2008). Decreased IL-1 RA concentration in
ascites is associated with a significant improvement in overall survival in ovarian
cancer. Cytokine *42*, 77–84. https://doi.org/10.1016/j.cyto.2008.01.011.

7. Ahmed, N., and Stenvers, K. (2013). Getting to Know Ovarian Cancer Ascites:
Opportunities for Targeted Therapy-Based Translational Research. Front. Oncol. *3*.
https://doi.org/10.3389/fonc.2013.00256.

8. Lineage tracing on transcriptional landscapes links state to fate during differentiation
| Science (2025). https://www.science.org/doi/10.1126/science.aaw3381.

9. Dasgupta, A., Lim, A.R., and Ghajar, C.M. (2017). Circulating and disseminated
tumor cells: harbingers or initiators of metastasis? Mol. Oncol. *11*, 40–61.
https://doi.org/10.1002/1878-0261.12022.

10. Borcherding, N., Vishwakarma, A., Voigt, A.P., Bellizzi, A., Kaplan, J., Nepple, K.,
Salem, A.K., Jenkins, R.W., Zakharia, Y., and Zhang, W. (2021). Mapping the
immune environment in clear cell renal carcinoma by single-cell genomics.
Commun. Biol. *4*, 122. https://doi.org/10.1038/s42003-020-01625-6.

11. Cellular barcoding to decipher clonal dynamics in disease | Science (2025).
https://www.science.org/doi/10.1126/science.abm5874?url_ver=Z39.88-
2003&rfr_id=ori:rid:crossref.org&rfr_dat=cr_pub%20%200pubmed.

12. CellRank for directed single-cell fate mapping | Nature Methods (2025).
https://www.nature.com/articles/s41592-021-01346-6.

13. Yu, G., Wang, L.-G., Han, Y., and He, Q.-Y. (2012). clusterProfiler: an R Package for
Comparing Biological Themes Among Gene Clusters. OMICS J. Integr. Biol. *16*,
284–287. https://doi.org/10.1089/omi.2011.0118.

14. Bahrami, S., and Drabløs, F. (2016). Gene regulation in the immediate-early
response process. Adv. Biol. Regul. *62*, 37–49.
https://doi.org/10.1016/j.jbior.2016.05.001.

15. Koivisto, E., Acosta, A.J., Moilanen, A.-M., Tokola, H., Aro, J., Pennanen, H.,
Säkkinen, H., Kaikkonen, L., Ruskoaho, H., and Rysä, J. (2014). Characterization of
the Regulatory Mechanisms of Activating Transcription Factor 3 by Hypertrophic
Stimuli in Rat Cardiomyocytes. PLOS ONE *9*, e105168.
https://doi.org/10.1371/journal.pone.0105168.

16. Zhao, J., Peng, H., Gao, J., Nong, A., Hua, H., Yang, S., Chen, L., Wu, X., Zhang,
H., and Wang, J. (2021). Current insights into the expression and functions of tumor-
derived immunoglobulins. Cell Death Discov. *7*, 148. https://doi.org/10.1038/s41420-
021-00550-9.

17. Mehta, A., and Merkel, O.M. (2020). Immunogenicity of Cas9 Protein. J. Pharm. Sci.
109, 62–67. https://doi.org/10.1016/j.xphs.2019.10.003.

18. Li, W., Xu, H., Xiao, T., Cong, L., Love, M.I., Zhang, F., Irizarry, R.A., Liu, J.S.,
Brown, M., and Liu, X.S. (2014). MAGeCK enables robust identification of essential
genes from genome-scale CRISPR/Cas9 knockout screens. Genome Biol. *15*, 554.
https://doi.org/10.1186/s13059-014-0554-4.

19. Shultz, D.B., Sandhya Rani, M.R., Fuller, J.D., Ransohoff, R.M., and Stark, G.R.
(2009). Roles of IKK-β, IRF1, and p65 in the Activation of Chemokine Genes by
Interferon-γ. J. Interferon Cytokine Res. *29*, 817–824.
https://doi.org/10.1089/jir.2009.0034.

20. Győrffy, B. (2023). Discovery and ranking of the most robust prognostic biomarkers
in serous ovarian cancer. GeroScience *45*, 1889–1898.
https://doi.org/10.1007/s11357-023-00742-4.

21. Rouette, A., Trofimov, A., Haberl, D., Boucher, G., Lavallée, V.-P., D'Angelo, G.,
Hébert, J., Sauvageau, G., Lemieux, S., and Perreault, C. (2016). Expression of
immunoproteasome genes is regulated by cell-intrinsic and –extrinsic factors in
human cancers. Sci. Rep. *6*, 34019. https://doi.org/10.1038/srep34019.

22. Ma, X., Li, Y., Wang, D., Niu, J., Li, Q., Chen, Y., Wang, M., Wen, J., Liao, C., Wang,
N., et al. (2025). PSMB10 maintains the stemness of chemotherapeutic drug-
resistant leukemia cells by inhibiting senescence and cytotoxic T lymphocyte-
mediated killing in a ubiquitinated degradation manner. J. Exp. Clin. Cancer Res. *44*,
170. https://doi.org/10.1186/s13046-025-03420-9.

23. Barbarulo, A., Iansante, V., Chaidos, A., Naresh, K., Rahemtulla, A., Franzoso, G.,
Karadimitris, A., Haskard, D.O., Papa, S., and Bubici, C. (2013). Poly(ADP-ribose)
polymerase family member 14 (PARP14) is a novel effector of the JNK2-dependent
pro-survival signal in multiple myeloma. Oncogene *32*, 4231–4242.
https://doi.org/10.1038/onc.2012.448.

24. Escors, D., Gato-Cañas, M., Zuazo, M., Arasanz, H., García-Granda, M.J., Vera, R.,
and Kochan, G. (2018). The intracellular signalosome of PD-L1 in cancer cells.
Signal Transduct. Target. Ther. *3*, 26. https://doi.org/10.1038/s41392-018-0022-9.

25. Jin, S., Guerrero-Juarez, C.F., Zhang, L., Chang, I., Ramos, R., Kuan, C.-H., Myung,
P., Plikus, M.V., and Nie, Q. (2021). Inference and analysis of cell-cell
communication using CellChat. Nat. Commun. *12*, 1088.
https://doi.org/10.1038/s41467-021-21246-9.

26. Chen, S., Saeed, A.F.U.H., Liu, Q., Jiang, Q., Xu, H., Xiao, G.G., Rao, L., and Duo,
Y. (2023). Macrophages in immunoregulation and therapeutics. Signal Transduct.
Target. Ther. *8*, 207. https://doi.org/10.1038/s41392-023-01452-1.

27. Ahmed, I., and Ismail, N. (2020). M1 and M2 Macrophages Polarization via
mTORC1 Influences Innate Immunity and Outcome of Ehrlichia Infection. J. Cell.
Immunol. *2*, 108–115. https://doi.org/10.33696/immunology.2.029.

28. O'Carroll, C., Fagan, A., Shanahan, F., and Carmody, R.J. (2014). Identification of a
Unique Hybrid Macrophage-Polarization State following Recovery from
Lipopolysaccharide Tolerance. J. Immunol. *192*, 427–436.
https://doi.org/10.4049/jimmunol.1301722.

29. Patterson, M.T., Firulyova, M.M., Xu, Y., Hillman, H., Bishop, C., Zhu, A., Hickok,
G.H., Schrank, P.R., Ronayne, C.E., Caillot, Z., et al. (2023). Trem2 promotes foamy
macrophage lipid uptake and survival in atherosclerosis. Nat. Cardiovasc. Res. *2*,
1015–1031. https://doi.org/10.1038/s44161-023-00354-3.

30. Binnewies, M., Pollack, J.L., Rudolph, J., Dash, S., Abushawish, M., Lee, T.,
Jahchan, N.S., Canaday, P., Lu, E., Norng, M., et al. (2021). Targeting TREM2 on
tumor-associated macrophages enhances immunotherapy. Cell Rep. *37*, 109844.
https://doi.org/10.1016/j.celrep.2021.109844.

31. Park, M.D., Reyes-Torres, I., LeBerichel, J., Hamon, P., LaMarche, N.M., Hegde, S.,
Belabed, M., Troncoso, L., Grout, J.A., Magen, A., et al. (2023). TREM2
macrophages drive NK cell paucity and dysfunction in lung cancer. Nat. Immunol.
24, 792–801. https://doi.org/10.1038/s41590-023-01475-4.

32. Iyer, S., Zhang, S., Yucel, S., Horn, H., Smith, S.G., Reinhardt, F., Hoefsmit, E.,
Assatova, B., Casado, J., Meinsohn, M.-C., et al. (2021). Genetically Defined
Syngeneic Mouse Models of Ovarian Cancer as Tools for the Discovery of
Combination Immunotherapy. Cancer Discov. *11*, 384–407.
https://doi.org/10.1158/2159-8290.CD-20-0818.

33. Zheng, X., Wang, X., Cheng, X., Liu, Z., Yin, Y., Li, X., Huang, Z., Wang, Z., Guo, W.,
Ginhoux, F., et al. (2023). Single-cell analyses implicate ascites in remodeling the
ecosystems of primary and metastatic tumors in ovarian cancer. Nat. Cancer *4*,
1138–1156. https://doi.org/10.1038/s43018-023-00599-8.

34. Patras, L., Shaashua, L., Matei, I., and Lyden, D. (2023). Immune determinants of the pre-metastatic niche. Cancer Cell *41*, 546–572. https://doi.org/10.1016/j.ccell.2023.02.018.

35. Peinado, H., Zhang, H., Matei, I.R., Costa-Silva, B., Hoshino, A., Rodrigues, G., Psaila, B., Kaplan, R.N., Bromberg, J.F., Kang, Y., et al. (2017). Pre-metastatic niches: organ-specific homes for metastases. Nat. Rev. Cancer *17*, 302–317. https://doi.org/10.1038/nrc.2017.6.

36. Hershey, B.J., Barozzi, S., Orsenigo, F., Pompei, S., Iannelli, F., Kamrad, S., Matafora, V., Pisati, F., Calabrese, L., Fragale, G., et al. (2023). Clonal cooperation through soluble metabolite exchange facilitates metastatic outgrowth by modulating Allee effect. Sci. Adv. *9*, eadh4184. https://doi.org/10.1126/sciadv.adh4184.

37. Śliwa, A., Szczerba, A., Pięta, P.P., Białas, P., Lorek, J., Nowak-Markwitz, E., and Jankowska, A. (2024). A Recipe for Successful Metastasis: Transition and Migratory Modes of Ovarian Cancer Cells. Cancers *16*, 783. https://doi.org/10.3390/cancers16040783.

38. Schrijver, W.A.M.E., Selenica, P., Lee, J.Y., Ng, C.K.Y., Burke, K.A., Piscuoglio, S., Berman, S.H., Reis-Filho, J.S., Weigelt, B., van Diest, P.J., et al. (2018). Mutation Profiling of Key Cancer Genes in Primary Breast Cancers and Their Distant Metastases. Cancer Res. *78*, 3112–3121. https://doi.org/10.1158/0008-5472.CAN-17-2310.

39. Müller, V., and Pantel, K. (2004). Bone marrow micrometastases and circulating tumor cells: current aspects and future perspectives. Breast Cancer Res. *6*, 258–261. https://doi.org/10.1186/bcr942.

40. Sänger, N., Effenberger, K.E., Riethdorf, S., Van Haasteren, V., Gauwerky, J., Wiegratz, I., Strebhardt, K., Kaufmann, M., and Pantel, K. (2011). Disseminated tumor cells in the bone marrow of patients with ductal carcinoma in situ. Int. J. Cancer *129*, 2522–2526. https://doi.org/10.1002/ijc.25895.

41. Wawrzyniak, P., and Hartman, M.L. (2025). Dual role of interferon-gamma in the response of melanoma patients to immunotherapy with immune checkpoint inhibitors. Mol. Cancer *24*, 89. https://doi.org/10.1186/s12943-025-02294-x.

42. Remsik, J., Tong, X., Kunes, R.Z., Li, M.J., Estrera, R., Snyder, J., Thomson, C., Osman, A.M., Chabot, K., Sener, U.T., et al. (2025). Interferon-γ orchestrates leptomeningeal anti-tumour response. Nature, 1–10. https://doi.org/10.1038/s41586-025-09012-z.

43. Jorgovanovic, D., Song, M., Wang, L., and Zhang, Y. (2020). Roles of IFN-γ in tumor progression and regression: a review. Biomark. Res. *8*, 49. https://doi.org/10.1186/s40364-020-00228-x.

44. Zheng, H., Guan, X., Meng, X., Tong, Y., Wang, Y., Xie, S., Guo, L., and Lu, R.
(2023). IFN-γ in ovarian tumor microenvironment upregulates HLA-E expression and
predicts a poor prognosis. J. Ovarian Res. *16*, 229. https://doi.org/10.1186/s13048-
023-01286-z.

45. Reddy, S.U., Sadia, F.Z., Vancura, A., and Vancurova, I. (2024). IFNγ-Induced Bcl3,
PD-L1 and IL-8 Signaling in Ovarian Cancer: Mechanisms and Clinical Significance.
Cancers *16*, 2676. https://doi.org/10.3390/cancers16152676.

46. Murphy, B., Miyamoto, T., Manning, B.S., Mirji, G., Ugolini, A., Kannan, T., Hamada,
K., Zhu, Y.P., Claiborne, D.T., Huang, L., et al. (2024). Myeloid activation clears
ascites and reveals IL27-dependent regression of metastatic ovarian cancer. J. Exp.
Med. *221*, e20231967. https://doi.org/10.1084/jem.20231967.

47. Ye, Z., Ai, X., Zhao, L., Fei, F., Wang, P., and Zhou, S. (2021). Phenotypic plasticity
of myeloid cells in glioblastoma development, progression, and therapeutics.
Oncogene *40*, 6059–6070. https://doi.org/10.1038/s41388-021-02010-1.

48. Schol, P., Elsas, M.J. van, Middelburg, J., Twilhaar, M.K.N., Hall, T. van, Sluis, T.C.
van der, and Burg, S.H. van der (2024). Myeloid effector cells in cancer. Cancer Cell
42, 1997–2014. https://doi.org/10.1016/j.ccell.2024.11.002.

49. Guilliams, M., and Svedberg, F.R. (2021). Does tissue imprinting restrict
macrophage plasticity? Nat. Immunol. *22*, 118–127. https://doi.org/10.1038/s41590-
020-00849-2.

50. Yarilina, A., Park-Min, K.-H., Antoniv, T., Hu, X., and Ivashkiv, L.B. (2008). TNF
activates an IRF1-dependent autocrine loop leading to sustained expression of
chemokines and STAT1-dependent type I interferon–response genes. Nat. Immunol.
9, 378–387. https://doi.org/10.1038/ni1576.

51. Toney, N.J., Opdenaker, L.M., Frerichs, L., Modarai, S.R., Ma, A., Archinal, H., Ajayi,
G.O., and Sims-Mourtada, J. (2025). B cells enhance IL-1 beta driven invasiveness
in triple negative breast cancer. Sci. Rep. *15*, 2211. https://doi.org/10.1038/s41598-
025-86064-1.

52. Luo, J.-L., Kamata, H., and Karin, M. (2005). IKK/NF-κB signaling: balancing life and
death – a new approach to cancer therapy. J. Clin. Invest. *115*, 2625–2632.
https://doi.org/10.1172/JCI26322.

53. Horowitz, J.C., Lee, D.Y., Waghray, M., Keshamouni, V.G., Thomas, P.E., Zhang, H.,
Cui, Z., and Thannickal, V.J. (2004). Activation of the pro-survival
phosphatidylinositol 3-kinase/AKT pathway by transforming growth factor-beta1 in
mesenchymal cells is mediated by p38 MAPK-dependent induction of an autocrine
growth factor. J. Biol. Chem. *279*, 1359–1367.
https://doi.org/10.1074/jbc.M306248200.

54. Xu, Y., Tabe, Y., Jin, L., Watt, J., McQueen, T., Ohsaka, A., Andreeff, M., and
Konopleva, M. (2008). TGF-beta receptor kinase inhibitor LY2109761 reverses the
anti-apoptotic effects of TGF-beta1 in myelo-monocytic leukaemic cells co-cultured
with stromal cells. Br. J. Haematol. *142*, 192–201. https://doi.org/10.1111/j.1365-
2141.2008.07130.x.

55. Wang, X., and Zheng, X. (2022). scRNA-seq data in ovarian cancer. *1*.
https://doi.org/10.17632/rc47y6m9mp.1.

1634

1635

Graphical abstract

Figure 1: Tracking Clonal Lineage in Ovarian Cancer Metastasis
(**A**) Schematic depicting main elements of the MetTag library barcoding construct. (**B**) Schematic of injection and tissue collection timeline for the staggered injection experimental model using MetTag-barcoded ID8 ovarian cell lines. Schematic was partially created with BioRender.com. (**C**) Genomic DNA (gDNA)-level barcode distributions in the omentum graphed as frequency under WT, NRG, and MAC depletion conditions. (**D-E**) gDNA-level barcode distributions in ascites and pelvic fat pads under WT and NRG conditions. In (C-E) panels, BC.ID1 = TP-1, BC.ID2 = TP-2, BC.ID3 = TP-3, BC.ID4 = TP-4, BC.ID5 = TP-5, BC.ID6 = TP-6 for clarity. (**F**) Injection schema for comparing one-time (400,000 total, n = 10) cell dose and fractionated (2,400,000 total, n = 10) cell injections (left) and representative BLI images at endpoint (right). (**G**) Bar graph depicting total flux emission in each group at experimental endpoint. (**H**) Schematic of scRNA-seq experiment conducted using a reversed BC.ID injection order. LARRY barcodes are readily captured by Chromium 10x Feature Barcoding. (**I**) Pie charts of RNA-level barcode frequencies in each location grouped as early (BC.ID5, BC.ID6), mid (BC.ID4, BC.ID3), and late (BC.ID2, BC.ID1). (**J**) Heatmap of sample level barcode frequencies, BC.ID6 = TP-1, BC.ID5 = TP-2, BC.ID4 = TP-3, BC.ID3 = TP-4, BC.ID2 = TP-5, BC.ID1 = TP-6. (**K**) Schematic (top) and MetTag BC.ID distribution (bottom) for independent TNBC lung metastasis model depicting enrichment of scRNA-seq derived early TP DTCs in the metastatic lung at endpoint. Data in Figure 1G was analyzed using a two-tailed Student's t test and graphed as mean with standard deviation (SD).

Figure 2: Clonal Analysis Reveals Shared Evolutionary Path Towards Inflammatory Signature
(**A**) Split UMAP depicting cluster distributions across the four experimental conditions and identifying cancer cell-enriched clusters in ascites and omentum. (**B**) Reclustered cancer cell UMAP colored by the two metastasis-bearing anatomical locations. (**C**) Stacked bar chart depicting the frequency of Seurat subclusters within each metastatic location. (**D**) Seurat cluster UMAP of reclustered cancer cells with transcriptional program labels. (**E**) Dot plot depicting Hallmark pathways across all subclusters used to define each cluster subset gene program in (D). (**F**) Schematic representation of the possible outcomes of LARRY clonal analysis based on transcriptome patterns of enriched clones. (**G**) Heatmap of normalized clonal LARRY barcode counts across the two anatomical locations and individual mice (HTOs). (**H**) UMAP overlay of top 3 expanded clones in the omentum. (**I**) Bar chart depicting the relative subcluster proportions present in each of the top 3 barcode clones. (**J**) scVelo-generated PAGA velocity graph indicating trajectory arrows and inferred cluster lineage. (**K**) scVelo and CellRank-based simulated path from start to endpoint clusters. (**L**) Violin plot depicting RNA-level contribution of each timepoint to the IFN-gamma signature. Data in Figure 2L was analyzed by first defining the IFN-gamma signature using canonical mouse genes, followed by directly applying the wilcox.test() function in Seurat to compare TP-1 and TP-6 expression.

Figure 3: Activation of IFNγ Response in Ascites is Necessary for Successful Omental Colonization
(**A**) DEG-based volcano plot of gene expression differences between OmMet and AscMet, where threshold for significance was defined at p <0.05 and log2FC < -1.0 for downregulation or >1.0 for upregulation. (**B**) Boxplots comparing IFN signature modules between AscMet and OmMet. (**C**) Schematic describing in vivo ovarian metastatic niche signature screen. Partially created with BioRender.com. (**D**) *In vivo* CRISPR screen RankView plots for all 100 genes generated using MAGeCKFlute. OmMet (n=5) and AscMet (n=4) samples were compared to the original pool of ID8 cells. (**E**) Ranks of individual sgRNAs targeting positive control Pten and depleted genes. (**F**) Lollipop plot of depleted pathways associated with depleted genes in Fig. 3E. (**G**) IFNγ treatment time course qPCR plots of select CRISPR hits and positive controls. (**H**) IFNγ treatment time course western blot probing for Stat1/pStat1 and GBP1. (**I**) Schematic of in vivo experimental metastasis study comparing sgNTC and sgIFNGR1.g1 groups. (**J**) Time course log10-transformed total emission/flux values (left). Two independent groups pooled; cohort 1 NTC; n = 8, KO; n = 7; cohort 2 NTC; n = 5, KO; n = 5. log10-transformed flux plot for week six with p value (right). (**K**) Kaplan Meier survival curve generated for cohort 1. (**L**) Experimental schematic for staggered sgIfngr1 (*n = 4*) or sgNTC1 (*n = 3*) followed by sgNTC2-cell injection. (**M**) Representative endpoint BLI images of metastasis bearing mice in each group (left) and endpoint BLI quantification comparing tumor burden between sgIfngr1 and sgNTC first injection (right). (**N**) Heatmap showing enrichment of first DTC seed - either sgNTC1 or sgIfngr1 - and 2nd DTC seed, sgNTC2. DEG data in Figure 3A was analyzed using the FindMarkers() function. Data in Figure 3B was analyzed by first defining custom IFN signatures, followed by applying wilcox.test() in Seurat to compare module scores between AscMet and OmMet. CRISPR screen related data (3D-3F) was analyzed using MAGeCK and MAGeCKFlute. Data in 3G, 3J, 3K, and 3M was analyzed and graphed in Prism. 3G, SJ, and 3M were analyzed using a standard t test; 3G and 3M data was plotted as mean with SD, while 3J was plotted as median. Survival data in 3K was computed using a standard log-rank test.

Figure 4: IFNγ Response Mediates Resistance to Anoikis and Activates Pro-survival Programs
(**A**) Cell-Titer Glo (CTG) assay luminescence values graphed as fold change for regular culture conditions (left) and ultra-low attachment (ULA) conditions (right), with and without IFNγ treatment. (**B**) Volcano plots generated from bulk RNA-seq depicting top DEGs between sgNTC (wild type) cells under regular culture conditions with or without IFNγ treatment (left) and under ULA conditions with and without IFNγ treatment (right). (**C**) Venn diagram showing 25 overlapping and 30 unique upregulated genes in sgNTC cells treated with IFNγ under ULA conditions. (**D**) Heatmap of vst-normalized gene counts for ULA-specific upregulated genes. (**E**) Bar plot depicting top 15 Reactome pathways upregulated under ULA conditions in cells treated with IFNγ. (**F-G**) CTG flux dot plots for Stat1-KO (F) and Irf1-KO (G) cells under ultra-low attachment, with or without IFNγ treatment. (**H**) Violin plots comparing expression level of immunoproteasome (Psmb10, Psmb9), DNA repair/apoptosis (Parp12, Parp14, Parp9), and immune checkpoint (Cd274) levels between OmMet and AscMet. (**I**) scRNA-seq based gating strategy used to identify Irf1^high and Irf1^low cells for module scoring. (**J**) Violin plots comparing expression level of *Cd274* (left), *Psmb10* (middle), and *Parp14* (right) between Irf1^high and Irf1^low cells. Fold change in F4A was determined by normalizing each cell line's CTG value to the regular attachment, without IFNγ treatment.CTG assay luminescence data (4A) was graphed as mean and SD, data in 4F and 4G was graphed as median values. scRNA-seq based transcript expression levels were compared between Irf1^high and Irf1^low cells and OmMet/AscMet cells by applying the wilcox.test() function in Seurat. RNA-seq data was analyzed using DESeq2 and visualized as variance-stabilized counts.

Figure 5: Niche-specific Transcriptional States of Omental and Ascites-Derived TAMs

(**A**) Violin plots depicting CRISPR screen hits expression level across Broad cell ID categories (cell types). (**B**) CellChat-generated circus plots of interaction strengths between Broad cell ID categories in metastatic omentums (left) and ascites (right). (**C**) Stacked bar chart of cell type proportions across the four experimental groups. (**D**) UMAPs of reclustered myeloid cells, split by experimental condition. (**E**) UMAP of subclustered myeloid cells overlayed with scVelo RNA velocity arrows. Arrows are pointing from ascites TAMs (clusters 1 and 5) to omentum TAMs (clusters 0 and 6). (**F**) DEG volcano plot comparing gene expression differences between ascites TAM cluster 1 and omentum TAM cluster 0. (**G**) Stacked bar chart of subsetted T/NK cells comparing cell proportions in ascites and omentum. (**H-I**) Dot plots depicting Reactome pathways enriched in ascital (H) and omentum-derived TAMs (I). Significance of DEGs in Figure 5F was evaluated by applying the FindMarkers() and Wilcoxon test in Seurat to compare ascites and omental Cd68^high/Adgre1^high macrophages.

Figure 6: Peritoneal Macrophages are Direct DTC Pro-Survival Interactors

(**A**) Liposomal chlodronate-based macrophage depletion strategy and timeline schematic with representative tissue images. Omental tumor burden quantification and flow-based ascites quantification (bottom). Control group; n= 7, and Chlodrosome; n = 7. (**B**) Schematic describing Cre-based NicheTracing labeling system where ID8 and E0771 cancer cells express Cre.ERT2. Upon nuclear entry of Cre.ERT2 into Ai6 reporter niche cells, labeled niche cells turn ZsGreen positive. (**C**) Representative gating strategy depicting the major cell population tagged with zsG. (**D**) Quantification of F4/80+ and TREM2 mean fluorescent intensity (MFI) of contacted, ZsGreen high cells. (**E**) Volcano plot comparing DEGs between metastasis-infiltrating Trem2-low and Trem2-high macrophages. (**F**) Schematic depicting mouse model for macrophage-specific Trem2 knockout. (**G**) Log transformed BLI-derived flux values in Trem2$^{fl/fl}$ (wild type; n = 5) and Lys2-Cre/Trem2$^{fl/fl}$ (knockout; n = 5) mice at metastasis endpoint. Mice were injected with ID8 p53-/- Luciferase cells. (**H**) Schematic of macrophage-cancer cell coculture layout (top) and representative coculture images for one cell line (bottom).(**I**) Plots of percent CD45 negative cancer cell viability (ZombieAqua negative) post coculture in three cell lines. (**J**) Stacked bar chart of main cell type proportions captured in human scRNA-seq dataset across five anatomical locations, using maintype-2 annotation provided by publisher. (**K**) Venn diagram of top NicheNet predicted ligands after filtering for robust expression and overlapping with top ligands in our murine dataset. (**L**) Heatmap comparing enrichment of a curated list of receptors in bulk RNA-seq comparing ULA-cultured cells treated with IFNγ and regular attachment conditions. Data in Figures 6D was plotted as mean with SD and differences between groups were compared via a two-tailed unpaired Student's t test. Data in Figures 6I was similarly analyzed using a Student's t test and plotted with median values. Data in Figure 6E was analyzed using the FindMarkers() function in Seurat and Wilcoxon test to compare gated Trem2-low and Trem2-high cells. RNA-seq heatmap data in 6L was analyzed using DESeq2 and visualized as variance-stabilized counts.