Obsessive-Compulsive Tendencies Shift the Balance Between Competitive Neurocognitive Functions

Bianka Brezóczki [1,2,3], Teodóra Vékony [4]*, Bence Csaba Farkas [5,6,7], Flóra Hann [1,2,3,8], Dezső Németh [4,9,10] *

[1] Doctoral School of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary
[2] Institute of Psychology, ELTE Eötvös Loránd University, Budapest, Hungary
[3] Brain, Memory and Language Research Group, Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary
[4] Gran Canaria Cognitive Research Center, Department of Education and Psychology, Atlántico Medio University, Las Palmas de Gran Canaria, Spain
[5] Institut du Psychotraumatisme de l'Enfant et de l'Adolescent, Conseil Départemental Yvelines et Hauts-de-Seine et Centre Hospitalier des Versailles, Versailles, France
[6] UVSQ, Inserm, Centre de Recherche en Epidémiologie et Santé des Populations, Université Paris-Saclay, Versailles, France
[7] LNC2, Département d'études Cognitives, École Normale Supérieure, INSERM, PSL Research University, Paris, France
[8] Institute of Experimental Medicine, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary
[9] BML-NAP Research Group, Institute of Psychology, Eötvös Loránd University and Institute of Cognitive Neuroscience and Psychology, HUN-REN Research Centre for Natural Sciences, Budapest, Hungary
[10] Centre de Recherche en Neurosciences de Lyon CRNL U1028 UMR5292, INSERM, CRNS, Université Claude Bernard Lyon 1, 69500 Bron, France

*These authors contributed equally to this work.

Correspondence: Correspondence should be addressed to Dezső Németh at dezso.nemeth[at]inserm.fr

Highlights

- OCD theories propose an imbalance between automatic and goal-directed control systems.
- We tested whether this imbalance emerges at a functional level, beyond reward-based learning.
- Statistical learning and cognitive flexibility showed an antagonistic relationship in a non-clinical sample.
- This inverse association weakened as obsessive–compulsive tendencies increased.
- OC tendencies alter the interaction between automatic learning and executive control even at subclinical levels.

Summary

Theoretical models of Obsessive–Compulsive Disorder (OCD) emphasize that symptoms may arise from an imbalance between habitual and goal-directed processes, characterized by increased reliance on habitual behavior and reduced efficiency of goal-directed control. However, it remains unclear whether similar alterations appear at a more general functional level, beyond reward-driven mechanisms. The present study, therefore, investigated the relationship between statistical learning (SL), an implicit, reward-independent mechanism that supports the detection of environmental regularities, and cognitive flexibility, defined as the capacity to adapt behavior and cognitive strategies to changing environmental demands. By adopting a dimensional approach to obsessive-compulsive (OC) tendencies in a non-clinical sample, we aimed to clarify how continuous symptom variability relates to the interaction of these neurocognitive processes. A total of 404 participants completed an online probabilistic sequence learning task assessing SL and a card-sorting task measuring cognitive flexibility. Results revealed an antagonistic relationship between SL and cognitive flexibility. Importantly, this inverse association weakened as OC tendencies increased, suggesting that OC tendencies may alter the typical balance between automatic and goal-directed functions even at undiagnosed levels.

Keywords: statistical learning, cognitive flexibility, obsessive-compulsive disorder, spectrum approach, competitive memory systems

64   Graphical Abstract

Introduction

A core challenge in human cognition involves maintaining a dynamic interplay between efficient automatic and flexible goal-directed functions to support adaptive behavior. Goal-directed behaviour relies on a broad set of executive functions – including sustained attention, cognitive flexibility, inhibitory control, and working memory updating [1–3]. In contrast, automatic behaviour depends on previously acquired associations and operates with reduced conscious engagement [4–6]. Disruption of this interplay can lead to behavioral difficulties, as seen in conditions such as Obsessive-Compulsive Disorder (OCD), where automaticity tends to dominate over goal-directed control, resulting in rigid, repetitive mental (obsessions) and behavioral rituals (compulsions)[7–12,58,59]. Indeed, varying degrees of obsessions and compulsive acts, collectively referred to as obsessive-compulsive (OC) tendencies, are distributed along a continuum in the general population and, when persistent and sufficiently severe, culminate in a clinical presentation of OCD [13–21]. Considering OC tendencies from a dimensional perspective provides a more precise and comprehensive identification of the neurocognitive mechanisms underlying OCD, effectively overcoming the limitations of traditional categorical diagnoses that often obscure the continuous and nuanced variability of these symptoms [11,13,22,23,27]. Therefore, the present study aimed to delineate the role of the balance between automatic and goal-directed functions underlying maladaptive symptoms of OCD by assessing OC tendencies as a dimensional spectrum within the non-clinical population.

Prominent theoretical models propose that the repetitive symptoms of OCD and transdiagnostic tendencies arise from diminished goal-directed control coupled with a corresponding dominance of habitual responding [8,9,12,25,26,28,32,58]. Crucially, much of the empirical support for this perspective derives from dual-task paradigms that impose substantial cognitive demands by requiring the simultaneous coordination of multiple processes under state-dependent conditions, thereby illustrating how goal-directed and habitual systems interact under high cognitive load [8,9,28,29]. Within such contexts, disruptions in the balance between controlled and habitual processing may emerge, reflecting context-dependent processing constraints rather than indicating a generalized alteration in cognitive functioning. However, these state-level findings leave open the question of whether compulsive tendencies also reflect a more stable, trait-like bias in the interaction between controlled and automatized learning mechanisms. Individuals with generally weaker cognitive control may rely more consistently on simplified or habitual processing strategies not only under high task demands but also as part of a broader compensatory organization of cognitive functioning across contexts. The present study therefore aims to examine whether individual differences are systematically associated with performance across tasks that differentially engage controlled and automatized processing, with the goal of identifying stable, trait-related neurocognitive patterns underlying OC tendencies.

From a habitual perspective, a key limitation of current frameworks becomes apparent. Evidence from outcome devaluation and related feedback-based paradigms consistently points to enhanced habitual responding in OCD and compulsivity[10,12,26]. Yet in these tasks, the balance between goal-directed and habitual control is tightly coupled to continuous external reward or feedback, making it difficult to isolate learning mechanisms

from reward-driven performance. Consequently, understanding OC requires moving beyond
reward-dependent paradigms to probe the underlying, feedback-independent mechanisms [20].
Addressing this question requires considering recent advances in learning theory, which
emphasize the crucial role of unsupervised processes in driving neural plasticity and
establishing the foundation for supervised learning [30.] From this perspective, supervised
learning emerges as a later process built upon earlier, reward-free, unsupervised mechanisms
[30]. Accordingly, to better understand the development and persistence of automatic,
compulsive behavioral characteristics of OCD and OC tendencies, it is essential to
disentangle feedback sensitivity and pure learning mechanisms [20] and to revisit the primary
role of unsupervised learning and its interaction with goal-directed functions.
One such unsupervised learning mechanism is statistical learning (SL), a fundamental
function that supports the acquisition of skills and habits through the incidental extraction of
novel probabilistic patterns from the environmental stream [6,31,33–35]. SL in clinical OCD
populations has shown both impairments and intact results (for review see: Brezóczki et al.
[36]), while non-clinical individuals with OC tendencies demonstrate intact or even faster
learning rates [20]. This highlights the need to better understand how other neurocognitive
functions contribute to such performance. Indeed, within the framework of interactive
memory systems, neurocognitive functions such as SL rarely operate in isolation; instead,
they engage in dynamic and often competitive interactions with goal-directed functions [6,34,37–
42] Typically, increased reliance on one system corresponds with reduced engagement of the
other, as demonstrated by the inverse relationship between SL and goal-directed functions. In
this context, cognitive flexibility, defined as the capacity to shift behavior, strategies, or
mental frameworks in response to changing environmental demands, is a particularly critical
goal-directed function [1–3], as maladaptive, rigid, and repetitive behavioral patterns of OC may
reflect its impairments [43,44]. If OCD and OC tendencies involve a disruption in the balance
between automatic and goal-directed processes, then investigating their interaction outside
the influence of supervision becomes particularly informative.

To provide a complementary perspective on the neurocognitive profile underlying
OCD and OC tendencies, this study focused on SL and goal-directed functions, specifically
cognitive flexibility. Therefore, in the present study, we recruited 404 adult participants from
university courses via an online experiment. SL was assessed using an unsupervised learning
paradigm, the Alternating Serial Reaction Time (ASRT) task [45,46], which is a probabilistic
learning task that lacks the supervision- and reinforcement-based factors on SL performance
[47]. Cognitive flexibility was operationalized as the number of perseverative errors on the Card
Sorting Task [48], a widely accepted and extensively used measure of cognitive flexibility [49–55].
The prevalence of undiagnosed OC tendencies was evaluated using the OCI-12
questionnaire[49.] The present study aimed to uncover the interplay between SL and cognitive
flexibility and to investigate how non-clinical OC tendencies may modulate this relationship.
Based on existing literature, we hypothesized an antagonistic relationship between SL and
cognitive flexibility, such that stronger performance on the ASRT task would be associated
with weaker performance on the Card Sorting Task. Crucially, we further hypothesized that
OC tendencies would modulate this interaction, with the relationship between SL and

cognitive flexibility varying across different levels of OC tendencies and potentially
diverging from a straightforward competitive pattern at higher levels of OC tendencies.

Results

How do OC tendencies modulate the interplay between cognitive flexibility and statistical
learning? - Reaction Time

We performed a linear mixed-effects model on blockwise median RT of the ASRT task, with
fixed effects for Time (Block 1-15), Triplet Type (high- vs. low-probability), OCI-12 score,
Number of Perseverative errors of the Card Sorting Task, and all higher-order interactions.
Random effects included participant-specific intercepts and a random slope for Time to
account for individual variability. The statistical parameters underlying the effects reported
above are summarized in Table 1, whereas the full set of statistical estimates (b, SE, 95% CI,
t, df, and p values) is provided in Supplementary Table S1.
The model reported a statistically significant main effect of Triplet Type, indicating
faster RTs for high-probability trials compared to low-probability trials, which evidences
statistical learning during the task. The main effect of Time indicated a decrease in RTs
throughout the task, demonstrating changes in visuomotor performance over time.
Importantly, the interaction between Triplet Type and Time suggested that statistical learning
developed gradually, as the difference between high- and low-probability trials increased
over time.
Cognitive flexibility was significantly associated with overall RT performance, as
reflected by a main effect of Perseverative errors, whereby a higher number of perseverative
errors, indicative of reduced cognitive flexibility, predicted slower responses. However, the
absence of significant interactions involving Perseverative errors suggests that cognitive
flexibility did not modulate visuomotor performance (Time $\times$ Perseverative error), or
statistical learning (Triplet Type $\times$ Perseverative error), nor the trajectory of statistical
learning (Time $\times$ Triplet Type $\times$ Perseverative error).
In contrast, OCI-12 scores showed no significant main effect and did not interact with
any of the task-related variables, indicating that OC tendencies were not associated with
baseline RT (main effect of OCI-12 score), cognitive flexibility (OCI-12 score $\times$
Perseverative error), visuomotor performance (Time $\times$ OCI-12 score), statistical learning
(Triplet Type $\times$ OCI), the trajectory of statistical learning (Time $\times$ Triplet Type $\times$ OCI-12
score). Moreover, OCI-12 scores did not modulate the relationship between cognitive
flexibility (Perseverative error $\times$ Triplet Type $\times$ OCI-12 score, see: Figure 1) and the
dynamics of statistical learning in the present study (Perseverative error $\times$ Time $\times$ Triplet
Type $\times$ OCI-12 score).
186
187

Table 1. Results of the Linear Mixed-Effects Model predicting blockwise median Reaction Time

Effect	df	F	p
Time	1, 399.98	125.06	**< .001**
Triplet Type	1, 11297.07	169.21	**< .001**
OCI-12 score	1, 400.01	3.15	.077
Perseverative error	1, 400.02	7.31	**.007**
Time × Triplet Type	1, 11297.10	10.42	**.001**
Time × OCI-12 score	1, 399.88	0.15	.699
Triplet Type × OCI-12 score	1, 11297.11	0.00	.926
Time × Perseverative error	1, 400.07	3.13	.078
Triplet Type × Perseverative error	1, 11297.07	0.44	.508
OCI-12 score × Perseverative error	1, 400.01	0.26	.600
Time × Triplet Type × OCI-12 score	1, 11297.18	0.09	.763
Time × Triplet Type × Perseverative error	1, 11297.1	0.92	.338
Time × OCI-12 score × Perseverative error	1, 400.00	3.61	.058
Triplet Type × OCI-12 score × Perseverative error	1, 11297.09	1.29	.257
Time × Triplet Type × OCI-12 score × Perseverative error	1, 11297.15	0.67	.412

Note. The table summarizes all main effects and interactions. Degrees of freedom *(df)* correspond to Satterthwaite approximations. F = F-statistic; p = significance value. Statistically significant results ($p < .05$) are shown in bold. Significant main effects were followed up with post hoc comparisons: The main effect of Triplet Type (High vs. Low) was significant ($b = -5.87$, SE = 0.45, z = -13.01, $p < .001$, 95% CI [-6.76, -4.99]), indicating faster responses to high- than to low-probability triplets. Time was also significant ($b = -1.1$, SE = 0.10, z = -11.18, $p < .001$, 95% CI [-1.29, -0.9]), showing that RT decreased as the task progressed. Statistically significant Perseverative Error indicated that more perseverative errors were associated with slower reaction times ($b = 1.44$, SE = 0.53, z = 2.70, $p = .007$, 95% CI [0.4, 2.49]). The significant Time × Triplet Type interaction ($b = -0.34$, SE = 0.10, z = -3.23, $p = .001$, 95% CI [-0.54, -0.13]) reflected gradual improvement in statistical learning over time. Time was treated as a continuous variable in all analyses; however, for the post hoc analyses, it was segmented into three time points ($Block_5$, $Block_{10}$, $Block_{15}$). $Block_5$ ($b = -4.86$, SE = 0.55, z = -8.84, $p < .001$, 95% CI [-5.94, -3.78]), $Block_{10}$ ($b = -6.55$, SE = 0.50, z = -13.16, $p < .001$, 95% CI [-7.52, -5.57]), and $Block_{15}$ ($b = -8.24$, SE = 0.86, z = -9.58, $p < .001$, 95% CI [-9.92, -6.55]). Sidak-adjusted pairwise comparisons revealed significant differences in statistical learning (High vs. Low differences) between $Block_5$–$Block_{10}$ ($b = 1.69$, SE = 0.52, z = 3.23, $p = .004$), $Block_5$–$Block_{15}$ ($b = 3.37$, SE = 1.05, z = 3.23, $p = .004$), and $Block_{10}$–$Block_{15}$ ($b = 1.69$, SE = 0.52, z = 3.23, $p = .004$).

Figure 1. The figure illustrates the relationship between cognitive flexibility, as measured by the Number of Perseverative errors on the Card Sorting Task along the x-axis, and statistical learning performance along the y-axis, which is operationalized as the difference in reaction time between high- and low-probability triplets. Each point represents an individual participant, with color encoding their total OCI-12 score along a continuous gradient from low (yellow) through intermediate (pink) to high (purple) obsessive-compulsive tendencies. A linear regression line with 95% confidence intervals is overlaid in black. While the OCI-12 score was treated as a continuous variable in all statistical analyses, for visualization purposes, the data are faceted by the levels of OC tendencies. These levels represent varying degrees of OCI-12 scores, approximating one standard deviation below the mean (Lower OC tendencies), at the mean (Medium OC tendencies), and one standard deviation above the mean (Higher OC tendencies), respectively. The color gradient used in the plot (yellow to pink to purple) visually reflects this increase in the level of OC tendencies. No relationship was observed between cognitive flexibility and RT-based statistical learning, and this pattern was consistent across the level of OC tendencies.

How do OC tendencies modulate the interplay between cognitive flexibility and statistical learning? – Accuracy

We performed a linear mixed-effects model on blockwise mean accuracy of the ASRT task, with fixed effects for Time (Block 1-15), Triplet Type (high- vs. low-probability), OCI-12 score, Number of Perseverative errors of the Card Sorting Task, and all higher-order interactions. Random effects included participant-specific intercepts and correlated slopes for Time and Triplet Type to account for individual variability. The statistical parameters underlying the effects reported above are summarized in Table 2, whereas the full set of statistical estimates (b, SE, 95% CI, t, df, and p values) is provided in Supplementary Table S2.

Participants were more accurate on high-probability trials than on low-probability trials, providing clear evidence of statistical learning (main effect of Triplet Type). Overall accuracy declined slightly over the course of the task (main effect of Time), but the difference between high- and low-probability trials increased, indicating a gradual improvement in statistical learning (Time × Triplet Type).

Cognitive flexibility influenced performance: a higher number of Perseverative errors
were associated with lower overall accuracy (main effect of Perseverative error), and a
modest decline in visuomotor performance over time (Time × Perseverative error). Moreover,
a significant Triplet Type × Perseverative error interaction revealed an antagonistic
relationship between statistical learning and cognitive flexibility (Triplet Type ×
Perseverative error), with greater statistical learning linked to reduced cognitive flexibility,
that is higher number of perseverative errors: ($b_{\text{Low perseverative errors}}$ = 1.95, 95% CI [1.59, 2.30];
$b_{\text{Mean perseverative errors}}$ = 2.22, 95% CI [1.97, 2.47]; $b_{\text{High perseverative errors}}$ = 2.50, 95% CI [2.13,
2.86]). Pairwise comparisons indicated that the statistical learning increased significantly
with perseverative errors ($b_{\text{Low perseverative errors}} \bar{\;} {}_{\text{Mean perseverative errors}}$ = −0.28, 95% CI [−0.53,
−0.02], p = .034; $b_{\text{Low perseverative errors}} \bar{\;} {}_{\text{High perseverative errors}}$ = −0.55, 95% CI [−1.06, −0.04], p =
.034; $b_{\text{Mean perseverative errors}} \bar{\;} {}_{\text{High perseverative errors}}$ = −0.28, 95% CI [−0.53, −0.02], p = .034). Thus,
individuals with poorer cognitive flexibility (higher numbers of perseverative errors)
exhibited enhanced statistical learning, except when considering the trajectory of learning
(Time × Triplet Type × Perseverative error), where this pattern was less pronounced.
OC tendencies did not significantly influence overall accuracy (main effect of OCI-12
score), visuomotor performance (Time × OCI-12 score), statistical learning (Triplet Type ×
OCI-12 score), or its improvement over time (Time × Triplet Type × OCI-12 score).
Similarly, no interactions were observed between OC tendencies and Perseverative errors for
overall accuracy or visuomotor performance (Time × Perseverative error × OCI-12 score).
However, a significant Triplet Type × Perseverative error × OCI-12 score interaction
indicated that OC tendencies modulate the antagonistic relationship between statistical
learning and cognitive flexibility (see: Figure 2). The post hoc analysis examining the trend
of statistical learning across perseverative error levels (M-SD, M, M+SD) revealed that
among individuals with lower OC tendencies, perseverative errors were positively associated
with statistical learning, with a significant slope (b = 0.145, SE = 0.048, z = 2.97, p = .002).
A similar but weaker positive relationship was observed at the medium level of OC
tendencies (b = 0.069, SE = 0.032, z = 2.11, p = .034), suggesting that poorer cognitive
flexibility was linked to enhanced statistical learning among lower and medium level of OC
tendencies. In contrast, this association disappeared among individuals with higher OC
tendencies, where the slope was non-significant (b = −0.006, SE = 0.040, z = −0.17, p =
.862). Overall, statistical learning increased as the number of perseverative errors rose for
individuals with lower to medium OC tendencies, reflecting an antagonistic pattern in which
reduced cognitive flexibility was accompanied by better statistical learning. This antagonistic
pattern weakened in participants with high OC tendencies, where statistical learning remained
moderate regardless of cognitive flexibility. Importantly, this modulation was not observed
when examining the full trajectory of learning over time (Triplet Type × Time ×
Perseverative error × OCI-12 score).
275

276 **Table 1. Results of the Linear Mixed-Effects Model predicting blockwise mean accuracy**

Effect	df	F	p
Time	1, 400.22	58.82	**< .001**
Triplet Type	1, 400.14	302.7	**< .001**
OCI-12 score	1, 399.97	0.00	.976
Perseverative error	1, 400.06	6.18	**.013**
Time × Triplet Type	1,10898.51	33.15	**< .001**
Time × OCI-12 score	1, 400.05	0.01	.935
Triplet Type × OCI-12 score	1, 399.91	0.89	.347
Time × Perseverative error	1, 400.39	4.41	**.036**
Triplet Type × Perseverative error	1, 400.52	4.48	**.035**
OCI-12 score × Perseverative error	1, 400.03	0.91	.341
Time × Triplet Type × OCI-12 score	1, 10898.03	0.98	.321
Time × Triplet Type × Perseverative error	1, 10899.01	0.04	.845
Time × OCI-12 score × Perseverative error	1, 400.26	2.30	.130
Triplet Type × OCI-12 score × Perseverative error	1, 400.30	6.21	**.013**
Time × Triplet Type × OCI-12 score × Perseverative error	1,10898.65	1.81	.178

Note. The table summarizes all main effects and interactions. Degrees of freedom *(df)* correspond to
Satterthwaite approximations. *F* = F-statistic; *p* = significance value. Statistically significant results *(p < .05)* are
shown in bold. Significant main effects were followed up with post hoc comparisons: The main effect of Triplet
Type indicated higher accuracy for high- than for low-probability triplets (b = 2.22, SE = 0.13, z = 17.40, *p* <
.001, 95% CI [1.97, 2.47]). Time was also significant (b = -0.15, SE = 0.02, z = -7.67, *p* < .001, 95% CI [-0.18, -
0.11]), showing that accuracy decreased as the task progressed. The main effect of Perseverative error indicated
that more perseverative errors were associated with lower accuracy (b = -0.11, SE = 0.04, z = -2.49, *p* = .013,
95% CI [-0.2, -0.02]) . Time × Triplet Type interaction revealed improvement in statistical learning over time.
Time was treated as a continuous variable in all analyses; however, for the post hoc analyses, it was segmented
into three time points (Block$_5$, Block$_{10}$, Block$_{15}$). Block$_5$ (b = 1.75, SE = 0.15, z = 11.56, *p* < .001, 95% CI [1.46,
2.05]), Block$_{10}$ *(b* = 2.54, SE = 0.14, z = 18.26, *p* < .001, 95% CI [2.26, 2.81]), and Block$_{15}$ (b = 3.32, SE = 0.23,
z = 14.48, *p* < .001, 95% CI [2.87, 3.77]). Sidak-adjusted pairwise comparisons revealed significant differences
between Block$_5$–Block$_{10}$ (b = -0.78, SE = 0.52, z = 3.23, p = .004), Block$_5$–Block$_{15}$ (b = -1.57, SE = 1.05, z =
3.23, *p* = .004), and Block$_{10}$–Block$_{15}$ (b = -0.78, SE = 1.05, z = 3.23, *p* = .004). For the Time × Perseverative
Error interaction, Sidak-adjusted pairwise comparisons showed significant declines over time at lower
perseverative error levels (b = -0.11, SE = 0.03, z = -3.92, *p* < .001, 95% CI [-0.16, -0.05]; *b* = -0.15, SE = 0.02,
z = -7.67, *p* < .001, 95% CI [-0.18, -0.11]) and at higher perseverative error levels (b = -0.19, SE = 0.03, z = -
6.79, *p* < .001, 95% CI [-0.24, -0.13]), indicating a decline in visuomotor performance over time. Perseverative
Error was treated as a continuous variables in all primary analyses; however, for the post hoc analyses they were
depicted in three categories (M–SD, M, M+SD).

Figure 2. The figure illustrates the relationship between cognitive flexibility, as measured by the number of perseverative errors on the Card Sorting Task along the x-axis, and statistical learning performance along the y-axis, which is operationalized as the difference in accuracy between high- and low-probability triplets. Each point represents an individual participant, with color encoding their total OCI-12 score along a continuous gradient from low (yellow) through intermediate (pink) to high (purple) obsessive-compulsive tendencies. A linear regression line with 95% confidence intervals is overlaid in black. While the OCI-12 score was treated as a continuous variable in all statistical analyses, for visualization purposes, the data are faceted by the levels of OC tendencies. These levels represent varying degrees of OCI-12 scores, approximating one standard deviation below the mean (Lower OC tendencies) at the mean (Medium OC tendencies), and one standard deviation above the mean (Higher OC tendencies), respectively. The color gradient used in the plot (yellow to pink to purple) visually reflects this increase in the level of OC tendencies. The figure highlights the moderating role of OC tendencies in the antagonistic association between statistical learning and cognitive flexibility. Specifically, the typical inverse relationship, where reduced cognitive flexibility is associated with enhanced statistical learning, was observed in participants with Lower and Medium OC tendencies. In contrast, this pattern was not present among individuals with Higher OC tendencies, for whom statistical learning remained relatively stable across the cognitive flexibility continuum. This suggests a disruption of the typical trade-off between these cognitive functions in the context of OC tendencies.

Discussion

In this study, we addressed a critical gap in understanding the relationship between goal-directed and automatic behaviour by examining how unsupervised SL interacts with cognitive flexibility. We specifically investigated how this relationship is modulated by OC tendencies within a non-clinical adult population with a focus on stable, trait-related organizational strategies in cognitive processing. We conducted online experiments on an undiagnosed population, measuring OC tendencies along a continuous spectrum from mild to elevated levels. SL was evaluated using an unsupervised learning task while cognitive flexibility was assessed with the Card Sorting Task. Our findings corroborate previous theories indicating a competitive interaction between SL and executive functions, specifically cognitive flexibility. Crucially, we found that this inverse relationship progressively

weakened as OC tendencies increased, suggesting that, even at undiagnosed levels, OC tendencies can disrupt the optimal balance between SL and goal-directed functions.

Our results build upon existing theories that highlight a disruption between automatic and goal-directed processes in both clinical OCD and subclinical OC tendencies or compulsivity [8–12,23,26] In particular, the typical antagonistic relationship between these functions, as emphasized by competitive memory system frameworks [6,34,37–39,41,42,56,57] appears to weaken with increasing levels of non-clinical OC tendencies. This pattern may appear partly inconsistent with the habit hypothesis [8,9,26,58,59], which primarily predicts a context-dependent shift toward habitual control. However, methodological differences may partly account for these discrepancies since they predominantly employed outcome devaluation or contingency degradation paradigms, typically assessing learning after extended instrumental training and evaluating goal-directed behavior based on participants' ability to adapt to altered rules and feedback within a single experimental context [8,9,26]. Such paradigms require the simultaneous integration of multiple neurocognitive processes under high cognitive demand, which may bias performance toward habitual responding. In contrast, the present study indicates that higher OC tendencies are associated with reduced cognitive flexibility, constraining statistical learning rather than enhancing. From this perspective, the observed interplay between SL performance and cognitive flexibility at higher levels of undiagnosed OC tendencies represents task-specific interference and reflects a stable organizational characteristic of behavioral adaptation across contexts.

Taken together, these findings suggest that the pattern observed at the trait level diverges from the state-dependent bias toward habitual responding described in prior work, raising the possibility of a dissociation between transient task-related effects and more stable neurocognitive organization across the OC spectrum. Whereas state-level paradigms have primarily revealed increased reliance on habitual control under high cognitive demand [8,9,26], the present results indicate that increasing OC tendencies are associated with a weakening of the relationship between executive control and statistical learning. Thus, individuals with weaker executive functions, specifically reduced cognitive flexibility, demonstrate enhanced acquisition of environmental statistical regularities at lower levels of OC tendencies, whereas at higher levels, the same reduction in flexibility is associated with impaired SL. We can speculate that the interaction between SL and cognitive flexibility may reflect an individual's position along the OC continuum: at lower levels of OC tendencies, reduced flexibility may facilitate the efficient extraction of statistical regularities, whereas at higher levels, increased rigidity appears to constrain learning. More broadly, these findings suggest that as OC tendencies increase, the typical antagonism between SL and executive functions may become attenuated, such that automatic functions are less able to compensate for diminished flexibility. Elucidating this potential dissociation will require future research to disentangle trait-level and state-dependent contributions to this interaction and to determine how it shapes the balance between learning mechanisms and executive control across the OC spectrum.

Beyond demonstrating a trait-dependent interaction between neurocognitive systems, our findings address an important gap by elucidating the interaction between unsupervised learning, a primary form of learning [30], and executive functions among OC tendencies. Consistent with prior findings, unsupervised learning was found to be preserved [20,60–63] suggesting this core learning mechanism remains robust across OC tendencies similar to

OCD, although not all prior findings are fully aligned with this pattern [64–68]. This preservation
is particularly noteworthy, as the excessive bias toward habitual behaviour in OCD may
partly stem from external reward-driven learning. In feedback- or reward-driven contexts, it
is difficult to disentangle whether performance improvements reflect sensitivity to external
reinforcement or strengthened stimulus–response associations. Moreover, individuals with
OCD and OC tendencies may reach a ceiling effect more rapidly due to heightened reward or
feedback sensitivity, thereby accelerating the shift toward enhanced automatic processes and
reducing the opportunity for flexible behavioral adaptation. Although the pure learning
mechanism remained intact, the interaction of SL with goal-directed functions became
atypical among higher OC tendencies. This suggests that the disruption lies not within the
pure learning mechanisms themselves, but in the dynamic interplay between these systems
among OC tendencies and possibly, in OCD [9,26,29,69.] Future research should further
investigate the state-dependent interaction between habitual and goal-directed processes in
relation to unsupervised learning mechanisms, particularly in the absence of external reward.
This would clarify the fundamental relationship between habitual and goal-directed functions
and their contribution to OC- and OCD-related behavioral manifestations.

On the other hand, previous approaches emphasizing impairments in goal-directed or
executive functions in OCD have primarily examined these processes as distinct functional
domains [43,60,61,65,72,74], without sufficiently considering how these alterations emerge within a
broader neurocognitive network. In contrast, our findings align with the perspective that
neither habitual, automatic functions nor cognitive inflexibility alone serve as robust
predictors of OC symptomatology; rather, it is the interaction between these factors,
specifically, reduced flexibility combined with heightened automaticity, that better accounts
for symptom presence [8,9,19,71,77,79]. But unlike a previous study, which measured generative
cognitive flexibility and assessed habitual tendencies via self-report questionnaires [19], our
study specifically examined reactive cognitive flexibility, which a recent meta-analysis has
challenged as being independently impaired in OCD [70]This may explain why, unlike previous
findings showing an inverse relationship, we observe a weakened rather than inverse
association between these processes at higher levels of OC tendencies. To advance
understanding of OC tendencies and the pathophysiology of OCD, future studies should
endeavor to replicate these findings using dual-task paradigms that concurrently integrate
learning mechanisms and cognitive flexibility within a dual-task design.

Our findings underscore the importance of investigating not only individual
neurocognitive functions but also their dynamic interplay in advancing our understanding of
the roots of OC symptomatology. Our results imply that the weakened interplay observed
among higher OC tendencies could serve as a potential predictor of OC symptomatology
[9,11,19,26,59]. Alongside theoretical frameworks that propose a bias toward automatic processes
in clinical OCD [9,10,19,25,26,29,59,71] other perspectives emphasize the potential emergence of
compensatory mechanisms, wherein impairments in implicit learning are offset by increased
engagement of controlled, explicit processes to maintain effective performance [36,60,61,64,73,75]
These findings were supported by both behavioral and neuroimaging evidence, showing that
during unsupervised behavioral tasks [60,61] neural activity was predominantly associated with
hippocampal and medial temporal lobe regions rather than striatal structures [62,63] While the
typical antagonistic interaction between these systems is maintained among lower levels of

OC tendencies, this pattern appears to weaken as tendencies increase within the non-clinical
OC spectrum. This raises the possibility that, in clinical populations, such a shift could result
in a more cooperative or compensatory way of interaction. Our findings align with this
perspective, as they suggest that the characteristics of both clinical OCD and subclinical OC
tendencies may stem from atypical interactions between neurocognitive systems [9,60,61,64,73,75].
Furthermore, it can be speculated that whether these systems engage in competitive or
cooperative interactions depends on the specific cognitive demands of the task, suggesting a
context-sensitive dynamic rather than a static relationship. Therefore, future research is
needed to determine whether this altered cognitive interplay persists in clinical OCD or
whether, as hypothesized, compensatory mechanisms reshape the dynamic toward a more
cooperative and functionally adaptive interaction in clinical OCD.
Another noteworthy finding is that the unrevealed interplay with cognitive flexibility
appeared only in the accuracy-based performance indicator of the ASRT task. This
dissociation is particularly interesting as it raises the question of which performance
components are more sensitive to OC tendencies. One possible explanation is that the card-
sorting task relies on accuracy without imposing temporal constraints [48]. Accuracy- and rule-
based learning generally require more effortful processing [76,78,80], which may account for the
antagonistic relationship between the two measures. Specifically, enhanced implicit
adaptation to rule-based structures may come at the cost of reduced performance in tasks
requiring explicit rule detection, an effect that aligns with the competition and cooperation
framework outlined above [6,34,38,39,41,57]. This pattern also reflects the clinical features of OCD,
where symptoms often involve an excessive focus on order, regularity, and precision, such as
checking, arranging, or striving for symmetry [7,81]. Consequently, accuracy-based measures
may be especially sensitive to the influence of OC tendencies. Taken together, these findings
suggest that OC symptoms may interfere more strongly with learning processes grounded in
precision and rule detection, offering a clearer picture of how cognitive rigidity manifests in
the disorder.
It is important to note that our study was conducted on an undiagnosed, so-called non-
clinical population of healthy adults. Therefore, while findings from clinical OCD research
provide an important theoretical background, our results should not be generalized to clinical
OCD populations but rather interpreted within a transdiagnostic framework. At the same
time, spectrum-based approaches offer valuable theoretical and methodological contributions
to the deeper understanding of clinical conditions, including the characteristics of OCD
[11,13,20,22,24,82]. Importantly, previous research has demonstrated that OC tendencies can
influence a broad range of neurocognitive functions, including statistical learning [20] inhibitory
control [83], cognitive flexibility[19],and decision-making [84,85] This can prove that this approach
serves as an effective methodological tool for identifying individual differences in
neurocognitive functions. The current study contributes to this growing body of literature by
highlighting how OC tendencies may influence not just cognitive domains in isolation but the
dynamic interplay between them, offering a more integrated understanding of the
neurocognitive profile associated with OC tendencies. Our findings indicate that while a
dynamic interplay among neurocognitive functions is maintained at lower levels of OC
tendencies, this interaction may potentially be absent in the clinical populations, resulting in
impaired adaptive behavior. This disruption could be anticipated by higher subclinical OC

tendencies. It is also possible that this inverse relationship between these functions persists at
the clinical level but manifests in an exaggerated, potentially nonlinear manner, consistent
with a U-shaped function across the OC spectrum. To clarify these dynamics, it is essential to
extend investigations to the clinical population in future research.

Some limitations should be noted in the present study. First, the relatively high
exclusion rate, a common challenge in online experimental designs [20,57,82,86–88], may have
affected the generalizability of the findings. Although rigorous data quality checks were
essential to obtain high-quality data, the inclusion criterion requiring the absence of any self-
reported psychiatric or neurological history likely limited the representation of individuals
with more pronounced OC tendencies. As a result, the upper range of the OC spectrum may
have been underrepresented, potentially narrowing the scope of our conclusions. We did not
control for other conditions that are frequently associated with elevated compulsivity and are
often comorbid with OC tendencies, including anxiety and depressive symptomatology
[11,89,90]. Consequently, the present results do not allow us to fully isolate effects specifically
related to the OC symptom dimension from those potentially driven by related affective or
psychopathological factors. Presenting the tasks in a fixed order may have introduced
potential order or fatigue effects. Future studies should therefore consider counterbalancing
task order to minimize such influences, as performance on later tasks may be affected by
declining attention or cognitive fatigue. In addition, no attention checks were included during
the administration of the OCI-12 questionnaire, which may have increased the likelihood of
inattentive responding [91]. Moreover, although cognitive flexibility was assessed using a card
sorting task, this measure has inherent methodological limitations [92,93]. Therefore, future
research would benefit from incorporating multiple measures of cognitive flexibility to obtain
a more comprehensive assessment of executive functioning. Finally, the present study was
conducted in a university-based adult sample, which may limit the generalizability of the
findings to the broader population. Accordingly, future studies should extend these
investigations to more diverse age groups and community samples to strengthen external
validity.

To conclude, our results support the competitive memory systems hypothesis, that
emphasize an antagonistic relationship between neurocognitive systems [6,34,37,41,42] by
elucidating a more general, trait-dependent dynamic interplay between SL and goal-directed
processes, here cognitive flexibility, and how their interaction is modulated by the prevalence
of OC tendencies in a non-clinical population. While lower prevalence of OC tendencies may
facilitate adaptive neurocognitive interactions, higher OC tendencies might represent early
neurocognitive markers predictive of increased vulnerability to OCD. These results
underscore the significance of conceptualizing OC tendencies as transdiagnostic traits to
enhance early identification and inform tailored interventions. Importantly, our study
highlights the necessity for cognitive neuroscience to transcend the examination of isolated
neurocognitive functions and instead prioritize the investigation of their interactive dynamics,
as the integrity of individual processes alone may mask critical dysfunctions emerging from
their interactions.

Resource availability

Lead contact

Further information and requests for resources should be directed to and will be fulfilled by Prof. Dezső Németh, dezso.nemeth@inserm.fr.

Materials availability

This study did not generate new unique reagents.

Data availability statement

Data and codes for analysis are available on the following link:

https://osf.io/fwb38/?view_only=03b2754e647742a2a8e5158a39174b45

Acknowledgement

ANR-24-CE37-5807) (to D.N.); the National Brain Research Program project NAP2022-I-2/2022 (to D.N.), EKÖP-24-3-II-ELTE-1159 (to B.B); the Spanish Ministry of Science, Innovation and Universities (MICIU), the State Research Agency (AEI), and the European Regional Development Fund (FEDER, UE) through the grant PID2024-160183NA-I00 (MICIU/AEI/10.13039/501100011033/FEDER, UE) (to T.V.)

Contributions

B. B.: Conceptualization, Software, Validation, Investigation, Data Curation, Formal analysis, Writing – Original Draft, Writing – Review & Editing, Visualization; T.V.: Conceptualization, Software, Validation, Investigation, Data Curation, Formal analysis, Resources, Writing – Review & Editing, Supervision; B. C. F.: Methodology, Validation, Writing – Review & Editing, F. H.: Writing – Review & Editing; D.N.: Conceptualization, Software, Validation, Resources, Writing – Review & Editing, Supervision, Funding.

Declaration of interests

The authors declare no competing interests.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work the author(s) used ChatGPT 4.0 to improve the writing style of our proposal. The author(s) reject using AI for scientific purposes to gain unethical benefits. However, the author(s) believe that it helps the equality of native and non-native English speakers to have the same opportunities. Thus, the author(s) do recommend using generative AI for lingual purposes (only). After using this tool/service, the author(s) reviewed and edited the content as needed and take(s) full responsibility for the content of the publication.

STAR METHODS

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

A total of 568 university students took part in the study through an online experiment in exchange for course credit. The sample shows partial participant overlap with those reported

in Brezóczki et al. [20]Hann et al. [94], and Nagy et al. [86]; however, the datasets are not identical,
as data were collected across multiple studies with different primary research aims and
measurement protocols. To maintain data integrity, rigorous quality control procedures were
implemented, resulting in the exclusion of 164 participants who failed to follow the
instructions. Our goal was to meet multiple data quality criteria, which are outlined in detail
in the 'Quality control of the data' section below. The final sample consisted of 404
participants (285 female, 111 male, and 3 non-specified; $M_{Age} = 22.06$ years; $SD_{Age} = 5.09$).
This study did not perform analyses based on sex or gender, which could limit the
generalizability of the findings. Participants provided informed consent, and the studies
received approval from the Research Ethics Committee of Eötvös Loránd University,
Budapest, Hungary (2021/504), following the principles of the Declaration of Helsinki.
559

560 **METHOD DETAILS**
561

562 *Alternating Serial Reaction Time task*
563

Statistical learning was measured by the Alternating Serial Reaction Time (ASRT) task [45,46].
The task was implemented in JavaScript using jsPsych version 6.1.0 [95]. Each trial presented a
visual stimulus, an illustration of a dog's head, in one of four horizontally aligned screen
positions (Figure 3A). Participants were instructed to respond as quickly and accurately as
possible by pressing the corresponding keyboard key ("S," "F," "J," or "L" from left to right).
The stimulus remained visible until a correct response was given, followed by a 120 ms
response-to-stimulus interval before the next trial. Incorrect responses resulted in the stimulus
remaining on the screen until corrected. Unbeknownst to participants, the stimuli followed a
probabilistic eight-element sequence alternating between pattern and random elements (e.g.,
2–r–4–r–3–r–1–r, where numbers denote fixed positions and "r" indicates a random one)
(Figure 3B). This alternating structure resulted in some three-element combinations (triplets)
occurring more frequently than others. High-probability triplets – such as 2–X–4, 4–X–3, 3–
X–1, and 1–X–2, where X is the variable middle element – occurred more often due to their
consistent alignment with the underlying sequence. High-probability triplets emerged either
when two pattern elements enclosed a random element (50% of trials) or when two random
elements enclosed a pattern element (12.5% of trials) (Figure 3C). Making up the remaining
37.5% of trials, low-probability triplets included less predictable element combinations based
on the sequence structure.
The ASRT task consisted of 15 blocks, each containing 10 repetitions of an eight-element
sequence, for a total of 80 stimuli per block. After each block, participants received feedback
on their average accuracy and reaction time, but not on their statistical learning performance.
Still, no information was provided about their statistical learning performance, as they were
unaware that the task involved detecting a hidden regularity.
587
588

589 *Card Sorting Task*
590

Cognitive flexibility was assessed using the Card Sorting Task (CST), adapted from the Wisconsin Card Sorting Test (WCST), a well-established measure of executive function [48,96,97] The CST evaluates an individual's ability to shift cognitive strategies in response to changing task demands. During the task, four reference cards are displayed simultaneously on the screen. Each card contains visual stimuli that differ along three dimensions: color (red, yellow, green, or blue), quantity (one to four items), and shape (triangle, star, diamond, or circle). In each trial, a separate target card appears beneath the reference cards (Figure 3D). Participants are instructed to match the target card with one of the four reference cards according to a common feature, either color, quantity, or shape, by selecting the corresponding card. Importantly, the correct matching rule is not disclosed to the participant. Instead, participants are required to infer the rule based on feedback provided after each response, which indicates whether the selected card was correct or incorrect (Figure 3E). The task comprises a total of sixty-four trials. The underlying categorization rule changes after every sequence of ten consecutive correct responses. The order of rule changes follows a fixed sequence: first based on color, then shape, followed by quantity, and then repeats in the same order. Cognitive flexibility is measured by the number of perseverative errors, defined as repeated use of a previously correct but no longer valid rule. A lower number of perseverative errors indicates greater cognitive flexibility.

Figure 3. Design of the Study. (A) In the Alternating Serial Reaction Time (ASRT) task, participants were instructed to respond by pressing keys corresponding to the spatial location of a target stimulus, depicted as a dog's head. Each trial followed an eight-element probabilistic sequence composed of alternating pattern and random elements. For example, a sequence such as 2–r–4–r–3–r–1–r consisted of fixed pattern trials at positions 2, 4, 3, and 1 (from left to right), interleaved with random elements, denoted by 'r', which could appear at any of the four possible locations. **(B)** Each trial was considered the third element in a sequence of three consecutive trials (a triplet). Pattern elements, which consistently appeared in fixed positions throughout the task, were displayed with red backgrounds, whereas random elements, selected randomly from the four possible locations, were shown with blue backgrounds. The structure of the sequence resulted in certain triplets occurring more frequently than others. These were categorized as high-probability triplets, while less frequent combinations were defined as low-probability triplets. **(C)** For illustrative purposes, three triplets are highlighted: 2(P)–1(R)–

4(P), a pattern-ending high-probability triplet; 2(R)–3(P)–4(R), a random-ending high-probability triplet; and
4(R)–1(P)–2(R), a random-ending low-probability triplet. High-probability triplets could occur in one of two
ways: either with two pattern trials and a random trial in the middle (accounting for 50% of all trials), or with
two random trials and a pattern trial in the middle (accounting for 12.5% of trials). Overall, 62.5% of trials
constituted the final element of a high-probability triplet, whereas the remaining 37.5% served as the final
element of a low-probability triplet. **(D)** Schematic illustration of the Card Sorting Task (CST). Four reference
cards are presented simultaneously, differing along three stimulus dimensions (color, shape, and quantity). On
each trial, a target card appears below the reference cards, and participants are instructed to match it to one of
the reference cards according to an sorting rule. **(E)** The rules are presented in a fixed, repeating order for up to
six categories: Color, Shape, Number. The sorting rule changes after ten consecutive correct responses.
Cognitive flexibility is indexed by the number of perseverative errors, defined as continued application of a
previously correct but currently invalid rule.

634
635

Obsessive-Compulsive Inventory

Obsessive-compulsive tendencies were assessed using the 12-item version of the Obsessive-
Compulsive Inventory (OCI-12, [98,99]), a self-report questionnaire measuring the severity of
symptoms on a 5-point Likert scale ranging from 0 (not at all) to 4 (extremely). Participants
reported how much each symptom had bothered them during the previous month. The OCI-
12 has demonstrated strong psychometric validity, including high internal consistency and
test-retest reliability, in both clinical and non-clinical samples [98]. The Cronbach's alpha in our
sample indicated good internal consistency ($\alpha = 0.89$). The total score can vary from 0 to 48.
In our final sample, the observed scores ranged from 0 to 43 (M = 12.58; SD = 9.02); see
Figure 4. For reference, the full 18-item OCI-R was also administered and is provided in the
Supplementary Materials.

649
Figure 4. The distribution of the participants' OC tendencies as measured by the OCI-12 total score in
the sample. The x-axis indicates the total OCI-12 score, while the y-axis indicates the number of participants.
The color gradient transitions from yellow, through pink, to purple, reflecting increasing levels of OC
tendencies.

Procedures

The experiment was conducted using the Gorilla Experiment Builder platform
(https://www.gorilla.sc) [100]. It began with participants reading an informational briefing and
providing informed consent. Subsequently, participants completed the ASRT task, which
started with two practice blocks followed by 15 task blocks. The practice blocks consisted
solely of random trials and were excluded from analysis. To assess participants' awareness of
any patterns, we administered post-task inquiries asking them to report any regularities they
may have noticed during the ASRT task. Specifically, participants were questioned about
their detection of hidden sequences or patterns and asked to elaborate if applicable. None
were able to accurately describe the alternating sequence, indicating that the task remained
implicit, consistent with prior research [34,101] Following the ASRT task, cognitive flexibility
was evaluated using the CST, and obsessive-compulsive tendencies were measured with the
OCI-12. Additionally, demographic data on socioeconomic status, sleep, health, and
handedness were collected. To account for potential confounding variables associated with
the online setting, participants provided concise details about their testing environment,
which are thoroughly outlined in the 'Quality Control of the Data' section.
672
673 **Quality control of the data**
674
Online data collection is known to present several methodological challenges, as outcomes
may be influenced not only by the psychological constructs of interest but also by variability
in participants' response styles, engagement levels, and task completion strategies [91]. To
address these concerns, we implemented data quality control procedures following
established recommendations for online experimental research [88]. From the initial sample, we
excluded participants who: 1) reported prior completion of the ASRT task (N = 6, 1.06%), 2)
restarted or quit the experiment at any point (N = 28, 4.93%); 3) failed the attention test
screening for non-compliance (N = 26, 4.40%); 4) performed their mean raw accuracy on the
ASRT task was below 80%, as such low performance was considered indicative of
insufficient task engagement rather than genuine performance (N = 24, 4.23%); 5) age
exceeding 55 years (N = 1, 0.17%), 6) reported neurological impairment or head injury, or
reported a diagnosis of any psychiatric condition (N = 58, 10.21%); 7) reported having
consumed alcohol within 6 hours, recreational drugs or psychoactive medications within 24
hours before the experiment (N = 38, 6.69%); 8) took long self-paced breaks between ASRT
blocks, as such extended interruptions may indicate reduced task engagement (N = 4, 0.7%),
9) random buttom pressing during ASRT task (N = 3, 0.53%) and 10) participants who
scored in the zero category on the CST, indicating random clicking during the task or
incomplete understanding of it (N = 19, 3.34%). As some participants met multiple exclusion
criteria, the combined criteria resulted in the exclusion of a total of 164 participants
(28.87%).
695
696 **QUANTIFICATION AND STATISTICAL ANALYSIS**
697
698 **Data preparation and analysis**
699

Data preprocessing, statistical analysis, and visualization were carried out in R version 4.4.2
[102] For the ASRT task, each trial was categorized according to whether it represented the final
element of a high- or low-probability triplet. Only the first response for each trial was
retained; subsequent responses were excluded to ensure response validity. Trials with
reaction times (RTs) below 150 ms or above 1000 ms, as well as those falling beyond three
absolute median deviations from each participant's median RT, were removed to control for
outliers (Brezóczki et al., 2025). Repetition trials (e.g., 2–2–2, 1–1–1) and trills (e.g., 2–1–2,
3–2–3) were excluded due to their potential to elicit prepotent motor responses unrelated to
statistical learning processes [34,103]. The first two trials of each block were also removed, as
they did not complete a triplet structure. Furthermore, incorrect responses were excluded
from RT analyses. Following these procedures, 17.54% of trials were excluded from
accuracy analyses and 24.61% from RT analyses. These exclusion rates are consistent with
those reported in online ASRT studies [20,57,82,86,87].
We utilized Linear Mixed Models (LMMs) for analyzing RT and accuracy performance.
LMMs were fit with the mixed function from the afex package [104].Regarding statistical
learning and visuomotor performance, we computed the median reaction times and mean
accuracy of the ASRT task for each participant in every block. These measures were used as
outcome variables in linear mixed models with the Time (Block 1–15), Triplet Type (last
element of a high- vs. low-probability triplet), Cognitive flexibility (Perseverative error), OC
tendencies (OCI-12 score), and their higher-order interactions as fixed effects in the model.
The Time, OCI-12 score, and Perseverative error factors were treated as mean-centered
continuous variables. The models also included participant-specific intercepts and correlated
slopes for the within-subject variables of Time and Triplet Type, treated as random effects.
For RTs, the final model achieved convergence using the participants as random intercepts
with by-participant correlated slopes for the Block. For accuracy, the final model achieved
convergence using the participants as random intercepts with by-participant correlated slopes
for Triplet Type and Time. Additional analyses conducted on further variables (Trials to
Complete First Category, Total Number of Correct Responses) are reported in the
Supplementary Information (Tables S5–S8).
Post hoc analyses involved the computation of estimated marginal means and
marginal trends for simple effects and simple slopes, utilizing the emmeans package [105].
Significant interactions involving continuous predictors, specifically the OCI-12 score and
Perseverative errors, were probed at three levels: the mean, as well as at ±1 standard
deviation from their means. For the Time factor, probing was performed at the end of the
5th, 10th, and 15th blocks. Figures were created with the ggplot2.
In reporting the results, the main effects of Time are interpreted as improvement in
visuomotor performance, both on RT and accuracy, regardless of probabilistic patterns.
Meanwhile, the main effect of Triplet Type is seen as an indicator of the amount of implicit
probabilistic learning, and the interaction between Time and Triplet Type represents the
temporal trajectory of statistical learning throughout the task. Please note that, although we
used the OCI-12 score as a continuous variable for all our analyses, for more illustrative data
visualization and clarified post hoc analysis, we depicted results at lower OC tendencies
(Mean OCI-12 score - SD), medium OC tendencies (Mean OCI-12 score) and higher OC
tendencies (Mean OCI-12 score + SD), based on standard deviations from the mean. For the

more complex interactions, the same approach was applied to the Number of Perseverative
errors.
746
747 Hence, our models were structured according to the following general form in *afex* syntax:
748
749 Reaction time:
750 *MedianRT ~ Time * Triplet Type * OCI-12 score * Perseverative error + (Time | Participant)*
751
752 Accuracy:
753 *MeanAccuracy ~ Time * Triplet Type * OCI-12 score * Perseverative error + (Time +*
754 *Triplet Type | Participant)*
755

References
1. Diamond, A. (2013). Executive Functions. Annu. Rev. Psychol. *64*, 135–168. https://doi.org/10.1146/annurev-psych-113011-143750.
2. Friedman, N.P., and Robbins, T.W. (2022). The role of prefrontal cortex in cognitive control and executive function. Neuropsychopharmacology *47*, 72–89. https://doi.org/10.1038/s41386-021-01132-0.
3. Miyake, A., Friedman, N.P., Emerson, M.J., Witzki, A.H., Howerter, A., and Wager, T.D. (2000). The Unity and Diversity of Executive Functions and Their Contributions to Complex "Frontal Lobe" Tasks: A Latent Variable Analysis. Cogn. Psychol. *41*, 49–100. https://doi.org/10.1006/cogp.1999.0734.
4. Ashby, F.G., Turner, B.O., and Horvitz, J.C. (2010). Cortical and basal ganglia contributions to habit learning and automaticity. Trends Cogn. Sci. *14*, 208–215. https://doi.org/10.1016/j.tics.2010.02.001.
5. Kaufman, S.B., DeYoung, C.G., Gray, J.R., Jiménez, L., Brown, J., and Mackintosh, N. (2010). Implicit learning as an ability. Cognition *116*, 321–340. https://doi.org/10.1016/j.cognition.2010.05.011.
6. Pedraza, F., Farkas, B.C., Vékony, T., Haesebaert, F., Phelipon, R., Mihalecz, I., Janacsek, K., Anders, R., Tillmann, B., Plancher, G., et al. (2024). Evidence for a competitive relationship between executive functions and statistical learning. NPJ Sci. Learn. *9*, 30. https://doi.org/10.1038/s41539-024-00243-9.
7. American Psychiatric Association (2022). Diagnostic and Statistical Manual of Mental Disorders (American Psychiatric Association Publishing) https://doi.org/10.1176/appi.books.9780890425787.
8. Gillan, C.M., Apergis-Schoute, A.M., Morein-Zamir, S., Urcelay, G.P., Sule, A., Fineberg, N.A., Sahakian, B.J., and Robbins, T.W. (2015). Functional Neuroimaging of Avoidance Habits in Obsessive-Compulsive Disorder. American Journal of Psychiatry *172*, 284–293. https://doi.org/10.1176/appi.ajp.2014.14040525.
9. Gillan, C.M., Papmeyer, M., Morein-Zamir, S., Sahakian, B.J., Fineberg, N.A., Robbins, T.W., and de Wit, S. (2011). Disruption in the Balance Between Goal-Directed Behavior and Habit Learning in Obsessive-Compulsive Disorder. American Journal of Psychiatry *168*, 718–726. https://doi.org/10.1176/appi.ajp.2011.10071062.
10. Gillan, C.M., and Robbins, T.W. (2014). Goal-directed learning and obsessive–compulsive disorder. Philosophical Transactions of the Royal Society B: Biological Sciences *369*, 20130475. https://doi.org/10.1098/rstb.2013.0475.
11. Gillan, C.M., Kosinski, M., Whelan, R., Phelps, E.A., and Daw, N.D. (2016). Characterizing a psychiatric symptom dimension related to deficits in goal-directed control. Elife *5*. https://doi.org/10.7554/eLife.11305.
12. Yu, Q., Gao, F., Li, C., Xia, J., Cao, Y., Wang, X., Xiao, C., Lu, J., Liu, Q., Fan, J., et al. (2024). Compulsion is associated with impaired goal-directed and habitual learning and responding in obsessive-compulsive disorder. International Journal of Clinical and Health Psychology *24*, 100531. https://doi.org/10.1016/j.ijchp.2024.100531.
13. Abramowitz, J.S., Fabricant, L.E., Taylor, S., Deacon, B.J., McKay, D., and Storch, E.A. (2014). The relevance of analogue studies for understanding obsessions and compulsions. Clin. Psychol. Rev. *34*, 206–217. https://doi.org/10.1016/j.cpr.2014.01.004.

14. Hajcak, G., Huppert, J.D., Simons, R.F., and Foa, E.B. (2004). Psychometric properties of the OCI-R in
a college sample. Behaviour Research and Therapy *42*, 115–123.
https://doi.org/10.1016/j.brat.2003.08.002.
15. Foa, E.B., Huppert, J.D., Leiberg, S., Langner, R., Kichic, R., Hajcak, G., and Salkovskis, P.M. (2002).
The Obsessive-Compulsive Inventory: Development and validation of a short version. Psychol. Assess.
14, 485–496. https://doi.org/10.1037/1040-3590.14.4.485.
16. Fullana, M.A., Vilagut, G., Rojas-Farreras, S., Mataix-Cols, D., de Graaf, R., Demyttenaere, K., Haro,
J.M., de Girolamo, G., Lépine, J.P., Matschinger, H., et al. (2010). Obsessive–compulsive symptom
dimensions in the general population: Results from an epidemiological study in six European countries.
J. Affect. Disord. *124*, 291–299. https://doi.org/10.1016/j.jad.2009.11.020.
17. Mataix-Cols, D., do Rosario-Campos, M.C., and Leckman, J.F. (2005). A Multidimensional Model of
Obsessive-Compulsive Disorder. American Journal of Psychiatry *162*, 228–238.
https://doi.org/10.1176/appi.ajp.162.2.228.
18. Mataix-Cols, D., Junqué, C., Sànchez-Turet, M., Vallejo, J., Verger, K., and Barrios, M. (1999).
Neuropsychological functioning in a subclinical obsessive-compulsive sample. Biol. Psychiatry *45*,
898–904. https://doi.org/10.1016/S0006-3223(98)00260-1.
19. Ramakrishnan, S., Robbins, T.W., and Zmigrod, L. (2022). Cognitive Rigidity, Habitual Tendencies,
and Obsessive-Compulsive Symptoms: Individual Differences and Compensatory Interactions. Front.
Psychiatry *13*. https://doi.org/10.3389/fpsyt.2022.865896.
20. Brezóczki, B., Farkas, B.C., Hann, F., Pesthy, O., Tóth-Fáber, E., Farkas, K., Csigó, K., Németh, D.,
and Vékony, T. (2025). Individual differences in probabilistic learning and updating predictive
representations in individuals with obsessive-compulsive tendencies. BMC Psychiatry *25*, 368.
https://doi.org/10.1186/s12888-025-06786-4.
21. Zermatten, A., Van der Linden, M., Jermann, F., and Ceschi, G. (2006). Validation of a French version
of the Obsessive–Compulsive Inventory-Revised in a non-clinical sample. European Review of Applied
Psychology *56*, 151–155. https://doi.org/10.1016/j.erap.2005.07.003.
22. Gillan, C.M., and Rutledge, R.B. (2021). Smartphones and the Neuroscience of Mental Health. Annu.
Rev. Neurosci. *44*, 129–151. https://doi.org/10.1146/annurev-neuro-101220-014053.
23. Gillan, C.M., Kalanthroff, E., Evans, M., Weingarden, H.M., Jacoby, R.J., Gershkovich, M., Snorrason,
I., Campeas, R., Cervoni, C., Crimarco, N.C., et al. (2020). Comparison of the Association Between
Goal-Directed Planning and Self-reported Compulsivity vs Obsessive-Compulsive Disorder Diagnosis.
JAMA Psychiatry *77*, 77. https://doi.org/10.1001/jamapsychiatry.2019.2998.
24. Gillan, C.M., Fineberg, N.A., and Robbins, T.W. (2017). A trans-diagnostic perspective on obsessive-
compulsive disorder. Psychol. Med. *47*, 1528–1548. https://doi.org/10.1017/S0033291716002786.
25. Graybiel, A.M., and Rauch, S.L. (2000). Toward a Neurobiology of Obsessive-Compulsive Disorder.
Neuron *28*, 343–347. https://doi.org/10.1016/S0896-6273(00)00113-6.
26. Gillan, C.M., Morein-Zamir, S., Urcelay, G.P., Sule, A., Voon, V., Apergis-Schoute, A.M., Fineberg,
N.A., Sahakian, B.J., and Robbins, T.W. (2014). Enhanced Avoidance Habits in Obsessive-Compulsive
Disorder. Biol. Psychiatry *75*, 631–638. https://doi.org/10.1016/j.biopsych.2013.02.002.
27. Colzato, L., Zhang, H., Roessner, V., Beste, C., and Hommel, B. (2025). Non-bivalent psychopathology:
Rethinking mental disorders through metacontrol. Neurosci. Biobehav. Rev. *176*, 106297.
https://doi.org/10.1016/j.neubiorev.2025.106297.
28. Gillan, C.M., Morein-Zamir, S., Urcelay, G.P., Sule, A., Voon, V., Apergis-Schoute, A.M., Fineberg,
N.A., Sahakian, B.J., and Robbins, T.W. (2014). Enhanced avoidance habits in obsessive-compulsive
disorder. Biol. Psychiatry *75*, 631–638.
29. Robbins, T.W., Vaghi, M.M., and Banca, P. (2019). Obsessive-Compulsive Disorder: Puzzles and
Prospects. Neuron *102*, 27–47. https://doi.org/10.1016/j.neuron.2019.01.046.
30. Zhong, L., Baptista, S., Gattoni, R., Arnold, J., Flickinger, D., Stringer, C., and Pachitariu, M. (2025).
Unsupervised pretraining in biological neural networks. Nature. https://doi.org/10.1038/s41586-025-
09180-y.
31. Aslin, R.N. (2017). Statistical learning: a powerful mechanism that operates by mere exposure. WIREs
Cognitive Science *8*. https://doi.org/10.1002/wcs.1373.

32. Banca, P., Herrojo Ruiz, M., Gonzalez-Zalba, M.F., Biria, M., Marzuki, A.A., Piercy, T., Sule, A.,
Fineberg, N.A., and Robbins, T.W. (2024). Action sequence learning, habits, and automaticity in
obsessive-compulsive disorder. Elife *12*. https://doi.org/10.7554/eLife.87346.4.

33. Conway, C.M. (2020). How does the brain learn environmental structure? Ten core principles for
understanding the neurocognitive mechanisms of statistical learning. Neurosci. Biobehav. Rev. *112*,
279–299. https://doi.org/10.1016/j.neubiorev.2020.01.032.

34. Horváth, K., Nemeth, D., and Janacsek, K. (2022). Inhibitory control hinders habit change. Sci. Rep. *12*,
8338. https://doi.org/10.1038/s41598-022-11971-6.

35. Theeuwes, J., Bogaerts, L., and van Moorselaar, D. (2022). What to expect where and when: how
statistical learning drives visual selection. Trends Cogn. Sci. *26*, 860–872.
https://doi.org/10.1016/j.tics.2022.06.001.

36. Brezóczki, B., Vékony, T., Pesthy, O., Tóth-Fáber, E., Csigó, K., Farkas, K., and Nemeth, D. (2023).
Unraveling sequence learning in obsessive–compulsive disorder. Curr. Opin. Behav. Sci. *54*, 101326.
https://doi.org/10.1016/j.cobeha.2023.101326.

37. Ambrus, G.G., Vékony, T., Janacsek, K., Trimborn, A.B.C., Kovács, G., and Nemeth, D. (2020). When
less is more: Enhanced statistical learning of non-adjacent dependencies after disruption of bilateral
DLPFC. J. Mem. Lang. *114*, 104144. https://doi.org/10.1016/j.jml.2020.104144.

38. Nemeth, D., Janacsek, K., Polner, B., and Kovacs, Z.A. (2013). Boosting Human Learning by Hypnosis.
Cerebral Cortex *23*, 801–805. https://doi.org/10.1093/cercor/bhs068.

39. Virag, M., Janacsek, K., Horvath, A., Bujdoso, Z., Fabo, D., and Nemeth, D. (2015). Competition
between frontal lobe functions and implicit sequence learning: evidence from the long-term effects of
alcohol. Exp. Brain Res. *233*, 2081–2089. https://doi.org/10.1007/s00221-015-4279-8.

40. Tóth-Fáber, E., Janacsek, K., Szőllősi, Á., Kéri, S., and Nemeth, D. (2021). Regularity detection under
stress: Faster extraction of probability-based regularities. PLoS One *16*, e0253123.
https://doi.org/10.1371/journal.pone.0253123.

41. Pedraza, F., Vékony, T., Farkas, B.C., Haesebaert, F., Phelipon, R., Mihalecz, I., Janacsek, K.,
Tillmann, B., Anders, R., Plancher, G., et al. (2024). The interplay between executive functions and
updating predictive representations. Preprint, https://doi.org/10.1101/2024.12.05.626969
https://doi.org/10.1101/2024.12.05.626969.

42. Poldrack, R.A., Clark, J., Paré-Blagoev, E.J., Shohamy, D., Creso Moyano, J., Myers, C., and Gluck,
M.A. (2001). Interactive memory systems in the human brain. Nature *414*, 546–550.
https://doi.org/10.1038/35107080.

43. Chamberlain, S.R., Fineberg, N.A., Menzies, L.A., Blackwell, A.D., Bullmore, E.T., Robbins, T.W., and
Sahakian, B.J. (2007). Impaired Cognitive Flexibility and Motor Inhibition in Unaffected First-Degree
Relatives of Patients With Obsessive-Compulsive Disorder. American Journal of Psychiatry *164*, 335–
338. https://doi.org/10.1176/ajp.2007.164.2.335.

44. Tomiyama, H., Nakao, T., Murayama, K., Nemoto, K., Ikari, K., Yamada, S., Kuwano, M., Hasuzawa,
S., Togao, O., Hiwatashi, A., et al. (2019). Dysfunction between dorsal caudate and salience network
associated with impaired cognitive flexibility in obsessive-compulsive disorder: A resting-state fMRI
study. Neuroimage Clin. *24*, 102004. https://doi.org/10.1016/j.nicl.2019.102004.

45. Farkas, B.C., Krajcsi, A., Janacsek, K., and Nemeth, D. (2023). The complexity of measuring reliability
in learning tasks: An illustration using the Alternating Serial Reaction Time Task. Behav. Res. Methods
56, 301–317. https://doi.org/10.3758/s13428-022-02038-5.

46. Howard, J.H., and Howard, D. V. (1997). Age differences in implicit learning of higher order
dependencies in serial patterns. Psychol. Aging *12*, 634–656. https://doi.org/10.1037/0882-
7974.12.4.634.

47. Nemeth, D., and Tóth-Fáber, E. (2026). Resolving Fundamental Debates in Statistical Learning: The
Power of Process Dissociation. Preprint, https://doi.org/10.31234/osf.io/uwa35_v1
https://doi.org/10.31234/osf.io/uwa35_v1.

48. Fox, C.J., Mueller, S.T., Gray, H.M., Raber, J., and Piper, B.J. (2013). Evaluation of a Short-Form of the
Berg Card Sorting Test. PLoS One *8*, e63885. https://doi.org/10.1371/journal.pone.0063885.

49.   Zhang, Z., Yang, L.-Z., Vékony, T., Wang, C., and Li, H. (2024). Split-half reliability estimates of an
898   online card sorting task in a community sample of young and elderly adults. Behav. Res. Methods *56*,
899   1039–1051.
900   50.   Baker, K.S., Gibson, S.J., Georgiou☐Karistianis, N., and Giummarra, M.J. (2018). Relationship
901   between self☐reported cognitive difficulties, objective neuropsychological test performance and
902   psychological distress in chronic pain. European Journal of Pain *22*, 601–613.
903   https://doi.org/10.1002/ejp.1151.
904   51.   Pesthy, Z.V., Berta, K., Vékony, T., Farkas, B.C., Németh, D., and Kun, B. (2026). Dissociating the
905   cognitive underpinnings of recreational cannabis use from problematic use. Compr. Psychiatry *147*,
906   152685. https://doi.org/10.1016/j.comppsych.2026.152685.
907   52.   Berta, K., Pesthy, Z.V., Vékony, T., Farkas, B.C., Németh, D., and Kun, B. (2023). The
908   neuropsychological profile of work addiction. Sci. Rep. *13*, 20090. https://doi.org/10.1038/s41598-023-
909   47515-9.
910   53.   Wollenhaupt, C., Wilke, L., Erim, Y., Rauh, M., Steins-Loeber, S., and Paslakis, G. (2019). The
911   association of leptin secretion with cognitive performance in patients with eating disorders. Psychiatry
912   Res. *276*, 269–277. https://doi.org/10.1016/j.psychres.2019.05.001.
913   54.   Gelonch, O., Garolera, M., Valls, J., Rosselló, L., and Pifarré, J. (2016). Executive function in
914   fibromyalgia: Comparing subjective and objective measures. Compr. Psychiatry *66*, 113–122.
915   https://doi.org/10.1016/j.comppsych.2016.01.002.
916   55.   Dickson, K.S., Ciesla, J.A., and Zelic, K. (2017). The Role of Executive Functioning in Adolescent
917   Rumination and Depression. Cognit. Ther. Res. *41*, 62–72. https://doi.org/10.1007/s10608-016-9802-0.
918   56.   Horváth, K., Nemeth, D., Janacsek, K., and Kóbor, A. Cooperative interaction between statistical
919   learning and inhibitory control.
920   57.   Vékony, T., Brezóczki, B., Csifcsák, G., Németh, D., and Simor, P. (2025). A functional trade-off
921   between executive control and implicit statistical learning is dynamically gated by mind wandering.
922   Preprint, https://doi.org/10.1101/2025.08.05.668618 https://doi.org/10.1101/2025.08.05.668618.
923   58.   Banca, P., Voon, V., Vestergaard, M.D., Philipiak, G., Almeida, I., Pocinho, F., Relvas, J., and Castelo-
924   Branco, M. (2015). Imbalance in habitual versus goal directed neural systems during symptom
925   provocation in obsessive-compulsive disorder. Brain *138*, 798–811.
926   https://doi.org/10.1093/brain/awu379.
927   59.   Gillan, C.M. (2021). Recent developments in the habit hypothesis of OCD and compulsive disorders.
928   The neurobiology and treatment of OCD: accelerating progress, 147–167.
929   60.   Soref, A., Liberman, N., Abramovitch, A., Poznanski, Y., and Dar, R. (2021). Intact capacity for
930   implicit learning in obsessive-compulsive disorder. J. Behav. Ther. Exp. Psychiatry *73*, 101667.
931   https://doi.org/10.1016/j.jbtep.2021.101667.
932   61.   Soref, A., Liberman, N., Abramovitch, A., and Dar, R. (2018). Explicit instructions facilitate
933   performance of OCD participants but impair performance of non-OCD participants on a serial reaction
934   time task. J. Anxiety Disord. *55*, 56–62. https://doi.org/10.1016/j.janxdis.2018.02.003.
935   62.   Rauch, S.L., Wedig, M.M., Wright, C.I., Martis, B., McMullin, K.G., Shin, L.M., Cannistraro, P.A., and
936   Wilhelm, S. (2007). Functional Magnetic Resonance Imaging Study of Regional Brain Activation
937   During Implicit Sequence Learning in Obsessive–Compulsive Disorder. Biol. Psychiatry *61*, 330–336.
938   https://doi.org/10.1016/j.biopsych.2005.12.012.
939   63.   Rauch, S.L., Savage, C.R., Alpert, N.M., Dougherty, D., Kendrick, A., Curran, T., Brown, H.D., Manzo,
940   P., Fischman, A., and Jenike, M.A. (1997). Probing striatal function in obsessive-compulsive disorder: a
941   PET study of implicit sequence learning. Journal of Neuropsychiatry and Clinical Neurosciences *9*,
942   568–573.
943   64.   Deckersbach, T., Savage, C.R., Curran, T., Bohne, A., Wilhelm, S., Baer, L., Jenike, M.A., and Rauch,
944   S.L. (2002). A Study of Parallel Implicit and Explicit Information Processing in Patients With
945   Obsessive-Compulsive Disorder. American Journal of Psychiatry *159*, 1780–1782.
946   https://doi.org/10.1176/appi.ajp.159.10.1780.

65.   Kathmann, N., Rupertseder, C., Hauke, W., and Zaudig, M. (2005). Implicit Sequence Learning in
Obsessive-Compulsive Disorder: Further Support for the Fronto-Striatal Dysfunction Model. Biol.
Psychiatry 58, 239–244. https://doi.org/10.1016/j.biopsych.2005.03.045.

66.   Goldman, B.L., Martin, E.D., Calamari, J.E., Woodard, J.L., Chik, H.M., Messina, M.G., Pontarelli,
N.K., Marker, C.D., Riemann, B.C., and Wiegartz, P.S. (2008). Implicit learning, thought-focused
attention and obsessive-compulsive disorder: A replication and extension. Behaviour Research and
Therapy 46, 48–61. https://doi.org/10.1016/j.brat.2007.10.004.

67.   Marker, C.D., Calamari, J.E., Woodard, J.L., and Riemann, B.C. (2006). Cognitive self-consciousness,
implicit learning and obsessive–compulsive disorder. J. Anxiety Disord. 20, 389–407.
https://doi.org/10.1016/j.janxdis.2005.03.003.

68.   Vloet, T.D., Marx, I., Kahraman-Lanzerath, B., Zepf, F.D., Herpertz-Dahlmann, B., and Konrad, K.
(2010). Neurocognitive Performance in Children with ADHD and OCD. J. Abnorm. Child Psychol. 38,
961–969. https://doi.org/10.1007/s10802-010-9422-1.

69.   Gruner, P., Anticevic, A., Lee, D., and Pittenger, C. (2016). Arbitration between Action Strategies in
Obsessive-Compulsive Disorder. The Neuroscientist 22, 188–198.
https://doi.org/10.1177/1073858414568317.

70.   Fradkin, I., Strauss, A.Y., Pereg, M., and Huppert, J.D. (2018). Rigidly Applied Rules? Revisiting
Inflexibility in Obsessive Compulsive Disorder Using Multilevel Meta-Analysis. Clinical Psychological
Science 6, 481–505. https://doi.org/10.1177/2167702618756069.

71.   Vaghi, M.M., Cardinal, R.N., Apergis-Schoute, A.M., Fineberg, N.A., Sule, A., and Robbins, T.W.
(2019). Action-Outcome Knowledge Dissociates From Behavior in Obsessive-Compulsive Disorder
Following Contingency Degradation. Biol. Psychiatry Cogn. Neurosci. Neuroimaging 4, 200–209.
https://doi.org/10.1016/j.bpsc.2018.09.014.

72.   Menzies, L., Achard, S., Chamberlain, S.R., Fineberg, N., Chen, C.-H., del Campo, N., Sahakian, B.J.,
Robbins, T.W., and Bullmore, E. (2007). Neurocognitive endophenotypes of obsessive-compulsive
disorder. Brain 130, 3223–3236. https://doi.org/10.1093/brain/awm205.

73.   Ullman, M.T. (2004). Contributions of memory circuits to language: the declarative/procedural model.
Cognition 92, 231–270. https://doi.org/10.1016/j.cognition.2003.10.008.

74.   Penadés, R., Catalán, R., Rubia, K., Andrés, S., Salamero, M., and Gastó, C. (2007). Impaired response
inhibition in obsessive compulsive disorder. European Psychiatry 22, 404–410.
https://doi.org/10.1016/j.eurpsy.2006.05.001.

75.   Ullman, M.T., and Pullman, M.Y. (2015). A compensatory role for declarative memory in
neurodevelopmental disorders. Neurosci. Biobehav. Rev. 51, 205–222.

76.   Nomura, E., Maddox, W., Filoteo, J., Ing, A., Gitelman, D., Parrish, T., Mesulam, M.-M., and Reber, P.
(2006). Neural Correlates of Rule-Based and Information-Integration Visual Category Learning.
Cerebral Cortex 17, 37–43. https://doi.org/10.1093/cercor/bhj122.

77.   Vaghi, M.M., Luyckx, F., Sule, A., Fineberg, N.A., Robbins, T.W., and De Martino, B. (2017).
Compulsivity Reveals a Novel Dissociation between Action and Confidence. Neuron 96, 348-354.e4.
https://doi.org/10.1016/j.neuron.2017.09.006.

78.   Feng, X., Perceval, G.J., Feng, W., and Feng, C. (2020). High Cognitive Flexibility Learners Perform
Better in Probabilistic Rule Learning. Front. Psychol. 11. https://doi.org/10.3389/fpsyg.2020.00415.

79.   Vaghi, M.M., Vértes, P.E., Kitzbichler, M.G., Apergis-Schoute, A.M., van der Flier, F.E., Fineberg,
N.A., Sule, A., Zaman, R., Voon, V., Kundu, P., et al. (2017). Specific Frontostriatal Circuits for
Impaired Cognitive Flexibility and Goal-Directed Planning in Obsessive-Compulsive Disorder:
Evidence From Resting-State Functional Connectivity. Biol. Psychiatry 81, 708–717.
https://doi.org/10.1016/j.biopsych.2016.08.009.

80.   Seger, C.A., and Miller, E.K. (2010). Category Learning in the Brain. Annu. Rev. Neurosci. 33, 203–
219. https://doi.org/10.1146/annurev.neuro.051508.135546.

81.   Bragdon, L.B., Gibb, B.E., and Coles, M.E. (2018). Does neuropsychological performance in OCD
relate to different symptoms? A meta-analysis comparing the symmetry and obsessing dimensions.
Depress. Anxiety 35, 761–774. https://doi.org/10.1002/da.22785.

82. Hann, F., Pesthy, O., Brezóczki, B., Vékony, T., Nagy, C.A., Sapey-Triomphe, L.-A., Tóth-Fáber, E., Farkas, B.C., Farkas, K., and Nemeth, D. (2024). Autistic Traits Relate to Speed/Accuracy Trade-off but not Statistical Learning and Updating. Preprint, https://doi.org/10.31219/osf.io/p8m37_v1 https://doi.org/10.31219/osf.io/p8m37_v1.

83. Dominke, C., Graham-Schmidt, K., Gentsch, A., and Schütz-Bosbach, S. (2021). Action inhibition in individuals with high obsessive-compulsive trait of incompleteness: An ERP study. Biol. Psychol. *159*, 108019. https://doi.org/10.1016/j.biopsycho.2021.108019.

84. Scholl, J., Trier, H.A., Rushworth, M.F.S., and Kolling, N. (2022). The effect of apathy and compulsivity on planning and stopping in sequential decision-making. PLoS Biol. *20*, e3001566. https://doi.org/10.1371/journal.pbio.3001566.

85. Hoven, M., Mulder, T., Denys, D., van Holst, R.J., and Luigjes, J. (2024). Lower confidence and increased error sensitivity in OCD patients while learning under volatility. Transl. Psychiatry *14*, 370. https://doi.org/10.1038/s41398-024-03042-3.

86. Nagy, C.A., Hann, F., Brezóczki, B., Farkas, K., Vékony, T., Pesthy, O., and Németh, D. (2025). Intact ultrafast memory consolidation in adults with autism and neurotypicals with autism traits. Brain Res. *1847*, 149299. https://doi.org/10.1016/j.brainres.2024.149299.

87. Vékony, T., Farkas, B.C., Brezóczki, B., Mittner, M., Csifcsák, G., Simor, P., and Németh, D. (2025). Mind wandering enhances statistical learning. iScience *28*, 111703. https://doi.org/10.1016/j.isci.2024.111703.

88. Rodd, J.M. (2024). Moving experimental psychology online: How to obtain high quality data when we can't see our participants. J. Mem. Lang. *134*, 104472. https://doi.org/10.1016/j.jml.2023.104472.

89. Choi, K.W., Kim, Y.-K., and Jeon, H.J. (2020). Comorbid Anxiety and Depression: Clinical and Conceptual Consideration and Transdiagnostic Treatment. In, pp. 219–235. https://doi.org/10.1007/978-981-32-9705-0_14.

90. Goodwin, G.M. (2015). The overlap between anxiety, depression, and obsessive-compulsive disorder. Dialogues Clin. Neurosci. *17*, 249–260. https://doi.org/10.31887/DCNS.2015.17.3/ggoodwin.

91. Sarna, N., Dar, R., and Mazor, M. (2025). Biased and Inattentive Responding Drive Apparent Metacognitive Biases in Mental Health. Preprint at PsyArXiv. https://doi. org/10.31234/osf. io/fzwu8_v1.

92. Ord, J.S., Greve, K.W., Bianchini, K.J., and Aguerrevere, L.E. (2010). Executive dysfunction in traumatic brain injury: The effects of injury severity and effort on the Wisconsin Card Sorting Test. J. Clin. Exp. Neuropsychol. *32*, 132–140. https://doi.org/10.1080/13803390902858874.

93. Miles, S., Howlett, C.A., Berryman, C., Nedeljkovic, M., Moseley, G.L., and Phillipou, A. (2021). Considerations for using the Wisconsin Card Sorting Test to assess cognitive flexibility. Behav. Res. Methods *53*, 2083–2091.

94. Hann, F., Pesthy, O., Brezóczki, B., Vékony, T., Nagy, C.A., Sapey-Triomphe, L.-A., Tóth-Fáber, E., Farkas, B.C., Farkas, K., and Németh, D. (2025). Autistic traits relate to speed/accuracy trade-off but not statistical learning and updating. Sci. Rep. *15*, 32001. https://doi.org/10.1038/s41598-025-16138-7.

95. Vékony, T. (2021). Alternating Serial Reaction Time Task created with jsPsych (Version 1.0.0). Preprint.

96. Vékony, T. (2022). Card Sorting Task created with jsPsych (Version 1.0.1). Preprint.

97. de Leeuw, J.R. (2015). jsPsych: A JavaScript library for creating behavioral experiments in a Web browser. Behav. Res. Methods *47*, 1–12. https://doi.org/10.3758/s13428-014-0458-y.

98. Abramovitch, A., Abramowitz, J.S., and McKay, D. (2021). The OCI-12: A syndromally valid modification of the obsessive-compulsive inventory-revised. Psychiatry Res. *298*, 113808. https://doi.org/10.1016/j.psychres.2021.113808.

99. Fulop, F., Demeter, G., Honbolygó, F., and Csigo, K. (2024). Assessing obsessive-compulsive symptoms in a subclinical and clinical sample: the development of the Hungarian version of the OCI-R. Hungar Assoc Psychopharmacol *26*, 144–152.

100. Anwyl-Irvine, A.L., Massonnié, J., Flitton, A., Kirkham, N., and Evershed, J.K. (2020). Gorilla in our midst: An online behavioral experiment builder. Behav. Res. Methods *52*, 388–407. https://doi.org/10.3758/s13428-019-01237-x.

1050   101.   Vékony, T., Ambrus, G.G., Janacsek, K., and Nemeth, D. (2022). Cautious or causal? Key implicit
1051   sequence learning paradigms should not be overlooked when assessing the role of DLPFC (Commentary
1052   on Prutean et al.). Cortex *148*, 222–226. https://doi.org/10.1016/j.cortex.2021.10.001.

1053   102.   R Core Team (2024). R: A language and environment for statistical computing (Version 4.4.2)
1054   [Computer software]. R Foundation for Statistical Computing. https://www.R-project.org/. Preprint.

1055   103.   Soetens, E., Melis, A., and Notebaert, W. (2004). Sequence learning and sequential effects.
1056   Psychological Research Psychologische Forschung *69*, 124–137. https://doi.org/10.1007/s00426-003-
1057   0163-4.

1058   104.   Singmann, H., Bolker, B., Højsgaard, S., Fox, J., Lawrence, M.A., and Mertens, U. (2024). afex:
1059   Analysis of Factorial Experiments [Computer software]. Preprint.

1060   105.   Lenth, R. emmeans: Estimated Marginal Means, aka Least-Squares Means [Computer software].
1061   https://CRAN.R-project.org/package=emmeans. Preprint.

1062