Inverse expression of Ten3 and Lphn2 across the developing mouse brain reveals a global strategy for circuit assembly

URee Chon[1,2,3], Daniel T. Pederick[1,3,4]*, Jun Ho Song[1], Yanbo Zhang[1], Isabella Rana[1,5], Liqun Luo[1]*

[1]Department of Biology, Howard Hughes Medical Institute, Stanford University, Stanford, CA, USA
[2]Neurosciences PhD Program, Stanford University, Stanford, CA, USA
[3]These authors contributed equally to this study
[4]Present address: Department of Neuroscience, Johns Hopkins University School of Medicine, Baltimore, MD, USA
[5]Present address: Neuroscience PhD Program, Columbia University, New York, NY, USA
*Corresponding authors: dpederi1@jh.edu, lluo@stanford.edu

SUMMARY

Precise wiring of neural circuits requires molecular strategies that ensure accurate target selection across diverse brain regions. Here, we identify inverse expression between a ligand–receptor pair, Teneurin-3 (Ten3) and Latrophilin-2 (Lphn2), throughout the developing mouse brain. Ten3 and Lphn2 exhibit inverse expression gradients along a retinotopic axis orthogonal to the ephrin-A and EphA gradients; along the tonotopic axis across multiple brainstem auditory nuclei; and along the dorsomedial–ventrolateral axis in striatum and pallidum. Their inverse expression also creates discrete domains of cerebellar Purkinje cells and cerebellar nuclei. Using conditional tag mice, we show that inverse Ten3 and Lphn2 expression patterns predict connectivity, following a 'Ten3→Ten3, Lphn2→Lphn2' rule in all above circuits, and that Lphn2 is required in executing this rule in Purkinje cells→cerebellar nuclei projection. Our findings reveal a global strategy of coordinating gene expression of key wiring molecules with circuit connectivity across the developing brain.

INTRODUCTION

Precise wiring of neural circuits during development is essential for proper information processing and animal behavior. The specificity of neuronal connectivity emerges from cell–cell interactions that rely on the differential expression of cell surface proteins (CSPs) to distinguish neuronal identity and guide the correct selection of synaptic partners.[1–5] However, while the human brain contains ~10^{11} neurons that form highly selective connections, the genome encodes only ~20,000 protein-coding genes, less than ~3,000 of which encode CSPs.[6,7] This disparity raises a fundamental question: how does the brain achieve such extraordinary wiring specificity with a limited molecular repertoire? Several mechanisms have been proposed to reconcile this disparity.[4] Graded expression enables different levels of the same CSP to specify different connections. Neurons may leverage combinatorial expression of multiple CSPs to generate unique molecular signatures. Alternative splicing and post-translational modifications can produce diverse isoforms from a single gene, expanding the functional repertoire of CSPs. Despite extensive research on individual or pairs of CSPs that act as guidance molecules for wiring of specific circuits, however, the spatial dynamics of any ligand–receptor pair at a whole brain scale, and how they cooperate to establish precise wiring of multiple circuits across development, have not yet been systematically examined.

Teneurins are evolutionarily conserved type II transmembrane protein that instructs synaptic partner matching by homophilic adhesion.[8–11] Additionally, we previously showed that Teneurin-3 (Ten3) and the adhesion G protein-coupled-receptor Latrophilin-2 (Lphn2) display inverse expression in two parallel hippocampal networks.[12] Ten3+ proximal CA1 axons connect with Ten3+ target neurons in distal subiculum, and Lphn2+ distal CA1 axons connect with Lphn2+ target neurons in proximal subiculum. Ten3 and Lphn2 cooperate for target selection in the CA1→subiculum projection through Ten3–Ten3 homophilic attraction and Ten3–

Lphn2 reciprocal heterophilic repulsion.[12,13] In a companion manuscript, we further demonstrate that the Ten3–Lphn2 molecular module is repeatedly used to instruct target selection across multiple nodes of the extended hippocampal network.[14] These studies raise the question of whether this ligand–receptor pair could be broadly deployed in establishing wiring specificity in other regions of the mammalian brain.

Here we report, using whole-brain expression at multiple developmental stages, that inverse expression of Ten3 and Lphn2 is present in many regions across the developing mouse brain. We focus our detailed analyses on four systems. These include the visual and auditory systems, in which nearby neurons in the input field project axons topographically to nearby neurons in the target field to form continuous retinotopic and tonotopic maps[15,16], as well as the basal ganglia and cerebellum where the connectivity organization is less well defined. Across these regions, we observe that inverse Ten3 and Lphn2 expression patterns predict wiring specificity, following a 'Ten3→Ten3, Lphn2→Lphn2' rule relating CSP expression to connectivity. In the visual system, Ten3–Lphn2 inverse expression defines retinotopy along the dorsal–ventral axis orthogonal to the anterior–posterior axis defined by inverse ephrin-A–EphA expression.[15,17] In the auditory system, Ten3–Lphn2 inverse expression aligns with tonotopy across multiple brainstem auditory nuclei, providing a logic for how information about sound frequency could be relayed faithfully across multiple synapses. In the basal ganglia, Ten3–Lphn2 inverse expression coincides with striatal-pallidal and cortical-striatal projections. In the cerebellum, Ten3–Lphn2 inverse expression matches Purkinje cell→cerebellar nuclei projection specificity, and we further demonstrate that Lphn2 is required for implementing the 'Ten3→Ten3' connectivity rule. These findings, along with our companion manuscripts[14,18], establish Ten3 and Lphn2 as a molecular module that organizes specific and precise connectivity across anatomically and functionally distinct brain and spinal cord regions, uncovering a global principle for circuit organization throughout the developing nervous system.

RESULTS AND DISCUSSION

Ten3 and Lphn2 exhibit inverse expression across many regions of the developing brain

Given the inverse Ten3 and Lphn2 expression in the extended hippocampus network,[12,14] we first asked whether similar patterns extend to other regions of the brain during development. We utilized knock-in alleles of Ten3-HA (**Figure S1A**) and Lphn2-mVenus[19] to characterize their distribution across the entire developing brain at embryonic day (E) 15 and 17, and postnatal day (P) 0, 2, 4, and 8. Immunostained brains were cleared and whole-mount imaged with light-sheet microscopy (**Figure 1A**).

Across development, we observed Ten3 and Lphn2 expression in broad, largely non-overlapping and often adjacent regions spanning the forebrain, midbrain, and hindbrain (**Figure 1B**; **Videos S1–S7**). For example, at E17, we observed differential expression in discrete brain regions, including the neocortex, striatum, hypothalamus, superior colliculus, cerebellar cortex, cochlear nucleus, spinal nucleus of the trigeminal, and nucleus of the solitary tract through coronal (**Figure 1C**), horizontal (**Figure 1D**), and sagittal (**Figure 1E**) sections. As examples, Ten3 was enriched medially and Lphn2 laterally in the superior colliculus (**Figure 1C₂, D₅, E₁₀, E₁₁**, white arrows). Conversely, Lphn2 was enriched medially and Ten3 laterally in multiple nuclei of the superior olivary complex (**Figure 1C₃**, magenta arrow), as well as the mammillary nucleus of the hypothalamus (**Figure 1D₈**, orange arrow). In the cochlear nucleus (**Figure 1C₃**, green arrow) and spinal trigeminal nucleus (**Figure 1C₄**, yellow arrow), Lphn2 expression was higher in dorsal and Ten3 higher in ventral regions. Ten3 and Lphn2 also exhibited inverse expression in discrete nuclei of the thalamus (**Figure 1D₇**, red arrow). In the neocortex, Ten3 was high posteriorly whereas Lphn2 was high anteriorly (**Figure 1E₉**, purple arrow); at a coronal section midway along the anterior–posterior axis, Lphn2 was high at lateral and medial edges whereas Ten3 was high in the intermediate zone (**Figure 1C₁**, purple arrow). To facilitate the identification of anatomical regions that express Ten3 and Lphn2 throughout the entire brain and across different timepoints, we developed an interactive, ImageJ-based visualization tool that enables users to actively navigate and examine anatomical regions of interest (**Supplemental Data 1**).

Together, these data show that Ten3 and Lphn2 proteins exhibit spatially segregated and largely inverse expression patterns across the developing brain. These inverse expression patterns do not generally follow fixed body axes but rather are specialized in each brain region. We focused further investigation on four systems that

exhibited robust and stereotyped Ten3–Lphn2 inverse expression: the visual system, auditory system, basal ganglia, and cerebellum, in the context of development and connectivity patterns of each system. Quantitative protein level analyses across multiple developmental stages examined revealed dynamic expression of Ten3 and Lphn2 across developmental stages, and that in systems where timing of target selection of axons has been determined, this timing overlaps with peak expression of Ten3 and Lphn2 (**Figure S2**).

Ten3 and Lphn2 expression aligns with retinotopy along the dorsal–ventral axis of the retina

Retinotopy is a fundamental organization of the visual system—retinal ganglion cells (RGCs) in nearby positions of the retina project axons to nearby neurons in the target regions including the superior colliculus (SC), such that the two-dimensional spatial map of the world in the retina is recapitulated in brain targets (**Figure 2A**). The orderly projection of RGC axons to the SC (equivalent of tectum in non-mammalian vertebrates) have been well described: RGCs from the anterior (nasal) retina project to posterior SC/tectum, posterior (temporal) retina to anterior SC/tectum, ventral retina to medial SC/tectum, and dorsal retina to lateral SC/tectum.[1,17] An extensive body of work has demonstrated that graded expression of cell-surface proteins ephrin-As and their receptors EphAs in both the retina and SC instructs retinotopic map along the anterior–posterior axis of the retina.[20–24] Much less is known about the mechanisms that control RGC axon targeting along the orthogonal dorsal–ventral axis, even though several candidates have been proposed, including ephrin-B and EphB, as well as the Wnt signaling pathway. Ephrin-Bs and EphBs appear to play different roles in frogs and mice, and exhibit shallow expression gradients and produce mild phenotypes with *EphB* disruption;[25,26] Wnt signaling lacks loss-of-function evidence to support a direct instructive role.[27] Interestingly, *Ten3* mRNA has been reported to be expressed in the ventral retina and was shown to have defects in retinocollicular circuit wiring when deleted.[28] These findings suggested Ten3 as a candidate guidance molecule that may contribute to dorsal–ventral retinotopic organization.

RGC projections and target selection at the SC begin during embryogenesis (**Figure S2A**).[29] Using immunostaining in a coronal section at E15, we found that Ten3 was highly expressed in the ventral retina whereas Lphn2 was highly expressed in the dorsal retina (**Figure 2B**). Quantification of normalized fluorescence intensity from flat-mount retina further revealed that Ten3 and Lphn2 proteins formed inverse gradients along the dorsal–ventral axis of the retina (**Figure 2C, D**). Three-dimensional reconstructions of cleared E15 whole brains visualized by light-sheet microscopy revealed that Ten3 and Lphn2 labeled discrete axon bundles in the optic tract, which comprises RGC axons connecting the eyes to brain targets (**Figure 2E**). Inverse expression was also evident in the SC, with preferential expression of Ten3 in medial SC, inversely to Lphn2 enrichment in lateral SC (**Figure 2F**; **Figure S2A**), forming opposing gradients (**Figure 2G**). Double *in situ* hybridization for *Ten3* and *Lphn2* mRNA confirmed local production of each CSP's mRNA in the SC (**Figure S3**), suggesting that the presence of Ten3 and Lphn2 protein in SC is likely contributed by both RGC axons and SC target neurons. Given the known topographic projections of ventral and dorsal RGCs target medial and lateral SC,[1] our findings indicate that RGC axon projections follow the 'Ten3→Ten3, Lphn2→Lphn2' rule, and suggest that these proteins could participate in establishing the retinocollicular maps, perhaps in collaboration ephrin-B/EphB and Wnt,[25–27] orthogonal to the ephrin-A/EphA-regulated axis.

Ten3 and Lphn2 expression aligns with tonotopy in multiple brainstem nuclei along the ascending auditory pathway

Tonotopy is a fundamental organizing principle of the auditory system.[30] Sounds of different frequencies are represented by hair cells at different locations in the cochlea. Through ordered axonal projections, this tonotopic organization is recapitulated along the central auditory pathway in brainstem nuclei (**Figure 3A**) and beyond. These tonotopic maps are fundamental to frequency discrimination, pitch perception, and language processing.[31,32] Little is known about the molecular mechanisms that instruct the formation of tonotopic connections.[16]

Immunostaining revealed that during embryonic and early postnatal periods, Ten3 and Lphn2 proteins were distributed in opposing gradients in all cochlear nuclei including the ventral cochlear nucleus (VCN; **Figure 3B, C**; **Figure S3B**), as well as medial nucleus of the trapezoid body (MNTB), superior paraolivary nucleus (SPN), and lateral superior olive (LSO) in the superior olivary complex (**Figure 3D, E**; **Figure S3C**). Strikingly, Ten3 expression always aligned with low-frequency regions, whereas Lphn2 with high-frequency regions (compare

Figure 3B–E with **Figure 3A**). Double *in situ* hybridization for *Ten3* and *Lphn2* mRNA confirmed local production of Ten3 and Lphn2 in all nuclei of the auditory brainstem (**Figure S4**), suggesting that the protein expression is likely contributed by both input axons and target neurons. Given the known topographic projections between low- and high-frequency regions, these expression patterns indicate that the projections of auditory neurons between different nuclei follow the 'Ten3→Ten3, Lphn2→Lphn2' connectivity rule, and suggest that these proteins could participate in establishing tonotopic maps prior to the onset of hearing.

Analyzing Ten3 expression separately in axons and targets in the VCN→MNTB projections

The 'Ten3→Ten3, Lphn2→Lphn2' connectivity rule implies that in each brain region that follows this rule, Ten3 or Lphn2 proteins are contributed by both input axons and target neurons. To directly visualize this and to quantify the contributions of proteins produced by axons and targets, we next aimed to dissect the contribution of Ten3 in axons and targets separately, using the VCN→MNTB projections as an example. To differentially label the same protein in axons and targets, we generated conditional tag (*cTAG*) mice for Ten3 (**Figure S1A**). In *Ten3^cTAG* mice, the endogenous Ten3 is tagged with an HA epitope tag, which is converted to a Halo-tag in the presence of Flp recombinase. Expression analysis in the hippocampus before and after introducing a germline-FlpO validated the fidelity of protein expression as previously reported[10,12] (**Figure S1A**).

Projection neurons from the VCN to the MNTB are excitatory neurons that express *Slc17a6* encoding the vesicular glutamate transporter VGLUT2, whereas neurons in the MNTB are exclusively inhibitory.[33,34] To selectively label Ten3+ excitatory projection neurons, we crossed *Ten3^cTAG* mice with *Slc17a6-Flp*, which expresses Flp in all VGLUT2+ glutamatergic neurons, including those in the VCN (**Figure S5B**). This strategy enabled us to independently examine Ten3^Halo from VCN axons and Ten3^HA produced by local MNTB neurons at the MNTB (**Figure 3F**).

In control *Ten3^cTAG* mice lacking Flp expression, Ten3^HA (but no Ten3^Halo) was detected in the VCN, consistent with the absence of recombination. In *Slc17a6-Flp;Ten3^cTAG* mice, we observed robust recombination throughout VCN, with Ten3^Halo completely replacing Ten3^HA by P0 (**Figure 3G**). At the target MNTB, *Ten3^cTAG* mice displayed Ten3^HA signal but no Ten3^Halo signal, consistent with lack of recombinase activity in axons and targets. By contrast, *Slc17a6-Flp;Ten3^cTAG* mice exhibited Ten3^Halo signal that overlapped with Ten3^HA expression in the lateral region of the MNTB (**Figure 3H**). Thus, axons of Ten3^Halo+ VCN excitatory neurons preferentially target to regions enriched for Ten3^HA+ inhibitory neurons in the MNTB.

Our data reveal that Ten3 is predominantly expressed in excitatory neurons in the VCN, as in *Slc17a6-Flp;Ten3^cTAG* mice, we did not detect any residual Ten3^HA signals (**Figure 3G**, right panels). Moreover, given the complete conversion of Ten3^HA to Ten3^Halo in VCN, Ten3^HA signals in the MNTB are not contributed by VCN axons and derive exclusively from local MNTB neurons. This allowed us to quantify the relative contribution of Ten3 proteins in axons and targets. Comparing Ten3^HA levels in the MNTB in *Ten3^cTAG* and *Slc17a6-Flp;Ten3^cTAG* mice indicates that the majority of Ten3 protein in the MNTB was contributed by target neurons (**Figure 3I**). Finally, Ten3^Halo+ VCN axons terminated at Ten3^HA region in the lateral MNTB, providing direct evidence supporting the 'Ten3→Ten3' targeting rule. Interestingly, Ten3^Halo+ VCN axons passed by Ten3− (and Lphn2+) regions in the medial MNTB to reach their destination (**Figure 3H**, middle row). This observation highlights the selectivity of VCN axon targeting and suggests that repulsion from Lphn2 may contribute to this selectivity, as previously described in the hippocampal network.[12,14]

Ten3 and Lphn2 display inverse expression in the basal ganglia

We next focused on the basal ganglia, a key regulator of motor output, action selection, and decision making.[35,36] GABAergic spiny projection neurons in the input nucleus of the basal ganglia, the caudate putamen (CP, also known as striatum) form a direct pathway projecting to the globus pallidus internal segment (GPi) and substantia nigra, and an indirect pathway that first projects to the globus pallidus external segment (GPe). We found that Ten3 and Lphn2 exhibited inverse expression in both CP and GPe (**Figure S2D**; **Video S1**), and thus focused our further analysis on the indirect (CP→GPe) pathway (**Figure 4A**).

Immunostaining of Ten3 and Lphn2 revealed preferential expression of Ten3 in dorsomedial CP and GPe globally, inversely to Lphn2 enrichment in ventrolateral CP and GPe (**Figure 4B, C**; **Figure S2D**). Quantification

across the dorsomedial–ventrolateral axis of the CP and GPe revealed inverse gradients of expression (**Figure 4D, F**). This inverse pattern was confirmed by Spearman's correlation analysis across animals, which revealed significant negative correlations between Ten3 and Lphn2 expression in both the CP (**Figure 4E**) and GPe (**Figure 4G**). Three-dimensional reconstructions of cleared whole brains visualized by light-sheet microscopy reinforced the consistent spatial segregation of Ten3 and Lphn2 in the CP and GPe (**Video S8**). Double *in situ* hybridization for *Ten3* and *Lphn2* mRNA showed that both *Ten3* and *Lphn2* mRNAs were expressed in the CP and GPe (**Figure S6A, B**). These experiments validated that at least a part of the proteins we visualized in both the CP and GPe were produced locally rather than exclusively contributed by incoming axons. Furthermore, within CP, both Ten3 and Lphn2 mRNAs were both co-expressed with *Tac1* or *Penk*, markers for spiny projection neurons in the direct and indirect pathways, respectively (**Figure S6C, D**).

Unlike the continuous gradients at embryonic stages (**Figure S2D**), Ten3[+] regions in the dorsomedial CP formed spatially clustered hotspots by P4, intermingled with cortical axon bundles (**Figure 4B**). Lphn2[+] regions maintained a broader distribution of expression throughout the ventrolateral CP. Near the central region of the CP where the global gradients of Ten3 and Lphn2 met, Lphn2[+] regions also formed hotspots interdigitating with the Ten3[+] hotspots (**Figure 4B**; **Figure S2D**). To test whether these hotspots are related to previously described striosome (patch) and matrix compartments,[37] we performed co-expression of Ten3 and Lphn2 with μ-opioid receptor (MOR) or calbindin, markers of striosomes and matrix, respectively.[37,38] We did not find clear correspondence between their expression patterns (**Figure S6E, F**), suggesting that heterogeneity created by Ten3 and Lphn2 hotspots is unrelated to striosomes and matrix.

Striatopallidal projections follow the Ten3→Ten3 rule

Given that the CP and GPe are both globally patterned by differential expression of Ten3 and Lphn2, we next asked whether the CP→GPe projections follow the 'Ten3→Ten3' rule. To examine this, we injected lentivirus expressing *FlpO-GFP* into the CP of *Ten3[cTAG/cTAG];Lphn2[mV/mV]* mice at P0 and examined projection patterns of Ten3[Halo+] CP axons in the GPe at P4 (**Figure 4H**). The robust Ten3[Halo] signal confirmed local production of Ten3 in the CP. We found that Ten3[Halo+] axons from the CP preferentially innervate the Ten3[HA] regions of the GPe, while avoiding Lphn2[mV] regions (**Figure 4I, J**). Quantification of axon-target overlap confirmed substantial overlap of Ten3[Halo+] CP axon signals within Ten3[HA] target regions but little overlap in the Lphn2[mV] target region (**Figure 4K**). Thus, the CP→GPe projections follow the 'Ten3→Ten3' rule.

Corticostriatal projections follow the Ten3→Ten3 rule

The CP receives excitatory input from across the neocortex, as well as specific thalamic regions, mainly the parafascicular nuclei.[39] We next examined whether these input axons contribute to Ten3 expression in the CP, and whether input projections also follow the 'Ten3→Ten3' rule. To do this we utilized our conditional tag allele in combination with cell-type-specific Flp drivers to differentially label Ten3 produced from input axons versus the target arbors.

Before describing the contribution of thalamic and cortical inputs of Ten3 to the CP, we performed two control experiments. First, in the absence of Flp recombinase expression, we did not detect Ten3[Halo] signal in the CP of *Ten3[cTAG]* mice (**Figure 5A**). Second, to visualize local Ten3 production in CP neurons, which are predominantly GABAergic, we crossed *Ten3[cTAG]* mice with *Slc32a1-Flp*, which expresses Flp in all GABAergic neurons (**Figure S5A**). We observed robust Ten3[Halo] signals in the CP (**Figure 5B₃**), confirming local production of Ten3 in the CP, as also indicated by mRNA *in situ* hybridization analysis (**Figure S6A**). Interestingly, some of the Ten3[Halo] signals overlapped with Ten3[HA] signals of the CP (**Figure 5B₄**). The persistence of Ten3[HA] could be contributed by (1) expression of Ten3 in non-GABAergic CP neurons, such as cholinergic interneurons[40]; (2) incomplete recombination in GABAergic CP neurons; and/or (3) input axons from cortex and/or thalamus.

Next, we examined whether thalamic excitatory neurons contribute Ten3 to the CP. To test this, we crossed *Ten3[cTAG]* mice with *Slc17a6-Flp*, which expresses Flp in most/all thalamic excitatory neurons[41,42] (**Figure S5B**). We did not detect Ten3[Halo] axon labeling in the CP (**Figure 5C**), suggesting that thalamic axons that innervate the CP do not express Ten3 at their axon terminals at this developmental stage (P4). This contrasts with previous studies suggesting that Ten3 is expressed in both the CP and parafascicular nuclei, where it was proposed to

regulate thalamo-striatal connectivity.[43,44] This discrepancy could reflect differential detecting thresholds for mRNA and protein or post-transcriptional regulation. Although prior studies reported *Ten3* mRNAs in the parafascicular nuclei, we did not detect high levels of Ten3 protein in this region throughout development (**Figure S6G**). While *Ten3* mRNA persist in many brain regions into adulthood[45], Ten3 protein levels in many brain regions sharply decline after postnatal day 4 (**Figure S2**), suggesting post-transcriptional regulation.

Finally, to test whether Ten3$^+$ cortical excitatory neurons project axons to the CP, we crossed *Ten3^{cTAG}* mice with *Slc17a7-Flp*, where Flp is expressed primarily in cortical excitatory neurons but not in any of the CP neurons nor thalamic neurons (**Figure S5C**). Ten3^{Halo} signal was robustly detected in the CP and coincided with Ten3^{HA}-expressing regions, including Ten3^{HA} hotspots at the intermediate zone between Ten3$^+$ and Lphn2$^+$ regions (**Figure 5D**). These data indicate that at least some Ten3$^+$ cortical neurons project to the CP, and such projections follow the 'Ten3→Ten3' rule. Interestingly, retrosplenial area and posterior parietal association areas of the cortex had high Ten3 expression (**Supplemental Data 1**) and are known to project to the dorsomedial CP.[46] These findings suggest that Ten3 expression may contribute to topographic cortical inputs that target distinct CP subregions.

Together, the inverse expression patterns of Ten3 and Lphn2 at CP, both as global gradients and segregated hotspots at the intermediate zone, could be used to organize cortical input and pallidal output of CP neurons following the 'Ten3→Ten3' connectivity rule.

Inverse expression patterns of Ten3 and Lphn2 in both Purkinje cells and cerebellar nuclei reveal new molecular organization of the cerebellum

The cerebellum has a highly organized and stereotyped architecture and plays key roles in motor coordination, sensorimotor integration, and cognition.[47–49] All information processing in the cerebellar cortex channels through a single layer of Purkinje cells (PCs) projecting their axons to the cerebellar nuclei (CN), which then broadcast to the rest of the brain. The PC→CN projections are established early in development and exhibit precise spatial organization,[47] yet the molecular mechanisms that guide PC axons to their appropriate nuclear targets remain poorly defined.

In coronal sections of the developing cerebellum, we found that Ten3 and Lphn2 proteins were inversely expressed in PCs in alternating segments across the medial–lateral axis (**Figure 6A**; **Figure S2E**), with sharp transitions (**Figure 6B**). Quantification of normalized fluorescence intensities along the PC layer (**Figure 6A**) revealed the reciprocity of Ten3 and Lphn2 across hemispheres in an individual mouse (**Figure 6C**). Spearman's correlation analysis confirmed an inverse relationship across different mice (**Figure 6D**). Inverse expression of Ten3 and Lphn2 was also evident in the CN (**Figure 6E**), with preferential expression of Ten3 in dorsomedial and ventrolateral CN, inversely to Lphn2 enrichment in intermediate CN in this coronal section (**Figure 6F**). This pattern was confirmed by Spearman's correlation analysis across multiple mice (**Figure 6G**). To further assess the stereotypy of Ten3 and Lphn2 expression, we quantified spatial similarity on whole-mount-stained cerebella. This analysis revealed minimal intra-hemispheric overlap between Ten3 and Lphn2, and high inter-hemispheric and inter-animal similarity for Ten3 or for Lphn2 (**Figure S7D**), to a similar degree as other regions with well-established, highly stereotyped expression patterns (**Figure S7E**). Double *in situ* hybridization for *Ten3* and *Lphn2* mRNA revealed similar inverse expression pattern throughout development in both PC layer and CN (**Figure S7A–C**), validating that at least some of the protein expression in both PC and CN is contributed by local neurons.

Previous studies of the cerebellar cortex have extensively characterized PC organization along the medial–lateral axis based on expression patterns of specific genes,[50] revealing alternating arrays of parasagittal zebrin II-positive (zebrin$^+$) and zebrin II-negative (zebrin$^-$) stripes that run vertically across the entire cerebellar cortex.[50–52] Given our finding of Ten3$^+$ and Lphn2$^+$ 'bands' distributed along the medial–lateral axis, we next asked whether these expression patterns correspond to the zebrin stripes. Co-staining of Ten3 and Lphn2 with either zebrin$^+$ or zebrin$^-$ molecular markers, aldolase C or PlcB4, respectively,[53,54] we did not find correspondence in their expression patterns (**Figure S7F, G**), indicating that Ten3$^+$ and Lphn2$^+$ 'bands' represent a distinct organization. Indeed, visualizing Ten3$^+$ and Lphn2$^+$ PCs in 3D revealed that many parasagittal bands of Ten3 and Lphn2 apparent from sections (e.g., **Figure 6A**) were in fact connected in 3D, forming continuous regions (**Figure S7H, Video S9**). The Ten3$^+$ and Lphn2$^+$ regions in 3D could nevertheless be segmented into five discrete domains per

hemisphere (**Figure 6H**). Notably, 3D analysis of expression in the CN also revealed five discrete domains spanning 3D organization of the CN (**Figure 6I$_1$, Video S10**). Furthermore, the five CN domains defined by Ten3 and Lphn2 expression do not align strictly with the three histologically separable cerebellar nuclei—namely, the fastigial, interposed, and dentate nucleus (**Figure 6I$_2$; Figure S7A**)—suggesting that Ten3 and Lphn2 expression may define a molecular organization distinct from the traditional cytoarchitectural boundaries.

The observed Ten3 and Lphn2 pattern in the PC layer stands in stark contrast to the column-like organization typically associated with parasagittal stripes across all axes (**Video S11**), as most well-known cerebellar molecular markers conform to a stripe-like pattern. The five broad, continuous domains in both PCs and CN defined by inverse Ten3 and Lphn2 expression represent novel molecular heterogeneity in the cerebellum, which may reflect previously unknown functional organization within the cerebellar circuits.

Purkinje cell→cerebellar nuclei projections follow the 'Ten3→Ten3, Lphn2→Lphn2' rule

Given the inverse expression of Ten3 and Lphn2 in both PCs and CN, we next tested whether the PC→CN projection follows the 'Ten3→Ten3, Lphn2→Lphn2' rule. We first tested whether Ten3$^+$ PC axons overlapped with Ten3$^+$ target region of CN. We injected lentivirus expressing *FlpO-GFP* into the cerebellar cortex of *Ten3$^{cTAG/cTAG}$;Lphn2$^{mV/mV}$* mice at P0 and examined Ten3^{Halo+} PC axons in the CN (**Figure 6J**). We found that Ten3^{Halo+} PC axons preferentially projected to the Ten3^{HA+} regions of the CN and avoided Lphn2^{mV+} regions of the CN (**Figure 6K–P**). Quantitative analysis confirmed significantly greater overlap between Ten3^{Halo+} axon signals with Ten3^{HA+} target regions compared to Lphn2^{mV+} target regions (**Figure 6Q**).

To test whether Lphn2$^+$ Purkinje cells project to Lphn2$^+$ regions of the CN, we utilized new Lphn2 conditional tag mice we generated (**Figure S1B**) and injected the *FlpO-GFP* lentivirus into the cerebellar cortex of *Ten3$^{cTAG/cTAG}$;Lphn2$^{cTAG/cTAG}$* mice at P0. This would switch Lphn2 expression from Lphn2^{FLAG} to Lphn2^{SNAP} in PC axons, while leaving Ten3^{HA} unrecombined in CN target region (**Figure 6R**). Indeed, Lphn2^{SNAP+} PC axons projected preferentially to Ten3$^-$ regions of the CN, avoiding Ten3^{HA+} territories (**Figure 6S–X**). Quantitative analysis confirmed this preference (**Figure 6Y**).

In summary, Ten3 and Lphn2 show an inverse expression pattern in both the PCs and CN, and PC→CN projections follow the 'Ten3→Ten3, Lphn2→Lphn2' rule. Together with our findings in the visual, auditory, and basal ganglia systems (**Figures 2–5**), as well as the extended hippocampal network,[12,14] these findings reveal a generalizable mechanism by which spatially segregated Ten3 and Lphn2 could guide the formation of precise long-range connections in parallel networks using the 'Ten3→Ten3, Lphn2→Lphn2' connectivity rule.

Lphn2 is required for targeting precision of PC→CN axons

Given our demonstration that Ten3-mediated homophilic attraction and Lphn2–Ten3–mediated heterophilic repulsion instruct target selection of the extended hippocampal circuit to follow the 'Ten3→Ten3, Lphn2→Lphn2' connectivity rule,[12,14] we asked whether Lphn2 plays a functional role in executing the 'Ten3→Ten3' projection rule observed in the cerebellum. Due to the complex nature of Ten3 connectivity in the cerebellum, traditional tracing approaches were not feasible, as Flp-driver lines that selectively target PCs during the embryonic window when Ten3 and Lphn2 expression begins were unavailable. Therefore, we used our conditional tag mice to examine if Ten3$^+$ PC axons would mistarget to Ten3$^-$ regions of CN when *Lphn2* was deleted.

We injected *FlpO-GFP* lentivirus into PCs of *En1-Cre;Lphn2$^{+/+}$;Ten3^{cTAG}* mice (control, **Figure 7A**) or *En1-Cre;Lphn2$^{fl/fl}$;Ten3^{cTAG}* mice (experimental, **Figure 7B**) at P0. *En1*, encoding Engrailed homeobox 1, is expressed in developing rhombomere I that gives rise to both PCs and CN. Recombination in both regions was confirmed with a Cre-dependent reporter (**Figure S5D**), which would result in *Lphn2* being deleted in both PCs and CN. In line with previous results, Ten3^{Halo+} axons from PCs selectively targeted Ten3^{HA} regions in the anterior, intermediate, and posterior CN in *Lphn2$^{+/+}$* control mice (**Figure 7C–F**). By contrast, Ten3^{Halo+} axons from PCs targeted more broadly into Ten3$^-$ CN regions (non-Ten3^{HA} regions) in *Lphn2^{cKO}* mice, indicative of mistargeting (**Figure 7G–J**, red arrows). This mistargeting was observed at all levels of the anterior–posterior axis, with variations in severity. Some Ten3^{Halo+} PC axons still innervated Ten3^{HA} domains of the CN, which could be contributed by Ten3–Ten3 homophilic attraction not affected by our experimental manipulation. Additionally, the order in which Ten3^{Halo+} axons encounter Ten3- or Lphn2-expressing domains may influence their targeting

as demonstrated in the companion paper[14]—contact with Ten3$^+$ regions of CN first may preferentially stabilize these projections, causing Ten3^{Halo+} axons innervation of Ten3^{HA} domains. The order of PC axons encountering Ten3$^+$ or Lphn2$^+$ CN zones may differ in different CN regions, contributing to heterogeneity of the phenotypes.

Nevertheless, quantification of axonal overlap overall revealed a significant reduction in Ten3^{Halo} alignment with Ten3^{HA} regions ($P < 0.01$) and a significant increase in Ten3^{Halo} overlap in Ten3$^-$ CN regions ($P < 0.01$) in *Lphn2$^{fl/fl}$* mice compared to *Lphn2$^{+/+}$* control mice (**Figure 7K**). These results indicate that *Lphn2* is required for PC→CN projections to follow the 'Ten3→Ten3' rule, likely caused by Lphn2 acting as a repulsive ligand[12,14] to restrict Ten3$^+$ PC axons to appropriate Ten3$^+$ CN targets.

CONCLUDING REMARKS

In this study, we found striking, inverse expression of cell-surface proteins Ten3 and Lphn2 across the visual, auditory, basal ganglia, and cerebellar circuits during development. These inverse expression patterns emerge in three distinct modes: 1) gradients in the retina–superior colliculus of the visual circuit and the cochlear nucleus–superior olivary complex of the auditory circuit (**Figure 2 and 3**); 2) discrete domains in the cerebellum (**Figure 6**) and the extended hippocampus described in the companion paper;[14] and 3) a combination of both in the striatal circuit (**Figure 4**). These distinct spatial expression modes may reflect different underlying organizational logics for specific circuits. In retinotopic and tonotopic maps, nearby neurons in the input field project to nearby neurons in the target field to preserve continuous spatial and frequency maps. Continuous gradients of wiring molecules enable their levels to instruct targeting specificity.[1,20–24] By contrast, expression in discrete domains may serve as independent modules for circuit segregation and parallel information processing, as in the case of the medial and lateral subnetworks in the extended hippocampal circuit.[55] The global gradients of Ten3 and Lphn2 expression in CP may reflect continuous representation of sensorimotor function of the basal ganglia.[35,39] The discrete hotspots in CP and discrete domains in the cerebellum of Ten3 and Lphn2 expression we discovered do not match with previous described striosomes/matrix or parasagittal zebrin+/zebrin– bands, suggesting novel molecular organizations, the functional significance of which remains to be discovered. These findings also suggest that detailed examination of Ten3 and Lphn2 expression in other circuits (**Supplemental Data 1**), similar to what we have done here for the four systems, may reveal previously unknown connection specificity that might lead to new insights into their functional organization.

The temporal dynamics of Ten3 and Lphn2 expression suggest a universal role in target selection. In systems where timing of target selection of axons has been determined, this timing overlaps with peak expression of Ten3 and Lphn2 (**Figure S2**). This exquisite spatiotemporal regulation of CSP expression allows a limited set of molecules to direct wiring of many circuits across diverse brain regions.

Interestingly, the inverse expression of Ten3 and Lphn2 is observed across a diverse array of brain regions, including the forebrain, midbrain, and hindbrain, despite their distinct developmental origins. This raises an intriguing question about whether a conserved regulatory mechanism governs this expression pattern across the brain, or if region-specific cues independently establish similar molecular expression patterns. Regardless of details of the regulatory mechanisms, our findings suggest that to accommodate the increasing complexity of the nervous system during evolution, tweaking expression patterns of key existing wiring molecules could be a strategy alternative to—and perhaps simpler than—reinventing new wiring molecules.

Together with our findings in the companion manuscripts demonstrating functional roles of Ten3 and Lphn2 in instructing target selection in the extended hippocampal circuit[14] and somatotopic map[18], these results define a conserved molecular logic for target recognition that is used repeatedly across the nervous system to establish precise neural circuits. The inverse expression of Ten3 and Lphn2, along with their molecular function as a ligand–receptor pair mediating both attraction and repulsion, reveals a global molecular strategy by which distinct circuits achieve precise circuit assembly throughout development.

Limitations of the study
While our study provides a foundational atlas of Ten3 and Lphn2 expression, several limitations remain. First, although we focused on four representative circuits, many other brain regions also exhibit inverse expression

patterns of Ten3 and Lphn2. These regions may likewise employ the 'Ten3→Ten3, Lphn2→Lphn2' rule to guide circuit assembly, which need to be investigated in future studies. Second, in addition to regulating target selection during development for the specific Ten3 and Lphn2 pair, there are other paralogs of teneurins (4 in total) and latrophilins (3 in total) in the mammalian genome that are implicated in synapse formation, organization and functional modulation.[56,57] Several of these paralogs are also expressed in non-neuronal subpopulations, suggesting roles beyond classical neuron-neuron communication. The broader expression across multiple cell types and overlapping expression patterns suggests that they may play broader roles in addition to spatially defined circuit assembly. Finally, although we described inverse expression of Ten3 and Lphn2 and the related connectivity rule in four systems, we only showed functional requirement of Lphn2 in the PC→CN connectivity. Future loss- and gain-of-function experiments will be required to demonstrate causal roles of Ten3 and Lphn2 in wiring the visual, auditory, and striatal systems. Given that Ten3 and Lphn2 are repeatedly used in the extended hippocampal network[12,14], the somatotopic map[18], and PC→CN connectivity (**Figure 7**), it seems likely that Ten3 and Lphn2 will play an analogous role to instruct target selection in other circuits that follow the 'Ten3→Ten3, Lphn2→Lphn2' rule, perhaps in collaboration with other CSPs.

ACKNOWLEDGEMENTS
We thank members of the Luo Lab, especially Ellen Gingrich, Tom Hindmarsh-Sten, Lijun Qi, Zhuoran Li, Alex Starr, Yunming Wu, and Cheng Lyu, as well as Julia Kaltschmidt, Jennifer Raymond, and Kang Shen for their support, expertise, and insights on this study. We also thank Caiying Guo and the Transgenic Facility at Janelia Research Campus for generating the cTAG mice; Yiming Zhang, Huyan Meng, and Zhigang He at the Boston Children's Hospital Viral Core for the *LV-FlpO-GFP* virus; Artur Kania and Kevin Sangster for feedback on the manuscript; Kenneth Magpusao and Carlota Manalac for animal assistance; David Luginbuhl and Mary Molacavage for administrative assistance. D.T.P. was supported by the American Australian Association Education Fund Scholarship. L.L. is an HHMI investigator. This work was supported by the National Institutes of Health (R01-NS050580 to L.L.), and the Simons Foundation (SFARI Fellows-to-Faculty Award to D.T.P.)

AUTHOR CONTRIBUTIONS
L.L., D.T.P., and U.C. conceived the project and designed all experiments. U.C. and D.T.P. performed majority of the experiments and data analyses. J.H.S. performed whole brain registration and Dice coefficient analyses, and Y.Z. conducted the *in situ* hybridization assays. I.R. assisted with the iDISCO experiments. L.L. and D.T.P. supervised the study. U.C., D.T.P., and L.L. wrote the manuscript.

DECLARATION OF INTERESTS
L.L. is on the advisory board of *Cell*.

REFERENCES

1. Sperry, R.W. (1963). Chemoaffinity in the orderly growth of nerve fiber patterns and connections*. Proceedings of the National Academy of Sciences *50*, 703–710. https://doi.org/10.1073/pnas.50.4.703.
2. Kolodkin, A.L., and Tessier-Lavigne, M. (2011). Mechanisms and Molecules of Neuronal Wiring: A Primer. Cold Spring Harb Perspect Biol *3*, a001727. https://doi.org/10.1101/cshperspect.a001727.
3. Sanes, J.R., and Yamagata, M. (2009). Many Paths to Synaptic Specificity. Annu. Rev. Cell Dev. Biol. *25*, 161–195. https://doi.org/10.1146/annurev.cellbio.24.110707.175402.
4. Luo, L. (2020). Principles of Neurobiology (Garland Science).
5. Dickson, B.J. (2002). Molecular Mechanisms of Axon Guidance. Science *298*, 1959–1964. https://doi.org/10.1126/science.1072165.
6. The human brain - The Human Protein Atlas https://www.proteinatlas.org/humanproteome/brain.
7. Bausch-Fluck, D., Goldmann, U., Müller, S., van Oostrum, M., Müller, M., Schubert, O.T., and Wollscheid, B. (2018). The in silico human surfaceome. Proceedings of the National Academy of Sciences *115*, E10988–E10997. https://doi.org/10.1073/pnas.1808790115.

8. Hong, W., Mosca, T.J., and Luo, L. (2012). Teneurins Instruct Synaptic Partner Matching in an Olfactory Map. Nature *484*, 201–207. https://doi.org/10.1038/nature10926.

9. Mosca, T.J., Hong, W., Dani, V.S., Favaloro, V., and Luo, L. (2012). Trans-synaptic Teneurin signalling in neuromuscular synapse organization and target choice. Nature *484*, 237–241. https://doi.org/10.1038/nature10923.

10. Berns, D.S., DeNardo, L.A., Pederick, D.T., and Luo, L. (2018). Teneurin-3 controls topographic circuit assembly in the hippocampus. Nature *554*, 328–333. https://doi.org/10.1038/nature25463.

11. Klein, R., and Pasterkamp, R.J. (2021). Recent advances in inter-cellular interactions during neural circuit assembly. Current Opinion in Neurobiology *69*, 25–32. https://doi.org/10.1016/j.conb.2020.12.004.

12. Pederick, D.T., Lui, J.H., Gingrich, E.C., Xu, C., Wagner, M.J., Liu, Y., He, Z., Quake, S.R., and Luo, L. (2021). Reciprocal repulsions instruct the precise assembly of parallel hippocampal networks. Science *372*, 1068–1073. https://doi.org/10.1126/science.abg1774.

13. Pederick, D.T., Perry-Hauser, N.A., Meng, H., He, Z., Javitch, J.A., and Luo, L. (2023). Context-dependent requirement of G protein coupling for Latrophilin-2 in target selection of hippocampal axons. eLife *12*, e83529. https://doi.org/10.7554/eLife.83529.

14. Gingrich, E.C., Pederick, D.T., Zhang, Y., and Luo, L. (2025). Ten3–Lphn2-mediated target selection across the extended hippocampal network demonstrates a repeated strategy for circuit assembly. Preprint at bioRxiv, https://doi.org/10.1101/2025.08.13.670207 https://doi.org/10.1101/2025.08.13.670207.

15. Luo, L., and Flanagan, J.G. (2007). Development of Continuous and Discrete Neural Maps. Neuron *56*, 284–300. https://doi.org/10.1016/j.neuron.2007.10.014.

16. Sitko, A.A., and Goodrich, L.V. (2021). Making sense of neural development by comparing wiring strategies for seeing and hearing. Science *371*, eaaz6317. https://doi.org/10.1126/science.aaz6317.

17. McLaughlin, T., and O'Leary, D.D.M. (2005). MOLECULAR GRADIENTS AND DEVELOPMENT OF RETINOTOPIC MAPS. Annual Review of Neuroscience *28*, 327–355. https://doi.org/10.1146/annurev.neuro.28.061604.135714.

18. Sangster, K.T., Zhang, X., Toro, D. del, Sarantopoulos, C., Moses, A.M., Mahasenan, S., Pederick, D.T., Perreault, S., Fallet-Bianco, C., Roome, R.B., et al. (2025). Teneurin-3 and latrophilin-2 are required for somatotopic map formation and somatosensory topognosis. Preprint at bioRxiv, https://doi.org/10.1101/2025.08.13.670179 https://doi.org/10.1101/2025.08.13.670179.

19. Anderson, G.R., Maxeiner, S., Sando, R., Tsetsenis, T., Malenka, R.C., and Südhof, T.C. (2017). Postsynaptic adhesion GPCR latrophilin-2 mediates target recognition in entorhinal-hippocampal synapse assembly. Journal of Cell Biology *216*, 3831–3846. https://doi.org/10.1083/jcb.201703042.

20. Drescher, U., Kremoser, C., Handwerker, C., Löschinger, J., Noda, M., and Bonhoeffer, F. (1995). In vitro guidance of retinal ganglion cell axons by RAGS, a 25 kDa tectal protein related to ligands for Eph receptor tyrosine kinases. Cell *82*, 359–370. https://doi.org/10.1016/0092-8674(95)90425-5.

21. Feldheim, D.A., Kim, Y.-I., Bergemann, A.D., Frisén, J., Barbacid, M., and Flanagan, J.G. (2000). Genetic Analysis of Ephrin-A2 and Ephrin-A5 Shows Their Requirement in Multiple Aspects of Retinocollicular Mapping. Neuron *25*, 563–574. https://doi.org/10.1016/S0896-6273(00)81060-0.

22. Feldheim, D.A., Nakamoto, M., Osterfield, M., Gale, N.W., DeChiara, T.M., Rohatgi, R., Yancopoulos, G.D., and Flanagan, J.G. (2004). Loss-of-Function Analysis of EphA Receptors in Retinotectal Mapping. J. Neurosci. *24*, 2542–2550. https://doi.org/10.1523/JNEUROSCI.0239-03.2004.

23. Rashid, T., Upton, A.L., Blentic, A., Ciossek, T., Knöll, B., Thompson, I.D., and Drescher, U. (2005). Opposing Gradients of Ephrin-As and EphA7 in the Superior Colliculus Are Essential for Topographic Mapping in the Mammalian Visual System. Neuron *47*, 57–69. https://doi.org/10.1016/j.neuron.2005.05.030.

24. Suetterlin, P., and Drescher, U. (2014). Target-Independent EphrinA/EphA-Mediated Axon-Axon Repulsion as a Novel Element in Retinocollicular Mapping. Neuron *84*, 740–752. https://doi.org/10.1016/j.neuron.2014.09.023.

25. Hindges, R., McLaughlin, T., Genoud, N., Henkemeyer, M., and O'Leary, D.D.M. (2002). EphB Forward Signaling Controls Directional Branch Extension and Arborization Required for Dorsal-Ventral Retinotopic Mapping. Neuron *35*, 475–487. https://doi.org/10.1016/S0896-6273(02)00799-7.

26. Mann, F., Ray, S., Harris, W.A., and Holt, C.E. (2002). Topographic Mapping in Dorsoventral Axis of the *Xenopus* Retinotectal System Depends on Signaling through Ephrin-B Ligands. Neuron *35*, 461–473. https://doi.org/10.1016/S0896-6273(02)00786-9.

27. Schmitt, A.M., Shi, J., Wolf, A.M., Lu, C.-C., King, L.A., and Zou, Y. (2006). Wnt–Ryk signalling mediates medial–lateral retinotectal topographic mapping. Nature *439*, 31–37. https://doi.org/10.1038/nature04334.

28. Dharmaratne, N., Glendining, K.A., Young, T.R., Tran, H., Sawatari, A., and Leamey, C.A. (2012). Ten-m3 Is Required for the Development of Topography in the Ipsilateral Retinocollicular Pathway. PLOS ONE *7*, e43083. https://doi.org/10.1371/journal.pone.0043083.

29. Osterhout, J.A., El-Danaf, R.N., Nguyen, P.L., and Huberman, A.D. (2014). Birthdate and Outgrowth Timing Predict Cellular Mechanisms of Axon Target Matching in the Developing Visual Pathway. Cell Reports *8*, 1006–1017. https://doi.org/10.1016/j.celrep.2014.06.063.

30. Kandler, K., Clause, A., and Noh, J. (2009). Tonotopic reorganization of developing auditory brainstem circuits. Nat Neurosci *12*, 711–717. https://doi.org/10.1038/nn.2332.

31. Oxenham, A.J., Bernstein, J.G.W., and Penagos, H. (2004). Correct tonotopic representation is necessary for complex pitch perception. Proc Natl Acad Sci U S A *101*, 1421–1425. https://doi.org/10.1073/pnas.0306958101.

32. Mengler, E.D., Hogben, J.H., Michie, P., and Bishop, D.V.M. (2005). Poor frequency discrimination is related to oral language disorder in children: a psychoacoustic study. Dyslexia *11*, 155–173. https://doi.org/10.1002/dys.302.

33. Ito, T., Bishop, D.C., and Oliver, D.L. (2011). Expression of glutamate and inhibitory amino acid vesicular transporters in the rodent auditory brainstem. Journal of Comparative Neurology *519*, 316–340. https://doi.org/10.1002/cne.22521.

34. Bonito, M.D., Narita, Y., Avallone, B., Sequino, L., Mancuso, M., Andolfi, G., Franzè, A.M., Puelles, L., Rijli, F.M., and Studer, M. (2013). Assembly of the Auditory Circuitry by a Hox Genetic Network in the Mouse Brainstem. PLOS Genetics *9*, e1003249. https://doi.org/10.1371/journal.pgen.1003249.

35. DeLong, M.R., and Georgopoulos, A.P. (1981). Motor Functions of the Basal Ganglia. In Comprehensive Physiology (John Wiley & Sons, Ltd), pp. 1017–1061. https://doi.org/10.1002/cphy.cp010221.

36. Parent, A., and Hazrati, L.-N. (1995). Functional anatomy of the basal ganglia. I. The cortico-basal ganglia-thalamo-cortical loop. Brain Research Reviews *20*, 91–127. https://doi.org/10.1016/0165-0173(94)00007-c.

37. Brimblecombe, K.R., and Cragg, S.J. (2017). The Striosome and Matrix Compartments of the Striatum: A Path through the Labyrinth from Neurochemistry toward Function. ACS Chem. Neurosci. *8*, 235–242. https://doi.org/10.1021/acschemneuro.6b00333.

38. Tajima, K., and Fukuda, T. (2013). Region-specific diversity of striosomes in the mouse striatum revealed by the differential immunoreactivities for mu-opioid receptor, substance P, and enkephalin. Neuroscience *241*, 215–228. https://doi.org/10.1016/j.neuroscience.2013.03.012.

39. Foster, N.N., Barry, J., Korobkova, L., Garcia, L., Gao, L., Becerra, M., Sherafat, Y., Peng, B., Li, X., Choi, J.-H., et al. (2021). The mouse cortico–basal ganglia–thalamic network. Nature *598*, 188–194. https://doi.org/10.1038/s41586-021-03993-3.

40. Ahmed, N.Y., Knowles, R., and Dehorter, N. (2019). New Insights Into Cholinergic Neuron Diversity. Front. Mol. Neurosci. *12*. https://doi.org/10.3389/fnmol.2019.00204.

41. Herzog, E., Bellenchi, G.C., Gras, C., Bernard, V., Ravassard, P., Bedet, C., Gasnier, B., Giros, B., and El Mestikawy, S. (2001). The Existence of a Second Vesicular Glutamate Transporter Specifies Subpopulations of Glutamatergic Neurons. J. Neurosci. *21*, RC181–RC181. https://doi.org/10.1523/JNEUROSCI.21-22-j0001.2001.

42. Fremeau, R.T., Voglmaier, S., Seal, R.P., and Edwards, R.H. (2004). VGLUTs define subsets of excitatory neurons and suggest novel roles for glutamate. Trends in Neurosciences *27*, 98–103. https://doi.org/10.1016/j.tins.2003.11.005.

43. Tran, H., Sawatari, A., and Leamey, C.A. (2023). Ten-m3 plays a role in the formation of thalamostriatal projections. Developmental Neurobiology *83*, 255–267. https://doi.org/10.1002/dneu.22927.

44. Tran, H., Sawatari, A., and Leamey, C.A. (2015). The glycoprotein Ten-m3 mediates topography and patterning of thalamostriatal projections from the parafascicular nucleus in mice. European Journal of Neuroscience *41*, 55–68. https://doi.org/10.1111/ejn.12767.

45. ISH Data :: Allen Brain Atlas: Mouse Brain https://mouse.brain-map.org/.

46. Hintiryan, H., Foster, N.N., Bowman, I., Bay, M., Song, M.Y., Gou, L., Yamashita, S., Bienkowski, M.S., Zingg, B., Zhu, M., et al. (2016). The mouse cortico-striatal projectome. Nat Neurosci *19*, 1100–1114. https://doi.org/10.1038/nn.4332.

47. Apps, R., Hawkes, R., Aoki, S., Bengtsson, F., Brown, A.M., Chen, G., Ebner, T.J., Isope, P., Jörntell, H., Lackey, E.P., et al. (2018). Cerebellar Modules and Their Role as Operational Cerebellar Processing Units. Cerebellum *17*, 654–682. https://doi.org/10.1007/s12311-018-0952-3.

48. Strick, P.L., Dum, R.P., and Fiez, J.A. (2009). Cerebellum and Nonmotor Function. Annu. Rev. Neurosci. *32*, 413–434. https://doi.org/10.1146/annurev.neuro.31.060407.125606.

49. Wagner, M.J., and Luo, L. (2020). Neocortex–Cerebellum Circuits for Cognitive Processing. Trends in Neurosciences *43*, 42–54. https://doi.org/10.1016/j.tins.2019.11.002.

50. Sillitoe, R.V., and Hawkes, R. (2013). Zones and Stripes: Development of Cerebellar Topography. In Handbook of the Cerebellum and Cerebellar Disorders (Springer, Dordrecht), pp. 43–59. https://doi.org/10.1007/978-94-007-1333-8_3.

51. Brochu, G., Maler, L., and Hawkes, R. (1990). Zebrin II: A polypeptide antigen expressed selectively by purkinje cells reveals compartments in rat and fish cerebellum. Journal of Comparative Neurology *291*, 538–552. https://doi.org/10.1002/cne.902910405.

52. Sillitoe, R.V., and Hawkes, R. (2002). Whole-mount Immunohistochemistry: A High-throughput Screen for Patterning Defects in the Mouse Cerebellum. J Histochem Cytochem. *50*, 235–244. https://doi.org/10.1177/002215540205000211.

53. Ahn, A.H., Dziennis, S., Hawkes, R., and Herrup, K. (1994). The cloning of zebrin II reveals its identity with aldolase C. Development *120*, 2081–2090. https://doi.org/10.1242/dev.120.8.2081.

54. Sarna, J.R., Marzban, H., Watanabe, M., and Hawkes, R. (2006). Complementary stripes of phospholipase Cβ3 and Cβ4 expression by Purkinje cell subsets in the mouse cerebellum. Journal of Comparative Neurology *496*, 303–313. https://doi.org/10.1002/cne.20912.

55. Igarashi, K.M., Ito, H.T., Moser, E.I., and Moser, M.-B. (2014). Functional diversity along the transverse axis of hippocampal area CA1. FEBS Letters *588*, 2470–2476. https://doi.org/10.1016/j.febslet.2014.06.004.

56. Araç, D., and Li, J. (2019). Teneurins and latrophilins: two giants meet at the synapse. Current Opinion in Structural Biology *54*, 141–151. https://doi.org/10.1016/j.sbi.2019.01.028.

57. Liakath-Ali, K., Refaee, R., and Südhof, T.C. (2024). Cartography of teneurin and latrophilin expression reveals spatiotemporal axis heterogeneity in the mouse hippocampus during development. PLOS Biology *22*, e3002599. https://doi.org/10.1371/journal.pbio.3002599.

58. Kimmel, R.A., Turnbull, D.H., Blanquet, V., Wurst, W., Loomis, C.A., and Joyner, A.L. (2000). Two lineage boundaries coordinate vertebrate apical ectodermal ridge formation. Genes Dev. *14*, 1377–1389. https://doi.org/10.1101/gad.14.11.1377.

59. Renier, N., Wu, Z., Simon, D.J., Yang, J., Ariel, P., and Tessier-Lavigne, M. (2014). iDISCO: A Simple, Rapid Method to Immunolabel Large Tissue Samples for Volume Imaging. Cell *159*, 896–910. https://doi.org/10.1016/j.cell.2014.10.010.

60. Chi, J., Crane, A., Wu, Z., and Cohen, P. (2018). Adipo-Clear: A Tissue Clearing Method for Three-Dimensional Imaging of Adipose Tissue. J Vis Exp, 58271. https://doi.org/10.3791/58271.

61. Berg, S., Kutra, D., Kroeger, T., Straehle, C.N., Kausler, B.X., Haubold, C., Schiegg, M., Ales, J., Beier, T., Rudy, M., et al. (2019). ilastik: interactive machine learning for (bio)image analysis. Nat Methods *16*, 1226–1232. https://doi.org/10.1038/s41592-019-0582-9.

62. Ma, J., He, Y., Li, F., Han, L., You, C., and Wang, B. (2024). Segment anything in medical images. Nat Commun *15*, 654. https://doi.org/10.1038/s41467-024-44824-z.

63.   Klein, S., Staring, M., Murphy, K., Viergever, M.A., and Pluim, J. (2010). elastix: A Toolbox for Intensity-Based Medical Image Registration. IEEE Trans. Med. Imaging *29*, 196–205. https://doi.org/10.1109/TMI.2009.2035616.

64.   Wang, F., Flanagan, J., Su, N., Wang, L.-C., Bui, S., Nielson, A., Wu, X., Vo, H.-T., Ma, X.-J., and Luo, Y. (2012). RNAscope: A Novel in Situ RNA Analysis Platform for Formalin-Fixed, Paraffin-Embedded Tissues. The Journal of Molecular Diagnostics *14*, 22–29. https://doi.org/10.1016/j.jmoldx.2011.08.002.

65.   Furman, M., Xu, H.-P., and Crair, M.C. (2013). Competition driven by retinal waves promotes morphological and functional synaptic development of neurons in the superior colliculus. J Neurophysiol *110*, 1441–1454. https://doi.org/10.1152/jn.01066.2012.

66.   Marrs, G.S., and Spirou, G.A. (2012). Embryonic assembly of auditory circuits: spiral ganglion and brainstem. The Journal of Physiology *590*, 2391–2408. https://doi.org/10.1113/jphysiol.2011.226886.

67.   Howell, D.M., Morgan, W.J., Jarjour, A.A., Spirou, G.A., Berrebi, A.S., Kennedy, T.E., and Mathers, P.H. (2007). Molecular guidance cues necessary for axon pathfinding from the ventral cochlear nucleus. J Comp Neurol *504*, 533–549. https://doi.org/10.1002/cne.21443.

68.   Nakamura, P.A., and Cramer, K.S. (2011). Formation and Maturation of the Calyx of Held. Hear Res *276*, 70–78. https://doi.org/10.1016/j.heares.2010.11.004.

69.   Hoffpauir, B.K., Kolson, D.R., Mathers, P.H., and Spirou, G.A. (2010). Maturation of synaptic partners: functional phenotype and synaptic organization tuned in synchrony. The Journal of Physiology *588*, 4365–4385. https://doi.org/10.1113/jphysiol.2010.198564.

70.   Leake, P.A., Snyder, R.L., and Hradek, G.T. (2002). Postnatal refinement of auditory nerve projections to the cochlear nucleus in cats. Journal of Comparative Neurology *448*, 6–27. https://doi.org/10.1002/cne.10176.

71.   Kim, G., and Kandler, K. (2003). Elimination and strengthening of glycinergic/GABAergic connections during tonotopic map formation. Nat Neurosci *6*, 282–290. https://doi.org/10.1038/nn1015.

72.   Sanes, D.H., and Siverls, V. (1991). Development and specificity of inhibitory terminal arborizations in the central nervous system. Journal of Neurobiology *22*, 837–854. https://doi.org/10.1002/neu.480220805.

73.   Kapfer, C., Seidl, A.H., Schweizer, H., and Grothe, B. (2002). Experience-dependent refinement of inhibitory inputs to auditory coincidence-detector neurons. Nat Neurosci *5*, 247–253. https://doi.org/10.1038/nn810.

74.   Pierce, E.T. (1973). Time of origin of neurons in the brain stem of the mouse. Prog Brain Res *40*, 53–65. https://doi.org/10.1016/S0079-6123(08)60679-2.

75.   Marrs, G.S., Morgan, W.J., Howell, D.M., Spirou, G.A., and Mathers, P.H. (2013). Embryonic Origins of the Mouse Superior Olivary Complex. Dev Neurobiol *73*, 384–398. https://doi.org/10.1002/dneu.22069.

76.   Shepard, A.R., Scheffel, J.L., and Yu, W.-M. (2019). Relationships between neuronal birthdates and tonotopic position in the mouse cochlear nucleus. J Comp Neurol *527*, 999–1011. https://doi.org/10.1002/cne.24575.

77.   Bayer, S.A., and Altman, J. (1987). Directions in neurogenetic gradients and patterns of anatomical connections in the telencephalon. Progress in Neurobiology *29*, 57–106. https://doi.org/10.1016/0301-0082(87)90015-3.

78.   Sohur, U.S., Padmanabhan, H.K., Kotchetkov, I.S., Menezes, J.R.L., and Macklis, J.D. (2014). Anatomic and Molecular Development of Corticostriatal Projection Neurons in Mice. Cerebral Cortex *24*, 293–303. https://doi.org/10.1093/cercor/bhs342.

79.   Mesías, R.E., Zaki, Y., Guevara, C.A., Friedman, L.G., Hussein, A., Therrien, K., Magee, A.R., Tzavaras, N., Valle, P.D., Baxter, M.G., et al. (2023). Development of prefrontal corticostriatal connectivity in mice. Preprint at bioRxiv, https://doi.org/10.1101/2023.03.14.532475 https://doi.org/10.1101/2023.03.14.532475.

Figure 1. Whole brain staining of Ten3 and Lphn2 across development.
(A) Overview of the procedure used to stain whole mouse brains for Ten3 and Lphn2 proteins.
(B) 3D staining patterns of Ten3 and Lphn2 proteins in the mouse brain across different stages of development.
E, embryonic day; P, postnatal day. Scale bar, 2 mm.

(C–E) Ten3 and Lphn2 protein expression in single optical sections in the coronal (C), horizontal (D), and sagittal (E) planes of the same 3D volume of an E17 brain imaged by light sheet microscopy. The number of each image is matched with the section location (line) in the schematic above. Superior colliculus, white arrows; superior olivary complex, magenta arrow; ventral cochlear nucleus, green arrow; spinal trigeminal nucleus, yellow arrow; thalamic nuclei, red arrow; mammillary nucleus of the hypothalamus, orange arrow; neocortex, purple arrows. Detailed expression patterns in brain regions of E17 and P2 mouse brains can be found in **Supplemental Data 1**. Scale bars, 1 mm.

In this and all subsequent figures, A, anterior; P, posterior; D, dorsal; V, ventral; M, medial; L, lateral. Unless otherwise mentioned, all Ten3 and Lphn2 proteins were visualized with anti-HA and anti-GFP antibodies in *Ten3$^{HA/HA}$;Lphn2$^{mV/mV}$* (mV for mVenus) knock-in mice and shown as cyan and yellow colors, respectively. See Figure S1 for additional data.

Figure 2. Inverse expression patterns of Ten3 and Lphn2 follow retinotopic connections
(A) Schematic diagram of retinotopic connections from the retina to the superior colliculus along the dorsal–ventral axis of the retina, with Ten3 (cyan) and Lphn2 (yellow) expression from our data superimposed.

(B) Ten3 and Lphn2 protein expression in a single coronal section of the retina. RGC axons, retinal ganglion cell axons. Scale bar, 200 μm.

(C) Flat whole mount E15 retina preparation stained for Ten3 and Lphn2 proteins. Red circle indicates optic nerve head, where RGC axons exit the retina. Scale bar, 200 μm.

(D) Quantification of Ten3 and Lphn2 proteins across the dorsal–ventral axis of the retina. See STAR Methods for details on regions analyzed (n = 4 mice; 1 retina/mouse). Mean ± SEM.

(E) 3D reconstruction of E15 optic tract from light sheet microscope images with Ten3 and Lphn2 proteins labeled. Scale bar, 400 μm. Green inset, zoom in of the optic tract (scale bar, 200 μm). Magenta inset, cross section of the optic tract (scale bar, 50 μm). Both insets reveal two discrete Ten3$^+$ bundles in the most anterior and posterior regions of the optic tract, along with a single Lphn2$^+$ bundle in the middle.

(F) Ten3 and Lphn2 protein expression in the superior colliculus in a coronal section of an E15 brain. Vertical dashed line indicates the midline. Dotted arrow indicates the axis used for quantification in (G). Scale bar, 200 μm.

(G) Quantification of Ten3 and Lphn2 protein across the medial–lateral axis of the superior colliculus (n = 3 mice). Mean ± SEM.

See Figures S2 and S3 for additional data.

Figure 3. Tonotopic expression of Ten3 and Lphn2 in auditory processing regions of the brainstem.
(A) Schematic diagram of auditory processing nuclei in the brainstem and tonotopic organization represented in a rainbow color spectrum. Cyan and yellow summarize expression patterns of Ten3 and Lphn2 from our data, which align with tonotopic maps across these nuclei. Tonotopic projections from VCN to MNTB are illustrated. DCN, dorsal cochlear nucleus; VCN, ventral cochlear nucleus; MSO, medial superior olive; LSO, lateral superior olive; SPN, superior paraolivary nucleus; MNTB, medial nucleus of the trapezoid body.
(B) Ten3 and Lphn2 protein expression in the VCN of a P0 brain. Scale bar, 200 μm.
(C) Quantification of Ten3 and Lphn2 protein across the dorsal–ventral axis of the VCN (n = 3 mice). Mean ± SEM.
(D) Ten3 and Lphn2 protein expression in the superior olivary complex of a P0 brain. Scale bar, 200 μm.
(E) Quantification of Ten3 and Lphn2 protein across the medial–lateral axis of the superior olivary complex (n = 3 mice). Mean ± SEM.

(F) Schematic diagram depicting spatial patterns of recombination of $Ten3^{cTAG}$ allele in the VCN→MNTB projection pathway in the absence (top) and presence (bottom) of $Slc17a6$-Flp.

(G, H) Ten3^{HA} and Ten3^{Halo} protein expression in the VCN (G) and MNTB (H) of $Ten3^{cTAG/cTAG}$ and $Slc17a6$-$Flp;Ten3^{cTAG/cTAG}$ P0 mice. VCN (G) and MNTB (H) are dash outlined. Scale bar, 100 μm.

(I) Quantification of relative HA staining in the MNTB of $Ten3^{cTAG/cTAG}$ (n = 5) and $Slc17a6$-$Flp;Ten3^{cTAG/cTAG}$ (n = 4) P0 mice. **, $p < 0.01$; unpaired t-test.

See Figures S2 and S4 for additional data.

Figure 4. Ten3 and Lphn2 display inverse expression in the CP and GPe

(A) Schematic diagram of connections from the caudate putamen (CP) to the globus pallidus external segment (GPe), constituting part of the indirect pathway of the basal ganglia.

(B, C) Ten3 and Lphn2 protein expression in the CP (B) and GPe (C) of a P4 brain, shown in coronal sections. CP and GPe are demarcated by the dashed outlines. Dotted lines indicate the axis used for quantification in (D) and (F). * indicate cortical axon bundles streaming through the CP. Scale bar, 200 µm.

(D) Quantification of Ten3 and Lphn2 protein across the dorsomedial–ventrolateral axis (dotted line in B) of the CP denoted.

(E) Spearman's correlation between Ten3 and Lphn2 expression in the CP (n = 3 mice). Each dot represents the normalized fluorescence intensity of Ten3 and Lphn2 in a single x-y spatial bin along the dorsomedial–

ventrolateral axis (n = 100 bins per brain). Dots of the same color are from the same mouse. See STAR Methods for details on analysis. Spearman's correlation, ****p < 0.0001.

(F, G) Same as (D, E), except for GPe. Spearman's correlation, ****p < 0.0001.

(H) Experimental design and summary of results for tracing CP→GPe connectivity pattern. LV, lentivirus. Recombination of *Ten3^{cTAG}* allele at the injection site causes Ten3^{HA+} CP neurons to express Ten3^{Halo}. Recombined Ten3^{Halo+} CP axons target GPe regions that express Ten3^{HA}.

(I) Injection site of *LV-FlpO-GFP* in the CP of *Ten3$^{cTAG/cTAG}$;Lphn2 $^{mV/mV}$* mice at P4. *LV-FlpO-GFP* injected CP neurons shown in green, Ten3^{HA} in cyan, and Ten3^{Halo} in magenta. Dotted lines outline the CP. Scale bar, 200 μm.

(J) Projection patterns of Ten3^{Halo+} CP axons (from injection in panel I) in the GPe. Ten3^{Halo+} CP axons target the Ten3^{HA+} region and avoid Lphn2^{mV+} region of the GPe. Dotted lines outline the GPe. Scale bar, 200 μm.

(K) Percent overlap of normalized fluorescence of Ten3^{Halo+} CP projections to Ten3^{HA+} or Lphn2^{mV+} GPe target regions in *Ten3$^{cTAG/cTAG}$;Lphn2 $^{mV/mV}$* mice at P4 (n = 7). Paired t-test; ****p < 0.0001.

Figure 5. Corticostriatal projections follow the Ten3→Ten3 rule

Left panels are two coronal sections of the mouse brain atlas, with red highlighting regions of recombination due to the expression patterns of the specific Flp driver. Ctx, cortex; TH, thalamus. Right panels are experimental data in P4 mice. CP is demarcated by the dashed outlines. Scale bar, 200 µm. Genotypes on the left.

(A) In the absence of Flp driver, *Ten3$^{cTAG/cTAG}$* mice express Ten3^{HA} but not Ten3^{Halo} protein in the CP (dash outlined).

(B) In the presence of *Slc32a1-Flp*, which is expressed in all GABAergic neurons, many Ten3$^+$ CP regions have strong Halo signal (yellow arrows). Some regions are high for both Halo and HA (white arrows).

(C) In the presence of *Slc17a6-Flp*, which is expressed in all excitatory neurons in the thalamus, the lack of Halo signal in CP suggests that thalamic excitatory projection neurons do not express Ten3 in their axon terminals at this stage.

(D) In the presence of *Slc17a7-Flp*, which is expressed in all excitatory neurons in the cortex, the overlap between Halo and HA signals in the CP (white arrows) indicates that Ten3$^+$ cortical neurons project their axons to Ten3$^+$ striatal regions.

Scale bar, 200 µm.

Figure 6. Inverse expression of Ten3 and Lphn2 in both Purkinje cells and cerebellar nuclei
(A) Ten3 and Lphn2 protein expression in the cerebellum of a P4 brain, shown in a coronal section. Dashed vertical line represents the midline. Scale bar, 500 µm.

(B) Enlarged view of the cerebellar cortical region outlined in (A, left), with different colored arrows indicating the transitions of Ten3- and Lphn2-expression bands in the Purkinje cell layer. Scale bar, 500 μm.

(C) Quantification of normalized Ten3 and Lphn2 protein fluorescence intensity along the PC layer across the left–right axis of the entire cerebellar cortex shown in (A). Different colored arrows indicate the same positions marked by the same colors in (B).

(D) Spearman's correlation between Ten3 and Lphn2 expression in the cerebellar cortex from data in (C); n = 3 mice. Each dot represents the normalized expression level of Ten3 and Lphn2 in a single x-y spatial bin (n = 100 bins per brain along the PC layer). Dots of the same color are from the same mouse (see STAR Methods for detail). Spearman's correlation, ****$p < 0.0001$.

(E–G) Same as (B–D), except for cerebellar nuclei. Spearman's correlation, ****$p < 0.0001$.

(H) Ten3 and Lphn2 protein expression of the right hemisphere of the cerebellar cortex at P4 in a 3D volume (H_1) and a 2D horizontal slab (H_2), after whole brain clearing and light-sheet imaging. Dashed lines denote each domain. Scale bar, 300 μm.

(I) Ten3 and Lphn2 protein expression of the right hemisphere of the cerebellar nuclei at P4 in a 3D volume (I_1). Ten3 and Lphn2 expression do not align strictly with the fastigial nucleus (FN; I_2 left), interposed nucleus (IP; I_2 middle), and dentate nucleus (DN; I_2 right). Dashed lines denote each domain. Scale bar, 300 μm.

(J) Experimental design (top) and summary of results (bottom) for tracing Ten3$^+$ PC→CN connectivity. LV, lentivirus. Recombination of $Ten3^{cTAG}$ allele at the injection site causes Ten3$^+$ PCs to switch from expressing Ten3^{HA} to Ten3^{Halo}. Recombined Ten3^{Halo+} PC axons target CN regions that express Ten3^{HA}.

(K, M, O) Low-mag coronal sections of $Ten3^{cTAG/cTAG};Lphn2^{mV/mV}$ cerebella visualized at P4. Each row indicates an individual mouse. The cerebellar cortices of these mice have been injected at P0 with $LV\text{-}FlpO\text{-}GFP$ (green in K). Ten3^{HA} is in cyan and Ten3^{Halo} in magenta.

(L, N, P) Ten3^{HA}, Lphn2^{mV}, and Ten3^{Halo} protein expression in the anterior (L), intermediate (N), and posterior (P) cerebellar nuclei of $Ten3^{cTAG/cTAG};Lphn2^{mV/mV}$ mice at P4. Panels L_{1-4}, N_{1-4}, and P_{1-4} correspond to dashed boxes in panels K, M, O, respectively. Dotted lines outline the CN.

(Q) Percent overlap of normalized fluorescence of Ten3^{Halo+} PC axons with Ten3^{HA}-positive or -negative, or Lphn2^{mV}-positive CN target regions in $Ten3^{cTAG/cTAG};Lphn2^{mV/mV}$ mice at P4 (n = 6). Paired t-test; ****$P < 0.0001$; ns, not significant.

(R) Experimental design (top) and summary of results (bottom) for tracing Lphn2$^+$ PC→CN connectivity. LV, lentivirus. Recombination of $Lphn2^{cTAG}$ allele at the injection site causes Lphn2$^+$ Purkinje cells to switch from expressing Lphn2^{FLAG} to Lphn2^{SNAP}. Recombined Lphn2^{SNAP+} PC axons target Ten3^{HA}-negative regions of the CN.

(S, U, W) Low-mag coronal sections of $Ten3^{cTAG/cTAG};Lphn2^{cTAG/cTAG}$ cerebella visualized at P4. Each row indicates an individual mouse. The cerebellar cortices of these mice have been injected at P0 with $LV\text{-}FlpO\text{-}GFP$ (green). Ten3^{HA} is in cyan and Lphn2^{SNAP} in orange.

(T, V, X) Lphn2^{SNAP+} PC axons target Ten3^{HA}-negative region in the anterior (T), intermediate (V), and posterior (X) cerebellar nuclei of $Ten3^{cTAG/cTAG};Lphn2^{cTAG/cTAG}$ mice at P4. Panels T_{1-3}, V_{1-3}, and X_{1-3} correspond to dotted box in panels S, U, W, respectively. Dotted lines outline the CN.

(Y) Percent overlap of normalized fluorescence of Lphn2^{SNAP+} PC axon projections to Ten3^{HA}-positive or Ten3^{HA}-negative CN target regions in $Ten3^{cTAG/cTAG};Lphn2^{cTAG/cTAG}$ mice at P4 (n = 9). Paired t-test; ****$P < 0.0001$. Scale bar, 200 μm for all panels unless specified.

Figure 7. Lphn2 is required for Ten3+ Purkinje cell axons to precisely target Ten3+ regions of the cerebellar nuclei.

(A, B) Experimental design and summary of results for examining Ten3→Ten3 projection in control (A) and *Lphn2* conditional knockout in the cerebellum (B). LV, lentivirus.

(C–F) Ten3HA and Ten3Halo protein expression in the anterior (C), intermediate (D), and posterior (E & F) CN of control (*En1-Cre;Lphn2+/+;Ten3cTAG/cTAG*) mice at P4.

(G–J) Ten3HA and Ten3Halo protein expression in the anterior (G), intermediate (H), and posterior (I & J) CN of *Lphn2* conditional knockout (*En1-Cre;Lphn2fl/fl;Ten3cTAG/cTAG*) mice at P4. Red arrows indicate mistargeting of Ten3Halo+ axons in a Ten3HA-negative region.

(K) Percent overlap of normalized fluorescence of Ten3Halo+ Purkinje cell axonal projections to Ten3HA-positive or -negative CN target regions in control (n = 6) and *Lphn2* conditional knockout mice (n = 7) at P4. Paired (within the same genotype) and unpaired (between genotypes) t-tests were performed. **p < 0.01. Mean ± SEM.

Scale bar, 200 μm for all panels.

EXPERIMENTAL MODEL AND STUDY PARTICIPANT DETAILS

Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rabbit anti-HA	Cell Signaling Technology	Cat# 3724S; RRID: AB_1549585
chicken anti-GFP	Aves Labs	Cat# GFP-1020
rabbit anti-Halo	Promega	Cat# G9281
rabbit anti-FLAG	Abcam	Cat# ab1162
guinea pig anti-Calbindin	Synaptic Systems	Cat# 214004
rabbit anti-MOR	ImmunoStar	Cat# 24216
rabbit anti-Aldolase C	Abcam	Cat# ab115212
mouse anti-PlcB4	Santa Cruz Biotechnology	Cat# sc166131
HA-tag–AF-555	ThermoFisher	Cat# 26183-A555
SNAP–Surface-549	NEB	Cat# S9112S
Bacterial and virus strains		
Lentivirus-FlpO-GFP	Modified from Pederick, et al. 2021	N/A
Critical commercial assays		
RNAscope® Multiplex Fluorescent Reagent Kit v2	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 323100
Experimental models: Organisms/strains		
Mouse: Ten3-conditional tag ($Ten3^{cTAG}$)	Generated from this study	
Mouse: Lphn2-conditional tag ($Lphn2^{cTAG}$)	Generated from this study	
Mouse: B6.Cg-$Slc32a1^{tm1.1(flpo)Hze}$/J		RRID: IMSR_JAX:029591
Mouse: B6;129S-$Slc17a6^{tm1.1(flpo)Hze}$/J		RRID: IMSR_JAX:030212
Mouse: B6;129S6-$Slc17a7^{em1(flpo)Tasic}$/J		RRID: IMSR_JAX:034422
Mouse: $En1^{tm2(cre)Wrst}$/J		RRID: IMSR_JAX:007916
Mouse: B6;129S6-$Adgrl2^{tm1Sud}$/J	Anderson, et al. 2017	RRID: IMSR_JAX:023401
Mouse: $Lphn2^{fl/fl}$	Anderson, et al. 2017	N/A
Oligonucleotides		
RNAscope® Probe- Mouse $Ten3$	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 411951
RNAscope® Probe- Mouse $Lphn2$-C1	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 319341

RNAscope® Probe- Mouse *Lphn2*-C2	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 319341-C2
RNAscope® Probe- Mouse *Tac1*-C1	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 410351-C2
RNAscope® Probe- Mouse *Penk*-C3	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 318761-C3
RNAscope® Probe- Mouse *Slc17a6*-C2	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 319171-C2
RNAscope® Probe- Mouse *Slc32a1*-C3	Advanced Cell Diagnostics (ACD), a Bio-Techne brand	Cat# 319191-C3
Software and algorithms		
Zen	Carl Zeiss	RRID: SCR_013672
Fiji	National Institutes of Health	RRID: SCR_002285
Illustrator	Adobe	RRID: SCR_010279
R	R Core Team	RRID: SCR_001905
R Studio	Posit, PBC	http://posit.co/
Prism 10	GraphPad	RRID:SCR_002798
Imaris	Oxford Instruments	RRID: SCR_007370
ImSpector	Miltenyi Biotec	RRID: SCR_015249
MATLAB	Mathworks	RRID: SCR_001622
Ilastik	Berg, et al. 2019	https://www.ilastik.org
Elastix	Klein, et al. 2010	https://elastix.dev/
MedSAM	Ma, et al. 2024	https://github.com/bowang-lab/MedSAM

RESOURCE AVAILBILITY

Materials Availability
Plasmids generated for the *LV-FlpO-GFP* virus will be deposited in Addgene. Mouse lines generated in this study will be deposited to the Jackson Laboratory.

Data and Code Availability
Source data are provided in the manuscript. Interactive platform (Supplementary Data 1) and raw images of whole brain staining of Ten3 and Lphn2 across development will be deposited. Code will be available on Github. All access will be released upon publication.

EXPERIMENTAL MODEL AND PARTICIPANT DETAILS

All animal procedures followed animal care guidelines approved by Stanford University's Administrative Panel on Laboratory Animal Care and Administrative Panel on Biosafety in accordance with NIH guidelines. Both male and female mice were used, where they were group housed with access to food and water *ad libitum*. Mice at embryonic day (E) 15 and 17, and postnatal day (P) 0, 2, 4, 8, 21, 42 were used.

Slc32a1-Flp (Jackson Laboratory, Stock 029591), *Slc17a6-Flp* (Jackson Laboratory, Stock 030212), and *Slc17a7-Flp* (Jackson Laboratory, Stock 034422) were bred to used *Ten3^cTAG* to switch Ten3 tags in specific cell types. *En1-Cre* (Jackson Laboratory, Stock 007916) mice[58] as well as *Lphn2^mVenus* and *Lphn2^fl/fl* mice[19] were on a C57BL/6 and 129 mixed background and were backcrossed to wild-type CD1 mice to improve pup survival rate. *Ten3^cTAG* and *Lphn3^cTAG* mice were generated in this study. All genotyping was performed as previously described[10,12,19,58].

METHOD DETAILS

<u>Generation of conditional tagged mice</u>
Ten3 and *Lphn2* conditional tag (cTAG) mice were generated by the Gene Targeting and Transgenics core at HHMI/Janelia Research Campus. Both mouse lines were generated by homologous recombination in ES cells using standard procedures. The *Ten3^cTAG* construct was designed such that exon 29 was fused in-frame with an FRT-HA tag, followed by an SV40 polyA signal and an FRT-HaloTag. A self-excising selection cassette containing a loxP2272-flanked RNA polymerase II promoter-NeoR and ACE-Cre was inserted downstream of the Halo tag. This cassette is removed automatically in F1 mice. Homology arms of 4298 bp (5') and 2231 bp (3') were retrieved from BAC clone RP23-369L1 using recombineering techniques. To enhance targeting efficiency, the CRISPR-Cas9 system was employed. The construct was co-electroporated into an F1 hybrid 129S6 × C57BL/6J ES cell line along with a guide RNA (gRNA) 5'-ATCGGCAAGAGGTAACCCCC-3' and Cas9 protein (Fisher Scientific, Cat# A36498). Forty-eight G418-resistant ES colonies were isolated and screened via nested PCR using primers located outside and inside the construct. The primers used for ES cell screening were as follows: **5' Arm:** Forward primers: *Ten3 Scr F1*: 5'- ATGGCTTTCGGAGTCACATGC-3; *Ten3 Scr F2*: 5'- TCTTAGTCTCCACCTGAGCA-3' and Reverse primers: *SV40 PA R*: 5'-GTGGTTTGTCCAAACTCATCA-3'; *Ten3 Scr 5R1*: 5'- CATCGTATGGGTAGAAGTTCC-3'; **3' Arm:** Forward primers: *ACE Scr R1*: 5'- ACAGCACCATTGTCCACTTG-3'; *ACE Scr R2*: 5'-GCTGGTAAGGGATATTTGCC-3' and Reverse primers: *Ten3 Scr R1*: 5'- AGGAGACTCTCCAGATCTGT-3'; *Ten3 Scr R2*: 5'- CTGTCATGTGGTACACGCTA-3.' Three of the 39 ES clones that tested positive for both homology arms were used to generate mice. Chimeric mice were produced by aggregating these ES cells with 8-cell-stage CD-1 embryos. Correct targeting was further confirmed by homozygosity testing in the progeny. All three ES clones contributed to the germline and produced homozygous pups. Chimeras were bred with wild-type C57BL/6J females.

The *Lphn2^cTAG* construct was designed such that exon 24 was fused in-frame with an FRT-Flag tag, followed by an SV40 polyA signal and an FRT-SNAP tag. A self-excising selection cassette containing a loxP2272-flanked RNA polymerase II promoter-NeoR and ACE-Cre was inserted downstream of the SNAP tag. This cassette is removed automatically in F1 mice. Homology arms of 3686 bp (5') and 2814 bp (3') were retrieved from BAC clone RP23-283E24 using recombineering techniques. To enhance targeting efficiency, a CRISPR-Cas9 system was employed. The construct was co-electroporated into an F1 hybrid 129S6 × C57BL/6J ES cell line along with a guide RNA (gRNA) 5'-AGTCTTTAATAGTACGGCTAAGG-3' and Cas9 protein (Fisher Scientific, Cat# A36498). Forty-eight G418-resistant ES colonies were isolated and screened via nested PCR using primers located outside and inside the construct. The primers used for ES cell screening were as follows: **5' Arm:** Forward primers: *Lphn2 Scr F1*: 5'-AGGACTGACATCACATGGTC-3'; *Lphn2 Scr F2*: 5'- GCTTCTGGAATGAGCTAGCT-3' and Reverse primers: *SV40 PA R*: 5'-GTGGTTTGTCCAAACTCATCA-3'; *Flag tag R*: 5'-CGTCGTCATCCTTGTAATCG-3'; **3' Arm:** Forward primers: *ACE Scr R1*: 5'- ACAGCACCATTGTCCACTTG-3'; *ACE Scr R2*: 5'-GCTGGTAAGGGATATTTGCC-3' and Reverse primers: *Lphn Scr R1*: 5'-ACACTTAAGAGTGCCTGTGC-3'; *Lphn Scr R2*: 5'-TTCACAGACATGTTTGCACC-3.' Three of the 25 ES clones that tested positive for both homology arms were used to generate mice. Chimeric mice were produced by aggregating these ES cells with 8-cell-stage CD-1 embryos. Correct targeting was further

confirmed by homozygosity testing in the progeny. All three ES clones contributed to the germline and produced homozygous pups. Chimeras were bred with wild-type C57BL/6J females.

Genotyping

Primers used for genotyping $Ten3^{HA}$ and $Ten3^{Halo}$ alleles include Ten3-HA-F (*CTACGATGGGTACTACGTAC*), Ten3-HA-R (*GTGGTTTGTCCAAACTCATCA*), Ten3-Halo-F (*GGTCTGAATCTGCTGCAAGA*), and Ten3-R (*TCCAGTCCAATTGAGTGCAG*). The following oligonucleotide combinations were used to detect the presence of different alleles: Ten3-HA-F/Ten3-HA-R/Ten3-R (*WT*: 329bp; $Ten3^{HA}$: 218bp), and Ten3-Halo-F/Ten3-R ($Ten3^{Halo}$: 361bp). Primers used for genotyping $Lphn2^{FLAG}$ and $Lphn2^{SNAP}$ alleles include Lphn2-FLAG-F (*AAGTATGCCCAACCTTGGAG*), Lphn2-FLAG-R (*CGTCGTCATCCTTGTAATCG*), Lphn2-SNAP-F (*GGCTGGGTTAATGTCGACAT*), and Lphn2-R (*TCACAGGACTGTGAGGTTCA*). The following oligonucleotide combinations were used to detect the presence of different alleles: Lphn2-FLAG-F/Lphn2-FLAG-R/Lphn2-R (WT: 324bp; $Lphn2^{FLAG}$: 221bp), and Lphn2-SNAP-F/Lphn2-R ($Lphn2^{SNAP}$: 223bp).

Whole brain immunolabeling and clearing

Mice were injected with 2.5% avertin and were transcardially perfused with cold phosphate buffered saline (PBS) and 4% paraformaldehyde (PFA). Dissected brains were post-fixed with 4% PFA for at room temperature for 1 hour. Whole brain immunolabeling and clearing was performed using a modified iDISCO protocol.[59,60] Brains were dehydrated through a graded methanol (MeOH) series in B1N buffer, at 20%, 40%, 60%, and 80% MeOH, followed by three washes in 100% MeOH. The incubation time for the dehydration steps were 45 minutes for embryonic brains and 1 hour for postnatal brains. Brains were then incubated overnight in a 2:1 dichloromethane (DCM):MeOH mixture at room temperature. On the next day, they were washed in 100% DCM and 100% MeOH (three times), before incubation in 1:5 H_2O_2/MeOH mixture at room temperature. The brains then went through rehydration in the reverse MeOH gradient in B1N buffer. The brains were then incubated in 5% DMSO/0.3M Glycine/PTxwH (10% Triton X–100, Tween-20, heparin, 5% NaN_3 in PBS) twice for efficient penetration of primary antibodies.

For immunostaining, brains were incubated with primary antibodies—rabbit anti-HA (1:300, Cell Signaling Technology, 3724S) and chicken anti-GFP (1:1000, Aves Labs, GFP-1020) in PTxwH for 5 days (embryonic brains, E15 and E17), 8 days (early postnatal brains, P0–P4), or 11 days (late postnatal and adult brains, P8–P42) brains, at 37°C. After three days of washing in PTxwH, the brains were then transferred into secondary antibodies– donkey anti-rabbit Alexa Fluor 555 (ThermoFisher, A32794) and donkey anti-chicken Alexa Fluor 647 (ThermoFisher, A78952) at 37°C for 4, 6, or 8 days for embryonic, early postnatal, or late postnatal/adult brains, respectively. Following secondary antibody incubation, the brains were washed in PTxwH for 3 days and PBS for 1 day.

For tissue clearing, brains were dehydrated through a graded MeOH series in water, washed three times in 100% MeOH, and incubated overnight in 2:1 DCM:MeOH mixture. Samples were then washed three times in 100% DCM before being cleared in dibenzyl ether (DBE). The cleared brains were subsequently washed daily in ethyl cinnamate for a minimum of three days prior to imaging.

Light–sheet imaging, registration, and dice coefficient analysis

Cleared whole brains were imaged using UltraMicroscope Blaze light–sheet microscope (Miltenyi Biotec) at 1× magnification. We acquired a series of images using three channels (488 nm for autofluorescence, 560 nm for Ten3, and 640 nm for Lphn2) with dynamic focusing for Ten3 and Lphn2 channels at a z-step size of 3 μm. Autofluorescence channel was imaged without dynamic focusing. Voxel sizes were 3.55 μm × 3.55 μm for E15 brains, and 5.91 μm × 5.91 μm for E17 and postnatal brains. Raw image stacks were first downsampled four times, resulting in a voxel size of 23.64 μm × 23.64 μm × 12 μm. Local Ten3- and Lphn2-positive voxels were binarized using Ilastik.[61] We note that the segmentation models were optimized for each region of interest (ROI) by training them with local signal information.

For registration, we first chose one brain from the seven imaged brains as our reference brain. This brain had the qualitatively cleanest signals (e.g., least artifacts and deformation). On the right hemisphere of the

reference brain, 3-dimensional ROI masks were prepared using a transformer-based segmentation tool[62] with a combination of the three imaging channels. The other six brains were then registered to this reference brain via non-linear transformation.[63] For inter-hemisphere analysis within the same brains, we reflected the brain with respect to its anterior–posterior axis and performed registration against the non-reflected brain to minimize variations in region boundaries. Finally, signal overlaps were quantified using Dice coefficients (MATLAB).

Immunohistochemistry

Mice were injected with 2.5% Avertin and were trascardially perfused with PBS followed by 4% PFA. Dissected brains were post-fixed with 4% PFA at room temperature for 1 hour, and then cryoprotected in 30% sucrose for 2-3 days. Brains were then embedded in OCT (Tissue-Tek), frozen in isopentane bath that was cooled on dry ice and stored at –20°C until sectioning. Using a cryostat, the frozen brains were sectioned at 60 μm thickness into PBS and stored at 4°C. Immunohistochemistry were performed as previously described.[12] Briefly, the sections were incubated in 10% NDST (0.3% Triton X–100 in PBS and 10% normal donkey serum) for two hours at room temperature, followed by incubation with primary antibodies in 5% NDST (0.3% Triton X–100 in PBS and 5% normal donkey serum) for two nights at 4°C. After multiple washes in PBST (0.3% Triton X–100 in PBS), the sections were incubated with secondary antibodies (1:500) in 5% NDST and DAPI. Sections were mounted with Fluoromount-G (SouthernBiotech, 0100-01) after multiple washes in PBST and PBS. Primary antibodies used were rabbit anti-HA (1:300, Cell Signaling Technology, 3724S), chicken anti-GFP (1:1000, Aves Labs, GFP-1020), rabbit anti-HaloTag (1:500, Promega, G9281), rabbit anti-FLAG-tag (1:1000, Abcam, ab1162), guinea pig anti-Calbindin (1:500, Synaptic Systems, 214004), rabbit anti-MOR (1:500, ImmunoStar, 24216), rabbit anti-Aldolase C (1:300, Abcam, ab115212), and mouse anti-PlcB4 (1:500, Santa Cruz Biotechnology, sc166131). Monoclonal antibody HA-tag–AF-555 (1:300, ThermoFisher, 26183-A555) and SNAP–Surface-549 (1:300, NEB, S9112S) were used. For rabbit anti-FLAG-tag, the antibody was pre–adsorbed on wild–type tissue prior to performing standard immunohistochemistry. Pre-adsorption followed the same immunohistochemistry procedure as stated above. Secondary antibodies conjugated to Alexa 488, 567, Cy3, or 647 (Jackson ImmunoResearch) were used at 1:500 from 50% glycerol stocks. Sections were imaged using the tile scan function on Zeiss LSM 900 confocal microscope with 10× magnification.

Double *in situ* hybridization

Pups of postnatal day 4 (P4) or younger were anesthetized on ice for 1–2 minutes, whereas older mice were injected with 2.5% Avertin. Mice were euthanized by decapitation and were rapidly dissected into cold PBS. The brains were then embedded into Optimum Cutting Temperature (OCT, Tissue–Tek), snap frozen in dry ice–cooled isopentane and stored in –20°C until sectioning. All samples were processed within 1 minute of dissection to preserve RNA integrity. The fresh–frozen brains were sectioned at 10 μm thickness using a cryostat and mounted onto Superfrost Plus slides (VWR). In situ hybridization was performed using the RNAscope® Multiplex Fluorescent Reagent Kit v2 (ACD Bio) following the manufacturer's protocol.[64] Briefly, sections were fixed in 4% PFA for 2 hours at RT, dehydrated in graded ethanol, and treated with Protease IV. Following target-specific probes from ACD Bio were used: *Ten3* (411951), *Lphn2*-C1 (319341), *Lphn2*-C2 (319341-C2), *Tac1* (410351-C2), *Penk* (318761-C3), *Slc17a6* (319171-C2), and *Slc32a1*(319191-C3). The sections were hybridized with these probes for 2 h at 40°C, followed by signal amplification and detection using Opal fluorophores (Akoya Biosciences; Opal 520, 570, and 650; 1:1500 dilution). Sections were counterstained with DAPI, washed, and mounted in ProLong Gold Antifade Mountant (ThermoFisher, P36930). Sections were imaged using the tile scan function on Zeiss LSM 900 confocal microscope (10× magnification) with identical acquisition settings across samples in the same batch of experiments.

Lentivirus production

Lentivirus expressing Cre-GFP under the ubiquitin-C (UBC) promoter that was originally generated by the Neuroscience Gene Vector and Virus core at Stanford University[12] was modified such that Cre-GFP was removed and FlpO-Green Lantern was inserted immediately downstream of the UBC promoter using Gibson assembly. Correct insertion of plasmid was verified by whole plasmid sequencing (Plasmidsaurus).

Recombinant vectors were generated by transient transfection of 293T cells. The supernatant was harvested 72 hours post-transfection, filtered through a 0.45 μm membrane, and concentrated via ultracentrifugation (18,000 rpm, 2 h, 4°C; SW 32 Ti rotor). Pellets were pooled, resuspended in 3.5 mL DPBS, and further purified by sucrose cushion ultracentrifugation (20% sucrose cushion, 18,000 rpm, 2 h, 4°C; SW 55 Ti rotor). Final pellets were resuspended in DPBS containing 0.001% pluronic F-68, aliquoted (10 μL), and stored at −80°C.

Stereotactic injections in neonatal mice

All stereotactic injections were performed on P0 mice. Pups were anesthetized by hypothermia on ice for 2–3 minutes, placed in the stereotaxic frame (Stoelting), and injected with 500 nL (striatum) or 300 uL (cerebellum) of virus per site at 100 nL/min using a pulled glass micropipette (Drummond). *LV-FlpO-GFP* was diluted 1:10 from the original viral stock (1.01×10^{13} copies per mL, Boston Children's Hospital Vector Core). Injection coordinates were determined relative to lambda. Striatum injections were 2.0 mm anterior, 1.3 mm lateral, and 1mm ventral from lambda. Cerebellum injections were 3 mm and 2.8 mm anterior (2 injections), 1.8 mm lateral, and 0.5 mm ventral from lambda. After injection, pups were placed on a warm heating pad for recovery before being returned to their home cage. All injections were verified by *post hoc* analysis.

QUANTIFICATION AND STATISTICAL ANALYSIS

Image Analysis

Fluorescence intensity profiles for Ten3 and Lphn2 were obtained from coronal sections of P4 brains in the CP and GPe (Figure 4B, C), as well as the Purkinje cell layer and the cerebellar nuclei (Figure 6B, E). For each image, ROIs were manually delineated based on anatomical boundaries visible in the DAPI channel. Axes used for quantification were indicated using the dotted lines in Figures 4B, 4C, and 6E. The Purkinje cell layer across the left–right axis of the entire cerebellar cortex was quantified. Mean fluorescence intensity values were extracted along these axes for both Ten3 and Lphn2 channels. To enable direct quantitative comparison across animals, intensity profiles were resampled into 100 equally spaced bins along the axis using a custom MATLAB code previously reported (MathWorks)[11]. Each bin represented the normalized mean fluorescence intensity within its corresponding spatial segment. These binned intensity values were then used for correlation analysis. Spearman's correlation coefficients were computed in R, with each brain region analyzed independently (n = 3 mice each). Data points in the scatterplots correspond to a single x–y bin, and individual mice are indicated by distinct colors.

For spatial quantification of Ten3 and Lphn2 in Figures 2G, 3C and 3E a 200-pixel-wide segmented line was drawn through the cell body layer from each end of the region of interest. These areas were defined by anatomical features or DAPI staining. Segmented lines were drawn, straightened using the Straighten function, background subtraction was performed using the Subtract function and intensity values were measured using the Plot Profile command (performed in Fiji). The intensity plots were resampled into 100 equal bins (MATLAB) and each individual trace was normalized to a maximum of 100. Traces from each mouse were averaged, normalized to a maximum of 100 again and then the normalized average trace from each mouse was combined to give the final distribution.

For relative expression analysis in Figure S2 the following method was used. Sections were stained in three batches, with each batch containing one animal of each age. All images from a region of interest in a batch were imaged using a Zeiss LSM 900 confocal microscope on the same day using the same settings. To quantify relative expression levels regions of interest were drawn using anatomical features, and fluorescent intensity values were obtained using the Histogram tool (performed in Fiji). The mean expression level was calculated for the top 10% of pixels in each image and averaged with all other values from the same region of that animal. Averages were then normalized to the age with the highest expression value. Normalized values were grouped by age and region and plotted in Prism 10 (GraphPad).

Ten3→Ten3, Lphn2→Lphn2 projection mapping analysis

Mice were only included if they met the following criteria: 1) lentivirus injections must be specifically in the striatum for experiments described in Figure 5, or the cerebellar cortex for experiments described in Figure 7; 2) axons projecting to the target region must be at the border of target Ten3–Lphn2 expression; 3) more than three

sections per animal needs to have sufficient expression of Ten3^{Halo} or Lphn2^{SNAP} axons at the target region for quantification. Mice that fulfilled these criteria are reported in Figures 5 and 7 and were included in quantifications. Confocal images of sections were acquired using tile function on the Zeiss LSM 900 confocal microscope with identical acquisition settings for all samples within an animal. To account for variability of injection sites between mice, exposure settings were adjusted individually to prevent signal saturation. Projection patterns were quantified using ImageJ/Fiji.

To quantify the overlap of Ten3^{Halo+} or Lphn2^{SNAP+} axons at Ten3^{HA+} target region, ROIs were manually delineated on the DAPI channel using anatomical landmarks. These ROIs were extracted using the clear outside function in ImageJ/Fiji. For each channel, false-positive values were subtracted, and signal thresholds were individually adjusted. A Gaussian blur (radius = 10) was applied, after which ROIs were converted to masks. Image calculator was then used to compute the number of overlapping pixels across all signal channels. Statistical analyses were performed using paired t-tests using Prism 10 (GraphPad).

Lphn2-mediated PC→CN wiring analysis

Overlap of Ten3^{Halo+} axons and Ten3^{HA+} target region in *En1-Cre;Lphn2$^{+/+}$;Ten3$^{cTAG/cTAG}$* and *En1-Cre;Lphn2$^{fl/fl}$;Ten3$^{cTAG/cTAG}$* mice were quantified using the same inclusion criteria, imaging procedures, and imaging processing methods described above. Following mask conversion, the Halo channel region was scaled down by 75% of its original size and shape from the center of the mask, and the outer 25% area along the Ten3–Lphn2 border was isolated and extracted using the clear function. An equivalent area—matching the pixel width of the outer 25% region—was also taken from the other side of the Ten3–Lphn2 border. Quantifications of axon innervation were performed using the areas on both sides of the Ten3–Lphn2 border. Corresponding areas in other channels were extracted, and Image Calculator was used to determine the number of overlapping pixels across all signal channels. Statistical analyses were performed as previously described, and all quantification was performed by an experimenter blinded to genotype.

Figure S1. Design of *Ten3^cTAG* and *Lphn2^cTAG* mice, related to Figures 1, 3, 5–7

(A) Schematic of the *Ten3^cTAG* mice. To generate the *Ten3* conditional tag mice, an HA epitope tag was added immediately upstream of the STOP codon (red octagon) of the *Ten3* open reading frame, flanked by two *FRT* sites (green triangle), followed by a Halo tag. In the absence of Flp recombinase, Ten3 protein is tagged with HA; in the presence of Flp recombinase, Ten3 protein is tagged with Halo. Bottom: Ten3 protein expression in a horizontal section across the hippocampal-entorhinal region in a postnatal day 8 (P8) mouse was visualized with anti-HA antibody in the hippocampus of *Ten3^cTAG/cTAG* mice (left), and with Halo tag ligand in the same area in *Pgk1-FlpO;Ten3^cTAG/cTAG* mice. Both samples are double stained with HA and Halo tag ligand, demonstrating lack of Halo signal in the absence of Flp and complete conversion of HA into Halo in the presence of Flp.

(B) Same as (A) except for the *Lphn2^cTAG* mice. In the absence of Flp recombinase, Lphn2 is tagged with a FLAG tag; in the presence of Flp recombinase, Lphn2 is tagged with a SNAP tag.

Scale bar, 200 μm.

Figure S2. Temporal patterns of Ten3 and Lphn2 expression across developing brain regions, related to Figures 2, 3, 4 and 6

Coronal sections across developmental stages were stained in the same batch of experiments under identical conditions and imaged under identical exposure settings to allow for comparative analysis of relative protein intensity across time. Quantifications of normalized relative levels of expression are shown in the rightmost panels. All quantification analyses were based on n = 3 mice, except cochlear nucleus (E17 and P0, n = 3; P4 and P8, n = 2). Scale bar = 200 µm for all images.

(A) Ten3 and Lphn2 protein expression in the superior colliculus (SC). Expression of both proteins peaked during embryogenesis at E15 or earlier, followed by a continuous decline after birth. The timing of peak expression coincided with the period when retinal ganglion cell (RGC) axons innervate the SC at E15,[29] and the subsequent

decline of expression corresponded to the onset of refinement of axon targeting after birth by cholinergic retinal waves.[65]

(B, C) Ten3 and Lphn2 protein expression in the ventral cochlear nucleus (B) and the nuclei of the superior olivary complex (C). In the ventral cochlear nucleus (VCN), peak expression levels of both Ten3 and Lphn2 were observed during embryogenesis at E17 or earlier, with a dramatic decrease in expression after P4. The peak expression of Ten3 and Lphn2 in the superior olivary complex (SOC) was at around birth with a sharp reduction at P4. These expression profiles aligned with the known developmental timeline of the auditory brainstem. During embryogenesis, peak expression in the VCN coincides with the period of initial axonal outgrowth at E15.5.[66–68] This is followed by peak expression at the target, the SOC, at P0, when axons mature into a calyx in the medial nucleus of the trapezoid body (MNTB)[69] and prune in the lateral superior olive (LSO), nuclei of the SOC. Expression declined by P4, a time when spontaneous cochlear activity begins to drive further synaptic refinement in the VCN and the SOC.[70–73] Notably, neurons of the VCN are born in a dorsal–ventral birth order gradient, whereas those in the SOC are born in a medial–lateral gradient.[74–76] Within both VCN and SOC, Lphn2+ regions corresponded to earlier born neurons, whereas Ten3+ regions corresponded to later-born neurons. Throughout development, Ten3 and Lphn2 maintained spatially inverse expression patterns.

(D) Ten3 and Lphn2 protein expression in the caudate putamen (CP, also known as striatum). Ten3 was highly enriched in dorsomedial CP, whereas Lphn2 was predominantly expressed in the ventrolateral CP, with highest expression of Lphn2 at E15 and Ten3 at P0. The peak expression of Lphn2 and Ten3 coincided with the timing of neuronal birth, where the spiny projection neurons constituting 95% of the CP neurons are born in a ventrolateral (E14-18) to dorsomedial (E18-P0) birth order gradient throughout the entire CP.[77] Here again, Lphn2+ regions corresponded to earlier born neurons, whereas Ten3+ regions corresponded to later-born neurons. Also, earlier in development, the intermediate zone where Ten3 and Lphn2 domains converged appeared relatively homogeneous. By P0, these regions resolved into discrete Ten3- and Lphn2-enriched clusters, forming spatially refined hotspots. This transition paralleled the onset of glutamatergic synapse innervation from both the thalamus[43] and cortex,[78,79] the maturation of synaptic connections, and dendritic spine growth in spiny projection neurons, during which bundles of corticostriatal axonal fibers begin to invade and penetrate the CP. * indicate cortical axon bundles passing through the CP, creating "holes" in the Ten3 and Lphn2 expression patterns.

(E) Ten3 and Lphn2 protein expression in the cerebellum. Inverse expression of Ten3 and Lphn2 was observed both in both Purkinje cells (PCs) and the cerebellar nuclei (CN) throughout the cerebellum. Peak expression of Lphn2 was observed E17, whereas peak Ten3 expression was at P0. As lobulation progressed postnatally, the emergence of stable, inversely patterned expression across the cerebellar cortex and nuclei along the medial–lateral axis was observed.

(F, G) Normalized relative Ten3 (F) and Lphn2 (G) protein expression in all areas throughout development. Abbreviations: SC, superior colliculus; SOC, superior oliveary complex; VCN, ventral cochlear nucleus; CP, caudate putamen.

Together, these findings reveal that Ten3 and Lphn2 exhibit inverse expression that are both spatially segregated and temporally dynamic across multiple brain systems. Peak Ten3 and Lphn2 expression was observed during embryogenesis in the SC and cochlear nucleus, whereas expression in the SOC, CP, and cerebellum peaked around birth. Notably, in the auditory, striatal, and cerebellar circuits, Lphn2 expression correlated with earlier-born and Ten3 with later-born neurons, suggesting Ten3 and Lphn2 may establish connectivity across temporally distinct neuronal populations through sequential molecular expression. In systems where timing of target selection of axons has been determined, this timing overlaps with peak expression of Ten3 and Lphn2.

Figure S3. *Ten3* and *Lphn2* mRNA display inverse expression in the superior colliculus, related to Figure 2

(A) *Ten3* and *Lphn2* mRNA expression in a single coronal section of the superior colliculus at P4.

(B, C) Co-expression analysis of *Ten3* mRNA (B) and *Lphn2* mRNA (C) with *VGLUT2* and *VGAT* mRNA, markers for excitatory and inhibitory neurons, respectively. In the high-magnification insets (correspond to the dashed rectangles to the left), arrows point to cells co-expressing *Ten3* or *Lphn2* and *VGLUT2* or *VGAT*. Side arrowheads point to cells only expressing either *Ten3* mRNA (B) or *Lphn2* mRNA (C). Upward arrowheads point to cells only expressing either *VGLUT2* mRNA (top two panels) or *VGAT* mRNA (bottom two panels). Both *Ten3* and *Lphn2* are expressed in both excitatory and inhibitory neurons.

Vertical dashed line indicates the midline. Scale bar, 200 μm (Inset 100 μm).

Figure S4. *Ten3* and *Lphn2* mRNA display inverse expression in the auditory brainstem nuclei, related to Figure 3

Ten3 and *Lphn2* mRNA expression in a single coronal section of the ventral cochlear nucleus (VCN; A) and the nuclei of the superior olivary complex (B) at E17. VCN, MNTB, SPN, and LSO are demarcated by the dashed outlines. MNTB, medial nucleus of the trapezoid body; SPN, superior paraolivary nucleus; LSO, lateral superior olive. Vertical dashed line indicates the midline. Scale bar, 200 μm.

Figure S5. Validation of Flp and Cre driver lines, related to Figures 3, 5, and 7
(A) Coronal section of *Slc32a1-Flp;Ai224* at P4 showing labeled GABAergic neurons in the CP. *Ai224* is a Flp-dependent nuclear tdTomato reporter line. A_2 is a higher-magnified view of the white dashed box in A_1, highlighting labeled GABAergic neurons within CP. Vertical dashed lines are midlines in this and all other panels.
(B) Coronal sections of *Slc17a6-Flp;Ai224* at P4 showing labeled excitatory neurons in the anterior thalamus (B_1, arrow) and posterior thalamus (B_2, arrow). Cells in the parafascicular nucleus (Pf, demarcated by dashed

circle in B_2) and ventral cochlear nucleus (B_3, arrow) are also labeled but cells in the medial nucleus of the trapezoid body of the superior olivary complex demarcated by the dashed outlines are not labeled (B_4).

(C) Coronal section of *Slc17a7-Flp;Ai224* at P4 showing labeled excitatory neurons in the cortex. C_2 is a higher-magnified view of the white dashed box in C_1. Arrowhead and arrow point to labeled excitatory neurons within the hippocampus and the cortex, respectively.

(D) Coronal section of *En1-Cre;Ai224* at P4 showing labeled Purkinje cells and the neurons in the cerebellar nuclei. D_2 is a higher-magnified view of the white dashed box in D_1, highlighting labeled neurons within the cerebellar nuclei (arrow) and the Purkinje cell layer (arrowhead).

CP, Caudate putamen; Ctx, Cortex. Dashed vertical line, midline. Scale bar, 500 μm (100 μm for B_3).

Figure S6. *Ten3* and *Lphn2* mRNA display inverse expression in CP and GPe, related to Figure 4

(A, B) *Ten3* and *Lphn2* mRNA expression in the CP (A) and GPe (B) of a P4 brain, shown in coronal sections. CP and GPe are demarcated by the dashed outlines based on DAPI co-staining (not shown). High-magnification insets correspond to the dashed rectangles in the left panel.

(C, D) Co-expression analysis of *Ten3* (E) and *Lphn2* (F) with *Tac1* (C_1; D_1) or *Penk* (C_2; D_2), markers of direct and indirect pathway spiny projection neurons, respectively. High-magnification insets correspond to the dashed rectangles in the left panel. Arrows point to cells co-expressing *Ten3* (C) or *Lphn2* (D) and $Tac1^+$ or $Penk^+$ CP regions. Side arrowheads point to cells exclusively expressing either *Ten3* mRNA (C) or *Lphn2* mRNA (D). Upward arrowheads point to cells exclusively expressing either *Tac1* mRNA (C_1; D_1) or *Penk* mRNA (C_2; D_2). Some $Ten3^+$ regions overlap with *Tac1* expression and others overlap with *Penk* expression; likewise for $Lphn2^+$ regions. Thus, both *Ten3* and *Lphn2* are expressed in spiny projection neurons in the direct and indirect pathways.

(E, F) Co-expression analysis of Ten3 and Lphn2 with μ-opioid receptor (MOR, E) or calbindin (F), markers of striosomes and extrastriosomal matrix, respectively. High-magnification insets correspond to the dashed rectangles in the left panel (E_4; F_4). Arrows point to cells co-expressing Ten3 or Lphn2 and MOR^+ (E) or $Calbindin^+$ (F) CP regions. As is evident, some $Ten3^+$ hotspots overlap with MOR expression and others overlap with calbindin expression; likewise for $Lphn2^+$ hotspots. Thus, Ten3 and Lphn2 hotspots do not correspond with the striosome/matrix division of the CP.

(G) Ten3 and Lphn2 protein expression in the thalamus at P0, 2 and 4. Note that parafascicular nucleus, a nucleus that is known to project axons to the CP, does not express high levels of Ten3 protein during development, while other thalamic nuclei, including the lateral posterior nucleus of the thalamus (LP), lateral geniculate nucleus (LGN), and ventral posterior nucleus of the thalamus (VPN), display high Ten3 expression. Scale bar, 200 μm.

Figure S7. *Ten3* and *Lphn2* mRNA and proteins co-expression analysis in the cerebellum, related to Figure 6

(A) *Ten3* and *Lphn2* mRNA expression in the cerebellum of a P4 brain, shown in a coronal section, showing their inverse expression along the medial–lateral axis. Notably, *Ten3* and *Lphn2* expression in the CN did not conform to the anatomical boundaries of the fastigial, interposed, and dentate nuclei of CN, demarcated by dashed outlines, suggesting their expression is not restricted to specific anatomical CN subdivisions but instead reveals a new anatomical organization. Vertical dashed line indicates the midline. FN, fastigial nucleus; IP, interposed nucleus; DN, dentate nucleus.

(B, C) Co-expression of *Ten3* mRNA (C) or *Lphn2* mRNA (D) with both *VGLUT2* or *VGAT* mRNA, markers for excitatory or inhibitory neurons, respectively, in the cerebellar nuclei. In high-magnification insets, arrows point to cells that co-express *Ten3* or *Lphn2* with either *VGLUT2* or *VGAT*. Side arrowheads point to cells only expressing either *Ten3* mRNA (C) or *Lphn2* mRNA (D). Upward arrowheads point to cells only expressing either *VGLUT2* mRNA (top two panels) or *VGAT* mRNA (bottom two panels). These data indicate that both *Ten3* and *Lphn2* are expressed in both glutamatergic and GABAergic neurons. PCL, Purkinje cell layer; CN, cerebellar nuclei.

(D) Quantification of spatial stereotypy of Ten3 and Lphn2 expression in 3D P0 cerebella (n = 7 brains) using the Dice similarity coefficient (DSC). DSC measures the accuracy of spatial overlap, where 0 represents complete spatial segregation and 1 represents complete spatial overlap. After signal segmentation and whole brain registration (see STAR Methods for detail), DSC analysis revealed minimal intra-hemispheric overlap between Ten3 and Lphn2 (top row)—consistent with their inverse expression patterns, and high inter-hemispheric (2^{nd} and 3^{rd} rows) and inter-animal similarity (4^{th} and 5^{th} rows) for either Ten3 or for Lphn2—indicating robust spatial stereotypy. Each dot represents a pairwise comparison; error bars represent mean ± SEM.

(E) As a comparision of spatial stereotypy of Ten3 and Lphn2 expression in the cerebellum (D), here we quantify spatial stereotypy of Ten3 and Lphn2 expression in the P0 subiculum, medial mammillary nucleus, and the superior colliculus using DSC. Comparisons include spatial overlap of Ten3 and Lphn2 within the same hemisphere (top), Ten3 or Lphn2 expression across hemispheres of the same brain (2^{nd} and 3^{rd} rows), and inter-animal comparisons of Ten3 and Lphn2 expression patterns (4^{th} and 5^{th} rows). Each dot represents a pairwise comparison; bar represent mean ± SEM.

(F, G) Ten3 and Lphn2 protein expression in the cerebellum of a P4 brain partially overlap with both aldolase C (F) and PlcB4 (G), zebrin$^+$ and zebrin$^-$ markers, respectively. Arrows indicate regions where either Ten3 or Lphn2 expression overlaps with zebrin$^+$ (F) or zebrin$^-$ (G) markers, and arrowheads indicate where either Ten3 or Lphn2 expression does not overlap with zebrin$^+$ (F) or zebrin$^-$ (G) markers. A subset of Aldolase C$^+$ regions overlapped with either Ten3$^+$ and Lphn2$^+$ domains, and similarly, a subset of Plcb4$^+$ regions comprised regions overlapping with either Ten3 or Lphn2 expression. These data indicate that both Ten3 and Lphn2 are expressed in both zebrin$^+$ and zebrin$^-$ neurons.

(H) Ten3 and Lphn2 protein expression in the cerebellar cortex visualized in serial horizontal sections following iDISCO clearing and whole brain light-sheet imaging. Magenta arrows in G_1 indicate two discrete Ten3-expressing domains that converge in more dorsal plane (H_2). Green arrows in G_2 mark another pair of Ten3 domains that merge further dorsally (G_3). In 3D, both Ten3 and Lphn2 expression form continuous, spatially distinct domains, producing four discrete domains per hemisphere. Scale bar, 700 μm.

Scale bar, 200 μm for all panels, unless specified.

SUPPLEMENTARY MATERIALS

Video S1. Three-dimensional reconstruction of whole brain Ten3 and Lphn2 expression at postnatal day 2 (P2)

Video S2. Horizontal section series showing whole brain Ten3 and Lphn2 expression at embryonic day 15 (E15). Anterior is to the left. The flythrough runs from dorsal to ventral.

Videos S3–S7. Same as Video S2 except they were from E17, P0, P2, P4, and P8, respectively.

Video S8. Three-dimensional reconstruction of Ten3 and Lphn2 expression in the caudate putamen (CP) and external segment of the globus pallidus (GPe) at P4

Video S9. Three-dimensional reconstruction of Ten3 and Lphn2 expression in the cerebellar cortex at P4

Video S10. Three-dimensional reconstruction of Ten3 and Lphn2 expression in the cerebellar nuclei at P4

Video S11. Three-dimensional reconstruction of zebrin expression in the adult cerebellar cortex.

Videos S1-10 display Ten3 in cyan and Lphn2 in yellow; for Video S11, zebrin is shown in green.

Supplementary Data 1. Interactive visualization of Ten3 and Lphn2 expression in horizonal sections of E17 and P2 mouse brains

To visualize the datasets, download all files. To visualize E17 data, open the file *Ten3_Lphn2_E17.tif* in Fiji by dragging it into the Fiji console. Next, launch the ROI manager in Fiji (*Analyze > Tools > ROI Manager*) and load the file *E17_ROIs.zip*. Each ROI corresponds to a defined region of interest: clicking on the ROI names will allow you to view Ten3 and Lphn2 expression across respective regions. Same for P2. Ten3 is in cyan and Lphn2 is in yellow.